Table 2. Method availability for cardiovascular target loci

What the standard analytic methods can and cannot deliver on these targets. Counts are loci or target-disease pairs, not patients.

Measure	Value	Of	Basis / rule
Colocalization availability at cardiovascular target loci			
Cardiovascular target loci screened	65	—	R15 rule, cardiovascular subset
Loci with a genome-wide significant disease signal in the gene window	3	of 65	disease GWAS p < 5×10^{-8}
Loci yielding a usable credible-set colocalization	2	of 65	coloc.susie, after the FIX 1 assertion
Colocalization UNAVAILABLE	**63 (96.9%)**	of 65	method availability, not evidence
Infrastructure attrition — disease GWAS cannot be region-queried			
Cardiovascular pairs lost (CV-EXTENDED, atlas definition)	**265 (59.2%)**	of 448	PRIMARY figure
Cardiovascular pairs lost (CV-PRIMARY, strict definition)	218 (54.4%)	of 401	robustness, Analysis 2
Part C-conditioned sub-rate	57 (55.3%)	of 103	SUB-RATE ONLY — conditioned on two unrelated Part C rules
The four accessions responsible for the 57 Part C-conditioned losses			
GCST90029021	10 pairs	of 57	plain gzip, no .tbi (HTTP 404) — not range-queryable
GCST90043777	9 pairs	of 57	plain gzip, no .tbi (HTTP 404) — not range-queryable
GCST90044006	23 pairs	of 57	plain gzip, no .tbi (HTTP 404) — not range-queryable
GCST90080708	15 pairs	of 57	plain gzip, no .tbi (HTTP 404) — not range-queryable

Fine-mapping diagnostics under the corrected locus-selection rule

Diagnostic	R14.1 — flawed rule (withdrawn)	R15 — corrected
Loci analysed	28 of 60 (32 superseded)	6 of 106 screened
Loci genome-wide significant on the disease side	1 / 28	6 / 6 (100%)
DENTIST residual-z RMSE, gene-expression side — median	4.46	**3.15**
DENTIST residual-z RMSE, gene-expression side — loci above 2	93%	**100% (6/6)**
DENTIST residual-z RMSE, disease side — median	0.71 (n=27)	1.70
DENTIST residual-z RMSE, disease side — loci above 2	0% (0/27)	33% (2/6)
Credible-set pair returned	2 / 28 (7%) WITHDRAWN	**4 / 6 (67%)**

R15 selection rule, declared before running: n_overlap ≥ 100; the disease GWAS has ≥ 1 variant with p < 5×10^{-8} inside the merged gene window; one locus per target by largest overlap; no cap.

R14.1 omitted the disease-significance criterion, so it admitted loci with no disease signal — its residual-z values on the disease side are an artefact of that and are withdrawn, shown only for comparison.

Of the R14.1 60-locus set, 28 completed and 32 are excluded as superseded. The residual-z diagnostic is DENTIST (Chen et al. 2021, Nat Commun 12:7117); the implementation is ours, the method is theirs.

Colocalization figures are method availability, not evidence: Gate D is closed as a documented negative result. One R14.1 locus (CA2) has a non-finite disease-side residual-z and is excluded from that row only, so its n is 27 against 28 elsewhere.

Reproduce: python src/make_table2.py Sources: cv_report.json, cv_robustness.json, d1f/partc_skipped_accessions.tsv, d1h/part3_results.tsv, d1j/r15_results.tsv