
Special Relativity

Postulates · Lorentz γ · $\Delta t = \gamma \Delta t_0$ · $L = L_0/\gamma$ · $E = \gamma mc^2$: exam-ready in 11 sheets



↑ the whole topic on one page: red = high-yield, green = the one number everything hangs on

➜ Syllabus coverage

Basic Physics lists seven lines; the third is "theory of special relativity". This notebook covers exactly that line. Photons and the quantum theory of light belong to the Quantum Physics lesson.

High-yield list

① $c = 3 \times 10^8$ m/s, same in all frames
② $\gamma = 1/\sqrt{(1 - v^2/c^2)}$, $\gamma \geq 1$
③ $\Delta t = \gamma \Delta t_0$ · $L = L_0/\gamma$
④ $u = (u' + v)/(1 + u'v/c^2)$
⑤ $E = \gamma mc^2$ · $K = (\gamma - 1)mc^2$
⑥ $E^2 = p^2c^2 + m_0^2c^4$
⑦ L_0, Δt_0 = proper length, proper time

Legend yellow = key term · red circle = must-memorise fact · **purple box** = memory trick · **red box** = exam trap · **green** = worked example

➜ Contents


✓ Frames, ether & Michelson–Morley … p. 2
✓ Einstein's two postulates (1905) … p. 3
✓ The Lorentz factor γ … p. 4
✓ Lorentz transformation … p. 5
✓ Velocity addition … p. 5
✓ Time dilation $\Delta t = \gamma \Delta t_0$ … p. 6
✓ Length contraction $L = L_0/\gamma$ … p. 7
✓ Simultaneity & the light cone … p. 8
✓ Mass, energy & momentum … p. 9
✓ Night-before p. 10 · Practice p. 11

1 Frames, the Ether & Michelson–Morley

experiment ★

Definition An **inertial frame** is one in which Newton's first law holds — a free body moves in a straight line at constant speed. Special relativity compares **only inertial frames**. Classical mechanics assumes **absolute time and length**: $t' = t$, $x' = x - vt$, correct only while $v \ll c$.

↙ Newton: "time flows the same everywhere"



equal perpendicular arms: only an "ether wind" could shift the fringes

Michelson–Morley, 1887. At the Case School of Applied Science, Cleveland. A beam is split by a **beam splitter** into two **equal, perpendicular** arms; mirrors return them to recombine. A stationary luminous ether should make the outward and return legs unequal, so rotating the apparatus should drag the fringe pattern.

Quantity	Value
Predicted fringe shift	≈ **0.4** fringe
Shift actually observed	**< 0.01** fringe
Ether drift still allowed	a few km/s at most

Memory trick The arms are **perpendicular and equal**, so in a real ether only the **round trip** differs — "parallel arm gains, perpendicular arm is dragged". The predicted shift is a **fraction of one fringe**, which is exactly why the test is so delicate.

Exam trap A **null result removes a hypothesis; it does not prove another.** Michelson won the **1907** Nobel Prize in Physics — the first American to do so — for precision optical instruments, **not** for relativity.

Worked example ✍ how big was the predicted shift? For an arm of length L and an ether drift v, $t_{\parallel} = L/(c-v) + L/(c+v) = 2Lc/(c^2-v^2)$, $t_{\perp} = 2L/\sqrt{c^2-v^2}$. With $v = 30$ km/s, $v/c = 10^{-4}$, so $1/(1-\beta^2) \approx 1 + \beta^2$ and $1/\sqrt{1-\beta^2} \approx 1 + \beta^2/2$: $\Delta t \approx (2L/c)(\beta^2/2) = L\beta^2/c$. For $L = 11$ m and $\lambda = 550$ nm, $N = c\Delta t/\lambda \approx$ **0.2 fringe**, the classical prediction of the order of 0.4 fringe — and the experiment saw less than a hundredth of it.

Still the same technique Splitting one beam into two paths and comparing them is how LIGO detects gravitational waves and how stepper lithography prints features a few nanometres wide. The null result ended the ether; the interferometer never went out of fashion.

2 Einstein's Two Postulates (1905) highest-yield ★

Why it was needed Maxwell's equations give one speed for light in vacuum and no other. Galilean kinematics plus an ether predicted the same laws everywhere **except** in the ether frame — so ether physics would be the one exception. Einstein's 1905 paper *On the Electrodynamics of Moving Bodies* removed the exception instead of explaining it.

The two postulates ① **Principle of relativity** — the laws of physics are identical in every inertial frame; no closed-laboratory experiment can detect uniform absolute motion.

② **Constancy of the speed of light** — light in vacuum travels at $c = 3 \times 10^8$ m/s in every inertial frame, whatever the motion of the source or of the observer.

Postulate	Rules out	Forces	Frequent MCQ hook
① Relativity principle	absolute rest, a privileged ether frame	all inertial frames are physically equivalent	Galilean mechanics is the $v \ll c$ limit
② c invariant	the ether drift of Michelson–Morley	no absolute time; γ finite only for $v < c$	c is exact in SI since 1983

Class	Galileo / Newton (1642–1727)	Einstein / Lorentz (1905)
Time	absolute: $t' = t$	mixed with space: $t' = \gamma(t - vx/c^2)$
Length	absolute: $x' = x - vt$	mixed with time: $x' = \gamma(x - vt)$
Velocity sum	$u = u' + v$	$u = (u' + v)/(1 + u'v/c^2)$
Speed limit	none	c

Worked example ✍ A pulse leaves a stationary source. Postulate 2 says every observer measures c. Postulate 1 says the laws hold in all frames, so the same equations that give the usual Doppler shift in air must give the **relativistic Doppler shift** in vacuum: for an observer receding at v, $f' = f\sqrt{((1-\beta)/(1+\beta))}$.

Memory trick "The **physics** is the same, **light** is the same." Postulate 1: all observers agree on the laws. Postulate 2: all observers agree on *one* speed — and that one speed is what mixes space with time.

Common misconception c is not "the speed of light" in general — it is the speed **in vacuum**, and since **1983** the metre is defined so that $c = 299\,792\,458$ m/s exactly. In glass, water or air the speed is c/n.


One line on the general theory The **general theory (1915)** drops the inertial-frame restriction and treats gravitation as curvature of spacetime. That is a separate lesson: here every frame is inertial and gravity is not in the equations at all.

3 The Lorentz Factor γ

MCQ staple

Definition Every formula in special relativity carries one number, the **Lorentz factor**: $\gamma = 1/\sqrt{(1 - v^2/c^2)} = 1/\sqrt{(1 - \beta^2)}$, with $\beta = v/c$. Properties: $\gamma \geq 1$ always, $\gamma = 1$ at $v = 0$, $\gamma \to \infty$ as $v \to c$, and $\gamma \approx 1 + \beta^2/2$ when $v \ll c$ (the Galilean limit).

v/c	β²	√(1−β²)	γ	1/γ
0	0	1.000	**1.000**	1.000
0.20	0.04	0.980	1.021	0.980
0.40	0.16	0.917	1.091	0.917
0.50	0.25	0.866	1.155	0.866
0.60	0.36	0.800	**1.250**	0.800
0.80	0.64	0.600	**1.667**	0.600
0.866 = √3/2	0.75	0.500	**2.000**	0.500
0.90	0.81	0.436	2.294	0.436
0.99	0.9801	0.141	**7.089**	0.141



flat near $v = 0$, and $\propto \infty$ at $v = c$

Worked example 1 ✍️ γ at 0.6c: $1/\sqrt{0.64} = 1/0.8 =$ **1.25**. at 0.8c: $1/\sqrt{0.36} = 1/0.6 =$ **5/3 = 1.67**. at 0.99c: $1/\sqrt{0.0199} =$ **7.09** LHC protons run at $\gamma \approx 7\,500$.

Memory trick γ is the number you **multiply time by** and the number you **divide length by**. $\sqrt{(1 - v^2/c^2)}$ is just **1/γ** — never γ.

Worked example 2: the inverse ✍️ "What speed gives $\gamma = 2$?" From $\gamma^2(1 - v^2/c^2) = 1 \to v = c\sqrt{(1 - 1/\gamma^2)}$. $\gamma = 2 \to v = c\sqrt{(3/4)} =$ **0.866c** Check the other way: $\gamma = 1.25 \to v = c\sqrt{(1 - 1/1.5625)} = c\sqrt{0.36} =$ **0.60c** ✓

Exam trap (i) $\gamma(0.6c) =$ **1.25**, not 1.67 — 1.67 belongs to 0.8c. (ii) γ can never be **less than 1**; if your answer is 0.8 you have computed $\sqrt{(1 - \beta^2)}$. (iii) $\gamma \to \infty$ does **not** forbid c: it forbids a body with mass reaching c.

All the results in one line each

Time: $\Delta t = \gamma \Delta t_0$ **Length:** $L = L_0/\gamma$

Mass: $m = \gamma m_0$ **Energy:** $E = \gamma m_0 c^2$

Momentum: $p = \gamma m_0 v$ **Velocity sum:** $u = (u'+v)/(1+u'v/c^2)$

IT link A Lorentz boost is a 2 × 2 matrix on (x, ct) — the form lattice gauge-theory codes and relativistic engines actually use:

```
gamma = 1 / sqrt(1 - v*v/(c*c));
Lambda = [[ gamma,     -gamma*v/c ],
          [ -gamma*v/c,  gamma ]];  /* boost on (x, ct) */
```

4 The Lorentz Transformation

written favourite

Definition The rules that take an event from the **rest frame S'** of an object to the lab frame **S** in which the object moves along x at speed v:

The four equations $x' = \gamma(x - vt)$ · $t' = \gamma(t - vx/c^2)$ · $y' = y$ · $z' = z$


Inverse (swap the primes, flip the sign of v): $x = \gamma(x' + vt')$ · $t = \gamma(t' + vx'/c^2)$

Derivation sketch ✍ why the coefficients must be γ

1. Space and time are homogeneous, so the transform is linear: $x' = a(x - vt)$, $t' = b(t - k \cdot v \cdot x)$
2. Postulate 2 for the ray $x = ct$, which must also have $x' = ct'$: $a(c-v) = b(c - kvc) \Rightarrow$ **$a = b$** and **$k = 1/c^2$**
3. The transform for $-v$ must be the inverse of the one for $+v$; with $a = b = A$ this gives $A^2(1 - v^2/c^2) = 1$, so $A = \gamma$

The **interval** $c^2t^2 - x^2 - y^2 - z^2$ is the same in every inertial frame, and so is c. That is the whole reason time and length cannot both stay absolute.

Frame-dependent	Form	Frame-invariant
Time, length, momentum	$\gamma\Delta t_0$ · L_0/γ · $\gamma m_0 v$	c, the interval $c^2t^2 - x^2 - y^2 - z^2$, m_0, m_0c^2, **charge**, and the causal order of timelike events
Energy	$\gamma m_0 c^2$	



the same 45° light ray in both frames — that is what fixes γ

5 Relativistic Velocity Addition

MCQ trap

Composition If an object moves at u' in S' and S' moves at v in S, then $u = (u' + v)/(1 + u'v/c^2)$. Putting $u' = c$ returns $u = c$ exactly, so c is the ceiling of every speed.

Worked example 1 ✍ Two ships head the same way, each seeing the other at 0.6c. Speed of one relative to Earth: $u = (0.6c + 0.6c)/(1 + 0.36) =$ **0.882c** — **not** 1.2c.

Worked example 2 ✍ Two bodies recede at 0.8c and 0.6c relative to Earth: $u = (0.8c + 0.6c)/(1 + 0.8 \times 0.6) =$ **0.946c**

Memory trick "γ in, γ out": every relativistic formula is the Newtonian one times γ (time, momentum, energy) or divided by γ (length).

Exam trap Galilean addition $u = u' + v$ is **wrong** except in the $v \ll c$ limit. 0.6c + 0.6c = 0.882c, never 1.2c.

6 Time Dilation $\Delta t' = \gamma \Delta t$


most tested ★

Definition Two events at the **same place in one inertial frame** are separated there by the proper time Δt_0 — the interval read by one clock present at both. Every other inertial frame measures a longer interval: $\Delta t' = \gamma \Delta t_0 = \Delta t_0 / \sqrt{(1 - v^2/c^2)}$, and always $\Delta t' \geq \Delta t_0$.

↙ books write it $\Delta t = \gamma \Delta t_0$ or $\Delta t' = \gamma \Delta t$ — check which symbol is the proper time

Derivation A: the light clock ✍ A pulse bounces between two mirrors a proper distance L_0 apart, in a clock moving at v.

1. Own frame S': straight up and down, so $\Delta t_0 = 2L_0/c$
2. Lab S: each leg is the hypotenuse of a right triangle with legs L_0 and $v\Delta t/2$
3. $c\Delta t/2 = \sqrt{(L_0^2 + v^2\Delta t^2/4)} \Rightarrow \Delta t^2(c^2 - v^2) = 4L_0^2 \Rightarrow$ **$\Delta t = \gamma \Delta t_0$**



each leg: hypotenuse $L_0\sqrt{2}$

the same two ticks; the tilted path takes $\sqrt{2}$ longer

Derivation B: from the Lorentz transform ✍ Both ticks are at the same place in S', so $\Delta x' = 0$.

1. $\Delta x' = \gamma(\Delta x + v\Delta t') = 0 \Rightarrow \Delta x = -v\Delta t'$
2. $\Delta t' = \gamma(\Delta t - v\Delta x/c^2) = \Delta t/\gamma \Rightarrow$ **$\Delta t = \gamma \Delta t'$**

Worked example 1: the muon ✍ Proper lifetime $\tau_0 = 2.2\ \mu s$ at $v = 0.8c$, $\gamma = 5/3$. $\Delta t = (5/3)(2.2\ \mu s) =$ **3.67 μs**; the lab path is $v\Delta t = 0.8c \times 3.67\ \mu s =$ **880 m**.

Worked example 2: cosmic rays ✍ At $v = 0.998c$, $\gamma = 1/\sqrt{0.004} =$ **15.8**, so $\Delta t = 15.8 \times 2.2\ \mu s =$ **34.8 μs** and the muon covers ≈ **10.4 km** — muons born high in the atmosphere reach the ground.


Memory trick "**Proper time is the shortest time.**" One clock in one place reads the smallest interval; every other frame over-measures it, because $\gamma \geq 1$.

Exam trap Both observers see the other's clock run slow — no contradiction, they are in **different** inertial frames. Only when one twin **turns round** do the readings differ, and then the stay-at-home twin is **older**.

Applications & IT link **Hafele–Keating (1971):** four caesium clocks flown round the world on jets came back shifted by exactly the amount γ and the difference in gravitational potential predict. **GPS:** satellite clocks gain ≈ **38 μs per day** overall.

7 Length Contraction $L' = L/\gamma$ high-yield ★

Definition A body moving along the line joining two observers has its length measured **smaller** than its **proper length**: $L = L_0\sqrt{1 - v^2/c^2} = L_0/\gamma$, where L_0 is the length in the body's own rest frame. Only the component **parallel to the motion** contracts; every transverse dimension is unchanged.



the shortened rod is read with **both ends at one instant of S**

Derivation ✍ (one line) The rod is at rest in S', so its ends differ by $\Delta x' = L_0$. In S both end positions must be recorded at the **same instant**, $\Delta t = 0$, so the time term vanishes and $x'_2 - x'_1 = \gamma(x_2 - x_1) = \gamma L$. Hence $L_0 = \gamma L \Rightarrow L = L_0/\gamma$.

Worked example ✍ A spaceship has proper length $L_0 = 120$ m and passes Earth at $0.8c$ ($\gamma = 5/3$): $L = 120 \times 0.6 =$ **72 m** in the Earth frame, while its height and width are unchanged — 72 m long, H high, W wide.

v	$0.20c$	$0.50c$	$0.60c$	$0.80c$	$0.866c$	$0.90c$	$0.99c$
γ	1.021	1.155	**1.25**	**1.667**	2.000	2.294	7.089
$L/L_0 = \sqrt{1-\beta^2}$	0.980	0.866	**0.800**	**0.600**	0.500	0.436	0.141

Memory trick "γ divides length, γ multiplies time, and it only touches the **along-motion** direction." Read $1/\gamma$ straight off the γ table: at $0.8c$ it is 0.6, so the rod keeps 60 % of its length.

Exam trap (i) Contraction is **not** an optical illusion of light arriving late — it survives unlimited resolution. (ii) It is **not** perpendicular to the motion. (iii) The **proper** length L_0 is what the object itself measures. (iv) Only **inertial** observers can quote $L = L_0/\gamma$.

Worked example ✍ the LHC ring The Large Hadron Collider ring is 27.4 km long in the laboratory frame. The protons circling it at $\gamma \approx 7\,500$ see their own orbit as only 27.4 km / 7 500 ≈ **3.7 m** of proper circumference. That is the "length-contracted universe" that makes particle physics convenient, and it is the same equation turned round.

Reading the sky Astrophysicists invert the same formula: a strongly redshifted galaxy is receding close to c, and its surface brightness, angular size and light-travel time must all be corrected with these factors. A quasar at $z = 6$ has been travelling since the Universe was about 1 Gyr old.

8 Relativity of Simultaneity, Spacetime & the Light Cone

conceptual ★


Definition Two events simultaneous in one frame ($\Delta t = 0$) are **not** simultaneous in a frame moving along the line joining them: $\Delta t' = -\gamma v \Delta x / c^2 \neq 0$. Their order is therefore relative. Spacetime is the single 4-dimensional entity (x, y, z, t); Minkowski (1908) showed its geometry is not Euclidean.

Separation	$\Delta s^2 = c^2\Delta t^2 - \Delta x^2 - \Delta y^2 - \Delta z^2$	All frames must agree that
Timelike ($\Delta x < c\Delta t$)	> 0	the cause comes first
Null / lightlike ($\Delta x = c\Delta t$)	$= 0$	only light does this
Spacelike ($\Delta x > c\Delta t$)	< 0	order **can** differ; no signal connects them


Worked example ✍ Two flashes occur 1 000 km apart along a rail. An observer S at the front records them **simultaneously**, so $\Delta t = 0$, $\Delta x = 1\,000$ km. A train passes at $v = 0.6c$ ($\gamma = 1.25$). In the train frame $\Delta t' = -\gamma v \Delta x / c^2 = -(1.25)(0.6c)(1.0 \times 10^6\ \text{m})/c^2 = -0.75 \times 10^6/(3 \times 10^8)\ \text{s} =$ **-2.5 ms**; the rear flash happens 2.5 ms **earlier** in the train frame.

Memory trick "Set **$\Delta t = 0$** in the transform and simultaneity splits." A pair of events one frame calls simultaneous gains a gap $\gamma v \Delta x / c^2$ in the other frame.

Exam trap Simultaneity is relative only along the relative motion: events side by side still agree ($\Delta x = 0 \Rightarrow \Delta t' = 0$). The twin paradox is **not** a violation — that worldline is not straight.



no timelike worldline leaves the 45° cone — that is why no mass reaches c




the twin paradox: the straight worldline carries the longest proper time

IT link Two datacentres call different events "simultaneous" — exactly the problem above. Light cones become wall diagrams, latency budgets become Δt, and the speed of the medium (fibre, $\approx 2 \times 10^8$ m/s) plays the part of c.

9 Relativistic Mass, Energy & Momentum $E = mc^2$ ★

Definition Total energy of a body of rest mass m_0 moving at v: $E = \gamma m_0 c^2$. Writing the historic **relativistic mass** $m = \gamma m_0$, the same statement is $E = mc^2$, i.e. $E = mc^2 + K$: the total energy is the rest energy plus the kinetic energy, and $K = (\gamma - 1)m_0c^2$.

Quantity	Formula	Notes
Mass: rest m_0, relativistic m	$m_0 \cdot \gamma m_0$	m_0 is invariant; $m \to \infty$ as $v \to c$
Rest energy E_0	m_0c^2	$1\ g \to 9 \times 10^{13}$ J
Total energy E	$\gamma m_0 c^2$	$= mc^2$ with $m = \gamma m_0$
Kinetic energy K	$(\gamma - 1)m_0c^2$	$K/E = 1 - 1/\gamma = 1 - \sqrt{(1-\beta^2)}$
Momentum p	$\gamma m_0 v$	$= mv$ only if $v \ll c$
Mass–energy relation	$E^2 = p^2c^2 + m_0^2c^4$	photon: $m_0 = 0 \Rightarrow E = pc$



E_0 stays; K grows with γ

Exam trap $E = mc^2$ is the **total** energy only when m means γm_0; with the invariant mass m_0 it reads $E = \gamma m_0 c^2$. "Mass increases with speed" is old relativistic-mass language — modern texts keep m_0 fixed. Charge and m_0 never change with speed, and K is never $\gamma m_0 c^2$.

v	γ	m/m_0	$K/(m_0c^2)$	$E/(m_0c^2)$	K/E	L/L_0	$\Delta t/\Delta t_0$
$0.6c$	1.25	1.25	0.25	1.25	0.20	0.800	1.25
$0.8c$	$5/3 = 1.67$	1.67	0.667	1.67	0.40	0.600	1.67

Memory trick "**E is the whole bar, K is the extra bit on top.**" At 0.6c a quarter of the rest energy has been added (20 % of the total); at 0.8c two thirds has (40 % of the total). Never write $K = \gamma mc^2$.

Worked example 1 ✍ electron at 0.8c

$m_0 = 9.11 \times 10^{-31}$ kg, $m_0c^2 = 0.511$ MeV, $\gamma = 5/3$. $E =$ **0.852 MeV**, $K =$ **0.341 MeV**, $m =$ **1.52×10^{-30} kg**, $p =$ **3.64×10^{-22} kg·m/s**

Worked example 2 ✍ proton at 0.6c

$m_0 = 1.673 \times 10^{-27}$ kg, $m_0c^2 = 938.3$ MeV, $\gamma = 1.25$. $E =$ **1173 MeV** $= 1.88 \times 10^{-10}$ J, $K =$ **234.6 MeV**, $p = \gamma m_0 v =$ **3.76×10^{-19} kg·m/s**

Worked example 3 ✍ mass–energy in bulk

1 g fully released: $E = 10^{-3} \times (3 \times 10^8)^2 =$ **9×10^{13} J** ≈ 21 500 t of TNT, about 1.4 times the 1945 Hiroshima yield.

Applications Nuclear reactors and weapons · PET imaging ($e^- + e^+ \to$ two 511 keV γ rays) · smoke detectors (^{241}Am α) · the LHC: two 7 TeV protons meet at 14 TeV · γ-ray astronomy, $p = E/c$.

Night-Before-the-Exam Sheet

Fact sheet


- $c = 3 \times 10^8$ m/s (exactly 299 792 458 m/s)
- $\gamma = 1/\sqrt{(1 - v^2/c^2)} \geq 1$, $\beta = v/c$
- $\Delta t = \gamma \Delta t_0$ (Δt_0 = proper time)
- $L = L_0/\gamma$, along the motion only
- $u = (u' + v)/(1 + u'v/c^2)$
- $\Delta t' = -\gamma v \Delta x/c^2$ if $\Delta t = 0$
- $E = \gamma m_0 c^2$, $E_0 = m_0 c^2$
- $K = (\gamma - 1) m_0 c^2$
- $m = \gamma m_0$ (historic), $p = \gamma m_0 v$
- $E^2 = p^2c^2 + m_0^2 c^4$, photon $E = pc$
- $\gamma(0.6c) = 1.25$, $\gamma(0.8c) = 1.667$
- $\gamma(0.866c) = 2$, $\gamma(0.99c) = 7.09$
- $1\ g \rightarrow 9 \times 10^{13}$ J
- 1 eV $= 1.602 \times 10^{-19}$ J

→ Top 12 one-liners for MCQs

1. Special relativity: Einstein, **1905**
2. Michelson–Morley ether test: **1887**
3. Postulate 2: c is the same for all inertial frames
4. $\gamma \geq 1$; $\gamma = 1$ only at $v = 0$
5. Proper time is the **shortest** interval
6. Contraction is only along the motion
7. Nothing with mass reaches c
8. Simultaneity is relative along the motion
9. Timelike events keep their order in every frame
10. $K = (\gamma - 1)m_0c^2$, never $\gamma m_0 c^2$
11. Photon: $m_0 = 0$ and $E = pc$
12. Michelson: Nobel Prize in Physics, **1907**

Top traps $\gamma(0.6c) = 1.25$, not 1.67 · $\sqrt{(1 - v^2/c^2)}$ is $1/\gamma$, never γ · L contracts along the motion, never across it · $0.6c + 0.6c = 0.882c$, never $1.2c$ · $K \neq \gamma mc^2$ · a null result removes a hypothesis, it does not establish one · c is the speed **in vacuum** only.

All tricks together "The physics is the same, light is the same" (postulates) · γ multiplies time, divides length, grows the energy bar · $\sqrt{(1 - \beta^2)}$ is the length fraction and $1/\gamma$ · $\Delta t = 0$ in the transform makes simultaneity split · proper time is shortest · E is the whole bar, K is the extra bit on top · only side-on lengths shrink.



two postulates in, five results out — every one of them is just γ

✓ Practice MCQs (exam pattern, 10 marks)

try first, then check

1. Speed of light in vacuum, for all frames:
(a) set by the source (b) larger when closing
(c) same for all frames (d) varies with the frame

2. The Lorentz factor at $v = 0.8c$ is:
(a) $5/3 = 1.67$ (b) 1.25
(c) 0.60 (d) $1/3$

3. $\gamma = 2$ corresponds to a speed of:
(a) $0.50c$ (b) $0.99c$
(c) $0.866c$ (d) $0.75c$

4. Proper length 400 m at $0.8c$. Length seen:
(a) 667 m (b) 240 m
(c) 480 m (d) 400 m

5. A light clock ticks 2.00 s apart. At $0.6c$:
(a) 1.60 s (b) 2.00 s
(c) 3.20 s (d) 2.50 s

6. The energy equivalent of 1 g of mass is:
(a) 9×10^{13} J (b) 9×10^{16} J
(c) 3×10^{5} J (d) 9×10^{10} J

7. Two ships at $0.6c$ each, same direction:
(a) $1.20c$ (b) $0.60c$
(c) $0.75c$ (d) $0.882c$

8. Einstein published it in:
(a) 1901 (b) 1915
(c) 1905 (d) 1921

9. The Michelson–Morley experiment (1887):
(a) shift proving ether (b) no shift at all
(c) light slows in water (d) c varies with speed

10. Total energy with factor γ ($E_0 = mc^2$):
(a) $(\gamma - 1)E_0$ (b) E_0
(c) E_0/γ (d) γE_0

Answer key → 1-c · 2-a · 3-c · 4-b · 5-d · 6-a · 7-d · 8-c · 9-b · 10-d

Written question State Einstein's two postulates. Using the Lorentz transformation, derive time dilation for a clock moving at $v = 0.8c$, and find the length of a 200 m rod moving along its length at $0.8c$. 10 marks

1. **P1** — the laws of physics are identical in every inertial frame. **P2** — light in vacuum travels at c for every inertial observer, whatever the source or observer does. 3
2. Lorentz transform: $x' = \gamma(x - vt)$, $t' = \gamma(t - vx/c^2)$, $\gamma = 1/\sqrt{(1 - v^2/c^2)}$. 2
3. Time dilation: both ticks are at one place in the clock's frame, so $\Delta x' = 0$ and $\Delta x = -v\Delta t'$; then $\Delta t' = \gamma(\Delta t - v\Delta x/c^2) = \Delta t/\gamma$, i.e. $\boldsymbol{\Delta t = \gamma \Delta t'}$. 3
4. At $0.8c$, $\gamma = 1/\sqrt{0.36} = \mathbf{5/3}$: 3.00 s on the clock measures 5.00 s in the lab. 1
5. Length: read both ends at one instant of the measuring frame, $\Delta t = 0$, so $\Delta x' = \gamma \Delta x = L_0$ and $\boldsymbol{L = L_0/\gamma = 120\ m}$. 1

→ Before the exam ✓

- ✓ Can state both postulates
- ✓ Can compute and invert γ
- ✓ Can derive $\Delta t = \gamma \Delta t_0$
- ✓ Knows when $\Delta t' = -\gamma v \Delta x/c^2$ works