

# A Centrifugation-Assisted Lateral Flow Assay Platform for Bioassay Sensitivity and Visualization Enhancement

Hang Yuan<sup>1,†</sup>, Ruiqi Yong<sup>1,†</sup>, Shuhe Liu<sup>1</sup>,  
Wenwen Yuan<sup>1,2</sup>, Sixuan Duan<sup>1,2</sup>, Miao Lu<sup>1</sup>, and Pengfei Song<sup>1,2\*</sup>

<sup>1</sup>School of Advanced Technology, Xi'an Jiaotong - Liverpool University, Suzhou, China

Contact:

<sup>2</sup>Department of Electrical and Electronic Engineering, University of Liverpool, Liverpool, UK    Pengfei.Song@xjtlu.edu.cn

## ABSTRACT

This study presents a centrifugation-assisted lateral flow assay (CLFA) platform that addresses the limited sensitivity and uncontrollable incubation time of traditional lateral flow assays (LFA). The CLFA platform generates a centrifugal force by controlling the motor to improve the flow rate controllability and optimize incubation time at the test line (T line). As a proof-of-concept, human chorionic gonadotropin (hCG) was chosen for quantification to validate the sensitivity and visualization enhancements. The centrifuged experiment group exhibits better linearity ( $R^2=0.9795$ ) with the logarithm of the concentration (0-2000 mIU/mL) than the uncentrifuged control group. And the T line intensity enhances by an average of 8.23%. Our CLFA platform potentially improves early screening and advances point-of-care testing development.

## INTRODUCTION

- ★ Lateral flow assay (LFA) is one of the most promising techniques for point-of-care testing (POCT) due to its user-friendliness and cost-effectiveness [1].
- ★ Limited sensitivity and uncontrollable incubation time are the main challenges of LFA [2, 3].
- ★ We developed a simple centrifugation-assisted LFA (CLFA) platform to improve flow rate controllability and optimize incubation time.
- ★ We demonstrated the low-level human chorionic gonadotropin (hCG) quantification on this platform to validate the improved sensitivity and visualization.

## RESULTS




Figure 2. The results of hCG test strips with different concentrations (n=5).

### Test strip results:

- Fig. 2(a) shows the test strip results of the control and experiment groups taken by the smartphone.
- Fig. 2(b) shows the ratio of the mean grayscale of the T line to that of the C line.

### Better linear relationship:


- The centrifuged experiment group indicates a better linear relationship with the logarithm of the concentration from 0 to 2000 mIU/mL ( $R^2=0.9795$ ) than the uncentrifuged control group.

### Visualization enhancement:

- The experiment group demonstrates an average increase of 8.23% in the T line intensity at the same hCG concentration.

### ACKNOWLEDGMENTS:

This work is supported by XJTLU Key Programme Special Fund – Exploratory Research Programme (KSF-E-39), and XJTLU Research Development Fund (RDF-18-02-20).



Xi'an Jiaotong-Liverpool University

西交利物浦大学

Download  
paper:



Hang Yuan

Undergraduate Student (Year 3)

Xi'an Jiaotong-Liverpool University

Email: Hang.Yuan20@student.xjtlu.edu.cn

Web: <https://enderhangyuan.github.io/>

Seeking a PhD Position



### REFERENCES:

- [1] D. Lee et al., "Capillary flow control in lateral flow assays via delaminating timers," *Science Advances*, vol. 7, pp. eabf9833, 2021.
- [2] D. M. Kainz et al., "Flow control for lateral flow strips with centrifugal microfluidics," *Lab on a Chip*, vol. 19, pp. 2718-2727, 2019.
- [3] M. Shen et al., "An enhanced centrifugation-assisted lateral flow immunoassay for the point-of-care detection of protein biomarkers," *Lab on a Chip*, vol. 20, pp. 2626-2634, 2020.