
Top Pfam domains by abundance per strain
Symbiodinium microadriaticum
Beta_helix 7.3%
Gp_dh_C 6.2%
Gp_dh_N 6.2%
Sacchrp_dh_NADP 3.0%
Enolase_C 3.0%
Enolase_N 3.0%
ABC_tran 2.1%
ABC_membrane 2.0%
HSP70 2.1%
PA14 7.5%
TIG 7.5%
G8 7.3%
Tubulin 3.2%
Tubulin_C 3.2%
Pkinase 6.5%
Annexin 6.1%
EF-hand_1 3.5%
EF-hand_7 2.8%
cNMP_binding 2.4%
TIP120 2.8%
0
10
20
30
Number of detected proteins
Symbiodinium natans
Beta_helix 3.1%
Inhibitor_I69 0.8%
Peptidase_C10 0.8%
Peptidase_M14 0.8%
Peptidase_M43 0.8%
Reprolysin_4 0.8%
TIG 4.0%
PA14 4.0%
G8 3.1%
fn3 2.4%
Sushi 0.9%
Lectin_C 0.9%
Cadherin 0.8%
I-set 1.5%
Fea1 3.8%
CBM_3 0.9%
Cohesin 0.9%
CBM_X2 0.9%
MAM 0.8%
DUF5011 0.8%
0
10
20
30
Number of detected proteins
Breviolum minutum
Gp_dh_C 7.6%
Gp_dh_N 7.6%
Enolase_C 3.0%
Enolase_N 3.0%
Beta_helix 2.6%
Sod_Fe_C 3.2%
Sod_Fe_N 3.2%
TIG 2.8%
fn3 2.6%
G8 2.6%
PA14 2.6%
CBM_1 2.4%
Tubulin 2.2%
Tubulin_C 2.2%
EF-hand_5 10.5%
Pkinase 4.4%
I-set 2.5%
Annexin 2.5%
EF-hand_1 2.3%
EF-hand_7 1.8%
0
10
20
30
Number of detected proteins
% detected signal (within strain)
12%
6%
1%
Category
Metabolic enzymes
Transport & membrane
Redox & oxidative stress
Structural / cytoskeletal
Signaling & regulation
Other / uncategorised