

## Rules for integrands involving inverse tangents and cotangents

1.  $\int u \operatorname{ArcTan}[a + b x^n] dx$

**1:**  $\int \operatorname{ArcTan}[a + b x^n] dx$

Derivation: Integration by parts

Rule:

$$\int \operatorname{ArcTan}[a + b x^n] dx \rightarrow x \operatorname{ArcTan}[a + b x^n] - b n \int \frac{x^n}{1 + a^2 + 2 a b x^n + b^2 x^{2n}} dx$$

Program code:

```
Int[ArcTan[a_+b_.*x_^n_],x_Symbol] :=
  x*ArcTan[a+b*x^n] -
  b*n*Int[x^n/(1+a^2+2*a*b*x^n+b^2*x^(2*n)),x] /;
FreeQ[{a,b,n},x]
```

```
Int[ArcCot[a_+b_.*x_^n_],x_Symbol] :=
  x*ArcCot[a+b*x^n] +
  b*n*Int[x^n/(1+a^2+2*a*b*x^n+b^2*x^(2*n)),x] /;
FreeQ[{a,b,n},x]
```

2.  $\int x^m \operatorname{ArcTan}[a + b x^n] dx$

**1:**  $\int \frac{\operatorname{ArcTan}[a + b x^n]}{x} dx$

Derivation: Algebraic expansion

Basis:  $\operatorname{ArcTan}[z] = \frac{1}{2} i \operatorname{Log}[1 - i z] - \frac{1}{2} i \operatorname{Log}[1 + i z]$

Rule:

$$\int \frac{\text{ArcTan}[a + b x^n]}{x} dx \rightarrow \frac{i}{2} \int \frac{\text{Log}[1 - i a - i b x^n]}{x} dx - \frac{i}{2} \int \frac{\text{Log}[1 + i a + i b x^n]}{x} dx$$

Program code:

```
Int[ArcTan[a_+b_*x_^n_]/x_,x_Symbol] :=
  I/2*Int[Log[1-I*a-I*b*x^n]/x,x] -
  I/2*Int[Log[1+I*a+I*b*x^n]/x,x] /;
FreeQ[{a,b,n},x]
```

```
Int[ArcCot[a_+b_*x_^n_]/x_,x_Symbol] :=
  I/2*Int[Log[1-I/(a+b*x^n)]/x,x] -
  I/2*Int[Log[1+I/(a+b*x^n)]/x,x] /;
FreeQ[{a,b,n},x]
```

**2:**  $\int x^m \text{ArcTan}[a + b x^n] dx$  when  $(m | n) \in \mathbb{Q} \wedge m+1 \neq 0 \wedge m+1 \neq n$

Reference: G&R 2.851, CRC 456, A&S 4.4.69

Reference: G&R 2.852, CRC 458, A&S 4.4.71

Derivation: Integration by parts

Rule: If  $(m | n) \in \mathbb{Q} \wedge m+1 \neq 0 \wedge m+1 \neq n$ , then

$$\int x^m \text{ArcTan}[a + b x^n] dx \rightarrow \frac{x^{m+1} \text{ArcTan}[a + b x^n]}{m+1} - \frac{b n}{m+1} \int \frac{x^{m+n}}{1 + a^2 + 2 a b x^n + b^2 x^{2n}} dx$$

Program code:

```
Int[x_^m_*ArcTan[a_+b_*x_^n_],x_Symbol] :=
  x^(m+1)*ArcTan[a+b*x^n]/(m+1) -
  b*n/(m+1)*Int[x^(m+n)/(1+a^2+2*a*b*x^n+b^2*x^(2*n)),x] /;
FreeQ[{a,b},x] && RationalQ[m,n] && m+1!=0 && m+1!=n
```

```

Int[x_^m_.*ArcCot[a_+b_.*x_^n_],x_Symbol] :=
  x^(m+1)*ArcCot[a+b*x^n]/(m+1) +
  b*n/(m+1)*Int[x^(m+n)/(1+a^2+2*a*b*x^n+b^2*x^(2*n)),x] /;
FreeQ[{a,b},x] && RationalQ[m,n] && m+1!=0 && m+1!=n

```

2.  $\int u \operatorname{ArcTan}[a + b f^{c+d x}] dx$

1:  $\int \operatorname{ArcTan}[a + b f^{c+d x}] dx$

Derivation: Algebraic expansion

Basis:  $\operatorname{ArcTan}[z] == \frac{1}{2} i \operatorname{Log}[1 - i z] - \frac{1}{2} i \operatorname{Log}[1 + i z]$

Rule:

$$\int \operatorname{ArcTan}[a + b f^{c+d x}] dx \rightarrow \frac{i}{2} \int \operatorname{Log}[1 - i (a + b f^{c+d x})] dx - \frac{i}{2} \int \operatorname{Log}[1 + i (a + b f^{c+d x})] dx$$

Program code:

```

Int[ArcTan[a_+b_.*f^(c_+d_.*x_)],x_Symbol] :=
  I/2*Int[Log[1-I*a-I*b*f^(c+d*x)],x] -
  I/2*Int[Log[1+I*a+I*b*f^(c+d*x)],x] /;
FreeQ[{a,b,c,d,f},x]

```

```

Int[ArcCot[a_+b_.*f^(c_+d_.*x_)],x_Symbol] :=
  I/2*Int[Log[1-I/(a+b*f^(c+d*x))],x] -
  I/2*Int[Log[1+I/(a+b*f^(c+d*x))],x] /;
FreeQ[{a,b,c,d,f},x]

```

**2:**  $\int x^m \operatorname{ArcTan}[a + b f^{c+dx}] dx$  when  $m \in \mathbb{Z} \wedge m > 0$

Derivation: Algebraic expansion

Basis:  $\operatorname{ArcTan}[z] == \frac{1}{2} i \operatorname{Log}[1 - iz] - \frac{1}{2} i \operatorname{Log}[1 + iz]$

Rule: If  $m \in \mathbb{Z} \wedge m > 0$ , then

$$\int x^m \operatorname{ArcTan}[a + b f^{c+dx}] dx \rightarrow \frac{i}{2} \int x^m \operatorname{Log}[1 - i(a + b f^{c+dx})] dx - \frac{i}{2} \int x^m \operatorname{Log}[1 + i(a + b f^{c+dx})] dx$$

Program code:

```
Int[x_^m_.*ArcTan[a_+b_*f^(c_+d_*x_)],x_Symbol] :=
  I/2*Int[x^m*Log[1-I*a-I*b*f^(c+d*x)],x] -
  I/2*Int[x^m*Log[1+I*a+I*b*f^(c+d*x)],x] /;
FreeQ[{a,b,c,d,f},x] && IntegerQ[m] && m>0
```

```
Int[x_^m_.*ArcCot[a_+b_*f^(c_+d_*x_)],x_Symbol] :=
  I/2*Int[x^m*Log[1-I/(a+b*f^(c+d*x))],x] -
  I/2*Int[x^m*Log[1+I/(a+b*f^(c+d*x))],x] /;
FreeQ[{a,b,c,d,f},x] && IntegerQ[m] && m>0
```

**3:**  $\int u \operatorname{ArcTan}\left[\frac{c}{a + b x^n}\right]^m dx$

Derivation: Algebraic simplification

Basis:  $\operatorname{ArcTan}[z] == \operatorname{ArcCot}\left[\frac{1}{z}\right]$

Rule:

$$\int u \operatorname{ArcTan}\left[\frac{c}{a + b x^n}\right]^m dx \rightarrow \int u \operatorname{ArcCot}\left[\frac{a}{c} + \frac{b x^n}{c}\right]^m dx$$

Program code:

```
Int[u_.*ArcTan[c_./(a_.+b_.*x_^n_.)]^m_.,x_Symbol] :=
  Int[u*ArcCot[a/c+b*x^n/c]^m,x] /;
FreeQ[{a,b,c,n,m},x]
```

```
Int[u_.*ArcCot[c_./(a_.+b_.*x_^n_.)]^m_.,x_Symbol] :=
  Int[u*ArcTan[a/c+b*x^n/c]^m,x] /;
FreeQ[{a,b,c,n,m},x]
```

**4.**  $\int u \operatorname{ArcTan}\left[\frac{c x}{\sqrt{a + b x^2}}\right] dx$  when  $b + c^2 == 0$

**1:**  $\int \operatorname{ArcTan}\left[\frac{c x}{\sqrt{a + b x^2}}\right] dx$  when  $b + c^2 == 0$

Derivation: Integration by parts

Basis: If  $b + c^2 == 0$ , then  $\partial_x \operatorname{ArcTan}\left[\frac{c x}{\sqrt{a + b x^2}}\right] == \frac{c}{\sqrt{a + b x^2}}$

Rule: If  $b + c^2 == 0$ , then

$$\int \text{ArcTan}\left[\frac{c x}{\sqrt{a+b x^2}}\right] dx \rightarrow x \text{ArcTan}\left[\frac{c x}{\sqrt{a+b x^2}}\right] - c \int \frac{x}{\sqrt{a+b x^2}} dx$$

Program code:

```
Int[ArcTan[c_*x_/Sqrt[a_+b_*x_^2]],x_Symbol] :=
  x*ArcTan[(c*x)/Sqrt[a+b*x^2]] - c*Int[x/Sqrt[a+b*x^2],x] /;
FreeQ[{a,b,c},x] && EqQ[b+c^2,0]
```

```
Int[ArcCot[c_*x_/Sqrt[a_+b_*x_^2]],x_Symbol] :=
  x*ArcCot[(c*x)/Sqrt[a+b*x^2]] + c*Int[x/Sqrt[a+b*x^2],x] /;
FreeQ[{a,b,c},x] && EqQ[b+c^2,0]
```

2.  $\int (d x)^m \text{ArcTan}\left[\frac{c x}{\sqrt{a+b x^2}}\right] dx$  when  $b+c^2=0$

1:  $\int \frac{\text{ArcTan}\left[\frac{c x}{\sqrt{a+b x^2}}\right]}{x} dx$  when  $b+c^2=0$

Derivation: Integration by parts

Basis: If  $b+c^2=0$ , then  $\partial_x \text{ArcTan}\left[\frac{c x}{\sqrt{a+b x^2}}\right] = \frac{c}{\sqrt{a+b x^2}}$

Rule: If  $b+c^2=0$ , then

$$\int \frac{\text{ArcTan}\left[\frac{c x}{\sqrt{a+b x^2}}\right]}{x} dx \rightarrow \text{ArcTan}\left[\frac{c x}{\sqrt{a+b x^2}}\right] \text{Log}[x] - c \int \frac{\text{Log}[x]}{\sqrt{a+b x^2}} dx$$

Program code:

```
Int[ArcTan[c_*x_/Sqrt[a_+b_*x_^2]]/x_,x_Symbol] :=
  ArcTan[c*x/Sqrt[a+b*x^2]]*Log[x] - c*Int[Log[x]/Sqrt[a+b*x^2],x] /;
FreeQ[{a,b,c},x] && EqQ[b+c^2,0]
```

```
Int[ArcCot[c_*x_/Sqrt[a_+b_*x_^2]]/x_,x_Symbol] :=
  ArcCot[c*x/Sqrt[a+b*x^2]]*Log[x] + c*Int[Log[x]/Sqrt[a+b*x^2],x] /;
FreeQ[{a,b,c},x] && EqQ[b+c^2,0]
```

**2:**  $\int (dx)^m \text{ArcTan}\left[\frac{cx}{\sqrt{a+bx^2}}\right] dx$  when  $b+c^2 = 0 \wedge m \neq -1$

Derivation: Integration by parts

Basis: If  $b+c^2 = 0$ , then  $\partial_x \text{ArcTan}\left[\frac{cx}{\sqrt{a+bx^2}}\right] = \frac{c}{\sqrt{a+bx^2}}$

Rule: If  $b+c^2 = 0 \wedge m \neq -1$ , then

$$\int (dx)^m \text{ArcTan}\left[\frac{cx}{\sqrt{a+bx^2}}\right] dx \rightarrow \frac{(dx)^{m+1} \text{ArcTan}\left[\frac{cx}{\sqrt{a+bx^2}}\right]}{d(m+1)} - \frac{c}{d(m+1)} \int \frac{(dx)^{m+1}}{\sqrt{a+bx^2}} dx$$

Program code:

```
Int[(d_*x_)^m_*ArcTan[c_*x_/Sqrt[a_+b_*x_^2]],x_Symbol] :=
  (d*x)^(m+1)*ArcTan[c*x/Sqrt[a+b*x^2]]/(d*(m+1)) - c/(d*(m+1))*Int[(d*x)^(m+1)/Sqrt[a+b*x^2],x] /;
FreeQ[{a,b,c,d,m},x] && EqQ[b+c^2,0] && NeQ[m,-1]
```

```
Int[(d_*x_)^m_*ArcCot[c_*x_/Sqrt[a_+b_*x_^2]],x_Symbol] :=
  (d*x)^(m+1)*ArcCot[c*x/Sqrt[a+b*x^2]]/(d*(m+1)) + c/(d*(m+1))*Int[(d*x)^(m+1)/Sqrt[a+b*x^2],x] /;
FreeQ[{a,b,c,d,m},x] && EqQ[b+c^2,0] && NeQ[m,-1]
```

3.  $\int \frac{\text{ArcTan}\left[\frac{cx}{\sqrt{a+bx^2}}\right]^m}{\sqrt{d+ex^2}} dx$  when  $b+c^2 = 0 \wedge bd-ae = 0$

$$1. \int \frac{\text{ArcTan}\left[\frac{c x}{\sqrt{a+b x^2}}\right]^m}{\sqrt{a+b x^2}} dx \text{ when } b + c^2 = 0$$

$$1: \int \frac{1}{\sqrt{a+b x^2} \text{ArcTan}\left[\frac{c x}{\sqrt{a+b x^2}}\right]} dx \text{ when } b + c^2 = 0$$

Derivation: Reciprocal rule for integration

Basis: If  $b + c^2 = 0$ , then  $\partial_x \text{ArcTan}\left[\frac{c x}{\sqrt{a+b x^2}}\right] = \frac{c}{\sqrt{a+b x^2}}$

Rule: If  $b + c^2 = 0$ , then

$$\int \frac{1}{\sqrt{a+b x^2} \text{ArcTan}\left[\frac{c x}{\sqrt{a+b x^2}}\right]} dx \rightarrow \frac{1}{c} \text{Log}\left[\text{ArcTan}\left[\frac{c x}{\sqrt{a+b x^2}}\right]\right]$$

Program code:

```
Int[1/(Sqrt[a_.+b_.*x_^2]*ArcTan[c_.*x_/Sqrt[a_.+b_.*x_^2]]),x_Symbol] :=
  1/c*Log[ArcTan[c*x/Sqrt[a+b*x^2]]] /;
FreeQ[{a,b,c},x] && EqQ[b+c^2,0]
```

```
Int[1/(Sqrt[a_.+b_.*x_^2]*ArcCot[c_.*x_/Sqrt[a_.+b_.*x_^2]]),x_Symbol] :=
  -1/c*Log[ArcCot[c*x/Sqrt[a+b*x^2]]] /;
FreeQ[{a,b,c},x] && EqQ[b+c^2,0]
```

$$2: \int \frac{\text{ArcTan}\left[\frac{c x}{\sqrt{a+b x^2}}\right]^m}{\sqrt{a+b x^2}} dx \text{ when } b + c^2 = 0 \wedge m \neq -1$$

Derivation: Power rule for integration

Basis: If  $b + c^2 = 0$ , then  $\partial_x \text{ArcTan}\left[\frac{c x}{\sqrt{a+b x^2}}\right] = \frac{c}{\sqrt{a+b x^2}}$



Rule: If  $b + c^2 = 0 \wedge m \neq -1$ , then

$$\int \frac{\text{ArcTan}\left[\frac{c x}{\sqrt{a+b x^2}}\right]^m}{\sqrt{a+b x^2}} dx \rightarrow \frac{\text{ArcTan}\left[\frac{c x}{\sqrt{a+b x^2}}\right]^{m+1}}{c (m+1)}$$

Program code:

```
Int[ArcTan[c_.*x_/Sqrt[a_.+b_.*x_^2]]^m_/Sqrt[a_.+b_.*x_^2],x_Symbol] :=
  ArcTan[c*x/Sqrt[a+b*x^2]]^(m+1)/(c*(m+1)) /;
FreeQ[{a,b,c,m},x] && EqQ[b+c^2,0] && NeQ[m,-1]
```

```
Int[ArcCot[c_.*x_/Sqrt[a_.+b_.*x_^2]]^m_/Sqrt[a_.+b_.*x_^2],x_Symbol] :=
  -ArcCot[c*x/Sqrt[a+b*x^2]]^(m+1)/(c*(m+1)) /;
FreeQ[{a,b,c,m},x] && EqQ[b+c^2,0] && NeQ[m,-1]
```

$$2: \int \frac{\text{ArcTan}\left[\frac{c x}{\sqrt{a+b x^2}}\right]^m}{\sqrt{d+e x^2}} dx \text{ when } b+c^2=0 \wedge b d-a e=0$$

Derivation: Piecewise constant extraction

Basis: If  $b d - a e = 0$ , then  $\partial_x \frac{\sqrt{a+b x^2}}{\sqrt{d+e x^2}} = 0$

Rule: If  $b+c^2=0 \wedge b d - a e = 0$ , then

$$\int \frac{\text{ArcTan}\left[\frac{c x}{\sqrt{a+b x^2}}\right]^m}{\sqrt{d+e x^2}} dx \rightarrow \frac{\sqrt{a+b x^2}}{\sqrt{d+e x^2}} \int \frac{\text{ArcTan}\left[\frac{c x}{\sqrt{a+b x^2}}\right]^m}{\sqrt{a+b x^2}} dx$$

Program code:

```
Int[ArcTan[c_.x_/Sqrt[a_.+b_.x_^2]]^m_/Sqrt[d_.+e_.x_^2],x_Symbol] :=
  Sqrt[a+b*x^2]/Sqrt[d+e*x^2]*Int[ArcTan[c*x/Sqrt[a+b*x^2]]^m/Sqrt[a+b*x^2],x] /;
FreeQ[{a,b,c,d,e,m},x] && EqQ[b+c^2,0] && EqQ[b*d-a*e,0]
```

```
Int[ArcCot[c_.x_/Sqrt[a_.+b_.x_^2]]^m_/Sqrt[d_.+e_.x_^2],x_Symbol] :=
  Sqrt[a+b*x^2]/Sqrt[d+e*x^2]*Int[ArcCot[c*x/Sqrt[a+b*x^2]]^m/Sqrt[a+b*x^2],x] /;
FreeQ[{a,b,c,d,e,m},x] && EqQ[b+c^2,0] && EqQ[b*d-a*e,0]
```

$$5: \int u \text{ArcTan}\left[v+s\sqrt{v^2+1}\right] dx \text{ when } s^2=1$$

Derivation: Algebraic simplification

Basis: If  $s^2=1$ , then  $\text{ArcTan}\left[z+s\sqrt{z^2+1}\right] = \frac{\pi s}{4} + \frac{\text{ArcTan}[z]}{2}$

Basis: If  $s^2 = 1$ , then  $\text{ArcCot}\left[z + s\sqrt{z^2 + 1}\right] = \frac{\pi s}{4} - \frac{\text{ArcTan}[z]}{2}$

Rule: If  $s^2 = 1$ , then

$$\int u \text{ArcTan}\left[v + s\sqrt{v^2 + 1}\right] dx \rightarrow \frac{\pi s}{4} \int u dx + \frac{1}{2} \int u \text{ArcTan}[v] dx$$

Program code:

```
Int[u_.*ArcTan[v_+s_.*Sqrt[w_] ],x_Symbol] :=
  Pi*s/4*Int[u,x] + 1/2*Int[u*ArcTan[v],x] /;
EqQ[s^2,1] && EqQ[w,v^2+1]
```

```
Int[u_.*ArcCot[v_+s_.*Sqrt[w_] ],x_Symbol] :=
  Pi*s/4*Int[u,x] - 1/2*Int[u*ArcTan[v],x] /;
EqQ[s^2,1] && EqQ[w,v^2+1]
```

6:  $\int \frac{f[x, \text{ArcTan}[a x]]}{1 + (a + b x)^2} dx$

Derivation: Integration by substitution

Basis:  $\frac{f[z]}{1+z^2} == f[\text{Tan}[\text{ArcTan}[z]]] \text{ArcTan}'[z]$

Basis:  $r + s x + t x^2 == -\frac{s^2-4 r t}{4 t} \left(1 - \frac{(s+2 t x)^2}{s^2-4 r t}\right)$

Basis:  $1 + \text{Tan}[z]^2 == \text{Sec}[z]^2$

Rule:

$$\int \frac{f[x, \text{ArcTan}[a + b x]]}{1 + (a + b x)^2} dx \rightarrow \frac{1}{b} \text{Subst}\left[\int f\left[-\frac{a}{b} + \frac{\text{Tan}[x]}{b}, x\right] dx, x, \text{ArcTan}[a + b x]\right]$$

Program code:

```

If[TrueQ[$LoadShowSteps],

Int[u_*v_^n_,x_Symbol] :=
  With[{tmp=InverseFunctionOfLinear[u,x]},
    ShowStep["", "Int[f[x,ArcTan[a+b*x]]/(1+(a+b*x)^2),x]",
      "Subst[Int[f[-a/b+Tan[x]/b,x],x],x,ArcTan[a+b*x]]/b",Hold[
        (-Discriminant[v,x]/(4*Coefficient[v,x,2]))^n/Coefficient[tmp[[1]],x,1]*
        Subst[Int[SimplifyIntegrand[SubstForInverseFunction[u,tmp,x]*Sec[x]^(2*(n+1)),x],x], x, tmp]]] /;
    Not[FalseQ[tmp]] && EqQ[Head[tmp],ArcTan] && EqQ[Discriminant[v,x]*tmp[[1]]^2+D[v,x]^2,0]] /;
    SimplifyFlag && QuadraticQ[v,x] && ILtQ[n,0] && NegQ[Discriminant[v,x]] && MatchQ[u,r_.*f_^w_ /; FreeQ[f,x]],

Int[u_*v_^n_,x_Symbol] :=
  With[{tmp=InverseFunctionOfLinear[u,x]},
    (-Discriminant[v,x]/(4*Coefficient[v,x,2]))^n/Coefficient[tmp[[1]],x,1]*
    Subst[Int[SimplifyIntegrand[SubstForInverseFunction[u,tmp,x]*Sec[x]^(2*(n+1)),x],x], x, tmp] /;
    Not[FalseQ[tmp]] && EqQ[Head[tmp],ArcTan] && EqQ[Discriminant[v,x]*tmp[[1]]^2+D[v,x]^2,0]] /;
    QuadraticQ[v,x] && ILtQ[n,0] && NegQ[Discriminant[v,x]] && MatchQ[u,r_.*f_^w_ /; FreeQ[f,x]]]

```

```

If[TrueQ[$LoadShowSteps],

Int[u_*v_^n_,x_Symbol] :=
  With[{tmp=InverseFunctionOfLinear[u,x]},
    ShowStep["", "Int[f[x,ArcCot[a+b*x]]/(1+(a+b*x)^2),x]",
      "-Subst[Int[f[-a/b+Cot[x]/b,x],x],x,ArcCot[a+b*x]]/b",Hold[
        (-Discriminant[v,x]/(4*Coefficient[v,x,2]))^n/Coefficient[tmp[[1]],x,1]*
        Subst[Int[SimplifyIntegrand[SubstForInverseFunction[u,tmp,x]*Csc[x]^(2*(n+1)),x],x], x, tmp]]] /;
    Not[FalseQ[tmp]] && EqQ[Head[tmp],ArcCot] && EqQ[Discriminant[v,x]*tmp[[1]]^2+D[v,x]^2,0]] /;
    SimplifyFlag && QuadraticQ[v,x] && ILtQ[n,0] && NegQ[Discriminant[v,x]] && MatchQ[u,r_.*f_^w_ /; FreeQ[f,x]],

Int[u_*v_^n_,x_Symbol] :=
  With[{tmp=InverseFunctionOfLinear[u,x]},
    (-Discriminant[v,x]/(4*Coefficient[v,x,2]))^n/Coefficient[tmp[[1]],x,1]*
    Subst[Int[SimplifyIntegrand[SubstForInverseFunction[u,tmp,x]*Csc[x]^(2*(n+1)),x],x], x, tmp] /;
    Not[FalseQ[tmp]] && EqQ[Head[tmp],ArcCot] && EqQ[Discriminant[v,x]*tmp[[1]]^2+D[v,x]^2,0]] /;
    QuadraticQ[v,x] && ILtQ[n,0] && NegQ[Discriminant[v,x]] && MatchQ[u,r_.*f_^w_ /; FreeQ[f,x]]]

```

$$7. \int u \operatorname{ArcTan}[c + d \operatorname{Tan}[a + b x]] dx$$

$$1. \int \operatorname{ArcTan}[c + d \operatorname{Tan}[a + b x]] dx$$

$$\text{1: } \int \operatorname{ArcTan}[c + d \operatorname{Tan}[a + b x]] dx \text{ when } (c + i d)^2 = -1$$

Derivation: Integration by parts

$$\text{Basis: If } (c + i d)^2 = -1, \text{ then } \partial_x \operatorname{ArcTan}[c + d \operatorname{Tan}[a + b x]] = \frac{i b}{c + i d + c e^{2 i a + 2 i b x}}$$

Rule: If  $(c + i d)^2 = -1$ , then

$$\int \operatorname{ArcTan}[c + d \operatorname{Tan}[a + b x]] dx \rightarrow x \operatorname{ArcTan}[c + d \operatorname{Tan}[a + b x]] - i b \int \frac{x}{c + i d + c e^{2 i a + 2 i b x}} dx$$

Program code:

```
Int[ArcTan[c_.+d_.*Tan[a_.+b_.*x_]],x_Symbol] :=
  x*ArcTan[c+d*Tan[a+b*x]] -
  I*b*Int[x/(c+I*d+c*E^(2*I*a+2*I*b*x)),x] /;
FreeQ[{a,b,c,d},x] && EqQ[(c+I*d)^2,-1]
```

```
Int[ArcCot[c_.+d_.*Tan[a_.+b_.*x_]],x_Symbol] :=
  x*ArcCot[c+d*Tan[a+b*x]] +
  I*b*Int[x/(c+I*d+c*E^(2*I*a+2*I*b*x)),x] /;
FreeQ[{a,b,c,d},x] && EqQ[(c+I*d)^2,-1]
```

```
Int[ArcTan[c_.+d_.*Cot[a_.+b_.*x_]],x_Symbol] :=
  x*ArcTan[c+d*Cot[a+b*x]] -
  I*b*Int[x/(c-I*d-c*E^(2*I*a+2*I*b*x)),x] /;
FreeQ[{a,b,c,d},x] && EqQ[(c-I*d)^2,-1]
```

```

Int[ArcCot[c_.+d_.*Cot[a_.+b_.*x_]],x_Symbol] :=
  x*ArcCot[c+d*Cot[a+b*x]] +
  I*b*Int[x/(c-I*d-c*E^(2*I*a+2*I*b*x)),x] /;
FreeQ[{a,b,c,d},x] && EqQ[(c-I*d)^2,-1]

```

**2:**  $\int \text{ArcTan}[c + d \text{Tan}[a + b x]] dx$  when  $(c + i d)^2 \neq -1$

Derivation: Integration by parts

Basis:  $\partial_x \text{ArcTan}[c + d \text{Tan}[a + b x]] = \frac{b(1+i c+d) e^{2 i a+2 i b x}}{1+i c-d+(1+i c+d) e^{2 i a+2 i b x}} - \frac{b(1-i c-d) e^{2 i a+2 i b x}}{1-i c+d+(1-i c-d) e^{2 i a+2 i b x}}$

Rule: If  $(c + i d)^2 \neq -1$ , then

$$\int \text{ArcTan}[c + d \text{Tan}[a + b x]] dx \rightarrow x \text{ArcTan}[c + d \text{Tan}[a + b x]] - b(1+i c+d) \int \frac{x e^{2 i a+2 i b x}}{1+i c-d+(1+i c+d) e^{2 i a+2 i b x}} dx + b(1-i c-d) \int \frac{x e^{2 i a+2 i b x}}{1-i c+d+(1-i c-d) e^{2 i a+2 i b x}} dx$$

Program code:

```

Int[ArcTan[c_.+d_.*Tan[a_.+b_.*x_]],x_Symbol] :=
  x*ArcTan[c+d*Tan[a+b*x]] -
  b*(1+I*c+d)*Int[x*E^(2*I*a+2*I*b*x)/(1+I*c-d+(1+I*c+d)*E^(2*I*a+2*I*b*x)),x] +
  b*(1-I*c-d)*Int[x*E^(2*I*a+2*I*b*x)/(1-I*c+d+(1-I*c-d)*E^(2*I*a+2*I*b*x)),x] /;
FreeQ[{a,b,c,d},x] && NeQ[(c+I*d)^2,-1]

```

```

Int[ArcCot[c_.+d_.*Tan[a_.+b_.*x_]],x_Symbol] :=
  x*ArcCot[c+d*Tan[a+b*x]] +
  b*(1+I*c+d)*Int[x*E^(2*I*a+2*I*b*x)/(1+I*c-d+(1+I*c+d)*E^(2*I*a+2*I*b*x)),x] -
  b*(1-I*c-d)*Int[x*E^(2*I*a+2*I*b*x)/(1-I*c+d+(1-I*c-d)*E^(2*I*a+2*I*b*x)),x] /;
FreeQ[{a,b,c,d},x] && NeQ[(c+I*d)^2,-1]

```

```

Int[ArcTan[c_.+d_.*Cot[a_.+b_.*x_]],x_Symbol] :=
  x*ArcTan[c+d*Cot[a+b*x]] +
  b*(1+I*c-d)*Int[x*E^(2*I*a+2*I*b*x)/(1+I*c-d-(1+I*c-d)*E^(2*I*a+2*I*b*x)),x] -
  b*(1-I*c+d)*Int[x*E^(2*I*a+2*I*b*x)/(1-I*c-d-(1-I*c+d)*E^(2*I*a+2*I*b*x)),x] /;
FreeQ[{a,b,c,d},x] && NeQ[(c+I*d)^2,-1]

```

```

Int[ArcCot[c_.+d_.*Cot[a_.+b_.*x_]],x_Symbol] :=
  x*ArcCot[c+d*Cot[a+b*x]] -
  b*(1+I*c-d)*Int[x*E^(2*I*a+2*I*b*x)/(1+I*c-d-(1+I*c-d)*E^(2*I*a+2*I*b*x)),x] +
  b*(1-I*c+d)*Int[x*E^(2*I*a+2*I*b*x)/(1-I*c-d-(1-I*c+d)*E^(2*I*a+2*I*b*x)),x] /;
FreeQ[{a,b,c,d},x] && NeQ[(c-I*d)^2,-1]

```

2.  $\int (e + f x)^m \text{ArcTan}[c + d \text{Tan}[a + b x]] dx$  when  $m \in \mathbb{Z}^+$

**1:**  $\int (e + f x)^m \text{ArcTan}[c + d \text{Tan}[a + b x]] dx$  when  $m \in \mathbb{Z}^+ \wedge (c + i d)^2 = -1$

Derivation: Integration by parts

Basis: If  $(c + i d)^2 = -1$ , then  $\partial_x \text{ArcTan}[c + d \text{Tan}[a + b x]] = \frac{i b}{c + i d + c e^{2 i a + 2 i b x}}$

Rule: If  $m \in \mathbb{Z}^+ \wedge (c + i d)^2 = -1$ , then

$$\int (e + f x)^m \text{ArcTan}[c + d \text{Tan}[a + b x]] dx \rightarrow \frac{(e + f x)^{m+1} \text{ArcTan}[c + d \text{Tan}[a + b x]]}{f (m+1)} - \frac{i b}{f (m+1)} \int \frac{(e + f x)^{m+1}}{c + i d + c e^{2 i a + 2 i b x}} dx$$

Program code:

```

Int[(e_.+f_.*x_)^m_.*ArcTan[c_.+d_.*Tan[a_.+b_.*x_]],x_Symbol] :=
  (e+f*x)^(m+1)*ArcTan[c+d*Tan[a+b*x]]/(f*(m+1)) -
  I*b/(f*(m+1))*Int[(e+f*x)^(m+1)/(c+I*d+c*E^(2*I*a+2*I*b*x)),x] /;
FreeQ[{a,b,c,d,e,f},x] && IGtQ[m,0] && EqQ[(c+I*d)^2,-1]

```



```

Int[(e_.+f_.**x_)^m_.*ArcCot[c_.+d_.*Tan[a_.+b_.**x_]],x_Symbol] :=
  (e+f*x)^(m+1)*ArcCot[c+d*Tan[a+b*x]]/(f*(m+1)) +
  I*b/(f*(m+1))*Int[(e+f*x)^(m+1)/(c+I*d+c*E^(2*I*a+2*I*b*x)),x] /;
FreeQ[{a,b,c,d,e,f},x] && IGtQ[m,0] && EqQ[(c+I*d)^2,-1]

```

```

Int[(e_.+f_.**x_)^m_.*ArcTan[c_.+d_.*Cot[a_.+b_.**x_]],x_Symbol] :=
  (e+f*x)^(m+1)*ArcTan[c+d*Cot[a+b*x]]/(f*(m+1)) -
  I*b/(f*(m+1))*Int[(e+f*x)^(m+1)/(c-I*d-c*E^(2*I*a+2*I*b*x)),x] /;
FreeQ[{a,b,c,d,e,f},x] && IGtQ[m,0] && EqQ[(c-I*d)^2,-1]

```

```

Int[(e_.+f_.**x_)^m_.*ArcCot[c_.+d_.*Cot[a_.+b_.**x_]],x_Symbol] :=
  (e+f*x)^(m+1)*ArcCot[c+d*Cot[a+b*x]]/(f*(m+1)) +
  I*b/(f*(m+1))*Int[(e+f*x)^(m+1)/(c-I*d-c*E^(2*I*a+2*I*b*x)),x] /;
FreeQ[{a,b,c,d,e,f},x] && IGtQ[m,0] && EqQ[(c-I*d)^2,-1]

```

$$2: \int (e + f x)^m \operatorname{ArcTan}[c + d \operatorname{Tan}[a + b x]] dx \text{ when } m \in \mathbb{Z}^+ \wedge (c + i d)^2 \neq -1$$

## Derivation: Integration by parts

$$\text{Basis: } \partial_x \operatorname{ArcTan}[c + d \operatorname{Tan}[a + b x]] = \frac{b(1+i c+d) e^{2 i a+2 i b x}}{1+i c-d+(1+i c+d) e^{2 i a+2 i b x}} - \frac{b(1-i c-d) e^{2 i a+2 i b x}}{1-i c+d+(1-i c-d) e^{2 i a+2 i b x}}$$

Rule: If  $m \in \mathbb{Z}^+ \wedge (c + i d)^2 \neq -1$ , then

$$\int (e + f x)^m \operatorname{ArcTan}[c + d \operatorname{Tan}[a + b x]] dx \rightarrow \frac{(e + f x)^{m+1} \operatorname{ArcTan}[c + d \operatorname{Tan}[a + b x]]}{f(m+1)} - \frac{b(1+i c+d)}{f(m+1)} \int \frac{(e + f x)^{m+1} e^{2 i a+2 i b x}}{1+i c-d+(1+i c+d) e^{2 i a+2 i b x}} dx + \frac{b(1-i c-d)}{f(m+1)} \int \frac{(e + f x)^{m+1} e^{2 i a+2 i b x}}{1-i c+d+(1-i c-d) e^{2 i a+2 i b x}} dx$$

## Program code:

```
Int[(e_+f_.*x_)^m_.*ArcTan[c_+d_.*Tan[a_+b_.*x_]],x_Symbol] :=
  (e+f*x)^(m+1)*ArcTan[c+d*Tan[a+b*x]]/(f*(m+1)) -
  b*(1+I*c+d)/(f*(m+1))*Int[(e+f*x)^(m+1)*E^(2*I*a+2*I*b*x)/(1+I*c-d+(1+I*c+d)*E^(2*I*a+2*I*b*x)),x] +
  b*(1-I*c-d)/(f*(m+1))*Int[(e+f*x)^(m+1)*E^(2*I*a+2*I*b*x)/(1-I*c+d+(1-I*c-d)*E^(2*I*a+2*I*b*x)),x] /;
FreeQ[{a,b,c,d,e,f},x] && IGtQ[m,0] && NeQ[(c+I*d)^2,-1]
```

```
Int[(e_+f_.*x_)^m_.*ArcCot[c_+d_.*Tan[a_+b_.*x_]],x_Symbol] :=
  (e+f*x)^(m+1)*ArcCot[c+d*Tan[a+b*x]]/(f*(m+1)) +
  b*(1+I*c+d)/(f*(m+1))*Int[(e+f*x)^(m+1)*E^(2*I*a+2*I*b*x)/(1+I*c-d+(1+I*c+d)*E^(2*I*a+2*I*b*x)),x] -
  b*(1-I*c-d)/(f*(m+1))*Int[(e+f*x)^(m+1)*E^(2*I*a+2*I*b*x)/(1-I*c+d+(1-I*c-d)*E^(2*I*a+2*I*b*x)),x] /;
FreeQ[{a,b,c,d,e,f},x] && IGtQ[m,0] && NeQ[(c+I*d)^2,-1]
```

```
Int[(e_+f_.*x_)^m_.*ArcTan[c_+d_.*Cot[a_+b_.*x_]],x_Symbol] :=
  (e+f*x)^(m+1)*ArcTan[c+d*Cot[a+b*x]]/(f*(m+1)) +
  b*(1+I*c-d)/(f*(m+1))*Int[(e+f*x)^(m+1)*E^(2*I*a+2*I*b*x)/(1+I*c-d-(1+I*c-d)*E^(2*I*a+2*I*b*x)),x] -
  b*(1-I*c+d)/(f*(m+1))*Int[(e+f*x)^(m+1)*E^(2*I*a+2*I*b*x)/(1-I*c-d-(1-I*c+d)*E^(2*I*a+2*I*b*x)),x] /;
FreeQ[{a,b,c,d,e,f},x] && IGtQ[m,0] && NeQ[(c-I*d)^2,-1]
```

```

Int[(e_.+f_.**x_)^m_.*ArcCot[c_.+d_.*Cot[a_.+b_.**x_]],x_Symbol] :=
  (e+f*x)^(m+1)*ArcCot[c+d*Cot[a+b*x]]/(f*(m+1)) -
  b*(1+I*c-d)/(f*(m+1))*Int[(e+f*x)^(m+1)*E^(2*I*a+2*I*b*x)/(1+I*c-d-(1+I*c-d)*E^(2*I*a+2*I*b*x)),x] +
  b*(1-I*c+d)/(f*(m+1))*Int[(e+f*x)^(m+1)*E^(2*I*a+2*I*b*x)/(1-I*c-d-(1-I*c+d)*E^(2*I*a+2*I*b*x)),x] /;
FreeQ[{a,b,c,d,e,f},x] && IGtQ[m,0] && NeQ[(c-I*d)^2,-1]

```

$$8. \int u \operatorname{ArcTan}[c + d \operatorname{Tanh}[a + b x]] \, dx$$

$$1. \int u \operatorname{ArcTan}[\operatorname{Tanh}[a + b x]] \, dx$$

$$\mathbf{1:} \int \operatorname{ArcTan}[\operatorname{Tanh}[a + b x]] \, dx$$

Derivation: Integration by parts

Basis:  $\partial_x \operatorname{ArcTan}[\operatorname{Tanh}[a + b x]] = b \operatorname{Sech}[2 a + 2 b x]$

Rule:

$$\int \operatorname{ArcTan}[\operatorname{Tanh}[a + b x]] \, dx \rightarrow x \operatorname{ArcTan}[\operatorname{Tanh}[a + b x]] - b \int x \operatorname{Sech}[2 a + 2 b x] \, dx$$

Program code:

```

Int[ArcTan[Tanh[a_.+b_.**x_]],x_Symbol] :=
  x*ArcTan[Tanh[a+b*x]] - b*Int[x*Sech[2*a+2*b*x],x] /;
FreeQ[{a,b},x]

```

```

Int[ArcCot[Tanh[a_.+b_.**x_]],x_Symbol] :=
  x*ArcCot[Tanh[a+b*x]] + b*Int[x*Sech[2*a+2*b*x],x] /;
FreeQ[{a,b},x]

```

```

Int[ArcTan[Coth[a_.+b_.**x_]],x_Symbol] :=
  x*ArcTan[Coth[a+b*x]] + b*Int[x*Sech[2*a+2*b*x],x] /;
FreeQ[{a,b},x]

```

```
Int[ArcCot[Coth[a_.+b_.*x_]],x_Symbol] :=
  x*ArcCot[Coth[a+b*x]] - b*Int[x*Sech[2*a+2*b*x],x] /;
FreeQ[{a,b},x]
```

$$2: \int (e + f x)^m \operatorname{ArcTan}[\operatorname{Tanh}[a + b x]] \, dx \text{ when } m \in \mathbb{Z}^+$$

Derivation: Integration by parts

Basis:  $\partial_x \operatorname{ArcTan}[\operatorname{Tanh}[a + b x]] = b \operatorname{Sech}[2 a + 2 b x]$

Rule: If  $m \in \mathbb{Z}^+$ , then

$$\int (e + f x)^m \operatorname{ArcTan}[\operatorname{Tanh}[a + b x]] \, dx \rightarrow \frac{(e + f x)^{m+1} \operatorname{ArcTan}[\operatorname{Tanh}[a + b x]]}{f (m + 1)} - \frac{b}{f (m + 1)} \int (e + f x)^{m+1} \operatorname{Sech}[2 a + 2 b x] \, dx$$

Program code:

```
Int[(e_.+f_.*x_)^m_.*ArcTan[Tanh[a_.+b_.*x_]],x_Symbol] :=
  (e+f*x)^(m+1)*ArcTan[Tanh[a+b*x]]/(f*(m+1)) - b/(f*(m+1))*Int[(e+f*x)^(m+1)*Sech[2*a+2*b*x],x] /;
FreeQ[{a,b,e,f},x] && IGtQ[m,0]
```

```
Int[(e_.+f_.*x_)^m_.*ArcCot[Tanh[a_.+b_.*x_]],x_Symbol] :=
  (e+f*x)^(m+1)*ArcCot[Tanh[a+b*x]]/(f*(m+1)) + b/(f*(m+1))*Int[(e+f*x)^(m+1)*Sech[2*a+2*b*x],x] /;
FreeQ[{a,b,e,f},x] && IGtQ[m,0]
```

```
Int[(e_.+f_.*x_)^m_.*ArcTan[Coth[a_.+b_.*x_]],x_Symbol] :=
  (e+f*x)^(m+1)*ArcTan[Coth[a+b*x]]/(f*(m+1)) + b/(f*(m+1))*Int[(e+f*x)^(m+1)*Sech[2*a+2*b*x],x] /;
FreeQ[{a,b,e,f},x] && IGtQ[m,0]
```

```
Int[(e_.+f_.*x_)^m_.*ArcCot[Coth[a_.+b_.*x_]],x_Symbol] :=
  (e+f*x)^(m+1)*ArcCot[Coth[a+b*x]]/(f*(m+1)) - b/(f*(m+1))*Int[(e+f*x)^(m+1)*Sech[2*a+2*b*x],x] /;
FreeQ[{a,b,e,f},x] && IGtQ[m,0]
```

$$2. \int u \operatorname{ArcTan}[c + d \operatorname{Tanh}[a + b x]] \, dx$$

$$1. \int \operatorname{ArcTan}[c + d \operatorname{Tanh}[a + b x]] \, dx$$

$$\mathbf{1:} \int \operatorname{ArcTan}[c + d \operatorname{Tanh}[a + b x]] \, dx \text{ when } (c - d)^2 = -1$$

Derivation: Integration by parts

$$\text{Basis: If } (c - d)^2 = -1, \text{ then } \partial_x \operatorname{ArcTan}[c + d \operatorname{Tanh}[a + b x]] = \frac{b}{c - d + c e^{2a + 2bx}}$$

Rule: If  $(c - d)^2 = -1$ , then

$$\int \operatorname{ArcTan}[c + d \operatorname{Tanh}[a + b x]] \, dx \rightarrow x \operatorname{ArcTan}[c + d \operatorname{Tanh}[a + b x]] - b \int \frac{x}{c - d + c e^{2a + 2bx}} \, dx$$

Program code:

```
Int[ArcTan[c_.+d_.*Tanh[a_.+b_.*x_]],x_Symbol] :=
  x*ArcTan[c+d*Tanh[a+b*x]] -
  b*Int[x/(c-d+c*E^(2*a+2*b*x)),x] /;
FreeQ[{a,b,c,d},x] && EqQ[(c-d)^2,-1]
```

```
Int[ArcCot[c_.+d_.*Tanh[a_.+b_.*x_]],x_Symbol] :=
  x*ArcCot[c+d*Tanh[a+b*x]] +
  b*Int[x/(c-d+c*E^(2*a+2*b*x)),x] /;
FreeQ[{a,b,c,d},x] && EqQ[(c-d)^2,-1]
```

```
Int[ArcTan[c_.+d_.*Coth[a_.+b_.*x_]],x_Symbol] :=
  x*ArcTan[c+d*Coth[a+b*x]] -
  b*Int[x/(c-d-c*E^(2*a+2*b*x)),x] /;
FreeQ[{a,b,c,d},x] && EqQ[(c-d)^2,-1]
```

```

Int[ArcCot[c_.+d_.*Coth[a_.+b_.*x_]],x_Symbol] :=
  x*ArcCot[c+d*Coth[a+b*x]] +
  b*Int[x/(c-d-c*E^(2*a+2*b*x)),x] /;
FreeQ[{a,b,c,d},x] && EqQ[(c-d)^2,-1]

```

**2:**  $\int \text{ArcTan}[c + d \tanh[a + b x]] dx$  when  $(c - d)^2 \neq -1$

Derivation: Integration by parts

Basis:  $\partial_x \text{ArcTan}[c + d \tanh[a + b x]] = -\frac{i b (i - c - d) e^{2 a + 2 b x}}{i - c + d + (i - c - d) e^{2 a + 2 b x}} + \frac{i b (i + c + d) e^{2 a + 2 b x}}{i + c - d + (i + c + d) e^{2 a + 2 b x}}$

Rule: If  $(c - d)^2 \neq -1$ , then

$$\int \text{ArcTan}[c + d \tanh[a + b x]] dx \rightarrow x \text{ArcTan}[c + d \tanh[a + b x]] + i b (i - c - d) \int \frac{x e^{2 a + 2 b x}}{i - c + d + (i - c - d) e^{2 a + 2 b x}} dx - i b (i + c + d) \int \frac{x e^{2 a + 2 b x}}{i + c - d + (i + c + d) e^{2 a + 2 b x}} dx$$

Program code:

```

Int[ArcTan[c_.+d_.*Tanh[a_.+b_.*x_]],x_Symbol] :=
  x*ArcTan[c+d*Tanh[a+b*x]] +
  I*b*(I-c-d)*Int[x*E^(2*a+2*b*x)/(I-c+d+(I-c-d)*E^(2*a+2*b*x)),x] -
  I*b*(I+c+d)*Int[x*E^(2*a+2*b*x)/(I+c-d+(I+c+d)*E^(2*a+2*b*x)),x] /;
FreeQ[{a,b,c,d},x] && NeQ[(c-d)^2,-1]

```

```

Int[ArcCot[c_.+d_.*Tanh[a_.+b_.*x_]],x_Symbol] :=
  x*ArcCot[c+d*Tanh[a+b*x]] -
  I*b*(I-c-d)*Int[x*E^(2*a+2*b*x)/(I-c+d+(I-c-d)*E^(2*a+2*b*x)),x] +
  I*b*(I+c+d)*Int[x*E^(2*a+2*b*x)/(I+c-d+(I+c+d)*E^(2*a+2*b*x)),x] /;
FreeQ[{a,b,c,d},x] && NeQ[(c-d)^2,-1]

```

```

Int[ArcTan[c_.+d_.*Coth[a_.+b_.*x_]],x_Symbol] :=
  x*ArcTan[c+d*Coth[a+b*x]] -
  I*b*(I-c-d)*Int[x*E^(2*a+2*b*x)/(I-c-d-(I-c-d)*E^(2*a+2*b*x)),x] +
  I*b*(I+c+d)*Int[x*E^(2*a+2*b*x)/(I+c-d-(I+c+d)*E^(2*a+2*b*x)),x] /;
FreeQ[{a,b,c,d},x] && NeQ[(c-d)^2,-1]

```

```

Int[ArcCot[c_.+d_.*Coth[a_.+b_.*x_]],x_Symbol] :=
  x*ArcCot[c+d*Coth[a+b*x]] +
  I*b*(I-c-d)*Int[x*E^(2*a+2*b*x)/(I-c-d-(I-c-d)*E^(2*a+2*b*x)),x] -
  I*b*(I+c+d)*Int[x*E^(2*a+2*b*x)/(I+c-d-(I+c+d)*E^(2*a+2*b*x)),x] /;
FreeQ[{a,b,c,d},x] && NeQ[(c-d)^2,-1]

```

$$2. \int (e + f x)^m \operatorname{ArcTan}[c + d \operatorname{Tanh}[a + b x]] \, dx$$

$$1: \int (e + f x)^m \operatorname{ArcTan}[c + d \operatorname{Tanh}[a + b x]] \, dx \text{ when } m \in \mathbb{Z}^+ \wedge (c - d)^2 = -1$$

Derivation: Integration by parts

Basis: If  $(c - d)^2 = -1$ , then  $\partial_x \operatorname{ArcTan}[c + d \operatorname{Tanh}[a + b x]] = \frac{b}{c - d + c e^{2a+2bx}}$

Rule: If  $m \in \mathbb{Z}^+ \wedge (c - d)^2 = -1$ , then

$$\int (e + f x)^m \operatorname{ArcTan}[c + d \operatorname{Tanh}[a + b x]] \, dx \rightarrow \frac{(e + f x)^{m+1} \operatorname{ArcTan}[c + d \operatorname{Tanh}[a + b x]]}{f(m+1)} - \frac{b}{f(m+1)} \int \frac{(e + f x)^{m+1}}{c - d + c e^{2a+2bx}} \, dx$$

Program code:

```

Int[(e_.+f_.*x_)^m_.*ArcTan[c_.+d_.*Tanh[a_.+b_.*x_]],x_Symbol] :=
  (e+f*x)^(m+1)*ArcTan[c+d*Tanh[a+b*x]]/(f*(m+1)) -
  b/(f*(m+1))*Int[(e+f*x)^(m+1)/(c-d+c*E^(2*a+2*b*x)),x] /;
FreeQ[{a,b,c,d,e,f},x] && IGtQ[m,0] && EqQ[(c-d)^2,-1]

```

```

Int[(e_.+f_.**x_)^m_.*ArcCot[c_.+d_.*Tanh[a_.+b_.**x_]],x_Symbol] :=
  (e+f*x)^(m+1)*ArcCot[c+d*Tanh[a+b*x]]/(f*(m+1)) +
  b/(f*(m+1))*Int[(e+f*x)^(m+1)/(c-d+C*E^(2*a+2*b*x)),x] /;
FreeQ[{a,b,c,d,e,f},x] && IGtQ[m,0] && EqQ[(c-d)^2,-1]

```

```

Int[(e_.+f_.**x_)^m_.*ArcTan[c_.+d_.*Coth[a_.+b_.**x_]],x_Symbol] :=
  (e+f*x)^(m+1)*ArcTan[c+d*Coth[a+b*x]]/(f*(m+1)) -
  b/(f*(m+1))*Int[(e+f*x)^(m+1)/(c-d-C*E^(2*a+2*b*x)),x] /;
FreeQ[{a,b,c,d,e,f},x] && IGtQ[m,0] && EqQ[(c-d)^2,-1]

```

```

Int[(e_.+f_.**x_)^m_.*ArcCot[c_.+d_.*Coth[a_.+b_.**x_]],x_Symbol] :=
  (e+f*x)^(m+1)*ArcCot[c+d*Coth[a+b*x]]/(f*(m+1)) +
  b/(f*(m+1))*Int[(e+f*x)^(m+1)/(c-d-C*E^(2*a+2*b*x)),x] /;
FreeQ[{a,b,c,d,e,f},x] && IGtQ[m,0] && EqQ[(c-d)^2,-1]

```



$$\text{2: } \int (e + f x)^m \operatorname{ArcTan}[c + d \operatorname{Tanh}[a + b x]] dx \text{ when } m \in \mathbb{Z}^+ \wedge (c - d)^2 \neq -1$$

Derivation: Integration by parts

$$\text{Basis: } \partial_x \operatorname{ArcTan}[c + d \operatorname{Tanh}[a + b x]] = -\frac{i b (i - c - d) e^{2 a + 2 b x}}{i - c + d + (i - c - d) e^{2 a + 2 b x}} + \frac{i b (i + c + d) e^{2 a + 2 b x}}{i + c - d + (i + c + d) e^{2 a + 2 b x}}$$

Rule: If  $m \in \mathbb{Z}^+ \wedge (c - d)^2 \neq -1$ , then

$$\int (e + f x)^m \operatorname{ArcTan}[c + d \operatorname{Tanh}[a + b x]] dx \rightarrow \frac{(e + f x)^{m+1} \operatorname{ArcTan}[c + d \operatorname{Tanh}[a + b x]]}{f (m + 1)} + \frac{i b (i - c - d)}{f (m + 1)} \int \frac{(e + f x)^{m+1} e^{2 a + 2 b x}}{i - c + d + (i - c - d) e^{2 a + 2 b x}} dx - \frac{i b (i + c + d)}{f (m + 1)} \int \frac{(e + f x)^{m+1} e^{2 a + 2 b x}}{i + c - d + (i + c + d) e^{2 a + 2 b x}} dx$$

Program code:

```
Int[(e_.+f_.x_)^m_.*ArcTan[c_.+d_.Tanh[a_.+b_.x_]],x_Symbol] :=
  (e+f*x)^(m+1)*ArcTan[c+d*Tanh[a+b*x]]/(f*(m+1)) +
  I*b*(I-c-d)/(f*(m+1))*Int[(e+f*x)^(m+1)*E^(2*a+2*b*x)/(I-c+d+(I-c-d)*E^(2*a+2*b*x)),x] -
  I*b*(I+c+d)/(f*(m+1))*Int[(e+f*x)^(m+1)*E^(2*a+2*b*x)/(I+c-d+(I+c+d)*E^(2*a+2*b*x)),x] /;
FreeQ[{a,b,c,d,e,f},x] && IGtQ[m,0] && NeQ[(c-d)^2,-1]
```

```
Int[(e_.+f_.x_)^m_.*ArcCot[c_.+d_.Tanh[a_.+b_.x_]],x_Symbol] :=
  (e+f*x)^(m+1)*ArcCot[c+d*Tanh[a+b*x]]/(f*(m+1)) -
  I*b*(I-c-d)/(f*(m+1))*Int[(e+f*x)^(m+1)*E^(2*a+2*b*x)/(I-c+d+(I-c-d)*E^(2*a+2*b*x)),x] +
  I*b*(I+c+d)/(f*(m+1))*Int[(e+f*x)^(m+1)*E^(2*a+2*b*x)/(I+c-d+(I+c+d)*E^(2*a+2*b*x)),x] /;
FreeQ[{a,b,c,d,e,f},x] && IGtQ[m,0] && NeQ[(c-d)^2,-1]
```

```
Int[(e_.+f_.x_)^m_.*ArcTan[c_.+d_.Coth[a_.+b_.x_]],x_Symbol] :=
  (e+f*x)^(m+1)*ArcTan[c+d*Coth[a+b*x]]/(f*(m+1)) -
  I*b*(I-c-d)/(f*(m+1))*Int[(e+f*x)^(m+1)*E^(2*a+2*b*x)/(I-c-d-(I-c-d)*E^(2*a+2*b*x)),x] +
  I*b*(I+c+d)/(f*(m+1))*Int[(e+f*x)^(m+1)*E^(2*a+2*b*x)/(I+c-d-(I+c+d)*E^(2*a+2*b*x)),x] /;
FreeQ[{a,b,c,d,e,f},x] && IGtQ[m,0] && NeQ[(c-d)^2,-1]
```

```

Int[(e_.+f_.**x_)^m_.*ArcCot[c_.+d_.*Coth[a_.+b_.**x_]],x_Symbol] :=
  (e+f*x)^(m+1)*ArcCot[c+d*Coth[a+b*x]]/(f*(m+1)) +
  I*b*(I-c-d)/(f*(m+1))*Int[(e+f*x)^(m+1)*E^(2*a+2*b*x)/(I-c-d-(I-c-d)*E^(2*a+2*b*x)),x] -
  I*b*(I+c+d)/(f*(m+1))*Int[(e+f*x)^(m+1)*E^(2*a+2*b*x)/(I+c-d-(I+c-d)*E^(2*a+2*b*x)),x] /;
FreeQ[{a,b,c,d,e,f},x] && IGtQ[m,0] && NeQ[(c-d)^2,-1]

```

9.  $\int v (a + b \operatorname{ArcTan}[u]) \, dx$  when **u is free of inverse functions**

**1:**  $\int \operatorname{ArcTan}[u] \, dx$  when **u is free of inverse functions**

Derivation: Integration by parts

Rule: If u is free of inverse functions, then

$$\int \operatorname{ArcTan}[u] \, dx \rightarrow x \operatorname{ArcTan}[u] - \int \frac{x \partial_x u}{1+u^2} \, dx$$

Program code:

```

Int[ArcTan[u_],x_Symbol] :=
  x*ArcTan[u] -
  Int[SimplifyIntegrand[x*D[u,x]/(1+u^2),x],x] /;
InverseFunctionFreeQ[u,x]

```

```

Int[ArcCot[u_],x_Symbol] :=
  x*ArcCot[u] +
  Int[SimplifyIntegrand[x*D[u,x]/(1+u^2),x],x] /;
InverseFunctionFreeQ[u,x]

```

**2:**  $\int (c + dx)^m (a + b \operatorname{ArcTan}[u]) dx$  when  $m \neq -1 \wedge u$  is free of inverse functions

Derivation: Integration by parts

Rule: If  $m \neq -1 \wedge u$  is free of inverse functions, then

$$\int (c + dx)^m (a + b \operatorname{ArcTan}[u]) dx \rightarrow \frac{(c + dx)^{m+1} (a + b \operatorname{ArcTan}[u])}{d(m+1)} - \frac{b}{d(m+1)} \int \frac{(c + dx)^{m+1} \partial_x u}{1 + u^2} dx$$

Program code:

```
Int[(c_.+d_.*x_)^m_.*(a_.+b_.*ArcTan[u_]),x_Symbol] :=
  (c+d*x)^(m+1)*(a+b*ArcTan[u])/(d*(m+1)) -
  b/(d*(m+1))*Int[SimplifyIntegrand[(c+d*x)^(m+1)*D[u,x]/(1+u^2),x],x] /;
FreeQ[{a,b,c,d,m},x] && NeQ[m,-1] && InverseFunctionFreeQ[u,x] && Not[FunctionOfQ[(c+d*x)^(m+1),u,x]] && FalseQ[PowerVariableExpn[u,m+1]]
```

```
Int[(c_.+d_.*x_)^m_.*(a_.+b_.*ArcCot[u_]),x_Symbol] :=
  (c+d*x)^(m+1)*(a+b*ArcCot[u])/(d*(m+1)) +
  b/(d*(m+1))*Int[SimplifyIntegrand[(c+d*x)^(m+1)*D[u,x]/(1+u^2),x],x] /;
FreeQ[{a,b,c,d,m},x] && NeQ[m,-1] && InverseFunctionFreeQ[u,x] && Not[FunctionOfQ[(c+d*x)^(m+1),u,x]] && FalseQ[PowerVariableExpn[u,m+1]]
```

**3:**  $\int v (a + b \operatorname{ArcTan}[u]) \, dx$  when  $u$  and  $\int v \, dx$  are free of inverse functions

Derivation: Integration by parts

Rule: If  $u$  is free of inverse functions, let  $w = \int v \, dx$ , if  $w$  is free of inverse functions, then

$$\int v (a + b \operatorname{ArcTan}[u]) \, dx \rightarrow w (a + b \operatorname{ArcTan}[u]) - b \int \frac{w \partial_x u}{1 + u^2} \, dx$$

Program code:

```
Int[v_*(a_.+b_.*ArcTan[u_]),x_Symbol] :=
  With[{w=IntHide[v,x]},
    Dist[(a+b*ArcTan[u]),w,x] - b*Int[SimplifyIntegrand[w*D[u,x]/(1+u^2),x],x] /;
    InverseFunctionFreeQ[w,x] /;
    FreeQ[{a,b},x] && InverseFunctionFreeQ[u,x] && Not[MatchQ[v, (c_.+d_.*x)^m_. /; FreeQ[{c,d,m},x]]] && FalseQ[FunctionOfLinear[v*(a+b*A
```

```
Int[v_*(a_.+b_.*ArcCot[u_]),x_Symbol] :=
  With[{w=IntHide[v,x]},
    Dist[(a+b*ArcCot[u]),w,x] + b*Int[SimplifyIntegrand[w*D[u,x]/(1+u^2),x],x] /;
    InverseFunctionFreeQ[w,x] /;
    FreeQ[{a,b},x] && InverseFunctionFreeQ[u,x] && Not[MatchQ[v, (c_.+d_.*x)^m_. /; FreeQ[{c,d,m},x]]] && FalseQ[FunctionOfLinear[v*(a+b*A
```

**10:**  $\int \frac{\text{ArcTan}[v] \text{Log}[w]}{a + b x} dx$  when  $\partial_x \frac{v}{a + b x} = 0 \wedge \partial_x \frac{w}{a + b x} = 0$

Derivation: Algebraic expansion

Basis:  $\text{ArcTan}[z] = \frac{i}{2} \text{Log}[1 - i z] - \frac{i}{2} \text{Log}[1 + i z]$

Rule: If  $\partial_x \frac{v}{a + b x} = 0 \wedge \partial_x \frac{w}{a + b x} = 0$ , then

$$\int \frac{\text{ArcTan}[v] \text{Log}[w]}{a + b x} dx \rightarrow \frac{i}{2} \int \frac{\text{Log}[1 - i v] \text{Log}[w]}{a + b x} dx - \frac{i}{2} \int \frac{\text{Log}[1 + i v] \text{Log}[w]}{a + b x} dx$$

Program code:

```
Int[ArcTan[v_]*Log[w_]/(a_+b_*x_),x_Symbol] :=
  I/2*Int[Log[1-I*v]*Log[w]/(a+b*x),x] - I/2*Int[Log[1+I*v]*Log[w]/(a+b*x),x] /;
FreeQ[{a,b},x] && LinearQ[v,x] && LinearQ[w,x] && EqQ[Simplify[D[v/(a+b*x),x]],0] && EqQ[Simplify[D[w/(a+b*x),x]],0]
```

11.  $\int u \operatorname{ArcTan}[v] \operatorname{Log}[w] \, dx$  when  $v$ ,  $w$  and  $\int u \, dx$  are free of inverse functions

**1:**  $\int \operatorname{ArcTan}[v] \operatorname{Log}[w] \, dx$  when  $v$  and  $w$  are free of inverse functions

Derivation: Integration by parts

Rule: If  $v$  and  $w$  are free of inverse functions, then

$$\int \operatorname{ArcTan}[v] \operatorname{Log}[w] \, dx \rightarrow x \operatorname{ArcTan}[v] \operatorname{Log}[w] - \int \frac{x \operatorname{Log}[w] \partial_x v}{1+v^2} \, dx - \int \frac{x \operatorname{ArcTan}[v] \partial_x w}{w} \, dx$$

Program code:

```
Int[ArcTan[v_]*Log[w_],x_Symbol] :=
  x*ArcTan[v]*Log[w] -
  Int[SimplifyIntegrand[x*Log[w]*D[v,x]/(1+v^2),x],x] -
  Int[SimplifyIntegrand[x*ArcTan[v]*D[w,x]/w,x],x] /;
InverseFunctionFreeQ[v,x] && InverseFunctionFreeQ[w,x]
```

```
Int[ArcCot[v_]*Log[w_],x_Symbol] :=
  x*ArcCot[v]*Log[w] +
  Int[SimplifyIntegrand[x*Log[w]*D[v,x]/(1+v^2),x],x] -
  Int[SimplifyIntegrand[x*ArcCot[v]*D[w,x]/w,x],x] /;
InverseFunctionFreeQ[v,x] && InverseFunctionFreeQ[w,x]
```

**2:**  $\int u \operatorname{ArcTan}[v] \operatorname{Log}[w] \, dx$  when  $v$ ,  $w$  and  $\int u \, dx$  are free of inverse functions

Derivation: Integration by parts

Rule: If  $v$  and  $w$  are free of inverse functions, let  $z = \int u \, dx$ , if  $z$  is free of inverse functions, then

$$\int u \operatorname{ArcTan}[v] \operatorname{Log}[w] \, dx \rightarrow z \operatorname{ArcTan}[v] \operatorname{Log}[w] - \int \frac{z \operatorname{Log}[w] \partial_x v}{1 + v^2} \, dx - \int \frac{z \operatorname{ArcTan}[v] \partial_x w}{w} \, dx$$

Program code:

```
Int[u_*ArcTan[v]*Log[w],x_Symbol] :=
  With[{z=IntHide[u,x]},
    Dist[ArcTan[v]*Log[w],z,x] -
    Int[SimplifyIntegrand[z*Log[w]*D[v,x]/(1+v^2),x],x] -
    Int[SimplifyIntegrand[z*ArcTan[v]*D[w,x]/w,x],x] /;
    InverseFunctionFreeQ[z,x] /;
    InverseFunctionFreeQ[v,x] && InverseFunctionFreeQ[w,x]
```

```
Int[u_*ArcCot[v]*Log[w],x_Symbol] :=
  With[{z=IntHide[u,x]},
    Dist[ArcCot[v]*Log[w],z,x] +
    Int[SimplifyIntegrand[z*Log[w]*D[v,x]/(1+v^2),x],x] -
    Int[SimplifyIntegrand[z*ArcCot[v]*D[w,x]/w,x],x] /;
    InverseFunctionFreeQ[z,x] /;
    InverseFunctionFreeQ[v,x] && InverseFunctionFreeQ[w,x]
```