

## Rules for integrands of the form $(a + b x)^m (c + d x)^n$

**0:**  $\int (a + b x)^m (c + d x) dx$  when  $a d - b c (m + 2) == 0$

Derivation: Algebraic expansion

Basis: If  $a d - b c (m + 2) == 0$ , then  $c + d x == \frac{d (a + b (m + 2) x)}{b (m + 2)}$

Rule 1.1.1.2.0: If  $a d - b c (m + 2) == 0$ , then

$$\int (a + b x)^m (c + d x) dx \rightarrow \frac{d}{b (m + 2)} \int (a + b x)^m (a + b (m + 2) x) dx \rightarrow \frac{d x (a + b x)^{m+1}}{b (m + 2)}$$

```
Int[(a_+b_.*x_)^m_.*(c_+d_.*x_),x_Symbol] :=
  d*x*(a+b*x)^(m+1)/(b*(m+2)) /;
FreeQ[{a,b,c,d,m},x] && EqQ[a*d-b*c*(m+2),0]
```

$$1. \int (a+bx)^m (c+dx)^n dx \text{ when } bc - ad \neq 0 \wedge m+n+2 \neq 0$$

$$1. \int \frac{1}{(a+bx)(c+dx)} dx \text{ when } bc - ad \neq 0$$

$$\textcolor{red}{1}: \int \frac{1}{(a+bx)(c+dx)} dx \text{ when } bc + ad = 0$$

Derivation: Algebraic simplification

Basis: If  $bc + ad = 0$ , then  $(a+bx)(c+dx) = ac + bdx^2$

Rule 1.1.1.2.1.1.1: If  $bc + ad = 0$ , then

$$\int \frac{1}{(a+bx)(c+dx)} dx \rightarrow \int \frac{1}{ac + bdx^2} dx$$

Program code:

```
Int[1/((a_+b_.*x_)*(c_+d_.*x_)),x_Symbol] :=
  Int[1/(a*c+b*d*x^2),x] /;
FreeQ[{a,b,c,d},x] && EqQ[b*c+a*d,0]
```

**2:**  $\int \frac{1}{(a+bx)(c+dx)} dx$  when  $bc - ad \neq 0$

Derivation: Algebraic expansion

Basis:  $\frac{1}{(a+bx)(c+dx)} = \frac{b}{(bc-ad)(a+bx)} - \frac{d}{(bc-ad)(c+dx)}$

Rule 1.1.1.2.1.1.2: If  $bc - ad \neq 0$ , then

$$\int \frac{1}{(a+bx)(c+dx)} dx \rightarrow \frac{b}{bc-ad} \int \frac{1}{a+bx} dx - \frac{d}{bc-ad} \int \frac{1}{c+dx} dx$$

Program code:

```
Int[1/((a_+b_.*x_)*(c_+d_.*x_)),x_Symbol] :=
  b/(b*c-a*d)*Int[1/(a+b*x),x] - d/(b*c-a*d)*Int[1/(c+d*x),x] /;
FreeQ[{a,b,c,d},x] && NeQ[b*c-a*d,0]
```

**2:**  $\int (a+bx)^m (c+dx)^n dx$  when  $bc - ad \neq 0 \wedge m+n+2 \neq 0 \wedge m \neq -1$

Reference: G&R 2.155, CRC 59a with  $m+n+2 \neq 0$

Reference: G&R 2.110.2 or 2.110.6 with  $k = 1$  and  $m+n+2 \neq 0$

Derivation: Linear recurrence 3 with  $m+n+2 \neq 0$

Rule 1.1.1.2.1.2: If  $bc - ad \neq 0 \wedge m+n+2 \neq 0 \wedge m \neq -1$ , then

$$\int (a+bx)^m (c+dx)^n dx \rightarrow \frac{(a+bx)^{m+1} (c+dx)^{n+1}}{(bc-ad)(m+1)}$$

Program code:

```
Int[(a_.+b_.*x_)^m_.*(c_.+d_.*x_)^n_,x_Symbol] :=
  (a+b*x)^(m+1)*(c+d*x)^(n+1)/((b*c-a*d)*(m+1)) /;
FreeQ[{a,b,c,d,m,n},x] && NeQ[b*c-a*d,0] && EqQ[m+n+2,0] && NeQ[m,-1]
```

2.  $\int (a+bx)^m (c+dx)^n dx$  when  $bc+ad = 0 \wedge n = m$

1:  $\int (a+bx)^m (c+dx)^m dx$  when  $bc+ad = 0 \wedge m + \frac{1}{2} \in \mathbb{Z}^+$

Derivation: Inverted integration by parts

Rule 1.1.1.2.2.1: If  $bc+ad = 0 \wedge m + \frac{1}{2} \in \mathbb{Z}^+$ , then

$$\int (a+bx)^m (c+dx)^m dx \rightarrow \frac{x(a+bx)^m (c+dx)^m}{2m+1} + \frac{2acm}{2m+1} \int (a+bx)^{m-1} (c+dx)^{m-1} dx$$

Program code:

```
Int[(a+b_.**x_)^m_*(c+d_.**x_)^m_,x_Symbol] :=
  x*(a+b*x)^m*(c+d*x)^m/(2*m+1) + 2*a*c*m/(2*m+1)*Int[(a+b*x)^(m-1)*(c+d*x)^(m-1),x] /;
FreeQ[{a,b,c,d},x] && EqQ[b*c+a*d,0] && IGtQ[m+1/2,0]
```

2.  $\int (a+bx)^m (c+dx)^m dx$  when  $bc+ad = 0 \wedge m + \frac{1}{2} \in \mathbb{Z}^-$

1:  $\int \frac{1}{(a+bx)^{3/2} (c+dx)^{3/2}} dx$  when  $bc+ad = 0$

Rule 1.1.1.2.2.2.1: If  $bc+ad = 0$ , then

$$\int \frac{1}{(a+bx)^{3/2} (c+dx)^{3/2}} dx \rightarrow \frac{x}{ac \sqrt{a+bx} \sqrt{c+dx}}$$

Program code:

```
Int[1/((a+b_.**x_)^(3/2)*(c+d_.**x_)^(3/2)),x_Symbol] :=
  x/(a*c*Sqrt[a+b*x]*Sqrt[c+d*x]) /;
FreeQ[{a,b,c,d},x] && EqQ[b*c+a*d,0]
```

**2:**  $\int (a+bx)^m (c+dx)^m dx$  when  $bc+ad = 0 \wedge m + \frac{3}{2} \in \mathbb{Z}^-$

Derivation: Integration by parts

Basis:  $(a+bx)^m (c+dx)^m = x^{2(m+1)+1} \frac{(a+bx)^m (c+dx)^m}{x^{2(m+1)+1}}$

Basis: If  $bc+ad = 0$ , then  $\int \frac{(a+bx)^m (c+dx)^m}{x^{2(m+1)+1}} dx = -\frac{(a+bx)^{m+1} (c+dx)^{m+1}}{x^{2(m+1)} 2ac(m+1)}$

Rule 1.1.1.2.2.2.2: If  $bc+ad = 0 \wedge m + \frac{3}{2} \in \mathbb{Z}^-$ , then

$$\int (a+bx)^m (c+dx)^m dx \rightarrow -\frac{x(a+bx)^{m+1} (c+dx)^{m+1}}{2ac(m+1)} + \frac{2m+3}{2ac(m+1)} \int (a+bx)^{m+1} (c+dx)^{m+1} dx$$

Program code:

```
Int[(a+b*x)^m*(c+d*x)^m,x_Symbol] :=
  -x*(a+b*x)^(m+1)*(c+d*x)^(m+1)/(2*a*c*(m+1)) +
  (2*m+3)/(2*a*c*(m+1))*Int[(a+b*x)^(m+1)*(c+d*x)^(m+1),x] /;
FreeQ[{a,b,c,d},x] && EqQ[b*c+a*d,0] && ILtQ[m+3/2,0]
```

**3:**  $\int (a+bx)^m (c+dx)^m dx$  when  $b c + a d = 0 \wedge (m \in \mathbb{Z} \vee a > 0 \wedge c > 0)$

Derivation: Algebraic simplification

Basis: If  $b c + a d = 0 \wedge (m \in \mathbb{Z} \vee a > 0 \wedge c > 0)$ , then  $(a+bx)^m (c+dx)^m = (a c + b d x^2)^m$

Rule 1.1.1.2.2.3: If  $b c + a d = 0 \wedge (m \in \mathbb{Z} \vee a > 0 \wedge c > 0)$ , then

$$\int (a+bx)^m (c+dx)^m dx \rightarrow \int (a c + b d x^2)^m dx$$

Program code:

```
Int[(a_+b_.*x_)^m_.*(c_+d_.*x_)^m_,x_Symbol] :=
  Int[(a*c+b*d*x^2)^m,x] /;
FreeQ[{a,b,c,d,m},x] && EqQ[b*c+a*d,0] && (IntegerQ[m] || GtQ[a,0] && GtQ[c,0])
```

**4:**  $\int (a+bx)^m (c+dx)^m dx$  when  $bc+ad = 0 \wedge 2m \notin \mathbb{Z}$

Derivation: Piecewise constant extraction

Basis: If  $bc+ad = 0$ , then  $\partial_x \frac{(a+bx)^m (c+dx)^m}{(a+bdx^2)^m} = 0$

Basis: If  $bc+ad = 0$ , then  $\frac{(a+bx)^m (c+dx)^m}{(a+bdx^2)^m} = \frac{(a+bx)^{\text{FracPart}[m]} (c+dx)^{\text{FracPart}[m]}}{(a+bdx^2)^{\text{FracPart}[m]}}$

Rule 1.1.1.2.2.4: If  $bc+ad = 0 \wedge 2m \notin \mathbb{Z}$ , then

$$\int (a+bx)^m (c+dx)^m dx \rightarrow \frac{(a+bx)^{\text{FracPart}[m]} (c+dx)^{\text{FracPart}[m]}}{(a+bdx^2)^{\text{FracPart}[m]}} \int (a+bdx^2)^m dx$$

Program code:

```
Int[(a+b_.**x_)^m_*(c+d_.**x_)^m_,x_Symbol] :=
  (a+b*x)^FracPart[m]*(c+d*x)^FracPart[m]/(a*c+b*d*x^2)^FracPart[m]*Int[(a*c+b*d*x^2)^m,x] /;
FreeQ[{a,b,c,d,m},x] && EqQ[b*c+a*d,0] && Not[IntegerQ[2*m]]
```



**3:**  $\int (a+bx)^m (c+dx)^n dx$  when  $b c - a d \neq 0 \wedge m \in \mathbb{Z}$

Derivation: Algebraic expansion

Rule 1.1.1.2.3: If  $b c - a d \neq 0 \wedge m \in \mathbb{Z}$ , then

$$\int (a+bx)^m (c+dx)^n dx \rightarrow \int \text{ExpandIntegrand}[(a+bx)^m (c+dx)^n, x] dx$$

Program code:

```
Int[(a_+b_.*x_)^m_.*(c_+d_.*x_)^n_,x_Symbol] :=
  Int[ExpandIntegrand[(a+b*x)^m*(c+d*x)^n,x],x] /;
FreeQ[{a,b,c,d,n},x] && NeQ[b*c-a*d,0] && IGtQ[m,0] &&
  (Not[IntegerQ[n]] || EqQ[c,0] && LeQ[7*m+4*n+4,0] || LtQ[9*m+5*(n+1),0] || GtQ[m+n+2,0])
```

```
Int[(a_+b_.*x_)^m_.*(c_+d_.*x_)^n_,x_Symbol] :=
  Int[ExpandIntegrand[(a+b*x)^m*(c+d*x)^n,x],x] /;
FreeQ[{a,b,c,d},x] && NeQ[b*c-a*d,0] && ILtQ[m,0] && IntegerQ[n] && Not[IGtQ[n,0] && LtQ[m+n+2,0]]
```

**4:**  $\int (a+bx)^m (c+dx)^n dx$  when  $bc - ad \neq 0 \wedge m+n+2 \in \mathbb{Z}^- \wedge m \neq -1$

Reference: G&R 2.155, CRC 59a

Reference: G&R 2.110.2 or 2.110.6 with  $k = 1$

Derivation: Linear recurrence 3

Derivation: Integration by parts

Basis:  $(a+bx)^m (c+dx)^n = (c+dx)^{m+n+2} \frac{(a+bx)^m}{(c+dx)^{m+2}}$

Rule 1.1.1.2.4: If  $bc - ad \neq 0 \wedge m+n+2 \in \mathbb{Z}^- \wedge m \neq -1$ , then

$$\int (a+bx)^m (c+dx)^n dx \rightarrow \frac{(a+bx)^{m+1} (c+dx)^{n+1}}{(bc-ad)(m+1)} - \frac{d(m+n+2)}{(bc-ad)(m+1)} \int (a+bx)^{m+1} (c+dx)^n dx$$

Program code:

```
Int[(a_.+b_.x_)^m_*(c_.+d_.x_)^n_,x_Symbol] :=
  (a+b*x)^(m+1)*(c+d*x)^(n+1)/((b*c-a*d)*(m+1)) -
  d*Simplify[m+n+2]/((b*c-a*d)*(m+1))*Int[(a+b*x)^Simplify[m+1]*(c+d*x)^n,x] /;
FreeQ[{a,b,c,d,m,n},x] && NeQ[b*c-a*d,0] && ILtQ[Simplify[m+n+2],0] && NeQ[m,-1] &&
  Not[LtQ[m,-1] && LtQ[n,-1] && (EqQ[a,0] || NeQ[c,0] && LtQ[m-n,0] && IntegerQ[n])] &&
  (SumSimplerQ[m,1] || Not[SumSimplerQ[n,1]])
```

5.  $\int (a+bx)^m (c+dx)^n dx$  when  $bc - ad \neq 0 \wedge n > 0$

1:  $\int (a+bx)^m (c+dx)^n dx$  when  $bc - ad \neq 0 \wedge n > 0 \wedge m < -1$

Reference: G&R 2.110.3 or 2.110.4 with  $k = 1$

Derivation: Linear recurrence 1

Note: If  $n \in \mathbb{Z}$  and  $m \notin \mathbb{Z}$ , there is no need to drive  $m$  toward 0 along with  $n$ .

Rule 1.1.1.2.5.1: If  $bc - ad \neq 0 \wedge n > 0 \wedge m < -1$ , then

$$\int (a+bx)^m (c+dx)^n dx \rightarrow \frac{(a+bx)^{m+1} (c+dx)^n}{b(m+1)} - \frac{dn}{b(m+1)} \int (a+bx)^{m+1} (c+dx)^{n-1} dx$$

Program code:

```
Int[1/((a_+b_.*x_)^(9/4)*(c_+d_.*x_)^(1/4)),x_Symbol] :=
  -4/(5*b*(a+b*x)^(5/4)*(c+d*x)^(1/4)) - d/(5*b)*Int[1/((a+b*x)^(5/4)*(c+d*x)^(5/4)),x] /;
FreeQ[{a,b,c,d},x] && EqQ[b*c+a*d,0] && NegQ[a^2*b^2]
```

```
Int[(a_+b_.*x_)^m_*(c_+d_.*x_)^n_,x_Symbol] :=
  (a+b*x)^(m+1)*(c+d*x)^n/(b*(m+1)) -
  d*n/(b*(m+1))*Int[(a+b*x)^(m+1)*(c+d*x)^(n-1),x] /;
FreeQ[{a,b,c,d},x] && NeQ[b*c-a*d,0] && GtQ[n,0] && LtQ[m,-1] && Not[IntegerQ[n] && Not[IntegerQ[m]]] &&
  Not[IntegerQ[m+n+2,0] && (FractionQ[m] || GeQ[2*n+m+1,0])] && IntLinearQ[a,b,c,d,m,n,x]
```

**2:**  $\int (a+bx)^m (c+dx)^n dx$  when  $bc - ad \neq 0 \wedge n > 0 \wedge m+n+1 \neq 0$

Reference: G&R 2.151, CRC 59b

Reference: G&R 2.110.1 or 2.110.5 with  $k = 1$

Derivation: Linear recurrence 2

Derivation: Inverted integration by parts

Rule 1.1.1.2.5.2: If  $bc - ad \neq 0 \wedge n > 0 \wedge m+n+1 \neq 0$ , then

$$\int (a+bx)^m (c+dx)^n dx \rightarrow \frac{(a+bx)^{m+1} (c+dx)^n}{b(m+n+1)} + \frac{n(bc-ad)}{b(m+n+1)} \int (a+bx)^m (c+dx)^{n-1} dx$$

Program code:

```
Int[1/((a+b_.x_)^(5/4)*(c+d_.x_)^(1/4)),x_Symbol] :=
  -2/(b*(a+b*x)^(1/4)*(c+d*x)^(1/4)) + c*Int[1/((a+b*x)^(5/4)*(c+d*x)^(5/4)),x] /;
FreeQ[{a,b,c,d},x] && EqQ[b*c+a*d,0] && NegQ[a^2*b^2]
```

```
Int[(a+b_.x_)^m_*(c+d_.x_)^n_,x_Symbol] :=
  (a+b*x)^(m+1)*(c+d*x)^n/(b*(m+n+1)) +
  2*c*n/(m+n+1)*Int[(a+b*x)^m*(c+d*x)^(n-1),x] /;
FreeQ[{a,b,c,d},x] && EqQ[b*c+a*d,0] && IGtQ[m+1/2,0] && IGtQ[n+1/2,0] && LtQ[m,n]
```

```
Int[(a_.+b_.x_)^m_*(c_.+d_.x_)^n_,x_Symbol] :=
  (a+b*x)^(m+1)*(c+d*x)^n/(b*(m+n+1)) +
  n*(b*c-a*d)/(b*(m+n+1))*Int[(a+b*x)^m*(c+d*x)^(n-1),x] /;
FreeQ[{a,b,c,d},x] && NeQ[b*c-a*d,0] && GtQ[n,0] && NeQ[m+n+1,0] &&
  Not[IGtQ[m,0] && (Not[IntegerQ[n]] || GtQ[m,0] && LtQ[m-n,0])] &&
  Not[ILtQ[m+n+2,0]] && IntLinearQ[a,b,c,d,m,n,x]
```

**6:**  $\int (a+bx)^m (c+dx)^n dx$  when  $bc - ad \neq 0 \wedge m < -1$

Reference: G&R 2.155, CRC 59a

Reference: G&R 2.110.2 or 2.110.6 with  $k = 1$

Derivation: Linear recurrence 3

Derivation: Integration by parts

Basis:  $(a+bx)^m (c+dx)^n = (c+dx)^{m+n+2} \frac{(a+bx)^m}{(c+dx)^{m+2}}$

Rule 1.1.1.2.6: If  $bc - ad \neq 0 \wedge m < -1$ , then

$$\int (a+bx)^m (c+dx)^n dx \rightarrow \frac{(a+bx)^{m+1} (c+dx)^{n+1}}{(bc-ad)(m+1)} - \frac{d(m+n+2)}{(bc-ad)(m+1)} \int (a+bx)^{m+1} (c+dx)^n dx$$

Program code:

```
Int[(a_.+b_.*x_)^m_*(c_.+d_.*x_)^n_,x_Symbol] :=
  (a+b*x)^(m+1)*(c+d*x)^(n+1)/((b*c-a*d)*(m+1)) -
  d*(m+n+2)/((b*c-a*d)*(m+1))*Int[(a+b*x)^(m+1)*(c+d*x)^n,x] /;
FreeQ[{a,b,c,d,n},x] && NeQ[b*c-a*d,0] && LtQ[m,-1] &&
  Not[LtQ[n,-1] && (EqQ[a,0] || NeQ[c,0] && LtQ[m-n,0] && IntegerQ[n])] && IntLinearQ[a,b,c,d,m,n,x]
```

7.  $\int (a+bx)^m (c+dx)^n dx$  when  $bc - ad \neq 0 \wedge -1 \leq m < 0 \wedge -1 < n < 0$

1.  $\int \frac{1}{\sqrt{a+bx} \sqrt{c+dx}} dx$  when  $bc - ad \neq 0$

**1:**  $\int \frac{1}{\sqrt{a+bx} \sqrt{c+dx}} dx$  when  $a+c = 0 \wedge b-d = 0 \wedge a > 0$

Rule 1.1.1.2.7.1.1: If  $a+c = 0 \wedge b-d = 0 \wedge a > 0$ , then

$$\int \frac{1}{\sqrt{a+bx} \sqrt{c+dx}} dx \rightarrow \frac{1}{b} \operatorname{ArcCosh}\left[\frac{bx}{a}\right]$$

Program code:

```
Int[1/(Sqrt[a+b_.**x_]*Sqrt[c+d_.**x_]),x_Symbol] :=
  ArcCosh[b*x/a]/b /;
FreeQ[{a,b,c,d},x] && EqQ[a+c,0] && EqQ[b-d,0] && GtQ[a,0]
```

**2:**  $\int \frac{1}{\sqrt{a+bx} \sqrt{c+dx}} dx$  when  $b+d=0 \wedge a+c>0$

Derivation: Algebraic simplification

Basis: If  $a+c>0$ , then  $(a+bx)^m (c-bx)^m = ((a+bx)(c-bx))^m = (ac-b(a-c)x-b^2x^2)^m$

Rule 1.1.1.2.7.1.2: If  $b+d=0 \wedge a+c>0$ , then

$$\int \frac{1}{\sqrt{a+bx} \sqrt{c+dx}} dx \rightarrow \int \frac{1}{\sqrt{ac-b(a-c)x-b^2x^2}} dx$$

Program code:

```
Int[1/(Sqrt[a_+b_.*x_]*Sqrt[c_+d_.*x_]),x_Symbol] :=
  Int[1/Sqrt[a*c-b*(a-c)*x-b^2*x^2],x] /;
FreeQ[{a,b,c,d},x] && EqQ[b+d,0] && GtQ[a+c,0]
```

$$\mathbf{3:} \int \frac{1}{\sqrt{a+bx} \sqrt{c+dx}} dx \text{ when } bc - ad > 0 \wedge b > 0$$

Derivation: Integration by substitution

$$\text{Basis: If } b > 0, \text{ then } \frac{1}{\sqrt{a+bx} \sqrt{c+dx}} = \frac{2}{\sqrt{b}} \text{Subst} \left[ \frac{1}{\sqrt{bc-ad+dx^2}}, x, \sqrt{a+bx} \right] \partial_x \sqrt{a+bx}$$

Rule 1.1.1.2.7.1.3: If  $bc - ad > 0 \wedge b > 0$ , then

$$\int \frac{1}{\sqrt{a+bx} \sqrt{c+dx}} dx \rightarrow \frac{2}{\sqrt{b}} \text{Subst} \left[ \int \frac{1}{\sqrt{bc-ad+dx^2}} dx, x, \sqrt{a+bx} \right]$$

Program code:

```
Int[1/(Sqrt[a_+b_.*x_]*Sqrt[c_+d_.*x_]),x_Symbol] :=
  2/Sqrt[b]*Subst[Int[1/Sqrt[b*c-a*d+d*x^2],x],x,Sqrt[a+b*x]] /;
FreeQ[{a,b,c,d},x] && GtQ[b*c-a*d,0] && GtQ[b,0]
```

$$2. \int \frac{1}{(a+bx)(c+dx)^{1/3}} dx \text{ when } bc - ad \neq 0$$

$$\mathbf{1:} \int \frac{1}{(a+bx)(c+dx)^{1/3}} dx \text{ when } \frac{bc-ad}{b} > 0$$

Derivation: Integration by substitution

$$\text{Basis: Let } q = \left( \frac{bc-ad}{b} \right)^{1/3}, \text{ then } \frac{1}{(a+bx)(c+dx)^{1/3}} = -\frac{1}{2q(a+bx)} - \text{Subst} \left[ \frac{3}{2bq(q-x)} - \frac{3}{2b(q^2+qx+x^2)}, x, (c+dx)^{1/3} \right] \partial_x (c+dx)^{1/3}$$

Rule 1.1.1.2.7.2.1: If  $\frac{bc-ad}{b} > 0$ , let  $q = \left( \frac{bc-ad}{b} \right)^{1/3}$ , then

$$\int \frac{1}{(a+bx)(c+dx)^{1/3}} dx \rightarrow$$



$$-\frac{\text{Log}[a+bx]}{2bq} - \frac{3}{2bq} \text{Subst}\left[\int \frac{1}{q-x} dx, x, (c+dx)^{1/3}\right] + \frac{3}{2b} \text{Subst}\left[\int \frac{1}{q^2+qx+x^2} dx, x, (c+dx)^{1/3}\right]$$

Program code:

```
Int[1/((a_+b_.*x_)*(c_+d_.*x_)^(1/3)),x_Symbol] :=
  With[{q=Rt[(b*c-a*d)/b,3]},
    -Log[RemoveContent[a+b*x,x]]/(2*b*q) -
    3/(2*b*q)*Subst[Int[1/(q-x),x],x,(c+d*x)^(1/3)] +
    3/(2*b)*Subst[Int[1/(q^2+q*x+x^2),x],x,(c+d*x)^(1/3)]] /;
  FreeQ[{a,b,c,d},x] && PosQ[(b*c-a*d)/b]
```

$$2: \int \frac{1}{(a+bx)(c+dx)^{1/3}} dx \text{ when } \frac{bc-ad}{b} \neq 0$$

Derivation: Integration by substitution

Basis: Let  $q = \left(-\frac{bc-ad}{b}\right)^{1/3}$ , then  $\frac{1}{(a+bx)(c+dx)^{1/3}} = \frac{1}{2q(a+bx)} - \text{Subst}\left[\frac{3}{2bq(q+x)} - \frac{3}{2b(q^2-qx+x^2)}, x, (c+dx)^{1/3}\right] \partial_x (c+dx)^{1/3}$

Rule 1.1.1.2.7.2.2: If  $\frac{bc-ad}{b} \neq 0$ , let  $q = \left(-\frac{bc-ad}{b}\right)^{1/3}$ , then

$$\int \frac{1}{(a+bx)(c+dx)^{1/3}} dx \rightarrow$$

$$\frac{\text{Log}[a+bx]}{2bq} - \frac{3}{2bq} \text{Subst}\left[\int \frac{1}{q+x} dx, x, (c+dx)^{1/3}\right] + \frac{3}{2b} \text{Subst}\left[\int \frac{1}{q^2-qx+x^2} dx, x, (c+dx)^{1/3}\right]$$

Program code:

```
Int[1/((a_+b_.*x_)*(c_+d_.*x_)^(1/3)),x_Symbol] :=
  With[{q=Rt[-(b*c-a*d)/b,3]},
    Log[RemoveContent[a+b*x,x]]/(2*b*q) -
    3/(2*b*q)*Subst[Int[1/(q+x),x],x,(c+d*x)^(1/3)] +
    3/(2*b)*Subst[Int[1/(q^2-q*x+x^2),x],x,(c+d*x)^(1/3)]] /;
  FreeQ[{a,b,c,d},x] && NegQ[(b*c-a*d)/b]
```

$$3. \int \frac{1}{(a+bx)(c+dx)^{2/3}} dx \text{ when } b c - a d \neq 0$$

$$1: \int \frac{1}{(a+bx)(c+dx)^{2/3}} dx \text{ when } \frac{bc-ad}{b} > 0$$

Derivation: Integration by substitution

Basis: Let  $q = \left(\frac{bc-ad}{b}\right)^{1/3}$ , then  $\frac{1}{(a+bx)(c+dx)^{2/3}} = -\frac{1}{2q^2(a+bx)} - \text{Subst}\left[\frac{3}{2bq^2(q-x)} + \frac{3}{2bq(q^2+qx+x^2)}, x, (c+dx)^{1/3}\right] \partial_x (c+dx)^{1/3}$

Rule 1.1.1.2.7.3.1: If  $\frac{bc-ad}{b} > 0$ , let  $q = \left(\frac{bc-ad}{b}\right)^{1/3}$ , then

$$\int \frac{1}{(a+bx)(c+dx)^{2/3}} dx \rightarrow -\frac{\text{Log}[a+bx]}{2bq^2} - \frac{3}{2bq^2} \text{Subst}\left[\int \frac{1}{q-x} dx, x, (c+dx)^{1/3}\right] - \frac{3}{2bq} \text{Subst}\left[\int \frac{1}{q^2+qx+x^2} dx, x, (c+dx)^{1/3}\right]$$

Program code:

```
Int[1/((a_.*b_.*x_)*(c_.*d_.*x_)^(2/3)),x_Symbol] :=
  With[{q=Rt[(b*c-a*d)/b,3]},
    -Log[RemoveContent[a+b*x,x]]/(2*b*q^2) -
    3/(2*b*q^2)*Subst[Int[1/(q-x),x],x,(c+d*x)^(1/3)] -
    3/(2*b*q)*Subst[Int[1/(q^2+q*x+x^2),x],x,(c+d*x)^(1/3)]] /;
  FreeQ[{a,b,c,d},x] && PosQ[(b*c-a*d)/b]
```

$$2: \int \frac{1}{(a+bx)(c+dx)^{2/3}} dx \text{ when } \frac{b c - a d}{b} \neq 0$$

Derivation: Integration by substitution

Basis: Let  $q = \left(-\frac{b c - a d}{b}\right)^{1/3}$ , then  $\frac{1}{(a+bx)(c+dx)^{2/3}} = -\frac{1}{2 q^2 (a+bx)} + \text{Subst}\left[\frac{3}{2 b q^2 (q+x)} + \frac{3}{2 b q (q^2 - q x + x^2)}, x, (c+dx)^{1/3}\right] \partial_x (c+dx)^{1/3}$

Rule 1.1.1.2.7.3.2: If  $\frac{b c - a d}{b} \neq 0$ , let  $q = \left(-\frac{b c - a d}{b}\right)^{1/3}$ , then

$$\int \frac{1}{(a+bx)(c+dx)^{2/3}} dx \rightarrow$$

$$-\frac{\text{Log}[a+bx]}{2 b q^2} + \frac{3}{2 b q^2} \text{Subst}\left[\int \frac{1}{q+x} dx, x, (c+dx)^{1/3}\right] + \frac{3}{2 b q} \text{Subst}\left[\int \frac{1}{q^2 - qx + x^2} dx, x, (c+dx)^{1/3}\right]$$

Program code:

```
Int[1/((a_+b_.*x_)*(c_+d_.*x_)^(2/3)),x_Symbol] :=
  With[{q=Rt[-(b*c-a*d)/b,3]},
    -Log[RemoveContent[a+b*x,x]]/(2*b*q^2) +
    3/(2*b*q^2)*Subst[Int[1/(q+x),x],x,(c+d*x)^(1/3)] +
    3/(2*b*q)*Subst[Int[1/(q^2-q*x+x^2),x],x,(c+d*x)^(1/3)]] /;
  FreeQ[{a,b,c,d},x] && NegQ[(b*c-a*d)/b]
```

$$4. \int \frac{1}{(a+bx)^{1/3} (c+dx)^{2/3}} dx \text{ when } bc - ad \neq 0$$

$$1: \int \frac{1}{(a+bx)^{1/3} (c+dx)^{2/3}} dx \text{ when } bc - ad \neq 0 \wedge \frac{d}{b} > 0$$

Rule 1.1.1.2.7.4.1: If  $bc - ad \neq 0 \wedge \frac{d}{b} > 0$ , let  $q = \left(\frac{d}{b}\right)^{1/3}$ , then

$$\int \frac{1}{(a+bx)^{1/3} (c+dx)^{2/3}} dx \rightarrow -\frac{\sqrt{3} q}{d} \text{ArcTan}\left[\frac{2 q (a+bx)^{1/3}}{\sqrt{3} (c+dx)^{1/3}} + \frac{1}{\sqrt{3}}\right] - \frac{q}{2d} \text{Log}[c+dx] - \frac{3q}{2d} \text{Log}\left[\frac{q (a+bx)^{1/3}}{(c+dx)^{1/3}} - 1\right]$$

Program code:

```
Int[1/((a_.+b_.*x_)^(1/3)*(c_.+d_.*x_)^(2/3)),x_Symbol] :=
  With[{q=Rt[d/b,3]},
    -Sqrt[3]*q/d*ArcTan[2*q*(a+b*x)^(1/3)/(Sqrt[3]*(c+d*x)^(1/3))+1/Sqrt[3]] -
    q/(2*d)*Log[c+d*x] -
    3*q/(2*d)*Log[q*(a+b*x)^(1/3)/(c+d*x)^(1/3)-1]] /;
  FreeQ[{a,b,c,d},x] && NeQ[b*c-a*d,0] && PosQ[d/b]
```

**2:**  $\int \frac{1}{(a+bx)^{1/3} (c+dx)^{2/3}} dx$  when  $b c - a d \neq 0 \wedge \frac{d}{b} \neq 0$

Rule 1.1.1.2.7.4.2: If  $b c - a d \neq 0 \wedge \frac{d}{b} \neq 0$ , let  $q = \left(-\frac{d}{b}\right)^{1/3}$ , then

$$\int \frac{1}{(a+bx)^{1/3} (c+dx)^{2/3}} dx \rightarrow \frac{\sqrt{3} q}{d} \operatorname{ArcTan}\left[\frac{1}{\sqrt{3}} - \frac{2 q (a+bx)^{1/3}}{\sqrt{3} (c+dx)^{1/3}}\right] + \frac{q}{2 d} \operatorname{Log}[c+dx] + \frac{3 q}{2 d} \operatorname{Log}\left[\frac{q (a+bx)^{1/3}}{(c+dx)^{1/3}} + 1\right]$$

Program code:

```
Int[1/((a_.*b_.*x_)^(1/3)*(c_.*d_.*x_)^(2/3)),x_Symbol] :=
  With[{q=Rt[-d/b,3]},
    Sqrt[3]*q/d*ArcTan[1/Sqrt[3]-2*q*(a+b*x)^(1/3)/(Sqrt[3]*(c+d*x)^(1/3))] +
    q/(2*d)*Log[c+d*x] +
    3*q/(2*d)*Log[q*(a+b*x)^(1/3)/(c+d*x)^(1/3)+1]] /;
FreeQ[{a,b,c,d},x] && NeQ[b*c-a*d,0] && NegQ[d/b]
```

**5:**  $\int (a+bx)^m (c+dx)^n dx$  when  $bc - ad \neq 0 \wedge -1 < m < 0 \wedge n = m \wedge 3 \leq \text{Denominator}[m] \leq 4$

Derivation: Piecewise constant extraction

Basis:  $\partial_x \frac{(a+bx)^m (c+dx)^m}{((a+bx)(c+dx))^m} = 0$

Rule 1.1.1.2.7.5: If  $bc - ad \neq 0 \wedge -1 < m < 0 \wedge 3 \leq \text{Denominator}[m] \leq 4$ , then

$$\int (a+bx)^m (c+dx)^n dx \rightarrow \frac{(a+bx)^m (c+dx)^m}{(ac + (bc+ad)x + bdx^2)^m} \int (ac + (bc+ad)x + bdx^2)^m dx$$

$$\int (a+bx)^m (c+dx)^n dx \rightarrow \frac{(a+bx)^m (c+dx)^m}{((a+bx)(c+dx))^m} \int (ac + (bc+ad)x + bdx^2)^m dx$$

Program code:

```
Int[(a_.+b_.x_)^m_*(c_.+d_.x_)^m_,x_Symbol] :=
  (a+b*x)^m*(c+d*x)^m/(a*c+(b*c+a*d)*x+b*d*x^2)^m*Int[(a*c+(b*c+a*d)*x+b*d*x^2)^m,x] /;
FreeQ[{a,b,c,d},x] && NeQ[b*c-a*d,0] && LtQ[-1,m,0] && LeQ[3,Denominator[m],4] && AtomQ[b*c+a*d]
```

```
Int[(a_.+b_.x_)^m_*(c_.+d_.x_)^m_,x_Symbol] :=
  (a+b*x)^m*(c+d*x)^m/((a+b*x)*(c+d*x))^m*Int[(a*c+(b*c+a*d)*x+b*d*x^2)^m,x] /;
FreeQ[{a,b,c,d},x] && NeQ[b*c-a*d,0] && LtQ[-1,m,0] && LeQ[3,Denominator[m],4]
```

**6:**  $\int (a+bx)^m (c+dx)^n dx$  when  $bc - ad \neq 0 \wedge -1 < m < 0 \wedge -1 \leq n < 0$

Derivation: Integration by substitution

Basis: If  $p \in \mathbb{Z}^+$ , then

$$(a+bx)^m (c+dx)^n = \frac{p}{b} \text{Subst} \left[ x^{p(m+1)-1} \left( c - \frac{ad}{b} + \frac{d}{b} x^p \right)^n, x, (a+bx)^{1/p} \right] \partial_x (a+bx)^{1/p}$$

Rule 1.1.1.2.7.7: If  $bc - ad \neq 0 \wedge -1 < m < 0 \wedge -1 \leq n < 0$ , let  $p = \text{Denominator}[m]$ , then

$$\int (a+bx)^m (c+dx)^n dx \rightarrow \frac{p}{b} \text{Subst} \left[ \int x^{p(m+1)-1} \left( c - \frac{ad}{b} + \frac{d}{b} x^p \right)^n dx, x, (a+bx)^{1/p} \right]$$

Program code:

```
Int[(a_.+b_.*x_)^m_*(c_.+d_.*x_)^n_,x_Symbol] :=
  With[{p=Denominator[m]},
    p/b*Subst[Int[x^(p*(m+1)-1)*(c-a*d/b+d*x^p/b)^n,x,(a+b*x)^(1/p)]] /;
  FreeQ[{a,b,c,d},x] && NeQ[b*c-a*d,0] && LtQ[-1,m,0] && LeQ[-1,n,0] && LeQ[Denominator[n],Denominator[m]] &&
  IntLinearQ[a,b,c,d,m,n,x]
```

H.  $\int (a+bx)^m (c+dx)^n dx$  when  $bc - ad \neq 0$

$$1. \int (bx)^m (c+dx)^n dx$$

**1:**  $\int (bx)^m (c+dx)^n dx$  when  $m \notin \mathbb{Z} \wedge (n \in \mathbb{Z} \vee c > 0)$

Rule 1.1.1.2.H.1.1: If  $m \notin \mathbb{Z} \wedge (n \in \mathbb{Z} \vee c > 0)$ , then

$$\int (bx)^m (c+dx)^n dx \rightarrow \frac{c^n (bx)^{m+1}}{b(m+1)} \text{Hypergeometric2F1}\left[-n, m+1, m+2, -\frac{dx}{c}\right]$$

Program code:

```
Int[(b_.**x_)^m_*(c_+d_.**x_)^n_,x_Symbol] :=
  c^n*(b*x)^(m+1)/(b*(m+1))*Hypergeometric2F1[-n,m+1,m+2,-d*x/c] /;
FreeQ[{b,c,d,m,n},x] && Not[IntegerQ[m]] && (IntegerQ[n] || GtQ[c,0] && Not[EqQ[n,-1/2] && EqQ[c^2-d^2,0] && GtQ[-d/(b*c),0]])
```

**2:**  $\int (bx)^m (c+dx)^n dx$  when  $n \notin \mathbb{Z} \wedge (m \in \mathbb{Z} \vee -\frac{d}{bc} > 0)$

Rule 1.1.1.2.H.1.2: If  $n \notin \mathbb{Z} \wedge (m \in \mathbb{Z} \vee -\frac{d}{bc} > 0)$ , then

$$\int (bx)^m (c+dx)^n dx \rightarrow \frac{(c+dx)^{n+1}}{d(n+1) \left(-\frac{d}{bc}\right)^m} \text{Hypergeometric2F1}\left[-m, n+1, n+2, 1+\frac{dx}{c}\right]$$

Program code:

```
Int[(b_.**x_)^m_*(c_+d_.**x_)^n_,x_Symbol] :=
  (c+d*x)^(n+1)/(d*(n+1)*(-d/(b*c))^m)*Hypergeometric2F1[-m,n+1,n+2,1+d*x/c] /;
FreeQ[{b,c,d,m,n},x] && Not[IntegerQ[n]] && (IntegerQ[m] || GtQ[-d/(b*c),0])
```



$$3. \int (bx)^m (c+dx)^n dx \text{ when } m \notin \mathbb{Z} \wedge n \notin \mathbb{Z} \wedge c \neq 0 \wedge -\frac{d}{bc} \neq 0$$

$$1: \int (bx)^m (c+dx)^n dx \text{ when } m \notin \mathbb{Z} \wedge n \notin \mathbb{Z} \wedge c \neq 0 \wedge -\frac{d}{bc} \neq 0 \wedge (m \in \mathbb{R} \vee n \notin \mathbb{R})$$

### Derivation: Piecewise constant extraction

$$\text{Basis: } \partial_x \frac{(c+dx)^n}{\left(1+\frac{dx}{c}\right)^n} = 0$$

Rule 1.1.1.2.H.1.3.1: If  $m \notin \mathbb{Z} \wedge n \notin \mathbb{Z} \wedge c \neq 0 \wedge -\frac{d}{bc} \neq 0 \wedge (m \in \mathbb{R} \vee n \notin \mathbb{R})$ , then

$$\int (bx)^m (c+dx)^n dx \rightarrow \frac{c^{\text{IntPart}[n]} (c+dx)^{\text{FracPart}[n]}}{\left(1+\frac{dx}{c}\right)^{\text{FracPart}[n]}} \int (bx)^m \left(1+\frac{dx}{c}\right)^n dx$$

### Program code:

```
Int[(b_.**x_)^m_*(c_+d_.**x_)^n_,x_Symbol] :=
  c^IntPart[n]*(c+d*x)^FracPart[n]/(1+d*x/c)^FracPart[n]*Int[(b*x)^m*(1+d*x/c)^n,x] /;
FreeQ[{b,c,d,m,n},x] && Not[IntegerQ[m]] && Not[IntegerQ[n]] && Not[GtQ[c,0]] && Not[GtQ[-d/(b*c),0]] &&
  (RationalQ[m] && Not[EqQ[n,-1/2] && EqQ[c^2-d^2,0]] || Not[RationalQ[n]])
```

**2:**  $\int (bx)^m (c+dx)^n dx$  when  $m \notin \mathbb{Z} \wedge n \notin \mathbb{Z} \wedge c \neq 0 \wedge -\frac{d}{bc} \neq 0 \wedge \neg (m \in \mathbb{R} \vee n \in \mathbb{R})$

Derivation: Piecewise constant extraction

Basis:  $\partial_x \frac{(bx)^m}{\left(-\frac{dx}{c}\right)^m} == 0$

Rule 1.1.1.2.H.1.3.2: If  $m \notin \mathbb{Z} \wedge n \notin \mathbb{Z} \wedge c \neq 0 \wedge -\frac{d}{bc} \neq 0 \wedge \neg (m \in \mathbb{R} \vee n \in \mathbb{R})$ , then

$$\int (bx)^m (c+dx)^n dx \rightarrow \frac{\left(-\frac{bc}{d}\right)^{\text{IntPart}[m]} (bx)^{\text{FracPart}[m]}}{\left(-\frac{dx}{c}\right)^{\text{FracPart}[m]}} \int \left(-\frac{dx}{c}\right)^m (c+dx)^n dx$$

Program code:

```
Int[(b_.*x_)^m_*(c_+d_.*x_)^n_,x_Symbol] :=
  (-b*c/d)^IntPart[m]*(b*x)^FracPart[m]/(-d*x/c)^FracPart[m]*Int[(-d*x/c)^m*(c+d*x)^n,x] /;
FreeQ[{b,c,d,m,n},x] && Not[IntegerQ[m]] && Not[IntegerQ[n]] && Not[GtQ[c,0]] && Not[GtQ[-d/(b*c),0]]
```

2.  $\int (a+bx)^m (c+dx)^n dx$  when  $bc-ad \neq 0 \wedge m \notin \mathbb{Z}$

1:  $\int (a+bx)^m (c+dx)^n dx$  when  $bc-ad \neq 0 \wedge m \notin \mathbb{Z} \wedge \left(n \in \mathbb{Z} \vee \frac{b}{bc-ad} > 0\right)$

Rule 1.1.1.2.H.2.2.1: If  $bc-ad \neq 0 \wedge m \notin \mathbb{Z} \wedge \left(n \in \mathbb{Z} \vee \frac{b}{bc-ad} > 0\right)$ , then

$$\int (a+bx)^m (c+dx)^n dx \rightarrow \frac{(a+bx)^{m+1}}{b(m+1) \left(\frac{b}{bc-ad}\right)^n} \text{Hypergeometric2F1}\left[-n, m+1, m+2, -\frac{d(a+bx)}{bc-ad}\right]$$

Program code:

```
Int[(a_+b_.**x_)^m_*(c_+d_.**x_)^n_,x_Symbol] :=
  (b*c-a*d)^n*(a+b*x)^(m+1)/(b^(n+1)*(m+1))*Hypergeometric2F1[-n,m+1,m+2,-d*(a+b*x)/(b*c-a*d)] /;
FreeQ[{a,b,c,d,m},x] && NeQ[b*c-a*d,0] && Not[IntegerQ[m]] && IntegerQ[n]
```

```
Int[(a_+b_.**x_)^m_*(c_+d_.**x_)^n_,x_Symbol] :=
  (a+b*x)^(m+1)/(b*(m+1)*(b/(b*c-a*d))^n)*Hypergeometric2F1[-n,m+1,m+2,-d*(a+b*x)/(b*c-a*d)] /;
FreeQ[{a,b,c,d,m,n},x] && NeQ[b*c-a*d,0] && Not[IntegerQ[m]] && Not[IntegerQ[n]] && GtQ[b/(b*c-a*d),0] &&
  (RationalQ[m] || Not[RationalQ[n]] && GtQ[-d/(b*c-a*d),0])
```

**2:**  $\int (a+bx)^m (c+dx)^n dx$  when  $bc-ad \neq 0 \wedge m \notin \mathbb{Z} \wedge n \notin \mathbb{Z} \wedge \frac{b}{bc-ad} \neq 0$

Derivation: Piecewise constant extraction

Basis:  $\partial_x \frac{(c+dx)^n}{\left(\frac{bc}{bc-ad} + \frac{bdx}{bc-ad}\right)^n} = 0$

Rule 1.1.1.2.H.2.2.2: If  $bc-ad \neq 0 \wedge m \notin \mathbb{Z} \wedge n \notin \mathbb{Z} \wedge \frac{b}{bc-ad} \neq 0$ , then

$$\int (a+bx)^m (c+dx)^n dx \rightarrow \frac{(c+dx)^{\text{FracPart}[n]}}{\left(\frac{b}{bc-ad}\right)^{\text{IntPart}[n]} \left(\frac{b(c+dx)}{bc-ad}\right)^{\text{FracPart}[n]}} \int (a+bx)^m \left(\frac{bc}{bc-ad} + \frac{bdx}{bc-ad}\right)^n dx$$

Program code:

```
Int[(a+b*x_)^m_*(c+d*x_)^n_,x_Symbol] :=
  (c+d*x)^FracPart[n]/((b/(b*c-a*d))^IntPart[n]*(b*(c+d*x)/(b*c-a*d))^FracPart[n])*
  Int[(a+b*x)^m*Simp[b*c/(b*c-a*d)+b*d*x/(b*c-a*d),x]^n,x] /;
FreeQ[{a,b,c,d,m,n},x] && NeQ[b*c-a*d,0] && Not[IntegerQ[m]] && Not[IntegerQ[n]] && (RationalQ[m] || Not[SimplerQ[n+1,m+1]])
```

**S:**  $\int (a + b u)^m (c + d u)^n dx$  when  $u = e + f x$

Derivation: Integration by substitution

Rule 1.1.1.2.S: If  $u = e + f x$ , then

$$\int (a + b u)^m (c + d u)^n dx \rightarrow \frac{1}{f} \text{Subst} \left[ \int (a + b x)^m (c + d x)^n dx, x, u \right]$$

Program code:

```
Int[(a_.+b_.*u_)^m_.*(c_.+d_.*u_)^n_,x_Symbol] :=
  1/Coefficient[u,x,1]*Subst[Int[(a+b*x)^m*(c+d*x)^n,x],x,u] /;
FreeQ[{a,b,c,d,m,n},x] && LinearQ[u,x] && NeQ[Coefficient[u,x,0],0]
```

```
(* IntLinearQ[a,b,c,d,m,n,x] returns True iff (a+b*x)^m*(c+d*x)^n is integrable wrt x in terms of non-hypergeometric functions. *)
IntLinearQ[a_,b_,c_,d_,m_,n_,x_] :=
  IGtQ[m,0] || IGtQ[n,0] || IntegersQ[3*m,3*n] || IntegersQ[4*m,4*n] || IntegersQ[2*m,6*n] || IntegersQ[6*m,2*n] || ILtQ[m+n,-1] || In
```