

Rules for integrands involving $(a + b \operatorname{ArcTan}[c x])^p$

1: $\int (a + b \operatorname{ArcTan}[c x])^p dx$ when $p \in \mathbb{Z}^+$

Derivation: Integration by parts

Basis: $\partial_x (a + b \operatorname{ArcTan}[c x])^p = b c p \frac{(a + b \operatorname{ArcTan}[c x])^{p-1}}{1 + c^2 x^2}$

Rule: If $p \in \mathbb{Z}^+$, then

$$\int (a + b \operatorname{ArcTan}[c x])^p dx \rightarrow x (a + b \operatorname{ArcTan}[c x])^p - b c p \int \frac{x (a + b \operatorname{ArcTan}[c x])^{p-1}}{1 + c^2 x^2} dx$$

Program code:

```
Int[(a_.+b_.*ArcTan[c_.*x_])^p_.,x_Symbol] :=
  x*(a+b*ArcTan[c*x])^p -
  b*c*p*Int[x*(a+b*ArcTan[c*x])^(p-1)/(1+c^2*x^2),x] /;
FreeQ[{a,b,c},x] && IGtQ[p,0]
```

```
Int[(a_.+b_.*ArcCot[c_.*x_])^p_.,x_Symbol] :=
  x*(a+b*ArcCot[c*x])^p +
  b*c*p*Int[x*(a+b*ArcCot[c*x])^(p-1)/(1+c^2*x^2),x] /;
FreeQ[{a,b,c},x] && IGtQ[p,0]
```

2. $\int (d x)^m (a + b \operatorname{ArcTan}[c x])^p dx$ when $p \in \mathbb{Z}^+$

1. $\int \frac{(a + b \operatorname{ArcTan}[c x])^p}{x} dx$ when $p \in \mathbb{Z}^+$

1: $\int \frac{a + b \operatorname{ArcTan}[c x]}{x} dx$

Derivation: Algebraic expansion

Basis: $\operatorname{ArcTan}[z] = \frac{i}{2} \operatorname{Log}[1 - iz] - \frac{i}{2} \operatorname{Log}[1 + iz]$

Basis: $\operatorname{ArcCot}[z] = \frac{i}{2} \operatorname{Log}[1 - \frac{i}{z}] - \frac{i}{2} \operatorname{Log}[1 + \frac{i}{z}]$

Rule:

$$\begin{aligned} \int \frac{a + b \operatorname{ArcTan}[c x]}{x} dx &\rightarrow a \int \frac{1}{x} dx + \frac{ib}{2} \int \frac{\operatorname{Log}[1 - icx]}{x} dx - \frac{ib}{2} \int \frac{\operatorname{Log}[1 + icx]}{x} dx \\ &\rightarrow a \operatorname{Log}[x] + \frac{ib}{2} \operatorname{PolyLog}[2, -icx] - \frac{ib}{2} \operatorname{PolyLog}[2, icx] \end{aligned}$$

Program code:

```
Int[(a_.+b_.*ArcTan[c_.*x_])/x_,x_Symbol] :=
  a*Log[x] + I*b/2*Int[Log[1-I*c*x]/x,x] - I*b/2*Int[Log[1+I*c*x]/x,x] /;
FreeQ[{a,b,c},x]
```

```
Int[(a_.+b_.*ArcCot[c_.*x_])/x_,x_Symbol] :=
  a*Log[x] + I*b/2*Int[Log[1-I/(c*x)]/x,x] - I*b/2*Int[Log[1+I/(c*x)]/x,x] /;
FreeQ[{a,b,c},x]
```

$$2: \int \frac{(a + b \operatorname{ArcTan}[c x])^p}{x} dx \text{ when } p - 1 \in \mathbb{Z}^+$$

Derivation: Integration by parts

$$\text{Basis: } \frac{1}{x} = 2 \partial_x \operatorname{ArcTanh}\left[1 - \frac{2}{1 + i c x}\right]$$

Rule: If $p - 1 \in \mathbb{Z}^+$, then

$$\int \frac{(a + b \operatorname{ArcTan}[c x])^p}{x} dx \rightarrow 2 (a + b \operatorname{ArcTan}[c x])^p \operatorname{ArcTanh}\left[1 - \frac{2}{1 + i c x}\right] - 2 b c p \int \frac{(a + b \operatorname{ArcTan}[c x])^{p-1} \operatorname{ArcTanh}\left[1 - \frac{2}{1 + i c x}\right]}{1 + c^2 x^2} dx$$

Program code:

```
Int[(a_.+b_.*ArcTan[c_.*x_])^p_/x_,x_Symbol] :=
  2*(a+b*ArcTan[c*x])^p*ArcTanh[1-2/(1+I*c*x)] -
  2*b*c*p*Int[(a+b*ArcTan[c*x])^(p-1)*ArcTanh[1-2/(1+I*c*x)]/(1+c^2*x^2),x] /;
FreeQ[{a,b,c},x] && IGtQ[p,1]
```

```
Int[(a_.+b_.*ArcCot[c_.*x_])^p_/x_,x_Symbol] :=
  2*(a+b*ArcCot[c*x])^p*ArcCoth[1-2/(1+I*c*x)] +
  2*b*c*p*Int[(a+b*ArcCot[c*x])^(p-1)*ArcCoth[1-2/(1+I*c*x)]/(1+c^2*x^2),x] /;
FreeQ[{a,b,c},x] && IGtQ[p,1]
```

$$2: \int (d x)^m (a + b \operatorname{ArcTan}[c x])^p dx \text{ when } p \in \mathbb{Z}^+ \wedge (p = 1 \vee m \in \mathbb{Z}) \wedge m \neq -1$$

Derivation: Integration by parts

$$\text{Basis: } \partial_x (a + b \operatorname{ArcTan}[c x])^p = \frac{b c p (a + b \operatorname{ArcTan}[c x])^{p-1}}{1 + c^2 x^2}$$

Rule: If $p \in \mathbb{Z}^+ \wedge (p = 1 \vee m \in \mathbb{Z}) \wedge m \neq -1$, then

$$\int (dx)^m (a + b \operatorname{ArcTan}[cx])^p dx \rightarrow \frac{(dx)^{m+1} (a + b \operatorname{ArcTan}[cx])^p}{d(m+1)} - \frac{bcp}{d(m+1)} \int \frac{(dx)^{m+1} (a + b \operatorname{ArcTan}[cx])^{p-1}}{1+c^2x^2} dx$$

Program code:

```
Int[(d_.**x_)^m_.*(a_.+b_.**ArcTan[c_.**x_])^p_,x_Symbol] :=
  (d*x)^(m+1)*(a+b**ArcTan[c*x])^p/(d*(m+1)) -
  b*c*p/(d*(m+1))*Int[(d*x)^(m+1)*(a+b**ArcTan[c*x])^(p-1)/(1+c^2*x^2),x] /;
FreeQ[{a,b,c,d,m},x] && IGtQ[p,0] && (EqQ[p,1] || IntegerQ[m]) && NeQ[m,-1]
```

```
Int[(d_.**x_)^m_.*(a_.+b_.**ArcCot[c_.**x_])^p_,x_Symbol] :=
  (d*x)^(m+1)*(a+b**ArcCot[c*x])^p/(d*(m+1)) +
  b*c*p/(d*(m+1))*Int[(d*x)^(m+1)*(a+b**ArcCot[c*x])^(p-1)/(1+c^2*x^2),x] /;
FreeQ[{a,b,c,d,m},x] && IGtQ[p,0] && (EqQ[p,1] || IntegerQ[m]) && NeQ[m,-1]
```

$$3. \int (d + ex)^q (a + b \operatorname{ArcTan}[cx])^p dx \text{ when } p \in \mathbb{Z}^+$$

$$1. \int \frac{(a + b \operatorname{ArcTan}[cx])^p}{d + ex} dx \text{ when } p \in \mathbb{Z}^+$$

$$1: \int \frac{(a + b \operatorname{ArcTan}[cx])^p}{d + ex} dx \text{ when } p \in \mathbb{Z}^+ \wedge c^2 d^2 + e^2 = 0$$

Derivation: Integration by parts

$$\text{Basis: } \frac{1}{d+ex} = -\frac{1}{e} \partial_x \operatorname{Log} \left[\frac{2}{1+\frac{ex}{d}} \right]$$

Rule: If $p \in \mathbb{Z}^+ \wedge c^2 d^2 + e^2 = 0$, then

$$\int \frac{(a + b \operatorname{ArcTan}[cx])^p}{d + ex} dx \rightarrow -\frac{(a + b \operatorname{ArcTan}[cx])^p \operatorname{Log} \left[\frac{2}{1+\frac{ex}{d}} \right]}{e} + \frac{bc p}{e} \int \frac{(a + b \operatorname{ArcTan}[cx])^{p-1} \operatorname{Log} \left[\frac{2}{1+\frac{ex}{d}} \right]}{1 + c^2 x^2} dx$$

Program code:

```
Int[(a_.+b_.*ArcTan[c_.*x_])^p_./(d_.+e_.*x_),x_Symbol] :=
  -(a+b*ArcTan[c*x])^p*Log[2/(1+e*x/d)]/e +
  b*c*p/e*Int[(a+b*ArcTan[c*x])^(p-1)*Log[2/(1+e*x/d)]/(1+c^2*x^2),x] /;
FreeQ[{a,b,c,d,e},x] && IGtQ[p,0] && EqQ[c^2*d^2+e^2,0]
```

```
Int[(a_.+b_.*ArcCot[c_.*x_])^p_./(d_.+e_.*x_),x_Symbol] :=
  -(a+b*ArcCot[c*x])^p*Log[2/(1+e*x/d)]/e -
  b*c*p/e*Int[(a+b*ArcCot[c*x])^(p-1)*Log[2/(1+e*x/d)]/(1+c^2*x^2),x] /;
FreeQ[{a,b,c,d,e},x] && IGtQ[p,0] && EqQ[c^2*d^2+e^2,0]
```

$$2. \int \frac{(a + b \operatorname{ArcTan}[c x])^p}{d + e x} dx \text{ when } p \in \mathbb{Z}^+ \wedge c^2 d^2 + e^2 \neq 0$$

$$1: \int \frac{a + b \operatorname{ArcTan}[c x]}{d + e x} dx \text{ when } c^2 d^2 + e^2 \neq 0$$

Derivation: Algebraic expansion and integration by parts

$$\text{Basis: } \frac{1}{d+e x} == \frac{c}{e (i+c x)} - \frac{c d-i e}{e (i+c x) (d+e x)}$$

$$\text{Basis: } \frac{1}{i+c x} == -\frac{1}{c} \partial_x \operatorname{Log}\left[\frac{2}{1-i c x}\right]$$

$$\text{Basis: } \frac{1}{(i+c x) (d+e x)} == -\frac{1}{c d-i e} \partial_x \operatorname{Log}\left[\frac{2 c (d+e x)}{(c d+i e) (1-i c x)}\right]$$

$$\text{Basis: } \partial_x (a + b \operatorname{ArcTan}[c x]) == \frac{b c}{1+c^2 x^2}$$

Rule: If $c^2 d^2 + e^2 \neq 0$, then

$$\begin{aligned} \int \frac{a + b \operatorname{ArcTan}[c x]}{d + e x} dx &\rightarrow \frac{c}{e} \int \frac{a + b \operatorname{ArcTan}[c x]}{i + c x} dx - \frac{c d - i e}{e} \int \frac{a + b \operatorname{ArcTan}[c x]}{(i + c x) (d + e x)} dx \rightarrow \\ &-\frac{(a + b \operatorname{ArcTan}[c x]) \operatorname{Log}\left[\frac{2}{1-i c x}\right]}{e} + \frac{b c}{e} \int \frac{\operatorname{Log}\left[\frac{2}{1-i c x}\right]}{1 + c^2 x^2} dx + \frac{(a + b \operatorname{ArcTan}[c x]) \operatorname{Log}\left[\frac{2 c (d+e x)}{(c d+i e) (1-i c x)}\right]}{e} - \frac{b c}{e} \int \frac{\operatorname{Log}\left[\frac{2 c (d+e x)}{(c d+i e) (1-i c x)}\right]}{1 + c^2 x^2} dx \rightarrow \\ &-\frac{(a + b \operatorname{ArcTan}[c x]) \operatorname{Log}\left[\frac{2}{1-i c x}\right]}{e} + \frac{i b \operatorname{PolyLog}\left[2, 1 - \frac{2}{1-i c x}\right]}{2 e} + \frac{(a + b \operatorname{ArcTan}[c x]) \operatorname{Log}\left[\frac{2 c (d+e x)}{(c d+i e) (1-i c x)}\right]}{e} - \frac{i b \operatorname{PolyLog}\left[2, 1 - \frac{2 c (d+e x)}{(c d+i e) (1-i c x)}\right]}{2 e} \end{aligned}$$

Program code:

```

Int[(a_.+b_.*ArcTan[c_.*x_])/(d_+e_.*x_),x_Symbol] :=
  -(a+b*ArcTan[c*x])*Log[2/(1-I*c*x)]/e +
  b*c/e*Int[Log[2/(1-I*c*x)]/(1+c^2*x^2),x] +
  (a+b*ArcTan[c*x])*Log[2*c*(d+e*x)/((c*d+I*e)*(1-I*c*x))]/e -
  b*c/e*Int[Log[2*c*(d+e*x)/((c*d+I*e)*(1-I*c*x))]/(1+c^2*x^2),x] /;
FreeQ[{a,b,c,d,e},x] && NeQ[c^2*d^2+e^2,0]

```

```

Int[(a_.+b_.*ArcCot[c_.*x_])/(d_+e_.*x_),x_Symbol] :=
  -(a+b*ArcCot[c*x])*Log[2/(1-I*c*x)]/e -
  b*c/e*Int[Log[2/(1-I*c*x)]/(1+c^2*x^2),x] +
  (a+b*ArcCot[c*x])*Log[2*c*(d+e*x)/((c*d+I*e)*(1-I*c*x))]/e +
  b*c/e*Int[Log[2*c*(d+e*x)/((c*d+I*e)*(1-I*c*x))]/(1+c^2*x^2),x] /;
FreeQ[{a,b,c,d,e},x] && NeQ[c^2*d^2+e^2,0]

```

$$2: \int \frac{(a + b \operatorname{ArcTan}[c x])^2}{d + e x} dx \text{ when } c^2 d^2 + e^2 \neq 0$$

Derivation: Algebraic expansion and integration by parts

$$\text{Basis: } \frac{1}{d+ex} = \frac{c}{e(i+cx)} - \frac{cd-ie}{e(i+cx)(d+ex)}$$

$$\text{Basis: } \frac{1}{i+cx} = -\frac{1}{c} \partial_x \operatorname{Log}\left[\frac{2}{1-icx}\right]$$

$$\text{Basis: } \frac{1}{(i+cx)(d+ex)} = -\frac{1}{cd-ie} \partial_x \operatorname{Log}\left[\frac{2c(d+ex)}{(cd+ie)(1-icx)}\right]$$

$$\text{Basis: } \partial_x (a + b \operatorname{ArcTan}[cx])^2 = \frac{2bc(a+b \operatorname{ArcTan}[cx])}{1+c^2 x^2}$$

Rule: If $c^2 d^2 + e^2 \neq 0$, then

$$\begin{aligned} \int \frac{(a + b \operatorname{ArcTan}[cx])^2}{d + ex} dx &\rightarrow \frac{c}{e} \int \frac{(a + b \operatorname{ArcTan}[cx])^2}{i + cx} dx - \frac{cd - ie}{e} \int \frac{(a + b \operatorname{ArcTan}[cx])^2}{(i + cx)(d + ex)} dx \rightarrow \\ &- \frac{(a + b \operatorname{ArcTan}[cx])^2 \operatorname{Log}\left[\frac{2}{1-icx}\right]}{e} + \frac{2bc}{e} \int \frac{(a + b \operatorname{ArcTan}[cx]) \operatorname{Log}\left[\frac{2}{1-icx}\right]}{1 + c^2 x^2} dx + \end{aligned}$$

$$\begin{aligned}
& \frac{(a + b \operatorname{ArcTan}[c x])^2 \operatorname{Log}\left[\frac{2 c (d+e x)}{(c d+i e) (1-i c x)}\right]}{e} - \frac{2 b c}{e} \int \frac{(a + b \operatorname{ArcTan}[c x]) \operatorname{Log}\left[\frac{2 c (d+e x)}{(c d+i e) (1-i c x)}\right]}{1 + c^2 x^2} dx \rightarrow \\
& - \frac{(a + b \operatorname{ArcTan}[c x])^2 \operatorname{Log}\left[\frac{2}{1-i c x}\right]}{e} + \frac{i b (a + b \operatorname{ArcTan}[c x]) \operatorname{PolyLog}\left[2, 1 - \frac{2}{1-i c x}\right]}{e} - \frac{b^2 \operatorname{PolyLog}\left[3, 1 - \frac{2}{1-i c x}\right]}{2 e} + \\
& \frac{(a + b \operatorname{ArcTan}[c x])^2 \operatorname{Log}\left[\frac{2 c (d+e x)}{(c d+i e) (1-i c x)}\right]}{e} - \frac{i b (a + b \operatorname{ArcTan}[c x]) \operatorname{PolyLog}\left[2, 1 - \frac{2 c (d+e x)}{(c d+i e) (1-i c x)}\right]}{e} + \frac{b^2 \operatorname{PolyLog}\left[3, 1 - \frac{2 c (d+e x)}{(c d+i e) (1-i c x)}\right]}{2 e}
\end{aligned}$$

Program code:

```

Int[(a_.+b_.*ArcTan[c_.*x_])^2/(d_+e_.*x_),x_Symbol] :=
- (a+b*ArcTan[c*x])^2*Log[2/(1-I*c*x)]/e +
I*b*(a+b*ArcTan[c*x])*PolyLog[2,1-2/(1-I*c*x)]/e -
b^2*PolyLog[3,1-2/(1-I*c*x)]/(2*e) +
(a+b*ArcTan[c*x])^2*Log[2*c*(d+e*x)/((c*d+I*e)*(1-I*c*x))]/e -
I*b*(a+b*ArcTan[c*x])*PolyLog[2,1-2*c*(d+e*x)/((c*d+I*e)*(1-I*c*x))]/e +
b^2*PolyLog[3,1-2*c*(d+e*x)/((c*d+I*e)*(1-I*c*x))]/(2*e) /;
FreeQ[{a,b,c,d,e},x] && NeQ[c^2*d^2+e^2,0]

```

```

Int[(a_.+b_.*ArcCot[c_.*x_])^2/(d_+e_.*x_),x_Symbol] :=
- (a+b*ArcCot[c*x])^2*Log[2/(1-I*c*x)]/e -
I*b*(a+b*ArcCot[c*x])*PolyLog[2,1-2/(1-I*c*x)]/e -
b^2*PolyLog[3,1-2/(1-I*c*x)]/(2*e) +
(a+b*ArcCot[c*x])^2*Log[2*c*(d+e*x)/((c*d+I*e)*(1-I*c*x))]/e +
I*b*(a+b*ArcCot[c*x])*PolyLog[2,1-2*c*(d+e*x)/((c*d+I*e)*(1-I*c*x))]/e +
b^2*PolyLog[3,1-2*c*(d+e*x)/((c*d+I*e)*(1-I*c*x))]/(2*e) /;
FreeQ[{a,b,c,d,e},x] && NeQ[c^2*d^2+e^2,0]

```


$$\text{3: } \int \frac{(a + b \operatorname{ArcTan}[c x])^3}{d + e x} dx \text{ when } c^2 d^2 + e^2 \neq 0$$

Derivation: Algebraic expansion and integration by parts

$$\text{Basis: } \frac{1}{d+e x} == \frac{c}{e (i+c x)} - \frac{c d-i e}{e (i+c x) (d+e x)}$$

$$\text{Basis: } \frac{1}{i+c x} == -\frac{1}{c} \partial_x \operatorname{Log}\left[\frac{2}{1-i c x}\right]$$

$$\text{Basis: } \frac{1}{(i+c x) (d+e x)} == -\frac{1}{c d-i e} \partial_x \operatorname{Log}\left[\frac{2 c (d+e x)}{(c d+i e) (1-i c x)}\right]$$

$$\text{Basis: } \partial_x (a + b \operatorname{ArcTan}[c x])^3 == \frac{3 b c (a+b \operatorname{ArcTan}[c x])^2}{1+c^2 x^2}$$

Rule: If $c^2 d^2 + e^2 \neq 0$, then

$$\begin{aligned} \int \frac{(a + b \operatorname{ArcTan}[c x])^3}{d + e x} dx &\rightarrow \frac{c}{e} \int \frac{(a + b \operatorname{ArcTan}[c x])^3}{i + c x} dx - \frac{c d - i e}{e} \int \frac{(a + b \operatorname{ArcTan}[c x])^3}{(i + c x) (d + e x)} dx \rightarrow \\ &\quad - \frac{(a + b \operatorname{ArcTan}[c x])^3 \operatorname{Log}\left[\frac{2}{1-i c x}\right]}{e} + \frac{3 b c}{e} \int \frac{(a + b \operatorname{ArcTan}[c x])^2 \operatorname{Log}\left[\frac{2}{1-i c x}\right]}{1 + c^2 x^2} dx + \\ &\quad \frac{(a + b \operatorname{ArcTan}[c x])^3 \operatorname{Log}\left[\frac{2 c (d+e x)}{(c d+i e) (1-i c x)}\right]}{e} - \frac{3 b c}{e} \int \frac{(a + b \operatorname{ArcTan}[c x])^2 \operatorname{Log}\left[\frac{2 c (d+e x)}{(c d+i e) (1-i c x)}\right]}{1 + c^2 x^2} dx \rightarrow \\ &\quad - \frac{(a + b \operatorname{ArcTan}[c x])^3 \operatorname{Log}\left[\frac{2}{1-i c x}\right]}{e} + \frac{3 i b (a + b \operatorname{ArcTan}[c x])^2 \operatorname{PolyLog}\left[2, 1 - \frac{2}{1-i c x}\right]}{2 e} - \\ &\quad \frac{3 b^2 (a + b \operatorname{ArcTan}[c x]) \operatorname{PolyLog}\left[3, 1 - \frac{2}{1-i c x}\right]}{2 e} - \frac{3 i b^3 \operatorname{PolyLog}\left[4, 1 - \frac{2}{1-i c x}\right]}{4 e} + \\ &\quad \frac{(a + b \operatorname{ArcTan}[c x])^3 \operatorname{Log}\left[\frac{2 c (d+e x)}{(c d+i e) (1-i c x)}\right]}{e} - \frac{3 i b (a + b \operatorname{ArcTan}[c x])^2 \operatorname{PolyLog}\left[2, 1 - \frac{2 c (d+e x)}{(c d+i e) (1-i c x)}\right]}{2 e} + \\ &\quad \frac{3 b^2 (a + b \operatorname{ArcTan}[c x]) \operatorname{PolyLog}\left[3, 1 - \frac{2 c (d+e x)}{(c d+i e) (1-i c x)}\right]}{2 e} + \frac{3 i b^3 \operatorname{PolyLog}\left[4, 1 - \frac{2 c (d+e x)}{(c d+i e) (1-i c x)}\right]}{4 e} \end{aligned}$$

Program code:

```

Int[(a_.+b_.*ArcTan[c_.*x_])^3/(d_.+e_.*x_),x_Symbol] :=
- (a+b*ArcTan[c*x])^3*Log[2/(1-I*c*x)]/e +
3*I*b*(a+b*ArcTan[c*x])^2*PolyLog[2,1-2/(1-I*c*x)]/(2*e) -
3*b^2*(a+b*ArcTan[c*x])*PolyLog[3,1-2/(1-I*c*x)]/(2*e) -
3*I*b^3*PolyLog[4,1-2/(1-I*c*x)]/(4*e) +
(a+b*ArcTan[c*x])^3*Log[2*c*(d+e*x)/((c*d+I*e)*(1-I*c*x))]/e -
3*I*b*(a+b*ArcTan[c*x])^2*PolyLog[2,1-2*c*(d+e*x)/((c*d+I*e)*(1-I*c*x))]/(2*e) +
3*b^2*(a+b*ArcTan[c*x])*PolyLog[3,1-2*c*(d+e*x)/((c*d+I*e)*(1-I*c*x))]/(2*e) +
3*I*b^3*PolyLog[4,1-2*c*(d+e*x)/((c*d+I*e)*(1-I*c*x))]/(4*e) /;
FreeQ[{a,b,c,d,e},x] && NeQ[c^2*d^2+e^2,0]

```

```

Int[(a_.+b_.*ArcCot[c_.*x_])^3/(d_.+e_.*x_),x_Symbol] :=
- (a+b*ArcCot[c*x])^3*Log[2/(1-I*c*x)]/e -
3*I*b*(a+b*ArcCot[c*x])^2*PolyLog[2,1-2/(1-I*c*x)]/(2*e) -
3*b^2*(a+b*ArcCot[c*x])*PolyLog[3,1-2/(1-I*c*x)]/(2*e) +
3*I*b^3*PolyLog[4,1-2/(1-I*c*x)]/(4*e) +
(a+b*ArcCot[c*x])^3*Log[2*c*(d+e*x)/((c*d+I*e)*(1-I*c*x))]/e +
3*I*b*(a+b*ArcCot[c*x])^2*PolyLog[2,1-2*c*(d+e*x)/((c*d+I*e)*(1-I*c*x))]/(2*e) +
3*b^2*(a+b*ArcCot[c*x])*PolyLog[3,1-2*c*(d+e*x)/((c*d+I*e)*(1-I*c*x))]/(2*e) -
3*I*b^3*PolyLog[4,1-2*c*(d+e*x)/((c*d+I*e)*(1-I*c*x))]/(4*e) /;
FreeQ[{a,b,c,d,e},x] && NeQ[c^2*d^2+e^2,0]

```

2: $\int (d+e x)^q (a+b \operatorname{ArcTan}[c x]) dx$ when $q \neq -1$

Derivation: Integration by parts

Rule: If $q \neq -1$, then

$$\int (d+e x)^q (a+b \operatorname{ArcTan}[c x]) dx \rightarrow \frac{(d+e x)^{q+1} (a+b \operatorname{ArcTan}[c x])}{e (q+1)} - \frac{b c}{e (q+1)} \int \frac{(d+e x)^{q+1}}{1+c^2 x^2} dx$$

Program code:

```
Int[(d+_e*_x_)^q_.*(a+_b_.*ArcTan[c*_x_]),x_Symbol] :=
  (d+e*x)^(q+1)*(a+b*ArcTan[c*x])/(e*(q+1)) -
  b*c/(e*(q+1))*Int[(d+e*x)^(q+1)/(1+c^2*x^2),x] /;
FreeQ[{a,b,c,d,e,q},x] && NeQ[q,-1]
```

```
Int[(d+_e*_x_)^q_.*(a+_b_.*ArcCot[c*_x_]),x_Symbol] :=
  (d+e*x)^(q+1)*(a+b*ArcCot[c*x])/(e*(q+1)) +
  b*c/(e*(q+1))*Int[(d+e*x)^(q+1)/(1+c^2*x^2),x] /;
FreeQ[{a,b,c,d,e,q},x] && NeQ[q,-1]
```

3: $\int (d+ex)^q (a+b \operatorname{ArcTan}[cx])^p dx$ when $p-1 \in \mathbb{Z}^+ \wedge q \in \mathbb{Z} \wedge q \neq -1$

Derivation: Integration by parts

Rule: If $p-1 \in \mathbb{Z}^+ \wedge q \in \mathbb{Z} \wedge q \neq -1$, then

$$\int (d+ex)^q (a+b \operatorname{ArcTan}[cx])^p dx \rightarrow \frac{(d+ex)^{q+1} (a+b \operatorname{ArcTan}[cx])^p}{e(q+1)} - \frac{bc p}{e(q+1)} \int (a+b \operatorname{ArcTan}[cx])^{p-1} \operatorname{ExpandIntegrand}\left[\frac{(d+ex)^{q+1}}{1+c^2 x^2}, x\right] dx$$

Program code:

```
Int[(d_+e_.*x_)^q_.*(a_.+b_.*ArcTan[c_.*x_])^p_,x_Symbol] :=
  (d+e*x)^(q+1)*(a+b*ArcTan[c*x])^p/(e*(q+1)) -
  b*c*p/(e*(q+1))*Int[ExpandIntegrand[(a+b*ArcTan[c*x])^(p-1), (d+e*x)^(q+1)/(1+c^2*x^2), x], x] /;
FreeQ[{a,b,c,d,e},x] && IGtQ[p,1] && IntegerQ[q] && NeQ[q,-1]
```

```
Int[(d_+e_.*x_)^q_.*(a_.+b_.*ArcCot[c_.*x_])^p_,x_Symbol] :=
  (d+e*x)^(q+1)*(a+b*ArcCot[c*x])^p/(e*(q+1)) +
  b*c*p/(e*(q+1))*Int[ExpandIntegrand[(a+b*ArcCot[c*x])^(p-1), (d+e*x)^(q+1)/(1+c^2*x^2), x], x] /;
FreeQ[{a,b,c,d,e},x] && IGtQ[p,1] && IntegerQ[q] && NeQ[q,-1]
```

4. $\int (f x)^m (d + e x)^q (a + b \operatorname{ArcTan}[c x])^p dx$ when $p \in \mathbb{Z}^+$

1. $\int \frac{(f x)^m (a + b \operatorname{ArcTan}[c x])^p}{d + e x} dx$ when $p \in \mathbb{Z}^+ \wedge c^2 d^2 + e^2 = 0$

1: $\int \frac{(f x)^m (a + b \operatorname{ArcTan}[c x])^p}{d + e x} dx$ when $p \in \mathbb{Z}^+ \wedge c^2 d^2 + e^2 = 0 \wedge m > 0$

Derivation: Algebraic expansion

Basis: $\frac{x}{d+e x} = \frac{1}{e} - \frac{d}{e (d+e x)}$

Rule: If $p \in \mathbb{Z}^+ \wedge c^2 d^2 + e^2 = 0 \wedge m > 0$, then

$$\int \frac{(f x)^m (a + b \operatorname{ArcTan}[c x])^p}{d + e x} dx \rightarrow \frac{f}{e} \int (f x)^{m-1} (a + b \operatorname{ArcTan}[c x])^p dx - \frac{d f}{e} \int \frac{(f x)^{m-1} (a + b \operatorname{ArcTan}[c x])^p}{d + e x} dx$$

Program code:

```
Int[(f_.*x_)^m_.*(a_.+b_.*ArcTan[c_.*x_])^p_./(d_+e_.*x_),x_Symbol] :=
  f/e*Int[(f*x)^(m-1)*(a+b*ArcTan[c*x])^p,x] -
  d*f/e*Int[(f*x)^(m-1)*(a+b*ArcTan[c*x])^p/(d+e*x),x] /;
FreeQ[{a,b,c,d,e,f},x] && IGtQ[p,0] && EqQ[c^2*d^2+e^2,0] && GtQ[m,0]
```

```
Int[(f_.*x_)^m_.*(a_.+b_.*ArcCot[c_.*x_])^p_./(d_+e_.*x_),x_Symbol] :=
  f/e*Int[(f*x)^(m-1)*(a+b*ArcCot[c*x])^p,x] -
  d*f/e*Int[(f*x)^(m-1)*(a+b*ArcCot[c*x])^p/(d+e*x),x] /;
FreeQ[{a,b,c,d,e,f},x] && IGtQ[p,0] && EqQ[c^2*d^2+e^2,0] && GtQ[m,0]
```

$$2. \int \frac{(f x)^m (a + b \operatorname{ArcTan}[c x])^p}{d + e x} dx \text{ when } p \in \mathbb{Z}^+ \wedge c^2 d^2 + e^2 = 0 \wedge m < 0$$

$$1: \int \frac{(a + b \operatorname{ArcTan}[c x])^p}{x (d + e x)} dx \text{ when } p \in \mathbb{Z}^+ \wedge c^2 d^2 + e^2 = 0$$

Derivation: Integration by parts

$$\text{Basis: } \frac{1}{x (d + e x)} = \frac{1}{d} \partial_x \operatorname{Log} \left[2 - \frac{2}{1 + \frac{e x}{d}} \right]$$

Rule: If $p \in \mathbb{Z}^+ \wedge c^2 d^2 + e^2 = 0$, then

$$\int \frac{(a + b \operatorname{ArcTan}[c x])^p}{x (d + e x)} dx \rightarrow \frac{(a + b \operatorname{ArcTan}[c x])^p \operatorname{Log} \left[2 - \frac{2}{1 + \frac{e x}{d}} \right]}{d} - \frac{b c p}{d} \int \frac{(a + b \operatorname{ArcTan}[c x])^{p-1} \operatorname{Log} \left[2 - \frac{2}{1 + \frac{e x}{d}} \right]}{1 + c^2 x^2} dx$$

Program code:

```
Int[(a_.+b_.*ArcTan[c_.*x_])^p_./(x_*(d_+e_.*x_)),x_Symbol] :=
  (a+b*ArcTan[c*x])^p*Log[2-2/(1+e*x/d)]/d -
  b*c*p/d*Int[(a+b*ArcTan[c*x])^(p-1)*Log[2-2/(1+e*x/d)]/(1+c^2*x^2),x] /;
FreeQ[{a,b,c,d,e},x] && IGtQ[p,0] && EqQ[c^2*d^2+e^2,0]
```

```
Int[(a_.+b_.*ArcCot[c_.*x_])^p_./(x_*(d_+e_.*x_)),x_Symbol] :=
  (a+b*ArcCot[c*x])^p*Log[2-2/(1+e*x/d)]/d +
  b*c*p/d*Int[(a+b*ArcCot[c*x])^(p-1)*Log[2-2/(1+e*x/d)]/(1+c^2*x^2),x] /;
FreeQ[{a,b,c,d,e},x] && IGtQ[p,0] && EqQ[c^2*d^2+e^2,0]
```

$$\text{2: } \int \frac{(f x)^m (a + b \operatorname{ArcTan}[c x])^p}{d + e x} dx \text{ when } p \in \mathbb{Z}^+ \wedge c^2 d^2 + e^2 = 0 \wedge m < -1$$

Derivation: Algebraic expansion

$$\text{Basis: } \frac{1}{d+e x} = \frac{1}{d} - \frac{e x}{d (d+e x)}$$

Rule: If $p \in \mathbb{Z}^+ \wedge c^2 d^2 + e^2 = 0 \wedge m < -1$, then

$$\int \frac{(f x)^m (a + b \operatorname{ArcTan}[c x])^p}{d + e x} dx \rightarrow \frac{1}{d} \int (f x)^m (a + b \operatorname{ArcTan}[c x])^p dx - \frac{e}{d f} \int \frac{(f x)^{m+1} (a + b \operatorname{ArcTan}[c x])^p}{d + e x} dx$$

Program code:

```
Int[(f_.*x_)^m_*(a_.*b_.*ArcTan[c_.*x_])^p_./(d_+e_.*x_),x_Symbol] :=
  1/d*Int[(f*x)^m*(a+b*ArcTan[c*x])^p,x] -
  e/(d*f)*Int[(f*x)^(m+1)*(a+b*ArcTan[c*x])^p/(d+e*x),x] /;
FreeQ[{a,b,c,d,e,f},x] && IGtQ[p,0] && EqQ[c^2*d^2+e^2,0] && LtQ[m,-1]
```

```
Int[(f_.*x_)^m_*(a_.*b_.*ArcCot[c_.*x_])^p_./(d_+e_.*x_),x_Symbol] :=
  1/d*Int[(f*x)^m*(a+b*ArcCot[c*x])^p,x] -
  e/(d*f)*Int[(f*x)^(m+1)*(a+b*ArcCot[c*x])^p/(d+e*x),x] /;
FreeQ[{a,b,c,d,e,f},x] && IGtQ[p,0] && EqQ[c^2*d^2+e^2,0] && LtQ[m,-1]
```

2: $\int (f x)^m (d + e x)^q (a + b \operatorname{ArcTan}[c x]) dx$ when $q \neq -1 \wedge 2m \in \mathbb{Z} \wedge ((m | q) \in \mathbb{Z}^+ \vee m + q + 1 \in \mathbb{Z}^- \wedge m q < 0)$

Derivation: Integration by parts

Rule: If $q \neq -1 \wedge 2m \in \mathbb{Z} \wedge ((m | q) \in \mathbb{Z}^+ \vee m + q + 1 \in \mathbb{Z}^- \wedge m q < 0)$, let $u \rightarrow \int (f x)^m (d + e x)^q dx$, then

$$\int (f x)^m (d + e x)^q (a + b \operatorname{ArcTan}[c x]) dx \rightarrow u (a + b \operatorname{ArcTan}[c x]) - b c \int \frac{u}{1 + c^2 x^2} dx$$

Program code:

```
Int[(f_.*x_)^m_.*(d_+e_.*x_)^q_.*(a_+b_.*ArcTan[c_.*x_]),x_Symbol] :=
  With[{u=IntHide[(f*x)^m*(d+e*x)^q,x]},
    Dist[(a+b*ArcTan[c*x]),u] - b*c*Int[SimplifyIntegrand[u/(1+c^2*x^2),x],x] /;
    FreeQ[{a,b,c,d,e,f,q},x] && NeQ[q,-1] && IntegerQ[2*m] && (IGtQ[m,0] && IGtQ[q,0] || ILtQ[m+q+1,0] && LtQ[m*q,0])
```

```
Int[(f_.*x_)^m_.*(d_+e_.*x_)^q_.*(a_+b_.*ArcCot[c_.*x_]),x_Symbol] :=
  With[{u=IntHide[(f*x)^m*(d+e*x)^q,x]},
    Dist[(a+b*ArcCot[c*x]),u] + b*c*Int[SimplifyIntegrand[u/(1+c^2*x^2),x],x] /;
    FreeQ[{a,b,c,d,e,f,q},x] && NeQ[q,-1] && IntegerQ[2*m] && (IGtQ[m,0] && IGtQ[q,0] || ILtQ[m+q+1,0] && LtQ[m*q,0])
```


3: $\int (f x)^m (d + e x)^q (a + b \operatorname{ArcTan}[c x])^p dx$ when $p - 1 \in \mathbb{Z}^+ \wedge c^2 d^2 + e^2 = 0 \wedge (m | q) \in \mathbb{Z} \wedge q \neq -1$

Derivation: Integration by parts

Rule: If $p - 1 \in \mathbb{Z}^+ \wedge c^2 d^2 + e^2 = 0 \wedge (m | q) \in \mathbb{Z} \wedge q \neq -1$, let $u \rightarrow \int (f x)^m (d + e x)^q dx$, then

$$\int (f x)^m (d + e x)^q (a + b \operatorname{ArcTan}[c x])^p dx \rightarrow$$

$$u (a + b \operatorname{ArcTan}[c x])^p - b c p \int (a + b \operatorname{ArcTan}[c x])^{p-1} \operatorname{ExpandIntegrand}\left[\frac{u}{1 + c^2 x^2}, x\right] dx$$

Program code:

```
Int[(f_.*x_)^m_.*(d_.+e_.*x_)^q_.*(a_.+b_.*ArcTan[c_.*x_])^p_,x_Symbol] :=
  With[{u=IntHide[(f*x)^m*(d+e*x)^q,x]},
    Dist[(a+b*ArcTan[c*x])^p,u] - b*c*p*Int[ExpandIntegrand[(a+b*ArcTan[c*x])^(p-1),u/(1+c^2*x^2),x],x] /;
  FreeQ[{a,b,c,d,e,f,q},x] && IGtQ[p,1] && EqQ[c^2*d^2+e^2,0] && IntegersQ[m,q] && NeQ[m,-1] && NeQ[q,-1] && ILtQ[m+q+1,0] && LtQ[m*q,0]
```

```
Int[(f_.*x_)^m_.*(d_.+e_.*x_)^q_.*(a_.+b_.*ArcCot[c_.*x_])^p_,x_Symbol] :=
  With[{u=IntHide[(f*x)^m*(d+e*x)^q,x]},
    Dist[(a+b*ArcCot[c*x])^p,u] + b*c*p*Int[ExpandIntegrand[(a+b*ArcCot[c*x])^(p-1),u/(1+c^2*x^2),x],x] /;
  FreeQ[{a,b,c,d,e,f,q},x] && IGtQ[p,1] && EqQ[c^2*d^2+e^2,0] && IntegersQ[m,q] && NeQ[m,-1] && NeQ[q,-1] && ILtQ[m+q+1,0] && LtQ[m*q,0]
```

4: $\int (f x)^m (d + e x)^q (a + b \operatorname{ArcTan}[c x])^p dx$ when $p \in \mathbb{Z}^+ \wedge q \in \mathbb{Z} \wedge (q > 0 \vee a \neq 0 \vee m \in \mathbb{Z})$

Rule: If $p \in \mathbb{Z}^+ \wedge q \in \mathbb{Z} \wedge (q > 0 \vee a \neq 0 \vee m \in \mathbb{Z})$, then

$$\int (f x)^m (d + e x)^q (a + b \operatorname{ArcTan}[c x])^p dx \rightarrow \int (a + b \operatorname{ArcTan}[c x])^p \operatorname{ExpandIntegrand}[(f x)^m (d + e x)^q, x] dx$$

Program code:

```
Int[(f_.*x_)^m_.*(d_+e_.*x_)^q_.*(a_+b_.*ArcTan[c_.*x_])^p_.,x_Symbol] :=
  Int[ExpandIntegrand[(a+b*ArcTan[c*x])^p,(f*x)^m*(d+e*x)^q,x],x] /;
FreeQ[{a,b,c,d,e,f,m},x] && IGtQ[p,0] && IntegerQ[q] && (GtQ[q,0] || NeQ[a,0] || IntegerQ[m])
```

```
Int[(f_.*x_)^m_.*(d_+e_.*x_)^q_.*(a_+b_.*ArcCot[c_.*x_])^p_.,x_Symbol] :=
  Int[ExpandIntegrand[(a+b*ArcCot[c*x])^p,(f*x)^m*(d+e*x)^q,x],x] /;
FreeQ[{a,b,c,d,e,f,m},x] && IGtQ[p,0] && IntegerQ[q] && (GtQ[q,0] || NeQ[a,0] || IntegerQ[m])
```

$$5. \int (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx$$

$$1. \int (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx \text{ when } e = c^2 d$$

$$1. \int (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx \text{ when } e = c^2 d \wedge q > 0$$

$$\text{1: } \int (d + e x^2)^q (a + b \operatorname{ArcTan}[c x]) dx \text{ when } e = c^2 d \wedge q > 0$$

Rule: If $e = c^2 d \wedge q > 0$, then

$$\int (d + e x^2)^q (a + b \operatorname{ArcTan}[c x]) dx \rightarrow -\frac{b (d + e x^2)^q}{2 c q (2 q + 1)} + \frac{x (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])}{2 q + 1} + \frac{2 d q}{2 q + 1} \int (d + e x^2)^{q-1} (a + b \operatorname{ArcTan}[c x]) dx$$

Program code:

```
Int[(d_+e_.*x_^2)^q_.*(a_.+b_.*ArcTan[c_.*x_]),x_Symbol] :=
  -b*(d+e*x^2)^q/(2*c*q*(2*q+1)) +
  x*(d+e*x^2)^q*(a+b*ArcTan[c*x])/(2*q+1) +
  2*d*q/(2*q+1)*Int[(d+e*x^2)^(q-1)*(a+b*ArcTan[c*x]),x] /;
FreeQ[{a,b,c,d,e},x] && EqQ[e,c^2*d] && GtQ[q,0]
```

```
Int[(d_+e_.*x_^2)^q_.*(a_.+b_.*ArcCot[c_.*x_]),x_Symbol] :=
  b*(d+e*x^2)^q/(2*c*q*(2*q+1)) +
  x*(d+e*x^2)^q*(a+b*ArcCot[c*x])/(2*q+1) +
  2*d*q/(2*q+1)*Int[(d+e*x^2)^(q-1)*(a+b*ArcCot[c*x]),x] /;
FreeQ[{a,b,c,d,e},x] && EqQ[e,c^2*d] && GtQ[q,0]
```

$$\text{2: } \int (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx \text{ when } e = c^2 d \wedge q > 0 \wedge p > 1$$

Rule: If $e = c^2 d \wedge q > 0 \wedge p > 1$, then

$$\int (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx \rightarrow$$

$$-\frac{b p (d+e x^2)^q (a+b \operatorname{ArcTan}[c x])^{p-1}}{2 c q (2 q+1)} + \frac{x (d+e x^2)^q (a+b \operatorname{ArcTan}[c x])^p}{2 q+1} +$$

$$\frac{2 d q}{2 q+1} \int (d+e x^2)^{q-1} (a+b \operatorname{ArcTan}[c x])^p dx + \frac{b^2 d p (p-1)}{2 q (2 q+1)} \int (d+e x^2)^{q-1} (a+b \operatorname{ArcTan}[c x])^{p-2} dx$$

Program code:

```
Int[(d+_e_.*x_^2)^q_.*(a_+_b_.*ArcTan[c_.*x_])^p_,x_Symbol] :=
  -b*p*(d+e*x^2)^q*(a+b*ArcTan[c*x])^(p-1)/(2*c*q*(2*q+1)) +
  x*(d+e*x^2)^q*(a+b*ArcTan[c*x])^p/(2*q+1) +
  2*d*q/(2*q+1)*Int[(d+e*x^2)^(q-1)*(a+b*ArcTan[c*x])^p,x] +
  b^2*d*p*(p-1)/(2*q*(2*q+1))*Int[(d+e*x^2)^(q-1)*(a+b*ArcTan[c*x])^(p-2),x] /;
FreeQ[{a,b,c,d,e},x] && EqQ[e,c^2*d] && GtQ[q,0] && GtQ[p,1]
```

```
Int[(d+_e_.*x_^2)^q_.*(a_+_b_.*ArcCot[c_.*x_])^p_,x_Symbol] :=
  b*p*(d+e*x^2)^q*(a+b*ArcCot[c*x])^(p-1)/(2*c*q*(2*q+1)) +
  x*(d+e*x^2)^q*(a+b*ArcCot[c*x])^p/(2*q+1) +
  2*d*q/(2*q+1)*Int[(d+e*x^2)^(q-1)*(a+b*ArcCot[c*x])^p,x] +
  b^2*d*p*(p-1)/(2*q*(2*q+1))*Int[(d+e*x^2)^(q-1)*(a+b*ArcCot[c*x])^(p-2),x] /;
FreeQ[{a,b,c,d,e},x] && EqQ[e,c^2*d] && GtQ[q,0] && GtQ[p,1]
```

$$2. \int (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx \text{ when } e = c^2 d \wedge q < 0$$

$$1. \int \frac{(a + b \operatorname{ArcTan}[c x])^p}{d + e x^2} dx \text{ when } e = c^2 d$$

$$\text{1: } \int \frac{1}{(d + e x^2) (a + b \operatorname{ArcTan}[c x])} dx \text{ when } e = c^2 d$$

Derivation: Reciprocal rule for integration

Rule: If $e = c^2 d$, then

$$\int \frac{1}{(d + e x^2) (a + b \operatorname{ArcTan}[c x])} dx \rightarrow \frac{\operatorname{Log}[a + b \operatorname{ArcTan}[c x]]}{b c d}$$

Program code:

```
Int[1/((d_+e_.*x_^2)*(a_.+b_.*ArcTan[c_.*x_])),x_Symbol] :=
  Log[RemoveContent[a+b*ArcTan[c*x],x]]/(b*c*d) /;
FreeQ[{a,b,c,d,e},x] && EqQ[e,c^2*d]
```

```
Int[1/((d_+e_.*x_^2)*(a_.+b_.*ArcCot[c_.*x_])),x_Symbol] :=
  -Log[RemoveContent[a+b*ArcCot[c*x],x]]/(b*c*d) /;
FreeQ[{a,b,c,d,e},x] && EqQ[e,c^2*d]
```

$$\text{2: } \int \frac{(a + b \operatorname{ArcTan}[c x])^p}{d + e x^2} dx \text{ when } e = c^2 d \wedge p \neq -1$$

Derivation: Power rule for integration

Rule: If $e = c^2 d \wedge p \neq -1$, then

$$\int \frac{(a + b \operatorname{ArcTan}[c x])^p}{d + e x^2} dx \rightarrow \frac{(a + b \operatorname{ArcTan}[c x])^{p+1}}{b c d (p+1)}$$

Program code:

```
Int[(a_.+b_.*ArcTan[c_.*x_])^p_./(d_+e_.*x_^2),x_Symbol] :=
  (a+b*ArcTan[c*x])^(p+1)/(b*c*d*(p+1)) /;
FreeQ[{a,b,c,d,e,p},x] && EqQ[e,c^2*d] && NeQ[p,-1]
```

```
Int[(a_.+b_.*ArcCot[c_.*x_])^p_./(d_+e_.*x_^2),x_Symbol] :=
  -(a+b*ArcCot[c*x])^(p+1)/(b*c*d*(p+1)) /;
FreeQ[{a,b,c,d,e,p},x] && EqQ[e,c^2*d] && NeQ[p,-1]
```

$$\begin{aligned} &2. \int \frac{(a + b \operatorname{ArcTan}[c x])^p}{\sqrt{d + e x^2}} dx \text{ when } e = c^2 d \wedge p \in \mathbb{Z}^+ \\ &1. \int \frac{(a + b \operatorname{ArcTan}[c x])^p}{\sqrt{d + e x^2}} dx \text{ when } e = c^2 d \wedge n \in \mathbb{Z}^+ \wedge d > 0 \\ &\text{1: } \int \frac{(a + b \operatorname{ArcTan}[c x])}{\sqrt{d + e x^2}} dx \text{ when } e = c^2 d \wedge d > 0 \end{aligned}$$

Derivation: Integration by substitution and algebraic simplification

Note: Although not essential, these rules return antiderivatives free of complex exponentials of the form $i e^{\operatorname{ArcTan}[c x]}$ and $e^{\operatorname{ArcCot}[c x]}$.

Basis: If $e = c^2 d \wedge d > 0$, then $\frac{1}{\sqrt{d+e x^2}} = \frac{1}{c \sqrt{d}} \operatorname{Sec}[\operatorname{ArcTan}[c x]] \partial_x \operatorname{ArcTan}[c x]$

Basis: If $e = c^2 d \wedge d > 0$, then $\frac{1}{\sqrt{d+e x^2}} = -\frac{1}{c \sqrt{d}} \sqrt{\operatorname{Csc}[\operatorname{ArcCot}[c x]]^2} \partial_x \operatorname{ArcCot}[c x]$

Rule: If $e = c^2 d \wedge d > 0$, then

$$\int \frac{a + b \operatorname{ArcTan}[c x]}{\sqrt{d + e x^2}} dx \rightarrow \frac{1}{c \sqrt{d}} \operatorname{Subst}[(a + b x) \operatorname{Sec}[x], x, \operatorname{ArcTan}[c x]]$$

$$\rightarrow -\frac{2 i (a + b \operatorname{ArcTan}[c x]) \operatorname{ArcTan}\left[\frac{\sqrt{1+i c x}}{\sqrt{1-i c x}}\right]}{c \sqrt{d}} + \frac{i b \operatorname{PolyLog}\left[2, -\frac{i \sqrt{1+i c x}}{\sqrt{1-i c x}}\right]}{c \sqrt{d}} - \frac{i b \operatorname{PolyLog}\left[2, \frac{i \sqrt{1+i c x}}{\sqrt{1-i c x}}\right]}{c \sqrt{d}}$$

Program code:

```
Int[(a_.+b_.*ArcTan[c_.*x_])/Sqrt[d_+e_.*x_^2],x_Symbol] :=
-2*I*(a+b*ArcTan[c*x])*ArcTan[Sqrt[1+I*c*x]/Sqrt[1-I*c*x]]/(c*Sqrt[d]) +
I*b*PolyLog[2,-I*Sqrt[1+I*c*x]/Sqrt[1-I*c*x]]/(c*Sqrt[d]) -
I*b*PolyLog[2,I*Sqrt[1+I*c*x]/Sqrt[1-I*c*x]]/(c*Sqrt[d]) /;
FreeQ[{a,b,c,d,e},x] && EqQ[e,c^2*d] && GtQ[d,0]
```

```
Int[(a_.+b_.*ArcCot[c_.*x_])/Sqrt[d_+e_.*x_^2],x_Symbol] :=
-2*I*(a+b*ArcCot[c*x])*ArcTan[Sqrt[1+I*c*x]/Sqrt[1-I*c*x]]/(c*Sqrt[d]) -
I*b*PolyLog[2,-I*Sqrt[1+I*c*x]/Sqrt[1-I*c*x]]/(c*Sqrt[d]) +
I*b*PolyLog[2,I*Sqrt[1+I*c*x]/Sqrt[1-I*c*x]]/(c*Sqrt[d]) /;
FreeQ[{a,b,c,d,e},x] && EqQ[e,c^2*d] && GtQ[d,0]
```

$$2. \int \frac{(a + b \operatorname{ArcTan}[c x])^p}{\sqrt{d + e x^2}} dx \text{ when } e = c^2 d \wedge p \in \mathbb{Z}^+ \wedge d > 0$$

$$1: \int \frac{(a + b \operatorname{ArcTan}[c x])^p}{\sqrt{d + e x^2}} dx \text{ when } e = c^2 d \wedge p \in \mathbb{Z}^+ \wedge d > 0$$

Derivation: Integration by substitution

Basis: If $e == c^2 d \wedge d > 0$, then $\frac{1}{\sqrt{d+e x^2}} == \frac{1}{c \sqrt{d}} \operatorname{Sec}[\operatorname{ArcTan}[c x]] \partial_x \operatorname{ArcTan}[c x]$

Rule: If $e == c^2 d \wedge p \in \mathbb{Z}^+ \wedge d > 0$, then

$$\int \frac{(a + b \operatorname{ArcTan}[c x])^p}{\sqrt{d + e x^2}} dx \rightarrow \frac{1}{c \sqrt{d}} \operatorname{Subst}\left[\int (a + b x)^p \operatorname{Sec}[x] dx, x, \operatorname{ArcTan}[c x]\right]$$

Program code:

```
Int[(a_.+b_.*ArcTan[c_.*x_])^p_/Sqrt[d_+e_.*x_^2],x_Symbol] :=
  1/(c*Sqrt[d])*Subst[Int[(a+b*x)^p*Sec[x],x],x,ArcTan[c*x] /;
  FreeQ[{a,b,c,d,e},x] && EqQ[e,c^2*d] && IGtQ[p,0] && GtQ[d,0]
```

$$\mathbf{2:} \int \frac{(a + b \operatorname{ArcCot}[c x])^p}{\sqrt{d + e x^2}} dx \text{ when } e == c^2 d \wedge p \in \mathbb{Z}^+ \wedge d > 0$$

Derivation: Integration by substitution and piecewise constant extraction

Basis: If $e == c^2 d \wedge d > 0$, then $\frac{1}{\sqrt{d+e x^2}} == -\frac{1}{c \sqrt{d}} \frac{\operatorname{Csc}[\operatorname{ArcCot}[c x]]^2}{\sqrt{\operatorname{Csc}[\operatorname{ArcCot}[c x]]^2}} \partial_x \operatorname{ArcCot}[c x]$

Basis: $\partial_x \frac{\operatorname{Csc}[x]}{\sqrt{\operatorname{Csc}[x]^2}} == 0$

Basis: $\frac{\operatorname{Csc}[\operatorname{ArcCot}[c x]]}{\sqrt{\operatorname{Csc}[\operatorname{ArcCot}[c x]]^2}} == \frac{c x \sqrt{1 + \frac{1}{c^2 x^2}}}{\sqrt{1 + c^2 x^2}}$

Rule: If $e == c^2 d \wedge p \in \mathbb{Z}^+ \wedge d > 0$, then

$$\int \frac{(a + b \operatorname{ArcCot}[c x])^p}{\sqrt{d + e x^2}} dx \rightarrow -\frac{1}{c \sqrt{d}} \operatorname{Subst}\left[\int \frac{(a + b x)^p \operatorname{Csc}[x]^2}{\sqrt{\operatorname{Csc}[x]^2}} dx, x, \operatorname{ArcCot}[c x]\right]$$

$$\rightarrow -\frac{x\sqrt{1+\frac{1}{c^2x^2}}}{\sqrt{d+ex^2}} \text{Subst}\left[\int (a+bx)^p \text{Csc}[x] dx, x, \text{ArcCot}[cx]\right]$$

Program code:

```
Int[(a_.+b_.*ArcCot[c_.*x_])^p_/Sqrt[d_+e_.*x_^2],x_Symbol] :=
  -x*Sqrt[1+1/(c^2*x^2)]/Sqrt[d+e*x^2]*Subst[Int[(a+b*x)^p*Csc[x],x],x,ArcCot[c*x] /;
FreeQ[{a,b,c,d,e},x] && EqQ[e,c^2*d] && IGtQ[p,0] && GtQ[d,0]
```

$$\mathbf{2:} \int \frac{(a+b \text{ArcTan}[cx])^p}{\sqrt{d+ex^2}} dx \text{ when } e = c^2 d \wedge p \in \mathbb{Z}^+ \wedge d \neq 0$$

Derivation: Piecewise constant extraction

Basis: If $e = c^2 d$, then $\partial_x \frac{\sqrt{1+c^2 x^2}}{\sqrt{d+ex^2}} = 0$

Rule: If $e = c^2 d \wedge p \in \mathbb{Z}^+ \wedge d \neq 0$, then

$$\int \frac{(a+b \text{ArcTan}[cx])^p}{\sqrt{d+ex^2}} dx \rightarrow \frac{\sqrt{1+c^2 x^2}}{\sqrt{d+ex^2}} \int \frac{(a+b \text{ArcTan}[cx])^p}{\sqrt{1+c^2 x^2}} dx$$

Program code:

```
Int[(a_.+b_.*ArcTan[c_.*x_])^p_/Sqrt[d_+e_.*x_^2],x_Symbol] :=
  Sqrt[1+c^2*x^2]/Sqrt[d+e*x^2]*Int[(a+b*ArcTan[c*x])^p/Sqrt[1+c^2*x^2],x] /;
FreeQ[{a,b,c,d,e},x] && EqQ[e,c^2*d] && IGtQ[p,0] && Not[GtQ[d,0]]
```

```
Int[(a_.+b_.*ArcCot[c_.*x_])^p_/Sqrt[d_+e_.*x_^2],x_Symbol] :=
  Sqrt[1+c^2*x^2]/Sqrt[d+e*x^2]*Int[(a+b*ArcCot[c*x])^p/Sqrt[1+c^2*x^2],x] /;
FreeQ[{a,b,c,d,e},x] && EqQ[e,c^2*d] && IGtQ[p,0] && Not[GtQ[d,0]]
```

$$3. \int (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx \text{ when } e = c^2 d \wedge q < -1$$

$$1: \int \frac{(a + b \operatorname{ArcTan}[c x])^p}{(d + e x^2)^2} dx \text{ when } e = c^2 d \wedge p > 0$$

Rule: If $e = c^2 d \wedge p > 0$, then

$$\int \frac{(a + b \operatorname{ArcTan}[c x])^p}{(d + e x^2)^2} dx \rightarrow \frac{x (a + b \operatorname{ArcTan}[c x])^p}{2 d (d + e x^2)} + \frac{(a + b \operatorname{ArcTan}[c x])^{p+1}}{2 b c d^2 (p+1)} - \frac{b c p}{2} \int \frac{x (a + b \operatorname{ArcTan}[c x])^{p-1}}{(d + e x^2)^2} dx$$

Program code:

```
Int[(a_.+b_.*ArcTan[c_.*x_])^p_./(d_+e_.*x_^2)^2,x_Symbol] :=
  x*(a+b*ArcTan[c*x])^p/(2*d*(d+e*x^2)) +
  (a+b*ArcTan[c*x])^(p+1)/(2*b*c*d^2*(p+1)) -
  b*c*p/2*Int[x*(a+b*ArcTan[c*x])^(p-1)/(d+e*x^2)^2,x] /;
FreeQ[{a,b,c,d,e},x] && EqQ[e,c^2*d] && GtQ[p,0]
```

```
Int[(a_.+b_.*ArcCot[c_.*x_])^p_./(d_+e_.*x_^2)^2,x_Symbol] :=
  x*(a+b*ArcCot[c*x])^p/(2*d*(d+e*x^2)) -
  (a+b*ArcCot[c*x])^(p+1)/(2*b*c*d^2*(p+1)) +
  b*c*p/2*Int[x*(a+b*ArcCot[c*x])^(p-1)/(d+e*x^2)^2,x] /;
FreeQ[{a,b,c,d,e},x] && EqQ[e,c^2*d] && GtQ[p,0]
```

$$2. \int (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx \text{ when } e = c^2 d \wedge q < -1 \wedge p \geq 1$$

$$1. \int (d + e x^2)^q (a + b \operatorname{ArcTan}[c x]) dx \text{ when } e = c^2 d \wedge q < -1$$

$$1: \int \frac{a + b \operatorname{ArcTan}[c x]}{(d + e x^2)^{3/2}} dx \text{ when } e = c^2 d$$

Rule: If $e = c^2 d$, then

$$\int \frac{a + b \operatorname{ArcTan}[c x]}{(d + e x^2)^{3/2}} dx \rightarrow \frac{b}{c d \sqrt{d + e x^2}} + \frac{x (a + b \operatorname{ArcTan}[c x])}{d \sqrt{d + e x^2}}$$

Program code:

```
Int[(a_.+b_.*ArcTan[c_.*x_])/(d_+e_.*x_^2)^(3/2),x_Symbol] :=
  b/(c*d*Sqrt[d+e*x^2]) +
  x*(a+b*ArcTan[c*x])/(d*Sqrt[d+e*x^2]) /;
FreeQ[{a,b,c,d,e},x] && EqQ[e,c^2*d]
```

```
Int[(a_.+b_.*ArcCot[c_.*x_])/(d_+e_.*x_^2)^(3/2),x_Symbol] :=
  -b/(c*d*Sqrt[d+e*x^2]) +
  x*(a+b*ArcCot[c*x])/(d*Sqrt[d+e*x^2]) /;
FreeQ[{a,b,c,d,e},x] && EqQ[e,c^2*d]
```

2: $\int (d + e x^2)^q (a + b \operatorname{ArcTan}[c x]) dx$ when $e = c^2 d \wedge q < -1 \wedge q \neq -\frac{3}{2}$

Rule: If $e = c^2 d \wedge q < -1 \wedge q \neq -\frac{3}{2}$, then

$$\int (d + e x^2)^q (a + b \operatorname{ArcTan}[c x]) dx \rightarrow \frac{b (d + e x^2)^{q+1}}{4 c d (q+1)^2} - \frac{x (d + e x^2)^{q+1} (a + b \operatorname{ArcTan}[c x])}{2 d (q+1)} + \frac{2 q + 3}{2 d (q+1)} \int (d + e x^2)^{q+1} (a + b \operatorname{ArcTan}[c x]) dx$$

Program code:

```
Int[(d_+e_.*x_^2)^q_*(a_.+b_.*ArcTan[c_.*x_]),x_Symbol] :=
  b*(d+e*x^2)^(q+1)/(4*c*d*(q+1)^2) -
  x*(d+e*x^2)^(q+1)*(a+b*ArcTan[c*x])/(2*d*(q+1)) +
  (2*q+3)/(2*d*(q+1))*Int[(d+e*x^2)^(q+1)*(a+b*ArcTan[c*x]),x] /;
FreeQ[{a,b,c,d,e},x] && EqQ[e,c^2*d] && LtQ[q,-1] && NeQ[q,-3/2]
```

```

Int[(d+e*x^2)^q*(a+b_*ArcCot[c_*x_]),x_Symbol] :=
  -b*(d+e*x^2)^(q+1)/(4*c*d*(q+1)^2) -
  x*(d+e*x^2)^(q+1)*(a+b_*ArcCot[c*x])/(2*d*(q+1)) +
  (2*q+3)/(2*d*(q+1))*Int[(d+e*x^2)^(q+1)*(a+b_*ArcCot[c*x]),x] /;
FreeQ[{a,b,c,d,e},x] && EqQ[e,c^2*d] && LtQ[q,-1] && NeQ[q,-3/2]

```

$$2. \int (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx \text{ when } e = c^2 d \wedge q < -1 \wedge p > 1$$

$$1: \int \frac{(a + b \operatorname{ArcTan}[c x])^p}{(d + e x^2)^{3/2}} dx \text{ when } e = c^2 d \wedge p > 1$$

Rule: If $e = c^2 d \wedge p > 1$, then

$$\int \frac{(a + b \operatorname{ArcTan}[c x])^p}{(d + e x^2)^{3/2}} dx \rightarrow \frac{b p (a + b \operatorname{ArcTan}[c x])^{p-1}}{c d \sqrt{d + e x^2}} + \frac{x (a + b \operatorname{ArcTan}[c x])^p}{d \sqrt{d + e x^2}} - b^2 p (p-1) \int \frac{(a + b \operatorname{ArcTan}[c x])^{p-2}}{(d + e x^2)^{3/2}} dx$$

Program code:

```

Int[(a_+b_*ArcTan[c_*x_])^p/(d+e_*x^2)^(3/2),x_Symbol] :=
  b*p*(a+b_*ArcTan[c*x])^(p-1)/(c*d*Sqrt[d+e*x^2]) +
  x*(a+b_*ArcTan[c*x])^p/(d*Sqrt[d+e*x^2]) -
  b^2*p*(p-1)*Int[(a+b_*ArcTan[c*x])^(p-2)/(d+e*x^2)^(3/2),x] /;
FreeQ[{a,b,c,d,e},x] && EqQ[e,c^2*d] && GtQ[p,1]

```

```

Int[(a_+b_*ArcCot[c_*x_])^p/(d+e_*x^2)^(3/2),x_Symbol] :=
  -b*p*(a+b_*ArcCot[c*x])^(p-1)/(c*d*Sqrt[d+e*x^2]) +
  x*(a+b_*ArcCot[c*x])^p/(d*Sqrt[d+e*x^2]) -
  b^2*p*(p-1)*Int[(a+b_*ArcCot[c*x])^(p-2)/(d+e*x^2)^(3/2),x] /;
FreeQ[{a,b,c,d,e},x] && EqQ[e,c^2*d] && GtQ[p,1]

```

2: $\int (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx$ when $e = c^2 d \wedge q < -1 \wedge p > 1 \wedge q \neq -\frac{3}{2}$

Rule: If $e = c^2 d \wedge q < -1 \wedge p > 1 \wedge q \neq -\frac{3}{2}$, then

$$\int (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx \rightarrow \frac{b p (d + e x^2)^{q+1} (a + b \operatorname{ArcTan}[c x])^{p-1}}{4 c d (q+1)^2} - \frac{x (d + e x^2)^{q+1} (a + b \operatorname{ArcTan}[c x])^p}{2 d (q+1)} + \frac{2 q + 3}{2 d (q+1)} \int (d + e x^2)^{q+1} (a + b \operatorname{ArcTan}[c x])^p dx - \frac{b^2 p (p-1)}{4 (q+1)^2} \int (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^{p-2} dx$$

Program code:

```
Int[(d_+e_.*x_^2)^q_*(a_.+b_.*ArcTan[c_.*x_])^p_,x_Symbol] :=
  b*p*(d+e*x^2)^(q+1)*(a+b*ArcTan[c*x])^(p-1)/(4*c*d*(q+1)^2) -
  x*(d+e*x^2)^(q+1)*(a+b*ArcTan[c*x])^p/(2*d*(q+1)) +
  (2*q+3)/(2*d*(q+1))*Int[(d+e*x^2)^(q+1)*(a+b*ArcTan[c*x])^p,x] -
  b^2*p*(p-1)/(4*(q+1)^2)*Int[(d+e*x^2)^q*(a+b*ArcTan[c*x])^(p-2),x] /;
FreeQ[{a,b,c,d,e},x] && EqQ[e,c^2*d] && LtQ[q,-1] && GtQ[p,1] && NeQ[q,-3/2]
```

```
Int[(d_+e_.*x_^2)^q_*(a_.+b_.*ArcCot[c_.*x_])^p_,x_Symbol] :=
  -b*p*(d+e*x^2)^(q+1)*(a+b*ArcCot[c*x])^(p-1)/(4*c*d*(q+1)^2) -
  x*(d+e*x^2)^(q+1)*(a+b*ArcCot[c*x])^p/(2*d*(q+1)) +
  (2*q+3)/(2*d*(q+1))*Int[(d+e*x^2)^(q+1)*(a+b*ArcCot[c*x])^p,x] -
  b^2*p*(p-1)/(4*(q+1)^2)*Int[(d+e*x^2)^q*(a+b*ArcCot[c*x])^(p-2),x] /;
FreeQ[{a,b,c,d,e},x] && EqQ[e,c^2*d] && LtQ[q,-1] && GtQ[p,1] && NeQ[q,-3/2]
```

$$\mathbf{3:} \int (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx \text{ when } e = c^2 d \wedge q < -1 \wedge p < -1$$

Derivation: Integration by parts

$$\text{Basis: If } e = c^2 d, \text{ then } \frac{(a+b \operatorname{ArcTan}[c x])^p}{d+e x^2} = \partial_x \frac{(a+b \operatorname{ArcTan}[c x])^{p+1}}{b c d (p+1)}$$

Rule: If $e = c^2 d \wedge q < -1 \wedge p < -1$, then

$$\int (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx \rightarrow \frac{(d + e x^2)^{q+1} (a + b \operatorname{ArcTan}[c x])^{p+1}}{b c d (p+1)} - \frac{2 c (q+1)}{b (p+1)} \int x (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^{p+1} dx$$

Program code:

```
Int[(d_+e_.*x_^2)^q_*(a_+b_.*ArcTan[c_.*x_])^p_,x_Symbol] :=
  (d+e*x^2)^(q+1)*(a+b*ArcTan[c*x])^(p+1)/(b*c*d*(p+1)) -
  2*c*(q+1)/(b*(p+1))*Int[x*(d+e*x^2)^q*(a+b*ArcTan[c*x])^(p+1),x] /;
FreeQ[{a,b,c,d,e},x] && EqQ[e,c^2*d] && LtQ[q,-1] && LtQ[p,-1]
```

```
Int[(d_+e_.*x_^2)^q_*(a_+b_.*ArcCot[c_.*x_])^p_,x_Symbol] :=
  -(d+e*x^2)^(q+1)*(a+b*ArcCot[c*x])^(p+1)/(b*c*d*(p+1)) +
  2*c*(q+1)/(b*(p+1))*Int[x*(d+e*x^2)^q*(a+b*ArcCot[c*x])^(p+1),x] /;
FreeQ[{a,b,c,d,e},x] && EqQ[e,c^2*d] && LtQ[q,-1] && LtQ[p,-1]
```

$$4. \int (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx \text{ when } e = c^2 d \wedge 2(q+1) \in \mathbb{Z}^-$$

$$1. \int (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx \text{ when } e = c^2 d \wedge 2(q+1) \in \mathbb{Z}^-$$

$$\mathbf{1:} \int (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx \text{ when } e = c^2 d \wedge 2(q+1) \in \mathbb{Z}^- \wedge (q \in \mathbb{Z} \vee d > 0)$$

Derivation: Integration by substitution

Basis: If $e = c^2 d \wedge 2(q+1) \in \mathbb{Z} \wedge (q \in \mathbb{Z} \vee d > 0)$, then $(d + e x^2)^q = \frac{d^q}{c \cos[\operatorname{ArcTan}[c x]]^{2(q+1)}} \partial_x \operatorname{ArcTan}[c x]$

Rule: If $e = c^2 d \wedge 2(q+1) \in \mathbb{Z}^- \wedge (q \in \mathbb{Z} \vee d > 0)$, then

$$\int (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx \rightarrow \frac{d^q}{c} \operatorname{Subst}\left[\int \frac{(a + b x)^p}{\cos[x]^{2(q+1)}} dx, x, \operatorname{ArcTan}[c x]\right]$$

Program code:

```
Int[(d_+e_.*x_^2)^q_*(a_.+b_.*ArcTan[c_.*x_])^p_.,x_Symbol] :=
  d^q/c*Subst[Int[(a+b*x)^p/Cos[x]^(2*(q+1)),x],x,ArcTan[c*x]] /;
FreeQ[{a,b,c,d,e,p},x] && EqQ[e,c^2*d] && ILtQ[2*(q+1),0] && (IntegerQ[q] || GtQ[d,0])
```

$$\mathbf{2:} \int (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx \text{ when } e = c^2 d \wedge 2 (q + 1) \in \mathbb{Z}^- \wedge \neg (q \in \mathbb{Z} \vee d > 0)$$

Derivation: Piecewise constant extraction

$$\text{Basis: If } e = c^2 d, \text{ then } \partial_x \frac{\sqrt{1+c^2 x^2}}{\sqrt{d+e x^2}} = 0$$

Rule: If $e = c^2 d \wedge 2 (q + 1) \in \mathbb{Z}^- \wedge \neg (q \in \mathbb{Z} \vee d > 0)$, then

$$\int (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx \rightarrow \frac{d^{q+\frac{1}{2}} \sqrt{1+c^2 x^2}}{\sqrt{d+e x^2}} \int (1+c^2 x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx$$

Program code:

```
Int[(d_+e_.*x_^2)^q_*(a_.+b_.*ArcTan[c_.*x_])^p_.,x_Symbol] :=
  d^(q+1/2)*Sqrt[1+c^2*x^2]/Sqrt[d+e*x^2]*Int[(1+c^2*x^2)^q*(a+b*ArcTan[c*x])^p,x] /;
FreeQ[{a,b,c,d,e,p},x] && EqQ[e,c^2*d] && ILtQ[2*(q+1),0] && Not[IntegerQ[q] || GtQ[d,0]]
```


$$2. \int (d + e x^2)^q (a + b \operatorname{ArcCot}[c x])^p dx \text{ when } e = c^2 d \wedge 2(q+1) \in \mathbb{Z}^-$$

$$1: \int (d + e x^2)^q (a + b \operatorname{ArcCot}[c x])^p dx \text{ when } e = c^2 d \wedge 2(q+1) \in \mathbb{Z}^- \wedge q \in \mathbb{Z}$$

Derivation: Integration by substitution

$$\text{Basis: If } e = c^2 d \wedge q \in \mathbb{Z}, \text{ then } (d + e x^2)^q = - \frac{d^q}{c \operatorname{Sin}[\operatorname{ArcCot}[c x]]^{2(q+1)}} \partial_x \operatorname{ArcCot}[c x]$$

Rule: If $e = c^2 d \wedge 2(q+1) \in \mathbb{Z}^- \wedge q \in \mathbb{Z}$, then

$$\int (d + e x^2)^q (a + b \operatorname{ArcCot}[c x])^p dx \rightarrow - \frac{d^q}{c} \operatorname{Subst} \left[\int \frac{(a + b x)^p}{\operatorname{Sin}[x]^{2(q+1)}} dx, x, \operatorname{ArcCot}[c x] \right]$$

Program code:

```
Int[(d_+e_.*x_^2)^q_*(a_.*b_.*ArcCot[c_.*x_])^p_,x_Symbol] :=
  -d^q/c*Subst[Int[(a+b*x)^p/Sin[x]^(2*(q+1)),x],x,ArcCot[c*x] /;
  FreeQ[{a,b,c,d,e,p},x] && EqQ[e,c^2*d] && ILtQ[2*(q+1),0] && IntegerQ[q]
```

$$2: \int (d + e x^2)^q (a + b \operatorname{ArcCot}[c x])^p dx \text{ when } e = c^2 d \wedge 2(q+1) \in \mathbb{Z}^- \wedge q \notin \mathbb{Z}$$

Derivation: Piecewise constant extraction and integration by substitution

$$\text{Basis: If } e = c^2 d, \text{ then } \partial_x \frac{x \sqrt{\frac{1+c^2 x^2}{c^2 x^2}}}{\sqrt{d+e x^2}} = 0$$

$$\text{Basis: If } 2(q+1) \in \mathbb{Z} \wedge q \notin \mathbb{Z}, \text{ then } x \sqrt{1 + \frac{1}{c^2 x^2}} (1 + c^2 x^2)^{q-\frac{1}{2}} = - \frac{1}{c^2 \operatorname{Sin}[\operatorname{ArcCot}[c x]]^{2(q+1)}} \partial_x \operatorname{ArcCot}[c x]$$

Rule: If $e = c^2 d \wedge 2(q+1) \in \mathbb{Z}^- \wedge q \notin \mathbb{Z}$, then

$$\int (d + e x^2)^q (a + b \operatorname{ArcCot}[c x])^p dx \rightarrow \frac{c^2 d^{q+\frac{1}{2}} x \sqrt{\frac{1+c^2 x^2}{c^2 x^2}}}{\sqrt{d + e x^2}} \int x \sqrt{1 + \frac{1}{c^2 x^2}} (1 + c^2 x^2)^{q-\frac{1}{2}} (a + b \operatorname{ArcCot}[c x])^p dx$$

$$\rightarrow - \frac{d^{q+\frac{1}{2}} x \sqrt{\frac{1+c^2 x^2}{c^2 x^2}}}{\sqrt{d + e x^2}} \operatorname{Subst}\left[\int \frac{(a + b x)^p}{\operatorname{Sin}[x]^{2(q+1)}} dx, x, \operatorname{ArcCot}[c x]\right]$$

Program code:

```
Int[(d_+e_.*x_^2)^q_*(a_.+b_.*ArcCot[c_.*x_])^p_,x_Symbol] :=
  -d^(q+1/2)*x*Sqrt[(1+c^2*x^2)/(c^2*x^2)]/Sqrt[d+e*x^2]*Subst[Int[(a+b*x)^p/Sin[x]^(2*(q+1)),x],x,ArcCot[c*x] /;
  FreeQ[{a,b,c,d,e,p},x] && EqQ[e,c^2*d] && ILtQ[2*(q+1),0] && Not[IntegerQ[q]]
```

$$2. \int \frac{a + b \operatorname{ArcTan}[c x]}{d + e x^2} dx$$

$$1: \int \frac{\operatorname{ArcTan}[c x]}{d + e x^2} dx$$

Derivation: Algebraic expansion

$$\text{Basis: } \operatorname{ArcTan}[z] == \frac{1}{2} i \operatorname{Log}[1 - i z] - \frac{1}{2} i \operatorname{Log}[1 + i z]$$

$$\text{Basis: } \operatorname{ArcCot}[z] == \frac{1}{2} i \operatorname{Log}\left[1 - \frac{i}{z}\right] - \frac{1}{2} i \operatorname{Log}\left[1 + \frac{i}{z}\right]$$

Rule:

$$\int \frac{\operatorname{ArcTan}[c x]}{d + e x^2} dx \rightarrow \frac{i}{2} \int \frac{\operatorname{Log}[1 - i c x]}{d + e x^2} dx - \frac{i}{2} \int \frac{\operatorname{Log}[1 + i c x]}{d + e x^2} dx$$

Program code:

```
Int[ArcTan[c_.*x_]/(d_+e_.*x_^2),x_Symbol] :=
  I/2*Int[Log[1-I*c*x]/(d+e*x^2),x] - I/2*Int[Log[1+I*c*x]/(d+e*x^2),x] /;
  FreeQ[{c,d,e},x]
```

```
Int[ArcCot[c_.*x_]/(d_+e_.*x_^2),x_Symbol] :=
  I/2*Int[Log[1-I/(c*x)]/(d+e*x^2),x] - I/2*Int[Log[1+I/(c*x)]/(d+e*x^2),x] /;
FreeQ[{c,d,e},x]
```

2: $\int \frac{a + b \operatorname{ArcTan}[c x]}{d + e x^2} dx$

Derivation: Algebraic expansion

Rule:

$$\int \frac{a + b \operatorname{ArcTan}[c x]}{d + e x^2} dx \rightarrow a \int \frac{1}{d + e x^2} dx + b \int \frac{\operatorname{ArcTan}[c x]}{d + e x^2} dx$$

Program code:

```
Int[(a_+b_.*ArcTan[c_.*x_])/(d_+e_.*x_^2),x_Symbol] :=
  a*Int[1/(d+e*x^2),x] + b*Int[ArcTan[c*x]/(d+e*x^2),x] /;
FreeQ[{a,b,c,d,e},x]
```

```
Int[(a_+b_.*ArcCot[c_.*x_])/(d_+e_.*x_^2),x_Symbol] :=
  a*Int[1/(d+e*x^2),x] + b*Int[ArcCot[c*x]/(d+e*x^2),x] /;
FreeQ[{a,b,c,d,e},x]
```

3: $\int (d + e x^2)^q (a + b \operatorname{ArcTan}[c x]) dx$ when $q \in \mathbb{Z} \vee q + \frac{1}{2} \in \mathbb{Z}^-$

Derivation: Integration by parts

Note: If $q \in \mathbb{Z}^+ \vee q + \frac{1}{2} \in \mathbb{Z}^-$, then $\int (d + e x^2)^q dx$ is expressible as an algebraic function not involving logarithms, inverse trig or inverse hyperbolic functions.

Rule: If $q \in \mathbb{Z} \vee q + \frac{1}{2} \in \mathbb{Z}^-$, let $u = \int (d + e x^2)^q dx$, then

$$\int (d + e x^2)^q (a + b \operatorname{ArcTan}[c x]) dx \rightarrow u (a + b \operatorname{ArcTan}[c x]) - b c \int \frac{u}{1 + c^2 x^2} dx$$

Program code:

```
Int[(d_+e_.*x_^2)^q_.*(a_+b_.*ArcTan[c_.*x_]),x_Symbol] :=
  With[{u=IntHide[(d+e*x^2)^q,x]},
    Dist[a+b*ArcTan[c*x],u,x] - b*c*Int[u/(1+c^2*x^2),x] /;
  FreeQ[{a,b,c,d,e},x] && (IntegerQ[q] || ILtQ[q+1/2,0])
```

```
Int[(d_+e_.*x_^2)^q_.*(a_+b_.*ArcCot[c_.*x_]),x_Symbol] :=
  With[{u=IntHide[(d+e*x^2)^q,x]},
    Dist[a+b*ArcCot[c*x],u,x] + b*c*Int[u/(1+c^2*x^2),x] /;
  FreeQ[{a,b,c,d,e},x] && (IntegerQ[q] || ILtQ[q+1/2,0])
```

4: $\int (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx$ when $q \in \mathbb{Z} \wedge p \in \mathbb{Z}^+$

Rule: If $q \in \mathbb{Z} \wedge p \in \mathbb{Z}^+$, then

$$\int (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx \rightarrow \int (a + b \operatorname{ArcTan}[c x])^p \operatorname{ExpandIntegrand}[(d + e x^2)^q, x] dx$$

Program code:

```
Int[(d_+e_.*x_^2)^q_.*(a_.+b_.*ArcTan[c_.*x_])^p_.,x_Symbol] :=
  Int[ExpandIntegrand[(a+b*ArcTan[c*x])^p,(d+e*x^2)^q,x],x] /;
FreeQ[{a,b,c,d,e},x] && IntegerQ[q] && IGtQ[p,0]
```

```
Int[(d_+e_.*x_^2)^q_.*(a_.+b_.*ArcCot[c_.*x_])^p_.,x_Symbol] :=
  Int[ExpandIntegrand[(a+b*ArcCot[c*x])^p,(d+e*x^2)^q,x],x] /;
FreeQ[{a,b,c,d,e},x] && IntegerQ[q] && IGtQ[p,0]
```

$$6. \int (f x)^m (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx$$

$$1. \int \frac{(f x)^m (a + b \operatorname{ArcTan}[c x])^p}{d + e x^2} dx$$

$$1: \int \frac{(f x)^m (a + b \operatorname{ArcTan}[c x])^p}{d + e x^2} dx \text{ when } p > 0 \wedge m > 1$$

Derivation: Algebraic expansion

$$\text{Basis: } \frac{x^2}{d+e x^2} = \frac{1}{e} - \frac{d}{e (d+e x^2)}$$

Rule: If $p > 0 \wedge m > 1$, then

$$\int \frac{(f x)^m (a + b \operatorname{ArcTan}[c x])^p}{d + e x^2} dx \rightarrow \frac{f^2}{e} \int (f x)^{m-2} (a + b \operatorname{ArcTan}[c x])^p dx - \frac{d f^2}{e} \int \frac{(f x)^{m-2} (a + b \operatorname{ArcTan}[c x])^p}{d + e x^2} dx$$

Program code:

```
Int[(f_.*x_)^m_*(a_.+b_.*ArcTan[c_.*x_])^p_./(d_+e_.*x_^2),x_Symbol] :=
  f^2/e*Int[(f*x)^(m-2)*(a+b*ArcTan[c*x])^p,x] -
  d*f^2/e*Int[(f*x)^(m-2)*(a+b*ArcTan[c*x])^p/(d+e*x^2),x] /;
FreeQ[{a,b,c,d,e,f},x] && GtQ[p,0] && GtQ[m,1]
```

```
Int[(f_.*x_)^m_*(a_.+b_.*ArcCot[c_.*x_])^p_./(d_+e_.*x_^2),x_Symbol] :=
  f^2/e*Int[(f*x)^(m-2)*(a+b*ArcCot[c*x])^p,x] -
  d*f^2/e*Int[(f*x)^(m-2)*(a+b*ArcCot[c*x])^p/(d+e*x^2),x] /;
FreeQ[{a,b,c,d,e,f},x] && GtQ[p,0] && GtQ[m,1]
```

$$2: \int \frac{(f x)^m (a + b \operatorname{ArcTan}[c x])^p}{d + e x^2} dx \text{ when } p > 0 \wedge m < -1$$

Derivation: Algebraic expansion

Basis: $\frac{1}{d+e x^2} == \frac{1}{d} - \frac{e x^2}{d (d+e x^2)}$

Rule: If $p > 0 \wedge m < -1$, then

$$\int \frac{(f x)^m (a + b \operatorname{ArcTan}[c x])^p}{d + e x^2} dx \rightarrow \frac{1}{d} \int (f x)^m (a + b \operatorname{ArcTan}[c x])^p dx - \frac{e}{d f^2} \int \frac{(f x)^{m+2} (a + b \operatorname{ArcTan}[c x])^p}{d + e x^2} dx$$

Program code:

```
Int[(f_.*x_)^m_*(a_.+b_.*ArcTan[c_.*x_])^p_./(d_+e_.*x_^2),x_Symbol] :=
  1/d*Int[(f*x)^m*(a+b*ArcTan[c*x])^p,x] -
  e/(d*f^2)*Int[(f*x)^(m+2)*(a+b*ArcTan[c*x])^p/(d+e*x^2),x] /;
FreeQ[{a,b,c,d,e,f},x] && GtQ[p,0] && LtQ[m,-1]
```

```
Int[(f_.*x_)^m_*(a_.+b_.*ArcCot[c_.*x_])^p_./(d_+e_.*x_^2),x_Symbol] :=
  1/d*Int[(f*x)^m*(a+b*ArcCot[c*x])^p,x] -
  e/(d*f^2)*Int[(f*x)^(m+2)*(a+b*ArcCot[c*x])^p/(d+e*x^2),x] /;
FreeQ[{a,b,c,d,e,f},x] && GtQ[p,0] && LtQ[m,-1]
```

$$\begin{aligned}
3. \int \frac{(f x)^m (a + b \operatorname{ArcTan}[c x])^p}{d + e x^2} dx & \text{ when } e = c^2 d \\
1. \int \frac{x (a + b \operatorname{ArcTan}[c x])^p}{d + e x^2} dx & \text{ when } e = c^2 d \\
1: \int \frac{x (a + b \operatorname{ArcTan}[c x])^p}{d + e x^2} dx & \text{ when } e = c^2 d \wedge p \in \mathbb{Z}^+
\end{aligned}$$

Derivation: Algebraic expansion and power rule for integration

Basis: If $e = c^2 d$, then $\frac{x}{d+e x^2} = -\frac{i c}{e (1+c^2 x^2)} - \frac{1}{c d (i - c x)}$

Rule: If $e = c^2 d \wedge p \in \mathbb{Z}^+$, then

$$\int \frac{x (a + b \operatorname{ArcTan}[c x])^p}{d + e x^2} dx \rightarrow -\frac{i (a + b \operatorname{ArcTan}[c x])^{p+1}}{b e (p+1)} - \frac{1}{c d} \int \frac{(a + b \operatorname{ArcTan}[c x])^p}{i - c x} dx$$

Program code:

```

Int[x_*(a_+b_.*ArcTan[c_*x_])^p_./(d_+e_*x_^2),x_Symbol] :=
  -I*(a+b*ArcTan[c*x])^(p+1)/(b*e*(p+1)) -
  1/(c*d)*Int[(a+b*ArcTan[c*x])^p/(I-c*x),x] /;
FreeQ[{a,b,c,d,e},x] && EqQ[e,c^2*d] && IGtQ[p,0]

```

```

Int[x_*(a_+b_.*ArcCot[c_*x_])^p_./(d_+e_*x_^2),x_Symbol] :=
  I*(a+b*ArcCot[c*x])^(p+1)/(b*e*(p+1)) -
  1/(c*d)*Int[(a+b*ArcCot[c*x])^p/(I-c*x),x] /;
FreeQ[{a,b,c,d,e},x] && EqQ[e,c^2*d] && IGtQ[p,0]

```


$$\mathbf{2:} \int \frac{x (a + b \operatorname{ArcTan}[c x])^p}{d + e x^2} dx \text{ when } e = c^2 d \wedge p \notin \mathbb{Z}^+ \wedge p \neq -1$$

Derivation: Integration by parts

$$\text{Basis: If } e = c^2 d, \text{ then } \frac{(a+b \operatorname{ArcTan}[c x])^p}{d+e x^2} = \partial_x \frac{(a+b \operatorname{ArcTan}[c x])^{p+1}}{b c d (p+1)}$$

Rule: If $e = c^2 d \wedge p \notin \mathbb{Z}^+ \wedge p \neq -1$, then

$$\int \frac{x (a + b \operatorname{ArcTan}[c x])^p}{d + e x^2} dx \rightarrow \frac{x (a + b \operatorname{ArcTan}[c x])^{p+1}}{b c d (p+1)} - \frac{1}{b c d (p+1)} \int (a + b \operatorname{ArcTan}[c x])^{p+1} dx$$

Program code:

```
Int[x*(a_.+b_.*ArcTan[c_.*x_])^p_/(d_+e_.*x_^2),x_Symbol] :=
  x*(a+b*ArcTan[c*x])^(p+1)/(b*c*d*(p+1)) -
  1/(b*c*d*(p+1))*Int[(a+b*ArcTan[c*x])^(p+1),x] /;
FreeQ[{a,b,c,d,e},x] && EqQ[e,c^2*d] && Not[IGtQ[p,0]] && NeQ[p,-1]
```

```
Int[x*(a_.+b_.*ArcCot[c_.*x_])^p_/(d_+e_.*x_^2),x_Symbol] :=
  -x*(a+b*ArcCot[c*x])^(p+1)/(b*c*d*(p+1)) +
  1/(b*c*d*(p+1))*Int[(a+b*ArcCot[c*x])^(p+1),x] /;
FreeQ[{a,b,c,d,e},x] && EqQ[e,c^2*d] && Not[IGtQ[p,0]] && NeQ[p,-1]
```

$$\mathbf{2:} \int \frac{(a + b \operatorname{ArcTan}[c x])^p}{x (d + e x^2)} dx \text{ when } e = c^2 d \wedge p > 0$$

Derivation: Algebraic expansion

$$\text{Basis: If } e = c^2 d, \text{ then } \frac{1}{x (d+e x^2)} = -\frac{i c}{d+e x^2} + \frac{i}{d x (i+c x)}$$

Rule: If $e = c^2 d \wedge p > 0$, then

$$\int \frac{(a + b \operatorname{ArcTan}[c x])^p}{x (d + e x^2)} dx \rightarrow -\frac{i (a + b \operatorname{ArcTan}[c x])^{p+1}}{b d (p+1)} + \frac{i}{d} \int \frac{(a + b \operatorname{ArcTan}[c x])^p}{x (i + c x)} dx$$

Program code:

```
Int[(a_.+b_.*ArcTan[c_.*x_])^p_./(x_*(d_+e_.*x_^2)),x_Symbol] :=
  -I*(a+b*ArcTan[c*x])^(p+1)/(b*d*(p+1)) +
  I/d*Int[(a+b*ArcTan[c*x])^p/(x*(I+c*x)),x] /;
FreeQ[{a,b,c,d,e},x] && EqQ[e,c^2*d] && GtQ[p,0]
```

```
Int[(a_.+b_.*ArcCot[c_.*x_])^p_./(x_*(d_+e_.*x_^2)),x_Symbol] :=
  I*(a+b*ArcCot[c*x])^(p+1)/(b*d*(p+1)) +
  I/d*Int[(a+b*ArcCot[c*x])^p/(x*(I+c*x)),x] /;
FreeQ[{a,b,c,d,e},x] && EqQ[e,c^2*d] && GtQ[p,0]
```

3: $\int \frac{(f x)^m (a + b \operatorname{ArcTan}[c x])^p}{d + e x^2} dx$ when $e = c^2 d \wedge p < -1$

Derivation: Integration by parts

Basis: If $e = c^2 d$, then $\frac{(a+b \operatorname{ArcTan}[c x])^p}{d+e x^2} = \partial_x \frac{(a+b \operatorname{ArcTan}[c x])^{p+1}}{b c d (p+1)}$

Rule: If $e = c^2 d \wedge p < -1$, then

$$\int \frac{(f x)^m (a + b \operatorname{ArcTan}[c x])^p}{d + e x^2} dx \rightarrow \frac{(f x)^m (a + b \operatorname{ArcTan}[c x])^{p+1}}{b c d (p+1)} - \frac{f m}{b c d (p+1)} \int (f x)^{m-1} (a + b \operatorname{ArcTan}[c x])^{p+1} dx$$

Program code:

```
Int[(f_.*x_)^m_*(a_.+b_.*ArcTan[c_.*x_])^p_./(d_+e_.*x_^2),x_Symbol] :=
  (f*x)^m*(a+b*ArcTan[c*x])^(p+1)/(b*c*d*(p+1)) -
  f*m/(b*c*d*(p+1))*Int[(f*x)^(m-1)*(a+b*ArcTan[c*x])^(p+1),x] /;
FreeQ[{a,b,c,d,e,f,m},x] && EqQ[e,c^2*d] && LtQ[p,-1]
```

```

Int[(f_.*x_)^m.*(a_.+b_.*ArcCot[c_.*x_])^p/(d_+e_.*x_^2),x_Symbol] :=
  -(f*x)^m*(a+b*ArcCot[c*x])^(p+1)/(b*c*d*(p+1)) +
  f*m/(b*c*d*(p+1))*Int[(f*x)^(m-1)*(a+b*ArcCot[c*x])^(p+1),x] /;
FreeQ[{a,b,c,d,e,f,m},x] && EqQ[e,c^2*d] && LtQ[p,-1]

```

4: $\int \frac{x^m (a + b \operatorname{ArcTan}[c x])}{d + e x^2} dx$ when $m \in \mathbb{Z} \wedge \neg (m == 1 \wedge a \neq 0)$

Derivation: Algebraic expansion

Rule: If $m \in \mathbb{Z} \wedge \neg (m == 1 \wedge a \neq 0)$, then

$$\int \frac{x^m (a + b \operatorname{ArcTan}[c x])}{d + e x^2} dx \rightarrow \int (a + b \operatorname{ArcTan}[c x]) \operatorname{ExpandIntegrand}\left[\frac{x^m}{d + e x^2}, x\right] dx$$

Program code:

```

Int[x_^m_.*(a_.+b_.*ArcTan[c_.*x_])/(d_+e_.*x_^2),x_Symbol] :=
  Int[ExpandIntegrand[(a+b*ArcTan[c*x]),x^m/(d+e*x^2),x],x] /;
FreeQ[{a,b,c,d,e},x] && IntegerQ[m] && Not[EqQ[m,1] && NeQ[a,0]]

```

```

Int[x_^m_.*(a_.+b_.*ArcCot[c_.*x_])/(d_+e_.*x_^2),x_Symbol] :=
  Int[ExpandIntegrand[(a+b*ArcCot[c*x]),x^m/(d+e*x^2),x],x] /;
FreeQ[{a,b,c,d,e},x] && IntegerQ[m] && Not[EqQ[m,1] && NeQ[a,0]]

```

$$2. \int (f x)^m (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx \text{ when } e = c^2 d$$

$$1. \int x (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx \text{ when } e = c^2 d$$

$$1: \int x (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx \text{ when } e = c^2 d \wedge p > 0 \wedge q \neq -1$$

Derivation: Integration by parts

Rule: If $e = c^2 d \wedge p > 0 \wedge q \neq -1$, then

$$\int x (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx \rightarrow \frac{(d + e x^2)^{q+1} (a + b \operatorname{ArcTan}[c x])^p}{2 e (q + 1)} - \frac{b p}{2 c (q + 1)} \int (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^{p-1} dx$$

Program code:

```
Int[x_*(d_+e_.*x_^2)^q_.*(a_.+b_.*ArcTan[c_.*x_])^p_.,x_Symbol] :=
  (d+e*x^2)^(q+1)*(a+b*ArcTan[c*x])^p/(2*e*(q+1)) -
  b*p/(2*c*(q+1))*Int[(d+e*x^2)^q*(a+b*ArcTan[c*x])^(p-1),x] /;
FreeQ[{a,b,c,d,e,q},x] && EqQ[e,c^2*d] && GtQ[p,0] && NeQ[q,-1]
```

```
Int[x_*(d_+e_.*x_^2)^q_.*(a_.+b_.*ArcCot[c_.*x_])^p_.,x_Symbol] :=
  (d+e*x^2)^(q+1)*(a+b*ArcCot[c*x])^p/(2*e*(q+1)) +
  b*p/(2*c*(q+1))*Int[(d+e*x^2)^q*(a+b*ArcCot[c*x])^(p-1),x] /;
FreeQ[{a,b,c,d,e,q},x] && EqQ[e,c^2*d] && GtQ[p,0] && NeQ[q,-1]
```

$$\text{2: } \int \frac{x (a + b \operatorname{ArcTan}[c x])^p}{(d + e x^2)^2} dx \text{ when } e = c^2 d \wedge p < -1 \wedge p \neq -2$$

Rule: If $e = c^2 d \wedge p < -1 \wedge p \neq -2$, then

$$\int \frac{x (a + b \operatorname{ArcTan}[c x])^p}{(d + e x^2)^2} dx \rightarrow \frac{x (a + b \operatorname{ArcTan}[c x])^{p+1}}{b c d (p+1) (d + e x^2)} - \frac{(1 - c^2 x^2) (a + b \operatorname{ArcTan}[c x])^{p+2}}{b^2 e (p+1) (p+2) (d + e x^2)} - \frac{4}{b^2 (p+1) (p+2)} \int \frac{x (a + b \operatorname{ArcTan}[c x])^{p+2}}{(d + e x^2)^2} dx$$

Program code:

```
Int[x*(a_.+b_.*ArcTan[c_.*x_])^p_/(d_+e_.*x_^2),x_Symbol] :=
  x*(a+b*ArcTan[c*x])^(p+1)/(b*c*d*(p+1)*(d+e*x^2)) -
  (1-c^2*x^2)*(a+b*ArcTan[c*x])^(p+2)/(b^2*e*(p+1)*(p+2)*(d+e*x^2)) -
  4/(b^2*(p+1)*(p+2))*Int[x*(a+b*ArcTan[c*x])^(p+2)/(d+e*x^2)^2,x] /;
FreeQ[{a,b,c,d,e},x] && EqQ[e,c^2*d] && LtQ[p,-1] && NeQ[p,-2]
```

```
Int[x*(a_.+b_.*ArcCot[c_.*x_])^p_/(d_+e_.*x_^2),x_Symbol] :=
  -x*(a+b*ArcCot[c*x])^(p+1)/(b*c*d*(p+1)*(d+e*x^2)) -
  (1-c^2*x^2)*(a+b*ArcCot[c*x])^(p+2)/(b^2*e*(p+1)*(p+2)*(d+e*x^2)) -
  4/(b^2*(p+1)*(p+2))*Int[x*(a+b*ArcCot[c*x])^(p+2)/(d+e*x^2)^2,x] /;
FreeQ[{a,b,c,d,e},x] && EqQ[e,c^2*d] && LtQ[p,-1] && NeQ[p,-2]
```

$$\text{2. } \int x^2 (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx \text{ when } e = c^2 d$$

$$\text{1: } \int x^2 (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx \text{ when } e = c^2 d \wedge q < -1$$

Rule: If $q = -\frac{5}{2}$, then better to use rule for when $m + 2q + 3 = 0$.

Rule: If $e = c^2 d \wedge q < -1$, then

$$\int x^2 (d + e x^2)^q (a + b \operatorname{ArcTan}[c x]) dx \rightarrow -\frac{b (d + e x^2)^{q+1}}{4 c^3 d (q+1)^2} + \frac{x (d + e x^2)^{q+1} (a + b \operatorname{ArcTan}[c x])}{2 c^2 d (q+1)} - \frac{1}{2 c^2 d (q+1)} \int (d + e x^2)^{q+1} (a + b \operatorname{ArcTan}[c x]) dx$$

Program code:

```
Int[x_^2*(d_+e_*x_^2)^q_*(a_+b_*ArcTan[c_*x_]),x_Symbol] :=
  -b*(d+e*x^2)^(q+1)/(4*c^3*d*(q+1)^2) +
  x*(d+e*x^2)^(q+1)*(a+b*ArcTan[c*x])/(2*c^2*d*(q+1)) -
  1/(2*c^2*d*(q+1))*Int[(d+e*x^2)^(q+1)*(a+b*ArcTan[c*x]),x] /;
FreeQ[{a,b,c,d,e},x] && EqQ[e,c^2*d] && LtQ[q,-1] && NeQ[q,-5/2]
```

```
Int[x_^2*(d_+e_*x_^2)^q_*(a_+b_*ArcCot[c_*x_]),x_Symbol] :=
  b*(d+e*x^2)^(q+1)/(4*c^3*d*(q+1)^2) +
  x*(d+e*x^2)^(q+1)*(a+b*ArcCot[c*x])/(2*c^2*d*(q+1)) -
  1/(2*c^2*d*(q+1))*Int[(d+e*x^2)^(q+1)*(a+b*ArcCot[c*x]),x] /;
FreeQ[{a,b,c,d,e},x] && EqQ[e,c^2*d] && LtQ[q,-1] && NeQ[q,-5/2]
```

2: $\int \frac{x^2 (a + b \operatorname{ArcTan}[c x])^p}{(d + e x^2)^2} dx$ when $e = c^2 d \wedge p > 0$

Rule: If $e = c^2 d \wedge p > 0$, then

$$\int \frac{x^2 (a + b \operatorname{ArcTan}[c x])^p}{(d + e x^2)^2} dx \rightarrow \frac{(a + b \operatorname{ArcTan}[c x])^{p+1}}{2 b c^3 d^2 (p+1)} - \frac{x (a + b \operatorname{ArcTan}[c x])^p}{2 c^2 d (d + e x^2)} + \frac{b p}{2 c} \int \frac{x (a + b \operatorname{ArcTan}[c x])^{p-1}}{(d + e x^2)^2} dx$$

Program code:

```
Int[x_^2*(a_+b_*ArcTan[c_*x_])^p_/(d_+e_*x_^2)^2,x_Symbol] :=
  (a+b*ArcTan[c*x])^(p+1)/(2*b*c^3*d^2*(p+1)) -
  x*(a+b*ArcTan[c*x])^p/(2*c^2*d*(d+e*x^2)) +
  b*p/(2*c)*Int[x*(a+b*ArcTan[c*x])^(p-1)/(d+e*x^2)^2,x] /;
FreeQ[{a,b,c,d,e},x] && EqQ[e,c^2*d] && GtQ[p,0]
```

```

Int[x_^2*(a_+b_.*ArcCot[c_.*x_])^p_./(d_+e_.*x_^2)^2,x_Symbol] :=
  -(a+b*ArcCot[c*x])^(p+1)/(2*b*c^3*d^2*(p+1)) -
  x*(a+b*ArcCot[c*x])^p/(2*c^2*d*(d+e*x^2)) -
  b*p/(2*c)*Int[x*(a+b*ArcCot[c*x])^(p-1)/(d+e*x^2)^2,x] /;
FreeQ[{a,b,c,d,e},x] && EqQ[e,c^2*d] && GtQ[p,0]

```

3. $\int (f x)^m (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx$ when $e = c^2 d \wedge m + 2q + 2 = 0$

1. $\int (f x)^m (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx$ when $e = c^2 d \wedge m + 2q + 2 = 0 \wedge q < -1 \wedge p \geq 1$

1: $\int (f x)^m (d + e x^2)^q (a + b \operatorname{ArcTan}[c x]) dx$ when $e = c^2 d \wedge m + 2q + 2 = 0 \wedge q < -1$

Rule: If $e = c^2 d \wedge m + 2q + 2 = 0 \wedge q < -1$, then

$$\int (f x)^m (d + e x^2)^q (a + b \operatorname{ArcTan}[c x]) dx \rightarrow \frac{b (f x)^m (d + e x^2)^{q+1}}{c d m^2} - \frac{f (f x)^{m-1} (d + e x^2)^{q+1} (a + b \operatorname{ArcTan}[c x])}{c^2 d m} + \frac{f^2 (m-1)}{c^2 d m} \int (f x)^{m-2} (d + e x^2)^{q+1} (a + b \operatorname{ArcTan}[c x]) dx$$

Program code:

```

Int[(f_.*x_)^m_*(d_+e_.*x_^2)^q_*(a_+b_.*ArcTan[c_.*x_]),x_Symbol] :=
  b*(f*x)^m*(d+e*x^2)^(q+1)/(c*d*m^2) -
  f*(f*x)^(m-1)*(d+e*x^2)^(q+1)*(a+b*ArcTan[c*x])/(c^2*d*m) +
  f^2*(m-1)/(c^2*d*m)*Int[(f*x)^(m-2)*(d+e*x^2)^(q+1)*(a+b*ArcTan[c*x]),x] /;
FreeQ[{a,b,c,d,e,f},x] && EqQ[e,c^2*d] && EqQ[m+2*q+2,0] && LtQ[q,-1]

```

```

Int[(f_.*x_)^m_*(d_+e_.*x_^2)^q_*(a_+b_.*ArcCot[c_.*x_]),x_Symbol] :=
  -b*(f*x)^m*(d+e*x^2)^(q+1)/(c*d*m^2) -
  f*(f*x)^(m-1)*(d+e*x^2)^(q+1)*(a+b*ArcCot[c*x])/(c^2*d*m) +
  f^2*(m-1)/(c^2*d*m)*Int[(f*x)^(m-2)*(d+e*x^2)^(q+1)*(a+b*ArcCot[c*x]),x] /;
FreeQ[{a,b,c,d,e,f},x] && EqQ[e,c^2*d] && EqQ[m+2*q+2,0] && LtQ[q,-1]

```

2: $\int (f x)^m (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx$ when $e = c^2 d \wedge m + 2q + 2 = 0 \wedge q < -1 \wedge p > 1$

Rule: If $e = c^2 d \wedge m + 2q + 2 = 0 \wedge q < -1 \wedge p > 1$, then

$$\int (f x)^m (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx \rightarrow$$

$$\frac{b p (f x)^m (d + e x^2)^{q+1} (a + b \operatorname{ArcTan}[c x])^{p-1}}{c d m^2} - \frac{f (f x)^{m-1} (d + e x^2)^{q+1} (a + b \operatorname{ArcTan}[c x])^p}{c^2 d m} -$$

$$\frac{b^2 p (p-1)}{m^2} \int (f x)^m (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^{p-2} dx + \frac{f^2 (m-1)}{c^2 d m} \int (f x)^{m-2} (d + e x^2)^{q+1} (a + b \operatorname{ArcTan}[c x])^p dx$$

Program code:

```
Int[(f_.x_)^m_*(d_+e_.x_^2)^q_*(a_.+b_.*ArcTan[c_.x_])^p_,x_Symbol] :=
  b*p*(f*x)^m*(d+e*x^2)^(q+1)*(a+b*ArcTan[c*x])^(p-1)/(c*d*m^2) -
  f*(f*x)^(m-1)*(d+e*x^2)^(q+1)*(a+b*ArcTan[c*x])^p/(c^2*d*m) -
  b^2*p*(p-1)/m^2*Int[(f*x)^m*(d+e*x^2)^q*(a+b*ArcTan[c*x])^(p-2),x] +
  f^2*(m-1)/(c^2*d*m)*Int[(f*x)^(m-2)*(d+e*x^2)^(q+1)*(a+b*ArcTan[c*x])^p,x] /;
FreeQ[{a,b,c,d,e,f,m},x] && EqQ[e,c^2*d] && EqQ[m+2*q+2,0] && LtQ[q,-1] && GtQ[p,1]
```

```
Int[(f_.x_)^m_*(d_+e_.x_^2)^q_*(a_.+b_.*ArcCot[c_.x_])^p_,x_Symbol] :=
  -b*p*(f*x)^m*(d+e*x^2)^(q+1)*(a+b*ArcCot[c*x])^(p-1)/(c*d*m^2) -
  f*(f*x)^(m-1)*(d+e*x^2)^(q+1)*(a+b*ArcCot[c*x])^p/(c^2*d*m) -
  b^2*p*(p-1)/m^2*Int[(f*x)^m*(d+e*x^2)^q*(a+b*ArcCot[c*x])^(p-2),x] +
  f^2*(m-1)/(c^2*d*m)*Int[(f*x)^(m-2)*(d+e*x^2)^(q+1)*(a+b*ArcCot[c*x])^p,x] /;
FreeQ[{a,b,c,d,e,f,m},x] && EqQ[e,c^2*d] && EqQ[m+2*q+2,0] && LtQ[q,-1] && GtQ[p,1]
```


2: $\int (f x)^m (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx$ when $e = c^2 d \wedge m + 2q + 2 = 0 \wedge p < -1$

Derivation: Integration by parts

Basis: If $e = c^2 d$, then $\frac{(a+b \operatorname{ArcTan}[c x])^p}{d+e x^2} = \partial_x \frac{(a+b \operatorname{ArcTan}[c x])^{p+1}}{b c d (p+1)}$

Basis: If $m + 2q + 2 = 0$, then $\partial_x (x^m (d + e x^2)^{q+1}) = c m x^{m-1} (d + e x^2)^q$

Rule: If $e = c^2 d \wedge m + 2q + 2 = 0 \wedge p < -1$, then

$$\int (f x)^m (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx \rightarrow \frac{(f x)^m (d + e x^2)^{q+1} (a + b \operatorname{ArcTan}[c x])^{p+1}}{b c d (p+1)} - \frac{f m}{b c (p+1)} \int (f x)^{m-1} (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^{p+1} dx$$

Program code:

```
Int[(f_.*x_)^m_.*(d_+e_.*x_^2)^q_.*(a_+b_.*ArcTan[c_.*x_])^p_,x_Symbol] :=
  (f*x)^m*(d+e*x^2)^(q+1)*(a+b*ArcTan[c*x])^(p+1)/(b*c*d*(p+1)) -
  f*m/(b*c*(p+1))*Int[(f*x)^(m-1)*(d+e*x^2)^q*(a+b*ArcTan[c*x])^(p+1),x] /;
FreeQ[{a,b,c,d,e,f,m,q},x] && EqQ[e,c^2*d] && EqQ[m+2*q+2,0] && LtQ[p,-1]
```

```
Int[(f_.*x_)^m_.*(d_+e_.*x_^2)^q_.*(a_+b_.*ArcCot[c_.*x_])^p_,x_Symbol] :=
  -(f*x)^m*(d+e*x^2)^(q+1)*(a+b*ArcCot[c*x])^(p+1)/(b*c*d*(p+1)) +
  f*m/(b*c*(p+1))*Int[(f*x)^(m-1)*(d+e*x^2)^q*(a+b*ArcCot[c*x])^(p+1),x] /;
FreeQ[{a,b,c,d,e,f,m,q},x] && EqQ[e,c^2*d] && EqQ[m+2*q+2,0] && LtQ[p,-1]
```

$$4: \int (f x)^m (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx \text{ when } e = c^2 d \wedge m + 2q + 3 = 0 \wedge p > 0 \wedge m \neq -1$$

Derivation: Integration by parts

■ Basis: If $m + 2q + 3 = 0$, then $x^m (d + e x^2)^q = \partial_x \frac{x^{m+1} (d + e x^2)^{q+1}}{d (m+1)}$

Rule: If $e = c^2 d \wedge m + 2q + 3 = 0 \wedge p > 0 \wedge m \neq -1$, then

$$\int (f x)^m (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx \rightarrow \frac{(f x)^{m+1} (d + e x^2)^{q+1} (a + b \operatorname{ArcTan}[c x])^p}{d f (m+1)} - \frac{b c p}{f (m+1)} \int (f x)^{m+1} (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^{p-1} dx$$

Program code:

```
Int[(f_.*x_)^m_.*(d_+e_.*x_^2)^q_.*(a_.+b_.*ArcTan[c_.*x_])^p_,x_Symbol] :=
  (f*x)^(m+1)*(d+e*x^2)^(q+1)*(a+b*ArcTan[c*x])^p/(d*f*(m+1)) -
  b*c*p/(f*(m+1))*Int[(f*x)^(m+1)*(d+e*x^2)^q*(a+b*ArcTan[c*x])^(p-1),x] /;
FreeQ[{a,b,c,d,e,f,m,q},x] && EqQ[e,c^2*d] && EqQ[m+2*q+3,0] && GtQ[p,0] && NeQ[m,-1]
```

```
Int[(f_.*x_)^m_.*(d_+e_.*x_^2)^q_.*(a_.+b_.*ArcCot[c_.*x_])^p_,x_Symbol] :=
  (f*x)^(m+1)*(d+e*x^2)^(q+1)*(a+b*ArcCot[c*x])^p/(d*f*(m+1)) +
  b*c*p/(f*(m+1))*Int[(f*x)^(m+1)*(d+e*x^2)^q*(a+b*ArcCot[c*x])^(p-1),x] /;
FreeQ[{a,b,c,d,e,f,m,q},x] && EqQ[e,c^2*d] && EqQ[m+2*q+3,0] && GtQ[p,0] && NeQ[m,-1]
```

5. $\int (f x)^m (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx$ when $e = c^2 d \wedge q > 0$

1: $\int (f x)^m \sqrt{d + e x^2} (a + b \operatorname{ArcTan}[c x]) dx$ when $e = c^2 d \wedge m \neq -2$

Rule: If $e = c^2 d \wedge m \neq -2$, then

$$\int (f x)^m \sqrt{d + e x^2} (a + b \operatorname{ArcTan}[c x]) dx \rightarrow \frac{(f x)^{m+1} \sqrt{d + e x^2} (a + b \operatorname{ArcTan}[c x])}{f (m+2)} - \frac{b c d}{f (m+2)} \int \frac{(f x)^{m+1}}{\sqrt{d + e x^2}} dx + \frac{d}{m+2} \int \frac{(f x)^m (a + b \operatorname{ArcTan}[c x])}{\sqrt{d + e x^2}} dx$$

Program code:

```
Int[(f_.**x_)^m_*Sqrt[d_+e_.**x_^2]*(a_.+b_.*ArcTan[c_.**x_]),x_Symbol] :=
  (f*x)^(m+1)*Sqrt[d+e*x^2]*(a+b*ArcTan[c*x])/(f*(m+2)) -
  b*c*d/(f*(m+2))*Int[(f*x)^(m+1)/Sqrt[d+e*x^2],x] +
  d/(m+2)*Int[(f*x)^m*(a+b*ArcTan[c*x])/Sqrt[d+e*x^2],x] /;
FreeQ[{a,b,c,d,e,f,m},x] && EqQ[e,c^2*d] && NeQ[m,-2]
```

```
Int[(f_.**x_)^m_*Sqrt[d_+e_.**x_^2]*(a_.+b_.*ArcCot[c_.**x_]),x_Symbol] :=
  (f*x)^(m+1)*Sqrt[d+e*x^2]*(a+b*ArcCot[c*x])/(f*(m+2)) +
  b*c*d/(f*(m+2))*Int[(f*x)^(m+1)/Sqrt[d+e*x^2],x] +
  d/(m+2)*Int[(f*x)^m*(a+b*ArcCot[c*x])/Sqrt[d+e*x^2],x] /;
FreeQ[{a,b,c,d,e,f,m},x] && EqQ[e,c^2*d] && NeQ[m,-2]
```

$$\text{2: } \int (f x)^m (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx \text{ when } e = c^2 d \wedge p \in \mathbb{Z}^+ \wedge q - 1 \in \mathbb{Z}^+$$

Rule: If $e = c^2 d \wedge p \in \mathbb{Z}^+ \wedge q - 1 \in \mathbb{Z}^+$, then

$$\int (f x)^m (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx \rightarrow \int \operatorname{ExpandIntegrand}[(f x)^m (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p, x] dx$$

Program code:

```
Int[(f_.*x_)^m_*(d_+e_.*x_^2)^q_*(a_+b_.*ArcTan[c_.*x_])^p_.,x_Symbol] :=
  Int[ExpandIntegrand[(f*x)^m*(d+e*x^2)^q*(a+b*ArcTan[c*x])^p,x],x] /;
FreeQ[{a,b,c,d,e,f,m},x] && EqQ[e,c^2*d] && IGtQ[p,0] && IGtQ[q,1] && (EqQ[p,1] || IntegerQ[m])
```

```
Int[(f_.*x_)^m_*(d_+e_.*x_^2)^q_*(a_+b_.*ArcCot[c_.*x_])^p_.,x_Symbol] :=
  Int[ExpandIntegrand[(f*x)^m*(d+e*x^2)^q*(a+b*ArcCot[c*x])^p,x],x] /;
FreeQ[{a,b,c,d,e,f,m},x] && EqQ[e,c^2*d] && IGtQ[p,0] && IGtQ[q,1] && (EqQ[p,1] || IntegerQ[m])
```

$$\text{3: } \int (f x)^m (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx \text{ when } e = c^2 d \wedge q > 0 \wedge p \in \mathbb{Z}^+$$

Derivation: Algebraic expansion

Basis: If $e = c^2 d$, then $(d + e x^2)^q = d (d + e x^2)^{q-1} + c^2 d x^2 (d + e x^2)^{q-1}$

Rule: If $e = c^2 d \wedge q > 0 \wedge p \in \mathbb{Z}^+$, then

$$\int (f x)^m (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx \rightarrow d \int (f x)^m (d + e x^2)^{q-1} (a + b \operatorname{ArcTan}[c x])^p dx + \frac{c^2 d}{f^2} \int (f x)^{m+2} (d + e x^2)^{q-1} (a + b \operatorname{ArcTan}[c x])^p dx$$

Program code:

```
Int[(f_.*x_)^m_*(d_+e_.*x_^2)^q_.*(a_+b_.*ArcTan[c_.*x_])^p_,x_Symbol] :=
  d*Int[(f*x)^m*(d+e*x^2)^(q-1)*(a+b*ArcTan[c*x])^p,x] +
  c^2*d/f^2*Int[(f*x)^(m+2)*(d+e*x^2)^(q-1)*(a+b*ArcTan[c*x])^p,x] /;
FreeQ[{a,b,c,d,e,f,m},x] && EqQ[e,c^2*d] && GtQ[q,0] && IGtQ[p,0] && (RationalQ[m] || EqQ[p,1] && IntegerQ[q])
```

```
Int[(f_.*x_)^m_*(d_+e_.*x_^2)^q_.*(a_+b_.*ArcCot[c_.*x_])^p_,x_Symbol] :=
  d*Int[(f*x)^m*(d+e*x^2)^(q-1)*(a+b*ArcCot[c*x])^p,x] +
  c^2*d/f^2*Int[(f*x)^(m+2)*(d+e*x^2)^(q-1)*(a+b*ArcCot[c*x])^p,x] /;
FreeQ[{a,b,c,d,e,f,m},x] && EqQ[e,c^2*d] && GtQ[q,0] && IGtQ[p,0] && (RationalQ[m] || EqQ[p,1] && IntegerQ[q])
```

$$6. \int (f x)^m (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx \text{ when } e = c^2 d \wedge q < 0$$

$$1. \int \frac{(f x)^m (a + b \operatorname{ArcTan}[c x])^p}{\sqrt{d + e x^2}} dx \text{ when } e = c^2 d$$

$$\text{1: } \int \frac{(f x)^m (a + b \operatorname{ArcTan}[c x])^p}{\sqrt{d + e x^2}} dx \text{ when } e = c^2 d \wedge p > 0 \wedge m > 1$$

Rule: If $e = c^2 d \wedge p > 0 \wedge m > 1$, then

$$\int \frac{(fx)^m (a+b \operatorname{ArcTan}[cx])^p}{\sqrt{d+ex^2}} dx \rightarrow \frac{f (fx)^{m-1} \sqrt{d+ex^2} (a+b \operatorname{ArcTan}[cx])^p}{c^2 d m} - \frac{b f p}{c m} \int \frac{(fx)^{m-1} (a+b \operatorname{ArcTan}[cx])^{p-1}}{\sqrt{d+ex^2}} dx - \frac{f^2 (m-1)}{c^2 m} \int \frac{(fx)^{m-2} (a+b \operatorname{ArcTan}[cx])^p}{\sqrt{d+ex^2}} dx$$

Program code:

```
Int[(f_.*x_)^m_*(a_.+b_.*ArcTan[c_.*x_])^p_./Sqrt[d_+e_.*x_^2],x_Symbol] :=
  f*(f*x)^(m-1)*Sqrt[d+e*x^2]*(a+b*ArcTan[c*x])^p/(c^2*d*m) -
  b*f*p/(c*m)*Int[(f*x)^(m-1)*(a+b*ArcTan[c*x])^(p-1)/Sqrt[d+e*x^2],x] -
  f^2*(m-1)/(c^2*m)*Int[(f*x)^(m-2)*(a+b*ArcTan[c*x])^p/Sqrt[d+e*x^2],x] /;
FreeQ[{a,b,c,d,e,f},x] && EqQ[e,c^2*d] && GtQ[p,0] && GtQ[m,1]
```

```
Int[(f_.*x_)^m_*(a_.+b_.*ArcCot[c_.*x_])^p_./Sqrt[d_+e_.*x_^2],x_Symbol] :=
  f*(f*x)^(m-1)*Sqrt[d+e*x^2]*(a+b*ArcCot[c*x])^p/(c^2*d*m) +
  b*f*p/(c*m)*Int[(f*x)^(m-1)*(a+b*ArcCot[c*x])^(p-1)/Sqrt[d+e*x^2],x] -
  f^2*(m-1)/(c^2*m)*Int[(f*x)^(m-2)*(a+b*ArcCot[c*x])^p/Sqrt[d+e*x^2],x] /;
FreeQ[{a,b,c,d,e,f},x] && EqQ[e,c^2*d] && GtQ[p,0] && GtQ[m,1]
```

$$2. \int \frac{(fx)^m (a+b \operatorname{ArcTan}[cx])^p}{\sqrt{d+ex^2}} dx \text{ when } e = c^2 d \wedge p > 0 \wedge m \leq -1$$

$$1. \int \frac{(a+b \operatorname{ArcTan}[cx])^p}{x \sqrt{d+ex^2}} dx \text{ when } e = c^2 d \wedge p \in \mathbb{Z}^+$$

$$1. \int \frac{(a+b \operatorname{ArcTan}[cx])^p}{x \sqrt{d+ex^2}} dx \text{ when } e = c^2 d \wedge p \in \mathbb{Z}^+ \wedge d > 0$$

$$1: \int \frac{(a+b \operatorname{ArcTan}[cx])}{x \sqrt{d+ex^2}} dx \text{ when } e = c^2 d \wedge d > 0$$

Derivation: Integration by substitution, piecewise constant extraction and algebraic simplification!

Note: Although not essential, these rules returns antiderivatives free of complex exponentials of the form $e^{\operatorname{ArcTan}[cx]}$ and $i e^{\operatorname{ArcCot}[cx]}$.

Basis: If $e = c^2 d \wedge d > 0$, then $\frac{1}{x \sqrt{d+e x^2}} = \frac{1}{\sqrt{d}} \text{Csc}[\text{ArcTan}[c x]] \partial_x \text{ArcTan}[c x]$

Basis: If $e = c^2 d \wedge d > 0$, then $\frac{1}{x \sqrt{d+e x^2}} = -\frac{1}{\sqrt{d}} \frac{\text{Csc}[\text{ArcCot}[c x]] \text{Sec}[\text{ArcCot}[c x]]}{\sqrt{\text{Csc}[\text{ArcCot}[c x]]^2}} \partial_x \text{ArcCot}[c x]$

Rule: If $e = c^2 d \wedge d > 0$, then

$$\int \frac{(a + b \text{ArcTan}[c x])}{x \sqrt{d + e x^2}} dx \rightarrow \frac{1}{\sqrt{d}} \text{Subst}\left[\int (a + b x) \text{Csc}[x] dx, x, \text{ArcTan}[c x]\right]$$

$$\rightarrow -\frac{2}{\sqrt{d}} (a + b \text{ArcTan}[c x]) \text{ArcTanh}\left[\frac{\sqrt{1 + i c x}}{\sqrt{1 - i c x}}\right] + \frac{i b}{\sqrt{d}} \text{PolyLog}\left[2, -\frac{\sqrt{1 + i c x}}{\sqrt{1 - i c x}}\right] - \frac{i b}{\sqrt{d}} \text{PolyLog}\left[2, \frac{\sqrt{1 + i c x}}{\sqrt{1 - i c x}}\right]$$

Program code:

```
Int[(a_.+b_.*ArcTan[c_.*x_])/(x_*Sqrt[d_+e_.*x_^2]),x_Symbol] :=
  -2/Sqrt[d]*(a+b*ArcTan[c*x])*ArcTanh[Sqrt[1+I*c*x]/Sqrt[1-I*c*x]] +
  I*b/Sqrt[d]*PolyLog[2,-Sqrt[1+I*c*x]/Sqrt[1-I*c*x]] -
  I*b/Sqrt[d]*PolyLog[2,Sqrt[1+I*c*x]/Sqrt[1-I*c*x]] /;
FreeQ[{a,b,c,d,e},x] && EqQ[e,c^2*d] && GtQ[d,0]
```

```
Int[(a_.+b_.*ArcCot[c_.*x_])/(x_*Sqrt[d_+e_.*x_^2]),x_Symbol] :=
  -2/Sqrt[d]*(a+b*ArcCot[c*x])*ArcTanh[Sqrt[1+I*c*x]/Sqrt[1-I*c*x]] -
  I*b/Sqrt[d]*PolyLog[2,-Sqrt[1+I*c*x]/Sqrt[1-I*c*x]] +
  I*b/Sqrt[d]*PolyLog[2,Sqrt[1+I*c*x]/Sqrt[1-I*c*x]] /;
FreeQ[{a,b,c,d,e},x] && EqQ[e,c^2*d] && GtQ[d,0]
```

$$2. \int \frac{(a + b \text{ArcTan}[c x])^p}{x \sqrt{d + e x^2}} dx \text{ when } e = c^2 d \wedge p \in \mathbb{Z}^+ \wedge d > 0$$

$$1: \int \frac{(a + b \text{ArcTan}[c x])^p}{x \sqrt{d + e x^2}} dx \text{ when } e = c^2 d \wedge p \in \mathbb{Z}^+ \wedge d > 0$$

Derivation: Integration by substitution

Basis: If $e == c^2 d \wedge d > 0$, then $\frac{1}{x \sqrt{d+e x^2}} == \frac{1}{\sqrt{d}} \text{Csc}[\text{ArcTan}[c x]] \partial_x \text{ArcTan}[c x]$

Rule: If $e == c^2 d \wedge p \in \mathbb{Z}^+ \wedge d > 0$, then

$$\int \frac{(a + b \text{ArcTan}[c x])^p}{x \sqrt{d + e x^2}} dx \rightarrow \frac{1}{\sqrt{d}} \text{Subst}\left[\int (a + b x)^p \text{Csc}[x] dx, x, \text{ArcTan}[c x]\right]$$

Program code:

```
Int[(a_.+b_.*ArcTan[c_.*x_])^p/(x_*Sqrt[d_+e_.*x_^2]),x_Symbol] :=
  1/Sqrt[d]*Subst[Int[(a+b*x)^p*Csc[x],x],x,ArcTan[c*x]] /;
FreeQ[{a,b,c,d,e},x] && EqQ[e,c^2*d] && IGtQ[p,0] && GtQ[d,0]
```

$$\mathbf{2:} \int \frac{(a + b \text{ArcCot}[c x])^p}{x \sqrt{d + e x^2}} dx \text{ when } e == c^2 d \wedge p \in \mathbb{Z}^+ \wedge d > 0$$

Derivation: Integration by substitution and piecewise constant extraction

Basis: If $e == c^2 d \wedge d > 0$, then $\frac{1}{x \sqrt{d+e x^2}} == -\frac{1}{\sqrt{d}} \frac{\text{Csc}[\text{ArcCot}[c x]] \text{Sec}[\text{ArcCot}[c x]]}{\sqrt{\text{Csc}[\text{ArcCot}[c x]]^2}} \partial_x \text{ArcCot}[c x]$

Basis: $\partial_x \frac{\text{Csc}[x]}{\sqrt{\text{Csc}[x]^2}} == 0$

Basis: $\frac{\text{Csc}[\text{ArcCot}[c x]]}{\sqrt{\text{Csc}[\text{ArcCot}[c x]]^2}} == \frac{c x \sqrt{1 + \frac{1}{c^2 x^2}}}{\sqrt{1 + c^2 x^2}}$

Rule: If $e == c^2 d \wedge p \in \mathbb{Z}^+ \wedge d > 0$, then

$$\int \frac{(a + b \text{ArcCot}[c x])^p}{x \sqrt{d + e x^2}} dx \rightarrow -\frac{1}{\sqrt{d}} \text{Subst}\left[\int \frac{(a + b x)^p \text{Csc}[x] \text{Sec}[x]}{\sqrt{\text{Csc}[x]^2}} dx, x, \text{ArcCot}[c x]\right]$$

$$\rightarrow -\frac{c x \sqrt{1 + \frac{1}{c^2 x^2}}}{\sqrt{d + e x^2}} \text{Subst}\left[\int (a + b x)^p \sec[x] dx, x, \text{ArcCot}[c x]\right]$$

Program code:

```
Int[(a_.+b_.*ArcCot[c_.*x_])^p_/(x_*Sqrt[d_+e_.*x_^2]),x_Symbol] :=
  -c*x*Sqrt[1+1/(c^2*x^2)]/Sqrt[d+e*x^2]*Subst[Int[(a+b*x)^p*Sec[x],x],x,ArcCot[c*x]] /;
FreeQ[{a,b,c,d,e},x] && EqQ[e,c^2*d] && IGtQ[p,0] && GtQ[d,0]
```

$$2: \int \frac{(a + b \text{ArcTan}[c x])^p}{x \sqrt{d + e x^2}} dx \text{ when } e = c^2 d \wedge p \in \mathbb{Z}^+ \wedge d \neq 0$$

Derivation: Piecewise constant extraction

Basis: If $e = c^2 d$, then $\partial_x \frac{\sqrt{1+c^2 x^2}}{\sqrt{d+e x^2}} = 0$

Rule: If $e = c^2 d \wedge p \in \mathbb{Z}^+ \wedge d \neq 0$, then

$$\int \frac{(a + b \text{ArcTan}[c x])^p}{x \sqrt{d + e x^2}} dx \rightarrow \frac{\sqrt{1 + c^2 x^2}}{\sqrt{d + e x^2}} \int \frac{(a + b \text{ArcTan}[c x])^p}{x \sqrt{1 + c^2 x^2}} dx$$

Program code:

```
Int[(a_.+b_.*ArcTan[c_.*x_])^p_/(x_*Sqrt[d_+e_.*x_^2]),x_Symbol] :=
  Sqrt[1+c^2*x^2]/Sqrt[d+e*x^2]*Int[(a+b*ArcTan[c*x])^p/(x*Sqrt[1+c^2*x^2]),x] /;
FreeQ[{a,b,c,d,e},x] && EqQ[e,c^2*d] && IGtQ[p,0] && Not[GtQ[d,0]]
```

```
Int[(a_.+b_.*ArcCot[c_.*x_])^p_/(x_*Sqrt[d_+e_.*x_^2]),x_Symbol] :=
  Sqrt[1+c^2*x^2]/Sqrt[d+e*x^2]*Int[(a+b*ArcCot[c*x])^p/(x*Sqrt[1+c^2*x^2]),x] /;
FreeQ[{a,b,c,d,e},x] && EqQ[e,c^2*d] && IGtQ[p,0] && Not[GtQ[d,0]]
```

$$2. \int \frac{(f x)^m (a + b \operatorname{ArcTan}[c x])^p}{\sqrt{d + e x^2}} dx \text{ when } e = c^2 d \wedge p > 0 \wedge m < -1$$

$$1: \int \frac{(a + b \operatorname{ArcTan}[c x])^p}{x^2 \sqrt{d + e x^2}} dx \text{ when } e = c^2 d \wedge p > 0$$

Derivation: Integration by parts

$$\text{Basis: } \frac{1}{x^2 \sqrt{d + e x^2}} = -\partial_x \frac{\sqrt{d + e x^2}}{d x}$$

Rule: If $e = c^2 d \wedge p > 0$, then

$$\int \frac{(a + b \operatorname{ArcTan}[c x])^p}{x^2 \sqrt{d + e x^2}} dx \rightarrow -\frac{\sqrt{d + e x^2} (a + b \operatorname{ArcTan}[c x])^p}{d x} + b c p \int \frac{(a + b \operatorname{ArcTan}[c x])^{p-1}}{x \sqrt{d + e x^2}} dx$$

Program code:

```
Int[(a_.+b_.*ArcTan[c_.*x_])^p_/ (x_^2*Sqrt[d_+e_.*x_^2]), x_Symbol] :=
  -Sqrt[d+e*x^2]*(a+b*ArcTan[c*x])^p/(d*x) +
  b*c*p*Int[(a+b*ArcTan[c*x])^(p-1)/(x*Sqrt[d+e*x^2]), x] /;
FreeQ[{a,b,c,d,e}, x] && EqQ[e, c^2*d] && GtQ[p, 0]
```

```
Int[(a_.+b_.*ArcCot[c_.*x_])^p_/ (x_^2*Sqrt[d_+e_.*x_^2]), x_Symbol] :=
  -Sqrt[d+e*x^2]*(a+b*ArcCot[c*x])^p/(d*x) -
  b*c*p*Int[(a+b*ArcCot[c*x])^(p-1)/(x*Sqrt[d+e*x^2]), x] /;
FreeQ[{a,b,c,d,e}, x] && EqQ[e, c^2*d] && GtQ[p, 0]
```

$$2: \int \frac{(f x)^m (a + b \operatorname{ArcTan}[c x])^p}{\sqrt{d + e x^2}} dx \text{ when } e = c^2 d \wedge p > 0 \wedge m < -1 \wedge m \neq -2$$

Rule: If $e = c^2 d \wedge p > 0 \wedge m < -1 \wedge m \neq -2$, then

$$\int \frac{(f x)^m (a + b \operatorname{ArcTan}[c x])^p}{\sqrt{d + e x^2}} dx \rightarrow$$

$$\frac{(f x)^{m+1} \sqrt{d + e x^2} (a + b \operatorname{ArcTan}[c x])^p}{d f (m+1)} - \frac{b c p}{f (m+1)} \int \frac{(f x)^{m+1} (a + b \operatorname{ArcTan}[c x])^{p-1}}{\sqrt{d + e x^2}} dx - \frac{c^2 (m+2)}{f^2 (m+1)} \int \frac{(f x)^{m+2} (a + b \operatorname{ArcTan}[c x])^p}{\sqrt{d + e x^2}} dx$$

Program code:

```
Int[(f_.*x_)^m_*(a_.+b_.*ArcTan[c_.*x_])^p_/Sqrt[d_+e_.*x_^2],x_Symbol] :=
  (f*x)^(m+1)*Sqrt[d+e*x^2]*(a+b*ArcTan[c*x])^p/(d*f*(m+1)) -
  b*c*p/(f*(m+1))*Int[(f*x)^(m+1)*(a+b*ArcTan[c*x])^(p-1)/Sqrt[d+e*x^2],x] -
  c^2*(m+2)/(f^2*(m+1))*Int[(f*x)^(m+2)*(a+b*ArcTan[c*x])^p/Sqrt[d+e*x^2],x] /;
FreeQ[{a,b,c,d,e,f},x] && EqQ[e,c^2*d] && GtQ[p,0] && LtQ[m,-1] && NeQ[m,-2]
```

```
Int[(f_.*x_)^m_*(a_.+b_.*ArcCot[c_.*x_])^p_/Sqrt[d_+e_.*x_^2],x_Symbol] :=
  (f*x)^(m+1)*Sqrt[d+e*x^2]*(a+b*ArcCot[c*x])^p/(d*f*(m+1)) +
  b*c*p/(f*(m+1))*Int[(f*x)^(m+1)*(a+b*ArcCot[c*x])^(p-1)/Sqrt[d+e*x^2],x] -
  c^2*(m+2)/(f^2*(m+1))*Int[(f*x)^(m+2)*(a+b*ArcCot[c*x])^p/Sqrt[d+e*x^2],x] /;
FreeQ[{a,b,c,d,e,f},x] && EqQ[e,c^2*d] && GtQ[p,0] && LtQ[m,-1] && NeQ[m,-2]
```

$$2. \int (f x)^m (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx \text{ when } e = c^2 d \wedge q < -1$$

$$1: \int x^m (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx \text{ when } e = c^2 d \wedge (m \mid p \mid 2q) \in \mathbb{Z} \wedge q < -1 \wedge m > 1 \wedge p \neq -1$$

Derivation: Algebraic expansion

$$\text{Basis: } \frac{x^2}{d+e x^2} = \frac{1}{e} - \frac{d}{e (d+e x^2)}$$

Rule: If $e = c^2 d \wedge (m \mid p \mid 2q) \in \mathbb{Z} \wedge q < -1 \wedge m > 1 \wedge p \neq -1$, then

$$\int x^m (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx \rightarrow \frac{1}{e} \int x^{m-2} (d + e x^2)^{q+1} (a + b \operatorname{ArcTan}[c x])^p dx - \frac{d}{e} \int x^{m-2} (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx$$

Program code:

```
Int[x_^m_*(d_+e_.*x_^2)^q_*(a_+b_.*ArcTan[c_.*x_])^p_.,x_Symbol] :=
  1/e*Int[x^(m-2)*(d+e*x^2)^(q+1)*(a+b*ArcTan[c*x])^p,x] -
  d/e*Int[x^(m-2)*(d+e*x^2)^q*(a+b*ArcTan[c*x])^p,x] /;
FreeQ[{a,b,c,d,e},x] && EqQ[e,c^2*d] && IntegersQ[p,2*q] && LtQ[q,-1] && IGtQ[m,1] && NeQ[p,-1]
```

```
Int[x_^m_*(d_+e_.*x_^2)^q_*(a_+b_.*ArcCot[c_.*x_])^p_.,x_Symbol] :=
  1/e*Int[x^(m-2)*(d+e*x^2)^(q+1)*(a+b*ArcCot[c*x])^p,x] -
  d/e*Int[x^(m-2)*(d+e*x^2)^q*(a+b*ArcCot[c*x])^p,x] /;
FreeQ[{a,b,c,d,e},x] && EqQ[e,c^2*d] && IntegersQ[p,2*q] && LtQ[q,-1] && IGtQ[m,1] && NeQ[p,-1]
```

$$\mathbf{2:} \int x^m (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx \text{ when } e = c^2 d \wedge (m \mid p \mid 2q) \in \mathbb{Z} \wedge q < -1 \wedge m < 0 \wedge p \neq -1$$

Derivation: Algebraic expansion

$$\text{Basis: } \frac{1}{d+e x^2} = \frac{1}{d} - \frac{e x^2}{d (d+e x^2)}$$

Rule: If $e = c^2 d \wedge (m \mid p \mid 2q) \in \mathbb{Z} \wedge q < -1 \wedge m < 0 \wedge p \neq -1$, then

$$\int x^m (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx \rightarrow \frac{1}{d} \int x^m (d + e x^2)^{q+1} (a + b \operatorname{ArcTan}[c x])^p dx - \frac{e}{d} \int x^{m+2} (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx$$

Program code:

```
Int[x_^m_*(d_+e_.*x_^2)^q_*(a_+b_.*ArcTan[c_.*x_])^p_,x_Symbol] :=
  1/d*Int[x^m*(d+e*x^2)^(q+1)*(a+b*ArcTan[c*x])^p,x] -
  e/d*Int[x^(m+2)*(d+e*x^2)^q*(a+b*ArcTan[c*x])^p,x] /;
FreeQ[{a,b,c,d,e},x] && EqQ[e,c^2*d] && IntegersQ[p,2*q] && LtQ[q,-1] && ILtQ[m,0] && NeQ[p,-1]
```

```
Int[x_^m_*(d_+e_.*x_^2)^q_*(a_+b_.*ArcCot[c_.*x_])^p_,x_Symbol] :=
  1/d*Int[x^m*(d+e*x^2)^(q+1)*(a+b*ArcCot[c*x])^p,x] -
  e/d*Int[x^(m+2)*(d+e*x^2)^q*(a+b*ArcCot[c*x])^p,x] /;
FreeQ[{a,b,c,d,e},x] && EqQ[e,c^2*d] && IntegersQ[p,2*q] && LtQ[q,-1] && ILtQ[m,0] && NeQ[p,-1]
```

$$\mathbf{3:} \int (f x)^m (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx \text{ when } e = c^2 d \wedge q < -1 \wedge p < -1 \wedge m + 2q + 2 \neq 0$$

Derivation: Integration by parts

Rule: If $e = c^2 d \wedge q < -1 \wedge p < -1 \wedge m + 2q + 2 \neq 0$, then

$$\int (f x)^m (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx \rightarrow \frac{(f x)^m (d + e x^2)^{q+1} (a + b \operatorname{ArcTan}[c x])^{p+1}}{b c d (p+1)} -$$

$$\frac{f m}{b c (p+1)} \int (f x)^{m-1} (d+e x^2)^q (a+b \operatorname{ArcTan}[c x])^{p+1} dx - \frac{c f (m+2 q+2)}{b (p+1)} \int (f x)^{m+1} (d+e x^2)^q (a+b \operatorname{ArcTan}[c x])^{p+1} dx$$

Program code:

```
Int[(f_.**x_)^m_.*(d_+e_.**x_^2)^q_*(a_+b_.*ArcTan[c_.**x_])^p_,x_Symbol] :=
  (f*x)^m*(d+e*x^2)^(q+1)*(a+b*ArcTan[c*x])^(p+1)/(b*c*d*(p+1)) -
  f*m/(b*c*(p+1))*Int[(f*x)^(m-1)*(d+e*x^2)^q*(a+b*ArcTan[c*x])^(p+1),x] -
  c*f*(m+2*q+2)/(b*(p+1))*Int[(f*x)^(m+1)*(d+e*x^2)^q*(a+b*ArcTan[c*x])^(p+1),x] /;
FreeQ[{a,b,c,d,e,f,m},x] && EqQ[e,c^2*d] && RationalQ[m] && LtQ[q,-1] && LtQ[p,-1] && NeQ[m+2*q+2,0]
```

```
Int[(f_.**x_)^m_.*(d_+e_.**x_^2)^q_*(a_+b_.*ArcCot[c_.**x_])^p_,x_Symbol] :=
  -(f*x)^m*(d+e*x^2)^(q+1)*(a+b*ArcCot[c*x])^(p+1)/(b*c*d*(p+1)) +
  f*m/(b*c*(p+1))*Int[(f*x)^(m-1)*(d+e*x^2)^q*(a+b*ArcCot[c*x])^(p+1),x] +
  c*f*(m+2*q+2)/(b*(p+1))*Int[(f*x)^(m+1)*(d+e*x^2)^q*(a+b*ArcCot[c*x])^(p+1),x] /;
FreeQ[{a,b,c,d,e,f,m},x] && EqQ[e,c^2*d] && RationalQ[m] && LtQ[q,-1] && LtQ[p,-1] && NeQ[m+2*q+2,0]
```

$$4. \int x^m (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx \text{ when } e == c^2 d \wedge m \in \mathbb{Z}^+ \wedge m + 2q + 1 \in \mathbb{Z}^-$$

$$1. \int x^m (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx \text{ when } e == c^2 d \wedge m \in \mathbb{Z}^+ \wedge m + 2q + 1 \in \mathbb{Z}^-$$

$$1: \int x^m (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx \text{ when } e == c^2 d \wedge m \in \mathbb{Z}^+ \wedge m + 2q + 1 \in \mathbb{Z}^- \wedge (q \in \mathbb{Z} \vee d > 0)$$

Derivation: Integration by substitution

Basis: If $e == c^2 d \wedge m \in \mathbb{Z} \wedge m + 2q + 1 \in \mathbb{Z} \wedge (q \in \mathbb{Z} \vee d > 0)$, then

$$x^m (d + e x^2)^q == \frac{d^q \operatorname{Sin}[\operatorname{ArcTan}[c x]]^m}{c^{m+1} \operatorname{Cos}[\operatorname{ArcTan}[c x]]^{m+2(q+1)}} \partial_x \operatorname{ArcTan}[c x]$$

Rule: If $e == c^2 d \wedge m \in \mathbb{Z}^+ \wedge m + 2q + 1 \in \mathbb{Z}^- \wedge (q \in \mathbb{Z} \vee d > 0)$, then

$$\int x^m (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx \rightarrow \frac{d^q}{c^{m+1}} \operatorname{Subst}\left[\int \frac{(a + b x)^p \operatorname{Sin}[x]^m}{\operatorname{Cos}[x]^{m+2(q+1)}} dx, x, \operatorname{ArcTan}[c x]\right]$$

Program code:

```
Int[x_^m.*(d_+e_.x_^2)^q_*(a_+b_.ArcTan[c_.x_])^p_,x_Symbol] :=
  d^q/c^(m+1)*Subst[Int[(a+b*x)^p*Sin[x]^m/Cos[x]^(m+2*(q+1)),x],x,ArcTan[c*x] ] /;
FreeQ[{a,b,c,d,e,p},x] && EqQ[e,c^2*d] && IGtQ[m,0] && ILtQ[m+2*q+1,0] && (IntegerQ[q] || GtQ[d,0])
```

$$2: \int x^m (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx \text{ when } e == c^2 d \wedge m \in \mathbb{Z}^+ \wedge m + 2q + 1 \in \mathbb{Z}^- \wedge \neg (q \in \mathbb{Z} \vee d > 0)$$

Derivation: Piecewise constant extraction

$$\text{Basis: If } e == c^2 d, \text{ then } \partial_x \frac{\sqrt{1+c^2 x^2}}{\sqrt{d+e x^2}} == 0$$

Rule: If $e == c^2 d \wedge m \in \mathbb{Z}^+ \wedge m + 2q + 1 \in \mathbb{Z}^- \wedge \neg (q \in \mathbb{Z} \vee d > 0)$, then

$$\int x^m (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx \rightarrow \frac{d^{q+\frac{1}{2}} \sqrt{1+c^2 x^2}}{\sqrt{d+e x^2}} \int x^m (1+c^2 x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx$$

Program code:

```
Int[x_^m_.*(d_+e_.x_^2)^q_*(a_+b_.*ArcTan[c_.x_])^p_,x_Symbol] :=
  d^(q+1/2)*Sqrt[1+c^2*x^2]/Sqrt[d+e*x^2]*Int[x^m*(1+c^2*x^2)^q*(a+b*ArcTan[c*x])^p,x] /;
FreeQ[{a,b,c,d,e,p},x] && EqQ[e,c^2*d] && IGtQ[m,0] && ILtQ[m+2*q+1,0] && Not[IntegerQ[q] || GtQ[d,0]]
```

2. $\int x^m (d + e x^2)^q (a + b \operatorname{ArcCot}[c x])^p dx$ when $e = c^2 d \wedge m \in \mathbb{Z}^+ \wedge m + 2q + 1 \in \mathbb{Z}^-$

1: $\int x^m (d + e x^2)^q (a + b \operatorname{ArcCot}[c x])^p dx$ when $e = c^2 d \wedge m \in \mathbb{Z}^+ \wedge m + 2q + 1 \in \mathbb{Z}^- \wedge q \in \mathbb{Z}$

Derivation: Integration by substitution

Basis: If $e = c^2 d \wedge m \in \mathbb{Z} \wedge q \in \mathbb{Z}$, then $x^m (d + e x^2)^q = -\frac{d^q \cos[\operatorname{ArcCot}[c x]]^m}{c^{m+1} \sin[\operatorname{ArcCot}[c x]]^{m+2(q+1)}} \partial_x \operatorname{ArcCot}[c x]$

Rule: If $e = c^2 d \wedge m \in \mathbb{Z}^+ \wedge m + 2q + 1 \in \mathbb{Z}^- \wedge q \in \mathbb{Z}$, then

$$\int x^m (d + e x^2)^q (a + b \operatorname{ArcCot}[c x])^p dx \rightarrow -\frac{d^q}{c^{m+1}} \operatorname{Subst}\left[\int \frac{(a + b x)^p \cos[x]^m}{\sin[x]^{m+2(q+1)}} dx, x, \operatorname{ArcCot}[c x]\right]$$

Program code:

```
Int[x_^m_.*(d_+e_.x_^2)^q_*(a_+b_.*ArcCot[c_.x_])^p_,x_Symbol] :=
  -d^q/c^(m+1)*Subst[Int[(a+b*x)^p*cos[x]^m/sin[x]^(m+2*(q+1)),x],x,ArcCot[c*x]] /;
FreeQ[{a,b,c,d,e,p},x] && EqQ[e,c^2*d] && IGtQ[m,0] && ILtQ[m+2*q+1,0] && IntegerQ[q]
```

2: $\int x^m (d + e x^2)^q (a + b \operatorname{ArcCot}[c x])^p dx$ when $e = c^2 d \wedge m \in \mathbb{Z}^+ \wedge m + 2q + 1 \in \mathbb{Z}^- \wedge q \notin \mathbb{Z}$

Derivation: Piecewise constant extraction and integration by substitution

Basis: If $e = c^2 d$, then $\partial_x \frac{x \sqrt{\frac{1+c^2 x^2}{c^2 x^2}}}{\sqrt{d+e x^2}} = 0$

Basis: If $m \in \mathbb{Z} \wedge m+2q+1 \in \mathbb{Z} \wedge q \notin \mathbb{Z}$, then

$$x^{m+1} \sqrt{1 + \frac{1}{c^2 x^2}} (1 + c^2 x^2)^{q-\frac{1}{2}} = -\frac{\cos[\text{ArcCot}[c x]]^m}{c^{m+2} \sin[\text{ArcCot}[c x]]^{m+2(q+1)}} \partial_x \text{ArcCot}[c x]$$

Rule: If $e = c^2 d \wedge m \in \mathbb{Z}^+ \wedge m+2q+1 \in \mathbb{Z}^- \wedge q \notin \mathbb{Z}$, then

$$\begin{aligned} \int x^m (d+e x^2)^q (a+b \text{ArcCot}[c x])^p dx &\rightarrow \frac{c^2 d^{q+\frac{1}{2}} x \sqrt{\frac{1+c^2 x^2}{c^2 x^2}}}{\sqrt{d+e x^2}} \int x^{m+1} \sqrt{1 + \frac{1}{c^2 x^2}} (1 + c^2 x^2)^{q-\frac{1}{2}} (a+b \text{ArcCot}[c x])^p dx \\ &\rightarrow -\frac{d^{q+\frac{1}{2}} x \sqrt{\frac{1+c^2 x^2}{c^2 x^2}}}{c^m \sqrt{d+e x^2}} \text{Subst}\left[\int \frac{(a+b x)^p \cos[x]^m}{\sin[x]^{m+2(q+1)}} dx, x, \text{ArcCot}[c x]\right] \end{aligned}$$

Program code:

```
Int[x_^m_.*(d_+e_.x_^2)^q_*(a_+b_.ArcCot[c_.x_])^p_.,x_Symbol] :=
  -d^(q+1/2)*x*Sqrt[(1+c^2*x^2)/(c^2*x^2)]/(c^m*Sqrt[d+e*x^2])*Subst[Int[(a+b*x)^p*cos[x]^m/Sin[x]^(m+2*(q+1)),x],x,ArcCot[c*x]] /;
FreeQ[{a,b,c,d,e,p},x] && EqQ[e,c^2*d] && IGtQ[m,0] && ILtQ[m+2*q+1,0] && Not[IntegerQ[q]]
```

$$3. \int (f x)^m (d+e x^2)^q (a+b \text{ArcTan}[c x]) dx \text{ when } (q \in \mathbb{Z}^+ \wedge \neg (\frac{m-1}{2} \in \mathbb{Z}^- \wedge m+2q+3 > 0)) \vee (\frac{m+1}{2} \in \mathbb{Z}^+ \wedge \neg (q \in \mathbb{Z}^- \wedge m+2q+3 > 0)) \vee (\frac{m+2q+1}{2} \in \mathbb{Z}^- \wedge \frac{m-1}{2} \notin \mathbb{Z}^-)$$

$$1: \int x (d+e x^2)^q (a+b \text{ArcTan}[c x]) dx \text{ when } q \neq -1$$

Derivation: Integration by parts

$$\text{Basis: } x (d+e x^2)^q = \partial_x \frac{(d+e x^2)^{q+1}}{2e(q+1)}$$

Rule: If $q \neq -1$, then

$$\int x (d + e x^2)^q (a + b \operatorname{ArcTan}[c x]) dx \rightarrow \frac{(d + e x^2)^{q+1} (a + b \operatorname{ArcTan}[c x])}{2 e (q + 1)} - \frac{b c}{2 e (q + 1)} \int \frac{(d + e x^2)^{q+1}}{1 + c^2 x^2} dx$$

Program code:

```
Int[x*(d_+e_*x_^2)^q_.*(a_+b_.*ArcTan[c_*x_]),x_Symbol] :=
  (d+e*x^2)^(q+1)*(a+b*ArcTan[c*x])/(2*e*(q+1)) -
  b*c/(2*e*(q+1))*Int[(d+e*x^2)^(q+1)/(1+c^2*x^2),x] /;
FreeQ[{a,b,c,d,e,q},x] && NeQ[q,-1]
```

```
Int[x*(d_+e_*x_^2)^q_.*(a_+b_.*ArcCot[c_*x_]),x_Symbol] :=
  (d+e*x^2)^(q+1)*(a+b*ArcCot[c*x])/(2*e*(q+1)) +
  b*c/(2*e*(q+1))*Int[(d+e*x^2)^(q+1)/(1+c^2*x^2),x] /;
FreeQ[{a,b,c,d,e,q},x] && NeQ[q,-1]
```

2:

$$\int (f x)^m (d + e x^2)^q (a + b \operatorname{ArcTan}[c x]) dx \text{ when } (q \in \mathbb{Z}^+ \wedge \neg (\frac{m-1}{2} \in \mathbb{Z}^- \wedge m+2q+3 > 0)) \vee (\frac{m+1}{2} \in \mathbb{Z}^+ \wedge \neg (q \in \mathbb{Z}^- \wedge m+2q+3 > 0)) \vee (\frac{m+2q+1}{2} \in \mathbb{Z}^- \wedge \frac{m-1}{2} \notin \mathbb{Z}^-)$$

Derivation: Integration by parts

Note: If $(q \in \mathbb{Z}^+ \wedge \neg (\frac{m-1}{2} \in \mathbb{Z}^- \wedge m+2q+3 > 0)) \vee$

$$(\frac{m+1}{2} \in \mathbb{Z}^+ \wedge \neg (q \in \mathbb{Z}^- \wedge m+2q+3 > 0)) \vee (\frac{m+2q+1}{2} \in \mathbb{Z}^- \wedge \frac{m-1}{2} \notin \mathbb{Z}^-)$$

then $\int (f x)^m (d + e x^2)^q dx$ is expressible as an algebraic function not involving logarithms, inverse trig or inverse hyperbolic functions.

Rule: If $(q \in \mathbb{Z}^+ \wedge \neg (\frac{m-1}{2} \in \mathbb{Z}^- \wedge m+2q+3 > 0)) \vee$, let $u = \int (f x)^m (d + e x^2)^q dx$, then

$$(\frac{m+1}{2} \in \mathbb{Z}^+ \wedge \neg (q \in \mathbb{Z}^- \wedge m+2q+3 > 0)) \vee (\frac{m+2q+1}{2} \in \mathbb{Z}^- \wedge \frac{m-1}{2} \notin \mathbb{Z}^-)$$

$$\int (f x)^m (d + e x^2)^q (a + b \operatorname{ArcTan}[c x]) dx \rightarrow u (a + b \operatorname{ArcTan}[c x]) - b c \int \frac{u}{1 + c^2 x^2} dx$$

Program code:

```
Int[(f_.x_)^m_.*(d_.+e_.x_^2)^q_.*(a_.+b_.*ArcTan[c_.x_]),x_Symbol] :=
  With[{u=IntHide[(f*x)^m*(d+e*x^2)^q,x]},
    Dist[a+b*ArcTan[c*x],u,x] - b*c*Int[SimplifyIntegrand[u/(1+c^2*x^2),x],x]] /;
FreeQ[{a,b,c,d,e,f,m,q},x] && (
  IGtQ[q,0] && Not[ILtQ[(m-1)/2,0] && GtQ[m+2*q+3,0]] ||
  IGtQ[(m+1)/2,0] && Not[ILtQ[q,0] && GtQ[m+2*q+3,0]] ||
  ILtQ[(m+2*q+1)/2,0] && Not[ILtQ[(m-1)/2,0]] )
```

```

Int[(f_.*x_)^m_.*(d_+e_.*x_^2)^q_.*(a_+b_.*ArcCot[c_.*x_]),x_Symbol] :=
  With[{u=IntHide[(f*x)^m*(d+e*x^2)^q,x]},
    Dist[a+b*ArcCot[c*x],u,x] + b*c*Int[SimplifyIntegrand[u/(1+c^2*x^2),x],x] /;
  FreeQ[{a,b,c,d,e,f,m,q},x] && (
    IGtQ[q,0] && Not[ILtQ[(m-1)/2,0] && GtQ[m+2*q+3,0]] ||
    IGtQ[(m+1)/2,0] && Not[ILtQ[q,0] && GtQ[m+2*q+3,0]] ||
    ILtQ[(m+2*q+1)/2,0] && Not[ILtQ[(m-1)/2,0]] )

```

4: $\int \frac{x (a + b \operatorname{ArcTan}[c x])^p}{(d + e x^2)^2} dx$ when $p \in \mathbb{Z}^+$

Derivation: Algebraic expansion

$$\text{Basis: } \frac{x}{(d+ex^2)^2} = \frac{1}{4d^2 \sqrt{-\frac{e}{d}} \left(1 - \sqrt{-\frac{e}{d}} x\right)^2} - \frac{1}{4d^2 \sqrt{-\frac{e}{d}} \left(1 + \sqrt{-\frac{e}{d}} x\right)^2}$$

Rule: If $p \in \mathbb{Z}^+$, then

$$\int \frac{x (a + b \operatorname{ArcTan}[c x])^p}{(d + e x^2)^2} dx \rightarrow \frac{1}{4d^2 \sqrt{-\frac{e}{d}}} \int \frac{(a + b \operatorname{ArcTan}[c x])^p}{\left(1 - \sqrt{-\frac{e}{d}} x\right)^2} dx - \frac{1}{4d^2 \sqrt{-\frac{e}{d}}} \int \frac{(a + b \operatorname{ArcTan}[c x])^p}{\left(1 + \sqrt{-\frac{e}{d}} x\right)^2} dx$$

Program code:

```

Int[x_*(a_+b_.*ArcTan[c_.*x_])^p_./(d_+e_.*x_^2),x_Symbol] :=
  1/(4*d^2*Rt[-e/d,2])*Int[(a+b*ArcTan[c*x])^p/(1-Rt[-e/d,2]*x)^2,x] -
  1/(4*d^2*Rt[-e/d,2])*Int[(a+b*ArcTan[c*x])^p/(1+Rt[-e/d,2]*x)^2,x] /;
  FreeQ[{a,b,c,d,e},x] && IGtQ[p,0]

```

```

Int[x*(a_.+b_.*ArcCot[c_.*x_])^p_./(d_+e_.*x_^2)^2,x_Symbol] :=
  1/(4*d^2*Rt[-e/d,2])*Int[(a+b*ArcCot[c*x])^p/(1-Rt[-e/d,2]*x)^2,x] -
  1/(4*d^2*Rt[-e/d,2])*Int[(a+b*ArcCot[c*x])^p/(1+Rt[-e/d,2]*x)^2,x] /;
FreeQ[{a,b,c,d,e},x] && IGtQ[p,0]

```

5: $\int (f x)^m (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx$ when $q \in \mathbb{Z} \wedge p \in \mathbb{Z}^+ \wedge (p = 1 \vee m \in \mathbb{Z})$

Derivation: Algebraic expansion

Rule: If $q \in \mathbb{Z} \wedge p \in \mathbb{Z}^+ \wedge (p = 1 \vee m \in \mathbb{Z})$, then

$$\int (f x)^m (d + e x^2)^q (a + b \operatorname{ArcTan}[c x])^p dx \rightarrow \int (a + b \operatorname{ArcTan}[c x])^p \operatorname{ExpandIntegrand}[(f x)^m (d + e x^2)^q, x] dx$$

Program code:

```

Int[(f_.*x_)^m_.*(d_+e_.*x_^2)^q_.*(a_.+b_.*ArcTan[c_.*x_])^p_.,x_Symbol] :=
  With[{u=ExpandIntegrand[(a+b*ArcTan[c*x])^p,(f*x)^m*(d+e*x^2)^q,x]},
    Int[u,x] /;
    SumQ[u] /;
    FreeQ[{a,b,c,d,e,f,m},x] && IntegerQ[q] && IGtQ[p,0] && (EqQ[p,1] && GtQ[q,0] || IntegerQ[m])

```

```

Int[(f_.*x_)^m_.*(d_+e_.*x_^2)^q_.*(a_.+b_.*ArcCot[c_.*x_])^p_.,x_Symbol] :=
  With[{u=ExpandIntegrand[(a+b*ArcCot[c*x])^p,(f*x)^m*(d+e*x^2)^q,x]},
    Int[u,x] /;
    SumQ[u] /;
    FreeQ[{a,b,c,d,e,f,m},x] && IntegerQ[q] && IGtQ[p,0] && (EqQ[p,1] && GtQ[q,0] || IntegerQ[m])

```

$$6: \int (f x)^m (d + e x^2)^q (a + b \operatorname{ArcTan}[c x]) dx$$

Derivation: Algebraic expansion

Rule:

$$\int (f x)^m (d + e x^2)^q (a + b \operatorname{ArcTan}[c x]) dx \rightarrow a \int (f x)^m (d + e x^2)^q dx + b \int (f x)^m (d + e x^2)^q \operatorname{ArcTan}[c x] dx$$

Program code:

```
Int[(f_.*x_)^m_.*(d_+e_.*x_^2)^q_.*(a_+b_.*ArcTan[c_.*x_]),x_Symbol] :=
  a*Int[(f*x)^m*(d+e*x^2)^q,x] + b*Int[(f*x)^m*(d+e*x^2)^q*ArcTan[c*x],x] /;
FreeQ[{a,b,c,d,e,f,m,q},x]
```

```
Int[(f_.*x_)^m_.*(d_+e_.*x_^2)^q_.*(a_+b_.*ArcCot[c_.*x_]),x_Symbol] :=
  a*Int[(f*x)^m*(d+e*x^2)^q,x] + b*Int[(f*x)^m*(d+e*x^2)^q*ArcCot[c*x],x] /;
FreeQ[{a,b,c,d,e,f,m,q},x]
```

$$7. \int \frac{u (a + b \operatorname{ArcTan}[c x])^p}{d + e x^2} dx \text{ when } e = c^2 d$$

$$1: \int \frac{(f + g x)^m (a + b \operatorname{ArcTan}[c x])^p}{d + e x^2} dx \text{ when } p \in \mathbb{Z}^+ \wedge e = c^2 d \wedge m \in \mathbb{Z}^+$$

Derivation: Algebraic expansion

Rule: If $p \in \mathbb{Z}^+ \wedge e = c^2 d \wedge m \in \mathbb{Z}^+$, then

$$\int \frac{(f + g x)^m (a + b \operatorname{ArcTan}[c x])^p}{d + e x^2} dx \rightarrow \int \frac{(a + b \operatorname{ArcTan}[c x])^p}{d + e x^2} \operatorname{ExpandIntegrand}[(f + g x)^m, x] dx$$

Program code:

```
Int[(f_+g_.**x_)^m_.*(a_.+b_.*ArcTan[c_.**x_])^p_./(d_+e_.**x_^2),x_Symbol] :=
  Int[ExpandIntegrand[(a+b*ArcTan[c*x])^p/(d+e*x^2),(f+g*x)^m,x],x] /;
FreeQ[{a,b,c,d,e,f,g},x] && IGtQ[p,0] && EqQ[e,c^2*d] && IGtQ[m,0]
```

```
Int[(f_+g_.**x_)^m_.*(a_.+b_.*ArcCot[c_.**x_])^p_./(d_+e_.**x_^2),x_Symbol] :=
  Int[ExpandIntegrand[(a+b*ArcCot[c*x])^p/(d+e*x^2),(f+g*x)^m,x],x] /;
FreeQ[{a,b,c,d,e,f,g},x] && IGtQ[p,0] && EqQ[e,c^2*d] && IGtQ[m,0]
```

$$2. \int \frac{\text{ArcTanh}[u] (a + b \text{ArcTan}[c x])^p}{d + e x^2} dx \text{ when } p \in \mathbb{Z}^+ \wedge e = c^2 d$$

$$1: \int \frac{\text{ArcTanh}[u] (a + b \text{ArcTan}[c x])^p}{d + e x^2} dx \text{ when } p \in \mathbb{Z}^+ \wedge e = c^2 d \wedge u^2 = \left(1 - \frac{2I}{I+cx}\right)^2$$

Derivation: Algebraic expansion

$$\text{Basis: } \text{ArcTanh}[z] = \frac{1}{2} \text{Log}[1+z] - \frac{1}{2} \text{Log}[1-z]$$

$$\text{Basis: } \text{ArcCoth}[z] = \frac{1}{2} \text{Log}\left[1 + \frac{1}{z}\right] - \frac{1}{2} \text{Log}\left[1 - \frac{1}{z}\right]$$

Rule: If $p \in \mathbb{Z}^+ \wedge e = c^2 d \wedge u^2 = \left(1 - \frac{2I}{I+cx}\right)^2$, then

$$\int \frac{\text{ArcTanh}[u] (a + b \text{ArcTan}[c x])^p}{d + e x^2} dx \rightarrow \frac{1}{2} \int \frac{\text{Log}[1+u] (a + b \text{ArcTan}[c x])^p}{d + e x^2} dx - \frac{1}{2} \int \frac{\text{Log}[1-u] (a + b \text{ArcTan}[c x])^p}{d + e x^2} dx$$

Program code:

```
Int[ArcTanh[u_]*(a_+b_.*ArcTan[c_.*x_])^p_./(d_+e_.*x_^2),x_Symbol] :=
  1/2*Int[Log[1+u]*(a+b*ArcTan[c*x])^p/(d+e*x^2),x] -
  1/2*Int[Log[1-u]*(a+b*ArcTan[c*x])^p/(d+e*x^2),x] /;
FreeQ[{a,b,c,d,e},x] && IGtQ[p,0] && EqQ[e,c^2*d] && EqQ[u^2-(1-2*I/(I+c*x))^2,0]
```

```
Int[ArcCoth[u_]*(a_+b_.*ArcCot[c_.*x_])^p_./(d_+e_.*x_^2),x_Symbol] :=
  1/2*Int[Log[SimplifyIntegrand[1+1/u,x]]*(a+b*ArcCot[c*x])^p/(d+e*x^2),x] -
  1/2*Int[Log[SimplifyIntegrand[1-1/u,x]]*(a+b*ArcCot[c*x])^p/(d+e*x^2),x] /;
FreeQ[{a,b,c,d,e},x] && IGtQ[p,0] && EqQ[e,c^2*d] && EqQ[u^2-(1-2*I/(I+c*x))^2,0]
```


$$2: \int \frac{\text{ArcTanh}[u] (a + b \text{ArcTan}[c x])^p}{d + e x^2} dx \text{ when } p \in \mathbb{Z}^+ \wedge e = c^2 d \wedge u^2 = \left(1 - \frac{2I}{1-cx}\right)^2$$

Derivation: Algebraic expansion

$$\text{Basis: } \text{ArcTanh}[z] = \frac{1}{2} \text{Log}[1+z] - \frac{1}{2} \text{Log}[1-z]$$

$$\text{Basis: } \text{ArcCoth}[z] = \frac{1}{2} \text{Log}\left[1 + \frac{1}{z}\right] - \frac{1}{2} \text{Log}\left[1 - \frac{1}{z}\right]$$

Rule: If $p \in \mathbb{Z}^+ \wedge e = c^2 d \wedge u^2 = \left(1 - \frac{2I}{1-cx}\right)^2$, then

$$\int \frac{\text{ArcTanh}[u] (a + b \text{ArcTan}[c x])^p}{d + e x^2} dx \rightarrow \frac{1}{2} \int \frac{\text{Log}[1+u] (a + b \text{ArcTan}[c x])^p}{d + e x^2} dx - \frac{1}{2} \int \frac{\text{Log}[1-u] (a + b \text{ArcTan}[c x])^p}{d + e x^2} dx$$

Program code:

```
Int[ArcTanh[u_]*(a_+b_*ArcTan[c_*x_])^p_/(d_+e_*x_^2),x_Symbol] :=
  1/2*Int[Log[1+u]*(a+b*ArcTan[c*x])^p/(d+e*x^2),x] -
  1/2*Int[Log[1-u]*(a+b*ArcTan[c*x])^p/(d+e*x^2),x] /;
FreeQ[{a,b,c,d,e},x] && IGtQ[p,0] && EqQ[e,c^2*d] && EqQ[u^2-(1-2*I/(1-c*x))^2,0]
```

```
Int[ArcCoth[u_]*(a_+b_*ArcCot[c_*x_])^p_/(d_+e_*x_^2),x_Symbol] :=
  1/2*Int[Log[SimplifyIntegrand[1+1/u,x]]*(a+b*ArcCot[c*x])^p/(d+e*x^2),x] -
  1/2*Int[Log[SimplifyIntegrand[1-1/u,x]]*(a+b*ArcCot[c*x])^p/(d+e*x^2),x] /;
FreeQ[{a,b,c,d,e},x] && IGtQ[p,0] && EqQ[e,c^2*d] && EqQ[u^2-(1-2*I/(1-c*x))^2,0]
```

$$3. \int \frac{(a + b \operatorname{ArcTan}[c x])^p \operatorname{Log}[u]}{d + e x^2} dx \text{ when } p \in \mathbb{Z}^+ \wedge e = c^2 d$$

$$1: \int \frac{(a + b \operatorname{ArcTan}[c x])^p \operatorname{Log}[f + g x]}{d + e x^2} dx \text{ when } p \in \mathbb{Z}^+ \wedge e = c^2 d \wedge c^2 f^2 + g^2 = 0$$

Derivation: Integration by parts

Basis: If $e = c^2 d$, then $\frac{(a+b \operatorname{ArcTan}[c x])^p}{d+e x^2} = \partial_x \frac{(a+b \operatorname{ArcTan}[c x])^{p+1}}{b c d (p+1)}$

Rule: If $p \in \mathbb{Z}^+ \wedge e = c^2 d \wedge c^2 f^2 + g^2 = 0$, then

$$\int \frac{(a + b \operatorname{ArcTan}[c x])^p \operatorname{Log}[f + g x]}{d + e x^2} dx \rightarrow \frac{(a + b \operatorname{ArcTan}[c x])^{p+1} \operatorname{Log}[f + g x]}{b c d (p+1)} - \frac{g}{b c d (p+1)} \int \frac{(a + b \operatorname{ArcTan}[c x])^{p+1}}{f + g x} dx$$

Program code:

```
Int[(a_.+b_.*ArcTan[c_.*x_])^p_.*Log[f_+g_.*x_]/(d_+e_.*x_^2),x_Symbol] :=
  (a+b*ArcTan[c*x])^(p+1)*Log[f+g*x]/(b*c*d*(p+1)) -
  g/(b*c*d*(p+1))*Int[(a+b*ArcTan[c*x])^(p+1)/(f+g*x),x] /;
FreeQ[{a,b,c,d,e,f,g},x] && IGtQ[p,0] && EqQ[e,c^2*d] && EqQ[c^2*f^2+g^2,0]
```

```
Int[(a_.+b_.*ArcCot[c_.*x_])^p_.*Log[f_+g_.*x_]/(d_+e_.*x_^2),x_Symbol] :=
  (a+b*ArcCot[c*x])^(p+1)*Log[f+g*x]/(b*c*d*(p+1)) -
  g/(b*c*d*(p+1))*Int[(a+b*ArcCot[c*x])^(p+1)/(f+g*x),x] /;
FreeQ[{a,b,c,d,e,f,g},x] && IGtQ[p,0] && EqQ[e,c^2*d] && EqQ[c^2*f^2+g^2,0]
```

$$2: \int \frac{(a + b \operatorname{ArcTan}[c x])^p \operatorname{Log}[u]}{d + e x^2} dx \text{ when } p \in \mathbb{Z}^+ \wedge e = c^2 d \wedge (1 - u)^2 = \left(1 - \frac{2I}{I + c x}\right)^2$$

Derivation: Integration by parts

Rule: If $p \in \mathbb{Z}^+ \wedge e = c^2 d \wedge (1 - u)^2 = \left(1 - \frac{2I}{I + c x}\right)^2$, then

$$\int \frac{(a + b \operatorname{ArcTan}[c x])^p \operatorname{Log}[u]}{d + e x^2} dx \rightarrow \frac{i (a + b \operatorname{ArcTan}[c x])^p \operatorname{PolyLog}[2, 1 - u]}{2 c d} - \frac{b p i}{2} \int \frac{(a + b \operatorname{ArcTan}[c x])^{p-1} \operatorname{PolyLog}[2, 1 - u]}{d + e x^2} dx$$

Program code:

```
Int[(a_.+b_.*ArcTan[c_.*x_])^p_.*Log[u_] / (d_+e_.*x_^2), x_Symbol] :=
  I*(a+b*ArcTan[c*x])^p*PolyLog[2,1-u]/(2*c*d) -
  b*p*I/2*Int[(a+b*ArcTan[c*x])^(p-1)*PolyLog[2,1-u]/(d+e*x^2), x] /;
FreeQ[{a,b,c,d,e}, x] && IGtQ[p, 0] && EqQ[e, c^2*d] && EqQ[(1-u)^2 - (1-2*I/(I+c*x))^2, 0]
```

```
Int[(a_.+b_.*ArcCot[c_.*x_])^p_.*Log[u_] / (d_+e_.*x_^2), x_Symbol] :=
  I*(a+b*ArcCot[c*x])^p*PolyLog[2,1-u]/(2*c*d) +
  b*p*I/2*Int[(a+b*ArcCot[c*x])^(p-1)*PolyLog[2,1-u]/(d+e*x^2), x] /;
FreeQ[{a,b,c,d,e}, x] && IGtQ[p, 0] && EqQ[e, c^2*d] && EqQ[(1-u)^2 - (1-2*I/(I+c*x))^2, 0]
```

3: $\int \frac{(a + b \operatorname{ArcTan}[c x])^p \operatorname{Log}[u]}{d + e x^2} dx$ when $p \in \mathbb{Z}^+ \wedge e = c^2 d \wedge (1-u)^2 = \left(1 - \frac{2I}{I-cx}\right)^2$

Derivation: Integration by parts

Rule: If $p \in \mathbb{Z}^+ \wedge e = c^2 d \wedge (1-u)^2 = \left(1 - \frac{2I}{I-cx}\right)^2$, then

$$\int \frac{(a + b \operatorname{ArcTan}[c x])^p \operatorname{Log}[u]}{d + e x^2} dx \rightarrow -\frac{i (a + b \operatorname{ArcTan}[c x])^p \operatorname{PolyLog}[2, 1 - u]}{2 c d} + \frac{b p i}{2} \int \frac{(a + b \operatorname{ArcTan}[c x])^{p-1} \operatorname{PolyLog}[2, 1 - u]}{d + e x^2} dx$$

Program code:

```
Int[(a_.+b_.*ArcTan[c_.*x_])^p_.*Log[u_] / (d_+e_.*x_^2), x_Symbol] :=
  -I*(a+b*ArcTan[c*x])^p*PolyLog[2,1-u]/(2*c*d) +
  b*p*I/2*Int[(a+b*ArcTan[c*x])^(p-1)*PolyLog[2,1-u]/(d+e*x^2), x] /;
FreeQ[{a,b,c,d,e}, x] && IGtQ[p, 0] && EqQ[e, c^2*d] && EqQ[(1-u)^2 - (1-2*I/(I-c*x))^2, 0]
```

```
Int[(a_+b_.*ArcCot[c_.*x_])^p_.*Log[u_]/(d_+e_.*x_^2),x_Symbol] :=
  -I*(a+b*ArcCot[c*x])^p*PolyLog[2,1-u]/(2*c*d) -
  b*p*I/2*Int[(a+b*ArcCot[c*x])^(p-1)*PolyLog[2,1-u]/(d+e*x^2),x] /;
FreeQ[{a,b,c,d,e},x] && IGtQ[p,0] && EqQ[e,c^2*d] && EqQ[(1-u)^2-(1-2*I/(I-c*x))^2,0]
```

$$4. \int \frac{(a + b \operatorname{ArcTan}[c x])^p \operatorname{PolyLog}[k, u]}{d + e x^2} dx \text{ when } p \in \mathbb{Z}^+ \wedge e = c^2 d$$

$$1: \int \frac{(a + b \operatorname{ArcTan}[c x])^p \operatorname{PolyLog}[k, u]}{d + e x^2} dx \text{ when } p \in \mathbb{Z}^+ \wedge e = c^2 d \wedge u^2 = \left(1 - \frac{2I}{I+cx}\right)^2$$

Derivation: Integration by parts

Rule: If $p \in \mathbb{Z}^+ \wedge e = c^2 d \wedge u^2 = \left(1 - \frac{2I}{I+cx}\right)^2$, then

$$\int \frac{(a + b \operatorname{ArcTan}[c x])^p \operatorname{PolyLog}[k, u]}{d + e x^2} dx \rightarrow -\frac{i (a + b \operatorname{ArcTan}[c x])^p \operatorname{PolyLog}[k+1, u]}{2 c d} + \frac{b p i}{2} \int \frac{(a + b \operatorname{ArcTan}[c x])^{p-1} \operatorname{PolyLog}[k+1, u]}{d + e x^2} dx$$

Program code:

```
Int[(a_+b_.*ArcTan[c_.*x_])^p_.*PolyLog[k_,u_]/(d_+e_.*x_^2),x_Symbol] :=
  -I*(a+b*ArcTan[c*x])^p*PolyLog[k+1,u]/(2*c*d) +
  b*p*I/2*Int[(a+b*ArcTan[c*x])^(p-1)*PolyLog[k+1,u]/(d+e*x^2),x] /;
FreeQ[{a,b,c,d,e,k},x] && IGtQ[p,0] && EqQ[e,c^2*d] && EqQ[u^2-(1-2*I/(I+c*x))^2,0]
```

```
Int[(a_+b_.*ArcCot[c_.*x_])^p_.*PolyLog[k_,u_]/(d_+e_.*x_^2),x_Symbol] :=
  -I*(a+b*ArcCot[c*x])^p*PolyLog[k+1,u]/(2*c*d) -
  b*p*I/2*Int[(a+b*ArcCot[c*x])^(p-1)*PolyLog[k+1,u]/(d+e*x^2),x] /;
FreeQ[{a,b,c,d,e,k},x] && IGtQ[p,0] && EqQ[e,c^2*d] && EqQ[u^2-(1-2*I/(I+c*x))^2,0]
```

$$2: \int \frac{(a + b \operatorname{ArcTan}[c x])^p \operatorname{PolyLog}[k, u]}{d + e x^2} dx \text{ when } p \in \mathbb{Z}^+ \wedge e = c^2 d \wedge u^2 = \left(1 - \frac{2I}{1-cx}\right)^2$$

Derivation: Integration by parts

Rule: If $p \in \mathbb{Z}^+ \wedge e = c^2 d \wedge u^2 = \left(1 - \frac{2I}{1-cx}\right)^2$, then

$$\int \frac{(a + b \operatorname{ArcTan}[c x])^p \operatorname{PolyLog}[k, u]}{d + e x^2} dx \rightarrow \frac{\frac{1}{2} (a + b \operatorname{ArcTan}[c x])^p \operatorname{PolyLog}[k+1, u]}{2 c d} - \frac{b p \frac{1}{2}}{2} \int \frac{(a + b \operatorname{ArcTan}[c x])^{p-1} \operatorname{PolyLog}[k+1, u]}{d + e x^2} dx$$

Program code:

```
Int[(a_.+b_.*ArcTan[c_.*x_])^p_.*PolyLog[k_,u_]/(d_+e_.*x_^2),x_Symbol] :=
  I*(a+b*ArcTan[c*x])^p*PolyLog[k+1,u]/(2*c*d) -
  b*p*I/2*Int[(a+b*ArcTan[c*x])^(p-1)*PolyLog[k+1,u]/(d+e*x^2),x] /;
FreeQ[{a,b,c,d,e,k},x] && IGtQ[p,0] && EqQ[e,c^2*d] && EqQ[u^2-(1-2*I/(1-c*x))^2,0]
```

```
Int[(a_.+b_.*ArcCot[c_.*x_])^p_.*PolyLog[k_,u_]/(d_+e_.*x_^2),x_Symbol] :=
  I*(a+b*ArcCot[c*x])^p*PolyLog[k+1,u]/(2*c*d) +
  b*p*I/2*Int[(a+b*ArcCot[c*x])^(p-1)*PolyLog[k+1,u]/(d+e*x^2),x] /;
FreeQ[{a,b,c,d,e,k},x] && IGtQ[p,0] && EqQ[e,c^2*d] && EqQ[u^2-(1-2*I/(1-c*x))^2,0]
```

$$5. \int \frac{(a + b \operatorname{ArcCot}[c x])^q (a + b \operatorname{ArcTan}[c x])^p}{d + e x^2} dx \text{ when } e = c^2 d$$

$$1: \int \frac{1}{(d + e x^2) (a + b \operatorname{ArcCot}[c x]) (a + b \operatorname{ArcTan}[c x])} dx \text{ when } e = c^2 d$$

Rule: If $e = c^2 d$, then

$$\int \frac{1}{(d + e x^2) (a + b \operatorname{ArcCot}[c x]) (a + b \operatorname{ArcTan}[c x])} dx \rightarrow \frac{-\operatorname{Log}[a + b \operatorname{ArcCot}[c x]] + \operatorname{Log}[a + b \operatorname{ArcTan}[c x]]}{b c d (2 a + b \operatorname{ArcCot}[c x] + b \operatorname{ArcTan}[c x])}$$

Program code:

```
Int[1/((d_+e_.x_^2)*(a_+b_.ArcCot[c_.x_])*(a_+b_.ArcTan[c_.x_])),x_Symbol] :=
  (-Log[a+b*ArcCot[c*x]]+Log[a+b*ArcTan[c*x]])/(b*c*d*(2*a+b*ArcCot[c*x]+b*ArcTan[c*x])) /;
FreeQ[{a,b,c,d,e},x] && EqQ[e,c^2*d]
```

$$2: \int \frac{(a + b \operatorname{ArcCot}[c x])^q (a + b \operatorname{ArcTan}[c x])^p}{d + e x^2} dx \text{ when } e = c^2 d \wedge (p \mid q) \in \mathbb{Z} \wedge 0 < p \leq q$$

Derivation: Integration by parts

Rule: If $e = c^2 d \wedge (p \mid q) \in \mathbb{Z} \wedge 0 < p \leq q$, then

$$\int \frac{(a + b \operatorname{ArcCot}[c x])^q (a + b \operatorname{ArcTan}[c x])^p}{d + e x^2} dx \rightarrow -\frac{(a + b \operatorname{ArcCot}[c x])^{q+1} (a + b \operatorname{ArcTan}[c x])^p}{b c d (q + 1)} + \frac{p}{q + 1} \int \frac{(a + b \operatorname{ArcCot}[c x])^{q+1} (a + b \operatorname{ArcTan}[c x])^{p-1}}{d + e x^2} dx$$

Program code:

```
Int[(a_+b_.ArcCot[c_.x_])^q_.*(a_+b_.ArcTan[c_.x_])^p_./(d_+e_.x_^2),x_Symbol] :=
  -(a+b*ArcCot[c*x])^(q+1)*(a+b*ArcTan[c*x])^p/(b*c*d*(q+1)) +
  p/(q+1)*Int[(a+b*ArcCot[c*x])^(q+1)*(a+b*ArcTan[c*x])^(p-1)/(d+e*x^2),x] /;
FreeQ[{a,b,c,d,e},x] && EqQ[e,c^2*d] && IGtQ[p,0] && IGeQ[q,p]
```

```

Int[(a_+b_*ArcTan[c_*x_])^q_.*(a_+b_*ArcCot[c_*x_])^p_./(d_+e_*x_^2),x_Symbol] :=
  (a+b*ArcTan[c*x])^(q+1)*(a+b*ArcCot[c*x])^p/(b*c*d*(q+1)) +
  p/(q+1)*Int[(a+b*ArcTan[c*x])^(q+1)*(a+b*ArcCot[c*x])^(p-1)/(d+e*x^2),x] /;
FreeQ[{a,b,c,d,e},x] && EqQ[e,c^2*d] && IGtQ[p,0] && IGeQ[q,p]

```

8: $\int \frac{\text{ArcTan}[a x]}{c + d x^n} dx$ when $n \in \mathbb{Z} \wedge \neg (n = 2 \wedge d = a^2 c)$

Derivation: Algebraic expansion

Basis: $\text{ArcTan}[z] = \frac{1}{2} i \text{Log}[1 - i z] - \frac{1}{2} i \text{Log}[1 + i z]$

Basis: $\text{ArcCot}[z] = \frac{1}{2} i \text{Log}[1 - \frac{i}{z}] - \frac{1}{2} i \text{Log}[1 + \frac{i}{z}]$

Rule: If $n \in \mathbb{Z} \wedge \neg (n = 2 \wedge d = a^2 c)$, then

$$\int \frac{\text{ArcTan}[a x]}{c + d x^n} dx \rightarrow \frac{i}{2} \int \frac{\text{Log}[1 - i a x]}{c + d x^n} dx - \frac{i}{2} \int \frac{\text{Log}[1 + i a x]}{c + d x^n} dx$$

Program code:

```

Int[ArcTan[a_*x_]/(c_+d_*x_^n_),x_Symbol] :=
  I/2*Int[Log[1-I*a*x]/(c+d*x^n),x] -
  I/2*Int[Log[1+I*a*x]/(c+d*x^n),x] /;
FreeQ[{a,c,d},x] && IntegerQ[n] && Not[EqQ[n,2] && EqQ[d,a^2*c]]

```

```

Int[ArcCot[a_*x_]/(c_+d_*x_^n_),x_Symbol] :=
  I/2*Int[Log[1-I/(a*x)]/(c+d*x^n),x] -
  I/2*Int[Log[1+I/(a*x)]/(c+d*x^n),x] /;
FreeQ[{a,c,d},x] && IntegerQ[n] && Not[EqQ[n,2] && EqQ[d,a^2*c]]

```

9. $\int \frac{\text{Log}[d x^m] (a + b \text{ArcTan}[c x^n])}{x} dx$

$$1: \int \frac{\text{Log}[d x^m] \text{ArcTan}[c x^n]}{x} dx$$

Derivation: Algebraic expansion

$$\text{Basis: } \text{ArcTan}[c x^n] = \frac{i}{2} \text{Log}[1 - i c x^n] - \frac{i}{2} \text{Log}[1 + i c x^n]$$

Rule:

$$\int \frac{\text{Log}[d x^m] \text{ArcTan}[c x^n]}{x} dx \rightarrow \frac{i}{2} \int \frac{\text{Log}[d x^m] \text{Log}[1 - i c x^n]}{x} dx - \frac{i}{2} \int \frac{\text{Log}[d x^m] \text{Log}[1 + i c x^n]}{x} dx$$

Program code:

```
Int[Log[d_.**x_^m_.]*ArcTan[c_.**x_^n_.]/x_,x_Symbol] :=
  I/2*Int[Log[d*x^m]*Log[1-I*c*x^n]/x,x] - I/2*Int[Log[d*x^m]*Log[1+I*c*x^n]/x,x] /;
FreeQ[{c,d,m,n},x]
```

```
Int[Log[d_.**x_^m_.]*ArcCot[c_.**x_^n_.]/x_,x_Symbol] :=
  I/2*Int[Log[d*x^m]*Log[1-I/(c*x^n)]/x,x] - I/2*Int[Log[d*x^m]*Log[1+I/(c*x^n)]/x,x] /;
FreeQ[{c,d,m,n},x]
```


2: $\int \frac{\text{Log}[d x^m] (a + b \text{ArcTan}[c x^n])}{x} dx$

Derivation: Algebraic expansion

Rule:

$$\int \frac{\text{Log}[d x^m] (a + b \text{ArcTan}[c x^n])}{x} dx \rightarrow a \int \frac{\text{Log}[d x^m]}{x} dx + b \int \frac{\text{Log}[d x^m] \text{ArcTan}[c x^n]}{x} dx$$

Program code:

```
Int[Log[d_.**x_^m_.]*(a_+b_.*ArcTan[c_.**x_^n_.])/x_,x_Symbol] :=
  a*Int[Log[d*x^m]/x,x] + b*Int[(Log[d*x^m]*ArcTan[c*x^n])/x,x] /;
FreeQ[{a,b,c,d,m,n},x]
```

```
Int[Log[d_.**x_^m_.]*(a_+b_.*ArcCot[c_.**x_^n_.])/x_,x_Symbol] :=
  a*Int[Log[d*x^m]/x,x] + b*Int[(Log[d*x^m]*ArcCot[c*x^n])/x,x] /;
FreeQ[{a,b,c,d,m,n},x]
```

$$10. \int u (d + e \operatorname{Log}[f + g x^2]) (a + b \operatorname{ArcTan}[c x])^p dx$$

$$1: \int (d + e \operatorname{Log}[f + g x^2]) (a + b \operatorname{ArcTan}[c x]) dx$$

Derivation: Integration by parts

Rule:

$$\int (d + e \operatorname{Log}[f + g x^2]) (a + b \operatorname{ArcTan}[c x]) dx \rightarrow$$

$$x (d + e \operatorname{Log}[f + g x^2]) (a + b \operatorname{ArcTan}[c x]) - 2 e g \int \frac{x^2 (a + b \operatorname{ArcTan}[c x])}{f + g x^2} dx - b c \int \frac{x (d + e \operatorname{Log}[f + g x^2])}{1 + c^2 x^2} dx$$

Program code:

```
Int[(d_.+e_.*Log[f_.+g_.*x^2])*(a_.+b_.*ArcTan[c_.*x_]),x_Symbol] :=
  x*(d+e*Log[f+g*x^2])*(a+b*ArcTan[c*x]) -
  2*e*g*Int[x^2*(a+b*ArcTan[c*x])/(f+g*x^2),x] -
  b*c*Int[x*(d+e*Log[f+g*x^2])/(1+c^2*x^2),x] /;
FreeQ[{a,b,c,d,e,f,g},x]
```

```
Int[(d_.+e_.*Log[f_.+g_.*x^2])*(a_.+b_.*ArcCot[c_.*x_]),x_Symbol] :=
  x*(d+e*Log[f+g*x^2])*(a+b*ArcCot[c*x]) -
  2*e*g*Int[x^2*(a+b*ArcCot[c*x])/(f+g*x^2),x] +
  b*c*Int[x*(d+e*Log[f+g*x^2])/(1+c^2*x^2),x] /;
FreeQ[{a,b,c,d,e,f,g},x]
```

$$2. \int x^m (d + e \operatorname{Log}[f + g x^2]) (a + b \operatorname{ArcTan}[c x]) dx$$

$$1. \int \frac{(d + e \operatorname{Log}[f + g x^2]) (a + b \operatorname{ArcTan}[c x])}{x} dx$$

$$1. \int \frac{\operatorname{Log}[f + g x^2] (a + b \operatorname{ArcTan}[c x])}{x} dx$$

$$1. \int \frac{\text{Log}[f + g x^2] \text{ArcTan}[c x]}{x} dx \text{ when } c^2 f + g = 0$$

$$1: \int \frac{\text{Log}[f + g x^2] \text{ArcTan}[c x]}{x} dx \text{ when } g = c^2 f$$

Derivation: Piecewise constant extraction and algebraic simplification

Basis: If $g = c^2 f$, then $\partial_x (\text{Log}[f + g x^2] - \text{Log}[1 - i c x] - \text{Log}[1 + i c x]) = 0$

Basis: $(\text{Log}[1 - i c x] + \text{Log}[1 + i c x]) \text{ArcTan}[c x] = \frac{i}{2} \text{Log}[1 - i c x]^2 - \frac{i}{2} \text{Log}[1 + i c x]^2$

Rule: If $g = c^2 f$, then

$$\int \frac{\text{Log}[f + g x^2] \text{ArcTan}[c x]}{x} dx \rightarrow$$

$$(\text{Log}[f + g x^2] - \text{Log}[1 - i c x] - \text{Log}[1 + i c x]) \int \frac{\text{ArcTan}[c x]}{x} dx + \int \frac{(\text{Log}[1 - i c x] + \text{Log}[1 + i c x]) \text{ArcTan}[c x]}{x} dx \rightarrow$$

$$(\text{Log}[f + g x^2] - \text{Log}[1 - i c x] - \text{Log}[1 + i c x]) \int \frac{\text{ArcTan}[c x]}{x} dx + \frac{i}{2} \int \frac{\text{Log}[1 - i c x]^2}{x} dx - \frac{i}{2} \int \frac{\text{Log}[1 + i c x]^2}{x} dx$$

```
Int[Log[f_.+g_.*x^2]*ArcTan[c_.*x]/x_,x_Symbol] :=
  (Log[f+g*x^2]-Log[1-I*c*x]-Log[1+I*c*x])*Int[ArcTan[c*x]/x,x] + I/2*Int[Log[1-I*c*x]^2/x,x] - I/2*Int[Log[1+I*c*x]^2/x,x] /;
FreeQ[{c,f,g},x] && EqQ[g,c^2*f]
```

$$2: \int \frac{\text{Log}[f + g x^2] \text{ArcCot}[c x]}{x} dx \text{ when } g = c^2 f$$

Derivation: Piecewise constant extraction and algebraic simplification

Basis: If $g = c^2 f$, then $\partial_x (\text{Log}[f + g x^2] - \text{Log}[c^2 x^2] - \text{Log}[1 - \frac{i}{c x}] - \text{Log}[1 + \frac{i}{c x}]) = 0$

Basis: $(\text{Log}[c^2 x^2] + \text{Log}[1 - \frac{i}{c x}] + \text{Log}[1 + \frac{i}{c x}]) \text{ArcCot}[c x] = \text{Log}[c^2 x^2] \text{ArcCot}[c x] + \frac{i}{2} \text{Log}[1 - \frac{i}{c x}]^2 - \frac{i}{2} \text{Log}[1 + \frac{i}{c x}]^2$

Rule: If $g = c^2 f$, then

$$\int \frac{\text{Log}[f + g x^2] \text{ArcCot}[c x]}{x} dx \rightarrow$$

$$\left(\text{Log}[f + g x^2] - \text{Log}[c^2 x^2] - \text{Log}\left[1 - \frac{i}{c x}\right] - \text{Log}\left[1 + \frac{i}{c x}\right] \right) \int \frac{\text{ArcCot}[c x]}{x} dx + \int \frac{\left(\text{Log}[c^2 x^2] + \text{Log}\left[1 - \frac{i}{c x}\right] + \text{Log}\left[1 + \frac{i}{c x}\right] \right) \text{ArcCot}[c x]}{x} dx \rightarrow$$

$$\left(\text{Log}[f + g x^2] - \text{Log}[c^2 x^2] - \text{Log}\left[1 - \frac{i}{c x}\right] - \text{Log}\left[1 + \frac{i}{c x}\right] \right) \int \frac{\text{ArcCot}[c x]}{x} dx + \int \frac{\text{Log}[c^2 x^2] \text{ArcCot}[c x]}{x} dx + \frac{i}{2} \int \frac{\text{Log}\left[1 - \frac{i}{c x}\right]^2}{x} dx - \frac{i}{2} \int \frac{\text{Log}\left[1 + \frac{i}{c x}\right]^2}{x} dx$$

Program code:

```
Int[Log[f_+g_*x_^2]*ArcCot[c_*x_]/x_,x_Symbol] :=
  (Log[f+g*x^2]-Log[c^2*x^2]-Log[1-I/(c*x)]-Log[1+I/(c*x)])*Int[ArcCot[c*x]/x,x] +
  Int[Log[c^2*x^2]*ArcCot[c*x]/x,x] +
  I/2*Int[Log[1-I/(c*x)]^2/x,x] -
  I/2*Int[Log[1+I/(c*x)]^2/x,x] /;
FreeQ[{c,f,g},x] && EqQ[g,c^2*f]
```

$$\text{2: } \int \frac{\text{Log}[f + g x^2] (a + b \text{ArcTan}[c x])}{x} dx$$

Derivation: Algebraic expansion

Rule:

$$\int \frac{\text{Log}[f + g x^2] (a + b \text{ArcTan}[c x])}{x} dx \rightarrow a \int \frac{\text{Log}[f + g x^2]}{x} dx + b \int \frac{\text{Log}[f + g x^2] \text{ArcTan}[c x]}{x} dx$$

Program code:

```
Int[Log[f_+g_*x_^2]*(a_+b_*ArcTan[c_*x_])/x_,x_Symbol] :=
  a*Int[Log[f+g*x^2]/x,x] + b*Int[Log[f+g*x^2]*ArcTan[c*x]/x,x] /;
FreeQ[{a,b,c,f,g},x]
```

```
Int[Log[f_.*g_.*x_^2]*(a_+b_.*ArcCot[c_.*x_])/x_,x_Symbol] :=
  a*Int[Log[f+g*x^2]/x,x] + b*Int[Log[f+g*x^2]*ArcCot[c*x]/x,x] /;
FreeQ[{a,b,c,f,g},x]
```

$$2: \int \frac{(d + e \operatorname{Log}[f + g x^2]) (a + b \operatorname{ArcTan}[c x])}{x} dx$$

Derivation: Algebraic expansion

Rule:

$$\int \frac{(d + e \operatorname{Log}[f + g x^2]) (a + b \operatorname{ArcTan}[c x])}{x} dx \rightarrow d \int \frac{a + b \operatorname{ArcTan}[c x]}{x} dx + e \int \frac{\operatorname{Log}[f + g x^2] (a + b \operatorname{ArcTan}[c x])}{x} dx$$

Program code:

```
Int[(d_+e_.*Log[f_.*g_.*x_^2])*(a_+b_.*ArcTan[c_.*x_])/x_,x_Symbol] :=
  d*Int[(a+b*ArcTan[c*x])/x,x] + e*Int[Log[f+g*x^2]*(a+b*ArcTan[c*x])/x,x] /;
FreeQ[{a,b,c,d,e,f,g},x]
```

```
Int[(d_+e_.*Log[f_.*g_.*x_^2])*(a_+b_.*ArcCot[c_.*x_])/x_,x_Symbol] :=
  d*Int[(a+b*ArcCot[c*x])/x,x] + e*Int[Log[f+g*x^2]*(a+b*ArcCot[c*x])/x,x] /;
FreeQ[{a,b,c,d,e,f,g},x]
```

2: $\int x^m (d + e \operatorname{Log}[f + g x^2]) (a + b \operatorname{ArcTan}[c x]) dx$ when $\frac{m}{2} \in \mathbb{Z}^-$

Derivation: Integration by parts

Rule: If $\frac{m}{2} \in \mathbb{Z}$, then

$$\int x^m (d + e \operatorname{Log}[f + g x^2]) (a + b \operatorname{ArcTan}[c x]) dx \rightarrow \frac{x^{m+1} (d + e \operatorname{Log}[f + g x^2]) (a + b \operatorname{ArcTan}[c x])}{m+1} - \frac{2 e g}{m+1} \int \frac{x^{m+2} (a + b \operatorname{ArcTan}[c x])}{f + g x^2} dx - \frac{b c}{m+1} \int \frac{x^{m+1} (d + e \operatorname{Log}[f + g x^2])}{1 + c^2 x^2} dx$$

Program code:

```
Int[x_^m_.*(d_+e_.*Log[f_+g_.*x_^2])*(a_+b_.*ArcTan[c_.*x_]),x_Symbol] :=
  x^(m+1)*(d+e*Log[f+g*x^2])*(a+b*ArcTan[c*x])/(m+1) -
  2*e*g/(m+1)*Int[x^(m+2)*(a+b*ArcTan[c*x])/(f+g*x^2),x] -
  b*c/(m+1)*Int[x^(m+1)*(d+e*Log[f+g*x^2])/(1+c^2*x^2),x] /;
FreeQ[{a,b,c,d,e,f,g},x] && ILtQ[m/2,0]
```

```
Int[x_^m_.*(d_+e_.*Log[f_+g_.*x_^2])*(a_+b_.*ArcCot[c_.*x_]),x_Symbol] :=
  x^(m+1)*(d+e*Log[f+g*x^2])*(a+b*ArcCot[c*x])/(m+1) -
  2*e*g/(m+1)*Int[x^(m+2)*(a+b*ArcCot[c*x])/(f+g*x^2),x] +
  b*c/(m+1)*Int[x^(m+1)*(d+e*Log[f+g*x^2])/(1+c^2*x^2),x] /;
FreeQ[{a,b,c,d,e,f,g},x] && ILtQ[m/2,0]
```

3: $\int x^m (d + e \operatorname{Log}[f + g x^2]) (a + b \operatorname{ArcTan}[c x]) dx$ when $\frac{m+1}{2} \in \mathbb{Z}^+$

Derivation: Integration by parts

Rule: If $\frac{m+1}{2} \in \mathbb{Z}^+$, let $u = \int x^m (d + e \operatorname{Log}[f + g x^2]) dx$, then

$$\int x^m (d + e \operatorname{Log}[f + g x^2]) (a + b \operatorname{ArcTan}[c x]) dx \rightarrow u (a + b \operatorname{ArcTan}[c x]) - b c \int \frac{u}{1 + c^2 x^2} dx$$

Program code:

```
Int[x^m.*(d_.+e_.*Log[f_.+g_.*x_^2])*(a_.+b_.*ArcTan[c_.*x_]),x_Symbol] :=
  With[{u=IntHide[x^m*(d+e*Log[f+g*x^2]),x]},
    Dist[a+b*ArcTan[c*x],u,x] - b*c*Int[ExpandIntegrand[u/(1+c^2*x^2),x],x] /;
  FreeQ[{a,b,c,d,e,f,g},x] && IGtQ[(m+1)/2,0]
```

```
Int[x^m.*(d_.+e_.*Log[f_.+g_.*x_^2])*(a_.+b_.*ArcCot[c_.*x_]),x_Symbol] :=
  With[{u=IntHide[x^m*(d+e*Log[f+g*x^2]),x]},
    Dist[a+b*ArcCot[c*x],u,x] + b*c*Int[ExpandIntegrand[u/(1+c^2*x^2),x],x] /;
  FreeQ[{a,b,c,d,e,f,g},x] && IGtQ[(m+1)/2,0]
```

4: $\int x^m (d + e \operatorname{Log}[f + g x^2]) (a + b \operatorname{ArcTan}[c x]) dx$ when $m \in \mathbb{Z}$

Derivation: Integration by parts

Rule: If $m \in \mathbb{Z}$, let $u = \int x^m (a + b \operatorname{ArcTan}[c x]) dx$, then

$$\int x^m (d + e \operatorname{Log}[f + g x^2]) (a + b \operatorname{ArcTan}[c x]) dx \rightarrow u (d + e \operatorname{Log}[f + g x^2]) - 2 e g \int \frac{x u}{f + g x^2} dx$$

Program code:

```
Int[x_^m_.*(d_+e_.*Log[f_+g_.*x_^2])*(a_+b_.*ArcTan[c_.*x_]),x_Symbol] :=
  With[{u=IntHide[x^m*(a+b*ArcTan[c*x]),x]},
    Dist[d+e*Log[f+g*x^2],u,x] - 2*e*g*Int[ExpandIntegrand[x*u/(f+g*x^2),x],x] /;
    FreeQ[{a,b,c,d,e,f,g},x] && IntegerQ[m] && NeQ[m,-1]
```

```
Int[x_^m_.*(d_+e_.*Log[f_+g_.*x_^2])*(a_+b_.*ArcCot[c_.*x_]),x_Symbol] :=
  With[{u=IntHide[x^m*(a+b*ArcCot[c*x]),x]},
    Dist[d+e*Log[f+g*x^2],u,x] - 2*e*g*Int[ExpandIntegrand[x*u/(f+g*x^2),x],x] /;
    FreeQ[{a,b,c,d,e,f,g},x] && IntegerQ[m] && NeQ[m,-1]
```

3: $\int x (d + e \operatorname{Log}[f + g x^2]) (a + b \operatorname{ArcTan}[c x])^2 dx$ when $g = c^2 f$

Derivation: Integration by parts

Basis: $x (d + e \operatorname{Log}[f + g x^2]) = \partial_x \left(\frac{(f+g x^2) (d+e \operatorname{Log}[f+g x^2])}{2 g} - \frac{e x^2}{2} \right)$

Rule: If $g = c^2 f$, then

$$\int x (d + e \operatorname{Log}[f + g x^2]) (a + b \operatorname{ArcTan}[c x])^2 dx \rightarrow \frac{(f + g x^2) (d + e \operatorname{Log}[f + g x^2]) (a + b \operatorname{ArcTan}[c x])^2}{2 g} - \frac{e x^2 (a + b \operatorname{ArcTan}[c x])^2}{2} -$$

$$\frac{b}{c} \int (d + e \operatorname{Log}[f + g x^2]) (a + b \operatorname{ArcTan}[c x]) dx + b c e \int \frac{x^2 (a + b \operatorname{ArcTan}[c x])}{1 + c^2 x^2} dx$$

Program code:

```
Int[x_*(d_+e_.*Log[f_+g_.*x_^2])*(a_+b_.*ArcTan[c_.*x_])^2,x_Symbol] :=
  (f+g*x^2)*(d+e*Log[f+g*x^2])*(a+b*ArcTan[c*x])^2/(2*g) -
  e*x^2*(a+b*ArcTan[c*x])^2/2 -
  b/c*Int[(d+e*Log[f+g*x^2])*(a+b*ArcTan[c*x]),x] +
  b*c*e*Int[x^2*(a+b*ArcTan[c*x])/(1+c^2*x^2),x] /;
FreeQ[{a,b,c,d,e,f,g},x] && EqQ[g,c^2*f]
```

```
Int[x_*(d_+e_.*Log[f_+g_.*x_^2])*(a_+b_.*ArcCot[c_.*x_])^2,x_Symbol] :=
  (f+g*x^2)*(d+e*Log[f+g*x^2])*(a+b*ArcCot[c*x])^2/(2*g) -
  e*x^2*(a+b*ArcCot[c*x])^2/2 +
  b/c*Int[(d+e*Log[f+g*x^2])*(a+b*ArcCot[c*x]),x] -
  b*c*e*Int[x^2*(a+b*ArcCot[c*x])/(1+c^2*x^2),x] /;
FreeQ[{a,b,c,d,e,f,g},x] && EqQ[g,c^2*f]
```

U: $\int u (a + b \operatorname{ArcTan}[c x])^p dx$

Rule:

$$\int u (a + b \operatorname{ArcTan}[c x])^p dx \rightarrow \int u (a + b \operatorname{ArcTan}[c x])^p dx$$

Program code:

```
Int[u_.*(a_.+b_.*ArcTan[c_.*x_])^p_.,x_Symbol] :=
  Unintegrable[u*(a+b*ArcTan[c*x])^p,x] /;
FreeQ[{a,b,c,p},x] && (EqQ[u,1] ||
  MatchQ[u,(d_.+e_.*x)^q_./; FreeQ[{d,e,q},x]] ||
  MatchQ[u,(f_.*x)^m_.*(d_.+e_.*x)^q_./; FreeQ[{d,e,f,m,q},x]] ||
  MatchQ[u,(d_.+e_.*x^2)^q_./; FreeQ[{d,e,q},x]] ||
  MatchQ[u,(f_.*x)^m_.*(d_.+e_.*x^2)^q_./; FreeQ[{d,e,f,m,q},x]])
```

```
Int[u_.*(a_.+b_.*ArcCot[c_.*x_])^p_.,x_Symbol] :=
  Unintegrable[u*(a+b*ArcCot[c*x])^p,x] /;
FreeQ[{a,b,c,p},x] && (EqQ[u,1] ||
  MatchQ[u,(d_.+e_.*x)^q_./; FreeQ[{d,e,q},x]] ||
  MatchQ[u,(f_.*x)^m_.*(d_.+e_.*x)^q_./; FreeQ[{d,e,f,m,q},x]] ||
  MatchQ[u,(d_.+e_.*x^2)^q_./; FreeQ[{d,e,q},x]] ||
  MatchQ[u,(f_.*x)^m_.*(d_.+e_.*x^2)^q_./; FreeQ[{d,e,f,m,q},x]])
```

Rules for integrands involving $(a + b \operatorname{ArcTan}[c x^n])^p$

$$1. \int (a + b \operatorname{ArcTan}[c x^n])^p dx$$

$$1: \int \operatorname{ArcTan}[c x^n] dx$$

Derivation: Integration by parts

$$\text{Basis: } \partial_x \operatorname{ArcTan}[c x^n] = \frac{c n x^{n-1}}{1+c^2 x^{2n}}$$

Rule:

$$\int \operatorname{ArcTan}[c x^n] dx \rightarrow x \operatorname{ArcTan}[c x^n] - c n \int \frac{x^n}{1+c^2 x^{2n}} dx$$

Program code:

```
Int[ArcTan[c_.*x_^n_],x_Symbol] :=
  x*ArcTan[c*x^n] - c*n*Int[x^n/(1+c^2*x^(2*n)),x] /;
FreeQ[{c,n},x]
```

```
Int[ArcCot[c_.*x_^n_],x_Symbol] :=
  x*ArcCot[c*x^n] + c*n*Int[x^n/(1+c^2*x^(2*n)),x] /;
FreeQ[{c,n},x]
```

2: $\int (a + b \operatorname{ArcTan}[c x^n])^p dx$ when $p \in \mathbb{Z}^+ \wedge n \in \mathbb{Z}$

Derivation: Algebraic expansion

Basis: $\operatorname{ArcTan}[z] == \frac{i \operatorname{Log}[1-i z]}{2} - \frac{i \operatorname{Log}[1+i z]}{2}$

Basis: $\operatorname{ArcCot}[z] == \frac{i \operatorname{Log}[1-i z^{-1}]}{2} - \frac{i \operatorname{Log}[1+i z^{-1}]}{2}$

Rule: If $p \in \mathbb{Z}^+ \wedge n \in \mathbb{Z}$, then

$$\int (a + b \operatorname{ArcTan}[c x^n])^p dx \rightarrow \int \operatorname{ExpandIntegrand}\left[\left(a + \frac{i b \operatorname{Log}[1-i c x^n]}{2} - \frac{i b \operatorname{Log}[1+i c x^n]}{2}\right)^p, x\right] dx$$

Program code:

```
Int[(a_.+b_.*ArcTan[c_.*x_^n_.])^p_.,x_Symbol] :=
  Int[ExpandIntegrand[(a+(I*b*Log[1-I*c*x^n])/2-(I*b*Log[1+I*c*x^n])/2)^p,x],x] /;
FreeQ[{a,b,c,n},x] && IGtQ[p,0] && IntegerQ[n]
```

```
Int[(a_.+b_.*ArcCot[c_.*x_^n_.])^p_.,x_Symbol] :=
  Int[ExpandIntegrand[(a+(I*b*Log[1-I*x^(-n)/c])/2-(I*b*Log[1+I*x^(-n)/c])/2)^p,x],x] /;
FreeQ[{a,b,c,n},x] && IGtQ[p,0] && IntegerQ[n]
```

2. $\int (d x)^m (a + b \operatorname{ArcTan}[c x^n])^p dx$

1: $\int \frac{(a + b \operatorname{ArcTan}[c x^n])^p}{x} dx$ when $p \in \mathbb{Z}^+$

Derivation: Integration by substitution

Rule: If $p \in \mathbb{Z}^+$, then

$$\int \frac{(a + b \operatorname{ArcTan}[c x^n])^p}{x} dx \rightarrow \frac{1}{n} \operatorname{Subst} \left[\int \frac{(a + b \operatorname{ArcTan}[c x])^p}{x} dx, x, x^n \right]$$

Program code:

```
Int[(a_.+b_.*ArcTan[c_.*x_^n_])^p_/x_,x_Symbol] :=
  1/n*Subst[Int[(a+b*ArcTan[c*x])^p/x,x],x,x^n] /;
FreeQ[{a,b,c,n},x] && IGtQ[p,0]
```

```
Int[(a_.+b_.*ArcCot[c_.*x_^n_])^p_/x_,x_Symbol] :=
  1/n*Subst[Int[(a+b*ArcCot[c*x])^p/x,x],x,x^n] /;
FreeQ[{a,b,c,n},x] && IGtQ[p,0]
```

2: $\int (d x)^m (a + b \operatorname{ArcTan}[c x^n]) dx$ when $m \neq -1$

Derivation: Integration by parts

Basis: $\partial_x (a + b \operatorname{ArcTan}[c x^n]) = b c n \frac{x^{n-1}}{1+c^2 x^{2n}}$

Rule: If $m \neq -1$, then

$$\int (d x)^m (a + b \operatorname{ArcTan}[c x^n]) dx \rightarrow \frac{(d x)^{m+1} (a + b \operatorname{ArcTan}[c x^n])}{d (m+1)} - \frac{b c n}{d (m+1)} \int \frac{x^{n-1} (d x)^{m+1}}{1 + c^2 x^{2n}} dx$$

Program code:

```
Int[(d_.*x_)^m_.*(a_.+b_.*ArcTan[c_.*x_^n_]),x_Symbol] :=
  (d*x)^(m+1)*(a+b*ArcTan[c*x^n])/(d*(m+1)) -
  b*c*n/(d*(m+1))*Int[x^(n-1)*(d*x)^(m+1)/(1+c^2*x^(2*n)),x] /;
FreeQ[{a,b,c,d,m,n},x] && NeQ[m,-1]
```

```

Int[(d_.**x_)^m_.*(a_.+b_.**ArcCot[c_.**x_^n_]),x_Symbol] :=
  (d*x)^(m+1)*(a+b*ArcCot[c*x^n])/(d*(m+1)) +
  b*c*n/(d*(m+1))*Int[x^(n-1)*(d*x)^(m+1)/(1+c^2*x^(2*n)),x] /;
FreeQ[{a,b,c,d,m,n},x] && NeQ[m,-1]

```

3: $\int (d x)^m (a + b \operatorname{ArcTan}[c x^n])^p dx$ when $p \in \mathbb{Z}^+ \wedge m \in \mathbb{Z} \wedge n \in \mathbb{Z}$

Derivation: Algebraic expansion

$$\text{Basis: } \operatorname{ArcTan}[z] == \frac{i \operatorname{Log}[1-i z]}{2} - \frac{i \operatorname{Log}[1+i z]}{2}$$

$$\text{Basis: } \operatorname{ArcCot}[z] == \frac{i \operatorname{Log}[1-i z^{-1}]}{2} - \frac{i \operatorname{Log}[1+i z^{-1}]}{2}$$

Rule: If $p \in \mathbb{Z}^+ \wedge m \in \mathbb{Z} \wedge n \in \mathbb{Z}$, then

$$\int (d x)^m (a + b \operatorname{ArcTan}[c x^n])^p dx \rightarrow \int \operatorname{ExpandIntegrand}\left[(d x)^m \left(a + \frac{i b \operatorname{Log}[1-i c x^n]}{2} - \frac{i b \operatorname{Log}[1+i c x^n]}{2}\right)^p, x\right] dx$$

Program code:

```

Int[(d_.**x_)^m_.*(a_.+b_.**ArcTan[c_.**x_^n_])^p_,x_Symbol] :=
  Int[ExpandIntegrand[(d*x)^m*(a+(I*b*Log[1-I*c*x^n])/2-(I*b*Log[1+I*c*x^n])/2)^p,x],x] /;
FreeQ[{a,b,c,d,m,n},x] && IGtQ[p,0] && IntegerQ[m] && IntegerQ[n]

```

```

Int[(d_.**x_)^m_.*(a_.+b_.**ArcCot[c_.**x_^n_])^p_,x_Symbol] :=
  Int[ExpandIntegrand[(d*x)^m*(a+(I*b*Log[1-I*x^(-n)/c])/2-(I*b*Log[1+I*x^(-n)/c])/2)^p,x],x] /;
FreeQ[{a,b,c,d,m,n},x] && IGtQ[p,0] && IntegerQ[m] && IntegerQ[n]

```

U: $\int u (a + b \operatorname{ArcTan}[c x^n])^p dx$

Rule:

$$\int u (a + b \operatorname{ArcTan}[c x^n])^p dx \rightarrow \int u (a + b \operatorname{ArcTan}[c x^n])^p dx$$

Program code:

```
Int[u_.*(a_.+b_.*ArcTan[c_.*x_^n_])^p_.,x_Symbol] :=
  Unintegrable[u*(a+b*ArcTan[c*x^n])^p,x] /;
FreeQ[{a,b,c,n,p},x] && (EqQ[u,1] || MatchQ[u,(d_.*x)^m_./; FreeQ[{d,m},x]])
```

```
Int[u_.*(a_.+b_.*ArcCot[c_.*x_^n_])^p_.,x_Symbol] :=
  Unintegrable[u*(a+b*ArcCot[c*x^n])^p,x] /;
FreeQ[{a,b,c,n,p},x] && (EqQ[u,1] || MatchQ[u,(d_.*x)^m_./; FreeQ[{d,m},x]])
```