

Ruminant Farm System Model

Animal Module

June 23, 2025



Contents

I. Preface: A Day in the Life on a RuFaS Farm	5
II. Module Introduction	6
A. Animal Life Cycle	7
III. Herd-level Processes and Management	8
A. Herd Initialization	8
B. Pens and Animal Grouping	12
C. Ration Formulation	15
a. Energy and Nutrition Requirements	15
b. Energy and Nutrition Supply	39
c. Automated Ration Formulation	47
d. User-defined ration formulation	49
D. Herd Size Management	50
IV. Animal-level Processes and Management	52
A. RuFaS Bodyweight and Growth	54
B. Animal Reproduction	58
C. Milk Production	70
D. Lactation Curve: Parameters and Definitions	78
E. Culling	84
V. Methane Emission and Manure Excretion	92
A. Manure Excretion	92
B. Enteric Methane Emission	104
C. Enteric Methane Mitigation	107
VI. Abbreviations	110
VII. References	112

List of Figures

1	There are 5 life stages from calf to cow and a cull stage as outlined in this figure.	53
2	This is a high level schematic to illustrate the methodology behind the heifer breeding and re-breeding in RuFaS Animal Module.	64
3	This is a high level schematic to illustrate the methodology behind the cow breeding and re-breeding in RuFaS Animal Module.	66
4	A schematic illustrating the flow of milk reduction in ration formulation logic.	77

List of Tables

1	Overview of the distinct phases by which animals are created and introduced to the herd. . . .	8
2	User inputs in the animal input file that influence the generation of the Animal Population (pre-animal-population). The inputs of this table are found in the Herd Initialization Section	9
3	User inputs in the animal input file that influence the generation of the Animal Population (pre-animal-population). The inputs of this table are found in the Herd Initialization Section. . . .	9
4	The user must define at least 3 pens and for each pen, key inputs are:	13
6	A summary of animal types	14
7	Key variables used in protein requirement calculations.	21
8	Key variables used in mineral requirement calculations.	24
9	Key variables used in DMI requirement calculations.	25
10	Key variables used in protein requirement calculations.	31
11	Key variables used in mineral requirement calculations.	34
12	Key variables used in DMI estimation calculations.	35
13	Essential and non-essential amino acids	36
14	Key variables used in calculation of AA.	38
15	Key variables used in calculation of AA.	41
16	Key variables used in calculation of AA.	43
17	Key variables used in calculation of AA.	44
18	Key inputs found in the animal input file.	47
19	Key inputs found in the feed input file.	47
20	Key inputs found in the feed input file specific to user defined ration methodology. All are found in the section user_defined_ration_percentages	49
21	Key inputs found in the feed input file specific to user defined ration methodology. All are found in the section user_defined_ration_percentages	50
22	Below is a summary of the user inputs that are available and their definitions.	54
23	Below are the key user inputs that determine the heifer reproductive breeding program and impact reproductive efficiency and age at first calving.	60
24	The following inputs determine reproductive efficiency of cows and influence culling, calving interval, and calf output.	61
25	The following inputs can be used to determine if there is a decrease in the conception rate with parity and rebreeding attempts.	62
26	Important functions for the reproduction section.	68
27	The following user inputs are used to estimate customized herd-level lactation curve parameters. See the Methodology section of this document and the Lactation Curve Scientific Documentation for more detail about how these inputs are used with relation to daily milk production.	71
28	The following inputs are used to determine the milk fat and milk protein percentages for each lactating cow	72
29	The following inputs are used to constrain how much an average cow's daily milk production will be reduced due to an inadequate ration. See the Methodology section of this document and the Ration Formulation scientific documentation for more detail about how these inputs are used with relation to daily milk production.	73
30	Wood's Lactation Curve Base Parameters – These inputs provide the baseline values and variability for Wood's lactation curve parameters as derived from Li, Rosa, et al. (2022).	79
32	Adjustment Factors – These factors adjust the base parameters based on geographical, temporal, management, and parity considerations.	80
34	Farm Specific 305-Day Milk Production — These inputs help tailor the lactation curve to the farm's production targets and herd composition.	81
35	Summary of important functions for the reproduction section.	82
36	The following inputs are used to determine the probability and timing of cow death and sales events.	86



37	Cumulative distribution function (CDF) values associated with the conditional probability that a sold animal is removed due to each group of issues and/or the conditional probability of death over time. Corresponds to the “probability” value for each cull reason in the animal input json file. The 'cull_day_count' array defines the segments of the CDF	91
38	These inputs define basic animal and diet properties used to calculate nutrient intake and manure production.	93
39	These inputs are used when modeling cows and are particularly relevant for lactating animals.	93
40	These inputs define phosphorus-related outputs relevant to lactating cows.	94
41	Expected outputs from this section.	94
42	These inputs define basic animal and diet properties used to calculate nutrient intake and manure production.	104
43	These inputs are used when modeling cows and are particularly relevant for lactating animals.	104
44	Inputs specific to the methane emissions model and selected mitigation strategies.	105
45	Expected outputs from this section.	105
46	General dietary fiber and fat concentration inputs.	108
47	Feed-based methane mitigation strategies.	108
48	Dosage of methane mitigation additives.	108
49	Expected outputs from this section.	108
50	Common abbreviations used throughout this document	110



I. Preface: A Day in the Life on a RuFaS Farm

This document provides a high-level overview of the daily operations of a ruminant farm, as modeled by RuFaS. The overview will be through the eyes of Dr. Rue Fas. One day, Dr. Rue woke up and decided the life of a computer scientist wasn't for her. So she bought a dairy farm, and this is how she operates it every day.

Feeding the Animals Second thing in the morning, after coffee, Dr. Rue checks the calendar to see if it's time to formulate a new ration for her animals.

If a new ration needs to be formulated, Dr. Rue heads over to the barn where she keeps all her forages and feed. While there, she checks how well all the forages are holding up. After considering the feeds available, the quality of each one, and how much each one costs, she decides on how productive she wants her animals to be. It is a difficult balancing act, but eventually a ration recipe is found that will keep the animals happy while keeping costs low and sustainability high until the calendar says the ration recipe needs to be reformulated.

When Dr. Rue doesn't need to formulate a new ration, she takes the last ration recipe she made and grabs all the feed and forage the animals need for the day out of the feed barn. If there isn't enough for the animals, then a new ration is formulated using the steps above.

Taking Care of the Animals After making sure all the animals' feed is taken care of, Dr. Rue moves on to taking care of all the other animal tasks. This includes milking the animals, making sure they're healthy, getting them pregnant, and numerous other tasks.

As dawn breaks on the RuFaS farm, the animals awaken after a restful night. Joe, the farm master, prepares a large whiteboard to track HerdReproductionStatistics, ensuring he has all the data he needs for the day's activities.

Daily Routine - After leaving their pens, each animal lines up in front of the barn. Julie, the diligent farm veterinarian, examines them one by one. This daily check includes updates on the following areas:

- **Nutrients:** Each animal begins its day with a quick phosphorus requirement assessment. Julie records these details and updates the animal's phosphorus-related metrics.
- **Digestive System:** With nutrient information in hand, Julie shifts focus to each animal's digestive system, estimating manure output and enteric methane production for that day.
- **Milk Production:** All milking cows are assessed to gauge daily milk yield and milk component content (fat and protein). Any necessary adjustments are noted.
- **Animal Growth:** Around midday, Julie returns to each animal to evaluate daily weight changes. She updates each animal's body weight accordingly.
- **Reproduction:** HeiferII and Cow stage animals undergo a reproduction check. Julie closely monitors their reproductive progress and updates the HerdReproductionStatistics details on Joe's whiteboard.
- **Life Stage Evaluation:** Once the daily routines are complete, Julie determines if any animal is ready to progress to the next life stage. Graduating animals are marked as "graduated" to be moved to their new pens. Animals that must leave the herd—due to health, productivity, or other issues—are labeled as "sold" or "dead" to be removed from the farm population.
- **Herd Size Maintenance:** After every animal has been evaluated, Joe counts the herd to ensure it remains at a healthy size. If it is too large, he arranges the sale of some animals. If the herd is too small, he may opt to purchase additional heiferIIIs to keep numbers at the ideal level.
- **Herd Structure Management:** Joe also manages pen assignments. Newly graduated or newborn animals are guided to suitable pens, while animals marked for removal (sold or deceased) are taken out of the herd and removed from the farm records.



Output Reporting At day's end, Joe and Julie review the updated HerdReproductionStatistics on the white-board. Joe then compiles a concise report summarizing the day's herd stats. He forwards this information to the OutputManager before wrapping up another successful day on the RuFaS farm.

Dealing with the Manure Once all the animal chores are finished, Dr. Rue moves on to handling the manure. There are a lot of different ways that manure is collected from the animals: it is flushed out of the milking parlor with water, some of the pens need to have it shoveled out along with the bedding in the pen, and some pens have grates that allow the manure to go straight into a pit beneath the barn.

There are a lot of different places where manure is processed and stored on the farm. Dr. Rue goes from place to place on the farm, moving the manure from one place to the next. The manure chores are finished after moving all of the manure into her lagoons, compost piles, and other storages. As this final manure chore is completed, she writes down all the information about how much manure is in all these storages, and some information about what that manure is made of. How much water, nitrogen, phosphorus, other solids, etc. are in each storage.

Managing the Fields Now that Dr. Rue is done making sure all the manure has been handled and stored properly, she can take care of all the field tasks. She has a lot of fields on her farm, each with its own unique schedule. Her field management routine starts with checking each field's schedule to see if any of them are due to have manure applied, and if so, how much they should have applied. After noting all the manure that is supposed to go onto her fields, these amounts are checked against the amounts of manure that are in her manure storage. Dr. Rue then applies manure to each field on the schedule that day, trying to match the amount applied to the amount on the schedule as best she can.

After making sure every manure item on each farm's schedule has been taken care of for the day, the schedule for each field is rechecked, one field at a time. In each schedule she checks to see if that field needs to be tilled, if any chemical fertilizer needs to be applied, and if any crops need to be planted or harvested. After all these activities are taken care of, Mother Nature goes to work. The sun shines and rain falls, providing the field with water and energy that grows crops and soaks the soil. After taking care of each field one by one, Dr. Rue drives back to the main farm with all of the crops that she harvested on that day.

Storing the Feeds Now that she is back at the barns, all the crops that were just harvested can be stored. Fortunately she is very organized, so all her harvested crops are separate from one another and each has a label, indicating where it will be stored inside the feed barn. As each feed is stored, it is labeled with the current date. Dr. Rue is exhausted from another long day. She goes back to the farmhouse and rests up, preparing to do the whole thing over again tomorrow. Her daily routine might seem repetitive, but each day is a crucial part of the larger, continuous process that keeps her farm running efficiently and sustainably.

II. Module Introduction

The Animal Module simulates individual animals on a daily basis throughout each stage of life from birth until removal from the herd. In addition to the dynamic Monte-Carlo models that determine the occurrence of events throughout each animal's life, the Animal Module uses process-based, dynamic models to simulate the following processes:



- Individual animal growth
- Gestation and milk production
- Estimation of nutrient requirements and feed intake
- Enteric emission and manure excretion

Animals in the RuFaS model are divided into 5 life stages and a culling stage that were established based on the different management and physiological processes that need to be simulated for each class. The 5



classes and culling stage are described in more detail later in this document, but include: calves, heifers I, heifers II, heifers III, cow, and culled.

The attributes that define the animal characteristics and management are provided by the user and are comprehensively described in inputs section of this module. When a simulation begins, the herd is populated by a semi-random draw of animals from a large, existing set of animal objects. This process and the process for regenerating the population of animals used to initialize the herd are described in detail in the [Herd Initialization Section](#). The initially selected animal objects are then grouped into pens based on their life stage and the user-provided pen information. Once the herd has been set up, a regular progression of functions is called each day to simulate and report outputs of the processes required to manage the herd and the processes that define each animal's development and production. These daily updates are grouped into two classes in the RuFaS model: herd level daily routines (HerdManager) and animal level daily routines (AnimalManager). At the start of each day, the Herd-level daily update cycles through all the animals to call their daily update functions. Then based on the changes for each animal that resulted from the animal level daily updates, the HerdManager changes the animal life-stages, moves them to new pens, and removes them from the herd. If the herd size has decreased too much, the HerdManager will purchase new Heifer III animals to maintain the adult herd size. The HerdManager then checks if it is time to formulate a new diet (at a user-specified interval), summarizes and reports the herd statistics, and gathers and reports all of the manure excreted by the animals to pass to the Manure Module.

Daily outputs from the Animal module are provided at a variety of scales and time steps that range from reports at the individual animal level (e.g. animal event histories), animal class level (e.g. average dry-matter intake for Heifer II's), at the pen level (e.g. average net energy requirement or intake for Pen 3), and at the herd level (i.e. number of animals sold each day). A complete list of potential outputs from the Animal Module is provided in this document.

A. Animal Life Cycle

The animal life cycle is driven by Monte Carlo stochastic processes that simulate the growth, production, reproduction, and culling of individual dairy cattle animals and shares aggregated outcomes with the herd management sub-module to manage herd dynamics on a daily basis.

The life cycle module represents the variation in animal performance through the use of Monte Carlo Simulation methods (Reuven and Kroese, 2017). Monte Carlo simulation relies on random draws from probability distributions rather than assigning fixed values. In this model, two main strategies are used:

- Event-based, in which a random draw from a uniform distribution ($U(0,1)$) is compared to the probability of an event occurring to simulate if that event occurs, and
- Population-based, in which a random draw from a known distribution of attribute values is assigned to an animal at instantiation. Probability distributions used for that purpose are Normal Gaussian or Empirical.

Each animal within the RuFaS herd progresses through five main life stages (calf, Heifer I, Heifer II, Heifer III, and Cow) as illustrated in the figure below.

- Calves progress to Heifer I when their age reaches the user-defined input for weaning day
- Heifer I progresses to Heifer II when their age reaches the user-defined input for breeding start day
- Heifer II animals that successfully conceive become Heifer III's when the number of days left in their pregnancy is equal to the user-defined input for prefresh days (default 21)
- When a Heifer III calves, she becomes a Cow for the rest of her life on the RuFaS farm
- Cows cycle through lactating and non-lactating phases until they are removed from the herd due to a failure to conceive or a user-provided probability for sale or death within each parity (lactation).

When a HeiferIII or Cow reaches full gestation length, they start lactating and a new calf object is created. The calf is assigned a birthweight according to a user-informed random distribution. All male calves are



recorded and removed from the farm on day 1 of their life. Female calves either remain on the farm or are sold on day 1, determined by an event-based Monte Carlo method informed by the user-defined percent of female calves that are kept.

III. Herd-level Processes and Management

A. Herd Initialization

Important Note

In RuFaS code, the term “herd initialization” is used to refer to both the process of generating an animal population file and of selecting from that population to create a specific herd at the start of a simulation. In this doc, those two processes will be differentiated by calling them “Generation” and “Selection”, respectively.

“Animal population” can refer to both groups of animals (the pool of animals generated, and the herd of animals selected). In the code, the population is sometimes referred to as “pre_animal_population” and “post_animal_population” to differentiate whether the specified group of animals exists before or after the random selection of individuals to create a herd.

Introduction Every simulation begins with a herd consisting of calves, heifers, and dry and lactating cows. This herd comes to be through a two-step process:

1. A population of animals is generated through simulation or loaded from data (pre-animal-population). The animals in that herd should reflect the attributes of animals in the herd to be modeled.
2. Animals are randomly selected with replacement from the pre-animal-population to be in the initial herd, based on group numbers and characteristics that reflect the distribution of animals within the herd being modeled. Note that the animals in the herd on day one will have attributes of the pre-animal-population and may deviate from the desired herd to be modeled.

For the rest of the simulation, animals enter the herd through birth as newborn calves or purchased replacements (heiferIII's).

Table 1: Overview of the distinct phases by which animals are created and introduced to the herd.

Before the simulation	At the initiation of simulation	During the simulation
Generate (simulate) or load an Animal Population file (pre-animal-population). The animals present at the start of the simulation, and those available in the replacement market, are drawn from this file.	Herd is established by random selection of animals from the Animal Population to match the user-specified group sizes and age/stage distribution (post-animal-population). Management attributes—including lactation-curve parameters—are assigned as each animal enters the herd.	New animals enter via (a) birth or (b) purchase of heifers from the replacement market. Purchased animals inherit two attributes from the population file: • Breed • Mature body weight

Required User Inputs

Table 2: User inputs in the animal input file that influence the generation of the Animal Population (pre-animal-population). The inputs of this table are found in the Herd Initialization Section

Variable	Definition	Default	Min	Max
Initial_animal_num	The initial number of animals that serve as a starting point for the population generation simulation	10000	0	-
Simulation_days	The number of days to simulate for the population generation simulation	5000	0	-

Table 3: User inputs in the animal input file that influence the generation of the Animal Population (pre-animal-population). The inputs of this table are found in the Herd Initialization Section.

Variable	Definition	Default	Min	Max
Calf_num	Number of Calves (head) – The initial number of pre-weaned calves	8	2	–
HeiferI_num	Number of HeiferI animals (head) – The initial number of heifers that are weaned but not yet bred	44	2	–
HeiferII_num	Number of HeiferII animals (head) – The initial number of heifers that are either eligible for breeding (based on user-inputted Breeding Start Day for heifers), or in early pregnancy	38	2	–
HeiferIII_num_springers	Number of HeiferIII animals (head) – The initial number of close-up heifers (within the user-defined close-up period, e.g., Prefresh Day)	5	2	–
Cow_num	Number of Cows (head) – The initial number of dry and lactating cows	100	6	–

Continued on next page

Table continued from previous page

Variable	Definition	Default	Min	Max
Replace_num	Replacements (head) – Number of replacement animals available in the replacement market	5000	50	–
Lactating_fraction	Fraction of cows that are lactating. Used during herd initialization	0.8356	0	1
Parity_fractions: 1, 2, 3, 4, 5	Fractions of the milking animal population that are parity 1, 2, 3, 4, and 5+. The sum must equal 1.0	0.36, 0.26, 0.18, 0.11, 0.09	0	1

**Relevant Outputs**

After the animals have been selected from the animal population to create the initial herd, a summary of the Population (pre-selection pool, aka pre-animal-population) and of the Initial herd (selected animals, aka post-animal-population) is provided to the output manager.

Class and function: `AnimalModuleReporter.report_animal_population_statistics` Output variables:

- `.population....`
- `.initial....`

... includes: breed, number of animals of each class (calf, heiferI, heiferII, heiferIII, cow), number of cows by parity and lactation status, average age of animals within each class, distribution of age of animals within each class, average body weight of animals within each class, average cow days in milk, days in preg, parity, and calving interval.

Methodology When a simulation begins, the `HerdFactory` class generates or loads the pre-animal-population, from which it selects and compiles the post-animal-population with an appropriate distribution of animal ages and stages to serve as the starting herd for simulation.

Designation of an Animal Population

Option 1: Creating a new Animal Population An animal population can be generated as a stand-alone simulation task that creates and saves a pre-animal-population file as the output based on the user Animal Module inputs. Or, the animal population can be generated as the first step of a regular simulation task which then uses that generated pre-animal-population to select from and create the post-animal-population to use for the subsequent herd simulation. See the documentation about Task Manager for more information about how to generate a new pre-animal-population.

The `_generate_animals()` method creates the pre_animal_population, with the option to save those animals to file at the end of the generation process.

Calves are created according to the number specified by the user (`initial_animal_num`), some are sold based on user inputs, and those remaining proceed through select daily updates (growth, reproduction, milk, and transition through the appropriate life stages) for the user-specified number of days (`simulation_days`). The simulation occurs similarly to the regular life cycle simulation process, but with an extended period and large number of animals. There are a few notable distinctions:

- After 3000 days of simulation (specified in `animal_constants`), a copy of each heiferIII is saved to the replacement herd. The copy is made on the day when she is ready to transition to a lactating cow (ready to give birth), so that when the copy enters the herd as a purchased heiferIII during the actual simulation, she will give birth the very next day.
- If a cow has made it to parity six without being culled, she is removed from the herd. (The maximum parity of cows kept in the generated herd is five).

After the `simulation_days` number of days, the full herd contains a mixture of calves, heifers, cows, and replacements. The core status information and life history of each animal are either saved to file or passed for random sampling to create the post-animal-population.

Option 2: Point to an existing Animal Population Alternatively, the pre-animal-population will be loaded from a specified Animal Population input file consisting of calf, heifer, cow, and replacement objects.

Herd Selection The `_random_sample_with_replacement()` method selects animals from each of the following groups: calf, heiferI, heiferII, heiferIII, replacement, lactating cows by parity for parities 1-5, and dry cows by parity for parities 1-5. Animals are sampled randomly with replacement from the pre-animal-population according to the numbers, parity fractions, and milking cow fraction specified in the animal input file. The resulting list of animals is returned as the post-animal-population and summaries of the pre- and post-animal-populations are reported to the output manager.



B. Pens and Animal Grouping

Introduction All animals in RuFaS are assigned to a pen which is used to summarize nutrient requirements for diet formulation and feeding, aggregate enteric methane and manure excretions for reporting and passage to the Manure Module. Each pen tracks and updates daily a list of animals in the pen, the stocking density, the average nutrient requirement of the animals in the pen, the ration being fed to that pen, the manure excreted, and the methane emitted. In addition, constant attributes of each pen include the type of pen it is (freestall, tiestall, open lot, compost bedded pack barn), the type of animals that can be in that pen, the number stalls, the manure management practices associated with that pen, and the distance to the milking parlor for lactating cows. Each day after the HerdManager has updated the status of each animal and determined which animals need to be bought or sold, the animals are added and removed from pens as needed.

Required User Inputs

Table 4: The user must define at least 3 pens and for each pen, key inputs are:

Variable	Definition	Default	Min	Max
id	The index of the pen			
pen_name	The name or identifier of a given pen			
animal_combination	The valid combinations of animal types that can be in a pen. CALF = Calves; GROWING = HeiferI's and HeiferII's; CLOSE_UP = HeiferIII's and Dry Cows; LAC_COWS = Lactating Cows		CALF, GROWING	CLOSE_UP, LAC_COW
vertical_dist_to_milking_parlor	The elevation gain (in meters) between the pen and the milking parlor	0.1	0	NA
horizontal_dist_to_milking_parlor	The distance (in meters) between the pen and the milking parlor	10	0	NA
number_of_stalls	The number of stalls in the pen	1000	0	NA
pen_type	The type of pen (freestall, tiestall, open lot, compost bedded pack barn)	freestall	freestall, tiestall, open lot, compost bedded pack barn	
max_stocking_density	The maximum ratio of the number of animals in the pen to the number of stalls in the pen	1.2	0.5	5
manure_management_scenario_id	The ID number of the manure management practices that are used for this pen (scenarios are defined in manure management schema)	0	0	NA



Relevant Outputs

- **number_of_animals_in_pen_[id]_[AnimalCombination]** reports the number of animals in each pen each day.
- **ration_per_animals_for_pen_[id]_[AnimalCombination]** - reports average dry matter intake of an animal in the pen at the time of the ration formulation.
- **ration_per_animals_for_pen_[id]_[AnimalCombination].[RUFAS_feed_id]** - reports the amount of each feed that is delivered per animal in the pen
- **ration_nutrient_amount_for_pen_[AnimalCombination].[nutrient]** - is reporting the amount of each of the following nutrients that is provided by the ration per animal in the pen (all reported in kilograms unless otherwise indicated):

Methodology *Animal Grouping* To translate from the AnimalTypes needed to track an animal's life stages and the combinations of animals that are allowed to be grouped into a single pen, a new variable called AnimalCombination is created that combines some AnimalTypes to facilitate sorting into pens. The AnimalCombination class is defined in enums.py and includes the following mapping from AnimalTypes to AnimalCombinations:

Table 6: A summary of animal types

Animal Combination	Animal Types
CALF	Calves
GROWING	Heifer I and Heifer II
CLOSE_UP	Heifer III and Cows (non-lactating)
LAC_COW	Cows (lactating)

Currently there are only two options for assigning AnimalCombinations to pens.

- *Option 1* - The first and most commonly used option for grouping animals is to assign one AnimalCombination to each pen such that there is at least one pen for the 4 AnimalCombinations: CALF, GROWING, CLOSE_UP, and LAC_COW.
- *Option 2* - In some cases, and especially in very small farms, a user might wish to represent a scenario where all growing animals, pregnant heifers, and non-lactating cows are housed together. In this case, there is an option to have a pen that houses both GROWING and CLOSE_UP animals such that there is at least one pen with each of the 3 AnimalCombination inputs: CALF, GROWING_AND_CLOSE_UP, and LAC_COW.

Pen Sizing Due to oscillations in the exact number of animals in each animal type, it is strongly recommended that the stocking density is at least 120% of the expected number of animals in each pen.

Class

Pen Pens are initiated from the Pen class by the HerdManager at the beginning of the simulation and, if needed, an additional pen is created to accommodate animals if the animal numbers have exceeded the max stocking density.

Noteworthy Functions

Remove_animals_by_ids This function is called when the HerdManager calls for: remove_animal_from_pen_and_id_map. It is pen specific and removes the animal from the pen and the animal ID from the animal ID map that tracks the animals in the pen.



_add_new_animals This function is called by `Pen.update_animals` which is in turn called by `HerdManager.add_animal_to_pen_and_id_map`. Thus, when `HerdManager` initiates addition of animals to each pen, this function executes the addition of those animals.

Set animal nutritional requirements

C. Ration Formulation

The diet recipes in RuFaS are either formulated through least-cost formulation (Li, Rosa, et al., 2022) or provided by the user for each of 4 animal categories (calves, growing heifers, dry and close-up cows, and lactating cows). The composition of the available feeds are provided by a feed library that was adapted from the National Academies of Sciences, Engineering, and Medicine, 2021. The amount of feed required by the farm is tracked by pen and multiple pens of each animal category can be simulated; however, at this time, the RuFaS model can only accept 1 user-defined diet per animal category even if there are multiple simulated pens for that animal category.

In practice, animal feeding on dairy farms is a constantly evolving process that responds to fluctuations in feed availability, animal requirements and responses, and management goals. Although the feed delivery algorithms will make small modifications to the diet recipe in response to animal nutrient requirements and adjust the total amount of feed delivered based on the number of animals and their requirements, the diversity and degree of fluctuation in the diets is less than what is expected on a commercial dairy. The RuFaS feed library can also be modified or expanded on by changing the compositions of the existing feeds or by adding new feeds.

Regardless of whether or not a user chooses to provide their own ration or have RuFaS build and feed a ration optimized for least-cost, RuFaS needs a set of ingredients to work from. The RuFaS feed library should be consulted for the full list of ingredients and ingredients that are in both the NASEM and NRC libraries can be used. Users interested in feeding a ration that they define, need to provide a list of ingredients and their relative % dry matter intake for each animal class (pre-weaned calves, growing heifers, close-up animals, and lactating cows).

If a user would like RuFaS to use the list of ingredients to formulate a least-cost ration, it is imperative that proper costs are inputted for each ingredient listed, but the % dry matter intake in the user defined ration percentages does not need to be included. Feed costs need to be entered in \$ / kg DM. RuFaS will then use all the ingredients available (based on costs and inventory available if the feed is homegrown) to create a ration that meets the nutritional requirements of the animal class and is cost-optimized.

Important Note

For the time being, only one set of feeds and ration formula can be provided for each animal class.

a. Energy and Nutrition Requirements

NRC: Calculating Nutrient Requirements of Dairy Cattle Below are the calculations utilized based on the energy and nutrient requirements of dairy cattle as reported by the NRC and more recently the NASEM (National Research Council, 2001; National Academies of Sciences, Engineering, and Medicine, 2021).

Calculations are included for the following requirement categories:

- Energy (Maintenance, Activity, Growth, Pregnancy, Lactation)
- Protein (Maintenance, Growth, Pregnancy, Lactation)
- Minerals (Maintenance, Growth, Pregnancy, Lactation)
- Dry Matter Intake
- Amino Acids (NASEM only)

Where appropriate, requirement calculations are differentiated by animal age and reproductive status.

Nutrient Requirements of Dairy Cattle (NRC)



Energy Requirements For details about the variables included in each calculation, please refer to the energy requirements key.

- **Maintenance** - The maintenance requirement is calculated based on metabolic body weight (body weight^{0.75}).

Lactating and Dry Cows

$$CBW = MW * 0.06275$$

[AN.NRC.1]

$$CW = (18 + (DOP - 190) * 0.665) * (\frac{CBW}{45}), if DOP > 190.0$$

[AN.NRC.2]

Otherwise

$$NE_{maint} = 0.08 * (BW - CW)^{0.75}$$

[AN.NRC.3]

Heifers -The maintenance requirement is calculated based on metabolic body weight (body weight^{0.75}), body condition score, and the previous month's temperature.

$$CBW = MW * 0.06275$$

[AN.NRC.1]

$$CW = (18 + (DOP - 190) * 0.665) * (\frac{CBW}{45}), if DOP > 190.0$$

[AN.NRC.2]

Otherwise

$$BCS9 = (BCS5 - 1) * 2 + 1$$

[AN.NRC.6]

$$NE_{maint} = (BW - CW)^{0.75} * (0.086 * (0.8 + (BCS9 - 5) * 0.05) + 0.0007 * (20 - PrevTemp))$$

[AN.NRC.7]



- **Activity** - Activity requirement is proportional to body weight and daily walking distance. A grazing system and hilly topography will cost additional energy.

Lactating and Dry Cows

$$NEa1 = 0.0012 * BW \text{ if Housing} = \text{Grazing } 0$$

[AN.NRC.8]

Otherwise

$$NEa2 = 0.006 * BW \text{ if Topography} = \text{Hilly } 0$$

[AN.NRC.9]

Otherwise

$$NEa2 = 0.Distance * 0.00045 * BW + NEa1 + NEa2$$

[AN.NRC.10]

Heifers

$$NEa1 = 0.0009 * BW + 0.0016 * BW \text{ if Housing} = \text{Grazing } 0$$

[AN.NRC.11]

Otherwise

$$NEa2 = 0.006 * BW \text{ if Topography} = \text{Hilly } 0$$

[AN.NRC.12]

Otherwise

$$NEa2 = 0.Distance * 0.00045 * BW + NEa1 + NEa2$$

[AN.NRC.10]

- **Growth**

Lactating and Dry Cows Cows will continue to grow toward their mature body weight until the end of their second lactation.

$$MSBW = 0.96 * MW$$



[AN.NRC.13]

$$SBW = 0.96 * BW$$

[AN.NRC.14]

$$EBW = 0.96 * SBW$$

[AN.NRC.15]

$$EQSBW = (SBW - CW) * \frac{478}{MSBW} SBW$$

[AN.NRC.16]

$$ADG = \frac{(0.92-0.82)*MSBW}{CI}, if Parity = 1 \frac{(1-0.92)*MSBW}{CI}, if Parity = 2.0, if Parity > 2$$

[AN.NRC.17]

$$EQEBG = 0.956 * ADG$$

[AN.NRC.18]

$$EQEBW = 0.891 * EQSBW$$

[AN.NRC.19]

$$NEg = 0.0635 * EQEBW^{0.75} * EQEBG^{1.097}$$

[AN.NRC.20]

Heifers

$$ADG = \frac{0.55*MSBW-SBW}{(Age1stBred-Age)*30.4}, before breeding \frac{0.82*MSBW-SBW}{(Age1st-Age)*30.4}, after breeding$$

[AN.NRC.21]

$$EQEBG = 0.956 * ADG$$



[AN.NRC.22]

$$EQEBW = 0.891 * EQSBW$$

[AN.NRC.23]

$$NEg = 0.0635 * EQEBW^{0.75} * EQEBG^{1.097}$$

[AN.NRC.24]

- **Pregnancy**

$$ME_{preg} = (2 * 0.00159 * DOP - 0.0352) * \frac{CBW}{45} * 0.14, \text{ if } DOP > 190.0$$

Otherwise

[AN.NRC.25]

$$NE_{preg} = ME_{preg} * 0.664$$

[AN.NRC.26]

- **Lactation**

$$Milk_{Energy} = 0.0929 * Fat_{Milk} + \frac{0.0563}{0.93} * TP_{Milk} + 0.0395 * Lactose_{Milk}$$

[AN.NRC.27]

$$NE_{lact} = Milken * Milk$$

[AN.NRC.28]

Protein Requirements Reporting of protein requirements is divided into 4 components: maintenance, growth, pregnancy, and lactation (all in metabolizable protein). For details about the variables included in each calculation, please refer to the [protein requirements key](#).

- **Total Protein**

$$MP_{req} = MP_m + MP_g + MP_{preg} + MP_{lact}$$

[AN.NRC.36]



- **Maintenance**

Lactating and Dry Cows

$$MPm = 0.3 * (BW - CW)^{0.6} + 4.1 * (BW - CW)^{0.5} + (DMI * 1000 * 0.03 - 0.5 * (\frac{MPbact}{0.8} - MPbact)) + 0.4 * 11.8 * \frac{DMI}{0.67}$$

[AN.NRC.29]

- **Growth**

Lactating and Dry Cows

$$NPg = 0, if ADG = 0, ADG * (268 - 29.4 * \frac{NEg}{ADG})$$

[AN.NRC.30]

$$EffMP_NPg = \frac{(83.4 - 0.114 * EQSBW)}{100}, if EQSBW \leq 478, 0.28908$$

[AN.NRC.31]

Otherwise

$$MPg = \frac{NPg}{EffMP_NPg}$$

[AN.NRC.32]

Heifers

$$Npg = 0, if ADG = 0, ADG * (268 - 29.4 * \frac{NEg}{ADG})$$

[AN.NRC.33]

$$EffMP_NPg = \frac{(83.4 - 0.114 * EQSBW)}{100}, if EQSBW \leq 478, 0.28908$$

[AN.NRC.34]

Otherwise

$$MPg = \frac{NPg}{EffMP_NPg}$$

[AN.NRC.35]

$$MPreq = MPm + MPg + MPpreg + MPlact$$



[AN.NRC.36]

- **Pregnancy**

$$MP_{preg} = (0.69 * DOP - 69.2) * \frac{CBW}{45 * 0.33}, \text{ if } DOP > 190.0$$

[AN.NRC.37]

- **Lactation**

$$MPlact = Milk * \frac{TP_{Milk}}{100} * \frac{1000}{0.67}$$

[AN.NRC.38]

Table 7: Key variables used in protein requirement calculations.

Variable	Description	Units
EffMP_NPg	Efficiency of converting metabolizable protein or net protein	
MPg	Metabolizable protein requirement for growth	g
MPlact	Metabolizable protein requirement for lactation	g
MPpreg	Metabolizable protein requirement for pregnancy	g
NPg	Net protein requirement for growth	g

Mineral Requirements

Important Note

Only calcium (Ca) and phosphorus (P) requirements are considered currently. For details about the variables included in each calculation, please refer to the [mineral requirements key](#).

- **Maintenance**

Lactating and Dry Cows

Calcium Equations

$$Ca_{main} = 0.031 * BW + 0.08 * \frac{BW}{100}, \text{ for lactating cows}$$

$$Ca_{main} = 0.0154 * BW + 0.08 * \frac{BW}{100}, \text{ for dry cows}$$



[AN.NRC.39]

Phosphorus Equations

$$P_{\text{main}} = 1 * DMI + 0.002 * BW, \text{ for lactating cows}$$

$$P_{\text{main}} = 0.8 * DMI + 0.002 * BW, \text{ for dry cows}$$

[AN.NRC.40]

Heifers

Calcium Equation

$$Ca_{\text{main}} = 0.0154 * BW + 0.08 * \frac{BW}{100},$$

[AN.NRC.41]

Phosphorus Equation

$$P_{\text{main}} = 0.8 * DMI + 0.002 * BW,$$

[AN.NRC.42]

• **Growth**

Calcium Equation

$$Ca_{\text{growth}} = 9.83 * MW^{0.22} * BW^{-0.22} * \frac{ADG}{0.96}$$

[AN.NRC.43]

Phosphorus Equation

$$P_{\text{growth}} = (1.2 + 4.635 * MW^{0.22} * BW^{-0.22}) * \frac{ADG}{0.96}$$

[AN.NRC.44]

• **Pregnancy**

Calcium Equation

$$Ca_{\text{preg}} = 0.02456 * \exp(0.05581 - 0.00007 * DOP * DOP - 0.02456 * \exp(0.05581 - 0.00007 * -1 * DOP - 1, \text{ if } DOP > 190.0, \text{ else}$$



[AN.NRC.45]

Phosphorus Equation

$$P_{\text{preg}} = 0.02743 * \exp(0.05527 - 0.000075 * DOP * DOP) - 0.02743 * \exp(0.05527 - 0.000075 * DOP - 1 * (DOP - 1)),$$

If: $DOP > 190.0$ else

[AN.NRC.46]

- **Lactation**

Calcium Equation

$$Ca_{\text{lact}} = 1.22 * Milk$$

[AN.NRC.47]

Phosphorus Equation

$$P_{\text{lact}} = 0.9 * Milk$$

[AN.NRC.48]

- **Total Minerals**

Calcium Equation

$$Ca_{\text{req}} = Ca_{\text{main}} + Ca_{\text{growth}} + Ca_{\text{preg}} + Ca_{\text{lact}}$$

[AN.NRC.49]

Phosphorus Equation

$$P_{\text{req}} = P_{\text{main}} + P_{\text{growth}} + P_{\text{preg}} + P_{\text{lact}}$$

[AN.NRC.50]



Table 8: Key variables used in mineral requirement calculations.

Variable	Description	Units
Ca_{main}	Calcium maintenance requirement	g
Ca_{growth}	Calcium growth requirement	g
Ca_{lact}	Calcium lactation requirement	g
Ca_{preg}	Calcium pregnancy requirement	g
Ca_{req}	Total calcium requirement	g
DOP	Days of pregnancy	d
P_{growth}	Phosphorus growth requirement	g
P_{lact}	Phosphorus lactation requirement	g
P_{main}	Phosphorus maintenance requirement	g
P_{preg}	Phosphorus pregnancy requirement	g
P_{req}	Total phosphorus requirement	g

Dry Matter Intake (DMI) Estimation

Estimations are different for lactating cows and dry cows. For details about the variables included in each calculation, please refer to the DMI requirements key.

The sum of DMI of each feed is assumed to be less than the estimated DMI:

$$\sum_i Feed_i < DMI_{est}$$

- **Lactating Cows**

$$FCM = (0.4 * Milk) + (15 * Fat_Milk * \frac{Milk}{100})$$

[AN.NRC.51]

$$DMI_{est} = (0.372 * FCM + 0.0968 * BW^{0.75}) * (1 - exp - 0.192 * WOL + 3.67)$$

[AN.NRC.52]

- **Dry Cows**



$$DM_{Iest} = \left(\frac{(1.97 - 0.75 * \exp(0.16 * (DOP - 280)))}{100} \right) * BW$$

[AN.NRC.53]

- **Heifers**

If the heifer is **less** than 365 days old:

$$DM_{Iest} = (BW^{0.75} * \frac{(0.2435 * NE_{adj} - 0.0466 * NE_{adj}^2 - 0.0869)}{NE_{adj}}) - PREG_{adj}$$

[AN.NRC.54]

If the heifer is **more** than 365 days old:

$$DM_{Iest} = (BW^{0.75} * \frac{(0.2435 * NE_{adj} - 0.0466 * NE_{adj}^2 - 0.1128)}{NE_{adj}}) - PREG_{adj}$$

[AN.NRC.55]

Table 9: Key variables used in DMI requirement calculations.

Variable	Description	Units
DM _{Iest}	Dry matter intake estimation	kilogram (kg)
DOP	Days of pregnancy	days
NE _{adj}	the adjustment for the net energy (NE) concentration of the diet	if the NE concentration ≤ 1 mcal/kg DM, then = 0.95 if the concentration > 1 mcal/kg DM, then is = to the NE concentration
PREG _{adj}	the adjustment for pregnancy status	if DOP > 210 1 + ((210 - DOP) * 0.0025)
WOL	Week of lactation	Units are the integer part of days $\frac{DIM}{7}$

NASEM: Calculating Nutrient Requirements of Dairy Cattle: Eighth Revised Edition **Energy Requirements** The maintenance requirement is calculated based on metabolic body weight (MBW, body weight^{0.75}) adjusted for the tissues associated with pregnancy. For details about the variables included in each calculation, please refer to the energy requirements key.

- **Maintenance**

Lactating and Dry Cows For Lactating cows, the body weight is adjusted both for the gravid uterine tissues and the uterine tissues for normal activity in confinement conditions, otherwise include adjustments for activity requirements under grazing conditions.



$$NE_{maint} = 0.10 * (MBW - GrUterW - UterW)^{0.75}$$

[AN.NSM.1]

$$GrUterW = (CBW * 1.825) * \exp^{-(0.0243 - (0.0000245 * DOP)) * (280 - DOP)}$$

[AN.NSM.2]

$$UterW = ((CBW * 0.2288 - 0.204) * \exp^{-0.2 * DyLact} + 0.204)$$

[AN.NSM.3]

$$CW = inputvariable,$$

Otherwise

$$CBW = MW * 0.06275$$

[AN.NSM.4]

Heifers For heifers, the maintenance energy requirement is only adjusted for the gravid uterine weight:

$$NE_{main} = 0.10 * (BW - GrUterW)^{0.75}$$

[AN.NSM.5]

- **Activity** - Activity requirement is proportional to body weight and daily walking distance. A grazing system and hilly topography will cost additional energy.

$$NE_{a1} = \frac{(Dt_PastIn)}{DtDMI_n} \text{ if } < 0.005, \text{ then } 0$$

[AN.NSM.6]

Otherwise

$$NE_{a1} = 0.0075 * BW^{0.75} * \left(\frac{(600 - 12 * DtPastSupplIn)}{600} \right) \text{ if } \geq 0.005, \text{ then } 0$$

[AN.NSM.7]

Energy requirements for walking to and from the milking parlor are estimated as follows:



$$NEa2 = 0.00035 * \frac{(DistParlor)}{1000} * TripParlor * BW$$

[AN.NSM.8]

$$NEa3 = 0.0067 * \frac{(EnvTopoParlor)}{1000} * BW$$

[AN.NSM.9]

$$NEa = NEa1 + NEa2 + NEa3$$

[AN.NSM.10]

- **Growth** In National Academies of Sciences, Engineering, and Medicine (2021), body frame gain (fat + protein) corresponds to the true growth and it is part of the calculation which is further partitioned into body reserves or condition gain (or loss), and pregnancy-associated gain (considered a pregnancy requirement).

The value shown below was calculated but was never utilized in any further calculations in the model's current version. Thus, it has been omitted from the model and remains solely as a reference to past versions of the model.

$$EBW = 0.85 * BW$$

[AN.NSM.11]

$$EBG = 0.85 * ADG$$

[AN.NSM.12]

$$FatADG = (0.067 + 0.375 * (\frac{BW}{MW})) * \frac{EBG}{ADG}$$

[AN.NSM.13]

$$ProtADG = (0.201 + 0.081 * (\frac{BW}{MW})) * \frac{EBG}{ADG}$$

[AN.NSM.14]

$$REFADG = 9.4 * FatADG + 5.55 * ProtADG$$

[AN.NSM.15]



$$MEFrameADG = \frac{REFADG}{0.4}$$

[AN.NSM.16]

$$NEFrameADG = \frac{REFADG}{0.61}$$

[AN.NSM.17]

- **Pregnancy** - Daily rates of wet tissue deposition (kg/d) are derived from equations [AN.NSM.2] and [AN.NSM.3] as defined above.

During Gestation

$$GrUterWGain = (0.0243 - (0.0000245 * DayGest)) * GrUterW$$

[AN.NSM.18]

During Involution

$$MEpreg = (2 * 0.00159 * DOP - 0.0352) * \frac{CBW}{45} * 0.14, \text{ if } DOP > 190.0$$

[AN.NSM.19]

Otherwise

$$NEpreg = MEpreg * 0.64$$

[AN.NSM.20]

- **Lactation**

$$MilkEn = (0.0929 * Fat_Milk + 0.0547 * Prot_Milk + 0.0395 * Lact_Milk)$$

[AN.NSM.21]

If true protein, (TP_Milk) is measured, its energy content is adjusted to account for the energy of Non-Protein-Nitrogen (NPN), then use the following equation.

$$MilkEn = (0.0929 * Fat_Milk + 0.0585 * TP_Milk + 0.0395 * Lact_Milk)$$

[AN.NSM.22]



Protein Requirements are divided into 4 components: maintenance, growth, pregnancy, and lactation (all in MP). The last is defined as the sum of rumen undegraded protein (RUP + microbial protein (MICP)). Requirements for EEA are also included in National Academies of Sciences, Engineering, and Medicine (2021) but have not yet been implemented.

Each physiological function is quantified in terms of TP (true protein), instead of CP (crude protein).

To convert both net protein and net amino acid values to a metabolizable basis, net values have to be divided into the calculated efficiencies for each physiological function. Efficiencies can be either fixed or combined for lactating cows. For simplicity, requirements were included on a fixed basis. For details about the variables included in each calculation, please refer to the [protein requirements key](#).

- **Maintenance** - The protein requirements for non-productive functions (the quantification of protein secretion and accretion), include: scurf + endogenous urinary loss + metabolic fecal protein.

1. Scurf Protein

$$NP_{scurf} = 0.20 * BW^{0.60} * 0.85$$

[AN.NSM.23]

$$NetAA_{scurf} = NP_{scurf} * \frac{AACorrScurf}{100}$$

[AN.NSM.24]

2. Endogenous Protein

$$NP_{EndUrin} = 53 * 6.25 * BW * 0.001$$

[AN.NSM.25]

$$NetAA_{EndUrin} = 0.010 * 6.25 * BW * \frac{AACorrWhEmpBody}{100}$$

[AN.NSM.26]

- #### 3. Metabolic Fecal Protein
- The daily loss of CP as metabolic fecal protein (MFP) is estimated using the following equation.

$$CP_{MFP} = (11.62 + 0.134 * NDF) * DMI$$

[AN.NSM.27]

$$NP_{MFP} = CP_{MFP} * 0.73$$

[AN.NSM.28]



$$NPAAMFP = NPMFP * \frac{AACorrMFP}{100}$$

[AN.NSM.29]

- **Growth** There are two important concepts: frame growth and BW changes related to the mobilization of body reserves.

The following are the protein requirements associated with growth associated with increased frame size:

$$NPGrowth_{Frame} = FrameWG * 0.11 * 0.86$$

[AN.NSM.30]

If the change in BW is not frame growth but rather a change in body reserves, then the following equation is used because the protein content is assumed to be 8.0 percent protein.

$$NPGrowth_{Tissue} = TissueWG * 0.08 * 0.86$$

[AN.NSM.31]

The following equation is used to translate the protein requirement into the requirement for AA:

$$NetAAGrowth = NPGrowth * \frac{AACorrWEBW}{100}$$

[AN.NSM.32]

- **Pregnancy**

$$NPGest = GainGrUter * 125$$

[AN.NSM.33]

$$NetAAGest = NPGest * \frac{AACorrWEBW}{100}$$

[AN.NSM.34]

- **Lactation**

$$NetAAMilk = NPMilk * \frac{AacalcMilk}{100}$$



[AN.NSM.35]

- **Total Requirements** -Total MP requirements are expressed in grams per day. This corresponds to the sum of all maintenance and other physiological functions.

Lactating Cows

$$RecomMPSupply1 = \left[\frac{(NetAAScurf + NetAAMFP + NetAAMilk + NetAAGrowth)}{TargetEffMP} \right] + \frac{NPGest}{0.33} + NPEndUrin$$

[AN.NSM.36]

$$RecomMPAASupply1 = \frac{NPScurf + NPMFP + NPMilk + NPGrowth}{TargetEffAA} + \frac{NetAAGest}{0.33} + NetAAEndUrin$$

[AN.NSM.37]

Late Gestation Heifers and Cows

$$RecomMPSupply2 = \frac{(NPScurf + NPMFP)}{TargetEffMP} + \frac{NPGest}{0.33} + \frac{NPGrowth}{0.4} + NPEndUrin$$

[AN.NSM.38]

$$RecomMPAASupply2 = \frac{(NetAAScurf + NetAAMFP)}{NetAAMFP} + \frac{NetAAGest}{0.33} + \frac{NetAAGrowth}{0.4} + NPEndUrin$$

[AN.NSM.39]

Table 10: Key variables used in protein requirement calculations.

Variable	Description	Units
125 (coefficient)	Daily gain in mass of the gravid uterus	estimated constant 125 grams (g) of protein per kilogram (kg) of wet weight.
AAcalcMilk	Estimated AA composition of the true protein (TP) fraction involved in the estimation of AA Supply and recommendations	g of AA per 100 g of TP.
AACorrMFP	AA corrected for MFP	Fixed Value*
AACorrScurf	Each AA excreted as scurf	Fixed Value*

Continued on next page



Variable	Description	Units
AACorrWEBW	Each AA excreted as assuming whole empty body weight	Fixed Value*
AACorrWEBW	Each AA excreted assuming whole empty body composition	Fixed Value*
BW	body weight	kg
CPMFP	Crude protein (CP) in MFP	grams per day (g/d)
DMI	Dry matter intake	kg/d
FrameWG	Daily increase in body weight due to increased frame size	kg/d
GainGrUter	Daily gain in mass of the gravid uterus	kg/d
NetAA-Milk	Net AA yield in milk	g/d
NetAAGest	AA composition of protein accretion associated with pregnancy	g/d
NetAAGrowth	Net AA requirement for body frame weight gain	g/d
NetAAMFP	Net AA requirement for metabolic fecal protein	g/d
NetAAEndUrin	Net AA requirement for endogenous urinary excretion	g/d
NetAAscurf	Net AA demand for scurf excretion	g/d
NDF	Neutral detergent fiber in the diet	% DM
NPGest	Net protein requirement for pregnancy	g/d
NPGrowth	Net protein requirement for body frame weight gain	g/d
NPMFP	Net protein requirement for metabolic fecal protein i	g/d; assuming 73 % of TP in MFP
NPMilk	Milk's TP yield	g/d
NPscurf	Net protein requirement for scurf	g/d
PEndUrin	Net protein requirement for endogenous urinary excretion	g/d
TargetEffMP and TargetEffAA	Proposed target efficiencies of converting metabolizable protein (MP) and essential amino acids (EAA) to export proteins and body gain	Fixed Value*

Note: AA = Amino Acids; MFP= Metabolic Fecal Protein; *Fixed values are establish by National Academies of Sciences, Engineering, and Medicine (2021)

Mineral Requirements Only calcium and phosphorus requirements are included in the code at present. For details about the variables included in each calculation, please refer to the [mineral requirements key](#).

– Maintenance



Calcium Equations

$$Ca_{\text{main}} = 0.90 * DMI$$

[AN.NSM.40]

Phosphorus Equations

$$P_{\text{main}} = 1.0 * DMI + 0.0006 * BW$$

[AN.NSM.41]

– **Growth**

Calcium Equations

$$Ca_{\text{growth}} = 9.83 * MW^{-0.22} * BW^{-0.22} * ADG$$

[AN.NSM.42]

Phosphorus Equations

$$P_{\text{growth}} = (1.2 + 4.635 * MW^{0.22} * BW^{-0.22} + ADG)$$

[AN.NSM.43]

– **Pregnancy**

Calcium Equations

$$Ca_{\text{preg}} = 0.02456 * \exp((0.05581 - 0.00007 * DOP) * DOP) - 0.02456 * \exp((0.05581 - 0.00007 * (DOP - 1)) * (DOP - 1)) * (\frac{BW}{715}), \text{ if } DOP > 190; 0$$

[AN.NSM.44]

Phosphorus Equations

$$P_{\text{preg}} = 0.02743 * \exp((0.05527 - 0.000075 * DOP) * DOP) - 0.02743 * \exp((0.05527 - 0.000075 * (DOP - 1)) * (DOP - 1)) * (\frac{BW}{715}), \text{ if } DOP > 190; 0$$

[AN.NSM.45]

The denominator, 715 kg, is the average cow weight in the study by House and Bell (1993); therefore, the pregnancy (gestation) requirement is scaled to that BW. For details see Page 107 in the Minerals chapter of National Academies of Sciences, Engineering, and Medicine (2021). IF $DOP < 190$; then it is assumed that Ca_{Preg} is equal to zero.

**– Lactation**

Calcium Equations

$$Ca_{lact} = (0.295 + 0.239 * MilkTP) * Milk$$

[AN.NSM.46]

Phosphorus Equations

$$P_{lact} = Milk * (0.496 + 0.13 * TPMilk); Milk * 0.90$$

[AN.NSM.47]

– Total

Calcium Equations

$$Ca_{req} = Ca_{main} + Ca_{growth} + Ca_{preg} + Ca_{lact}$$

[AN.NSM.48]

Phosphorus Equations

$$P_{lact} = P_{main} + P_{growth} + P_{preg} + P_{lact}$$

[AN.NSM.49]

Table 11: Key variables used in mineral requirement calculations.

Variable	Description	Units
ADG	average daily BW gain	grams/day (g/d)
ADG	average daily gain	g/d
BW	Body weight	kilograms (kg)
BW	Body weight	kg
Ca_Req	Total calcium requirement	g/d
Ca_growth	Calcium requirement for growth	g/d
Ca_lact	Calcium requirement for lactation	g/d
Ca_main	Calcium requirement for maintenance	g/d
Ca_preg	Calcium pregnancy requirement	g/d
DMI	dry matter intake	kg/d
DOP	days of pregnancy	d

Continued on next page



Variable	Description	Units
Milk	Milk yield	kg/d
MilkTP	Milk true protein	%
MW	mature body weight	kg
P_lact	Phosphorus requirement for lactation	g/d
P_Req	Total phosphorus requirement	g/d
P_growth	Phosphorus requirement for growth	g/d
P_main	Phosphorus requirement for maintenance	g/d
P_preg	Phosphorus requirement for pregnancy	g

Dry Matter Intake Estimation Dry matter intake estimation is different for lactating cows and dry cows. The sum of the dry matter intake of each feed is assumed to be less than the dry matter intake estimation. For details about the variables included in each calculation, please refer to the [DMI estimation key](#).

$$\sum_i Feed_i < DM_{Est}$$

Lactation Cows

$$DM_{Est} = ((3.7 + Parity * 5.7) + 0.305 * MilkE + 0.022 * BW + (-0.689 - 1.87 * Parity) * BCS) * (1 - (0.212 + Parity * 0.136 * \exp(-0.053 * DIM)))$$

[AN.NSM.50]

Dry Cows and Growing Heifers

$$DM_{Est} = (0.0226 * MatBW * (1 - \exp(*1.47 * (\frac{BW}{MatBW})))) - (0.082 * (NDF - (23.1 + 56 * (\frac{BW}{MatBW}) - 30.6 * (\frac{BW}{MatBW})^2)))$$

[AN.NSM.51]

Table 12: Key variables used in DMI estimation calculations.

Variable	Description	Units
BCS	Body condition score	Scale of 1 to 5
BW	Body weight	kilograms (kg)

Continued on next page



Variable	Description	Units
DIM	Days in milk	days (d)
DMlest	Dry matter intake estimation	kg/d
MatBW	Mature body weight	kg
MilkE	Milk energy	Mcal/d
NDF	Neutral detergent fiber in diet	%
Parity	adjustment factor	primiparous (0) or multiparous (1)

NASEM: Calculating Amino Acid Requirements The methods for estimating the amino acids requirements for lactating cows based on the National Academies of Sciences, Engineering, and Medicine (2021), into RuFaS are described below.

- Amino acids are divided into essential (EAA) and non-essential (NEAA).
- This section will focus on calculating the requirements for the essential amino acids. The EAA and NEAA are listed in the next table.
- Find the equations in RuFaS in amino_acid.py

Table 13: Essential and non-essential amino acids

Essential (EAA)	Non-essential (NEAA)
Arginine	Alanine
Histidine	Aspartic acid
Isoleucine	Asparagine
Leucine	Cysteine
Lysine	Glutamic acid
Methionine	Glutamine
Phenylalanine	Glycine
Theonine	Proline
Tryptophan	Serine
Valine	Tyrosine

EAA Calculations - The AA requirements for lactating cows are categorized into maintenance, growth, pregnancy, and lactation. Notice that the first variable from each equation comes from the outputs of the metabolizable protein equations that are already in the RuFaS model. The second variable corresponds to the composition of AA used or secreted for each metabolic state (maintenance, growth, pregnancy, and lactation) of the cow, we can find those values in Table 6-2 (page 79) of National Academies of Sciences, Engineering, and Medicine (2021). The subscript "i" in the equations represents each specific amino acid being calculated; for example, "TotalAArequirements_i" refers to the total requirements for individual amino acids such as methionine, lysine, arginine, and others.

- **Maintenance**

1. Scurf Protein

$$NetAAscurf_i = NP_{scurf} * \frac{AACorrScurf_i}{100}$$

2. Endogenous Urinary Protein

$$NetAAEndUrine_i = 0.010 * 6.25 * BW * \frac{AACorrWholeEmptyBody_i}{100}$$

3. Metabolic Fecal Protein

$$NetAAMFP_i = NPMFP * \frac{AACorrMFP_i}{100}$$

• **Growth**

$$NetAAGrowth_i = NPGrowth * \frac{AACorrWholeEmptyBody_i}{100}$$

• **Pregnancy**

$$NetAAGest_i = NPGest * \frac{AACorrWholeEmptyBody_i}{100}$$

• **Lactation**

$$NetAAMilk_i = NPMilk * \frac{AAcalcMilk_i}{100}$$

• **Total**

Here are the equations for calculating the total AA_i requirements for lactating cows, late-gestation cows, and heifers.

The variables needed to calculate these equations come from the outputs generated in the AA requirements equations for scurf, endogenous urine, metabolic fecal protein, growth, gestation, and milk production (see formulas above).

The “ $TargetEfficiencyAA_i$ ” variable, refers to the target efficiency required to convert the EAA to build proteins and support body gain. It is a fixed value, and it can be found in Table 6-4 (page 88) of National Academies of Sciences, Engineering, and Medicine (2021).

1. Lactating Cows

$$TotalAArequirements_i = \left(\frac{NetAAscurf_i + NetAAMFP_i + NetAAGrowth_i + NetAAMilk_i}{TargetEfficiencyAA_i} \right) + \left(\frac{NetAAGest_i}{TargetEfficiencyGest} \right) + NetAAEndUrine_i$$

2. Late Gestation Cows and Heifers

$$TotalAArequirements_i = \left(\frac{NetAAscurf_i + NetAAMFP_i}{TargetEfficiencyAA_i} \right) + \left(\frac{NetAAGrowth_i}{TargetEfficiencyGrowth} \right) + \left(\frac{NetAAGest_i}{TargetEfficiencyGest} \right) + NetAAEndUrine_i$$

**Important Note**

If there are any questions regarding calculations regarding calculations of maintenance AA requirements for Lactating Cows and/or Total AA requirement of Lactating or Late Gestation Cows and Heifers, please refer to Tables 6-2 and 6-4 in National Academies of Sciences, Engineering, and Medicine, [2021](#).

Table 14: Key variables used in calculation of AA.

Variable	Description	Units
NetAAscurf	Net demand for AAi in scurf excretion	g/d
NPscurf	Net total protein requirement for scurf	g/d
AACorrScurfi ¹	For each amino acid, a fixed value for the proportion of the scurf in each AA.	-
NetAAEndUrinei	Net AAi requirement for endogenous urinary excretion	g/d
BW: Body weight	kg	
AACorrWholeEmptyBodyi ¹	For each amino acid, a fixed value for each AA within the tissues of the animal's whole empty body	g/100g TP
NetAAMFPI	Net AAi requirement for metabolic fecal protein	(MFP) g/d.
NPMFP	Net protein requirement for metabolic fecal protein	g/d. <i>assumes 73% of TP in MFP.</i>
AACorrMFPi ¹	Amino acids corrected for MFP. For each amino acid, a fixed value for each AA is excreted as metabolic fecal protein	g/100g TP
NetAAGrowthi	Net AAi requirement for body frame weight gain	g/d.
NPGrowth: Net protein requirement for body frame weight gain	g/d.	
AACorrWholeEmptyBodyi ¹	For each amino acid, a fixed value for each AA within the tissues of the animal's whole empty body	g/100g TP
NetAAGesti	Amino acids composition of protein accretion associated with gestation	g/d.
NPGest	Net protein requirement for gestation	g/d.
AACorrWholeEmptyBodyi ¹	For each amino acid, a fixed value for each AA within the tissues of the animal's whole empty body	g/100g TP
NetAAMilki	Net amino acids yield in milk	g/d.

Continued on next page



Variable	Description	Units
NPMilk	Net protein in milk. This refers to milk TP yield	g/d.
AAcalcMilki ¹	Estimated AA composition of the true protein (TP) fraction involved in the estimation of AA Supply and recommendations. Fixed values for each individual amino acid in milk	-
TotalAArequirementsi	Total AA requirement	g/d.
NetAAscurfi	Net demand for AAi in scurf excretion	g/d
NetAAMFPi	Net AAi requirement for metabolic fecal protein	g/d
NetAAGrowthi	Net AAi requirement for body frame weight gain	g/d
NetAAMilki	Net amino acids yield in milk	g/d
NetAAGesti	Amino acids composition of protein accretion associated with pregnancy	g/d
NetAAEndUrinei	Net AAi requirement for endogenous urinary excretion	g/d
TargetEfficiencyGest	Proposed target efficiency of using MP for gestation.	Value = 0.33
TargetEfficiencyAAi ¹	Proposed target efficiencies of essential amino acids (EAA) to export proteins and body gain	g/100g TP
TotalAArequirementsi	Total AA requirement	g/d
NetAAscurfi	Net demand for AAi in scurf excretion	g/d
NetAAMFPi	Net AAi requirement for metabolic fecal protein	g/d
NetAAGrowthi	Net AAi requirement for body frame weight gain	g/d
NetAAGesti	Amino acids composition of protein accretion associated with pregnancy	g/d
NetAAEndUrinei	Net AAi requirement for endogenous urinary excretion	g/d
TargetEfficiencyGest	Proposed target efficiency of using MP for gestation.	Value = 0.33
TargetEfficiencyGrowth	Proposed target efficiency of using MP for growth.	Value = 0.40
TargetEfficiencyAAi ¹	Proposed target efficiencies of essential amino acids (EAA) to export proteins and body gain	-

¹ National Academies of Sciences, Engineering, and Medicine, [2021](#)

b. Energy and Nutrition Supply

General Nutrient Supply



Energy Supply Energy supply from the feed includes NEm, NEI, and NEg. NEm provides energy for maintenance and activity requirements, NEI provides energy for lactation and pregnancy, and NEg provides energy for growth. Feed actual energy contents are corrected from standard values provided by National Research Council (2001). Minerals do not provide any energy.

- **Total Digestible Nutrients**

$$TDN_{conc} = \frac{TotalTDN}{DMI} * 100$$

[AN.SUP.1]

$$DMI_{to\,maint} = 1, if TotalTDN < 0.035 * BW^{0.75}, \frac{TotalTDN}{0.035 * BW^{0.75}}$$

[AN.SUP.2]

Otherwise:

$$Discount = 1, if TDN_{conc} < 60 \frac{(TDN_{conc} - (0.18 * TDN_{conc} - 10.3) * (DMI_{to\,maint} - 1))}{TDN_{conc}}$$

[AN.SUP.3]

$$TDN_{act_i} = TDN_i * Discount$$

[AN.SUP.4]

- **Digestible Energy**

$$DE_{act_i} = DE_i * Discount$$

[AN.SUP.4]

- **Metabolizable Energy**

$$ME_{act_i} = 1.01 * DE_{act_i} - 0.45 + 0.0046 * E_{Ei} - 3, if feed type is not fat and fat content is \geq 3\%$$

$$ME_{act_i} = 1.01 * DE_{act_i} - 0.45, if feed type is not fat and fat content is \leq 3\%$$



$$ME_{acti} = 1.01 * DE_i, \text{ if feed type is fat}$$

[AN.SUP.5]

• **Net Energy**

1. Maintenance

$$NE_{macti} = 1.37 * ME_{acti} - 0.138 * ME_{acti}^2 + 0.0105 * ME_{acti}^3 - 1.12, \text{ if feed type is not fat}$$

$$NE_{macti} = ME_{acti} * 0.8, \text{ if feed type is fat}$$

[AN.SUP.7]

2. Lactation

$$NE_{lacti} = 0.703 * ME_{acti} - 0.19 + 0.097 * ME_{acti} + 0.1997 * EE_i - 3, \text{ if feed type is not fat and fat content is } \geq 3\%$$

$$NE_{lacti} = 0.703 * ME_{acti} - 0.19, \text{ if feed type is not fat and fat content is } \leq 3\%$$

$$NE_{lacti} = 0 DE_{acti} * 0.8, \text{ if feed type is fat}$$

[AN.SUP.6]

3. Activity

$$NE_{gacti} = 1.42 * ME_{acti} - 0.174 * ME_{acti}^2 + 0.0122 * ME_{acti}^3 - 1.65, \text{ if feed type is not fat}$$

$$NE_{gacti} = ME_{acti} * 0.55, \text{ if feed type is fat}$$

[AN.SUP.8]

Table 15: Key variables used in calculation of AA.

Variable	Description	Units
NE _{gacti}	Actual net energy for growth of feed i	Mcal/kg

Continued on next page



Variable	Description	Units
NEmacti	Actual net energy for maintenance of feed i	Mcal/kg
NElacti	Actual net energy for lactation of feed i	Mcal/kg
MEacti	Actual metabolizable energy of feed i	Mcal/kg
EEi	Feed fat (ether extract) content	%
DEacti	Actual digestible energy of feed i	Mcal/kg
DEi	Standard DE feed i,	Mcal/kg
TDNacti	Actual TDN content of feed i	%
TDNi	Standard TDN content of feed i	%
Discount	TDN digestibility decrease caused by DMI and TDNconc	
DMI_to_maint	The amount of intake needed to meet the maintenance requirement	dimensionless
TDNconc	TDN concentration	%
TotalTDN ¹	Dietary TDN content	

¹ National Research Council, 2001

Protein Supply - Protein supply from the feed includes 3 parts: digestible RUP, digestible MCP, which is produced through RDP, and endogenous protein supply from sloughed cells and enzyme secretion into the gut lumen. MCP production requires a nitrogen source and an energy source, so MCP is either nitrogen-limited (when energy is sufficient) or energy-limited (when nitrogen is sufficient). Minerals do not provide any protein.

$$MP_{\text{supply}} = MP_{\text{bact}} + RUP_{\text{diet}} + EndP$$

[AN.SUP.14]

• Microbial Protein Supply

$$Kpi = 2.904 + 1.375 * \frac{DMI}{BW} * 100 - 0.02 * PercentConc, \text{ if feed is concentrate}$$

$$Kpi = 3.362 + 0.479 * \frac{DMI}{BW} * 100 - 0.17 * NDFi - 0.007 * PercentConc, \text{ if feed is forage}$$

$$Kpi = 3.054 + 0.614 * \frac{DMI}{BW} * 100, \text{ if feed is wet forage}$$

[AN.SUP.9]

$$RDPi = \frac{Kdi}{Kdi + Kpi} * \frac{NBi}{100} * Cpi + \frac{NAi}{100} * Cpi, \text{ if } Kpi + Kdi > 0$$



$$RDP_i = 0, \text{ if } K_{pi} + K_{di} \leq 0$$

[AN.SUP.10]

$$MP_{bact} = 0.64 * \min(1000 * 0.13 * TDN_{act_{diet}}, 1000 * 0.85 * RDP_{diet})$$

[AN.SUP.11]

• **Rumen Undegradable Protein Supply**

$$RUP_i = CP_i - RDP_i$$

[AN.SUP.12]

$$RUP_i = \sum_i Feed_i * RUP_i - dRUP_i$$

[AN.SUP.13]

• **Endogenous Protein Supply**

$$EndP = 0.4 * 11.8DMI$$

Table 16: Key variables used in calculation of AA.

Variable	Description	Units
EndP	the endogenous protein supply in	g/d
RUP _{diet}	Dietary digestible RUP,	g
feed _i	Dry matter intake of feed i, g	
dRUP _i	RUP digestibility of feed i,	% of RUP
RUP _i	Rumen undegradable protein of feed i,	% of DM
MP _{bact}	Metabolizable bacterial protein production,	g
TDN _{act_{diet}}	Dietary actual TDN,	kg
RDP _{diet}	Dietary RDP,	kg
RDP _i	Rumen degradable protein of feed i,	% of DM

Continued on next page



Variable	Description	Units
Kdi	Protein degradation rate of feed i,	%/h
NAi	Fraction A of protein of feed i,	% of CP
NBi	Fraction B of protein of feed i,	% of CP
Kpi	Protein passage rate of feed i,	%/h
PercentConc	Dietary concentrate percentage,	% of DM
NDFi	Neutral detergent fiber content of feed i,	%
MPsupply	Total metabolizable protein supply,	g/d
MPbact	Metabolizable bacterial protein production,	g/d
RUPdiet	Dietary digestible RUP,	g/d

Mineral Supply We will use calcium as an example, but phosphorus calculations will be similar.

$$Ca_{supply} = \sum_i Feed_i * Ca_i * \frac{dCa_i}{100}$$

[AN.SUP.15]

Table 17: Key variables used in calculation of AA.

Variable	Description
dCai	Ca digestibility of feed i
dCa	30% for forage, 60% for concentrate, 95% for mineral
dP	(P digestibility) = 64% for forage, 70% for concentrate, 80% for mineral

AA Nutrient Supply This section is divided into several subsections including Rumen Degraded and Undegraded Protein, Dry Matter Intake, Dietary Nutrient Intake, Dietary Nutrient Concentration, Digested Nutrients, Digested Rumen Undegradable Protein (RUP), Microbial Nitrogen Growth, Ruminal Nitrogen Outflow, and lastly Amino Acid Supply.

Energy Supply - Energy supply from the feed includes NEm, NEI, and NEg. NEm provides energy for maintenance and activity requirements, NEI provides energy for lactation and pregnancy, and NEg provides energy for growth. Feed actual energy contents are corrected from standard values provided by National Research Council (2001). Minerals do not provide any energy.

Total Digestible Nutrients

$$TDNconc = \frac{TotalTDN}{DMI} * 100$$

[AN.SUP.16]



$$DMI_{to_maint} = 1, \text{ If TotalTDN} < 0.035 * BW^{0.75}, \frac{TotalTDN}{0.035 * BW^{0.75}},$$

[AN.SUP.17]

Otherwise

$$Discount = 1, \text{ IF } TDN_{conc} < 60 \frac{(TDN_{conc} - (0.18 * TDN_{conc} - 10.3 * (DMI_{to_maint} - 1)))}{TDN_{conc}}$$

[AN.SUP.18]

Otherwise

$$TDN_{act_i} = TDN_i * Discount$$

[AN.SUP.19]

Digestible Energy

$$DE_{act_i} = DE_i * Discount$$

[AN.SUP.20]

Metabolizable Energy

$$\begin{aligned} ME_{act_i} &= 1.01 * DE_{act_i} - 0.45 + 0.0046 * EE_i - 3, \text{ if feed type is not fat and fat content is } \geq 3\% \\ ME_{act_i} &= 1.01 * DE_{act_i} - 0.45, \text{ if feed type is not fat and fat content is } \leq 3\% \\ ME_{act_i} &= DE_{act_i}, \text{ if feed type is fat} \end{aligned}$$

[AN.SUP.21]

Net Energy**Maintenance**

$$\begin{aligned} NE_{mact_i} &= 1.37 * ME_{act_i} - 0.138 * ME_{act_i}^2 + 0.0105 * ME_{act_i}^3 - 1.12, \text{ if feed type is not fat} \\ NE_{mact_i} &= ME_{act_i} * 0.8, \text{ if feed type is fat} \end{aligned}$$

[AN.SUP.22]

Lactation

$$\begin{aligned} NE_{lact_i} &= \\ 0.703 * ME_{act_i} - 0.19 + 0.097 * ME_{act_i} + 0.1997 * EE_i - 3, &\text{ if feed type is not fat and fat content is } \geq 3\% \\ NE_{lact_i} &= 0.703 * ME_{act_i} - 0.19, \text{ if feed type is not fat and fat content is } \leq 3\% \\ NE_{lact_i} &= DE_{act_i} * 0.8, \text{ if feed type is fat} \end{aligned}$$

[AN.SUP.23]

Growth



$$NE_{gacti} = 1.42 * ME_{acti} - 0.174 * ME_{acti}^2 + 0.0122 * ME_{acti}^3 - 1.65, \text{ if feed type is not fat}$$

$$NE_{gacti} = ME_{acti} * 0.55, \text{ if feed type is fat}$$

[AN.SUP.24]

Protein Supply - Protein supply from the feed includes 3 parts: digestible RUP, digestible MCP, which is produced through RDP, and endogenous protein supply from sloughed cells and enzyme secretion into the gut lumen. MCP production requires a nitrogen source and an energy source, so MCP is either nitrogen-limited (when energy is sufficient) or energy-limited (when nitrogen is sufficient). Minerals do not provide any protein.

$$MP_{supply} = MP_{bact} + RUP_{diet} + EndP$$

[AN.SUP.25]

Microbial Protein Supply

$$Kpi = 2.904 + 1.375 * \frac{DMI}{BW} * 100 - 0.02 * PercentConc, \text{ if feeding concentrate}$$

$$Kpi = 3.362 + 0.479 * \frac{DMI}{BW} * 100 - 0.017 * NDFi - 0.007 * PercentConc, \text{ if feeding forage}$$

$$Kpi = 3.054 + 0.614 * \frac{DMI}{BW} * 100, \text{ if feeding wet forage}$$

[AN.SUP.26]

$$RDPi = \frac{Kdi}{Kdi + Kpi} * (\frac{NBi}{100}) * Cpi + (\frac{NAi}{100}) * Cpi, \text{ if } Kpi + Kdi > 0, RDPi = 0, \text{ if } Kpi + Kdi \leq 0,$$

[AN.SUP.27]

$$MP_{bact} = 0.64 * \min(1000 * 0.13 * TDN_{act_{diet}}, 1000 * 0.85 * RDP_{diet})$$

[AN.SUP.28]

Rumen Undegradable Protein Supply

$$RUP_i = CP_i - RDP_i$$

[AN.SUP.29]

$$RUP_{diet} == \sum_i feed_i * RUP_i * dRUP_i$$

[AN.SUP.30]

Endogenous Protein Supply



$$EndP = 0.4 * 11.8DMI$$

[31]

Mineral Supply

$$Ca_{supply} = \sum_i feed_i * Ca_i * \frac{dCa_i}{100}$$

[AN.SUP.32]

c. Automated Ration Formulation

Introduction Automated ration formulation in RuFaS optimizes a ration for a single objective (least cost), utilizing sequential least squares quadratic programming, as described in Li, Rosa, et al., 2022. This process is carried out on an individual pen level, and uses the mean animal requirements and body weight during formulation. The ration formulation process includes information not only from the Animal Module's inputs (Table RF1), but the primary Feed input file (Table RF2), and Feed ingredient inclusion rates defined in the Feed Storage module's input files

Required User Inputs

Table 18: Key inputs found in the animal input file.

Variable	Definition	Default	Min	Max
ration.user_input	User decision to use the user-defined ration formulation or the automated ration formulation methods (true or false, respectively).	TRUE	-	-
ration.formulation_interval	Ration formulation interval (days).	30	1	-
ration.phosphorus_requirement_buffer	Increase in phosphorus nutrient requirement in calculation of animal requirements (percentage).	75	0	-

Table 19: Key inputs found in the feed input file.

Section	Variable	Definition	Default	Min	Max
-	calf_feeds or growing_feeds or close_up_feeds or lac_cow_feeds	Each section is comprised of a list of feeds (RuFaS Feed IDs).	-	-	-
purchased_feeds	purchased_feed	RuFaS Feed ID	-	-	-
	purchased_feed_cost	Price per feed (dollars/kg DM).	-	-	-
	calves/growing/close_up/lac_cow.ration_percentages	Percentage of feed ID, amount of a ration that is a given feed (percentage).	-	0	100
User_defined_ration_percentages					



Section	Variable	Definition	Default	Min
	tolerance	Allowable +/- percentage variance in each of the defined ration inclusion percentage values (fraction).	0.1	0
	milk_reduction_maximum	Allowable amount of milk reduction (kg) when dietary nutrient supply cannot meet animal requirements.	15	0

Relevant Outputs

- Average nutrient requirements, on a per animal basis, for each pen.
 - *Class and function:* AnimalModuleReporter.report_ration_interval_data
 - *Output variable:* avg_rqmts_pen_pen.id_pen.animal_combination.name
- Each formulation interval's mean animal ration, on a per animal basis, for each pen.
 - *Class and function:* AnimalModuleReporter.report_ration_interval_data
 - *Output variable:* ration_per_animal_for_pen_pen.id_pen.animal_combination.name
- Nutrient and energetic composition of ration, on a per animal basis, for each pen.
 - *Class and function:* AnimalModuleReporter.report_ration_interval_data
 - *Output variable:* ration_nutrient_amount_pen_pen.id_pen.animal_combination.name
- The total amount of Feeds, per pen.
 - *Class and function:* AnimalModuleReporter.report_daily_ration
 - *Output variable:* ration_daily_feed_totals_for_pen_pen.id_pen.animal_combination.name
- In cases of ration formulation failure, a summary of the constraints that failed, and the attempted ration and its supply will be supplied in a report.
 - *Class and function:* RationOptimizer.handle_failed_constraints
 - *Output variable:* failed_constraint_summary_for_pen_pen.id

Methodology Ration formulation is carried out at the pen level on a user-defined formulation interval (Table RF1). The mean requirements for a given pen are calculated prior to first formulation, as described in [REF IV. Animal Nutrition and Feeding, A. Calculations of Animal Energy and Nutrient Requirement]. This process then proceeds to utilize a series of constraints methods (Table RF3) that compares the animal requirements against the supply of the optimizer's ration attempt, as described in [REF IV. Animal Nutrition and Feeding, B. Calculations of Animal Energy and Nutrient Supply], all while abiding feed ingredient-specific inclusion rates, as defined in [REFERENCE TO NRC/NASEM LIBRARY].

If a ration cannot be formulated that meets the constraints given the pen-level requirements and limits to inclusion rates, then one of the following things will happen.

For lactating cow pens

- Record failed constraints and formulation attempt to OutputManager.
- Milk production will be reduced.
 - If average milk production for the pen drops below the values defined in [REFERENCE TO 5. Milk Production and Reduction], then the simulation halts with an error.



- Recalculation of the nutritional requirements for all animals in the pen.
- Ration formulation reattempt (repeat).

For growing and close-up pens

- Record failed constraints and formulation attempt to OutputManager.
- IF a successful ration from a prior formulation is available, said ration will be used.
- IF no previously successful ration is available, the simulation will stop.

Note: A graphical overview of this process can be found in Figure 4 in II Animal Management, C. Milk Production

The generalized equation for sequential quadratic programming is as follows (Li, Rosa, et al., 2022) and $g_i(x)$ is constraint i , which must be twice continuously differentiable with respect to all x_j in x .”).

INSERT EQUATION HERE

Table 20: Key inputs found in the feed input file specific to user defined ration methodology. All are found in the section `user_defined_ration_percentages`

Nutrient or Energy	Min	Max	Animal Groups
Dry matter intake	80% of requirement	120% of requirement	All except calves
Metabolizable protein	100% of requirement	150% of requirement	All except calves
Net energy: lactation	100% of requirement		Lactating pen only
Net energy: maintenance and activity	100% of requirement		All except calves
Net energy: growth	100% of requirement		All except calves
Total net energy	100% of requirement		Lactating pen only
Calcium	100% of requirement		All except calves
Phosphorus	100% of requirement		All except calves
Fat		7% of ration	All except calves
NDF	25% of ration	45% of ration	All except calves
Forage NDF	15% of ration		All except calves

d. User-defined ration formulation

Introduction The user-defined ration methodology follows the same process as described in the Automated ration formulation methodology, with the addition of the inputs found in Table RF4, as described in the Methodology below.

Required User Inputs



Table 21: Key inputs found in the feed input file specific to user defined ration methodology. All are found in the section `user_defined_ration_percentages`

Variable	Definition	Default	Min	Max
calves or growing or close-up or <code>lac_cow.ration_percentage</code>	For each feed ID, amount of a ration that is a given feed (percentage).		0	100
tolerance	Allowable +/- percentage variance in each of the defined ration inclusion percentage values (fraction).	0.1	0	1
milk_reduction_maximum	Allowable amount of milk reduction (kg) when dietary nutrient supply cannot meet animal requirements.	15	0	50

Methodology The key difference in the methodology for the user-defined ration methodology is that the feed inclusion rates are bounded by the `user_defined_ration_percentage` values defined for each feed item (Table RF4). Those percentages are bounded by $\pm$ the tolerance value (Table RF4) as a fraction of the percentage.

The ration formulation will proceed as described in the [Automated Methodology](#) above, but when formulation fails, different outcomes will occur.

For lactating cow pens

- Record failed constraints and formulation attempt to OutputManager.
- Milk production will be reduced.
 - IF average milk production for the pen drops below the values defined in [REFERENCE TO 5. Milk Production and Reduction], then the simulation halts with an error.
 - IF if the `milk_reduction_maximum` (Table RF4) is exceeded, then the simulation proceeds by using the user-defined ration percentages exactly.
- Recalculation of the nutritional requirements for all animals in the pen.
- Ration formulation reattempt (repeat).

For growing and close-up pens

- Record failed constraints and formulation attempt to OutputManager.
- Recording of which constraints were unable to be met.
- The simulation proceeds by using the user-defined ration percentages exactly.

D. Herd Size Management

If the number of close-up heifers is more than the herd needs, heifers in the Heifer III group are sold. If the number of close-up is less than the herd needs, a replacement is bought from the market. This is evaluated daily by comparing the number of (cows + Heifer III) to the herd num specified by the user. Details of this method are described in the Culling section of Animal-level processes and management.

Herd Size Management: Heifer III Sales

If the number of the heifers is more than the herd needs, heifers in the Heifer III group are sold. If the number of Heifer III is less than the herd needs, a replacement is bought from the market. This is evaluated daily by comparing the number of (cows + Heifer III) to the herd.num specified by the user.

User Inputs and Constants



- `Herd_num`: the target number of cows (dry + lactating) to maintain
- `SELLING_THRESHOLD` = 1.03 – hard-coded value (animal constant) used to determine threshold above `herd_num` at which heiferIII's should be sold.
- `BUYING_THRESHOLD` = 1.01 – hard-coded value used to determine threshold below `herd_num` at which heiferIII's should be bought

Parameters and Equations

- `HerdManager.check_if_heifers_need_to_be_sold()`. Evaluates if the current number of Heifer III and cows exceeds a specified threshold (defined as 3% of the herd statistics' target herd size). If the threshold is surpassed, Heifer IIIs are removed from the herd until herd size falls within acceptable range.
 - **Called by** `HerdManager.daily_routines()`
 - Returns a list of animals removed
 - Logic - While the number of (Heifer IIIs + cows) is greater than `herd_num * selling_threshold`:
 - * The last heifer III is removed and is added to `animals_removed`
 - * Information about that heifer III is added to `sold_heiferIIIs_info`
 - * The number of heifer III's is decremented by 1
 - * The number of `sold_heiferIII_oversupply_num` is incremented by 1
 - This cycle iterates until the number of (Heifer IIIs + cows) is equal to `herd_num * selling_threshold`
- `HerdManager.check_if_replacement_heifers_needed()` Determines whether additional Heifer IIIs need to be added to the herd based on the current herd size, purchase thresholds, and the availability of heifers in the replacement market.
 - **Called by** `HerdManager.daily_routines()`
 - Returns a list of animals purchased
 - While the number of (Heifer IIIs + cows + `bought_heifer_num`) is less than `herd_num * buying_threshold`:
 - * The first animal in the `replacement_market` enters the herd, assigned a phosphorus requirement and net merit value, and added to the `animals_added` list
 - * Number of `bought_heifer_num` is incremented by 1
 - This cycle iterates until the number of (Heifer IIIs + cows + `bought_heifer_num`) is equal to `herd_num * buying_threshold`

Reproductive Culls: Heifer II Sales

If the heifer does not have a successful conception or pregnancy and age (days born) exceeds heifer repro cull time, this heifer is culled

User Inputs and Constants

- `Heifer_repro_cull_time`: days old when a heifer would be culled for failure to become pregnant (Default: 500 d)

Parameters and Equations

- `Animal.evaluate_heiferII_for_culling()`
 - **Called by** `Animal.daily_routines()` → `Animal.animal_life_stage_update()`
 - This method determines whether a heifer should be culled based on its pregnancy status and age, returning *True* if the heifer is not pregnant and has surpassed the specified culling age.



Calf Sales

All live male calves are sold immediately after birth. Female calves are sold/kept according to a roll of the dice evaluated against the user-defined rate of keeping female calves.

User Inputs and Constants

- `Keep_female_calf_rate`: the percentage of female calves kept and raised on-farm (Default: 1)

Parameters and Equations

- `Animal._initialize_newborn_calf()` to `self.sold`
 - **Called by** `HerdManager._create_newborn_calf()` in `_perform_daily_routines_for_animals()`
 - * Creates a newborn calf instance
 - Contains logic that determines whether a calf should be sold, returning True if the calf is a male or if a random draw generates a number higher than the `keep_female_calf_rate`
 - If assigned to be sold, `sold_at_day = simulation_day`

Calf Losses (Deaths) A portion of calves are born dead according to the user-defined death rate.

User Inputs and Constants

- `Still_birth_rate`: rate of stillbirths (percent, 0-1) (Default: 0.065)

Parameters and Equations

- `Animal._initialize_newborn_calf()`
 - **Called by** `HerdManager._create_newborn_calf()` in `_perform_daily_routines_for_animals()`
 - * Creates a newborn calf instance
 - If a random draw is less than the `still_birth_rate`, then `self.sold = True` and `STILL_BIRTH` is added to `self.events`
 - If assigned to be sold, `sold_at_day = simulation_day`

IV. Animal-level Processes and Management

The methods in the RuFaS model that simulate individual animals' life events and the effects of herd management on individual animals are referred to as the Animal Life Cycle Sub-module. This part of the model is driven by a Monte Carlo stochastic process that simulates the growth, production, reproduction, and culling of individual dairy cattle animals and shares aggregated outcomes with the Herd Management Sub-Module to manage herd dynamics on a daily basis.

The life cycle module represents the variation in animal performance through the use of Monte Carlo Simulation methods (Reuven and Kroese, 2017). Monte Carlo simulation relies on random draws from probability distributions rather than assigning fixed values. In this model, two main strategies are used:

- **Monte Carlo Simulation (Event-Based):** Comparing a random draw from $U(0, 1)\rho_{Uniform}$ to the probability of an event occurring to simulate if that event occurs or did not (RandomUni in this document representing a random draw from $U(0, 1)\rho_{Uniform}$).
- **Random Population Simulation:** Selecting a random draw from a known distribution of animal attributes and assigning that value to the instantiation of an individual animal. (Random in this document representing a random draw from the following distribution) Probability distributions used are:
 - **Normal Gaussian:** $N(\mu, \sigma)$ where μ is the distribution mean, and σ is the standard deviation.
 - **Empirical:** $F(x) = \frac{1}{n} \sum_{i=1}^n 1_{\{x_i \leq x\}}$: where x is the observations from the sample and an indicator function equal to 1 if and otherwise.



Flow of information For each animal object, the daily updates are called according to the order described in the Day in the Life for Animal on a RuFaS Farm. This is:

DigestiveSystem → NutrientInputs → MilkProduction → BodyweightChange → Reproduction → LifeStage Change → Parturition

The processes that determine significant events in each animal's life cycle occur in the

1. Reproduction,
2. LifeStage Change, and
3. Parturition updates.

Within the Reproduction update methods, there are separate methods to simulate breeding management for heifers and cows. All of the Reproduction update methods use an event-based Monte Carlo model to stochastically simulate the probability that each animal will conceive, the date they conceive, and the success of the pregnancy. In addition to recording success and failures to conceive, the Reproduction update records hormone deliveries and number and success of pregnancy checks. All methods are based on protocols described by the CDCB and are described in more detail in the Reproduction Section.

Each animal within the RuFaS herd will progress through the 5 main life stages from calf, to Heifer I, to Heifer II, to Heifer III, to Cow followed by a cull stage as illustrated in the figure below:




Figure 1: There are 5 life stages from calf to cow and a cull stage as outlined in this figure.

Progression Between Life Stages: Calves progress to Heifer I when their age reaches the user-defined input for weaning day. Heifer I animals progress to Heifer II animals when their age reaches the user-defined input for breeding start day. Heifer II animals that successfully conceive progress to the Heifer III stage when they are close to calving. Specifically, Heifers II become Heifers III when the number of days left in their pregnancy is equal to the user-defined input for the prefresh days. When a Heifer III animal calves for the first time, she becomes a Cow and will remain a cow for the rest of her life on the RuFaS farm.

When a Heifer III calves for the first time and becomes a Cow, then that animal will cycle through lactating and non-lactating phases until they are removed from the herd due to a failure to conceive or a user-provided probability for culling within each parity. The methods that determine when a cow will leave the herd are described in detail in the Herd Exits section.

Parturition and New Calves: When a Heifer III or Cow animal reaches the end of their pregnancy and starts lactating, a new calf animal object is created and assigned a birthweight according to a user-informed random distribution. All male calves are recorded and removed from the farm on day 1 of their life. Female calves remain on the farm determined by an event-based Monte Carlo method informed by the user-defined percent of female calves that are kept.



A. RuFaS Bodyweight and Growth

Introduction An animal's bodyweight is initialized at birth according to a user defined normal distribution. Depending on her life stage, her bodyweight is then updated daily by adding their daily growth (commonly referred to as average daily gain), conceptus weight change, and tissue accretion and depletion associated with tissue mobilization to support lactation. The changes associated with each life stage are:

- Calves increase bodyweight by calf specific estimate for daily growth from birth to weaning.
- Non-pregnant heifers increase bodyweight by non-pregnant heifer specific estimate for daily growth from weaning until conception.
- Pregnant heifers increase their bodyweight by the pregnant heifer specific estimate for daily growth and an estimate for conceptus growth.
- Cows in parity 1 and 2 increase their bodyweight according to a parity and pregnancy status specific estimate for daily growth, an estimate for conceptus growth, and an estimate for body tissue changes that occur in lactation.
- Cows in parities 3 and above change their bodyweight according to estimates for conceptus growth and body tissue changes that occur in lactation.

These methods are further described by Li, Reed, et al. (2023)

Required User Inputs

Table 22: Below is a summary of the user inputs that are available and their definitions.

Variable	Definition (units)	Default	Min	Max
birth_weight_avg_ho	Average Holstein Birth Weight (kg/head)	42.9	1	NA
birth_weight_std_ho	Holstein Birth Weight Standard Deviation (kg/head)	69	0	NA
birth_weight_avg_je	Average Jersey Birth Weight (kg/head)	25.2	1	NA
birth_weight_std_je	Jersey Birth Weight Standard Deviation (kg/head)	4.4	0	NA
mature_body_weight_avg	The average mature body weight of cows (kg/head)	740.1	0	NA
mature_body_weight_std	The standard deviation of mature body weight of cows (kg/head)	73.5	0	NA
wean_day	The days of age at which calves are fully weaned from milk or milk replacer (days)	60	0	NA
target_heifer_preg_day	The target age (in days) for heifers to become pregnant - for adjusting heifer body weight (days)	420	NA	NA

Expected Outputs

- herd_statistics.avg_cow_body_weight (kg): the daily average bodyweight of all cows in the herd
- Pen_[pen ID]_[pen animal combination].avg_BW (kg): the daily average bodyweight of all animals in each pen



- sold_weight (kg)

Methodology Most methods determining the bodyweight of animals in RuFaS are implemented in growth.py.

The function evaluate_body_weight_change in growth.py calls the appropriate bodyweight change functions for each animal depending on their life stage and sums the appropriate changes into a single variable called daily_growth.

After setting the daily bodyweight change in the daily_growth variable, the evaluate_body_weight_change function updates each animal's bodyweight by adding it to the current body_weight. Thus for all animals, the final updated body_weight is set as:

$$\text{body_weight}_{\text{updated}} = \text{body_weight}_{\text{current}} + \text{daily_growth}$$

- **Calf Birth Weight** is assigned in reproduction.py using a random draw from a truncated normal distribution based on user inputs for the breed-specific average and standard deviation of the birthweight. The distribution is truncated at +/- 2 SD from the mean to prevent extremely small or large calves.

$$\text{Birth Weight} = \text{Random } N(\text{average_birth_weight_by_breed}, \text{std_std_birth_by_breed})$$

*Note - truncated at 2 standard deviations

[AN.BWT.6]

- **Calf Growth** Calves are assumed to double their weight between birth and weaning based on recommendations in the National Research Council, 2001. Thus the target average daily gain is estimated as:

$$\text{target_average_daily_gain} = \frac{\text{birth_weight}}{\text{wean_day}}$$

[AN.BWT.7]

- **Non Pregnant Heifer Growth** The daily growth for non-pregnant heifers is estimated through an average daily gain to reach 55% of their mature bodyweight by the time they are pregnant. Because the actual age at pregnancy is not known, a user input for the target_heifer_preg_day is used to estimate the target average daily growth or ADG. A minimum daily growth rate of 0.5 kg/d is enforced so if the estimated daily_growth is less than 0.5 kg/d, the minimum value is used.

$$\text{target_average_daily_gain} = \min\left(\frac{0.55\text{MSBW} - \text{SBW}}{\text{abs}(\text{target_heifer_pregnant_age} - \text{age})}, 0.5\right)$$

[AN.BWT.8]

Where MSBW is mature shrunk bodyweight (0.96 x mature bodyweight) and SBW is shrunk bodyweight (0.96 x bodyweight)



- **Pregnant Heifer Growth** Heifers are targeted to grow to 82% of their mature body weight by the time of their first calving. A gestation length is set at conception. The ADG of pregnant heifers is calculated based on the target heifer's body weight, gestation length, and days in pregnancy. The gestation length for each animal is set at conception by `calculate_gestation_length` in `reproduction.py`.

$$target_average_daily_gain = \frac{0.82MSBW - SBW}{gestationlength - daysinpregnancy}$$

[AN.BWT.9]

- **Parity 1 and 2 Cow Growth** The daily growth of cows is calculated based on the target of reaching 92% of the mature body weight by the end of the 1st lactation, and full mature body weight at the end of the 2nd lactation. Before pregnancy, the daily growth is estimated based on the calving interval and during pregnancy the daily growth is calculated based on days until calving.

Parity 1 Animals: *If not pregnant:*

$$target_average_daily_gain = \frac{(0.92 - 0.82)MSBW}{calving_interval}$$

[AN.BWT.10]

If pregnant:

$$target_average_daily_gain = \frac{0.92MSBW - bodyweight}{gestation_length - DIP}$$

[AN.BWT.11]

Parity 2 Animals *If not pregnant:*

$$target_average_daily_gain = \frac{(1 - 0.92)MSBW}{calving_interval}$$

[AN.BWT.12]

If pregnant:

$$target_average_daily_gain = \frac{(MSBW - bodyweight)}{gestation_length - DIP}$$



[AN.BWT.13]

The gestation length for cows is estimated with the same methods that are used for heifers and the calving interval is set either as the average herd calving interval or the animal's own calving interval which is calculated as the age at her second calving minus her age at first calving as part of the `daily_reproduction_update` in `animal.py`.

- **Conceptus Growth** is estimated the same way for both heifers and cows using a non-linear model for animals that are greater than 50 days in pregnancy based on the methods developed by Korver, Arendonk, and Koops (1985).

If the heifer or cow days in pregnancy is greater than 50 and less than the gestation length:

First the expected total conceptus weight is estimated using an empirical equation based on the calf birth weight and the gestation length:

$$conceptus_total_weight = (0.0148 * gestation_length) calf_birth_weight$$

[AN.BWT.14]

Then an animal specific conceptus parameter is calculated:

$$conceptus_parameter = \frac{total_conceptus_weight^{\frac{1}{3}}}{gestation_length - 50}$$

[AN.BWT.15]

Finally, the conceptus growth is estimated as:

$$conceptus_growth = 3 * conceptus_parameter^3 * (DIP - 50)^2$$

[AN.BWT.16]

When `days_in_pregnancy = gestation_length` and the animal calves, then conceptus growth is set to the negative value of the current `conceptus_weight` such that the weight of the calf and placenta are subtracted from the cow's bodyweight when her bodyweight is updated.

$$conceptus_growth = conceptus_weight$$

[AN.BWT.17]

- **Tissue Change Due to Lactation** Lactation related tissue changes are estimated by a non-linear model presented by Galvão et al. (2013) that estimates the daily tissue mobilized during lactation.



$$bodyweight_tissue_change = \frac{P_1}{P_2} * \exp\left(1 - \frac{DIM}{P_2}\right) + DIM * \exp\left(1 - \frac{DIM}{P_2}\right)$$

[AN.BWT.18]

Where P_1 and P_2 are parameters with distinct values for primiparous and multiparous cows:

Parameter	Primiparous	Multiparous
P_1	20	65
P_2	40	70

During the dry period, the net tissue change is assumed to be restored during the dry period. The net tissue change is estimated on the last day of lactation as:

$$tissue_changed = P_1 * \frac{DIM_{last}}{P_2} * \exp\left(1 - \frac{DIM_{last}}{P_2}\right) + DIM * \exp\left(1 - \frac{DIM}{P_2}\right)$$

[AN.BWT.19]

$$bodyweight_tissue_change = \frac{tissue_changed}{gestation\ length - DIP_{dry}}$$

[AN.BWT.20]

B. Animal Reproduction

Introduction The reproduction component of the RuFaS model is designed based on reproductive program options recommended by the Dairy Cattle Reproduction Council (Dairy Cattle Reproduction Council, 2018) and experts in the field. It offers three pre-defined reproductive program options for nulliparous heifers (estrus detection [ED], timed artificial insemination [TAI], and synchronized estrus detection [Synch-ED]) and three options for lactating cows (ED, TAI, and a combined ED-TAI approach). Users can select appropriate reproductive protocols and specify reproductive performance variables for both nulliparous heifers and lactating cows.

This reproduction component simulates key reproductive events, including estrus occurrence, estrus detection, hormone administration, artificial insemination (AI), pregnancy checks, and pregnancy outcomes (abortion or successful delivery). Reproductive culling decisions are also modeled, based on an animal's inability to conceive or sustain a pregnancy. These processes are modeled using Monte Carlo simulations to represent variability in reproductive events.

The simulation generates and tracks detailed reproductive performance metrics at the animal level. These outputs provide insights into estrus detection, synchronization treatments, breeding efficiency, and pregnancy outcomes. The key recorded variables include:

- Estrus Detection (ED) Metrics: Number of days an animal participates in the ED program (ED_days) and the total estrous events detected (estrus_count).
- Hormonal Treatments: Counts of synchronization protocol injections, including GnRH (GnRH_injections), PGF (PGF_injections), and CIDR (CIDR_injections).



- Breeding Interventions: Number of semen straws used (semen_number) and total AI procedures performed (AI_times).
- Reproductive Efficiency: Time from first breeding attempt to pregnancy confirmation (breeding_to_pregnancy_time) and the interval from calving to confirmed pregnancy (calving_to_pregnancy_time).
- Pregnancy Monitoring: Total number of pregnancy checks performed (pregnancy_diagnoses).

Required User Inputs

Table 23: Below are the key user inputs that determine the **heifer reproductive breeding program** and impact reproductive efficiency and age at first calving.

Section	Variable	Definition	Default	Min	Max
management_decisions	breeding_start_day_h	Age when RuFaS initiates estrous protocol (days)	-	380	0
	heifer_repro_method	TAI (Timed AI), ED (Estrus Detection), Synch-ED (Synchronized ED)	TAI	TAI, ED, Synch-ED	
	heifer_repro_cull_time	Age when a heifer is culled if not pregnant (days)	-	500	0
farm_level_heifers	estrus_detection_rate	Proportion of estrus detected in ED protocol	0.9	0	1
	estrus_conception_rate	Conception rate in ED protocol	0.6	0	1
	repro_sub_protocol	Hormone protocol used; varies by method (e.g., 5dCG2P, CP, N/A)	5dCG2P	5dCG2P, 5dCGP, 2P, CP, N/A	
	conception_rate	Conception rate for sub-protocol	0.6	0	1
	estrus_detection_rate	Detection rate in sub-protocol	0.6	0	1

Table 24: The following inputs determine reproductive efficiency of cows and influence culling, calving interval, and calf output.

Section	Variable	Definition	Default	Min	Max
management_decisions	cow_repro_method	TAI, ED, ED-TAI protocols	TAI		TAI, ED, ED-TAI
	do_not_breed_time	Days after calving to stop breeding if not pregnant	185	0	-
farm_level	voluntary_waiting_period	Days before breeding starts for ED/ED-TAI; ignored in TAI	50	0	-
	estrus_detection_protocol	Estrous protocol type	Double OvSynch	Double OvSynch, PreSynch, G6G, None	
farm_level_cows	estrus_conception_rate	Conception rate for ED/ED-TAI	0.5	0	-
	presynch_program	Presynch method used in TAI	OvSynch 56	OvSynch 48, OvSynch 56, CoSynch 72, 5d CoSynch, None	
	presynch_program_start_day	Day of first hormone injection for presynch	70	0	-
	ovsynch_program	OvSynch protocol used in TAI or ED-TAI	0.6	0	1
	ovsynch_program_start_day	Start day for OvSynch protocol	TAIafterPD	TAIafterPD, TAIbeforePD, PGFatPD, None	
	ovsynch_program_conception_rate	Conception rate for OvSynch program	-	-	-
	resynch_program	Resynch protocol around pregnancy check	-	-	-

NOTE: Anytime an insemination is performed after an OvSynch program, the conception rate will be changed to the ovsynch_program_conception_rate.

Table 25: The following inputs can be used to determine if there is a **decrease in the conception rate** with parity and rebreeding attempts.

Section	Variable	Definition	Default	Min	Max
farm_level	conception_rate_decrease	Percent decrease per breeding attempt after first	0.026	0.1	0
	decrease_conception_rate_in_rebreeding	Whether to decrease rate with rebreeding	FALSE	-	-
	decrease_conception_rate_by_parity	Whether to decrease rate by parity	FALSE	-	-



Relevant Outputs

Overall reporting for reproductive measures can be summarized and split into scheduled events and updates on an individual basis for heifers and cows.

These include:

- Events for Estrus (detection and scheduling)
- Hormone Administration
- Artificial Insemination (AI) (scheduling and status)
- Pregnancy Checks and Status
- Days Carry Calf
- Abortion Events
- Updates to Lactation
- Updates to Estrous
- Updates to Pregnancy
- Updates to Reproductive Status or Culling Decisions
- Updates to Protocols

Methodology *Heifer Breeding*

Heifer II - The `heiferII_reproduction_update` represents the breeding-eligible heifer stage, beginning when the animal enters a reproductive management program and ending when she reaches a user-defined day in pregnancy that marks the start of the pre-fresh period (`prefresh_day`). The breeding process follows the assigned reproductive management program and is simulated as described below (Figure 2). Each program includes a user-defined conception rate (`conception_rate`), determining the probability of pregnancy initiation at the time of insemination.




Figure 2: This is a high level schematic to illustrate the methodology behind the heifer breeding and re-breeding in RuFaS Animal Module.

Pregnancy diagnosis and reconfirmation occur after each insemination [heifer_pregnancy_update]. If a heifer remains nonpregnant beyond a user-defined age threshold (heifer_repro_cull_time, e.g., 500 days), she is culled due to reproductive failure. At a user-defined day in pregnancy (prefresh_day, e.g., one month before calving), the animal either transitions to the Heifer III stage or is sold as a replacement heifer.

The first-service reproductive management strategy for a heifer follows one of three strategies and are described in more detail below: estrus detection (ED), synchronized estrus detection (Synch-ED), or timed artificial insemination (TAI).

1. **ED [execute_heifer_ed_protocol]:** The ED strategy simulates spontaneous estrous cycles. Estrous cycle length is determined using a random draw from a normal distribution (De Vries et al., 2006). On the day of estrus predicted by the model, a random draw from a uniform distribution $U(0,1)$ is compared with the estrus detection rate (estrus_detection_rate). If the drawn value is lower than the detection rate, AI is scheduled on that day. If estrus is not detected, the next estrus event is scheduled based on the normal estrous cycle. On the scheduled AI day, pregnancy success is determined using a random draw from a uniform distribution $U(0,1)$, which is compared against the conception rate (conception_rate).
2. **Synch-ED [execute_heifer_synch_ed_protocol]:** In the Synch-ED strategy, hormonal treatment dates are scheduled according to the assigned program (repro_sub_protocol, 2P or CP), and estrus is simulated based on a program-specific estrous cycle following the hormonal treatment. Estrus detection and AI scheduling follow the same procedure as in the ED program.
3. **TAI [execute_heifer_tai_protocol]:** The TAI strategy simulates hormonal treatments administered on scheduled days from the start of the synchronization protocol. The timing of treatments and AI is modeled according to the assigned specific program (5dCGP or 5dCG2P) defined as a sub-protocol in the user inputs (repro_sub_protocol). On the scheduled AI day, pregnancy success is determined using a random draw from a uniform distribution $U(0,1)$, which is compared against the conception rate (conception_rate).



For all heifers who fail to conceive after the first insemination or experience pregnancy loss [open_heifer], the ED strategy is applied for subsequent breeding attempts.

Cow Breeding The breeding process for cows begins after calving and the voluntary waiting period (VWP), following one of three reproductive management strategies available for cows which are similar to those described in the heifer section and are pictured in Figure 3: ED, TAI, or ED-TAI. The program assigned to a cow determines the timing of insemination and the probability of pregnancy initiation.

- **ED [execute_cow_ed_protocol]:** In the ED strategy, spontaneous estrous cycles are simulated. The first estrous cycle after calving is determined on the calving date using a random draw from a normal distribution with user-specified mean and standard deviation (Kalantari, 2015). The subsequent estrous cycles are modeled similarly. If the scheduled estrus occurs before the VWP, the estrus is ignored. If the scheduled estrus occurs after the VWP, estrus detection is simulated following the same procedure as in the heifer ED strategy - a random draw from a uniform distribution $U(0,1)$ is compared with the estrus detection rate (estrus_detection_rate). If the drawn value is lower than the detection rate, AI is scheduled on that day. If estrus is not detected, the next estrus event is scheduled according to the estrous cycle length. On the scheduled AI day, pregnancy success is determined using a random draw from a uniform distribution $U(0,1)$, which is compared against the conception rate (ED_conception_rate).
- **TAI [execute_cow_tai_protocol]:** The TAI strategy simulates hormonal treatments administered on scheduled days according to the assigned synchronization protocol after the start of the breeding date (ovsynch_program_start_day). The timing of treatments and AI follow the predefined protocols (ovsynch_program, OvSynch-48, OvSynch-56, CoSynch-72, or 5d CoSynch). These protocols can be implemented alone or in combination with presynchronization protocols (presynch_program, PreSynch, Double OvSynch, or G6G). On the scheduled AI day, pregnancy success is determined using a random draw from a uniform distribution $U(0,1)$, compared against the conception rate.
- **ED-TAI [execute_cow_ed_tai_protocol]:** The ED-TAI strategy integrates ED with TAI-based synchronization programs. Estrus detection is simulated before TAI begins. If estrus is detected after VWP (voluntary_waiting_period) and before the initiation of the TAI (ovsynch_program_start_day), AI is scheduled according to the estrus detection protocol. If no estrus is detected, the cow follows the TAI protocol for insemination.

Important Note

For rebreedings, cows can be assigned to any of the three rebreeding strategies (resynch_program, TAIbeforePD, TAIafterPD, or PGFatPD) [open_cow].

- **TAIbeforePD:** The TAIbeforePD rebreeding strategy initiates a timed AI synchronization protocol before the first pregnancy diagnosis (PD). Cows that are due for pregnancy diagnosis but have not been confirmed pregnant yet start one of the predefined synchronization protocols (OvSynch-48, OvSynch-56, CoSynch-72, or 5-day CoSynch) six days before PD. This approach ensures that if a cow is found open (not pregnant) on PD day, she is already progressing through a synchronization protocol, reducing the waiting period before the next AI attempt.
- **TAIafterPD:** The TAIafterPD rebreeding strategy schedules synchronization for cows that are diagnosed as open on the PD day. If a cow is not pregnant, she is immediately enrolled in one of the available timed AI protocols (OvSynch-48, OvSynch-56, CoSynch-72, or 5-day CoSynch). This ensures timely rebreeding but does not initiate synchronization in advance of PD.
- **PGFatPD:** The PGFatPD rebreeding strategy uses prostaglandin (PGF) administration following PD for open cows to induce estrus. After PGF treatment (day 0), cows are monitored for natural estrus. If estrus is detected, AI is performed accordingly. If no estrus is detected within seven days, the cow is enrolled in a timed AI protocol (OvSynch-48, OvSynch-56, CoSynch-72, or 5-day CoSynch) to ensure insemination. This program balances estrus detection with timed AI, allowing for natural breeding opportunities before initiating synchronization.




Figure 3: This is a high level schematic to illustrate the methodology behind the cow breeding and re-breeding in RuFaS Animal Module.



Pregnancy Diagnosis (PD) and Pregnancy Loss For both heifers and lactating cows [heifer_pregnancy_update, cow_pregnancy_update], pregnancy diagnoses are scheduled at three intervals after AI (preg_check_day_1, preg_check_day_2, preg_check_day_3). The default settings simulate the first at 32 d after insemination, which, if confirmed positive for pregnancy, is followed at 60 d and then at 200 d (Wiltbank et al., 2016); however, the number and day of pregnancy checks can be adjusted by the model user according to farm practices. Between AI and three pregnancy checks, pregnancy losses are simulated to occur according to random draws from a U (0,1) compared with thresholds according to the expected pregnancy losses in each of those periods (preg_loss_rate_1, preg_loss_rate_2, preg_loss_rate_3). The default values are 0.02, 0.096, and 0.017 (Wiltbank et al., 2016). Users can adjust these numbers according to their herd performance.

On the next page are some important functions for the reproduction section of this document

Reproduction Functions

Table 26: Important functions for the reproduction section.

Function	Description	Parameters	Input Origins	Outputs
reproduction_update	Updates the reproductive status of all animals daily, including estrus occurrence, detection, AI scheduling, pregnancy status, and culling decisions.	day: (int) Current simulation day. animal_list: (list) List of all animals in the herd.	day: Updated by the simulation engine. animal_list: Provided as model input, updated dynamically.	Updated reproductive status for all animals in the herd. Scheduled events for estrus, AI, and pregnancy checks.
heiferII_reproduction_update	Manages reproduction processes for Heifer II animals, including estrus detection, breeding attempts, and pregnancy checks.	heifer: (object) Individual Heifer II animal.	heifer: Passed from the herd's current Heifer II group.	Updated estrus, AI status, and pregnancy details for the heifer.
cow_reproduction_update	Handles reproduction for lactating cows, including estrus detection, AI, and transitioning reproductive status.	cow: (object) Individual cow object. day: (int) Current simulation day.	cow: Passed from the herd's lactating group. day: Updated by the simulation engine.	Updated estrus, AI, pregnancy status, and culling decisions for the cow.
cow_give_birth	Manages calving events, updates the reproductive status, and initializes the lactation process.	cow: (object) Individual cow object. day: (int) Current simulation day.	cow: Passed from the pregnant group. day: Updated by the simulation engine.	Updated lactation and reproductive status for the cow.
execute_heifer_ed_protocol	Executes the estrus detection protocol for heifers, simulating natural estrus occurrence and detection probability.	heifer: (object) Individual Heifer II animal. day: (int) Current simulation day.	heifer: Passed dynamically during estrus updates. day: Updated by the simulation engine.	Estrus detection and AI scheduling for the heifer.
execute_heifer_tai_protocol	Manages the TAI protocol for heifers, simulating hormone administration and AI scheduling.	heifer: (object) Individual Heifer II animal. protocol: (dict) User-defined TAI protocol details.	heifer: Passed dynamically during protocol execution. protocol: Provided as user input.	AI scheduling and protocol updates for the heifer.
execute_heifer_synch_ed_protocol	Executes synchronized estrus detection protocol for heifers, combining hormonal synchronization and estrus detection.	heifer: (object) Individual Heifer II animal. protocol: (dict) User-defined Synch-ED protocol details.	heifer: Passed dynamically during protocol execution. protocol: Provided as user input.	Estrus detection, hormone administration, and AI scheduling for the heifer.

Function	Description	Parameters	Input Origins	Outputs
open_heifer	Handles open status updates for heifers, reassigning protocols or marking for culling if reproduction fails.	heifer: (object) Individual Heifer II animal.	heifer: Passed dynamically during daily updates.	Updated reproduction status or culling decision for the heifer.
heifer_pregnancy_update	Updates pregnancy status for heifers, including pregnancy checks and abortion events.	heifer: (object) Individual Heifer II animal. day: (int) Current simulation day.	heifer: Passed dynamically during daily updates. day: Updated by the simulation engine.	Pregnancy status, days in pregnancy, or abortion event for the heifer.
execute_cow_ed_protocol	Executes estrus detection protocol for cows, simulating natural estrus occurrence and detection probability.	cow: (object) Individual cow object. day: (int) Current simulation day.	cow: Passed dynamically during estrus updates. day: Updated by the simulation engine.	Estrus detection and AI scheduling for the cow.
execute_cow_tai_protocol	Manages the TAI protocol for cows, simulating hormone administration and AI scheduling.	cow: (object) Individual cow object. protocol: (dict) User-defined TAI protocol details.	cow: Passed dynamically during protocol execution. protocol: Provided as user input.	AI scheduling and protocol updates for the cow.
execute_cow_ed_tai_protocol	Executes ED-TAI protocol for cows, combining estrus detection with synchronized TAI scheduling.	cow: (object) Individual cow object. protocol: (dict) User-defined ED-TAI protocol details.	cow: Passed dynamically during protocol execution. protocol: Provided as user input.	Estrus detection, hormone administration, and AI scheduling for the cow.
cow_pregnancy_update	Updates pregnancy status for cows, including pregnancy checks and abortion events.	cow: (object) Individual cow object. day: (int) Current simulation day.	cow: Passed dynamically during daily updates. day: Updated by the simulation engine.	Pregnancy status, days in pregnancy, or abortion event for the cow.
open_cow	Handles open status updates for cows, reassigning protocols or marking for culling if reproduction fails.	cow: (object) Individual cow object.	cow: Passed dynamically during daily updates.	Updated reproduction status or culling decision for the cow.



C. Milk Production

Introduction This section describes how RuFaS determines the amount of milk and milk components produced by a particular cow on a particular day.

Milk yield is largely determined by lactation curve parameters. Wood's curve is the only lactation model implemented in RuFaS currently, but there is flexibility to potentially include other models as well. At the beginning of the simulation, the herd is assigned distributions for the Wood's curve parameters l , m , and n for each parity of cows (see more information in the [Lactation Curve](#) section). At the beginning of each new lactation, each cow is assigned a set of parameters for that lactation based on the herd's distributions for her current parity.

When the cow starts lactating, she produces milk from the calving day until dry or culled according to the assigned lactation curve model. Each day, each cow's milk production is calculated and recorded. Milk production is calculated from the Wood's curve equation using the given lactation curve parameters and days in milk, and then adjusted by applying random variation and any milk reduction required by other parts of the model. Milk components (protein, fat, and lactose) are calculated based on milk production and percent of each component for each cow.

Currently, the only source of milk reduction is if there is no possible ration using the feeds available that would support the baseline level of milk production. Ration formulation is attempted with reduced milk production levels until reaching an acceptable ration (that meets the nutrient requirements for the pen's average cow) or reaching a threshold (floor) milk production defined by some combination of user inputs and pre-defined constants.

Required User Inputs

Table 27: The following user inputs are used to estimate customized herd-level lactation curve parameters. See the Methodology section of this document and the Lactation Curve Scientific Documentation for more detail about how these inputs are used with relation to daily milk production.

Section	Variable	Definition	Default	Min	Max
config_properties	end_date	The year and Julian day on which the simulation will end	2009:100	- (cutoff at 2006)	– (cutoff at 2016)
	FIPS_county_code	Unique 5-digit code that represents a specific US county	-	1000	56045
animal_properties	cow_times_milked_per_day	Number of Milkings (per day) – The average or most common number of times cows are milked per day (1, 2, or 3 times daily)	3	0	-
	parity_fractions	Fractions of the milking animal population that are parity 1, 2, and 3 and beyond. The sum of these fractions must be 1.0	1: 0.346; 2: 0.272; 3: 0.379	1	0
	annual_milk_yield	The total milk yield generated by the farm in one year (kg). If this information is not available, it can be input as null	null	-	0
lactation_properties	milking_cow_fraction	Fraction of cows assumed to be in milk at any given time	0.8365 (305/365)	1	0

Table 28: The following inputs are used to determine the milk fat and milk protein percentages for each lactating cow

Section	Variable	Definition	Default	Min	Max
animal_properties	milk_fat_percent	The average or most common milk fat percentage in cow milk	3.5	0	
	milk_protein_percent	The average or most common milk protein percentage in cow milk	3.2	0	

Table 29: The following inputs are used to constrain how much an average cow's daily milk production will be reduced due to an inadequate ration. See the Methodology section of this document and the Ration Formulation scientific documentation for more detail about how these inputs are used with relation to daily milk production.

Section	Variable	Definition	Default	Min	Max
feed_properties	milk_reduction_maximum	Allowable amount of milk reduction (kg) when dietary nutrient supply cannot meet animal requirements		0	50
	tolerance	Allowable +/- percentage variance in each of the defined ration inclusion percentage values. <i>This variable defaults to 0.0 and currently has no effect due to no implementation of an automatic ration optimizer. RuFaS is hard-coded to use the user-defined ration at this time.</i>	0	0	1



Relevant Outputs

- `report_life_cycle_manager_data`:
 - Information added daily from herd statistics
 - `daily_milk_production`
 - `herd_milk_fat_percent`
 - `herd_milk_fat_kg`
 - `herd_milk_protein_percent`
 - `herd_milk_protein_kg`
- `report_average_nutrient_requirements`:
 - Reports the average ration per animal of each pen each time a new ration is formulated
 - `avg_milk_production_reduction_pen` (kg per animal)

Methodology

1. *Herd Level Lactation Curve Parameters*

At the beginning of the simulation, the herd is assigned distributions for the lactation curve parameters l , m , and n for each parity of cows. It calculates Wood's lactation curve parameter mean values by parity, adjusted based on the location, production, management practices, +/- reported milk production of the farm being simulated.

User Inputs and Constants

The following inputs are used to generate the herd's values for mean lactation curve parameters by parity.

- `year`: `end_date` in the input `config.json`. Uses 2016 if the end year is 2016 or later.
- `region`: assigns the appropriate region based on `FIPS_county_code` entered in the input `config.json`
- `milking frequency`: `cow_times_milk_per_day` in input `animal.json`
- `parity distribution`: `parity_fractions` in input `animal.json`
- `annual milk yield (optional)`: `annual_milk_yield` in `animal.json`
- `Milking_cow_fraction`: 0.8356 default value
 - `ls` in `lactation_curve_adjustment_inputs.json`
 - Fraction of lactating cows in adult cows, between 0 and 1.
 - Used to calculate adjusted lactation curve parameters to match a herd's reported annual milk production

Parameters and Equations

These are comprehensively outlined in the [Lactation Curve Parameters Determination](#) Section.

`LactationCurve.set_lactation_parameters()` is called by:

- `HerdFactory.initialize_herd()` to `Animal.setup_lactation_curve_parameters()`
- `HerdManager._init_()`



2. Cow Level Lactation Curve Parameters

Each cow is assigned a set of parameters for each lactation based on the herd's distributions of l , m , and n for her current parity. This occurs when a cow is initialized at the beginning of the simulation, when a heiferIII (close-up heifer) transitions into a cow, and each time a cow has a calf. It selects Wood's curve parameters for the cow for each parameter for each parity from the distributions based on the means calculated for the herd and constant standard deviations.

User Inputs and Constants

No additional inputs to consider.

Parameters and Equations

- LactationCurve.get_wood_parameters() is called by
 - Animal.initialize_cow
 - Animal.daily_reproduction_update
 - Animal.transition_heiferIII_to_cow
- Parity groupings are as follows: 1, 2, and 3+
- Immediately followed in each case by self.milk_production.set_wood_parameters() which assigns the l , m , and n values.

3. Daily Milk Production

Each day, each cow's milk production is calculated and recorded. Milk production is calculated from the Wood's curve equation using the given lactation curve parameters and days in milk. The calculated production is then adjusted by applying random variation and any milk reduction required by other parts of the model.

User Inputs and Constants

No additional inputs to consider but constants are below:

- DAILY_MILK_VARIATION_MEAN: 0 kg/d
- DAILY_MILK_VARIATION_STD_DEV: 1.0 kg/d

Parameters and Equations

$$\text{Calculated milk production} = l * t^m * e^{-nt}$$

[AN.MLK.9]

- l , m , n = lactation curve parameters
- t = DIM

Adjusted milk production = maximum of (calculated milk production + milk production variance - milk production reduction) or 0.

- MilkProduction.calculate_daily_milk_production()
 - **Called by** HerdManager.daily_routines() to Animal.daily_routines() to MilkProduction.perform_daily_milking_update()
 - **Parameters**
 - * Days_in_milk: Argument supplied from MilkProductionInputs (updated in Animal.daily_routines)



- * `L_param`, `m_param`, `n_param`: `Wood_l`, `wood_m`, `wood_n` supplied as arguments and assigned to individual on the day of calving via `set_wood_parameters()`
- **Returns** calculated milk yield on the provided day in kg
- `MilkProduction.daily_milk_produced()`
 - **Inputs**
 - * `_Daily_milk_produced` (calculated above)
 - * `Milk_production_variance` is generated based on the mean and standard deviation for daily milk variation (stored as constants in `AnimalModuleConstants`)
 - * `Milk_production_reduction`
 - **Returns** `adjusted_milk_production` (unless the milk yield calculated above was 0, in which case it returns 0)

4. Daily Components Production

Milk components (true protein, fat, and lactose) are calculated daily based on that day's milk production and percent of each component.

User Inputs and Constants

- `milk_fat_percent` – single constant value for all cows in the herd
- `milk_protein_percent` – single constant value for all cows in the herd
- `AnimalModuleConstants.MILK_LACTOSE`, current value 4.85

Parameters and Equations

- *Daily nutrient amount in milk = milk production * nutrient percentage * 0.01*
- `MilkProduction._calculate_nutrient_content()`
 - **Called by** `HerdManager.daily_routines()` to `Animal.daily_routines()` to `MilkProduction.perform_daily_milking_update()`
 - **Parameters**
 - * `Days_milk_produced` as calculated in the Daily Milk Production section above
 - * `Nutrient_percentage`: cow's percent true protein, fat, or lactose for each of the respective calculations
 - **Returns** Amount of a given nutrient in milk in kg

Please note that currently, true protein is the only milk protein currently used by the model. .

5. Milk Production and Reduction

Currently, the only source of milk reduction is if there is no possible ration using the feeds available that would support the baseline level of milk production.

Each time the ration is formulated, the nutrition requirements of the pen are calculated and used to determine the ration given to each animal. The ration is checked against the nutrition requirements of the pen. If the pen contains lactating cows and the ration does not meet the average cow's requirements, then each cow's milk production is reduced until one of three conditions is met:

- The ration meets the animal's requirement (i.e. no additional reduction).
- The milk production of every animal is reduced by the maximum allowed (a user-defined value with default of 0.5 kg/head/d). Then the final ration and milk reduction values are used.
- The average milk production of the pen falls below the minimum allowable average milk production. Then the final ration and milk reduction values are used.




Figure 4: A schematic illustrating the flow of milk reduction in ration formulation logic.

User Inputs and Constants

- `Milk_reduction_maximum`: Allowable amount of milk reduction (kg) when dietary nutrient supply cannot meet animal requirements
- `Tolerance`: Allowable +/- percentage variance in each of the defined ration inclusion percentage values. This variable is default to 0.0 and it currently has no effect due to no implementation of automatic ration optimizer.
- `Milk_reduction_kg` is set to 0.25 kg: Milk reduction amount for each failed ration attempt, kg.
- `Minimum_avg_pen_milk` is set to 15 kg: Minimum allowable average milk production, for a given pen, as used in ration formulation, kg/animal

Parameters and Equations

- `Pen.reduce_milk_production()`
 - **Called by** `Pen.use_user_defined_ration()` when `NutritionEvaluator` determines that the ration is inadequate to support milk production
 - Attempts to reduce the milk production of all animals in the pen; returns False if all animals have already reached their maximum
- `Animal.reduce_milk_production()`
 - **Called by** `Pen.reduce_milk_production` for each animal in the pen
 - Attempts to reduce milk production of an animal in a pen; returns True if reduction was successful and False otherwise. When reduction is successful, increments `milk_production_reduction` by the value



Currently, all cows in a pen will have the same milk reduction amount. This function allows future flexibility if desired to have individual reductions rather than applying the average..

6. *Starting and Ending Lactation*

Starting Lactation [Animal.transition_heiferIII_to_cow]: When a heifer reaches the point where her days pregnant is equal to gestation length, she is transitioned to a lactating cow and configuration data for a newborn calf is returned.

[Reproduction.cow_give_birth] and **[Animal.daily_reproduction_update]** When a cow's days pregnant is equal to gestation length, she gives birth and a newborn calf is born. Her parity is incremented plus one, her days in milk is reset to 1, and she is assigned new Wood's curve parameters for the new lactation.

For more information, refer to the [Reproduction](#) section of this document.

Ending Lactation [MilkProduction.perform_daily_milking_update]: When a cow's days pregnant is equal to her dry off days of pregnancy, her milk production and days in milk are set to 0 and remain there until completing her gestation. Cows with days_in_milk = 0 are considered not to be milking and do not have any milk production values calculated.

For more details on milk changes and reductions, go to the Milk Production and Reduction portion of this section. .

D. Lactation Curve: Parameters and Definitions

Introduction The Lactation Curve Parameter Determination module calculates and adjusts Wood's lactation curve parameters (l, m, n) based on farm-specific attributes such as geographic location, management practices, and production levels. These parameters shape the milk production curve for cows across different parities (1st, 2nd, 3rd + lactations).

This module is part of the Animal Module and integrates with the [Milk Production](#) calculations. The outputs influence various downstream models, including nutrient requirements, manure excretion, and greenhouse gas emission calculations.

The primary class in this module is:

LactationCurve: Manages and adjusts Wood's lactation curve parameters.

Required User Inputs

The module retrieves input data from simulation configuration files and the Input Manager. These inputs are organized into groups based on their functional roles. Each input is detailed by its parameter name, definition (including units where applicable), and any default values or constraints. There are three categories on inputs that should be considered in this section:

1. [Wood's Lactation Curve Base Parameters](#)
2. [Adjustment Factors](#)
3. [Farm Specific 305 d Milk Production](#)

Lactation Curve Parameters and Inputs

Table 30: Wood's Lactation Curve Base Parameters – These inputs provide the baseline values and variability for Wood's lactation curve parameters as derived from Li, Rosa, et al. (2022).

Variable	Definition	Units	Default
parameter_l_mean	Baseline value for parameter l of Wood's lactation curve	Unitless	19.9
parameter_m_mean	Baseline value for parameter m of Wood's lactation curve	Unitless	0.247
parameter_n_mean	Baseline value for parameter n of Wood's lactation curve	Unitless	0.003376
parameter_l_std_dev	Standard deviation for parameter l	Unitless	Parity 1: 0.28; Parity 2: 0.54; Parity 3+: 0.51
parameter_m_std_dev	Standard deviation for parameter m	Unitless	Parity 1: 0.0046; Parity 2: 0.0064; Parity 3+: 0.0060
parameter_n_std_dev	Standard deviation for parameter n	Unitless	Parity 1: 3.77e-5; Parity 2: 5.82e-5; Parity 3+: 5.54e-5

Lactation Curve Adjustment Factors

Table 32: Adjustment Factors – These factors adjust the base parameters based on geographical, temporal, management, and parity considerations.

Variable	Definition	Units	Default
region_adjustments	Regional adjustments based on FIPS codes ('fips.code') in the configuration file	Unitless	–
year_adjustments	Yearly adjustments based on calving year, defined by simulation time period ('time.end_date.year')	Unitless	2006–2016
milking_frequency_adjustments	Adjustments based on milking frequency ('cow.times.milked.per.day') in the animal input file	Unitless	2 or 3 (<i>Classified as twice-daily (< 2.5 milkings/day) or thrice-daily (≥ 2.5 milkings/day)</i>)
parity_adjustments	Adjustments to lactation curve based on parity (first, second, third+)	Unitless	1–3

Farm Specific 305-Day Milk Production Inputs

Table 34: Farm Specific 305-Day Milk Production — These inputs help tailor the lactation curve to the farm’s production targets and herd composition.

Variable	Definition	Units	Default
parity_fractions	Fraction of cows by lactation parity	Fraction	Parity 1: 0.346; Parity 2: 0.272; Parity 3+: 0.379
annual_milk_yield	Target annual milk yield to fit the lactation curve	kg	User-defined, default: null
milking_cow_fraction	Fraction of cows that are milking	Fraction	0.8365 (User-defined)



Relevant Outputs

Table 35: Summary of important functions for the reproduction section.

Parameter	Unit	Description
l	Unitless	Farm-specific scaling factor of the lactation curve by parity.
m	Unitless	Farm-specific rate of milk yield increase post-calving by parity.
n	Unitless	Farm-specific rate of milk yield decline after peak lactation by parity.

Methodology

1. Determining Base Lactation Curve Parameters (`set_lactation_parameters`). The Wood's lactation curve parameters are initialized with base values retrieved from `lactation_inputs["parameter_mean_values"]` in `LACTATION_CURVE_ADJUSTMENT_INPUTS.json`.
2. Adjusting parameters (`_calculate_adjusted_wood_parameters`). Wood's lactation curve parameters are adjusted based on temporal, geographical, and management factors (Li, Rosa, et al., 2022)

Each parameter (l , m , n) is adjusted using a series of modifications:

$$l_{adjusted} = l + \sum adjustments$$

[AN.MLK.1]

$$m_{adjusted} = m + scalingfactor_m * \sum adjustments$$

[AN.MLK.2]

$$n_{adjusted} = n + scaling factor_n * \sum adjustments$$



[AN.MLK.3]

Where:

$$scaling\ factor_m = 10^{-2}$$

$$scaling\ factor_n = 10^{-4}$$

Adjustments are applied based on:

- Region (using FIPS codes): `_get_region_adjustments`
 - Year of simulation (adjusted for trends in milk production): `_get_year_adjustments`
 - Milking frequency (twice vs. thrice daily): `_get_milking_frequency_adjustments`
 - Parity (different lactation curves for different parities): `animal_parity`
3. Fitting Parameters to Target Milk Yield (`_adjust_lactation_curve_to_milk_yield`). If `annual_milk_yield` is provided, the `l` parameter is optimized to match the expected 305-day milk yield:
- Estimate Herd Average 305-day Milk Yield:

$$Milk\ Yield_{305} = \frac{Annual\ Milk\ Yield}{Number\ of\ Milking\ Cows} * \frac{305}{Days\ per\ Year}$$

[AN.MLK.4]

- Estimate Parity Specific 305-day Milk Yield: (`_estimate_305_day_milk_yield_by_parity`)

$$M305_{Herd} = M305_{P1} * P1\% + M305_{P2} * P2\% + M305_{P3} * P3\%$$

[AN.MLK.5]

$$M305_{P2} = M305_{P1} * 118\%$$

[AN.MLK.6]

$$M305_{P3+} = M305_{P1} * 125\%$$

[AN.MLK.7]

118% and 125% are assumptions made based on the work by Li, Rosa, et al. (2022).

- Fit `l` Parameter Using Optimization: (`_fit_wood_l_param_to_milk_yield`). The error function is defined as: (`_calculate_305_day_milk_yield_error`)



$$Error = Predicted MilkYield305 - Target MilkYield305$$

[AN.MLK.8]

Using Scipy's minimize function, the l parameter is iteratively adjusted to minimize this error.

4. Final Lactation Curve Parameter Mapping. Once adjustments are made, parameters are stored in:

```
_parity_to_parameter_mapping = 1: "l": ..., "m": ..., "n": ...,
2: "l": ..., "m": ..., "n": ...,
3: "l": ..., "m": ..., "n": ...
```

Equations below are not in the LactationCurve class, but they are useful to understanding the bigger picture of how lactation curve parameter estimates are generated in RuFaS. Please see the [Milk Production](#) section for more details on these equations (Wood, 1967).

Wood's Lactation Curve

$$Y_t = l * t^m * e^{-nt}$$

[AN.MLK.9]

Where:

- Y(t): Daily milk yield.
- t: Time in days.
- l, m, n: Parameters adjusted for farm-specific conditions.

305-Day Milk Yield

$$\int_l^{305} l t^m e^{-nt} dt$$

[AN.MLK.10]

E. Culling

Introduction This section describes the processes of animals leaving the herd in RuFaS. Reasons for leaving the herd are death or sale.

Sale may occur of:



- Calves (all male calves; female calves based on `keep_female_calf_rate`)
- Heifer II's (if the heifer does not have a successful conception or pregnancy and age (days born) exceeds `heifer_repro_cull_time`)
- Heifer III's (if cow herd does not have room for additional heifers)
- Cows (through two possible mechanisms):
 - Disease-related culling, determined by a statistical distribution of occurrence, and
 - Production-related culling, which occurs when a cow has been labeled “Do Not Breed” due to reproductive failure past a designated number of days in milk and then her milk production drops below a designated threshold)

Deaths that may occur:

- Cows, determined by a statistical distribution of occurrence
- Stillborn calves
 - *Note:* stillborn calves are currently lumped together with calves sold at birth and reported as “calves sold”

Information about death and sale events are tracked and aggregated daily, including the day of death or sale, body weight, cull reason, days in milk, and parity.

Required User Inputs

Table 36: The following inputs are used to determine the probability and timing of cow death and sales events.

Variable	Definition	Default	Min	Max
parity_death_prob	Death probability by parity group (1st, 2nd, 3rd, 4th+ lactation)	-	0	1
parity_cull_prob	Cull probability by parity group (1st, 2nd, 3rd, 4th+ lactation)	-	0	1
parity_day_prob*	CDF values for likelihood of death over time; used with 'cull_day_count'	-	0	1
cull_day_count*	Defines time segments (days in milk) for the CDF curve	-	0	-
probability*	Conditional probability of culling due to each of six reasons; sums to 1	-	0	1
cull_day_prob*	CDF values for likelihood of [reason] culling over time	-	0	1
do_not_breed_time	Days after calving a cow will be culled if not pregnant	0	185	-
cull_milk_production	Milk threshold (kg/d) below which cows are culled	0	30	-
herd_num	Target number of dry and lactating cows on the farm	100	-	6
heifer_repro_cull_time	Max age (days) before a heifer is culled for not getting pregnant	500	0	-
keep_female_calf_rate	Fraction of female calves kept and raised on-farm	1	0	1
male_calf_rate_sexed_semen	Male calf rate when using sexed semen	0.1	0	1
male_calf_rate_conventional_semen	Male calf rate when using conventional semen	0.53	0	1
still_birth_rate	Stillbirth rate	0.065	0	1

Note: Variables marked with * are commonly left at default values. Adjusting them changes the timing of cow exit from the herd but does not affect the total number of exits.



Relevant Outputs

Information about death and sales events are tracked and aggregated daily by HerdManager, including the animal type, day of death or sale, body weight, cull reason, days in milk, and parity.

All of the following are reported by AnimalModuleReporter:

- **Report_life_cycle_manager_data** - is reported on a daily basis for the numbers of calves, HeiferII, HeiferIII, and Cow sold (calves born deceased tracked as newborns sold). It also reports HeiferIII purchased and documents other cow culls (i.e. sold and died)
Culling data is also reported here documenting the average age of cows and HeiferII culled and the reason (e.g. death, poor production, lameness, injury, mastitis, disease, udder, or unknown/other).
- **Report_sold_animal_information** - this data is reported at the end of a simulation. It creates a list of all animals sold and reports their ID number, type, body weight, day sold (simulation day), cull reason, days in milk, and parity.
- **Report_sold_animal_information_sort_by_sell_day** - also reported at the end of a simulation to document the overall number and weight of animals sold on that simulation day. It reports zero's for days on which no animal of that type was sold. This enables using report filters to slice and summarize sales statistics by animal type for a specified time period (e.g., number and body weight of cows sold over the last 365 days of the simulation).

Methodology *Cow Deaths*

At the beginning of each lactation, each cow “rolls the dice” to determine if +/- when she will die in the current lactation.

Probability of death associated with each lactation (1, 2, 3, 4+) is specified at the beginning of the simulation as a user input. At the beginning of each lactation for each cow, it is determined whether she will die during that lactation with a random draw compared with the appropriate probability. Then, for individuals assigned to die during that lactation, an empirical Cumulative Distributive Function (CDF) is used to determine on which day the death will occur based on the probability of death occurring during each interval of the lactation.

The values for the CDF determining probability of death by day of lactation are located in the animal input json file (death_day_prob; See Table 4).

User Inputs and Constants

- Parity death probability: “parity_death_prob”
 - Probability of death during the current lactation for cows beginning a new lactation, grouped into lactation groups 1, 2, 3, and 4+
- Death day probability: “death_day_prob”
 - Values to construct the CDF to determine future death day, given that death is going to occur (default values in the table below)
 - Values in the array correspond to the days in cull_day_count (another user input; default values are the “days” in the table below)

Parameters and Equations

Determination of death date (once already determined that death will occur this lactation):



Death day of Cow X is as follows

$$X = x_{i-1} + a_i * (R - c_{i-1})$$

[AN.ANM.1]

Where:

R = random float number between 0-1

c_{i-1}, c_i (the upper and lower values around R in the death day prob CDF),

x_{i-1}, x_i (days, correspond to c_{i-1}, c_i)

Empirical CDF: $a_i = \frac{x_{i-1}, x_i}{c_{i-1}, c_i}$

- Animal.determine_future_death_date()
 - **Called by** Animal.daily_routines() at the beginning of each new lactation (on the day a new calf is born).
 - **Returns** calculated future death date in simulation days or sys.maxsize (effectively no death date).
 - Logic: Draw random value between 0-1 and compare it to probability of death in that lactation group (parity_death_probability).
 - * If drawn value $\leq$ parity_death_probability, then go on to select date that death will occur - draws another random value between 0-1
 - * Compares random draw to the CDF defined by death_day_prob array to determine number of days into the future at which death will occur - adds that number of days to current days born to determine future death date .
 - * If drawn value $>$ parity_death_probability, then future death date is assigned as sys.maxsize (a really really big number – effectively, she will never reach her assigned future death date).
- Animal.animal_life_stage_update()
 - **Called by** HerdManager.daily_routines() → Animal.daily_routines()
 - Updates the animal status to DEAD if days born == future death date

Cow Sales

1. Disease Related

At the beginning of each lactation, each cow “rolls the dice” to determine if +/- when and for what attributed reason she will be sold for disease-related reasons (i.e., involuntary, not directly for production/reproduction reasons) in the current lactation.

Probability of disease sales associated with each lactation (1, 2, 3, 4+) is specified at the beginning of the simulation as a user input. At the beginning of each lactation for each cow, it is determined whether she will be sold for disease during that lactation with a random draw compared with the appropriate



probability. Then, for individuals assigned to be sold during that lactation, another random draw determines the reason for that sale from one of six possibilities (See Table 4). Finally, an empirical CDF is used to determine the day of sale based on the probability of sale occurring during each interval of the lactation for that particular sale reason.

User Inputs and Constants

- Parity cull probability: “parity_cull_prob”
 - Probability of sale for disease during the current lactation for cows beginning a new lactation, grouped into lactation groups 1, 2, 3, and 4+
- PFor each of the six reasons:
 - “Probability” - Conditional probability that a sold animal is removed due to each group of issues. This probability is used to determine the reason for sale after it has been decided that the animal will be sold during the current lactation. The sum of probabilities for the six culling reasons equals 1. See Table 2 below for default values used to construct the empirical PDF.
 - “Cull_day_prob”: Values to construct the CDF to determine future sale day, given that disease-related sale is going to occur. Values in the array correspond to the days in cull_day_count (another user input; default values are the “days”; default values in Table 4)

Parameters and Equations

Determination of sale date (once already determined that death will occur this lactation):

We compute the sale day of Cow X as follows

$$X = x_{i-1} + a_i * (R - c_{i-1})$$

[AN.ANM.2]

Where:

R = random float number between 0-1

c_{i-1}, c_i (the upper and lower values around R in the sale day prob CDF),

x_{i-1}, x_i (days, correspond to c_{i-1}, c_i)

Empirical CDF: $a_i = \frac{x_{i-1}, x_i}{c_{i-1}, c_i}$

- Animal.determine_future_cull_date()
 - **Called by** by Animal.daily_routines() at the beginning of each new lactation (on the day a new calf is born)
 - **Returns** a tuple with future cull (sale) date in simulation days and reason for culling
 - Logic: Draw random value between 0-1 and compare it to probability of death in that lactation group (parity_death_probability)
 - * If drawn value $\leq$ parity_death_probability, then go on to select date that death will occur - draws another random value between 0-1
 - * Compares random draw to the CDF defined by death_day_prob array to determine number of days into the future at which death will occur - adds that number of days to current days born to determine future death date



- * If drawn value > parity_death_probability, then future death date is assigned as sys.maxsize (a really really big number – effectively, she will never reach her assigned future death date)
- Animal.life_stage.update()
 - **Called by** HerdManager.daily_routines() → Animal.daily_routines()
 - Updates the animal status to SOLD if days born == future cull date

2. Production Related

If a cow is not pregnant when she reaches the days in milk cutoff specified by the user, she will be marked as Do Not Breed and no longer participate in reproductive protocols. When her milk production drops below the production cutoff specified by the user, she is sold for the reason of low production. Note that the only cows who are sold for “low production” are those flagged as DNB for reproductive failure; no low-producing cow who is pregnant or has fewer DIM than the cutoff will be sold for low production.

User Inputs and Constants

- Do_not_breed_time: The length of the breeding period after parturition for cows after which reproduction protocols stop if they fail to get pregnant
- Cull_milk_production: Milk production threshold at which 'do not breed' cows are culled if they fall below (in kg/d)

Parameters and Equations

- Reproduction.check_do_not_breed_flag()
 - Checks if a cow should be marked as do-not-breed if not pregnant beyond the user-defined breeding window
 - Called by cow reproduction update() in Reproduction.py
 - If not pregnant and DIM > do_not_breed_time and not already do_not_breed:
 - * Updates reproduction_data_stream with do not breed event
 - * Updates self.do_not_breed to true
- Animal.life_stage.update()
 - Updates the life stage of an animal based on its type and current simulation time
 - Called by Animal.daily_routines()
 - Includes the following logic:
 - * If a cow is do_not_breed and daily milk produced is less than user-defined cull_milk_production:
 - Sold_at_day = simulation_day
 - Cull_reason = low_prod_cull
 - Animal_status = sold



Table 37: Cumulative distribution function (CDF) values associated with the conditional probability that a sold animal is removed due to each group of issues and/or the conditional probability of death over time. Corresponds to the “probability” value for each cull reason in the animal input json file. The ‘cull_day_count’ array defines the segments of the CDF

Variable	0	5	15	45	90	135	180	225	270	330	380	430	480	530
Cumulative Probability of	0	0.18	0.32	0.42	0.48	0.54	0.60	0.65	0.70	0.77	0.83	0.89	0.95	1
Death														
Mastitis CDF	0	0.06	0.12	0.19	0.30	0.43	0.56	0.68	0.78	0.85	0.90	0.94	0.97	1
Feet & Leg	0	0.03	0.08	0.16	0.25	0.36	0.48	0.59	0.69	0.78	0.85	0.90	0.95	1
CDF														
Injury CDF	0	0.08	0.18	0.28	0.38	0.47	0.56	0.64	0.71	0.78	0.85	0.90	0.95	1
Disease CDF	0	0.04	0.12	0.24	0.34	0.42	0.50	0.57	0.64	0.72	0.81	0.89	0.95	1
Udder CDF	0	0.12	0.24	0.33	0.41	0.48	0.55	0.62	0.68	0.76	0.82	0.89	0.95	1
Unknown	0	0.05	0.11	0.18	0.27	0.37	0.45	0.54	0.62	0.70	0.77	0.84	0.95	1
CDF														



If a cow is not pregnant when she reaches the days in milk cutoff specified by the user, she will be marked as Do Not Breed and no longer participate in reproductive protocols. When her milk production drops below the production cutoff specified by the user, she is sold for the reason of low production. Note that the only cows who are sold for “low production” are those flagged as DNB for reproductive failure; no low-producing cow who is pregnant or has fewer DIM than the cutoff will be sold for low production.

V. Methane Emission and Manure Excretion

A. Manure Excretion

Introduction The Manure Excretion Calculator module computes manure excretion for different animal categories, including calves, growing heifers, lactating cows, and dry cows. It calculates the quantities of total mass and nutrient composition in manure. The implementation is designed for flexibility, enabling integration with other RuFaS modules for modeling farm management scenarios. This module includes:

- *Classes:* ManureExcretionCalculator
- *Key Methods:*
 - calculate_calf_manure
 - calculate_heifer_manure
 - calculate_cow_manure
 - _calculate_lactating_cow_manure
 - _calculate_dry_cow_manure
 - _calculate_phosphorus_excretion_values

This module integrates with the process_digestion method in the DigestiveSystem class to calculate manure excretion on a daily and per-animal basis. It plays a critical role in linking animal-level digestion processes with downstream environmental and nutrient management models.

The module takes inputs that include general animal properties (e.g., body weight, nutrient intake, and nutrient compositions), milk production characteristics (e.g., fat and protein content), specific nutrient properties (e.g., urine phosphorus requirements), and animal categories (e.g., CALF, HEIFER, COW). The outputs from this module are detailed manure excretion metrics, such as total manure mass, nutrient content, and volatile solids. These outputs are used by the Manure Module for downstream calculations, including manure methane (CH₄) emissions and nutrient balance tracking.

Required User Inputs

The Manure Excretion Calculator module computes manure excretion metrics for various animal categories. Inputs are drawn from pen ration files and animal module constants. They are organized into the following groups:

1. [General Animal Inputs](#)
2. [Cow Specific Inputs](#)
3. [Phosphorus Specific Inputs](#)

**General Inputs** Manure Excretion

Table 38: These inputs define basic animal and diet properties used to calculate nutrient intake and manure production.

Variable	Definition	Units
body_weight	Body weight of the current animal	kg
dry_matter_intake	Dry matter intake from pen ration	kg
dry_matter_concentration	Dry matter concentration of the diet	%
crude_protein_conc	Crude protein concentration in the diet	%
potassium_concentration	Potassium concentration in the diet	%
ash_diet_content	Ash content of the diet	%
neutral_detergent_fiber_concentration	Neutral detergent fiber concentration	%
acid_detergent_fiber_concentrations	Acid detergent fiber concentration	%
nutrient_concentrations	Nutrient concentrations (e.g., CP, NDF, ADF, ash)	%

Cow Specific Inputs – Manure Excretion

Table 39: These inputs are used when modeling cows and are particularly relevant for lactating animals.

Variable	Definition	Units	Default	Note
is_lactating	Indicates the cow's lactating status	Boolean	-	-
days_in_milk	Days post-parturition; for lactating cows only	days	-	Intermediate output
milk_protein	Milk protein content; for lactating cows only	%	3.2	-
daily_milk_production	Daily milk production; for lactating cows only	kg	-	-
body_weight	Body weight of the cow	kg	-	-
milk_fat	Milk fat content	%	3.5	-
metabolizable_energy_intake	Metabolizable energy intake per kg DM	Mcal/kg DM	-	-

Phosphorus Excretion Inputs – Manure Excretion



Table 40: These inputs define phosphorus-related outputs relevant to lactating cows.

Variable	Definition	Units
fecal_phosphorus	Amount of fecal phosphorus excreted by the current animal; Intermediate model output	g
urine_phosphorus_required	Phosphorus required for urine production; Intermediate model output	g
total_manure_excreted	Total manure excreted by the animal; Intermediate model output	kg

Expected Outputs

The outputs from the Manure Excretion Calculator are organized into two main categories: manure excretion values and manure nutrient composition. These outputs are consistently generated across animal categories.

Table 41: Expected outputs from this section.

Category	Parameter	Definition	Units
Manure Excretion	Urea	Concentration of urea in the manure	g/L
	Urine	Amount of urine excreted	kg
	Manure Mass	Total mass of manure excreted	kg
	Total Solids	Dry matter content of the manure	kg
	Degradable Volatile Solids	Portion of volatile solids that are degradable	kg
	Non-degradable Volatile Solids	Portion of volatile solids that are non-degradable	kg
Manure Nutrient Composition	Manure Total Ammoniacal Nitrogen	Total ammoniacal nitrogen excreted in the manure	kg
	Urine Nitrogen	Amount of nitrogen excreted via urine	kg
	Manure Nitrogen	Total nitrogen excreted in manure	kg
	Inorganic Phosphorus Fraction	Fraction of manure phosphorus that is inorganic	-
	Organic Phosphorus Fraction	Fraction of manure phosphorus that is organic	-
	Non-water Inorganic Phosphorus Fraction	Portion of inorganic phosphorus not water-extractable	-

Continued on next page



Table continued from previous page

Category	Parameter	Definition	Units
	Non-water Organic Phosphorus Fraction	Portion of organic phosphorus not water-extractable	-
	Total Phosphorus Excreted	Total phosphorus excreted by the animal	g
	Phosphorus Fraction	Overall fraction of phosphorus in the manure	-
	Potassium	Amount of potassium excreted in the manure	g

Methodology

• Calf Manure Excretion (calculate_calf_manure)

Manure excretions Manure: amount of feces and urine excreted daily by a calf (kg/d)

$$total_manure_excreted = 3.45 * dry_matter_intake$$

[AN.EXC.1]

Urine: amount of urine excreted in kg. This is an assumption.

$$urine = 2.0$$

[AN.EXC.2]

Manure total solids: amount of dry material excreted by the calf (kg/d).

$$total_solids = 0.393 * dry_matter_intake$$

[AN.EXC.3]

Total volatile solids (kg/d)

$$total_volatile_solids = 0.0023 * body_weight$$



[AN.EXC.4]

Degradable volatile solids excretion (kg/d)

$$degradable_volatile_solids = 0.9 * total_volatile_solids$$

[AN.EXC.5]

Non-degradable volatile solids excretion (kg/d)

$$non_degradable_volatile_solids = total_volatile_solids - degradable_volatile_solids$$

[AN.EXC.6]

Manure nutrient composition Manure N excretion (kg/d):

$$manure_nitrogen = \frac{(112.55 * dry_matter_intake * \frac{crudeprotein.concentration}{100})}{1000}$$

[AN.EXC.7]

Urine N excretion (kg/d)

$$urine_nitrogen = 0.45 * manure_nitrogen$$

[AN.EXC.8]

Manure total ammoniacal nitrogen (kg/d)

$$manure_total_ammoniacal_nitrogen = urine_nitrogen$$

[AN.EXC.9]

- **Heifer Manure Excretion (calculate_heifer_manure)**



Manure excretions Urine excretion (kg/d)

$$urine = 9.0$$

[AN.EXC.10]

Total manure excretion (kg/d)

$$total_manure_excreted = 4.158 * dry_matter_intake - 0.0246 * body_weight$$

[AN.EXC.11]

Because the equations that predict total manure excreted and total solids excreted rely on different inputs, in some extreme cases, the total mass of manure excreted can be predicted to be very close to the predicted total solids excreted which is unrealistically dry. For this reason, we set a minimum manure dry matter content of 15% for heifers and 12% for dry cows. If the mass predicted by equation [AN.EXC.11] is not greater than (total solids/minimum dry matter) then,

$$total_manure_excreted = \frac{total_solids}{dry_matter_manure}$$

[AN.EXC.#]

Total solids excretion (kg/d) - this is the same equation used for dry cows.

$$total_solids = 0.178 * dry_matter_intake + 2.733$$

[AN.EXC.12]

Total volatile solids excretion (kg/d)

$$total_volatile_solids = 0.0073 * body_weight$$

[AN.EXC.13]

Degradable volatile solids excretion (kg/d)

$$degradable_volatile_solids = 0.9 * total_volatile_solids$$



[AN.EXC.14]

Non-degradable volatile solids excretion (kg/d)

$$non_degradable_volatile_solids = total_volatile_solids - degradable_volatile_solids$$

[AN.EXC.6]

Manure nutrient composition Manure N excretion (kg/d)

$$manure_nitrogen = (15.1 + 0.83 * \frac{(dry_matter_intake * 1000) * \frac{crude_protein_concentration}{6.25}}{100})$$

[AN.EXC.14]

Fecal N excretion (kg/d)

$$fecal_nitrogen = (0.345 + 0.317 * \frac{(dry_matter_intake * 1000) * \frac{crude_protein_concentration}{6.25}}{100})$$

[AN.EXC.15]

Urine N excretion (kg/d)

$$urine_{nitrogen} = manure_{nitrogen} - fecal_{nitrogen}$$

[AN.EXC.16]

Manure total ammoniacal nitrogen (kg/d)

$$manure_total_ammoniacal_nitrogen = urine_nitrogen$$

[AN.EXC.9]

• **Heifer Manure Excretion (calculate_heifer_manure)**

Manure K excretion (g/d)



$$potassium = 1000 * dry_matter_intake * \frac{potassium_concentration}{100}$$

[AN.EXC.17]

- **Lactating Cow Manure Excretion (.calculate_lactating_cow_manure)**

Manure excretions Fecal water (kg/d): Calculate the amount of fecal water excreted (kg/d)

$$fecal_water = 1.987 * dry_matter_intake + 0.348 * acid_detergent_fiber_concentrations - 0.412 * crude_protein_concentration - 0.074 * dry_matter_concentration - 0.0057 * days_in_milk$$

[AN.EXC.18]

Total solids/Fecal dry matter (kg/d): Calculate the amount of fecal dry matter excreted (kg/d). Fecal dry matter is assumed to be equivalent to total solids.

$$fecal_solids = 0.576 + 0.370 * dry_matter_intake - 0.075 * crude_protein_concentration + 0.059 * acid_detergent_fiber_concentrations$$

[AN.EXC.19]

Urine (kg/d)

$$urine = -7.742 + 0.388 * dry_matter_intake + 0.726 * crude_protein_concentration + 2.066 * milk_protein$$

[AN.EXC.20]

Total manure excretion (kg/d)

$$total_manure_excreted = fecal_water + fecal_solids + urine$$

[AN.EXC.21]

Organic matter intake (kg/d)

$$organic_matter_intake = dry_matter_intake - ash_diet_content$$



[AN.EXC.22]

Degradable volatile solids (kg/d)

$$\text{degradable_volatile_solids} = -1.017 + 0.364 * \text{organic_matter_intake} + 0.029 * \text{neutral_detergent_fiber_concentration} - 0.023 * \text{crude_protein_concentration}$$

[AN.EXC.23]

Total volatile solids excreted by a lactating cow (kg/d)

$$\text{total_volatile_solids} = -1.201 + 0.402 * \text{organic_matter_intake} + 0.036 * \text{neutral_detergent_fiber_concentration} - 0.024 * \text{crude_protein_concentration}$$

[AN.EXC.24]

Non-degradable volatile solids excretion (kg/d)

$$\text{non_degradable_volatile_solids} = \text{total_volatile_solids} - \text{degradable_volatile_solids}$$

[AN.EXC.6]

Manure nutrient composition Total manure nitrogen (kg/d)

$$\text{manure_nitrogen} = (20.3 + 0.654 * \frac{(\text{dry_matter_intake} * 1000) * \frac{\text{crude_protein_concentration}}{\frac{6.25}{100}}}{1000})$$

[AN.EXC.25]

Fecal nitrogen (kg/d)

$$\text{fecal_nitrogen} = \frac{(10.1 * \text{dry_matter_intake} - 18.5)}{1000}$$

[AN.EXC.26]

Urine N excretion (kg/d)

$$\text{urine}_{\text{nitrogen}} = \text{manure}_{\text{nitrogen}} - \text{fecal}_{\text{nitrogen}}$$



[AN.EXC.16]

Manure total ammoniacal nitrogen (kg/d)

$$\text{manure_total_ammoniacal_nitrogen} = \text{urine_nitrogen}$$

[AN.EXC.9]

Manure K excretion (g/d)

$$\text{Potassium} = 7.21 * \text{dry_matter_intake} + 15944 * \frac{\text{potassium_concentration}}{100} - 164.5$$

[AN.EXC.27]

- **Dry Cow Manure Excretion (_calculate_dry_cow_manure)**

Manure excretions Urine excretion (kg/d): Amount of urine excreted in kg with assumption that 1.038 kg urine is approximately 1 L. Due to lack of information, average excretion rate from dry cows is assumed.

$$\text{urine} = 15.4$$

[AN.EXC.28]

Total manure excretion (kg/d): Amount of feces and urine excreted daily by dry cows (kg/d).

$$\text{total_manure_excreted} = 0.00711 * \text{body_weight} + 0.324 * \text{crude_protein_concentration} + 0.259 * \text{neutral_detergent_fiber_concentration} + 8.05$$

[AN.EXC.29]

Because the equations that predict total manure excreted and total solids excreted rely on different inputs, in some extreme cases, the total mass of manure excreted can be predicted to be very close to the predicted total solids excreted which is unrealistically dry. For this reason, we set a minimum manure dry matter content of 15% for heifers and 12% for dry cows. If the mass predicted by equation AN.EXC.11 is not greater than (total solids/minimum dry matter) then,

$$\text{total_manure_excreted} = \frac{\text{total_solids}}{\text{dry_matter_manure}}$$

Total solids excretion (kg/d) - this is the same equation used for dry cows.

$$\text{total_solids} = 0.178 * \text{dry_matter_intake} + 2.733$$



[AN.EXC.12]

Organic matter intake (kg/d)

$$\text{organic_matter_intake} = \text{dry_matter_intake} - \text{ash_diet_content}$$

[AN.EXC.22]

Degradable volatile solids (kg/d)

$$\text{degradable_volatile_solids} = -1.017 + 0.364 * \text{organic_matter_intake} + 0.029 * \text{neutral_detergent_fiber_concentration} - 0.023 * \text{crude_protein_concentration}$$

[AN.EXC.23]

Total volatile solids excreted by a lactating cow (kg/d)

$$\text{total_volatile_solids} = -1.201 + 0.402 * \text{organic_matter_intake} + 0.036 * \text{neutral_detergent_fiber_concentration} - 0.024 * \text{crude_protein_concentration}$$

[AN.EXC.24]

Non-degradable volatile solids excretion (kg/d)

$$\text{non_degradable_volatile_solids} = \text{total_volatile_solids} - \text{degradable_volatile_solids}$$

[AN.EXC.6]

- **Manure Phosphorus Excretion (_calculate_phosphorus_excretion_manure)**

Total phosphorus fraction of feces

$$\text{manure_phosphorus_fraction} = \frac{\text{fecal_phosphorus} + \text{urine_phosphorus_required}}{\text{total_manure_excreted} * 1000 \text{g/kg}}$$

[AN.EXC.30]

Inorganic phosphorus fraction



$$\text{inorganic_phosphorus_fraction} = 0.50 * \text{manure_phosphorus_fraction}$$

[AN.EXC.31]

Organic phosphorus fraction

$$\text{organic_phosphorus_fraction} = 0.05 * \text{manure_phosphorus_fraction}$$

[AN.EXC.32]

Milk phosphorus

$$\text{phosphorus_in_milk} = 0.0009 * \text{daily_milk_production} * 1000\text{g/kg}$$

[AN.EXC.33]

Manure P excreted by a cow (g/d)

$$\text{manure_phosphorus_excreted} = \text{fecal_phosphorus} + \text{urine_phosphorus_required}$$

[AN.EXC.34]

Total P excreted by a cow (g/d)

$$\begin{aligned} \text{total_phosphorus_excreted} = \\ \text{phosphorus_in_milk} + \text{fecal_phosphorus} + \text{urine_phosphorus_required} \end{aligned}$$

[AN.EXC.35]

Important Note

All references may be found at the end of this document or are based on expert communication with V. Sempali from 2023 to 2024.



B. Enteric Methane Emission

Introduction The Enteric Methane Calculator module calculates daily enteric methane emissions for calves, heifers, lactating cows, and dry cows. Methane emissions are estimated based on animal characteristics, dietary intake, and a variety of animal category specific methane models. The IPCC, 2006 model is available for all animal classes except calves. The equation from Mills et al., 2003 is also available for dry and lactating cows and the Niu, E. Kebreab, Hristov, et al., 2018 equation is available for lactating cows. Finally, a separate equation published by Pattanaik et al., 2003 is used for calves. Simulation of enteric methane mitigation via mitigation supplements 3-NOP or monensin are also available for lactating cows. The module integrates with the DigestiveSystem class and supports methane mitigation strategies using feed additives.

This module includes:

- **Classes:** EntericMethaneCalculator
- **Key Methods:**
 - calculate_calf_methane
 - calculate_heifer_methane
 - calculate_cow_methane
 - The lactating and dry cow methods are called within the calculate_cow_methane method
 - .calculate_lactating_cow_manure
 - .calculate_dry_cow_manure

The calculated enteric methane emissions are directly output to the Output Manager to generate the final reports.

Required User Inputs General Inputs

Table 42: These inputs define basic animal and diet properties used to calculate nutrient intake and manure production.

Variable	Definition	Units
dry_matter_intake	Dry matter intake from pen ration	kg
nutrient_concentrations	Nutrient concentrations (e.g., CP, NDF, ADF, ash)	%

Cow Specific Inputs

Table 43: These inputs are used when modeling cows and are particularly relevant for lactating animals.

Variable Name	Definition	Units	Default	Max
body_weight	Body weight of the current animal	kg	-	-
is_lactating	Indicates the cow's lactating status	Boolean	-	-

Continued on next page



Table continued from previous page

Variable Name	Definition	Units	Default	Max
milk_fat	Milk fat content	%	3.5	-
metabolizable_energy_intake	Metabolizable energy intake per kg DM	Mcal/kg DM	-	-

Methane Model-Specific and Mitigation Inputs

Table 44: Inputs specific to the methane emissions model and selected mitigation strategies.

Variable Name	Definition	Default and Range
methane_model	Selected methane model for emission estimation	"Mutian", "Mills", "IPCC"
methane_mitigation_method	Selected feed additive for methane mitigation	"3-NOP", "Monensin", "Essential Oils", "Seaweed"
methane_mitigation_additive_amount	Amount of feed additive used for methane mitigation (mg/kg DMI)	3-NOP (70; 40–100), Monensin (24; 20–36), Essential Oils (0; 0–100), Seaweed (0; 0–100)

Expected Outputs

The module predicts the daily enteric methane emissions for individual animals:

Table 45: Expected outputs from this section.

Category	Parameter	Definition	Units
Methane Emissions	Daily enteric methane emissions	Estimated methane production per animal per day	g CH ₄ /head/day

Methodology

• Required Diet Composition Inputs

Based on dry matter intake and gross energy concentration from nutrient composition:

$$\text{soluble_residue} = 100 - \text{ash_concentration} - \text{neutral_detergent_fiber_concentration} - \text{crude_protein_concentration} - \text{ethyl_ester_concentration}$$

[AN.MET.1]

$$\text{gross_energy_concentration} = 0.263 * \text{crude_protein_concentration} + 0.522 * \text{ethyl_ester_concentration} + 0.198 * \text{neutral_detergent_fiber_concentration} + 0.160 * \text{soluble_residue}$$



[AN.MET.2]

$$\text{starch_to_acid_detergent_fiber_concentration_ratio} = -0.0011 * \frac{\text{starch_concentration}}{\text{acid_detergent_fiber_concentrations}}$$

[AN.MET.3]

- **Calf Methane Emissions (calculate_calf_methane)** Methane emissions are calculated using body weight. The selected methane model for other animal categories does not apply here. Pattanaik et al., 2003 reported no difference in calf methane production in crossbred calves. Methane emissions by calves are estimated from mean values observed in that study.

$$\text{methane_emission} = \frac{0.013 * \text{body_weight}^{0.75} * 4.184 \text{ MJ/Mcal}}{0.05565}$$

[AN.MET.4]

- **Heifer Methane Emissions (calculate_heifer_methane)**

Methane Emission

$$\text{methane_emission} = \frac{(0.065 * \text{gross_energy_concentration} * \text{dry_matter_intake})}{0.05565}$$

[AN.MET.5]

- **Lactating Cow Methane Emissions (calculate_cow_methane)**

For lactating cows, the Mutian, Mills, or IPCC model is applied.

Mutian Model

$$\text{methane_emission} = -126 + 11.3 * \text{dry_matter_intake} + 2.30 * \text{neutral_detergent_fiber_concentration} + 28.8 * \text{milk_fat} + 0.148 * \text{body_weight}$$

[AN.MET.6]

Mills Model Mills et al., 2003 parameterized Mitscherlich equations for a nonlinear model of methane emissions. The CNCPS and IFSM models use the Mitscherlich 3 equation.

$$\text{methane_emission} = \frac{45.98 * (1 - \exp(-[(\text{starch_to_acid_detergent_fiber_concentration_ratio} + 0.0045) * \text{metabolizable_energy_intake} * 4.184]))}{0.05565}$$

[AN.MET.7]

**IPCC Model**

$$\text{methane_emission} = \frac{(0.065 * \text{gross_energy_concentration} * \text{dry_matter_intake})}{0.05565}$$

[AN.MET.5]

- **Dry Cow Methane Emissions (_calculate_dry_cow_manure)**

Mills Model Mills et al., 2003 parameterized Mitscherlich equations for a nonlinear model of methane emissions. The CNCPS and IFSM models use the Mitscherlich 3 equation.

$$\text{methane_emission} = \frac{45.98 * (1 - \exp(-[(\text{starch_to_acid_detergent_fiber_concentration_ratio}) + 0.0045] * \text{metabolizable_energy_intake} * 4.184))}{0.05565}$$

[AN.MET.7]**IPCC Model**

$$\text{methane_emission} = \frac{(0.065 * \text{gross_energy_concentration} * \text{dry_matter_intake})}{0.05565}$$

[AN.MET.5]

- **Methane Mitigation**

- Methane mitigation strategies are applied exclusively to lactating and dry cows. However, caution is advised when interpreting the outcomes for dry cows, as the equations used were developed based on datasets from lactating cows.
- Currently, the implemented methane mitigation feed additives include 3-NOP and monensin, with essential oils and seaweed planned for future integration.
- The detailed methodology for methane mitigation calculations will be provided in a separate document titled Methane Mitigation Calculator.

C. Enteric Methane Mitigation

Introduction The Methane Mitigation Calculator module evaluates the reduction in enteric methane yield due to the inclusion of specific feed additives in animal diets. This module calculates methane mitigation effects on a per-animal basis by considering dietary composition, feed additive dosage, and the selected mitigation strategy. Currently, the module supports the following mitigation strategies:

- 3-NOP (3-Nitrooxypropanol)
- Monensin

Future updates will include additional strategies such as essential oils and seaweed. The mitigation effects are first calculated as a percentage reduction in methane yield (methane production / DMI). Then this module integrates with the Enteric Methane Calculator to adjust the initially predicted enteric methane.

Required User Inputs



The Methane Mitigation Calculator module evaluates the reduction in enteric methane yield based on dietary composition and the use of specific feed additives. The inputs are organized into the following groups:

Dietary Composition

Table 46: General dietary fiber and fat concentration inputs.

Variable Name	Definition	Default and Range
neutral_detergent_fiber_concentration	Neutral detergent fiber concentration, % DM	-
ethyl_ester_concentration	Ether extract (crude fat) concentration, % DM	-
starch_concentration	Starch concentration in the diet, % DM	-

Table 47: Feed-based methane mitigation strategies.

Variable Name	Definition	Defaults and Units	Min	Max
methane_mitigation_method	Selected feed additive for methane mitigation	“3-NOP”, “Monensin”	-	-

Table 48: Dosage of methane mitigation additives.

Variable Name	Definition	Default and Range
methane_mitigation_additive_amount	Dosage per kg of DMI	3-NOP: 70 (40–100); Monensin: 24 (20–36) mg/kg DMI

Table 49: Expected outputs from this section.

Parameter	Definition	Units
Methane Yield Reduction	Change in methane production per unit DMI; negative indicates reduction	%

M Methodology

• 3-NOP Methane Yield Reduction

Reduction is modeled as a function of additive dosage, NDF, EE, and starch concentrations.

$$\text{methane_yield_reduction} = -30.8 - 0.226 * (\text{methane_mitigation_additive_amount} - 70.5) + 0.906 * \text{neutral_detergent_fiber_concentration} - 32.9 + 3.871 * \text{ethyl_ester_concentration} - 4.2 - 0.337 * (\text{starch_concentration} - 21.1)$$



[AN.MET.8]

- **Monensin Methane Yield Reduction** Reduction is based on additive dosage and starch concentration.

$$\text{methane_yield_reduction} = 6.35 - 0.277\text{methane_mitigation_additive_amount} - 0.182 * \text{starch_concentration}$$

[AN.MET.9]

- **Default Case** If no valid mitigation method is selected, the reduction defaults to 0%.
- **Adjust the Initial Predicted Enteric Methane (EntericMethaneCalculator.calculate_cow_methane)**
First, the initially predicted methane yield of the basal diet is calculated

$$\text{methane_yield} = \frac{\text{methane_emission}}{\text{dry_matter_intake}} * 100$$

[AN.MET.10]

Then the adjusted of methane production is calculated based on methane yield reduction (%)

$$\text{adjusted_methane_emission} = \text{methane_yield} * \left(1 + \frac{\text{methane_yield_reduction}}{100}\right) * \text{dry_matter_intake}$$

[AN.MET.11]



VI. Abbreviations

Table 50: Common abbreviations used throughout this document

Abbreviation	Full Term or Phrase
ADF	Acid detergent fiber (% of DM)
ADG	Average daily weight gain (g/d)
Age1stBred	First breeding age (month)
BCS5	Body condition score (1–5 basis)
BW	Body weight (kg)
Ca	Calcium content (% of DM)
CBW	Calf birth weight (kg)
CI	Calving interval (d)
cm	centimeter
Cost	Feed cost (\$/kg of DM)
CP	Crude protein (% of DM)
d	days
DE	Digestible energy (standard value, Mcal/kg)
DIM	Days in milk
DM	Dry matter (% of as-fed basis)
DMI	Dry matter intake
DOP	Days of pregnancy (d)
dRUP	RUP degradability (% of RUP)
EE	Ether extract, in crude fat (% DM)
EndCP	Endogenous CP
EndN	Endogenous N
EQSBW	Equivalent shrunk body weight (kg)
FA	Fatty acids
FIPS	Federal Information Processing Standards
g	grams
GE	Gross energy (standard value, Mcal/kg)
GrUterW	Gravid uterine weight (kg)
Kd	Rumen protein degradation rate (%/h)
kg	kilograms
kJ	kilojoule
L	liters
m	meter
max	maximum
Mcal	Megacalorie
MCP	Microbial Crude Protein
ME	Metabolizable energy (standard value, Mcal/kg)

Continued on next page

*Table continued from previous page*

Abbreviation	Full Term or Phrase
MICP	Microbial CP
min	minimum
mL	milliliters
MN	Microbial N
MP	Metabolizable protein
MTP	Microbial TP
MW	Mature body weight (kg)
MY	Milk yield
N	Nitrogen (% of DM)
NASEM	National Academies of Sciences, Engineering, and Medicine
NDF	Neutral detergent fiber (% of DM)
NEG	Net energy for growth (standard value, Mcal/kg)
NEL	Net energy for lactation (standard value, Mcal/kg)
NEM	Net energy for maintenance (standard value, Mcal/kg)
nNDF	Nondigestible Neutral Detergent Fiber
NPN	Non-protein Nitrogen
NRC	National Research Council (updated in 2021 to NASEM)
OM	Organic matter (% of DM)
P	Phosphorus content (% of DM)
PAF	Processing adjustment factor (not used)
Parity	Number of lactations
PrevTemp	Average daily temperature of last month (°C)
RDP	Ruminally degraded CP
ROM	Residual OM (% of DM)
RUP	Ruminally undegraded CP
s	seconds
TAN	Total ammoniacal N
TDN	Total digestible nutrient (% of DM)
TP	True protein (% of DM)
USDA	United States Department of Agriculture
UterW	Uterine weight (kg)
WSC	Water Soluble Carbohydrates



VII. References

References

- Appuhamy, J.A.D.R.N., L.E. Moraes, C. Wagner-Riddle, D.P. Casper, J. France, et al. (2014). "Development of mathematical models to predict volume and nutrient composition of fresh manure from lactating Holstein cows". In: *Animal Production Science* 54.12.
- Appuhamy, J.A.D.R.N., L.E. Moraes, C. Wagner-Riddle, D.P. Casper, and E. Kebreab (2018). "Predicting manure volatile solid output of lactating dairy cows". In: *Journal of Dairy Science* 101.1, pp. 820–829.
- ASABE (2005). *Manure Production and Characteristics*. St. Joseph, MN: ASABE, pp. 1–32.
- Boadi, D. et al. (2004). "Mitigation strategies to reduce enteric methane emissions from dairy cows: Update review". In: *Canadian Journal of Animal Science* 84.3, pp. 319–335. DOI: [10.4141/A03-109](https://doi.org/10.4141/A03-109).
- Boer, I.J. de et al. (2002). "Prediction of ammonia emission from dairy barns using feed characteristics part 1: Relation between feed characteristics and urinary urea concentration". In: *Journal of Dairy Science* 85.12, pp. 3382–3388. DOI: [10.3168/jds.S0022-0302\(02\)74425-1](https://doi.org/10.3168/jds.S0022-0302(02)74425-1).
- Dairy Cattle Reproduction Council (2018). *Reproductive Management Strategies for Dairy Cows and Heifers*. Accessed: 2023-06-17. URL: <https://www.dccouncil.org/protocols/>.
- De Vries, A. et al. (2006). "Economic comparison of timed artificial insemination and exogenous progesterone as treatments for ovarian cysts." In: *J Dairy Sci* 89.8, pp. 3028–3037.
- Galvão, K. N. et al. (2013). "Economic comparison of reproductive programs for dairy herds using estrus detection, timed artificial insemination, or a combination". In: *J Dairy Sci* 96.4, pp. 2681–2693.
- IPCC (2006). *Guidelines for National Greenhouse Gas Inventories*. <https://www.ipcc-nggip.iges.or.jp/public/2006gl/>. Accessed 2025-04-17.
- Johnson, A.C.B., K.F. Reed, and E. Kebreab (2016). "Short communication: Evaluation of nitrogen excretion equations from cattle". In: *Journal of Dairy Science* 99.9, pp. 7669–7678. DOI: [10.3168/jds.2015-10730](https://doi.org/10.3168/jds.2015-10730).
- Kalantari, Afshin Samia (2015). *Using mathematical modeling techniques for optimized dairy herd management and decision making*. The University of Wisconsin-Madison.
- Kebreab, Ermias et al. (2023). "A meta-analysis of effects of 3-nitrooxypropanol on methane production, yield, and intensity in dairy cattle". In: *Journal of Dairy Science* 106.2, pp. 927–936. DOI: [10.3168/jds.2022-22211](https://doi.org/10.3168/jds.2022-22211).
- Korver, S., J.A.M. van Arendonk, and W.J. Koops (1985). "A function for live-weight change between two calvings in dairy cattle". In: *Anim Sci* 40.2, pp. 233–241.
- Li, M., K.F. Reed, et al. (2023). "A stochastic animal life cycle simulation model for a whole dairy farm system model: Assessing the value of combined heifer and lactating dairy cow reproductive management programs". In: *J Dairy Sci* 106.5, pp. 3246–3267.
- Li, M., G.J.M. Rosa, et al. (2022). "Investigating the effect of temporal, geographic, and management factors on US Holstein lactation curve parameters". In: *J Dairy Sci* 105.9, pp. 7525–7538.
- Mills, J.A.N. et al. (2003). "Alternative approaches to predicting methane emissions from dairy cows". In: *Journal of Animal Science* 81.12, pp. 3141–3150. DOI: [10.2527/2003.81123141x](https://doi.org/10.2527/2003.81123141x).
- Monteny, G.J. et al. (2002). "Prediction of ammonia emission from dairy barns using feed characteristics part II: Relation between urinary urea concentration and ammonia emission". In: *Journal of Dairy Science* 85.12, pp. 3389–3394. DOI: [10.3168/jds.S0022-0302\(02\)74426-3](https://doi.org/10.3168/jds.S0022-0302(02)74426-3).
- National Academies of Sciences, Engineering, and Medicine (2021). *Nutrient Requirements of Dairy Cattle: Eighth Revised Edition*. Washington, DC: The National Academies Press.
- National Research Council (2001). *Nutrient Requirements of Dairy Cattle*. Washington, D.C.: National Academies Press.
- Nennich, T.D. et al. (2005). "Prediction of manure and nutrient excretion from dairy cattle". In: *Journal of Dairy Science* 88.10, pp. 3721–3733. DOI: [10.3168/jds.S0022-0302\(05\)73058-7](https://doi.org/10.3168/jds.S0022-0302(05)73058-7).
- Niu, M., E. Kebreab, A.N. Hristov, et al. (2018). "Prediction of enteric methane production, yield, and intensity in dairy cattle using an intercontinental database". In: *Global Change Biology* 24, pp. 3368–3389. DOI: [10.1111/gcb.14094](https://doi.org/10.1111/gcb.14094).
- Pattanaik, A.K. et al. (2003). "Influence of grain processing and dietary protein degradability on nitrogen metabolism, energy balance and methane production in young calves". In: *Asian-Australasian Journal of Animal Sciences* 16.10, pp. 1443–1450. DOI: [10.5713/ajas.2003.1443](https://doi.org/10.5713/ajas.2003.1443).



- Reed, K.F. et al. (2015). "Predicting nitrogen excretion from cattle". In: *Journal of Dairy Science* 98.5, pp. 3025–3035.
- Reuven, Y.R. and D.P. Kroese (2017). *Simulation and the Monte Carlo Method*. In: *Wiley Series in Probability and Statistics*. John Wiley and Sons Inc.
- Vadas, P.A. et al. (2007). "A model for phosphorus transformation and runoff loss for surface-applied manures". In: *Journal of Environmental Quality* 36.1, pp. 324–332.
- Wilkerson, V. A., D. R. Mertens, and D. P. Casper (1997). "Prediction of excretion of manure and nitrogen by Holstein dairy cattle". In: *Journal of Dairy Science* 80.12, pp. 3193–3204.
- Wiltbank, M.C. et al. (2016). "Pivotal periods for pregnancy loss during the first trimester of gestation in lactating dairy cows". In: *Theriogenology* 86.1, pp. 239–253.
- Wood, P. D. P. (1967). "Algebraic Model of the Lactation Curve in Cattle". In: *Nature* 216.5111, pp. 164–165.
DOI: [10.1038/216164a0](https://doi.org/10.1038/216164a0).