

Probability

Notes taken by Runqiu Ye
Lectures by Konstantin Tikhomirov
Carnegie Mellon University

Spring 2026

Contents

1	Measure theory review	3
1.1	Measurable space and mapping	3
1.2	Measure space	4
1.3	π - λ theorem	4
1.4	Extension theorems	5

1 Measure theory review

1.1 Measurable space and mapping

Definition (σ -field). A collection of subsets $\Sigma \subset 2^\Omega$ is a σ -field if

- $\emptyset \in \Sigma$.
- If $A \in \Sigma$, then $A^c \in \Sigma$.
- If $\{A_i\}_{i=1}^\infty \subset \Sigma$, then $\bigcup_{i=1}^\infty A_i \in \Sigma$.

The pair (Ω, Σ) is called a measurable space.

Definition (atom). Let Σ be a σ -field. Say $A \in \Sigma$ is an atom if for all $B \in \Sigma$ either $A \subset B$ or $A \cap B = \emptyset$.

Proposition. For all $\omega \in \Omega$, there exists atom $A \in \Sigma$ containing ω if Ω is finite or countable.

Proof. Define $\tilde{A} = \bigcap \{B \in \Sigma : \omega \in B\}$. We can check that $\tilde{A} \in \Sigma$ and \tilde{A} is an atom containing ω . \square

Corollary. If Ω is finite or countable, there exists a partition $\Omega = \bigsqcup_i \Omega_i$, where each Ω_i is an atom of Σ . With this partition, Σ is just the power set with respect to $\{\Omega_i\}_i$.

Definition. If $F \subset 2^\Omega$, then the σ -field generated by F is the smallest σ -field containing all elements of F . Write this σ -field as $\sigma(F)$.

Example. Let $\Omega = \{1, 2, 3, 4, 5\}$ and $F = \{\{2, 3\}, \{3, 4\}\}$. Construct σ -field Σ generated by F . Σ is all possible union of sets from the collection $\{\{2\}, \{3\}, \{4\}, \{1, 5\}\}$.

Definition (measurable mapping). Given two measurable spaces (Ω, Σ) and $(\tilde{\Omega}, \tilde{\Sigma})$. Then $f : \Omega \rightarrow \tilde{\Omega}$ is measurable if $f^{-1}(B) \in \Sigma$ for all $B \in \tilde{\Sigma}$.

Definition (Borel σ -field). Let (T, τ) be a topological space. Then the Borel σ -field $\mathcal{B}(T, \tau)$ is defined as the smallest σ -field containing all open sets.

Definition (product measurable space). Given two measurable spaces (Ω, Σ) and $(\tilde{\Omega}, \tilde{\Sigma})$. We can define the product measurable space as follows: let the ground set be $\Omega \times \tilde{\Omega}$, and let $\Sigma \otimes \tilde{\Sigma}$ be the smallest σ -field containing all rectangles $B \times \tilde{B}$ where $B \in \Sigma$ and $\tilde{B} \in \tilde{\Sigma}$.

More generally, let Λ be an index set and $(\Omega_\lambda, \Sigma_\lambda)_{\lambda \in \Lambda}$. Define the product σ -field $\bigotimes_{\lambda \in \Lambda} \Sigma_\lambda$ be the smallest σ -field containing all elements in the form of $\prod_{\lambda \in \Lambda} B_\lambda$ where $B_\lambda \in \Sigma_\lambda$ and $B_\lambda = \Omega_\lambda$ for all but countably many indices.

Proposition. Let $(\Omega_i, \Sigma_i)_{i=1}^n$ be measurable spaces and $(\prod_{i=1}^n \Omega_i, \bigotimes_{i=1}^n \Sigma_i)$ be the product space. Let (Ω, Σ) be the domain and $f = (f_1, \dots, f_n) : (\Omega, \Sigma) \rightarrow (\prod_{i=1}^n \Omega_i, \bigotimes_{i=1}^n \Sigma_i)$. Suppose f is measurable, then every coordinate projection $f_i : \Omega \rightarrow \Omega_i$ is measurable.

This is also true for arbitrary index set.

Proposition. If $f : (\Omega, \Sigma) \rightarrow (\Omega_f, \Sigma_f)$ and $g : (\Omega, \Sigma) \rightarrow (\Omega_g, \Sigma_g)$, then the concatenation (f, g) is measurable w.r.t. the product space $(\Omega_f \times \Omega_g, \Sigma_f \otimes \Sigma_g)$.

Proof. Let $A \times B$ be such that $A \in \Sigma_f$ and $B \in \Sigma_g$. Then the preimage

$$(f, g)^{-1}(A \times B) = f^{-1}(A) \cap g^{-1}(B) \in \Sigma.$$

By definition, the product σ -field is generated by rectangles, so the proof is complete. \square

1.2 Measure space

Definition (measure). Let (Ω, Σ) be a measurable space. Then $\mu : \Sigma \rightarrow [0, \infty]$ is a measure if

- $\mu(\emptyset) = 0$.
- If $A_i \in \Sigma$ is pairwise disjoint then $\mu(\bigcup_{i=1}^{\infty} A_i) = \sum_{i=1}^{\infty} \mu(A_i)$.

Proposition (continuity of measure). If $A_1 \subset A_2 \subset \dots$ is a nested sequence of elements of Σ and μ be any measure on (Ω, Σ) . Then

$$\mu\left(\bigcup_{i=1}^{\infty} A_i\right) = \lim_{i \rightarrow \infty} \mu(A_i).$$

If $A_1 \supset A_2 \supset \dots$ is a nested sequence of elements of Σ and $\mu(A_n) < \infty$ for some n . Then

$$\mu\left(\bigcap_{i=1}^{\infty} A_i\right) = \lim_{i \rightarrow \infty} \mu(A_i).$$

Definition. Let (Ω, Σ, μ) be a measure space.

Say μ is σ -finite if there is a representation $\Omega = \bigcup_{i=1}^{\infty} \Omega_i$ where $\Omega_i \in \Sigma$ and $\mu(\Omega_i) < \infty$.

Say μ is a probability measure if $\mu(\Omega) = 1$.

Definition (completion of measure space). Let (Ω, Σ, μ) be a measure space. Let

$$\tilde{\Sigma} = \{A \cup B : A \in \Sigma, B \subset \Omega, \text{there exists } C \in \Sigma \text{ with } \mu(C) = 0 \text{ and } B \subset C\}.$$

We can check $\tilde{\Sigma}$ is a σ -field. If $\tilde{\mu}$ is a measure on $(\Omega, \tilde{\Sigma})$ which agrees with μ on Σ , then $(\Omega, \tilde{\Sigma}, \tilde{\mu})$ is called a completion of (Ω, Σ, μ) .

1.3 π - λ theorem

Definition (π -system). Let Ω be a set and \mathcal{P} be a collection of subsets of Ω . Then \mathcal{P} is a π -system if it is closed with respect to taking finite intersections. That is, $A, B \in \mathcal{P}$ implies $A \cap B \in \mathcal{P}$.

Example. On the real line \mathbb{R} , both $\mathcal{P}_1 = \{(a, b) : a < b\}$ and $\mathcal{P}_2 = \{(-\infty, a] : a \in \mathbb{R}\}$ are π -systems.

Definition (λ -system). Let Ω be a set and \mathcal{L} be a collection of subsets of Ω . Say \mathcal{L} is a λ -system if

- $\emptyset \in \mathcal{L}$.
- $A \in \mathcal{L}$ implies $A^c \in \mathcal{L}$.

- for all countable collection of disjoint elements $A_i \in \mathcal{L}$, we have $\bigcup_{i=1}^{\infty} A_i \in \mathcal{L}$.

Theorem (π - λ theorem). Let Ω be a set, \mathcal{P} be a π -system and \mathcal{L} be a λ -system. Also suppose $\mathcal{P} \subset \mathcal{L}$, then $\sigma(\mathcal{P}) \subset \mathcal{L}$.

Proof. Let $\ell(\mathcal{P})$ be the smallest λ -system on Ω containing \mathcal{P} . The goal is to show that $\ell(\mathcal{P})$ is a σ -field. We need to show that if $A_i \in \ell(\mathcal{P})$ for $1 \leq i < \infty$, then $\bigcup_{i=1}^{\infty} A_i \in \ell(\mathcal{P})$. Note that

$$\bigcup_{i=1}^{\infty} A_i = \bigcup_{i=1}^{\infty} \left(A_i \setminus \bigcup_{j=1}^{i-1} A_j \right),$$

so it suffices to show that $A, B \in \ell(\mathcal{P})$ implies $A \cap B \in \ell(\mathcal{P})$.

For $A \in \ell(\mathcal{P})$ we define

$$W_A = \{B \subset \Omega : A \cap B \in \ell(\mathcal{P})\}.$$

It can be directly verified that W_A is a λ -system.

Take $A \in \mathcal{P}$, then for any $B \in \mathcal{P}$ we have $A \cap B \in \mathcal{P} \subset \ell(\mathcal{P})$. Hence, $\mathcal{P} \subset W_A$ and thus $\ell(\mathcal{P}) \subset W_A$ for all $A \in \mathcal{P}$, as $\ell(\mathcal{P})$ is the smallest λ -system on Ω containing \mathcal{P} . Now take $A \in \ell(\mathcal{P})$, we have $A \in W_B$ for all $B \in \mathcal{P}$. It follows that $A \cap B \in \ell(\mathcal{P})$ and thus $B \in W_A$. Hence similarly $\ell(\mathcal{P}) \subset W_A$ for all $A \in \ell(\mathcal{P})$.

Now for any pair $B, C \in \ell(\mathcal{P})$, we have $C \in W_B$ and thus $B \cap C \in \ell(\mathcal{P})$. This completes the proof. \square

1.4 Extension theorems

Definition (semi-field). A collection of subsets $S \subset 2^{\Omega}$ is a semi-field if

- $\emptyset \in S$ and $\Omega \in S$.
- $A, B \in S$ implies $A \cap B \in S$.
- If $A \in S$, then A^c is a finite disjoint union of sets in S .

Theorem (Caratheodory's extension theorem). Let S be a semi-field and let μ be a non-negative function on S satisfying:

- $\mu(\emptyset) = 0$.
- If A_1, \dots, A_n are disjoint and $\bigcup_{i=1}^n A_i \in S$, then $\mu(\bigcup_{i=1}^n A_i) = \sum_{i=1}^n \mu(A_i)$.
- If A_1, A_2, \dots are such that $\bigcup_{i=1}^{\infty} A_i \in S$, then $\mu(\bigcup_{i=1}^{\infty} A_i) \leq \sum_{i=1}^{\infty} \mu(A_i)$.

Then μ admits a unique extension $\bar{\mu}$ which is a measure on \bar{S} , the field (algebra) generated by S . Moreover, if $\bar{\mu}$ is σ -finite then $\bar{\mu}$ admits a unique extension $\tilde{\mu}$ to $\sigma(S)$.

Definition. Let T be any set. Write

$$\mathbb{R}^T = \{(\omega_t)_{t \in T} : \omega_t \in \mathbb{R}\}.$$

Also write \mathcal{R}^T as the σ -field generated by rectangles of the form $\prod_{t \in T} I_t$, where for each $t \in T$, I_t is either a semi-open interval of the form $(a, b]$ with $a < b$ or $I_t = \mathbb{R}$, and $I_t = \mathbb{R}$ for all but finitely many $t \in T$.

Theorem (Kolmogorov's extension theorem). For each finite non-empty subset $J \subset T$, let μ_J be a Borel probability measure in \mathbb{R}^J , and assume that the measures $(\mu_J)_{J \subset T, |J| < \infty}$ are compatible, in the sense that whenever $J_1 \subset J_2 \subset T$ with $0 \leq |J_1| \leq |J_2| < \infty$, $I_j \subset \mathbb{R}$ with $j \in J_1$ are Borel subsets of \mathbb{R} , and

$$\tilde{I}_j = \begin{cases} I_j & (j \in J_1) \\ \mathbb{R} & (j \in J_2 \setminus J_1), \end{cases}$$

one has

$$\mu_{J_2} \left(\prod_{j \in J_2} \tilde{I}_j \right) = \mu_{J_1} \left(\prod_{j \in J_1} I_j \right).$$

Then there exists a unique probability measure μ on $(\mathbb{R}^T, \mathcal{R}^T)$ consistent with $(\mu_J)_{J \subset T, |J| < \infty}$. That is, one has

$$\mu \left(\prod_{t \in T} I_t \right) = \mu_J \left(\prod_{j \in J} I_j \right)$$

whenever $J \subset T$ with $|J| < \infty$ and $I_t = \mathbb{R}$ for all $t \notin J$.