

Problem Set #1: Supervised Learning

Problem 1 Linear Classifiers (Logistic Regression and GDA)

Consider two datasets provided in the following files:

- i. data/ds1_{train,valid}.csv
- ii. data/ds2_{train,valid}.csv

Each file contains m examples, one example per row. The i -th row contains columns $x_0^{(i)} \in \mathbb{R}$, $x_1^{(i)} \in \mathbb{R}$ and $y^{(i)} \in \{0, 1\}$. Use logistic regression and GDA to perform binary classification.

(a) Average empirical loss for logistic regression:

$$J(\theta) = -\frac{1}{m} \sum_{i=1}^m y^{(i)} \log(h_{\theta}(x^{(i)})) + (1 - y^{(i)}) \log(1 - h_{\theta}(x^{(i)})),$$

where $y^{(i)} \in \{0, 1\}$, $h_{\theta}(x^{(i)}) = g(\theta^T x)$ and $g(z) = 1/(1 + e^{-z})$.

The gradient of the function

$$\frac{\partial J}{\partial \theta_j} = -\frac{1}{m} \sum_{i=1}^m (y^{(i)} - h_{\theta}(x^{(i)})) x_j^{(i)}.$$

It follows that

$$\frac{\partial^2 J}{\partial \theta_k \partial \theta_j} = \frac{1}{m} \sum_{i=1}^m h_{\theta}(x^{(i)}) (1 - h_{\theta}(x^{(i)})) x_k^{(i)} x_j^{(i)}.$$

Hence, The Hessian H of this function is

$$H = \frac{1}{m} \sum_{i=1}^m h_{\theta}(x^{(i)}) (1 - h_{\theta}(x^{(i)})) x^{(i)} (x^{(i)})^T.$$

Now, for any vector z , using Einstein's summation, we have

$$\begin{aligned} z^T H z &= \frac{1}{m} \sum_{i=1}^m h_{\theta}(x^{(i)}) (1 - h_{\theta}(x^{(i)})) z_k x_k^{(i)} x_j^{(i)} z_j \\ &= \frac{1}{m} \sum_{i=1}^m h_{\theta}(x^{(i)}) (1 - h_{\theta}(x^{(i)})) (x^T z)^2 \\ &\geq 0 \end{aligned}$$

This shows that H is PSD, and J is convex.

(b) **Coding problem.**

(c) To show that GDA results in a classifier that has a linear decision boundary, we want to show

$$p(y = 1 \mid x; \phi, \mu_0, \mu_1, \Sigma) = \frac{1}{1 + \exp(-(\theta^T x + \theta_0))}$$

for some $\theta \in \mathbb{R}^n$ and $\theta_0 \in \mathbb{R}$ as functions of ϕ , Σ , μ_0 , and μ_1 . We have

$$\begin{aligned} p(y = 1 \mid x) &= \frac{p(x \mid y = 1)p(y = 1)}{p(x \mid y = 1)p(y = 1) + p(x \mid y = 0)p(y = 0)} \\ &= \frac{\phi \exp(-\frac{1}{2}(x - \mu_1)^T \Sigma^{-1}(x - \mu_1))}{\phi \exp(-\frac{1}{2}(x - \mu_1)^T \Sigma^{-1}(x - \mu_1)) + (1 - \phi) \exp(-\frac{1}{2}(x - \mu_0)^T \Sigma^{-1}(x - \mu_0))} \\ &= \frac{1}{1 + \frac{1-\phi}{\phi} \exp(-\frac{1}{2}(x - \mu_0)^T \Sigma^{-1}(x - \mu_0) + \frac{1}{2}(x - \mu_1)^T \Sigma^{-1}(x - \mu_1))} \\ &= \frac{1}{1 + \frac{1-\phi}{\phi} \exp(-((\mu_1 - \mu_0)^T \Sigma^{-1}x + \frac{1}{2}(\mu_0^T \Sigma^{-1} \mu_0 - \mu_1^T \Sigma^{-1} \mu_1)))}. \end{aligned}$$

This is the desired form, where

$$\begin{aligned} \theta &= \Sigma^{-1}(\mu_1 - \mu_0), \\ \theta_0 &= \frac{1}{2}(\mu_0^T \Sigma^{-1} \mu_0 - \mu_1^T \Sigma^{-1} \mu_1) - \log \frac{1 - \phi}{\phi}. \end{aligned}$$

(d) The log-likelihood of the data is

$$\begin{aligned} \ell(\phi, \mu_0, \mu_1, \Sigma) &= \log \prod_{i=1}^m p(x^{(i)} \mid y^{(i)}; \mu_0, \mu_1, \Sigma) p(y^{(i)}; \phi) \\ &= \sum_{i=1}^m 1\{y^{(i)} = 1\} \left(-\frac{1}{2}(x^{(i)} - \mu_1)^T \Sigma^{-1}(x^{(i)} - \mu_1) + \log \phi \right) \\ &\quad + \sum_{i=1}^m 1\{y^{(i)} = 0\} \left(-\frac{1}{2}(x^{(i)} - \mu_0)^T \Sigma^{-1}(x^{(i)} - \mu_0) + \log(1 - \phi) \right) \\ &\quad - \frac{m}{2} \log |\Sigma| + C, \end{aligned}$$

where C is some constant independent of the parameters.

Let $\nabla_{\phi} \ell = 0$, we have

$$\phi = \frac{1}{m} \sum_{i=1}^m 1\{y^{(i)} = 1\}.$$

Let $\nabla_{\mu_1} \ell = 0$, we have

$$\sum_{i=1}^m 1\{y^{(i)} = 1\} \Sigma^{-1} x^{(i)} = \sum_{i=1}^m 1\{y^{(i)} = 1\} \Sigma^{-1} \mu_1,$$

and thus

$$\mu_1 = \frac{\sum_{i=1}^m 1\{y^{(i)} = 1\}x^{(i)}}{\sum_{i=1}^m 1\{y^{(i)} = 1\}}, \quad \mu_0 = \frac{\sum_{i=1}^m 1\{y^{(i)} = 0\}x^{(i)}}{\sum_{i=1}^m 1\{y^{(i)} = 0\}}.$$

To derive Σ , recall that $\nabla_A \log |A| = (A^{-1})^T$, so we have

$$\nabla_{\Sigma^{-1}} \ell = -\frac{m}{2} \Sigma^{-1} + \frac{1}{2} \sum_{i=1}^m (x^{(i)} - \mu_{y^{(i)}})(x^{(i)} - \mu_{y^{(i)}})^T.$$

Hence,

$$\Sigma = \frac{1}{m} \sum_{i=1}^m (x^{(i)} - \mu_{y^{(i)}})(x^{(i)} - \mu_{y^{(i)}})^T.$$

We conclude that the maximum likelihood estimates of the parameters are given by

$$\begin{aligned} \phi &= \frac{1}{m} \sum_{i=1}^m 1\{y^{(i)} = 1\}, \\ \mu_0 &= \frac{\sum_{i=1}^m 1\{y^{(i)} = 0\}x^{(i)}}{\sum_{i=1}^m 1\{y^{(i)} = 0\}}, \\ \mu_1 &= \frac{\sum_{i=1}^m 1\{y^{(i)} = 1\}x^{(i)}}{\sum_{i=1}^m 1\{y^{(i)} = 1\}}, \\ \Sigma &= \frac{1}{m} \sum_{i=1}^m (x^{(i)} - \mu_{y^{(i)}})(x^{(i)} - \mu_{y^{(i)}})^T. \end{aligned}$$

(e) **Coding problem.**

(f) See jupyter notebook for plots.

(g) See jupyter notebook for plots. On Dataset 1 GDA perform worse than logistic regression. This might be the case because for Dataset 1, the distribution of features are not quite multivariate normal.

(h) *** TO-DO ***

■

Problem 2 Incomplete, Positive-Only Labels

Dataset without full access to labels. In particular, we have labels only for a subset of positive examples. All negative examples and the rest of positive examples are unlabeled. Assume dataset $\{(x^{(i)}, t^{(i)}, y^{(i)})\}_{i=1}^m$ where $t^{(i)} \in \{0, 1\}$ is true label and where

$$y^{(i)} = \begin{cases} 1 & x^{(i)} \text{ is labeled} \\ 0 & \text{otherwise.} \end{cases}$$

All labeled examples are positive, which is to say $p(t^{(i)} = 1 \mid y^{(i)} = 1) = 1$. Goal is to construct a binary classifier h of true label t which only access to partial labels y . That is, construct h such that $h(x^{(i)}) \approx p(t^{(i)} = 1 \mid x^{(i)})$ as closely as possible, using only x and y .

- (a) Suppose each $y^{(i)}$ and $x^{(i)}$ conditionally independent given $t^{(i)}$:

$$p(y^{(i)} = 1 \mid t^{(i)} = 1, x^{(i)}) = p(y^{(i)} = 1 \mid t^{(i)} = 1).$$

That is, labeled examples are selected uniformly at random from positive examples.

Want to show $p(t^{(i)} = 1 \mid x^{(i)}) = p(y^{(i)} = 1 \mid x^{(i)})/\alpha$ for some $\alpha \in \mathbb{R}$. As $p(\cdot \mid x^{(i)})$ is a conditional measure, we have

$$\begin{aligned} p(y^{(i)} = 1 \mid x^{(i)}) &= p(y^{(i)} = 1 \mid t^{(i)} = 1, x^{(i)})p(t^{(i)} = 1 \mid x^{(i)}) \\ &\quad + p(y^{(i)} = 1 \mid t^{(i)} = 0, x^{(i)})p(t^{(i)} = 0 \mid x^{(i)}) \\ &= p(y^{(i)} = 1 \mid t^{(i)} = 1, x^{(i)})p(t^{(i)} = 1 \mid x^{(i)}) \\ &= p(y^{(i)} = 1 \mid t^{(i)} = 1)p(t^{(i)} = 1 \mid x^{(i)}). \end{aligned}$$

Hence, $p(t^{(i)} = 1 \mid x^{(i)}) = p(y^{(i)} = 1 \mid x^{(i)})/\alpha$ where $\alpha = p(y^{(i)} = 1 \mid t^{(i)} = 1)$.

- (b) Estimate α using a trained classifier h and a held-out validation set V . Let $V_+ = \{x^{(i)} \in V \mid y^{(i)} = 1\}$. Assuming $h(x^{(i)}) \approx p(y^{(i)} = 1 \mid x^{(i)})$ for all $x^{(i)}$. Want to show

$$h(x^{(i)}) \approx \alpha \text{ for all } x^{(i)} \in V_+.$$

May assume that $p(t^{(i)} = 1 \mid x^{(i)}) \approx 1$ when $x^{(i)} \in V_+$.

We have

$$\begin{aligned} h(x^{(i)}) &\approx p(y^{(i)} = 1 \mid x^{(i)}) \\ &= p(y^{(i)} = 1 \mid t^{(i)} = 1, x^{(i)})p(t^{(i)} = 1 \mid x^{(i)}) \\ &\approx \alpha. \end{aligned}$$

- (c) **Coding problem.**

- (d) **Coding problem.**

- (e) **Coding problem.** Estimate the constant α using validation set.

$$\alpha \approx \frac{1}{|V_+|} \sum_{x^{(i)} \in V_+} h(x^{(i)}).$$

To plot the decision boundary, we need to calculate the rescaled θ , write θ_* . The new decision boundary is given by $\frac{1}{\alpha} \frac{1}{1+\exp(-\theta^T x)} = \frac{1}{2}$. We have

$$\theta^T x + \log\left(\frac{2}{\alpha} - 1\right) = 0.$$

This is equivalent to $\theta_*^T x = 0$. This shows that θ_* and θ differs only in the 0-th index by a constant $\log\left(\frac{2}{\alpha} - 1\right)$. ■

Problem 3 Poisson Regression

- (a) The poisson distribution parametrized by λ is

$$p(y; \lambda) = \frac{e^{-\lambda} \lambda^y}{y!}.$$

Therefore, we have

$$p(y; \lambda) = \frac{1}{y!} \exp(-\lambda + y \log \lambda).$$

Compare with $p(y; \eta) = b(y) \exp(\eta^T T(y) - a(\eta))$, we conclude that the poisson distribution is in the exponential family, with

$$\begin{aligned} b(y) &= \frac{1}{y!}, \\ T(y) &= y, \\ \eta &= \log \lambda, \\ a(\eta) &= e^\eta. \end{aligned}$$

- (b) The canonical response function for the family

$$\mathbb{E}[T(y); \eta] = \mathbb{E}[T(y); \eta] = \lambda = e^\eta.$$

- (c) For a general linear model and a training set, the log likelihood

$$\begin{aligned} \log p(y^{(i)} | x^{(i)}; \eta) &= \log b(y) \exp(\eta^T T(y) - a(\eta)) \\ &= \log b(y) + \eta^T T(y) - a(\eta). \end{aligned}$$

For our model with poisson responses y , we have

$$\ell = \log p(y^{(i)} \mid x^{(i)}; \theta) = -\log y! + (\theta^T x^{(i)}) y^{(i)} - \exp(\theta^T x^{(i)}).$$

Taking the derivative with respect to θ_j , we have

$$\frac{\partial \ell}{\partial \theta_j} = (y^{(i)} - \exp(\theta^T x^{(i)})) x_j^{(i)}$$

Hence, the stochastic gradient ascent update rule for learning using a GLM model with poisson response y is

$$\begin{aligned} \theta_j &:= \theta_j + \alpha \frac{\partial \ell}{\partial \theta_j} \\ &:= \theta_j + \alpha (y^{(i)} - \exp(\theta^T x^{(i)})) x_j^{(i)}. \end{aligned}$$

- (d) **Coding problem.** To predict the dataset, recall that the hypothesise function for our model with poisson response y is

$$h_\theta(x) = \mathbb{E}[y \mid x] = e^\eta = e^{\theta^T x}.$$

■