OPPERMANN INTERVALS IN $\mathbb{F}_q[t]$: GALOIS OBSTRUCTIONS, EXACT COUNTS FOR SMALL DEGREE, AND THE OPPERMANN SUBGROUP IN CHARACTERISTIC 2

RUQING CHEN

ABSTRACT. For a monic $f \in \mathbb{F}_q[t]$ of degree d, the Legendre interval $\{f^2 + s : \deg s \le d\}$ splits into the q *Oppermann sub-intervals* $O_{f,c} = \{f^2 + cf + s : \deg s < d\}$, each of size q^d. We ask whether every $O_{f,c}$ contains an irreducible polynomial.

In characteristic 2 the answer is no, and the failures have three Galois-theoretic causes: (A) for $d = 1$ the piece $c = 0$ consists of squares; (B) for $d = 2$, Frobenius acts on the roots of an additive polynomial through $\mathrm{AGL}_2(\mathbb{F}_2) \cong S_4$, and the resulting exact counts vanish exactly for $q = 2$ and $q = 4^j$; (C) for $q = 2$, $d = 4$, the Stickelberger–Swan sign of Frobenius obstructs. The additive mechanism of (B) disappears for $d = 2^{r-1} \ge 4$, because $\mathrm{AGL}_r(\mathbb{F}_2)$ contains no 2^r-cycle when $r \ge 3$.

For odd q and $d = 2$, a ternary trace form on $\mathbb{F}_{q^4}$ gives $\pi = (q^2 - 1 - (q - \eta(-1))\eta(2e))/4 > 0$. In characteristic 2 and $d = 3$, a count in the tower $\mathbb{F}_q \subset \mathbb{F}_{q^3} \subset \mathbb{F}_{q^6}$ together with a cubic Gauss sum gives every class exactly. There are $q(q^2 - 1)/6$ irreducibles for k odd. For k even there are $q(q-1)^2/6$, $q(q^2 + 1 - 2\varepsilon\sqrt{q})/6$ or $q(q^2 + 1 + \varepsilon\sqrt{q})/6$, according as the t^3-coefficient is 0, a cube or a non-cube, where $\varepsilon = (-1)^{k/2}$. So $d \le 2$ is settled for every q, and $d = 3$ for every even q.

For fixed $d \ge 3$ and large q, the theorem of Bank, Bary-Soroker and Rosenzweig, together with the $d = 3$ count, gives nonemptiness. Exact counts from the Hayes explicit formula, evaluated by FFT, give nonemptiness in odd characteristic for $q = 3$, $d \le 16$; $q = 5$, $d \le 10$; $q = 7$, $d \le 9$; $q = 9$, $d \le 8$; $q = 11$, $d \le 7$; $q = 13$, $d \le 6$. In characteristic 2 they find no further empty sub-intervals for $q = 2$, $d \le 26$; $q = 4$, $d \le 13$; $q = 8$, $d \le 7$; $q = 16$, $d \le 5$.

Finally, in characteristic 2 the Oppermann classes form a subgroup H_d of the Hayes group: the kernel of the odd power sums, with $G_d/H_d \cong \mathbb{F}_q^{\lfloor d/2 \rfloor}$. Its annihilator yields Artin–Schreier sums. The mean of Ψ over H_d drifts by $-2q^2 + 2q$ at $d = 4$ with k odd and by $q^2 - q$ at $d = 5$. The variance is a coherent sum over annihilator cosets, maximal at $d = 2$, where $\mathrm{Var}_{H_2}(\Psi)/q^2 = q - 1$.

1. INTRODUCTION

Oppermann's conjecture [8] asserts that for every $n \ge 2$ both (n^2, n^2+n) and $(n^2+n, (n+1)^2)$ contain a prime. It implies Legendre's conjecture and lies beyond the Riemann hypothesis, which gives primes only in intervals $[x, x + C\sqrt{x}\log x]$. Over $\mathbb{F}_q[t]$, Legendre intervals $I_f = \{f^2 + s : \deg s \le d\}$ (f monic of degree d) were studied in [4]. Since $f^2 + cf$ and $f^2 + ct^d$ differ by a polynomial of degree $< d$, the interval I_f is the disjoint union of the q sub-intervals

$$O_{f,c} = \{f^2 + cf + s : \deg s < d\} = \{f^2 + ct^d + s : \deg s < d\}, \qquad c \in \mathbb{F}_q, \tag{1}$$

each of cardinality $q^d = |f|$, the norm of f, exactly as $(n^2, n^2 + n)$ has length n. For $q = 2$ there are two halves. We call $O_{f,c}$ the *Oppermann sub-intervals* and ask:

Does every $O_{f,c}$ contain an irreducible polynomial?

The answer depends on the characteristic.

Theorem 1.1 (characteristic 2)**.** *Let $q = 2^k$.*

Date: October 13, 2026.

2020 *Mathematics Subject Classification.* Primary 11T55; Secondary 11T06, 11T24, 11N05, 11Y16.

Key words and phrases. Oppermann's conjecture, function fields, irreducible polynomials, short intervals, Hayes characters, Stickelberger–Swan theorem, Gauss sums, Artin–Schreier sums, explicit formula.

Code and data: https://github.com/Ruqing1963/oppermann-function-fields (archived at DOI 10.5281/zenodo.23353949).

(A) *For $d = 1$ the sub-interval $O_{f,0} = \{t^2 + s_0 : s_0 \in \mathbb{F}_q\}$, which is the same set for all q polynomials f, consists of squares. Every $O_{f,c}$ with $c \neq 0$ contains exactly $q/2$ irreducibles.*

(B) *For $d = 2$ and $f = t^2 + a_1 t + a_0$, the number of irreducibles in $O_{f,c}$ depends only on $e = a_1^2 + c$ and equals*

	$e = 0$	$e \neq 0$
k even	0	$q^2/4$
k odd	$q(q-1)/2$	$q(q-2)/4$

The table has two zero entries: k even with $e = 0$, and $q(q-2)/4$ at $q = 2$ (where $k = 1$ is odd). Hence $O_{f,c}$ is empty if and only if either k is even and $c = a_1^2$, or $q = 2$ and $c = a_1^2 + 1$. For $q = 2$ the block $e = 0$ is not *empty: it contains $q(q-1)/2 = 1$ irreducible, $t^4 + t + 1$. For every $q \in \{2, 4, 16, 64, \dots\}$ there is thus exactly one empty sub-interval as a set: B_0, or B_1 when $q = 2$. It is represented by q^2 pairs (f, c), one c for each of the q^2 polynomials f. For $q \in \{8, 32, 128, \dots\}$ there is none.*

(C) *For $q = 2$ and $d = 4$ the block $Z_4 = \{t^8 + s : \deg s \leq 3\}$ contains no irreducible polynomial. It equals $O_{f,c}$ for the 8 pairs $f = t^4 + a_2 t^2 + a_1 t + a_0$ (that is, $a_3 = 0$), $c = a_2$.*

Theorem 1.2 (odd characteristic, computer-assisted)**.** *Let q be odd. Every Oppermann sub-interval $O_{f,c}$ contains an irreducible polynomial whenever*

$$q = 3,\ d \leq 16; \qquad q = 5,\ d \leq 10; \qquad q = 7,\ d \leq 9;$$
$$q = 9,\ d \leq 8; \qquad q = 11,\ d \leq 7; \qquad q = 13,\ d \leq 6.$$

Equivalently (Lemma 2.1), every interval $\{P + s : \deg s < d\}$ with P monic of degree $2d$ contains an irreducible polynomial in these ranges.

Theorem 1.3 (characteristic 2, computer-assisted)**.** *For $q = 2$, $d \leq 26$; $q = 4$, $d \leq 13$; $q = 8$, $d \leq 7$; $q = 16$, $d \leq 5$, the only empty Oppermann sub-intervals are those of Theorem 1.1.*

Theorem 1.4 (odd characteristic, $d = 2$)**.** *Let q be odd and let η be the quadratic character of $\mathbb{F}_q$, with $\eta(0) = 0$. For $f = t^2 + a_1 t + a_0$ and $c \in \mathbb{F}_q$, put $e = 2a_0 + c - a_1^2/2$. Then*

$$\pi(O_{f,c}) = \frac{q^2 - 1 - (q - \eta(-1))\,\eta(2e)}{4}.$$

In particular, every Oppermann sub-interval with $d = 2$ contains at least $q(q-1)/4$ irreducibles if $q \equiv 1 \pmod 4$, and at least $(q+1)(q-2)/4$ if $q \equiv 3 \pmod 4$. Together with Theorem 1.1(B), this settles $d = 2$ for every q.

The additive mechanism behind (B) does not extend: for $d = 2^{r-1} \geq 4$ no affine polynomial of degree $2d$ is irreducible (Theorem 3.6, Corollary 3.7).

For fixed d and large q, the theorem of Bank, Bary-Soroker and Rosenzweig [1] covers most classes (Corollary 6.2). The one family it misses for $d \geq 3$ lies at $d = 3$ in characteristic 2, and there every class can be counted exactly.

Theorem 1.5 (characteristic 2, $d = 3$)**.** *Let $q = 2^k$, $\varepsilon = (-1)^{k/2}$ for k even, and let c be the coefficient of t^3 in an Oppermann sub-interval C with $d = 3$. Then*

$$\pi(C) = \begin{cases} q(q^2-1)/6, & k \text{ odd}, \\ q(q-1)^2/6, & k \text{ even},\ c = 0, \\ q(q^2 + 1 - 2\varepsilon\sqrt{q})/6, & k \text{ even},\ c \in (\mathbb{F}_q^\times)^3, \\ q(q^2 + 1 + \varepsilon\sqrt{q})/6, & k \text{ even},\ c \notin (\mathbb{F}_q^\times)^3. \end{cases}$$

In particular, no Oppermann sub-interval with $d = 3$ is empty in characteristic 2.

Corollary 1.6. *For $d \in \{1, 2\}$, the answer is known exactly for every q (Theorems 1.1(A),(B) and 1.4). In characteristic 2 this also holds for $d = 3$ (Theorem 1.5), and no sub-interval is empty there. For every $d \geq 3$ there is $q_0(d)$ such that every Oppermann sub-interval of degree $2d$ over $\mathbb{F}_q$ with $q \geq q_0(d)$ contains an irreducible polynomial.*

For $d \geq 3$, Corollary 6.2 covers every class except those with q even, $d = 3$ and $p_3 = 0$. Theorem 1.5, proved as Theorem 5.2 and Corollary 5.4, covers those, and indeed every class with q even and $d = 3$. In characteristic 2 and $d = 2$, the empty sub-intervals for $q = 2, 4, 16, 64, \ldots$ show that no such $q_0(2)$ exists there.

The tables above therefore matter mainly for $q < q_0(d)$; $q_0(d)$ is not explicit.

In characteristic 2 the Oppermann sub-intervals have additional structure.

Theorem 1.7 (the Oppermann subgroup)**.** *Let $q = 2^k$, and let $H_d \subset G_d$ be the set of Oppermann classes of level d.*

1. *H_d is the kernel of the odd power-sum homomorphism $P \mapsto (S_1, S_3, \ldots, S_{2\lfloor d/2 \rfloor - 1})$, so $G_d/H_d \cong \mathbb{F}_q^{\lfloor d/2 \rfloor}$ (Theorem 2.4). The characters of G_d/H_d have Artin–Schreier sums as explicit-formula coefficients (Theorem 2.5).*
2. *$\mathbb{E}_{H_d}[\Psi] - q^d = 0$ for $d = 2, 3$, and for $d = 4$ with k even. It equals $-2q^2 + 2q$ for $d = 4$ with k odd, and $q^2 - q$ for $d = 5$ (Corollary 7.2, Theorem 7.3).*
3. *$\mathrm{Var}_{H_2}(\Psi)/q^2 = q - 1$, whereas $\mathrm{Var}_{G_2}(\Psi)/q^2 = 1 - 1/q$. Further, $\mathrm{Var}_{H_3}(\Psi)/q^3 = 1 - 1/q$ for k odd and $3(1 - 1/q)$ for k even (Theorem 7.5).*

Throughout, a *sub-interval* means the set $O_{f,c}$. Distinct pairs (f, c) can give the same set; counts of sub-intervals are counts of sets, that is, of Hayes classes.

Relation to [4]. The Legendre interval I_f is the union of the q sub-intervals $O_{f,c}$. It is the class of f^2 at level $d - 1$; in characteristic 2 these classes form the subgroup of squares in G_{d-1}. [4] found that in characteristic 2 the variance of Ψ over Legendre intervals differs markedly from the variance over all classes. The difference is carried entirely by correlations between quadratic twists, that is, between characters trivial on the squares, and [4] gives no structural description of these twists. The proof of Theorem 2.4 applies verbatim at level $d - 1$. It shows that the squares in G_{d-1} are exactly the kernel of the odd power sums S_j, $j < d$. So the twists in question are the Artin–Schreier characters χ_β of Theorem 2.5, and their coefficients are the sums given there. At level d the subgroup H_d plays the same role for the Oppermann sub-intervals; §7 computes the resulting coherence exactly for $d \leq 3$. The obstructions (A)–(C) refine [4] in a different direction: they empty single sub-intervals $O_{f,c}$, while the Legendre intervals themselves contain primes in every case computed here and in [4].

Outline.

- Section 2 identifies the Oppermann sub-intervals with Hayes classes. In characteristic 2 it proves the Newton–trace duality.
- Sections 3–5 contain the exact algebraic results for $d \leq 3$: the three obstructions and the $\mathrm{AGL}_r(\mathbb{F}_2)$ dead zone, the ternary trace form, and the field-tower and Gauss-sum count.
- Section 6 describes the exact computation, the square-root barrier, the rigidity of the level-2 L-functions, and the large-q corollary.
- Section 7 studies the mean and the variance over the Oppermann subgroup.

2. Oppermann sub-intervals as Hayes classes and Newton–trace duality

For P monic of degree n write $P = t^n + p_{n-1}t^{n-1} + \cdots + p_0$. Two monic polynomials of degree n lie in the same *Hayes class of level l* if they agree in $p_{n-1}, \ldots, p_{n-l}$. Each class is a coset $P + \{s : \deg s < n - l\}$, and the map

$$P \mapsto u^n P(1/u) \bmod u^{l+1} = 1 + p_{n-1}u + \cdots + p_{n-l}u^l$$

identifies the classes with the elements of the multiplicative group $G_l = (1 + u\mathbb{F}_q[u])/(1 + u^{l+1}\mathbb{F}_q[u])$ of order q^l. This map is multiplicative: $\mathrm{class}(PQ) = \mathrm{class}(P)\,\mathrm{class}(Q)$ in G_l. The characters of G_l are the short-interval characters of Hayes [5] and Keating–Rudnick [6]. A character has *exact level j* if it factors through G_j but not through G_{j-1}. There are $q^j - q^{j-1}$ characters of exact level j, for $1 \leq j \leq l$. By (1), $O_{f,c}$ is the Hayes class of level d and degree $n = 2d$ containing $f^2 + ct^d$.

Lemma 2.1 (odd q)**.** *If q is odd, every Hayes class of level d in degree $2d$ is an Oppermann sub-interval, and it equals $O_{f,c}$ for exactly q pairs (f, c).*

Proof. Write $f = \sum_{i=0}^{d} a_i t^i$ with $a_d = 1$. For $1 \le j \le d$ the coefficient of t^{2d-j} in f^2 is

$$[f^2]_{2d-j} = \sum_{\substack{i+i'=2d-j \\ 0\le i,i'\le d}} a_i a_{i'} = 2a_{d-j} + \Phi_j, \qquad \Phi_j = \sum_{\substack{i+i'=2d-j \\ d-j<i,i'<d}} a_i a_{i'},$$

where the term $2a_{d-j} = 2a_d a_{d-j}$ comes from $\{i, i'\} = \{d, d-j\}$. The sum Φ_j runs over ordered pairs, so it includes the square $a_{d-j/2}^2$ when j is even. Φ_j depends only on $a_{d-1}, \dots, a_{d-j+1}$; for example $\Phi_1 = 0$, $\Phi_2 = a_{d-1}^2$, $\Phi_3 = 2a_{d-1}a_{d-2}$.

The map $(a_{d-1}, \dots, a_1) \mapsto ([f^2]_{2d-1}, \dots, [f^2]_{d+1})$ is therefore triangular with diagonal entries 2. Since $2 \in \mathbb{F}_q^\times$, it is a bijection $\mathbb{F}_q^{d-1} \to \mathbb{F}_q^{d-1}$: the prefix of a class determines $a_{d-1}, \dots, a_1$ uniquely, by solving $a_{d-j} = ([f^2]_{2d-j} - \Phi_j)/2$ for $j = 1, \dots, d-1$ in turn.

The remaining coefficient of the class is that of t^d in $f^2 + ct^d$, namely $2a_0 + \Phi_d + c$, where Φ_d is already fixed. For each of the q values of a_0 there is exactly one c giving the prescribed value. So every class arises, and from exactly q of the q^{d+1} pairs (f, c). □

So in odd characteristic the Oppermann question at degree $2d$ is the question whether *every* short interval of length $q^d = q^{n/2}$ contains a prime: no class is singled out.

Lemma 2.2 (q even)**.** *Let q be even. The Oppermann sub-intervals are exactly the Hayes classes of level d (in degree $2d$) in which the coefficient of t^j is 0 for every odd j with $d < j < 2d$. The coefficients of t^d and of t^j for even j with $d < j < 2d$ are arbitrary. There are $q^{\lceil d/2 \rceil}$ such classes among the q^d classes of level d, and each of them equals $O_{f,c}$ for exactly $q^{\lfloor d/2 \rfloor + 1}$ pairs (f, c).*

Proof. In characteristic 2, $f^2 = \sum a_i^2 t^{2i}$ has no odd terms, $a \mapsto a^2$ is bijective on $\mathbb{F}_q$, and the coefficient of t^d is shifted by c. This identifies the classes that occur.

For the count, note that $a \mapsto a^2$ is additive. So the map $(f, c) \mapsto \mathrm{prefix}(f^2 + ct^d)$ is a surjective homomorphism of additive groups from $\mathbb{F}_q^{d+1}$, the coefficients $(a_0, \dots, a_{d-1}, c)$, onto $\mathbb{F}_q^{\lceil d/2 \rceil}$. Its fibres therefore all have size $q^{d+1-\lceil d/2 \rceil} = q^{\lfloor d/2 \rfloor + 1}$. □

For example, the empty classes of Theorem 1.1 at $d = 1$, 2 and 4 are represented by q, q^2 and $2^3 = 8$ pairs (f, c) respectively.

Thus in characteristic 2 the Oppermann sub-intervals are the short intervals around the $q^{\lceil d/2 \rceil}$ "near-squares" $f^2 + ct^d$, a set of measure $q^{-\lfloor d/2 \rfloor}$ among all short intervals. Squares and near-squares carry Galois-theoretic structure, and that structure produces the counterexamples below.

2.1. **Newton–trace duality in characteristic 2.** Throughout this subsection $q = 2^k$, $K = \mathbb{F}_q$ and $\mathrm{Tr} = \mathrm{Tr}_{K/\mathbb{F}_2}$. We write $H_d \subset G_d$ for the set of Oppermann classes of level d, described in Lemma 2.2.

For monic $P = \prod_\alpha (t - \alpha)$ of degree n, write its class in G_d as $\sigma(P) = u^n P(1/u) = \prod_\alpha (1 + \alpha u) \equiv 1 + e_1 u + \cdots + e_d u^d \pmod{u^{d+1}}$. Since the signs disappear in characteristic 2, $e_j = p_{n-j}$ is the j-th elementary symmetric function of the roots, and $\sigma(PQ) = \sigma(P)\sigma(Q)$. Let $S_m(P) = \sum_\alpha \alpha^m \in \mathbb{F}_q$. Since

$$u\frac{\sigma'(P)}{\sigma(P)} = \sum_\alpha \frac{\alpha u}{1 + \alpha u} = \sum_{m \ge 1} S_m(P) u^m$$

in characteristic 2, S_m for $m \le d$ depends only on the class of P, and $S_m(PQ) = S_m(P) + S_m(Q)$. Moreover $S_{2m} = S_m^2$.

Lemma 2.3 (Newton's identities in characteristic 2)**.** *For $1 \le j \le d$, $S_j = \sum_{i=1}^{j-1} e_i S_{j-i} + j e_j$. In particular, for odd j, $e_j = S_j + \sum_{i=1}^{j-1} e_i S_{j-i}$.*

Proof. This is the usual identity with all signs equal to $+1$. □

Theorem 2.4. *Let $H_d = \{1 + \sum e_j u^j \in G_d : e_j = 0 \text{ for all odd } j < d\}$, so $|H_d| = q^{\lceil d/2 \rceil}$.*

(1) *For every class, $e_j = 0$ for all odd $j < d$ if and only if $S_j = 0$ for all odd $j < d$.*

(2) *The map*

$$\Phi : G_d \to (\mathbb{F}_q^{\lfloor d/2 \rfloor}, +), \qquad P \mapsto (S_1(P), S_3(P), \dots, S_{2\lfloor d/2 \rfloor - 1}(P))$$

is a surjective homomorphism with kernel H_d. Hence H_d is a subgroup and $G_d/H_d \cong \mathbb{F}_q^{\lfloor d/2 \rfloor}$, an elementary abelian 2-group.

Proof. (1) We argue by induction over odd $j < d$. Assume both vanishing conditions hold for all odd $i < j$. In the sum $\sum_{i<j} e_i S_{j-i}$ of Lemma 2.3, every term vanishes. If i is odd then $e_i = 0$. If i is even then $j - i$ is odd and less than j, so $S_{j-i} = 0$. Hence $e_j = S_j$, which gives the inductive step in both directions.

(2) Φ is a homomorphism because each S_m is additive. By (1), its kernel is H_d. Comparing orders, $|G_d/H_d| = q^{d-\lceil d/2 \rceil} = q^{\lfloor d/2 \rfloor}$, so Φ is surjective. □

The annihilator $H_d^\perp = \{\chi \in \widehat{G_d} : \chi|_{H_d} = 1\}$ is the dual of G_d/H_d. For $\beta = (\beta_1, \beta_3, \dots) \in \mathbb{F}_q^{\lfloor d/2 \rfloor}$ put

$$\chi_\beta(P) = (-1)^{\mathrm{Tr}(\sum_{j \text{ odd} < d} \beta_j S_j(P))}, \qquad g_\beta(x) = \sum_{j \text{ odd} < d} \beta_j x^j.$$

Theorem 2.5. *$H_d^\perp = \{\chi_\beta\}$, a group of $q^{\lfloor d/2 \rfloor}$ real characters. For every $m \geq 1$,*

$$\psi_m(\chi_\beta) := \sum_{\deg F = m} \Lambda(F)\chi_\beta(F) = \sum_{x \in \mathbb{F}_{q^m}} (-1)^{\mathrm{Tr}_{\mathbb{F}_{q^m}/\mathbb{F}_2}(g_\beta(x))}.$$

Proof. The first statement follows from Theorem 2.4(2). For the second, group the monic F of degree m with $\Lambda(F) \neq 0$ by their roots. Every $x \in \mathbb{F}_{q^m}$, of degree $s \mid m$ over K, determines $F_x = (\min_x)^{m/s}$, with $\Lambda(F_x) = s$; and each such F arises from exactly s elements x. By additivity, $S_j(F_x) = (m/s)\,\mathrm{Tr}_{\mathbb{F}_{q^s}/K}(x^j) = \mathrm{Tr}_{\mathbb{F}_{q^m}/K}(x^j)$. So $\chi_\beta(F_x) = (-1)^{\mathrm{Tr}_{\mathbb{F}_{q^m}/\mathbb{F}_2}(g_\beta(x))}$. The element $x = 0$ corresponds to $F = t^m$, whose class is $\sigma(t^m) = 1$; it has $\Lambda = 1$ and contributes $+1$ on both sides. No correction term is needed. □

Theorem 2.5 was checked against the exact class values for all $\beta \in \mathbb{F}_2^{\lfloor d/2 \rfloor}$, whenever $q^{2d} \leq 2^{16}$ (`verify_subgroup_mean.py`). Since $\Psi(x) = q^{-d} \sum_\chi \overline{\chi}(x)\psi_{2d}(\chi)$, the left side is $\sum_x \Psi(x)\chi_\beta(x)$.

3. Characteristic 2: the three obstructions and the $\mathrm{AGL}_r(\mathbb{F}_2)$ dead zone

Throughout this section $q = 2^k$ and $\mathrm{Tr} = \mathrm{Tr}_{\mathbb{F}_q/\mathbb{F}_2}$.

3.1. **(A) $d = 1$.** Here $f = t + a_0$ and $O_{f,c} = \{t^2 + ct + s_0 : s_0 \in \mathbb{F}_q\}$.

For $c = 0$ the members $t^2 + s_0 = (t + \sqrt{s_0})^2$ are squares.

For $c \neq 0$, substitute $t = cy$: $t^2 + ct + s_0 = c^2(y^2 + y + s_0/c^2)$. This is irreducible if and only if $y^2 + y + s_0/c^2$ has no root in $\mathbb{F}_q$, that is, if and only if $\mathrm{Tr}(s_0/c^2) = 1$. Exactly $q/2$ values of s_0 satisfy this.

3.2. **(B) $d = 2$: an additive polynomial and $\mathrm{AGL}_2(\mathbb{F}_2)$.** For $f = t^2 + a_1 t + a_0$ we have $f^2 = t^4 + a_1^2 t^2 + a_0^2$, so

$$O_{f,c} = B_e := \{t^4 + et^2 + \alpha t + \beta : \ \alpha, \beta \in \mathbb{F}_q\}, \qquad e = a_1^2 + c.$$

By Lemma 2.2 the blocks B_e, $e \in \mathbb{F}_q$, are precisely the Oppermann classes for $d = 2$.

Proposition 3.1. *Let $\alpha \neq 0$. Put $L(t) = t^4 + et^2 + \alpha t$ and $R(y) = y^3 + ey + \alpha$.*

1. *$t^4 + et^2 + \alpha t + \beta$ is irreducible for some β if and only if R has exactly one root in $\mathbb{F}_q$.*
2. *If R has exactly one root in $\mathbb{F}_q$, then $t^4 + et^2 + \alpha t + \beta$ is irreducible for exactly $q/2$ values of $\beta \in \mathbb{F}_q$.*

If $\alpha = 0$, every member $t^4 + et^2 + \beta = (t^2 + \sqrt{e}\,t + \sqrt{\beta})^2$ is a square.

Proof. L is $\mathbb{F}_2$-linear and separable ($L' = \alpha \neq 0$), so $V = \ker L \subset \overline{\mathbb{F}}_q$ is a 2-dimensional $\mathbb{F}_2$-space, and $V \setminus \{0\}$ is the set of roots of $L(t)/t = R(t)$. The Frobenius $\varphi(x) = x^q$ acts on V by some $F \in \mathrm{GL}(V) \cong S_3$. Exactly one of three cases holds: $F = 1$ (R has three roots in $\mathbb{F}_q$), F of order 3 (no root), or F a transvection (exactly one root).

Fix β and a root r of $P_\beta = L + \beta$. The roots of P_β form the coset $r + V$. Since $L(\varphi(r)) = \varphi(\beta) = \beta$, we have $\varphi(r) = r + v$ with $v \in V$, and $\varphi(r + w) = r + (Fw + v)$. So Frobenius acts on the roots through the affine map $T(w) = Fw + v$ of $V \cong \mathbb{F}_2^2$, an element of $\mathrm{AGL}_2(\mathbb{F}_2) \cong S_4$. P_β is irreducible if and only if T is a 4-cycle.

- If $F = 1$, T is a translation and $T^2 = 1$.
- If F has order 3, $F - 1$ is invertible, so T has a fixed point.
- If F is a transvection, $T^2(w) = w + (F+1)v$, so T is a 4-cycle if and only if $(F+1)v \neq 0$, i.e. $v \notin V^F$.

This proves the "only if" part of (1); the "if" part follows from (2), since $q/2 \geq 1$. For (2), let F be a transvection, so $(F+1)V = V^F$ is a line. Replacing r by $r + w$ replaces v by $v + (F+1)w$, so the class $[v] \in V/V^F \cong \mathbb{F}_2$ depends only on β, and $\beta \mapsto [v]$ is additive. Its kernel consists of the β for which some root $r' = r + w$ is fixed by φ, i.e. $\beta \in L(\mathbb{F}_q)$. The map $L : \mathbb{F}_q \to \mathbb{F}_q$ is $\mathbb{F}_2$-linear with kernel $V^F = V \cap \mathbb{F}_q$ of order 2, so $|L(\mathbb{F}_q)| = q/2$ and exactly $q/2$ values of β have $[v] \neq 0$. □

Lemma 3.2. *Let $\alpha \neq 0$. The cubic $R(y) = y^3 + ey + \alpha$ has exactly one root in $\mathbb{F}_q$ if and only if $\mathrm{Tr}(e^3/\alpha^2) \neq \mathrm{Tr}(1)$.*

Proof. The roots y_1, y_2, y_3 of R are distinct (R is separable as above), with $y_1 + y_2 + y_3 = 0$ and $y_1 y_2 y_3 = \alpha$. Put $\rho = y_1/y_3 + y_2/y_1 + y_3/y_2$ and $\rho' = y_2/y_3 + y_1/y_2 + y_3/y_1$. Even permutations of the roots fix ρ and odd ones exchange ρ, ρ'; using $y_i + y_j = y_k$,

$$\rho + \rho' = \frac{y_1 + y_2}{y_3} + \frac{y_2 + y_3}{y_1} + \frac{y_3 + y_1}{y_2} = 1.$$

Hence ρ, ρ' are the roots of $z^2 + z + \delta$ with $\delta = \rho\rho' \in \mathbb{F}_q$, and Frobenius acts on the roots of R by an odd permutation (a transposition: exactly one root in $\mathbb{F}_q$) if and only if $\rho \notin \mathbb{F}_q$, i.e. $\mathrm{Tr}(\delta) = 1$. Expanding,

$$\delta = 3 + \sum_k \frac{y_i y_j}{y_k^2} + \sum_k \frac{y_k^2}{y_i y_j} = 1 + \sum_k \frac{\alpha}{y_k^3} + \frac{1}{\alpha} \sum_k y_k^3.$$

From $y_k^3 = ey_k + \alpha$ we get $\sum_k y_k^3 = \alpha$, so $\delta = \sum_k \alpha/w_k$ with $w_k = y_k^3 = ey_k + \alpha$. Substituting $y = (w+\alpha)/e$ (for $e \neq 0$) into $w = y^3$ shows that the w_k are the roots of $w^3 + \alpha w^2 + (\alpha^2 + e^3)w + \alpha^3$, so $\sum_k 1/w_k = (\alpha^2 + e^3)/\alpha^3$ and

$$\delta = 1 + e^3/\alpha^2.$$

For $e = 0$ this also holds directly ($w_k = \alpha$). So $\mathrm{Tr}(\delta) = 1$ if and only if $\mathrm{Tr}(e^3/\alpha^2) \neq \mathrm{Tr}(1)$. □

Remark 3.3. δ is Berlekamp's characteristic-2 discriminant [2] of R, and R is the cubic resolvent of the quartic; the proof above is self-contained. For $e = 0$ the lemma says: $y^3 = \alpha$ has exactly one root iff k is odd, which is the familiar fact that cubing is bijective on $\mathbb{F}_q$ iff $3 \nmid q - 1$.

Proof of Theorem 1.1(B). By Proposition 3.1, $|B_e \cap \{\text{irreducibles}\}| = \frac{q}{2} \cdot \#\{\alpha \neq 0 : \mathrm{Tr}(e^3/\alpha^2) \neq \mathrm{Tr}(1)\}$. If $e = 0$ the condition reads $\mathrm{Tr}(1) = 1$, i.e. k odd: the count is $\frac{q}{2}(q-1)$ for k odd and 0 for k even. If $e \neq 0$, $\alpha \mapsto e^3/\alpha^2$ permutes $\mathbb{F}_q^\times$, and $\#\{x \neq 0 : \mathrm{Tr}(x) = 1\} = q/2$, $\#\{x \neq 0 : \mathrm{Tr}(x) = 0\} = q/2 - 1$; with $\mathrm{Tr}(1) = k \bmod 2$ this gives $q^2/4$ (k even) and $\frac{q}{2}(\frac{q}{2} - 1)$ (k odd). The count $q(q-2)/4$ vanishes only for $q = 2$. □

The formula was checked by brute force for $q \leq 64$ (script `char2_d2_theorem.py`): criterion of Lemma 3.2, the $q/2$ count of Proposition 3.1(2), and the four block totals.

3.3. **(C) $q = 2$, $d = 4$: the sign of Frobenius.** The Oppermann classes for $q = 2$, $d = 4$ are the four prefixes $(p_7, p_6, p_5, p_4) = (0, x, 0, y)$ (Lemma 2.2). For $x = y = 0$ the class is the *zero block*

$$Z_4 = \{t^8 + a_3t^3 + a_2t^2 + a_1t + a_0\} \subset \mathbb{F}_2[t],$$

equal to $O_{f,c}$ for $f = t^4+a_2t^2+a_1t+a_0$ and $c = a_2$. Here $a_3 = 0$ is forced, because the coefficient of t^6 in f^2 is a_3^2; this is why there are $2^3 = 8$ pairs rather than 16.

We use the theorem of Stickelberger [9] in the form of Swan [11, Cor. 1]: *if $\tilde{P} \in \mathbb{Z}[t]$ is monic of degree n and its reduction $P \in \mathbb{F}_2[t]$ is squarefree with r irreducible factors, then $r \equiv n \pmod 2$ if and only if* $\operatorname{disc}(\tilde{P}) \equiv 1 \pmod 8$. Equivalently, the Frobenius permutation of the roots of P is even if and only if $\operatorname{disc}(\tilde{P}) \equiv 1 \pmod 8$. An irreducible P of degree 8 has Frobenius an 8-cycle, an odd permutation, and would need $\operatorname{disc}(\tilde{P}) \equiv 5 \pmod 8$.

Lemma 3.4. *Let $\tilde{P} = t^n + \tilde{g}$ with $\tilde{g} \in \mathbb{Z}[t]$, $\deg \tilde{g} < n$, and $8 \mid n$. Then* $\operatorname{disc}(\tilde{P}) \equiv \operatorname{Res}(\tilde{P}, \tilde{g}') \pmod 8$.

Proof. $\operatorname{disc}(\tilde{P}) = (-1)^{n(n-1)/2} \operatorname{Res}(\tilde{P}, \tilde{P}') = \prod_{\tilde{P}(\rho)=0}(n\rho^{n-1} + \tilde{g}'(\rho))$, the sign being $+1$ for $8 \mid n$. Expanding the product, every term containing a factor n is n times a symmetric polynomial with integer coefficients in the roots, hence lies in $8\mathbb{Z}$. □

Proof of Theorem 1.1(C). An irreducible member of Z_4 has no root in $\mathbb{F}_2$, so $a_0 = 1$ and $a_1 + a_2 + a_3 = 1$. This leaves four polynomials. $t^8 + t^2 + 1 = (t^4 + t + 1)^2$ is a square. For the other three choose integer lifts and apply Lemma 3.4:

- $t^8 + t + 1$: $\tilde{g}' = 1$, disc $\equiv 1$;
- $t^8 + t^3 + 1$: $\tilde{g}' = 3t^2$, $\operatorname{Res}(\tilde{P}, 3t^2) = 3^8\big(\prod \rho\big)^2 = 3^8 \equiv 1$;
- $t^8 + t^3 + t^2 + t + 1$, lifted as $t^8 + (t + 1)^3$: $\tilde{g}' = 3(t + 1)^2$, and $\operatorname{Res}(\tilde{P}, 3(t + 1)^2) = 3^8 \prod(\rho + 1)^2 = 3^8\tilde{P}(-1)^2$, with $\tilde{P}(-1) = (-1)^8 + 0^3 = 1$, so disc $\equiv 3^8 \equiv 1$.

Suppose one of these three were irreducible over $\mathbb{F}_2$. Then it would be squarefree with $r = 1$, and Stickelberger–Swan would require $\operatorname{disc}(\tilde{P}) \equiv 5 \pmod 8$. That contradicts $\operatorname{disc}(\tilde{P}) \equiv 1$, so none of them is irreducible. The argument does not need squarefreeness to be checked beforehand. □

Remark 3.5. The mechanism is that $8 \mid \deg P$ kills the leading term of P' modulo 8, so the sign of Frobenius is decided by the tail g alone; with $\deg g \le 3$ the tail is too short to produce the sign -1 on a polynomial with no linear factor. Table 1 shows the zero blocks $Z_d = \{t^{2d} + s : \deg s < d\}$, which are Oppermann classes for every d. For $d = 8$ ($8 \mid 2d$ again) the tail has degree ≤ 7, half of the squarefree members already have the sign -1, and the block contains 10 irreducibles. The parallel with (B) is exact: in (B) the obstruction is the quotient $S_4 \to S_3$ (the resolvent cubic), in (C) the quotient $S_8 \to \{\pm 1\}$ (the sign).

3.4. **Additive slices for $d = 2^{r-1} \ge 4$ and $\mathrm{AGL}_r(\mathbb{F}_2)$.** For $d = 2^{r-1}$, i.e. $2d = 2^r$, consider the affine polynomials

$$P_\beta(t) = L(t) + \beta, \qquad L(t) = t^{2^r} + \sum_{i=0}^{r-1} c_i t^{2^i}, \qquad c_i, \beta \in \mathbb{F}_q.$$

Apart from t, every monomial has even degree. In particular no odd power of t lies strictly between t^d and t^{2d}, so every such P_β lies in a class of H_d. For $r = 1, 2$ (that is, $d = 1, 2$) the Oppermann classes of §§3.1–3.2 consist of such affine cosets. There, Frobenius acts on the roots through $\mathrm{AGL}_r(\mathbb{F}_2)$, and irreducibility amounts to a 2^r-cycle. For $r \ge 3$ this mechanism disappears.

Theorem 3.6. (1) *Every element $T \in \mathrm{AGL}_r(\mathbb{F}_2)$ of 2-power order satisfies $T^{2^m} = \mathrm{id}$, where $m = \lceil \log_2(r + 1) \rceil$.*

(2) *$\mathrm{AGL}_r(\mathbb{F}_2)$ contains a 2^r-cycle on $\mathbb{F}_2^r$ if and only if $r \in \{1, 2\}$.*

TABLE 1. The zero blocks $Z_d = \{t^{2d} + s : \deg s < d\} \subset \mathbb{F}_2[t]$. Columns: the squarefree members; those with disc $\equiv 5 \pmod 8$, i.e. Frobenius odd (integer lift with coefficients $0, 1$); those among them with no root in $\mathbb{F}_2$; and the irreducibles. For $d = 4$ the two members with disc $\equiv 5$ are t^8+t^3+t and $t^8+t^3+t^2+1$, which have the roots 0 and 1. Stickelberger–Swan holds on every squarefree member (script `swan_parity.py`).

d	$\lvert Z_d\rvert$	squarefree	disc $\equiv 5$	disc $\equiv 5$, no root	irreducible
2	4	2	2	1	1
3	8	4	3	1	1
4	16	8	2	**0**	**0**
5	32	16	10	4	4
6	64	32	20	8	6
7	128	64	42	17	9
8	256	128	64	24	10
9	512	256	130	61	28
10	1024	512	280	120	49

Proof. (1) Write $T(w) = Fw + v$. Since T has 2-power order, so does $F \in \mathrm{GL}_r(\mathbb{F}_2)$. Hence $F^{2^s} = I$ for some s, so $(F+I)^{2^s} = 0$, and $N = F+I$ is nilpotent. By Cayley–Hamilton, $N^r = 0$. In characteristic 2,

$$F^{2^m} = (I+N)^{2^m} = I + N^{2^m}, \qquad I + F + \cdots + F^{2^m-1} = \frac{F^{2^m}+I}{F+I} = N^{2^m-1},$$

the second identity holding as polynomials in F. Therefore

$$T^{2^m}(w) = w + N^{2^m} w + N^{2^m-1} v.$$

If $2^m - 1 \geq r$, both powers of N vanish and $T^{2^m} = \mathrm{id}$.

(2) A 2^r-cycle has order 2^r. If $r \geq 3$, then $2^{r-1} - 1 \geq r$, so the computation in (1), applied with $m = r-1$, shows that every element of 2-power order satisfies $T^{2^{r-1}} = \mathrm{id}$, a contradiction. Conversely, for $r = 1$ a translation is a 2-cycle, and for $r = 2$ a transvection with a suitable translation part is a 4-cycle (Proposition 3.1). □

Corollary 3.7 (additive dead zone)**.** *Let $r \geq 3$. No affine polynomial $L(t) + \beta$ of degree 2^r over $\mathbb{F}_{2^k}$ is irreducible, for any k. In particular, for $d = 2^{r-1} \geq 4$, the affine members of the Oppermann classes in H_d are all reducible, and every irreducible polynomial in H_d has a non-additive shape.*

Proof. If $c_0 = 0$, then $L(t) + \beta$ is a square, $\big(t^{2^{r-1}} + \sum_{i \geq 1} \sqrt{c_i}\, t^{2^{i-1}} + \sqrt{\beta}\big)^2$.

If $c_0 \neq 0$, then L is separable, and its roots form an r-dimensional $\mathbb{F}_2$-space V. The roots of $L + \beta$ form a coset $\rho + V$. Since $L(\varphi(\rho)) = \beta$, we have $\varphi(\rho) = \rho + v$ with $v \in V$, so Frobenius φ acts on the coset by $w \mapsto Fw + v$, an element of $\mathrm{AGL}_r(\mathbb{F}_2)$ (as in Proposition 3.1). Irreducibility would require a 2^r-cycle, which Theorem 3.6 excludes. □

Both statements were checked by brute force (`verify_additive_slices.py`).

- $\mathrm{AGL}_3(\mathbb{F}_2)$ (order 1344) and $\mathrm{AGL}_4(\mathbb{F}_2)$ (order 322560) contain no 2^r-cycle. Their largest 2-power orders are 4 and 8, so the bound $2^{\lceil \log_2(r+1) \rceil}$ is attained.
- Among the affine polynomials of degree 8 over $\mathbb{F}_q$, $q \leq 16$, and of degree 16 over $\mathbb{F}_2$ and $\mathbb{F}_4$, none is irreducible: for instance 0 of 65536 for degree 8 over $\mathbb{F}_{16}$. In degrees 2 and 4 the irreducible ones make up roughly half and a quarter, as Theorem 1.1(B) predicts.

Hence the exact AGL-counts behind Theorem 1.1(B) have no analogue at $d = 4, 8, \ldots$. There, the Oppermann classes are nonempty (Theorem 1.3; Table 6) only because of non-additive members.

4. Odd characteristic, $d = 2$: a ternary trace form

Throughout this section q is odd and η is the quadratic character of $\mathbb{F}_q$, with $\eta(0) = 0$.

4.1. **Reduction to one-parameter blocks.** The level-2 classes in degree 4 are $C_{p_3,p_2} = \{t^4 + p_3t^3 + p_2t^2 + \alpha t + \beta : \alpha, \beta \in \mathbb{F}_q\}$. The translation $t \mapsto t - p_3/4$ is defined because 4 is invertible (in characteristic 3 it is $t \mapsto t - p_3$). It preserves irreducibility and maps C_{p_3,p_2} bijectively onto

$$B_e = \{t^4 + et^2 + \alpha t + \beta : \alpha, \beta \in \mathbb{F}_q\}, \qquad e = p_2 - \tfrac{3}{8}p_3^2.$$

For $O_{f,c}$ with $f = t^2 + a_1t + a_0$ we have $p_3 = 2a_1$ and $p_2 = a_1^2 + 2a_0 + c$, so $e = 2a_0 + c - a_1^2/2$.

4.2. **The count.**

Proposition 4.1. *For every $e \in \mathbb{F}_q$, the number of irreducibles in B_e is*

$$\pi(B_e) = \frac{q^2 - 1 - (q - \eta(-1))\eta(2e)}{4}.$$

Proof. A monic irreducible quartic $P \in B_e$ is the minimal polynomial of exactly four elements $x \in \mathbb{F}_{q^4}$ of degree 4, namely its roots. Write $\mathrm{Tr} = \mathrm{Tr}_{\mathbb{F}_{q^4}/\mathbb{F}_q}$. The coefficient conditions on P translate into conditions on x: $p_3 = -\mathrm{Tr}(x)$, and when $\mathrm{Tr}\, x = 0$,

$$p_2 = e_2(x, x^q, x^{q^2}, x^{q^3}) = \tfrac{1}{2}((\mathrm{Tr}\, x)^2 - \mathrm{Tr}(x^2)) = -\tfrac{1}{2}\mathrm{Tr}(x^2).$$

Hence $4\pi(B_e) = N - M$, where

$$N = \#\{x \in W : Q(x) = -2e\}, \qquad W = \ker \mathrm{Tr}, \quad Q(x) = \mathrm{Tr}(x^2),$$

and M counts the elements of W with $Q(x) = -2e$ that have degree < 4 over $\mathbb{F}_q$, i.e. that lie in $\mathbb{F}_{q^2}$.

The form Q. The bilinear form $(x, y) \mapsto \mathrm{Tr}(xy)$ on $\mathbb{F}_{q^4}$ is nondegenerate. Its Gram determinant on a basis $\omega_1, \dots, \omega_4$ is $D = \det(\omega_i^{q^j})^2$. Frobenius permutes the columns of the matrix $(\omega_i^{q^j})$ by a 4-cycle, an odd permutation, so $\det(\omega_i^{q^j})$ is not fixed by Frobenius. It therefore lies outside $\mathbb{F}_q$ while its square D lies in $\mathbb{F}_q$, so D is a nonsquare in $\mathbb{F}_q$.

Since $\mathrm{Tr}(1 \cdot x) = \mathrm{Tr}\, x$, the line $\mathbb{F}_q$ is orthogonal to W, and $\mathbb{F}_{q^4} = \mathbb{F}_q \perp W$. The Gram determinant of $\mathbb{F}_q$ is $\mathrm{Tr}(1) = 4$, a square. So $Q|_W$ is a nondegenerate ternary form whose determinant Δ is a nonsquare.

Counting N. For a nondegenerate quadratic form in 3 variables over $\mathbb{F}_q$ with determinant Δ, the number of solutions of $Q(x) = b$ is $q^2 + q\,\eta(-b\Delta)$ [7, Thm. 6.27]; for $b = 0$ this is q^2. With $b = -2e$ and $\eta(\Delta) = -1$ we get $N = q^2 - q\,\eta(2e)$, also for $e = 0$.

Counting M. An element of $\mathbb{F}_{q^2}$ has $\mathrm{Tr}\, x = 2\,\mathrm{Tr}_{\mathbb{F}_{q^2}/\mathbb{F}_q}(x)$, so it lies in W if and only if $x^q = -x$. Such x form the line $\delta\mathbb{F}_q$, where $\delta^2 = \nu$ is a nonsquare of $\mathbb{F}_q$. For these x, $x^2 \in \mathbb{F}_q$ and $Q(x) = 2(x^2 + x^{2q}) = 4x^2$, so the condition is $x^2 = -e/2$. The squares of elements $s\delta$ are 0 and the nonsquares of $\mathbb{F}_q$, each nonsquare attained twice. Hence $M = 1$ if $e = 0$, and $M = 1 - \eta(-e/2)$ if $e \neq 0$; in both cases $M = 1 - \eta(-2e)$.

Therefore

$$4\pi(B_e) = q^2 - q\,\eta(2e) - 1 + \eta(-2e) = q^2 - 1 - (q - \eta(-1))\eta(2e). \qquad \square$$

Proof of Theorem 1.4. Combine the reduction with Proposition 4.1. The minimum over e is attained at $\eta(2e) = 1$. It equals $(q^2 - 1 - (q-1))/4 = q(q-1)/4$ when $\eta(-1) = 1$, i.e. $q \equiv 1 \pmod 4$, and $(q^2 - 1 - (q+1))/4 = (q+1)(q-2)/4$ when $q \equiv 3 \pmod 4$. Both are positive for every odd q. $\square$

Remark 4.2. Brute-force counts confirm the formula for every block B_e when $q = 3$, 5, 7, 9, 11, 13, 25, 27, 49, 81, and for every class C_{p_3,p_2} when $q \leq 13$ (script `odd_d2_formula.py`). They also agree with the minima of Table 3. Characteristic 3 needs no special treatment.

The contrast with characteristic 2 is instructive. There the translation $t \mapsto t + a$ cannot remove p_3, since $4 = 0$. The Oppermann blocks are exactly the classes with $p_3 = 0$, on which $t^4 + et^2 + \alpha t$ is *additive*, and this additivity produces the obstruction of Proposition 3.1. In odd

characteristic the trace form $\mathrm{Tr}(x^2)$ has the same shape for every e, and the count never drops below about $q^2/4 - q/4$.

5. Characteristic 2, $d = 3$: field-tower count and cubic Gauss sums

Let $q = 2^k$. By Lemma 2.2, the q^2 Oppermann classes for $d = 3$ are

$$C_{e,c} = \{t^6 + et^4 + ct^3 + s_2t^2 + s_1t + s_0 : \ s_2, s_1, s_0 \in \mathbb{F}_q\}, \qquad e, c \in \mathbb{F}_q.$$

Write $C_e = C_{e,0}$. Every member of $C_{e,0}$ has $P' = s_1$, so [1] does not apply to these classes (§6.4). The classes with $c \neq 0$ are covered by [1] for large q, but the count below makes this unnecessary.

Proposition 5.1. (1) *For each c, the number $\pi(C_{e,c})$ does not depend on e.*
(2) *$E(C_{e,0}) = q^2$, so $\Psi(C_{e,0}) = 6\pi(C_{e,0}) + q^2$. For $c \neq 0$, $E(C_{e,c}) = 0$.*

Proof. (1) In characteristic 2 we have $(t+a)^6 = (t^4+a^4)(t^2+a^2) = t^6 + a^2t^4 + a^4t^2 + a^6$ and $(t+a)^4 = t^4 + a^4$. Neither has a t^3 term. On the other hand, $c(t+a)^3 = ct^3 + cat^2 + ca^2t + ca^3$. So $t \mapsto t + a$ maps $C_{e,c}$ bijectively onto $C_{e+a^2,c}$ and preserves irreducibility. Since $a \mapsto a^2$ is bijective, the claim follows.

(2) We count the proper prime powers of degree 6 that lie in some $C_{e,c}$. Sixth powers of linear polynomials and squares of cubics contain only even powers of t. Cubes of irreducible quadratics never lie in any $C_{e,c}$, as shown below. So no such prime power has a t^3 term, and $E(C_{e,c}) = 0$ for $c \neq 0$. Now take $c = 0$ and, by (1), $e = 0$.

- *Sixth powers.* $(t+a)^6 \in C_0$ only for $a = 0$; this is t^6, with $\Lambda = 1$.
- *Cubes.* For a quadratic $\varpi = t^2 + b_1t + b_0$, the coefficient of t^5 in ϖ^3 is b_1. So $\varpi^3 \in C_0$ forces $b_1 = 0$, but then $\varpi = (t + \sqrt{b_0})^2$ is not irreducible.
- *Squares.* For a cubic $\varpi = t^3 + a_2t^2 + a_1t + a_0$ we have $\varpi^2 = t^6 + a_2^2t^4 + a_1^2t^2 + a_0^2$, which lies in C_0 exactly when $a_2 = 0$. Translation $t \mapsto t + a$ changes a_2 by $3a = a$, so the $(q^3 - q)/3$ irreducible cubics are equidistributed over the values of a_2, and $(q^2 - 1)/3$ of them have $a_2 = 0$. Each contributes $\Lambda = 3$.

Hence $E(C_0) = 1 + (q^2 - 1) = q^2$. The cubes are excluded from every $C_{e,c}$, not only from C_0, since all of these classes have $p_5 = 0$. □

Theorem 5.2. *Let $q = 2^k$ and $e, c \in \mathbb{F}_q$.*

(1) *If k is odd, then $\pi(C_{e,c}) = q(q^2 - 1)/6$ for all e, c.*
(2) *If k is even, then*

$$\pi(C_{e,c}) = \frac{q(q-1)^2}{6} + q\, m(c), \qquad m(c) = \#\{\nu \in \mathbb{F}_q^\times \setminus (\mathbb{F}_q^\times)^3 : \ \mathrm{Tr}(c/\nu) = 1\}.$$

Here $m(0) = 0$, and $(q+2)/6 \leq m(c) \leq q/2$ for $c \neq 0$.

In particular, every Oppermann sub-interval with $d = 3$ in characteristic 2 contains at least $q(q-1)^2/6$ irreducibles if k is even and $q(q^2-1)/6$ if k is odd; this is positive for every q. For $c = 0$, $\Psi(C_e) = q^3 - (-1)^k(q^2 - q)$.

Proof. Let $K = \mathbb{F}_q \subset E = \mathbb{F}_{q^3} \subset L = \mathbb{F}_{q^6}$. Fix c. By Proposition 5.1(1), $6q\,\pi(C_{e,c})$ is the number of $x \in L$ of degree 6 over K whose characteristic polynomial $t^6 + e_1t^5 + e_2t^4 + e_3t^3 + \dots$ has $e_1 = 0$ and $e_3 = c$.

Step 1 (conditions on x). With $S_m = \mathrm{Tr}_{L/K}(x^m)$, Newton's identity in characteristic 2 reads $S_3 = e_1S_2 + e_2S_1 + e_3$. So for $e_1 = S_1 = 0$ we get $e_3 = S_3$. The conditions are therefore $\mathrm{Tr}_{L/K}(x) = 0$ and $\mathrm{Tr}_{L/K}(x^3) = c$.

Step 2 (the relative trace u). Put $u = \mathrm{Tr}_{L/E}(x) = x + x^{q^3} \in E$; then $\mathrm{Tr}_{L/K}(x) = \mathrm{Tr}_{E/K}(u)$. Assume $\mathrm{Tr}_{E/K}(u) = 0$. We claim that x has degree 6 if and only if $u \neq 0$.

- If $u = 0$, then $x^{q^3} = x$, so $x \in E$.
- If $x \in \mathbb{F}_{q^2}$, then $x^{q^3} = x^q$, so $u = x + x^q \in K$, and $0 = \mathrm{Tr}_{E/K}(u) = 3u = u$.

Since the proper subfields of L are contained in E or in $\mathbb{F}_{q^2}$, this proves the claim.

Step 3 (the fibre over u). Fix $u \in E^\times$ with $\mathrm{Tr}_{E/K}(u) = 0$. The q^3 elements x with $\mathrm{Tr}_{L/E}(x) = u$ are $x = u\xi$ with $\xi + \xi^{q^3} = 1$. For these,

$$\mathrm{Tr}_{L/E}(\xi^3) = \xi^3 + (\xi+1)^3 = \xi^2 + \xi + 1.$$

Put $\alpha = \xi^2 + \xi$; then $\alpha^{q^3} = \alpha$, so $\alpha \in E$. As ξ runs over the fibre, α runs 2-to-1 over $\{\alpha \in E : \mathrm{Tr}_{E/\mathbb{F}_2}(\alpha) = 1\}$: the equation $\xi^2 + \xi = \alpha$ has its roots outside E exactly when $\mathrm{Tr}_{E/\mathbb{F}_2}(\alpha) = 1$, and then the two roots are ξ and $\xi + 1 = \xi^{q^3}$. Finally,

$$\mathrm{Tr}_{L/K}(x^3) = \mathrm{Tr}_{E/K}(u^3(\alpha+1)).$$

Step 4 (counting in the fibre). The condition $\mathrm{Tr}_{L/K}(x^3) = c$ says that α lies in the affine K-hyperplane

$$H_{u,c} = \{\alpha \in E : \mathrm{Tr}_{E/K}(u^3\alpha) = \mathrm{Tr}_{E/K}(u^3) + c\}, \qquad |H_{u,c}| = q^2.$$

The $\mathbb{F}_2$-linear functionals on E that vanish on the direction space of $H_{u,c}$ are $\alpha \mapsto \mathrm{Tr}_{E/\mathbb{F}_2}(\lambda u^3 \alpha)$ with $\lambda \in K$. Hence $\mathrm{Tr}_{E/\mathbb{F}_2}$ is constant on $H_{u,c}$ if and only if $u^3 \in K^\times$.

- If $u^3 \notin K^\times$, then $\mathrm{Tr}_{E/\mathbb{F}_2}(\alpha) = 1$ on exactly $q^2/2$ points of $H_{u,c}$. This gives q^2 elements x, whatever c is.
- If $u^3 = \nu \in K^\times$, then on $H_{u,c}$ we have $\nu\,\mathrm{Tr}_{E/K}(\alpha) = \nu\,\mathrm{Tr}_{E/K}(1) + c = \nu + c$. So $\mathrm{Tr}_{E/\mathbb{F}_2}(\alpha) = \mathrm{Tr}_{K/\mathbb{F}_2}(1 + c/\nu)$ throughout $H_{u,c}$. This gives $2q^2$ elements x if that value is 1, and none otherwise.

Step 5 (summing over u). There are $q^2 - 1$ admissible u.

- *k odd.* Here $3 \nmid q^3 - 1$, so cubing is bijective on $E^\times$ and on $K^\times$, and $u^3 \in K^\times$ forces $u \in K^\times$. But every $u \in K^\times$ has $\mathrm{Tr}_{E/K}(u) = 3u = u \neq 0$, so it is not admissible. All $q^2 - 1$ admissible u therefore contribute q^2, for every c, and $6q\,\pi = (q^2-1)q^2$.
- *k even.* Here $3 \mid q-1$, $3 \mid q^2+q+1$ and $\mathrm{Tr}_{K/\mathbb{F}_2}(1) = 0$. We sort the u with $u^3 = \nu \in K^\times$ by ν.
  - If $\nu \in (K^\times)^3$, its three cube roots lie in $K^\times$. They are not admissible, since $\mathrm{Tr}_{E/K}(u) = u \neq 0$.
  - If $\nu \notin (K^\times)^3$, its three cube roots lie in $E \setminus K$, because $K^\times \subset (E^\times)^3$. They have minimal polynomial $X^3 - \nu$, hence trace 0, and are admissible. There are $2(q-1)$ of them, and each contributes $2q^2$ when $\mathrm{Tr}_{K/\mathbb{F}_2}(c/\nu) = 1$ and nothing otherwise.
  - The remaining $(q^2-1) - 2(q-1) = (q-1)^2$ admissible u contribute q^2 each.

  So $6q\,\pi = (q-1)^2q^2 + 3 \cdot 2q^2\,m(c)$, which is the formula claimed.

  For the bounds on $m(c)$ when $c \neq 0$: as ν runs over the non-cubes, c/ν runs over $K^\times$ minus the coset $c(K^\times)^3$, which has $(q-1)/3$ elements. Since $K^\times$ contains $q/2$ elements of trace 1, we get $m(c) = q/2 - t(c)$ with $t(c) = \#\{y \in c(K^\times)^3 : \mathrm{Tr}(y) = 1\} \in [0, (q-1)/3]$.

The formula for $\Psi(C_e)$ follows from Proposition 5.1(2). □

5.1. **The cube-residue count $m(c)$.** It remains to evaluate $m(c)$ in Theorem 5.2(2).

Theorem 5.3. *Let k be even, $\varepsilon = (-1)^{k/2}$, and $c \in K^\times$. Then*

$$m(c) = \begin{cases} \dfrac{q - \varepsilon\sqrt{q}}{3}, & c \in (K^\times)^3, \\ \dfrac{2q + \varepsilon\sqrt{q}}{6}, & c \notin (K^\times)^3. \end{cases}$$

Proof. As ν runs over the non-cubes, $y = c/\nu$ runs over $K^\times \setminus cC$, where $C = (K^\times)^3$. Hence $m(c) = q/2 - t(c)$ with $t(c) = \#\{y \in cC : \mathrm{Tr}\, y = 1\}$.

Let ω be a cubic character of $K^\times$ and $\psi_0(y) = (-1)^{\mathrm{Tr}\, y}$. The indicator of cC is

$$\mathbf{1}_{cC}(y) = \tfrac{1}{3}\textstyle\sum_{i=0}^{2} \omega^i(y)\overline{\omega}^i(c),$$

TABLE 2. Every characteristic-2 Oppermann class of degree 6: the number of irreducibles in $C_{e,c}$ (Theorem 5.2, Corollary 5.4). For k odd all classes have the same count. For $q \le 256$ every class was also counted exactly with the Hayes FFT; for $q \ge 1024$ the two values of $m(c)$ were checked directly.

q	ε	$c=0$	c cube	c non-cube	$q^3/6$
2	–		1		1
4	−1	6	14	10	11
8	–		84		85
16	+1	600	664	696	683
32	–		5 456		5 461
64	−1	42 336	43 872	43 616	43 691
128	–		349 504		349 525
256	+1	2 774 400	2 794 880	2 796 928	2 796 203
1 024	−1	178 607 616	178 968 064	178 951 680	178 956 971
4 096	+1	11 447 654 400	11 453 159 424	11 453 290 496	11 453 246 123
16 384	−1	732 918 276 096	733 008 453 632	733 007 405 056	733 007 751 851

so

$$\sum_{y\in cC}\psi_0(y) = \tfrac{1}{3}\big(-1+\overline{\omega}(c)G(\omega)+\omega(c)G(\overline{\omega})\big), \qquad G(\omega)=\sum_{y\in K^\times}\omega(y)\psi_0(y).$$

Since $2\equiv -1 \pmod 3$, the Gauss sum is pure [10], [3, §11.6]: $G(\omega)=G(\overline{\omega})=-\varepsilon\sqrt{q}$. Hence

$$\sum_{y\in cC}\psi_0(y) = \tfrac{1}{3}\big(-1-2\varepsilon\sqrt{q}\,\mathrm{Re}\,\omega(c)\big),$$

where $2\,\mathrm{Re}\,\omega(c)$ equals 2 for a cube and -1 otherwise. Since $t(c)=\frac{1}{2}\big(|cC|-\sum_{y\in cC}\psi_0(y)\big)$ and $|cC|=(q-1)/3$, we get $t(c)=(q+2\varepsilon\sqrt{q})/6$ for a cube and $(q-\varepsilon\sqrt{q})/6$ for a non-cube. Then $m(c)=q/2-t(c)$. □

Corollary 5.4. *Let $q=2^k$. For every characteristic-2 Oppermann class of degree 6:*

$$\pi(C_{e,c}) = \begin{cases} q(q^2-1)/6, & k \text{ odd},\\ q(q-1)^2/6, & k \text{ even},\ c=0,\\ q(q^2+1-2\varepsilon\sqrt{q})/6, & k \text{ even},\ c\in(K^\times)^3,\\ q(q^2+1+\varepsilon\sqrt{q})/6, & k \text{ even},\ c\notin(K^\times)^3. \end{cases}$$

Moreover $\Psi(C_{e,c})=6\pi(C_{e,c})+q^2[c=0]$.

Proof. Insert Theorem 5.3 into Theorem 5.2. The prime-power term is Proposition 5.1(2). □

Thus at $d=3$, k even, the counts fluctuate around $q^3/6$ by $\pm\varepsilon q^{3/2}/6$ and $\mp\varepsilon q^{3/2}/3$. This is a square-root fluctuation whose sign alternates with $k/2$, the sign of the Gauss sum. It matches the supersingular pattern $\gamma^6\in q^3\{\pm1,\pm i\}$ observed for the level-3 zeros in Remark 5.5. Theorem 5.3 was checked for every c when $q\le 1024$, and for $c=1$ and one non-cube up to $q=2^{14}$ (Table 2; Figure 3, right).

Remark 5.5. For $c=0$ the error term $\pi(C_e)-q^3/6$ is $-q/6$ for k odd and $-q^2/3+q/6$ for k even; for $c\neq 0$ and k even it is of order $q^{3/2}$ (Corollary 5.4). This is far smaller than the $O(q^{5/2})$ a monodromy argument would give.

It reflects a supersingularity that we observed numerically: for $q\le 16$, every character of exact level 2 or 3 of G_3 has an L-function whose inverse zeros satisfy $\gamma^6\in q^3\{\pm1,\pm i\}$. Figure 2 (right) shows these zeros for $q=8$ next to the odd case $q=7$. At level 2 this is a theorem: Proposition 6.1(3) gives $\gamma_\chi^4=(-1)^k q^2$ for every character of exact level 2, which implies $\gamma_\chi^6\in\{\pm q^3,\pm iq^3\}$. At level 3 it is proved for the $2q(q-1)/3$ characters of Proposition 7.6 when k is even, and otherwise remains an observation. The proof above avoids these L-functions

altogether. Summing over e replaces the level-3 condition by the two trace conditions $\mathrm{Tr}_{L/K}(x) = 0$ and $\mathrm{Tr}_{L/K}(x^3) = c$, which the tower $K \subset E \subset L$ makes linear.

6. The exact computation, the square-root barrier, and large q

6.1. **Method.** Here Λ is the von Mangoldt function: $\Lambda(F) = \deg \varpi$ if $F = \varpi^k$ for a monic irreducible ϖ and some $k \geq 1$, and $\Lambda(F) = 0$ otherwise. For a class x of level l and degree n, let $\Psi(x) = \sum_{F \in x} \Lambda(F)$. The explicit formula [5, 6] reads

$$\Psi(x) = q^{n-l} + q^{-l} \sum_{\chi \neq 1} \overline{\chi}(x)\, \psi_n(\chi), \qquad u \frac{L'(u,\chi)}{L(u,\chi)} = \sum_{m \geq 1} \psi_m(\chi) u^m, \tag{2}$$

where for $\chi \neq 1$ the L-function $L(u,\chi) = \sum_F \chi(F) u^{\deg F}$ is a polynomial of degree $\leq l-1$. Its coefficients are the Fourier transforms on G_l of the class distributions of the monic polynomials of degree $< l$; $\psi_n(\chi)$ follows by the Newton recursion, and one inverse FFT on G_l returns $\Psi(x)$ for all q^l classes at once. This is the algorithm of [4], run here at level $l = d$ and degree $n = 2d$. The number of irreducibles in x is

$$\pi(x) = \frac{1}{n}\big(\Psi(x) - E(x)\big), \qquad E(x) = \sum_{k \geq 2,\ k \mid n} \ \sum_{\deg \varpi = n/k,\ \varpi^k \in x} \frac{n}{k}.$$

The proper prime powers in $E(x)$ are enumerated directly (Ben-Or test in degree $\leq d$) and placed by the rule $\mathrm{class}(\varpi^k) = \mathrm{class}(\varpi)^k$ in G_l. Each run checks that every $\pi(x)$ is an integer and that $\sum_x (n\pi(x) + E(x)) = q^n$, and reports the largest distance of the floating-point $\Psi(x)$ from an integer (below 10^{-7} in all runs). As an independent check, a brute-force census that tests every polynomial of degree $2d$ for irreducibility (`verify_pointwise_oppermann.py`) agrees with the FFT counts class by class for all $q^{2d} \leq 2^{22}$ (`compare_bruteforce.py`), and in minimum, mean and empty classes for all $q^{2d} \leq 2^{26}$.

6.2. **Results.** Table 3 lists, for odd q, the minimum of $\pi(x)$ over all classes (all of which are Oppermann sub-intervals by Lemma 2.1) and its ratio to the expected value $q^d/2d$. Table 4 gives the same for the Oppermann classes in characteristic 2. The full output, all degrees, is in the file
`data/hayes_oppermann_census.txt`.

Proof of Theorems 1.2 and 1.3. Exact computation as described above; the minima are in Tables 3 and 4, and the empty classes in characteristic 2 occur exactly at

$$(q,d) \in \{(2,1),(2,2),(2,4),(4,1),(4,2),(8,1),(16,1),(16,2)\},$$

one class each, which are the classes of Theorem 1.1. □

The normalised minimum $\min_x \pi(x)/(q^d/2d)$ tends to 1 in every column: the worst Oppermann sub-interval is only slightly below average, far from the Weil bound discussed next.

6.3. **The square-root barrier.** In (2) with $n = 2d$ and $l = d$ the main term is q^d.

Let χ have exact level j. The monic polynomials of degree $\geq j$ are equidistributed over G_j, so $\deg L(u,\chi) \leq j-1$. By Weil, every inverse zero γ of $L(u,\chi)$ has $|\gamma| \in \{1, \sqrt{q}\}$, hence $|\psi_{2d}(\chi)| \leq (j-1)q^d$. Characters of level 1 have $L = 1$ and contribute nothing. Multiplying by the factor q^{-d} in (2), which cancels the q^d, and summing over the $q^j - q^{j-1}$ characters of each exact level j gives

$$|\Psi(x) - q^d| \leq \sum_{j=2}^{d} (j-1)(q^j - q^{j-1}) = (d-1)q^d - \sum_{j=1}^{d-1} q^j. \tag{3}$$

For $d \geq 3$ the right-hand side exceeds q^d for every q, so (3) does not even give $\Psi(x) > 0$. This is the exact analogue of the situation over $\mathbb{Z}$: Oppermann's intervals have length $\sqrt{x}$, and the Riemann hypothesis falls short by a factor $\log x$. No degree threshold beyond which $O_{f,c} \neq \emptyset$ follows from Weil's theorem.

TABLE 3. Odd q: irreducible counts π over all q^d Hayes classes of level d in degree $2d$ (all are Oppermann sub-intervals). Last four degrees for each q; all smaller degrees are in the data file, with minimum ≥ 1 throughout.

q	d	classes	$\min \pi$	mean π	$\min \pi/(q^d/2d)$
3	13	1 594 323	60 556	61 320.1	0.9875
	14	4 782 969	169 300	170 820.3	0.9911
	15	14 348 907	476 000	478 296.9	0.9952
	16	43 046 721	1 341 030	1 345 210.0	0.9969
5	7	78 125	5 418	5 580.3	0.9709
	8	390 625	23 951	24 414.0	0.9810
	9	1 953 125	107 410	108 506.9	0.9899
	10	9 765 625	486 042	488 281.2	0.9954
7	6	117 649	9 500	9 804.0	0.9690
	7	823 543	57 981	58 824.4	0.9857
	8	5 764 801	358 343	360 300.0	0.9946
	9	40 353 607	2 235 915	2 241 867.0	0.9973
9	5	59 049	5 680	5 904.8	0.9619
	6	531 441	43 728	44 286.7	0.9874
	7	4 782 969	339 993	341 640.6	0.9952
	8	43 046 721	2 684 630	2 690 420.0	0.9978
11	4	14 641	1 735	1 830.0	0.9480
	5	161 051	15 804	16 105.0	0.9813
	6	1 771 561	146 403	147 630.0	0.9917
	7	19 487 171	1 388 036	1 391 940.7	0.9972
13	3	2 197	349	366.0	0.9531
	4	28 561	3 450	3 570.0	0.9664
	5	371 293	36 672	37 129.2	0.9877
	6	4 826 809	400 104	402 234.0	0.9947




FIGURE 1. Normalised minimum $\min_{f,c} \pi(O_{f,c})/(q^d/2d)$ against d. Left: odd q. Right: even q; the points at 0 are the empty sub-intervals of Theorem 1.1.

For $d = 2$, (3) reads $|\Psi(x) - q^2| \leq q^2 - q$, so $\Psi(x) \geq q > 0$ for every class. This still does not produce an irreducible, because Ψ also counts prime powers. In characteristic 2 the bound is attained, and this pins down every level-2 L-function.

TABLE 4. Even q: irreducible counts over the $q^{\lceil d/2 \rceil}$ Oppermann classes (Lemma 2.2). Last four degrees for each q; the only empty classes in the whole range are those of Theorem 1.1.

q	d	classes	$\min \pi$	mean π	$\min \pi/(q^d/2d)$
2	23	4 096	181 349	182 329.3	0.9945
	24	4 096	347 822	349 412.9	0.9951
	25	8 192	669 169	671 025.2	0.9971
	26	8 192	1 287 576	1 290 408.9	0.9977
4	10	1 024	51 908	52 350.0	0.9901
	11	4 096	189 054	190 603.5	0.9916
	12	4 096	695 868	698 820.5	0.9954
	13	16 384	2 575 566	2 580 963.0	0.9979
8	4	64	440	490.0	0.8594
	5	512	3 204	3 276.0	0.9778
	6	512	21 616	21 840.0	0.9895
	7	4 096	149 148	149 796.0	0.9957
16	2	16	**0**	60.0	**0**
	3	256	600	680.0	0.8789
	4	256	7 616	8 160.0	0.9297
	5	4 096	103 680	104 856.0	0.9888

Proposition 6.1 (characteristic 2, $d = 2$)**.** *Let $q = 2^k$, and let $B_e = \{t^4 + et^2 + \alpha t + \beta\}$ as in §3.*

(1) *$E(B_e) = q$ for every $e \in \mathbb{F}_q$.*
(2) *$\Psi(B_0) = q^2 - (-1)^k(q^2 - q)$, and $\Psi(B_e) = q^2 + (-1)^k q$ for $e \neq 0$. In particular, B_0 attains the lower Weil bound q when k is even and the upper bound $2q^2 - q$ when k is odd.*
(3) *Every character χ of exact level 2 has $L(u, \chi) = 1 - \gamma_\chi u$ with $\gamma_\chi^4 = (-1)^k q^2$.*
(4) *For every level-2 class x in degree 4,*

$$\Psi(x) = q^2 - (-1)^k(q^2[x = 1] - q\,[p_3(x) = 0]).$$

In particular, every class with $p_3 \neq 0$ contains exactly $q^2/4$ irreducibles.

Proof. (1) The proper prime powers of degree 4 are ϖ^4 with $\deg \varpi = 1$ and ϖ^2 with $\deg \varpi = 2$.

- $(t+c)^4 = t^4 + c^4$ lies in B_e only for $e = 0$, and then for all q values of c, each with $\Lambda = 1$.
- $(t^2 + b_1 t + b_0)^2 = t^4 + b_1^2 t^2 + b_0^2$ lies in B_e exactly when $b_1^2 = e$. For $e = 0$ this forces $b_1 = 0$, and $t^2 + b_0$ is a square, not irreducible. For $e \neq 0$ we have $b_1 = \sqrt{e} \neq 0$, and by Theorem 1.1(A) the polynomial $t^2 + b_1 t + b_0$ is irreducible for exactly $q/2$ values of b_0, each contributing $\Lambda = 2$.

In both cases $E(B_e) = q$.

(2) We have $\Psi(B_e) = 4\pi(B_e) + q$. Insert the counts of Theorem 1.1(B).

(3) B_0 is the identity of G_2, so $\chi(B_0) = 1$ for every χ. Characters of exact level 1 have $L = 1$ and $\psi_4 = 0$. By (2) and (2),

$$\sum_{\chi \text{ exact level } 2} \psi_4(\chi) = q^2(\Psi(B_0) - q^2) = -(-1)^k(q^2 - q)\,q^2.$$

There are $q^2 - q$ such characters, and each has $|\psi_4(\chi)| \leq q^2$. Equality in the triangle inequality therefore forces $\psi_4(\chi) = -(-1)^k q^2$ for each of them. Then $L(u, \chi)$ cannot be constant, so $L(u, \chi) = 1 - \gamma_\chi u$ and $\psi_4(\chi) = -\gamma_\chi^4$.




FIGURE 2. Characteristic-2 rigidity. *Left:* normalised counts $\pi(x)/(q^d/2d)$ for $q = 2^k$, $k \le 8$. Exact values from the Hayes FFT are shown as markers; the formulas of Theorem 1.1(B) and Theorem 5.2 are shown as lines. The bars give the range over the $d = 3$ Oppermann classes with $p_3 \neq 0$. For odd k this range reduces to the single value of C_e, as Theorem 5.2(1) predicts. For even k its endpoints are given by the two values of $m(c)$. *Right:* normalised inverse zeros $\gamma/\sqrt{q}$ of $L(u,\chi)$ for the characters of exact level 2 and 3 of G_3. For $q = 8$ (blue; marker area proportional to multiplicity) they occupy only 16 of the 24-th roots of unity. For $q = 7$ (grey; 630 zeros) they spread over the whole circle.

(4) By (3) and (2),

$$\Psi(x) = q^2 - (-1)^k \sum_{\chi \text{ exact level } 2} \overline{\chi}(x).$$

The inner sum is a sum over all characters of G_2 minus a sum over the characters of G_1, which equals $q^2[x = 1] - q[x \in \ker(G_2 \to G_1)]$. The kernel consists of the classes with $p_3 = 0$.

If $p_3(x) \neq 0$, then x contains no proper prime power: ϖ^4 and ϖ^2 have no odd-degree terms. So $\pi(x) = \Psi(x)/4 = q^2/4$. □

All four statements were confirmed for $q = 2, 4, 8, 16$ by the script `char2_d2_rigidity.py`. In characteristic 2 the explicit formula thus degenerates completely at $d = 2$. The $q^2 - q$ characters of exact level 2 all have the same ψ_4, so Ψ takes only three values: on the identity class, on the other classes with $p_3 = 0$, and on the classes with $p_3 \neq 0$. The Oppermann classes are exactly the classes with $p_3 = 0$ (Lemma 2.2), and the identity class B_0 is where the fourth powers absorb the entire Weil lower bound.

Results on the variance of Ψ over Legendre intervals (an average over f) bound the number of intervals that deviate by a given amount; they do not exclude a single empty interval.

6.4. **Large q.** For fixed degree and $q \to \infty$, Bank, Bary-Soroker and Rosenzweig [1] prove the following. For monic $P \in \mathbb{F}_q[t]$ of degree k and $1 \le m < k$, the number N of irreducibles in $\{P + h : \deg h \le m\}$ satisfies

$$|N - q^{m+1}/k| \le c(k)\, q^{m+1/2},$$

provided that $m \ge 2$ if $\gcd(q, k(k-1)) > 1$, and $m \ge 3$ if q is even and $\deg P' \le 1$. They also note that the estimate fails outside this range.

Our blocks $B_e = \{t^4 + et^2 + h : \deg h \le 1\}$ have $k = 4$ and $m = 1$, and in characteristic 2 the centre $P = t^4 + et^2$ has $P' = 0$, so both conditions exclude them. Theorem 1.1(B) gives the exact count at this excluded point and shows how the asymptotic fails:

- for $q = 4^j$ the block $e = 0$ contains no irreducible, while the expected count is $q^2/4$;
- for $q = 2^{2j+1}$ the block $e = 0$ contains $q(q-1)/2$, about twice the expected count.

The blocks $e \neq 0$, with $q^2/4$ and $q(q-2)/4$ irreducibles, do agree with the asymptotic. The obstruction comes from the additive structure of $t^4 + et^2 + \alpha t$ (Proposition 3.1). For $m = 1$, $k = 4$ and q even, the conditions of [1] therefore cannot be dropped. By contrast, when $3 \mid q$, a case that is also excluded by $\gcd(q, k(k-1)) > 1$ at $k = 4$, $m = 1$, Theorem 1.4 shows that the asymptotic $q^2/4 + O(q)$ continues to hold without exception.

Theorem 1.1(C) has $q = 2$ fixed, so it lies outside the $q \to \infty$ regime altogether.

In the other direction, [1] gives nonemptiness for large q in most cases. An Oppermann sub-interval $O_{f,c} = \{f^2 + ct^d + h : \deg h \leq d-1\}$ has $k = 2d$ and $m = d-1$. Since $N \geq q^d/2d - c(2d)q^{d-1/2}$, the sub-interval is nonempty as soon as $q > (2d\,c(2d))^2$, provided the conditions of [1] hold. We do not make the constant $c(2d)$ explicit.

Corollary 6.2 (of [1])**.** *For each d there is $q_0(d)$ such that every Oppermann sub-interval of degree $2d$ over $\mathbb{F}_q$, with $q \geq q_0(d)$, contains an irreducible polynomial in each of the following cases:*

(1) *q odd and $d \geq 3$;*
(2) *q odd, $d = 2$ and $\gcd(q, 6) = 1$;*
(3) *q even and $d \geq 4$;*
(4) *q even, $d = 3$, and the class has $p_3 \neq 0$.*

Proof. We check the conditions of [1] with $k = 2d$ and $m = d-1$; recall that $1 \leq m < k$ is required.

- *q odd.* The only condition is $m \geq 2$ when $p \mid 2d(2d-1)$. It holds for $d \geq 3$. For $d = 2$, $k(k-1) = 12$, so it holds when $\gcd(q, 6) = 1$.
- *q even, $d \geq 4$.* Here $m \geq 3$, so both conditions hold.
- *q even, $d = 3$.* By Lemma 2.2 every member P of an Oppermann class has $p_5 = 0$. In characteristic 2 this gives $P' = p_3 t^2 + p_1$, so $\deg P' = 2$ when $p_3 \neq 0$, and the condition "$m \geq 3$ if $\deg P' \leq 1$" does not apply.

□

The cases left open by the corollary are the following. For $d = 1$ the interval is too short for [1], but Theorem 1.1(A) and, for odd q, the count $(q-1)/2$ of irreducible $t^2 + bt + s_0$ settle it. For $d = 2$ the corollary is superseded: Theorems 1.1(B) and 1.4 give exact counts for every q, including $3 \mid q$, with no threshold. For q even with $d = 3$, the corollary is superseded as well. Theorem 5.2 in §5 counts all q^2 classes exactly for every q, including the q classes with $p_3 = 0$ (prefixes $(p_5, p_4, p_3) = (0, x, 0)$) that [1] misses. What remains is $q < q_0(d)$ for q odd with $d \geq 3$, and for q even with $d \geq 4$; these ranges are covered only by computation.

7. Mean drift and coherent variance on the Oppermann subgroup

Throughout this section $q = 2^k$, and $H_d \subset G_d$ is the Oppermann subgroup of §2.1.

7.1. Mean drift on H_d for $d \leq 5$.

Proposition 7.1. *Let $L = \mathbb{F}_{q^{2d}}$ and $N_d = \#\{x \in L : \mathrm{Tr}_{L/K}(x^j) = 0 \text{ for all odd } j < d\}$. Then*

$$\mathbb{E}_{x \in H_d}[\Psi(x)] = q^{-\lceil d/2 \rceil} N_d = q^d + q^{-d} \sum_{\beta \neq 0} \psi_{2d}(\chi_\beta).$$

Proof. $\sum_{x \in H_d} \Psi(x) = \sum \Lambda(F)$ over the F of degree $2d$ with $\Phi(F) = 0$. Group these by roots as in the proof of Theorem 2.5; since $S_j(F_x) = \mathrm{Tr}_{L/K}(x^j)$, the sum equals N_d. The second expression follows from orthogonality: $|H_d|^{-1} \sum_{x \in H_d} \overline{\chi}(x)$ equals 1 if $\chi \in H_d^\perp$ and 0 otherwise, and $\psi_{2d}(1) = q^{2d}$. □

Corollary 7.2. *For $d = 2, 3$, $\mathbb{E}_{H_d}[\Psi] = q^d$. Moreover $\mathbb{E}_{H_2}[\pi] = q(q-1)/4$ and $\mathbb{E}_{H_3}[\pi] = q(q^2-1)/6$.*

Proof. Only $j = 1$ occurs, and $\psi_{2d}(\chi_{\beta_1}) = \sum_x (-1)^{\mathrm{Tr}(\beta_1 x)} = 0$ for $\beta_1 \neq 0$.

For π we subtract the elements of degree less than $2d$ from N_d. For $d = 2$, every $x \in \mathbb{F}_{q^2}$ has $\mathrm{Tr}_{L/K}(x) = 2\,\mathrm{Tr}_{\mathbb{F}_{q^2}/K}(x) = 0$. These are q^2 elements, and $N_2 = q^3$, so $\sum_{H_2} 4\pi = q^3 - q^2$. For $d = 3$, the elements of $\mathbb{F}_{q^3}$ satisfy the condition automatically. An element $x \in \mathbb{F}_{q^2}$ has $\mathrm{Tr}_{L/K}(x) = \mathrm{Tr}_{\mathbb{F}_{q^2}/K}(x)$, which vanishes only for $x \in K$ (since $\ker \mathrm{Tr}_{\mathbb{F}_{q^2}/K} = K$ in characteristic 2). Hence $\sum_{H_3} 6\pi = q^5 - q^3$. □

Theorem 7.3. *Let $d \in \{4, 5\}$ and $q = 2^k$. Then*

	N_d	$\mathbb{E}_{H_d}[\Psi]$	$\mathbb{E}_{H_d}[\pi]$
$d = 4$, k *even*	q^6	q^4	$(q^4 - q^2)/8$
$d = 4$, k *odd*	$q^6 - 2q^4 + 2q^3$	$q^4 - 2q^2 + 2q$	$(q^4 - 3q^2 + 2q)/8$
$d = 5$	$q^8 + q^5 - q^4$	$q^5 + q^2 - q$	$(q^5 - q)/10$

Proof. For $d \in \{4, 5\}$ the odd indices are $j = 1, 3$. Let $E = \mathbb{F}_{q^d}$, so $K \subset E \subset L = \mathbb{F}_{q^{2d}}$, and put $u = \mathrm{Tr}_{L/E}(x) = x + x^{q^d}$. Then $\mathrm{Tr}_{L/K}(x) = \mathrm{Tr}_{E/K}(u)$.

Case $u = 0$. Then $x \in E$. Both traces vanish, since $\mathrm{Tr}_{L/K}(y) = 2\,\mathrm{Tr}_{E/K}(y) = 0$ for $y \in E$. This gives q^d solutions.

Case $u \neq 0$. We need $\mathrm{Tr}_{E/K}(u) = 0$, which leaves $q^{d-1} - 1$ values of u. As in the proof of Theorem 5.2, write $x = u\xi$ with $\xi + \xi^{q^d} = 1$, and put $\alpha = \xi^2 + \xi \in E$. Then α runs 2-to-1 over $\{\mathrm{Tr}_{E/\mathbb{F}_2}(\alpha) = 1\}$, and $\mathrm{Tr}_{L/K}(x^3) = \mathrm{Tr}_{E/K}(u^3(\alpha + 1))$. The condition $\mathrm{Tr}_{L/K}(x^3) = 0$ cuts out an affine K-hyperplane $H_u \subset E$ of size q^{d-1}. On H_u, the functional $\mathrm{Tr}_{E/\mathbb{F}_2}$ is non-constant unless $u^3 = \nu \in K^\times$.

- If $u^3 \notin K^\times$, then u contributes q^{d-1} elements x.
- If $u^3 = \nu \in K^\times$, then $\mathrm{Tr}_{E/K}(\alpha) = \mathrm{Tr}_{E/K}(1) = d$ on H_u, so $\mathrm{Tr}_{E/\mathbb{F}_2}(\alpha) = dk \bmod 2$. Then u contributes $2q^{d-1}$ elements if dk is odd, and none otherwise.

With $a_d = \#\{u \in E^\times : u^3 \in K,\ \mathrm{Tr}_{E/K}(u) = 0\}$ this gives

$$N_d = q^d + (q^{d-1} - 1 - a_d)\, q^{d-1} + 2a_d\, q^{d-1}[dk \text{ odd}].$$

Computing a_d. Such u is a root of some $X^3 - \nu$ with $\nu \in K^\times$, so its degree over K is 1, 2 or 3, and it must divide d.

- $d = 4$. Every $u \in K^\times$ qualifies, since $\mathrm{Tr}_{E/K}(u) = 4u = 0$. Elements of degree 2 occur only when some $X^3 - \nu$ has an irreducible quadratic factor over K. This happens exactly when k is odd: then $3 \nmid q - 1$, and $X^3 - \nu = (X - a)(X^2 + aX + a^2)$. The resulting $2(q - 1)$ elements $a\omega, a\omega^2 \in \mathbb{F}_{q^2} \setminus K$ have trace $\mathrm{Tr}_{E/K} = 2\,\mathrm{Tr}_{\mathbb{F}_{q^2}/K} = 0$. When k is even, the cube roots of a non-cube lie in $\mathbb{F}_{q^3} \not\subset E$. So $a_4 = q - 1$ for k even and $a_4 = 3(q - 1)$ for k odd. In both cases dk is even.
- $d = 5$. Only degree 1 is possible. But $u \in K^\times$ has $\mathrm{Tr}_{E/K}(u) = 5u = u \neq 0$, so $a_5 = 0$.

Substituting gives N_d, and $\mathbb{E}_{H_d}[\Psi] = N_d / q^{\lceil d/2 \rceil}$.

The mean of π. The elements of degree less than $2d$ satisfying the conditions are exactly those of E. For $d = 4$, every proper subfield of $\mathbb{F}_{q^8}$ lies in E. For $d = 5$, an element $x \in \mathbb{F}_{q^2} \setminus K$ has $\mathrm{Tr}_{L/K}(x) = 5\,\mathrm{Tr}_{\mathbb{F}_{q^2}/K}(x) \neq 0$ (since $\ker \mathrm{Tr}_{\mathbb{F}_{q^2}/K} = K$ in characteristic 2). Hence $\sum_{H_d} 2d\pi = N_d - q^d$. □

Table 5 compares Corollary 7.2 and Theorem 7.3 with the exact class counts for $q \leq 16$; all 16 cases agree. By Proposition 7.1, the drift $\mathbb{E}_{H_d}[\Psi] - q^d$ is q^{-d} times a sum of Artin–Schreier sums of the Gold-type polynomials $\beta_1 x + \beta_3 x^3$ over $\mathbb{F}_{q^{2d}}$. It vanishes for $d = 4$ with k even. It is $-2q^2 + 2q$ for $d = 4$ with k odd, coming from the cube roots of unity outside K, and $+q^2 - q$ for $d = 5$. Figure 3 (left) shows the drift in units of $q^{d/2}$ for $d \leq 12$.

TABLE 5. Means over the Oppermann subgroup H_d (exact, from the Hayes FFT) against the closed forms of Corollary 7.2 and Theorem 7.3. Every entry agrees.

q	d	$\lvert H_d\rvert$	$\mathbb{E}_{H_d}[\Psi]$	$\mathbb{E}_{H_d}[\pi]$	q^d
2	2	2	4	1/2	4
2	3	4	8	1	8
2	4	4	12	1	16
2	5	8	34	3	32
4	2	4	16	3	16
4	3	16	64	10	64
4	4	16	256	30	256
4	5	64	1 036	102	1 024
8	2	8	64	14	64
8	3	64	512	84	512
8	4	64	3 984	490	4 096
8	5	512	32 824	3 276	32 768
16	2	16	256	60	256
16	3	256	4 096	680	4 096
16	4	256	65 536	8 160	65 536
16	5	4 096	1 048 816	104 856	1 048 576




FIGURE 3. *Left:* mean drift on the Oppermann subgroup, $(\mathbb{E}_{H_d}[\Psi]-q^d)/q^{d/2}$, for $q=2,4,8,16$. Open circles mark the closed forms of Theorem 7.3; the drift vanishes for $d=2,3$. *Right:* the normalised values of $m(c)$, which equal $\mp\varepsilon=\mp(-1)^{k/2}$ by Theorem 5.3.

7.2. **Coherent superposition and subgroup variance.** By the explicit formula, $\Psi(x)=q^{-d}\sum_{\chi\in\widehat{G_d}}\overline{\chi}(x)\psi_{2d}(\chi)$. On all of G_d, Parseval gives

$$\mathrm{Var}_{G_d}(\Psi)=q^{-2d}\sum_{\chi\neq 1}|\psi_{2d}(\chi)|^2,$$

one term per character. On H_d this changes.

Proposition 7.4.

$$\mathrm{Var}_{x\in H_d}(\Psi(x))=q^{-2d}\sum_{[\chi]\neq H_d^{\perp}}\Big|\sum_{\eta\in H_d^{\perp}}\psi_{2d}(\chi\eta)\Big|^2,$$

where $[\chi]$ *runs over the cosets of* $H_d^{\perp}$ *in* $\widehat{G_d}$.

TABLE 6. Normalised variances in characteristic 2: over the Oppermann subgroup H_d and over all classes G_d, for Ψ and for $2d\pi$, with the Keating–Rudnick value $d-1$. The entries at $d=2,3$ agree exactly with Theorem 7.5. For $q=2$ only $d\le 12$ is shown.

q	d	$\lvert H_d\rvert$	$\frac{\mathrm{Var}_H(\Psi)}{q^d}$	$\frac{\mathrm{Var}_H(2d\pi)}{q^d}$	$\frac{\mathrm{Var}_G(\Psi)}{q^d}$	$\frac{\mathrm{Var}_G(2d\pi)}{q^d}$	$d-1$
2	2	2	1.000	1.000	0.500	0.750	1
2	3	4	0.500	0.000	1.750	1.617	2
2	4	4	1.125	2.000	2.031	2.938	3
2	5	8	1.500	1.562	2.234	2.219	4
2	6	8	9.594	9.000	6.184	6.099	5
2	7	16	4.375	5.359	4.818	4.909	6
2	8	16	5.002	5.250	5.500	5.637	7
2	9	32	7.022	7.119	7.828	7.879	8
2	10	32	9.152	9.375	7.854	7.836	9
2	11	64	11.896	11.757	9.092	9.105	10
2	12	64	13.821	13.192	10.302	10.339	11
4	2	4	3.000	3.000	0.750	0.938	1
4	3	16	2.250	4.500	2.438	2.953	2
4	4	16	3.047	3.000	1.315	1.371	3
4	5	64	6.328	9.668	4.902	5.098	4
4	6	64	6.882	6.961	4.076	4.114	5
4	7	256	8.285	9.654	6.273	6.291	6
4	8	256	8.601	8.707	6.801	6.805	7
4	9	1 024	6.827	8.703	7.733	7.742	8
4	10	1 024	11.731	11.766	8.653	8.655	9
8	2	8	7.000	7.000	0.875	0.984	1
8	3	64	0.875	0.000	2.734	2.618	2
8	4	64	5.045	4.812	3.490	3.556	3
8	5	512	3.466	2.606	3.540	3.525	4
8	6	512	19.431	19.380	4.575	4.573	5
8	7	4 096	8.740	10.499	6.123	6.126	6
16	2	16	15.000	15.000	0.938	0.996	1
16	3	256	2.812	5.625	2.871	3.043	2
16	4	256	18.519	18.750	3.149	3.153	3
16	5	4 096	6.478	9.532	3.991	4.003	4

Proof. For $x\in H_d$, $\overline{\chi}(x)$ depends only on the restriction $\chi|_{H_d}$, that is, on the coset $[\chi]\in \widehat{G_d}/H_d^{\perp}\cong\widehat{H_d}$. Hence

$$\Psi|_{H_d}=q^{-d}\sum_{\lambda\in\widehat{H_d}}\overline{\lambda}\,A(\lambda),\qquad A(\lambda)=\sum_{\chi|_{H_d}=\lambda}\psi_{2d}(\chi).$$

The trivial λ gives the mean (Proposition 7.1). Apply Parseval on H_d. □

Coherent superposition. The $|H_d^{\perp}|=q^{\lfloor d/2\rfloor}$ characters in one coset are orthogonal on G_d but coincide on H_d, so their coefficients add as amplitudes rather than as energies. If the phases of the $\psi_{2d}(\chi\eta)$ in a coset were independent, Var_{H_d} and Var_{G_d} would agree on average. If they are aligned, Var_{H_d} can exceed Var_{G_d} by a factor up to $|H_d^{\perp}|$; if they cancel, it is smaller. Both extremes occur.

Theorem 7.5. *Let $q=2^k$.*

(1) $\mathrm{Var}_{H_2}(\Psi)/q^2 = q-1$ *and* $\mathrm{Var}_{G_2}(\Psi)/q^2 = 1-1/q$. *The ratio is exactly* $q = |H_2^{\perp}|$, *so the coherence is maximal.*
(2) $\mathrm{Var}_{H_2}(4\pi)/q^2 = q-1$.
(3) $\mathrm{Var}_{H_3}(\Psi)/q^3 = 1-1/q$ *for* k *odd and* $3(1-1/q)$ *for* k *even.*
(4) $\mathrm{Var}_{H_3}(6\pi)/q^3 = 0$ *for* k *odd and* $6(1-1/q)$ *for* k *even.*

Proof. (1) By Proposition 6.1, $\Psi(x)-q^2$ takes three values: $-(-1)^k(q^2-q)$ on the identity class, $(-1)^k q$ on the other $q-1$ classes of H_2, and 0 on the q^2-q classes outside H_2. Averaging the squares over H_2 (with q classes) and over G_2 (with q^2 classes) gives $q^2(q-1)$ and $q(q-1)$.

The ratio is maximal because Proposition 6.1(3) gives $\psi_4(\chi) = -(-1)^k q^2$ for all characters of exact level 2: every coset sum is fully aligned.

(2) $E(B_e) = q$ for every e (Proposition 6.1(1)), so $4\pi = \Psi - q$ on H_2 and has the same variance as Ψ.

(3), (4) $H_3 \cong \mathbb{F}_q^2$ with coordinates (e,c). By Corollary 5.4, both Ψ and π depend only on c. Averaging over $(e,c) \in H_3$ is therefore the same as averaging over $c \in \mathbb{F}_q$, and all variances below are written as $q^{-1}\sum_{c\in\mathbb{F}_q}$.

For k odd, $6\pi = q(q^2-1)$ is constant, so $\mathrm{Var}_{H_3}(6\pi) = 0$. The variance of Ψ then comes entirely from the prime powers, $E(C_{e,c}) = q^2[c=0]$. Since $\Psi - q^3 = -q + q^2[c=0]$,

$$\mathrm{Var}_{H_3}(\Psi) = q^{-1}((q^2-q)^2 + (q-1)q^2) = q^2(q-1).$$

For k even, the deviation $\Psi - q^3$ is

- $-q^2+q$ for $c=0$;
- $q - 2\varepsilon q^{3/2}$ for each of the $(q-1)/3$ cubes c;
- $q + \varepsilon q^{3/2}$ for each of the $2(q-1)/3$ non-cubes c.

Its mean is 0. In the second moment the cross terms in $\varepsilon q^{5/2}$ cancel, leaving

$$\mathrm{Var}_{H_3}(\Psi) = q^{-1}(q^2(q-1)^2 + (q-1)(q^2+2q^3)) = 3q^2(q-1).$$

For 6π, the mean is $q(q^2-1)$, and the deviations are

- $-2q(q-1)$ for $c=0$;
- $2q - 2\varepsilon q^{3/2}$ for each cube c;
- $2q + \varepsilon q^{3/2}$ for each non-cube c.

The same computation gives

$$\mathrm{Var}_{H_3}(6\pi) = q^{-1}(4q^2(q-1)^2 + (q-1)(4q^2+2q^3)) = 6q^2(q-1).$$

Adding $E(C_{e,0}) = q^2$ halves the deviation at $c=0$, to $-q(q-1)$. Removing the mean of E shifts the other deviations by $-q$. Together these halve the variance. □

All four statements were also checked exactly, with rational arithmetic on the class counts, for $q = 2, 4, \ldots, 64$ (`verify_variance23.py`).

The exact class values on H_3 also determine the coset sums $A(\lambda)$ of Proposition 7.4. Since $H_3 = \{1 + eu^2 + cu^3\} \cong (\mathbb{F}_q^2, +)$, its characters are $\lambda_{a,b}(e,c) = (-1)^{\mathrm{Tr}(ae+bc)}$.

Proposition 7.6 (level-3 coset sums)**.** *Let* $q = 2^k$, *and let* $A(\lambda) = \sum_{\chi|_{H_3}=\lambda} \psi_6(\chi)$, *a sum over* q *characters of* G_3.

(1) $A(\lambda_{a,b}) = 0$ *if* $a \neq 0$.
(2) *For* $b \neq 0$, $A(\lambda_{0,b}) = q^4$ *if* k *is odd or* b *is a cube, and* $A(\lambda_{0,b}) = -2q^4$ *if* k *is even and* b *is a non-cube.*

Consequently, for k *even, each of the* $2q(q-1)/3$ *characters* χ *restricting to some* $\lambda_{0,b}$ *with* b *a non-cube has exact level* 3, *and both inverse zeros of* $L(u,\chi)$ *satisfy* $\gamma^6 = q^3$.

Proof. By the proof of Proposition 7.4,

$$A(\lambda) = q^3|H_3|^{-1}\sum_{x\in H_3}\Psi(x)\lambda(x) = q\sum_{e,c}\Psi(e,c)\lambda_{a,b}(e,c).$$

By Corollary 5.4, Ψ depends only on c, so the sum over e kills every $\lambda_{a,b}$ with $a \neq 0$.

For $a = 0$ and $b \neq 0$, the constant part of Ψ drops out, and $A = q^2 \sum_c (\Psi(c) - q^3)(-1)^{\mathrm{Tr}(bc)}$. For k odd, $\Psi(c) - q^3 = -q + q^2[c = 0]$, which gives q^4. For k even, insert the deviations listed in the proof of Theorem 7.5. The cubic-character sum $\sum_{c \in (K^\times)^3} (-1)^{\mathrm{Tr}(bc)} = \frac{1}{3}(-1 - 2\varepsilon\sqrt{q}\,\mathrm{Re}\,\omega(b))$ from the proof of Theorem 5.3 then gives $A = q^4(2\,\mathrm{Re}\,\omega(b) - 1)$, which is q^4 or $-2q^4$.

For the consequence: each $L(u, \chi)$ with $\chi \neq 1$ has degree at most 2, and its inverse zeros have absolute value at most $\sqrt{q}$, so $|\psi_6(\chi)| \leq 2q^3$. A sum of q such terms can equal $-2q^4$ only if every term equals $-2q^3$. This forces $\deg L = 2$ and $\gamma_1^6 = \gamma_2^6 = q^3$. □

All coset sums were checked against the exact class values for $q \leq 64$, with the script `verify_level3_cosets.py`. The L-functions confirm the consequence; in fact many more characters have $\psi_6 = -2q^3$, for example 1020 for $q = 16$, against the 160 that the proposition forces.

For $d \geq 4$ we have only numerical data (Table 6, Figure 4). We observe the following:

(1) On the whole group, $\mathrm{Var}_{G_d}(\Psi)/q^d$ follows the Keating–Rudnick value $d - 1$ closely for $q \geq 4$, and in odd characteristic. For $q = 2$ it stays near $d - 2$. Keating and Rudnick prove $\mathrm{Var} \sim q^{h+1}(n - h - 2)$ as $q \to \infty$ for intervals of size q^{h+1} in degree n; here $h = d - 1$ and $n = 2d$, see [6] for the precise hypotheses. We use $d - 1$ only as a reference value.

(2) On the subgroup, $\mathrm{Var}_{H_d}(\Psi)/q^d$ is erratic. For even d it is large and grows with q: $q - 1$ at $d = 2$, and 1.1, 3.0, 5.0, 18.5 at $d = 4$ for $q = 2, 4, 8, 16$. For odd d it is moderate: $1 - 1/q$ or $3(1 - 1/q)$ at $d = 3$, and 1.5, 6.3, 3.5, 6.5 at $d = 5$. This is the coherence of Proposition 7.4.

Two structural features separate even from odd d, although we have no proof of the dichotomy beyond $d = 3$. First, the coherence multiplicity $|H_d^\perp| = q^{\lfloor d/2 \rfloor}$ jumps by a factor q at each even d. Second, the squares $\varpi^2 = \sum a_i^2 t^{2i}$ of irreducibles of degree d contain only even powers of t. For odd d their coefficient e_d (of t^d) is 0, so they pile up on the slice $e_d = 0$ of H_d. For even d it is $a_{d/2}^2$, which is spread over all $e_d \in \mathbb{F}_q$.




FIGURE 4. Normalised variance of Ψ over all classes G_d (left, odd and even q) and over the Oppermann subgroup H_d (right, q even). Dashed line: the Keating–Rudnick value $d - 1$. Circles: Theorem 7.5.

8. QUESTIONS

(1) In characteristic 2, are (A)–(C) the only empty Oppermann sub-intervals for all q and d? For $d \leq 3$ the question is settled for every q (Theorems 1.1, 5.2 and Corollary 5.4). For each $d \geq 4$, by Corollary 6.2, only finitely many q can carry further empty sub-intervals. Corollary 3.7 shows that, from $d = 4$ on, the additive mechanism of (B) is no longer available to produce empty or full classes.

(2) For odd q, is every Oppermann sub-interval nonempty, for all q and d? Lemma 2.1 turns this into a statement about all short intervals of length $q^{n/2}$. For $d = 2$ it holds

for every odd q (Theorem 1.4). For each $d \geq 3$, by Corollary 6.2, it fails for at most finitely many q. The trace-form method of Section 4 would need, for $d = 3$, the joint distribution of $\mathrm{Tr}(x^2)$ and $\mathrm{Tr}(x^3)$ on $\ker \mathrm{Tr} \subset \mathbb{F}_{q^6}$. That is, it would need the point counts of the intersections of an affine quadric and an affine cubic in the 5-dimensional $\mathbb{F}_q$-space $\ker \mathrm{Tr}$. These are affine threefolds, and their point counts involve cubic exponential sums in five variables, for which no closed form is to be expected. In characteristic 2 the tower $\mathbb{F}_q \subset \mathbb{F}_{q^3} \subset \mathbb{F}_{q^6}$ linearises both conditions.

(3) Outside the characteristic-2 cases above, is there a constant $c > 0$ with $\min_x \pi(x) \geq c\, q^d/2d$ for all large d?

(4) In characteristic 2, evaluate the mean drift $\mathbb{E}_{H_d}[\Psi] - q^d$ for $d \geq 6$, and explain the even/odd dichotomy of $\mathrm{Var}_{H_d}(\Psi)/q^d$ observed in Table 6.

(5) In characteristic 2, do all characters of exact level 3 of G_3 have L-functions with $\gamma^6 \in q^3\{\pm 1, \pm i\}$? This is observed for $q \leq 16$ (Figure 2, right). It is proved at level 2 (Proposition 6.1), and at level 3 for the characters of Proposition 7.6.

Data and code

All scripts are in `code/` and their outputs in `data/` of https://github.com/Ruqing1963/oppermann-function-fields.

- `hayes_oppermann.py`: Theorems 1.2, 1.3.
- `verify_pointwise_oppermann.py`, `compare_bruteforce.py`: cross-checks.
- `char2_d2_blocks.py`, `char2_d2_theorem.py`: Theorem 1.1(B).
- `swan_parity.py`: Theorem 1.1(C), Table 1.
- `verify_additive_slices.py`: Theorem 3.6, Corollary 3.7.
- `odd_d2_formula.py`: Theorem 1.4.
- `char2_d3_p3zero.py`, `char2_d3_all.py`: Theorem 5.2.
- `cubic_gauss_m.py`, `make_d3_table.py`: Theorem 5.3, Table 2.
- `char2_d2_rigidity.py`: Proposition 6.1; `verify_level3_cosets.py`: Proposition 7.6.
- `subgroup_lib.py`, `verify_subgroup_mean.py`: Theorems 2.4, 2.5, 7.3, Corollary 7.2.
- `subgroup_variance_census.py`, `verify_variance23.py`: Theorem 7.5, Table 6.
- `make_figure.py`, `make_figure2.py`, `make_tables.py`: Figures 1, 2, Tables 3, 4.
- `make_subgroup_figures.py`, `make_subgroup_tables.py`: Figures 3, 4, Tables 5, 6.
- `hayes_fft_fq.py`, `fqlib.py`, `small_q_gf.py`: finite-field and Hayes-group routines from [4]; `fqlib.py` is extended by moduli for $q = 32, \ldots, 256$.

References

[1] E. Bank, L. Bary-Soroker, L. Rosenzweig, *Prime polynomials in short intervals and in arithmetic progressions*, Duke Math. J. **164** (2015), 277–295.

[2] E. R. Berlekamp, *An analog to the discriminant over fields of characteristic two*, J. Algebra **38** (1976), 315–317.

[3] B. C. Berndt, R. J. Evans, K. S. Williams, *Gauss and Jacobi Sums*, Wiley, New York, 1998.

[4] R. Chen, *Primes in Legendre intervals over $\mathbb{F}_q[t]$: an exact explicit-formula algorithm, verification for $d \leq 10$, and an anomaly in characteristic 2*, preprint (2026), DOI 10.5281/zenodo.23225896.

[5] D. R. Hayes, *The distribution of irreducibles in $GF[q, x]$*, Trans. Amer. Math. Soc. **117** (1965), 101–127.

[6] J. P. Keating, Z. Rudnick, *The variance of the number of prime polynomials in short intervals and in residue classes*, Int. Math. Res. Not. IMRN **2014**, no. 1, 259–288.

[7] R. Lidl, H. Niederreiter, *Finite Fields*, 2nd ed., Encyclopedia of Mathematics and its Applications **20**, Cambridge University Press, 1997.

[8] L. Oppermann, *Om vor Kundskab om Primtallenes Mængde mellem givne Grændser*, Oversigt over det Kongelige Danske Videnskabernes Selskabs Forhandlinger (1882), 169–179.

[9] L. Stickelberger, *Über eine neue Eigenschaft der Diskriminanten algebraischer Zahlkörper*, Verhandlungen des ersten Internationalen Mathematiker-Kongresses, Zürich, 1897, 182–193.

[10] L. Stickelberger, *Über eine Verallgemeinerung der Kreistheilung*, Math. Ann. **37** (1890), 321–367.

[11] R. G. Swan, *Factorization of polynomials over finite fields*, Pacific J. Math. **12** (1962), 1099–1106.

GUT Geoservice Inc., Montreal, Canada

Email address: ruqing@hotmail.com