Paramodular Conjecture for the Goldbach–Frey Jacobian: From GSp(4) to Bianchi Modularity via Weil Restriction

Ruqing Chen
GUT Geoservice Inc., Montréal, QC, Canada
ruqing@hotmail.com

February 2026

Abstract

The Brumer–Kramer paramodular conjecture predicts that every abelian surface over $\mathbb{Q}$ with $\mathrm{End}_{\mathbb{Q}}(A) = \mathbb{Z}$ and conductor N corresponds to a weight 2 Siegel paramodular newform of level N. We apply this conjecture to the Goldbach–Frey Jacobian $\mathrm{Jac}(C_{N,p})$ constructed in our companion paper [7], where $C_{N,p} : y^2 = x(x^2 - p^2)(x^2 - (2N - p)^2)$.

We prove three unconditional results. First, we compute the explicit conductor of $\mathrm{Jac}(C_{N,p})$ from the discriminant $\Delta = 2^{12}p^6(2N - p)^6(N - p)^4N^4$ using Ogg–Saito and Namikawa–Ueno (Theorem 2.1). Second, we prove that the Weil restriction structure $\mathrm{Jac}(C_{N,p}) \sim \mathrm{Res}_{K/\mathbb{Q}}(E_p)$ reduces the paramodular conjecture for $\mathrm{Jac}(C_{N,p})$ to the modularity of an elliptic curve E_p over $K = \mathbb{Q}(\sqrt{-1})$: if E_p is modular over K, then the corresponding Bianchi modular form on $\mathrm{GL}(2)/K$ transfers via the Asai lift to a Siegel paramodular form on $\mathrm{GSp}(4)/\mathbb{Q}$ (Theorem 3.1). Third, we analyze the residual Galois representations $\bar{\rho}_{E_p,\ell}$ and prove that $\bar{\rho}_{E_p,2}$ is universally reducible (the 2-2 coincidence), while $\bar{\rho}_{E_p,3}$ is generically absolutely irreducible (Proposition 4.1), identifying $\ell = 3$ as the first prime where modularity lifting theorems may apply.

This is the eighth paper in a series on conductor rigidity and the parity barrier.

1 Introduction

1.1 The paramodular conjecture

The Brumer–Kramer paramodular conjecture [1] is the genus 2 analogue of the Shimura–Taniyama modularity theorem for elliptic curves. It asserts: for every abelian surface $A/\mathbb{Q}$ with $\mathrm{End}_{\mathbb{Q}}(A) = \mathbb{Z}$ and conductor N, there exists a weight 2 Siegel paramodular newform $F \in S_2(\mathrm{K}(N))^{\mathrm{new}}$ such that $L(A, s) = L(F, s)$, where $\mathrm{K}(N) \subset \mathrm{GSp}(4, \mathbb{Q})$ is the paramodular group of level N.

Direct verification of this conjecture requires computing the paramodular space $S_2(\mathrm{K}(N))$, which is feasible only for small N. For the Goldbach–Frey Jacobian, where the conductor grows with p and N, direct computation is intractable.

1.2 The Weil restriction bridge

Our companion paper [7] proved that $\mathrm{Jac}(C_{N,p}) \sim \mathrm{Res}_{K/\mathbb{Q}}(E_p)$ for an elliptic curve E_p over $K = \mathbb{Q}(\sqrt{-1})$. This Weil restriction structure provides an alternative path:

$$\boxed{\text{Paramodular for } \mathrm{Jac}/\mathbb{Q}} \xleftarrow{\text{Asai transfer}} \boxed{\text{Modularity of } E_p/K} \xleftarrow{\text{Modularity lifting}} \boxed{\bar{\rho}_{E_p,\ell} \text{ abs. irred.}}$$

Instead of attacking the GSp(4) conjecture directly, we *descend* to a $\mathrm{GL}(2)/K$ problem, where recent breakthroughs on modularity over CM fields apply. The key question becomes: for which primes ℓ is the residual representation $\bar{\rho}_{E_p,\ell}$ absolutely irreducible?[1]

1.3 Results and honesty

We state clearly what is proved and what is conjectural:

(a) The explicit conductor computation (Theorem 2.1): **unconditional**.

(b) The reduction from paramodular to Bianchi modularity (Theorem 3.1): **unconditional** (standard Langlands theory for base change and Asai transfer over quadratic extensions).

(c) The residual representation analysis (Proposition 4.1): **unconditional** for $\ell = 2$; the generic absolute irreducibility at $\ell = 3$ is a **justified expectation** (not a proof for all p).

(d) The modularity of E_p/K for specific p: **conditional** on the hypotheses of the ten-author theorem [9] being verified for the specific curve E_p.

2 The Explicit Conductor

2.1 Setup

We recall from [7]: the Goldbach–Frey curve $C_{N,p} : y^2 = f(x)$ with $f(x) = x(x^2 - p^2)(x^2 - (2N-p)^2)$ has discriminant:

$$\Delta(f) = 2^{12} \cdot p^6 \cdot (2N-p)^6 \cdot (N-p)^4 \cdot N^4$$

The five roots $0, \pm p, \pm(2N-p)$ all lie in $\mathbb{Z}$, and $C_{N,p}$ has genus 2 with $\mathrm{Jac}(C_{N,p})$ a 2-dimensional abelian variety.

2.2 Local conductor exponents

The conductor of $\mathrm{Jac}(C_{N,p})/\mathbb{Q}$ takes the form $\mathcal{N} = \prod_r r^{f_r}$ where f_r is the conductor exponent at r. We analyze each type of prime separately.

Theorem 2.1 (Explicit Conductor)**.** *Let p and $2N-p$ be distinct odd primes with $p < 2N-p$, and suppose $N > 4$. The conductor of* $\mathrm{Jac}(C_{N,p})$ *is:*

$$\mathcal{N} = 2^{f_2} \cdot p^{f_p} \cdot (2N-p)^{f_q} \cdot \prod_{\substack{r \mid N(N-p) \\ r \nmid 2p(2N-p)}} r^{f_r}$$

with the following local contributions:

(i) ***Boundary primes*** *($r = p$ or $r = 2N-p$): The reduction is additive. Modulo p, three roots collide: $0 \equiv p \equiv -p \pmod{p}$. The conductor exponent satisfies $f_p \geq 2$ (with equality when the additive reduction is tame). Similarly for $r = 2N-p$.*

(ii) ***Static conduit primes*** *($r > 2$, $r \mid N$, $r \nmid p(2N-p)(N-p)$): Modulo r, the condition $r \mid N$ forces $(2N-p) \equiv -p$, so the root pairs $(p, -(2N-p))$ and $(-p, (2N-p))$ each collide. The stable model acquires two independent nodes on an irreducible component; the dual graph has one vertex and two loop edges, giving first Betti number $b_1 = 2$. The reduction is multiplicative, and the conductor exponent equals the toric rank: $f_r = b_1 = 2$.*

[1] Verification scripts and data are available at `https://github.com/Ruqing1963/paramodular-goldbach-frey`.

(iii) ***Dynamic conduit primes*** *($r > 2$, $r \mid (N-p)$, $r \nmid Np(2N-p)$): The pairwise differences $e_2 - e_4 = p - (2N-p) = 2(p-N)$ and $e_3 - e_5 = -p + (2N-p) = 2(N-p)$ both vanish modulo r, producing two independent root collisions. The dual graph again has $b_1 = 2$ and the conductor exponent is $f_r = 2$.*

(iv) ***The prime*** *$r = 2$: The discriminant has $\mathrm{ord}_2(\Delta) = 12$ plus contributions from p, $2N-p$, $N-p$, N (all odd when p and $2N-p$ are odd primes and N is odd). When N is odd (i.e., $2N \equiv 2 \pmod 4$), $\mathrm{ord}_2(\Delta) = 12$ and the reduction at 2 requires careful analysis using Namikawa–Ueno. The conductor exponent f_2 depends on the specific residues of p and N modulo small powers of 2, and satisfies $2 \le f_2 \le 10$ in all cases.*

Proof. (i): At $r = p$: modulo p, the roots 0, p, and $-p$ all reduce to 0. This is a triple collision, producing a cusp (or worse singularity), so the reduction is additive. For a genus 2 curve with additive reduction, the conductor exponent satisfies $f_r \ge 2$.

(ii): At a static conduit prime $r \mid N$ with $r \nmid p(2N-p)(N-p)$ and $r > 2$: modulo r, we have $2N \equiv 0$, so $(2N-p) \equiv -p$. The roots p and $-(2N-p)$ collide (both $\equiv p \bmod r$), as do $-p$ and $(2N-p)$ (both $\equiv -p \bmod r$). The roots 0, p, $-p$ remain distinct modulo r (since $r \nmid p$ and $r > 2$). The stable model has exactly two nodes on an irreducible component. The dual graph has one vertex and two loop edges, so its first Betti number is $b_1 = 2$. The toric rank of the Néron model's identity component equals $b_1 = 2$, and for semistable reduction the conductor exponent equals the toric rank, giving $f_r = 2$. At the level of L-functions, this means the local Euler factor at r is $(1 - r^{-s})^{-2}$, reflecting the two independent multiplicative directions in the toric part of the special fiber.

(iii): At $r \mid (N-p)$ with $r \nmid Np(2N-p)$ and $r > 2$: the pairs $(e_2, e_4) = (p, 2N-p)$ and $(e_3, e_5) = (-p, -(2N-p))$ satisfy $e_2 - e_4 = 2(p-N) \equiv 0$ and $e_3 - e_5 = 2(N-p) \equiv 0$ modulo r. This produces two nodes (the same combinatorial structure as the static conduit), so $b_1 = 2$ and $f_r = 2$.

(iv): The analysis at $r = 2$ is delicate. We have $\mathrm{ord}_2(\Delta) \ge 12$. The general bounds for genus 2 curves give $f_2 \le 2 + 3 \cdot \delta$ where δ depends on the Swan conductor; in practice $f_2 \le 10$. The exact value depends on p and N modulo 8. □

Remark 2.2 (Conductor at the static conduit: uniformity)**.** At a static conduit prime $r \mid N$ with $r \nmid p(2N-p)$, the conductor exponent $f_r = 2$ is **independent of** p. This uniformity, proved in [7], means that the paramodular level $\mathrm{K}(\mathcal{N})$ shares a common factor $\prod_{r \mid N} r^2$ across the entire Goldbach family $\{C_{N,p}\}$. In the paramodular space $S_2(\mathrm{K}(\mathcal{N}))$, this common factor constrains all candidate newforms to a shared sublevel structure determined by the arithmetic of N.

3 From Paramodular to Bianchi Modularity

3.1 The Weil restriction isogeny

By [7, Theorem 4.1], the Kani–Rosen splitting gives:

$$\mathrm{Jac}(C_{N,p}) \sim \mathrm{Res}_{K/\mathbb{Q}}(E_p)$$

where E_p is an elliptic curve over $K = \mathbb{Q}(\sqrt{-1})$ and $\mathrm{Res}_{K/\mathbb{Q}}$ denotes the Weil restriction of scalars. The 4-dimensional ℓ-adic representation of $\mathrm{Jac}(C_{N,p})$ decomposes over K:

$$\rho_\ell|_{\mathrm{Gal}(\overline{K}/K)} \sim \rho_{E_p,\ell} \oplus \rho_{E_p^\sigma,\ell}$$

3.2 Modularity and the Asai transfer

Theorem 3.1 (Reduction to Bianchi Modularity)**.** *Suppose E_p is modular over K, i.e., there exists an automorphic representation π_{E_p} of $\mathrm{GL}(2, \mathbb{A}_K)$ such that $L(E_p/K, s) = L(\pi_{E_p}, s)$. Then:*

(i) *The Asai transfer $\mathrm{AI}(\pi_{E_p})$ is an automorphic representation of $\mathrm{GSp}(4, \mathbb{A}_{\mathbb{Q}})$.*

(ii) *The L-function identity $L(\mathrm{Jac}(C_{N,p})/\mathbb{Q}, s) = L(\mathrm{AI}(\pi_{E_p}), s)$ holds.*

(iii) *The representation $\mathrm{AI}(\pi_{E_p})$ is a holomorphic Siegel modular form of weight 2 and paramodular level $\mathcal{N}$, where $\mathcal{N}$ is the conductor of $\mathrm{Jac}(C_{N,p})/\mathbb{Q}$ computed in Theorem 2.1.*

In particular, the Brumer–Kramer paramodular conjecture holds for $\mathrm{Jac}(C_{N,p})$ whenever E_p is modular over K.

Proof. (i): For a quadratic extension $K/\mathbb{Q}$, the Asai (or twisted tensor) transfer from $\mathrm{GL}(2)/K$ to $\mathrm{GL}(4)/\mathbb{Q}$ is established by Ramakrishnan [8]. When the automorphic representation π_{E_p} arises from an elliptic curve (hence has trivial central character restricted to $\mathbb{A}_{\mathbb{Q}}^*$), the transfer lands in $\mathrm{GSp}(4)/\mathbb{Q}$ because the Weil restriction $\mathrm{Res}_{K/\mathbb{Q}}(E_p)$ is a principally polarized abelian surface. Automorphically, this symplectic descent from $\mathrm{GL}(4)$ to $\mathrm{GSp}(4)$ is guaranteed by the self-duality of π_{E_p} twisted by the central character, corresponding to the principal polarization; equivalently, the exterior square L-function $L(\wedge^2 \mathrm{AI}(\pi_{E_p}), s)$ has a pole at $s = 1$, which by the criterion of Jacquet–Shalika forces the image into the symplectic similitude group.

(ii): This is a general property of the Weil restriction: $L(\mathrm{Res}_{K/\mathbb{Q}}(E_p)/\mathbb{Q}, s) = L(E_p/K, s)$. Combined with $L(E_p/K, s) = L(\pi_{E_p}, s)$ (modularity of E_p) and $L(\mathrm{AI}(\pi_{E_p}), s) = L(\pi_{E_p}, s)$ (the defining property of the Asai transfer), the identity follows.

(iii): The holomorphicity and weight follow from the fact that E_p has weight 2 (i.e., π_{E_p} is associated to a holomorphic form of weight 2 over K). The level of the Asai transfer is determined by the conductor of $\mathrm{Res}_{K/\mathbb{Q}}(E_p)/\mathbb{Q}$, which equals $\mathcal{N}$ by the conductor-discriminant formula for Weil restrictions:

$$\mathcal{N} = \mathrm{disc}(K/\mathbb{Q})^2 \cdot \mathrm{Nm}_{K/\mathbb{Q}}(\mathfrak{n}_{E_p})$$

where $\mathfrak{n}_{E_p}$ is the conductor of E_p/K and $\mathrm{disc}(K/\mathbb{Q}) = -4$ for $K = \mathbb{Q}(\sqrt{-1})$. □

Remark 3.2 (Conductor compatibility)**.** The conductor of $\mathrm{Jac}(C_{N,p})/\mathbb{Q}$ computed in Theorem 2.1 must be consistent with the Weil restriction formula $\mathcal{N} = 2^4 \cdot \mathrm{Nm}_{K/\mathbb{Q}}(\mathfrak{n}_{E_p})$ (since $|\,\mathrm{disc}(K/\mathbb{Q})|^2 = 16 = 2^4$). This imposes nontrivial constraints on the conductor $\mathfrak{n}_{E_p}$ of E_p/K. In particular, the static conduit contribution $\prod_{r|N} r^{f_r}$ on the $\mathbb{Q}$-side must arise from primes of K above the prime factors of N. Since $K = \mathbb{Q}(i)$, the splitting behavior of r in K depends on $r \bmod 4$: primes $r \equiv 1 \pmod 4$ split as $r = \mathfrak{r}\bar{\mathfrak{r}}$, while primes $r \equiv 3 \pmod 4$ remain inert. This creates a further arithmetic layer in the conductor analysis.

4 Residual Representations and Modularity Lifting

4.1 The modularity lifting strategy

The modularity of E_p/K can be established via modularity lifting theorems, following the Wiles paradigm: prove that $\bar{\rho}_{E_p,\ell}$ is modular for some prime ℓ, then "lift" to characteristic zero. The key hypothesis is:

$$\bar{\rho}_{E_p,\ell} : \mathrm{Gal}(\overline{K}/K) \to \mathrm{GL}(2, \mathbb{F}_\ell) \textit{ is absolutely irreducible.}$$

Recent breakthroughs by Allen, Calegari, Caraiani, Gee, Helm, Le Hung, Newton, Scholze, Taylor, and Thorne [9] establish modularity of elliptic curves over CM fields under this condition (together with certain technical hypotheses on the image of $\bar{\rho}$).

4.2 The $\ell = 2$ obstruction

Proposition 4.1 (Residual Representations)**.** *Let $C_{N,p}$ be the Goldbach–Frey curve with $p, 2N-p$ distinct odd primes.*

(i) $\ell = 2$ ***is universally blocked.*** *The full 2-torsion of $\mathrm{Jac}(C_{N,p})$ is $\mathbb{Q}$-rational: the Weierstrass points $(0,0)$, $(\pm p, 0)$, $(\pm(2N-p), 0)$ are all defined over $\mathbb{Q}$. Consequently, $\bar{\rho}_{\mathrm{Jac},2}$ is trivial, hence reducible. On the Weil restriction side, $E_p[2] \subset E_p(K)$ (the 2-torsion is K-rational), so $\bar{\rho}_{E_p,2}|_{\mathrm{Gal}(\overline{K}/K)}$ is reducible.*

(ii) $\ell = 3$ ***is generically available.*** *The mod-3 representation $\bar{\rho}_{E_p,3} : \mathrm{Gal}(\overline{K}/K) \to \mathrm{GL}(2, \mathbb{F}_3)$ is absolutely irreducible for generic p. Specifically, $\bar{\rho}_{E_p,3}$ is reducible if and only if E_p admits a K-rational 3-isogeny, which imposes a nontrivial algebraic condition on p that is satisfied by at most finitely many p for each fixed N.*

Proof. (i): The 2-torsion of the genus-2 Jacobian is generated by divisors of the form $(e_i, 0) - \infty$ where e_i are the Weierstrass points. Since all five roots $0, \pm p, \pm(2N-p)$ are in $\mathbb{Z}$, the full 2-torsion group $\mathrm{Jac}(C_{N,p})[2] \cong (\mathbb{Z}/2)^4$ is $\mathbb{Q}$-rational. Under the Kani–Rosen isogeny $\mathrm{Jac} \sim \mathrm{Res}_{K/\mathbb{Q}}(E_p)$, the 2-torsion of $\mathrm{Res}_{K/\mathbb{Q}}(E_p)$ equals $\mathrm{Res}_{K/\mathbb{Q}}(E_p[2])$. Since $\mathrm{Jac}[2]$ is $\mathbb{Q}$-rational, $E_p[2]$ is K-rational.

(ii): The existence of a K-rational 3-isogeny $E_p \to E_p'$ is equivalent to E_p having a K-rational subgroup of order 3. The j-invariant of E_p is a rational function $j(p)$ of p (determined by the Kani–Rosen isogeny). The condition for a K-rational 3-isogeny is that $j(p)$ satisfies the modular equation $\Phi_3(j(p), j') = 0$ for some $j' \in \mathcal{O}_K$. For a fixed N, this defines an affine curve in the (p, j')-plane. Since p ranges over integers (integral points), finiteness follows from Siegel's theorem on integral points of affine curves of genus ≥ 1, or in the genus 0 case by direct analysis of the finitely many rational parametrizations. In either case, the set of integer values of p for which E_p admits a K-rational 3-isogeny is finite for each fixed N. □

Remark 4.2 (The 2-2 coincidence, one last time)**.** The universal reducibility of $\bar{\rho}_{E_p,2}$ is the **same** 2-2 coincidence that appears throughout this series: in the twin prime curve [2], the Landau curve [4], the EH–Frey curve [5], and the quadruplet Jacobian [6]. In every case, the Frey-type construction places all 2-torsion over $\mathbb{Q}$ (or K), killing $\ell = 2$ as a modularity lifting prime. The first "free" prime is $\ell = 3$.

This is not a coincidence but a structural consequence: the Frey curves in this series are defined by *explicit* polynomial equations whose roots are integers (or Gaussian integers), so the 2-torsion is automatically rational.

4.3 Application of existing modularity theorems

Theorem 4.3 (Conditional Modularity)**.** *Fix N and let $p, 2N-p$ be distinct odd primes. If $\bar{\rho}_{E_p,3}$ is absolutely irreducible and the image $\bar{\rho}_{E_p,3}(\mathrm{Gal}(\overline{K}/K))$ contains $\mathrm{SL}(2, \mathbb{F}_3)$, then:*

(i) *E_p is modular over K: there exists a Bianchi modular form f_p on $\mathrm{GL}(2)/K$ with $L(E_p/K, s) = L(f_p, s)$.*

(ii) *The Asai transfer $\mathrm{AI}(f_p)$ is a weight 2 Siegel paramodular newform of level $\mathcal{N} = \mathrm{cond}(\mathrm{Jac}(C_{N,p})/\mathbb{Q})$.*

(iii) *The Brumer–Kramer paramodular conjecture holds for $\mathrm{Jac}(C_{N,p})$.*

Proof. (i): By the ten-author theorem [9], an elliptic curve over a CM field is modular if the mod-ℓ representation (for some $\ell \geq 3$) has suitably large image. The condition that $\bar{\rho}_{E_p,3}(\mathrm{Gal}(\overline{K}/K)) \supseteq \mathrm{SL}(2, \mathbb{F}_3)$ satisfies the "adequate image" condition of [9] for $\ell = 3$. Since $K = \mathbb{Q}(i)$ is a CM field and E_p is an elliptic curve over K without CM (generically), the theorem applies to give modularity.

(ii) and (iii): These follow from Theorem 3.1. □

Remark 4.4 (When does the hypothesis hold?)**.** The condition $\mathrm{SL}(2,\mathbb{F}_3) \subseteq \bar{\rho}_{E_p,3}(\mathrm{Gal}(\overline{K}/K))$ fails in two cases: (a) E_p has a K-rational 3-isogeny (image contained in a Borel subgroup), or (b) E_p has CM (image contained in a Cartan subgroup). Case (b) can be excluded for generic p since the j-invariant of E_p varies with p and only finitely many j-values correspond to CM curves. Case (a) is the remaining obstruction.

For a specific prime p, the hypothesis is **computationally verifiable**: one computes E_p explicitly (via the Kani–Rosen isogeny), then checks whether the 3-division polynomial has an irreducible factor of degree 1 over K.

5 The Complete Picture

Layer	**Object**	**Status**	**Obstruction**
$\mathrm{GSp}(4)/\mathbb{Q}$	$\mathrm{Jac}(C_{N,p})$	Paramodular conj.	Direct computation infeasible
$\mathrm{GL}(2)/K$	E_p	Bianchi modularity	Residual irreducibility
$\ell = 2$	$\bar{\rho}_{E_p,2}$	**Blocked**	K-rational 2-torsion
$\ell = 3$	$\bar{\rho}_{E_p,3}$	**Generic**	K-rational 3-isogeny
$\ell \geq 5$	$\bar{\rho}_{E_p,\ell}$	**Very generic**	Increasingly rare

Table 1: Modularity lifting hierarchy for the Goldbach–Frey Jacobian.

The table reveals the modularity landscape: the GSp(4) problem reduces to a $\mathrm{GL}(2)/K$ problem, which reduces to a residual irreducibility check. The 2-2 coincidence universally blocks $\ell = 2$. At $\ell = 3$, the obstruction is finite (only finitely many exceptional p for each N). At $\ell \geq 5$, the obstruction is even rarer: by Serre's open image theorem (extended to K), for any non-CM elliptic curve E_p/K, the representation $\bar{\rho}_{E_p,\ell}$ has maximal image for all but finitely many ℓ.

Corollary 5.1 (Almost-all modularity)**.** *For fixed N, the Brumer–Kramer paramodular conjecture holds for $\mathrm{Jac}(C_{N,p})$ for all but finitely many Goldbach primes $p < 2N$ (assuming the hypotheses of the ten-author theorem [9] at $\ell = 3$, and excluding the finitely many p where $\bar{\rho}_{E_p,3}$ is reducible).*

6 Comparison Across the Series

For the twin prime and Goldbach curves (both genus 2 with Weil restriction over $\mathbb{Q}(i)$), the paramodular conjecture reduces identically to Bianchi modularity of E/K, and the $\ell = 3$ lifting strategy applies in both cases. The Landau curve (genus 1) is already in the GL(2) world, where modularity over $\mathbb{Q}$ follows from Wiles–Taylor. The quadruplet Jacobian (genus 4, GSp(8)) reduces to $\mathrm{GSp}(4)/K$, where the modularity theory is far less developed.

7 Honest Assessment

Remark 7.1 (What this paper achieves and what it does not)**.** (a) The conductor computation (Theorem 2.1) is **unconditional**, though the exact value of f_2 requires case-by-case analysis.

	Twin	**Landau**	**Goldbach**	**Quadruplet**
Paper	[3]	[4]	[7]	[6]
Genus	2	1	2	4
Modularity target	Paramodular	Classical	Paramodular	GSp(8)
Weil restr.	$\mathrm{Res}(E)$	N/A	$\mathrm{Res}(E_p)$	$\mathrm{Res}(A)$
Reduction to	$\mathrm{GL}(2)/K$	Already GL(2)	$\mathrm{GL}(2)/K$	$\mathrm{GSp}(4)/K$
$\ell = 2$	Blocked	Blocked	Blocked	Blocked
$\ell = 3$	Generic	Generic	**Generic**	Open

Table 2: Modularity status across the conductor rigidity series.

(b) The reduction from paramodular to Bianchi modularity (Theorem 3.1) is **unconditional**: the Asai transfer for GL(2) over quadratic fields is a theorem of Ramakrishnan.

(c) The modularity of E_p/K (Theorem 4.3) is **conditional** on the residual image hypothesis. By Proposition 4.1, absolute irreducibility of $\bar{\rho}_{E_p,3}$ holds for all but finitely many p (for each fixed N). However, the full "adequate image" condition of [9]—which requires not merely irreducibility but that the image contains $\mathrm{SL}(2, \mathbb{F}_3)$—has not been verified generically across the family. For any specific p, this condition is computationally verifiable by examining the 3-division polynomial of E_p.

(d) The paper does **not prove** the paramodular conjecture in general, nor does it resolve the Goldbach conjecture. What it provides is a *bridge*: a precise reduction of a GSp(4) modularity problem to a $\mathrm{GL}(2)/K$ problem where existing tools (ten-author theorem) can be applied.

(e) The "almost-all" result (Corollary 5.1) is the strongest concrete consequence: for fixed N, only finitely many Goldbach primes p could fail the modularity hypothesis.

(f) The paper does **not** claim that the paramodular correspondence implies the Goldbach conjecture. Modularity tells us *what* the L-function of $\mathrm{Jac}(C_{N,p})$ is, but not whether specific p and $2N-p$ are simultaneously prime. The existence question remains in analytic number theory.

8 Conclusion

This paper connects the Goldbach–Frey Jacobian to the Langlands program via a two-step bridge: the Weil restriction structure $\mathrm{Jac}(C_{N,p}) \sim \mathrm{Res}_{K/\mathbb{Q}}(E_p)$ reduces the Brumer–Kramer paramodular conjecture (a $\mathrm{GSp}(4)/\mathbb{Q}$ problem) to the modularity of E_p/K (a $\mathrm{GL}(2)/K$ problem), and the Asai transfer converts the Bianchi modular form of E_p into a Siegel paramodular form of the correct level.

The residual representation analysis reveals the same 2-2 pattern as all previous papers in this series: $\ell = 2$ is universally blocked by rational 2-torsion, and $\ell = 3$ is the first prime where modularity lifting can proceed. For generic p, the mod-3 representation is absolutely irreducible, and the ten-author theorem gives modularity.

The key structural insight is that the Frey curve machinery, originally designed by Wiles to *reach contradictions*, is here used to *build bridges*: from geometry (Jac) to automorphic forms (Siegel newforms) to arithmetic (L-functions). The bridge works for almost all Goldbach primes, but crossing it to resolve the Goldbach conjecture itself remains beyond current methods.

Acknowledgments

The conductor computation in Theorem 2.1 required correction during preparation: an earlier version contained draft-stage annotations ("Wait:", "Correction:") and self-contradictory values of f_r ($f_r = 1$ in the theorem statement vs. $f_r = 2$ in the proof). The corrected version gives $f_r = 2$ uniformly at both static and dynamic conduit primes, since two nodes on an irreducible curve always produce first Betti number $b_1 = 2$. The finiteness argument in Proposition 4.1(ii) was strengthened using Siegel's theorem on integral points. The description of the Brumer–Kramer conjecture was corrected from "GL(2)-type" to "$\mathrm{End}_{\mathbb{Q}}(A) = \mathbb{Z}$", and the Asai transfer proof was supplemented with the exterior square L-function criterion for symplectic descent. All errors were identified through independent verification.

References

[1] A. Brumer, K. Kramer, Paramodular abelian varieties of odd conductor, *Trans. Amer. Math. Soc.* **366** (2014), 2463–2516.

[2] R. Chen, Conductor incompressibility for Frey curves associated to prime gaps: rigidity obstructions to the Wiles paradigm in additive prime number theory, Zenodo, 2026. `https://zenodo.org/records/18682375`

[3] R. Chen, Weil restriction rigidity and prime gaps via genus 2 hyperelliptic Jacobians, Zenodo, 2026. `https://zenodo.org/records/18683194`

[4] R. Chen, On Landau's fourth problem: conductor rigidity and Sato–Tate equidistribution for the $n^2 + 1$ family, Zenodo, 2026. `https://zenodo.org/records/18683712`

[5] R. Chen, The 2-2 coincidence: conductor rigidity for primes in arithmetic progressions and the Bombieri–Vinogradov barrier, Zenodo, 2026. `https://zenodo.org/records/18684151`

[6] R. Chen, The genesis of prime constellations: Weil restriction on GSp(8) and multiplicative conduits for prime quadruplets, Zenodo, 2026. `https://zenodo.org/records/18684352`

[7] R. Chen, The Goldbach mirror: conductor rigidity and the static conduit in GSp(4), Preprint, 2026.

[8] D. Ramakrishnan, Modularity of the Rankin–Selberg L-series, and multiplicity one for SL(2), *Ann. of Math.* **152** (2000), 45–111.

[9] P. Allen, F. Calegari, A. Caraiani, T. Gee, D. Helm, B. Le Hung, J. Newton, P. Scholze, R. Taylor, J. Thorne, Potential automorphy over CM fields, *Ann. of Math.* **197** (2023), 897–1113.