

Crossover from Discrete to Continuous Quantum Geometry: Running Spectral Dimensions, Noncommutative Spectral Triples, and Critical Phase Transitions

Ruqing Chen

GUT Geoservice Inc., Montreal, Quebec, Canada

ruqing@hotmail.com

DOI: [10.5281/zenodo.23195881](https://doi.org/10.5281/zenodo.23195881)

Abstract

The spectral dimension $d_s(\tau) = -2 \, d \ln P / d \ln \tau$ of a diffusion process is widely used to diagnose dimensional reduction in quantum gravity. We show that it is exactly twice an equipartition count: $d_s(\tau) = 2\tau \langle \lambda \rangle_\tau$, the Gibbs average of the Laplacian at inverse temperature τ . Two consequences make precise how cutoff effects are separated from physical scaling. First, any discretisation with a bounded spectrum gives $d_s \rightarrow 0$ as $\tau \rightarrow 0$, so physical values must be read from scaling plateaus between the cutoff and finite-volume regimes. Second, a finite graph gives $d_s \rightarrow 0$ as $\tau \rightarrow \infty$.

For the hypercubic lattice we obtain the exact running dimension $d_s = 2dx [1 - I_1(x)/I_0(x)]$, with $x = 2\tau/a^2$. It rises from zero, overshoots the topological dimension (maximum $1.2178d$) and approaches d from above as $d(1 + a^2/8\tau)$. For the higher-derivative dispersion $E = k^2(1 + \ell^2 k^2)$ in four dimensions, which is the scaling relevant to asymptotic safety, Hořava gravity and quadratic gravity, we give the global interpolation in closed form, $P \propto \tau^{-1} [1 - \sqrt{\pi} s e^{s^2} \operatorname{erfc}(s)]$ with $s = \sqrt{\tau}/2\ell$, which runs from $d_s = 2$ to $d_s = 4$. Dimensional splitting $d_s \neq d_H$ is traced to an anomalous walk dimension, $d_s = 2d_H/d_w$. On the Sierpinski gasket this gives $d_s = 2 \ln 3 / \ln 5$, with log-periodic oscillations of period $\ln 5$.

On the noncommutative side, we prove that the lattice Hodge–Dirac operator $d + d^*$ on the torus T^d converges to the continuum Hodge–Dirac operator in norm-resolvent sense with error $O(a)$, whereas the naive symmetric-difference Dirac operator does not converge, because of fermion doubling. For the incidence Dirac operator of a graph, the Connes distance is bounded between $a d_{\text{hop}} / \sqrt{\deg^+}$ (with $\deg^+$ the maximal in-degree) and $a d_{\text{hop}}$. On $\mathbb{Z}^d$ the hop metric tends to the ℓ^1 metric, not the Euclidean one, whereas a linear test function shows that $d_C \geq |p - q|_2$ exactly, for every lattice spacing.

All statements are checked numerically by exact diagonalisation.

Contents

1	Introduction	2
2	The spectral dimension as an equipartition count	3
2.1	General identity	3
2.2	The hypercubic lattice	3
2.3	Higher-derivative dispersion: a closed-form global interpolation	4
3	Fractal graphs and dimensional splitting	4
3.1	Walk dimension	4
3.2	Sierpinski gasket	5
3.3	Critical phase transitions and effective functionals	5

4 Noncommutative spectral triples: from lattice to continuum	6
4.1 Norm-resolvent convergence of the lattice Hodge–Dirac operator	6
4.2 Connes distance on graphs	6
5 Discussion and limitations	7
A Numerical methods	9

1 Introduction

Several approaches to quantum gravity find that the effective dimension of spacetime depends on the scale at which it is probed, and that it decreases towards two at short distances. Examples include causal dynamical triangulations (CDT) [1], asymptotic safety [2], Hořava–Lifshitz gravity [3], loop quantum gravity and spin foams [4], quantum spacetimes with fractal properties [5] and multifractional theories [6]; see [7] for a review. The usual diagnostic is the spectral dimension

$$d_s(\tau) = -2 \frac{d \ln P(\tau)}{d \ln \tau}, \quad P(\tau) = \frac{1}{V} \text{Tr } e^{-\tau \Delta}, \quad (1)$$

defined by the return probability of a diffusion process. Different definitions of the diffusion can give different answers [8], and the interpretation of d_s requires care.

This paper studies the crossover from discrete to continuous geometry in three complementary languages: heat kernels, noncommutative spectral triples [9, 10], and fractal graphs. Our main results are the following.

1. *Spectral dimension as equipartition* (Theorem 1). $d_s(\tau) = 2\tau \langle \lambda \rangle_\tau$ exactly. As consequences, the ultraviolet value $d_s \rightarrow 0$ found for any discretisation is a cutoff artefact, a finite graph has $d_s \rightarrow 0$ also in the infrared, and plateaus of d_s correspond to power laws in the spectral density.
2. *Exact running dimensions* (Propositions 2.1 and 2.2). For the hypercubic lattice the running is non-monotonic and overshoots d . For the dispersion $E = k^2(1 + \ell^2 k^2)$ we give a closed-form global interpolation from 2 to 4 in $d = 4$, and a general- d formula in terms of parabolic-cylinder functions.
3. *Dimensional splitting* (Section 3). $d_s \neq d_H$ arises from an anomalous walk dimension, $d_s = 2d_H/d_w$ [11, 12]. In quantum gravity this corresponds to an anomalous scaling of the kinetic operator at fixed volume scaling. Self-similar graphs add log-periodic oscillations [13].
4. *Spectral triples* (Section 4). We prove $O(a)$ norm-resolvent convergence of the lattice Hodge–Dirac operator on T^d , show that the naive lattice Dirac operator fails to converge because of fermion doubling [14], bound the Connes distance on graphs, and prove $d_C \geq |p - q|_2$ on $a\mathbb{Z}^d$.

We also make three points precise. A vanishing ultraviolet spectral dimension is generic for any discretisation with a bounded spectrum. This is well known in the CDT, spin-foam and tensor-model literature, where physical values are read from scaling plateaus or from continuum limits taken at fixed physical scale; Corollary 2.1 formalises that separation. The hop metric of a hypercubic lattice converges to the ℓ^1 metric, not to the Riemannian distance. Lattice corrections to heat-kernel coefficients are non-universal ultraviolet terms, not conformal anomalies.

This is the fourth paper of the author’s programme of scale-dependent analysis (“pan-scale analytics”), which also covers complex dimensions in holographic entanglement [19], multifractal entanglement spectra [20], and algorithmic complexity under scaling [21], and is continued by a thermodynamic analysis of scale [22]. Connections are discussed in Section 5.

2 The spectral dimension as an equipartition count

2.1 General identity

Theorem 1 (Equipartition form of the spectral dimension). *Let $\Delta \geq 0$ have spectral measure ρ with $P(\tau) = \int e^{-\tau\lambda} d\rho(\lambda) < \infty$ for $\tau > 0$. Then*

$$d_s(\tau) = 2\tau \langle \lambda \rangle_\tau, \quad \langle \lambda \rangle_\tau = \frac{\int \lambda e^{-\tau\lambda} d\rho(\lambda)}{\int e^{-\tau\lambda} d\rho(\lambda)}. \quad (2)$$

In words, $d_s/2$ is the mean energy, in units of the temperature $1/\tau$, of a canonical ensemble whose levels are the eigenvalues of Δ .

Proof. Differentiate under the integral: $-\tau d \ln P / d\tau = \tau \int \lambda e^{-\tau\lambda} d\rho / \int e^{-\tau\lambda} d\rho$. $\square$

For a continuum Laplacian, $\rho(\lambda) \propto \lambda^{d/2-1}$, and each of the d quadratic degrees of freedom contributes $1/(2\tau)$ to $\langle \lambda \rangle_\tau$, which gives $d_s = d$. This is equipartition. The same thermodynamic reading of heat-kernel and Rényi quantities underlies the companion paper [20].

Corollary 2.1. (i) *If $\text{spec } \Delta \subset [0, \lambda_{\max}]$, then $0 \leq d_s(\tau) \leq 2\tau\lambda_{\max}$, and so $d_s \rightarrow 0$ as $\tau \rightarrow 0$.*

(ii) *On a finite connected graph, $d_s(\tau) \leq 2\tau(N-1)\lambda_{\max}e^{-\tau\lambda_1}$, where λ_1 is the smallest non-zero eigenvalue. Hence $d_s \rightarrow 0$ as $\tau \rightarrow \infty$.*

(iii) *If $\rho([0, \lambda]) \asymp \lambda^{\delta/2}$ for $\lambda_- \ll \lambda \ll \lambda_+$, then $d_s(\tau) \approx \delta$ for $1/\lambda_+ \ll \tau \ll 1/\lambda_-$.*

Proof. (i) Use $\langle \lambda \rangle_\tau \leq \lambda_{\max}$. (ii) The zero mode has weight $1/N$ in P and contributes nothing to the numerator, which is at most $(N-1)\lambda_{\max}e^{-\tau\lambda_1}/N$. (iii) In the scaling window the Gibbs average of a power-law density gives $\langle \lambda \rangle_\tau = \delta/(2\tau)$, with corrections from the edges of the window suppressed exponentially or by powers of $\tau\lambda_{\pm}$. $\square$

Part (i) states in exact form what is familiar from lattice studies of quantum gravity: a vanishing spectral dimension at the shortest scales is produced by any regularisation with a bounded spectrum. Physically meaningful ultraviolet values, such as $d_s \approx 2$, are therefore identified as plateaus at intermediate τ , well above the cutoff scale and below the finite-volume scale of part (ii), or through continuum limits taken at fixed physical scale.

2.2 The hypercubic lattice

Proposition 2.1 (Exact lattice running). *For the graph Laplacian of $a\mathbb{Z}^d$, with dispersion $E(k) = \sum_i \frac{2}{a^2}(1 - \cos k_i a)$, the return probability per site is $P(\tau) = [e^{-x}I_0(x)]^d$ with $x = 2\tau/a^2$, and*

$$d_s(\tau) = 2dx \left(1 - \frac{I_1(x)}{I_0(x)}\right), \quad x = \frac{2\tau}{a^2}. \quad (3)$$

As $\tau \rightarrow 0$, $d_s = 2dx + O(x^2)$. As $\tau \rightarrow \infty$,

$$P(\tau) = (4\pi\tau)^{-d/2}a^d \left(1 + \frac{d a^2}{16\tau} + O(a^4/\tau^2)\right), \quad d_s(\tau) = d \left(1 + \frac{a^2}{8\tau} + O(a^4/\tau^2)\right). \quad (4)$$

Consequently d_s approaches d from above. Its maximum is $d_s^{\max} = 1.2178d$, attained at $\tau \simeq 0.84a^2$.

Proof. The one-dimensional factor is $\frac{1}{2\pi} \int_{-\pi}^{\pi} e^{-x(1-\cos\theta)} d\theta = e^{-x}I_0(x)$, which gives the product form. Then $\frac{d}{dx} \ln[e^{-x}I_0(x)] = I_1/I_0 - 1$, and (3) follows. The asymptotic forms follow from $I_\nu(x) = \frac{e^x}{\sqrt{2\pi x}} \left(1 - \frac{4\nu^2-1}{8x} + \dots\right)$ [17], which gives $1 - I_1/I_0 = \frac{1}{2x} + \frac{1}{8x^2} + O(x^{-3})$. The maximum was located numerically from (3). $\square$

Remark (Origin of the overshoot). Per dimension and with $a = 1$, the lattice density of states is $\rho_{\text{lat}}(\lambda) = 1/(\pi\sqrt{\lambda(4-\lambda)})$ on $(0, 4)$, while the continuum has $\rho_{\text{cont}}(\lambda) = 1/(2\pi\sqrt{\lambda})$. The ratio $\rho_{\text{lat}}/\rho_{\text{cont}} = 2/\sqrt{4-\lambda}$ is at least 1 and diverges at the zone boundary $\lambda = 4$, where the group velocity vanishes (a Van Hove singularity). Relative to the continuum, the lattice therefore places extra weight at large λ . For $\tau \sim a^2$ the Gibbs mean $\langle \lambda \rangle_\tau$ exceeds the equipartition value $d/(2\tau)$, so $d_s > d$, until the bounded spectrum takes over for $\tau \ll a^2$.

The a^2/τ term is a non-universal ultraviolet correction to the heat-kernel coefficients. It is not a conformal anomaly: the anomaly is controlled by the coefficient $a_{d/2}$ of the Seeley–DeWitt expansion, which is a curvature invariant and vanishes on a flat torus.

2.3 Higher-derivative dispersion: a closed-form global interpolation

Effective dispersions $E \propto k^2(1 + \ell^2 k^2)$ arise from the transverse-traceless propagator of quadratic gravity [15]. Their ultraviolet scaling $E \propto k^4$ reproduces $d_s = 2$ in four dimensions, as found in asymptotic safety from the anomalous dimension $\eta_N = -2$ [2] and in Hořava gravity with $z = 3$ [3].

Proposition 2.2 (Global interpolation). *For $E(k) = k^2(1 + \ell^2 k^2)$ in $\mathbb{R}^d$ with flat measure,*

$$P(\tau) \propto \int_0^\infty u^{\frac{d}{2}-1} e^{-\tau u - \tau \ell^2 u^2} du = \Gamma(\frac{d}{2}) (2\tau \ell^2)^{-d/4} e^{\tau/(8\ell^2)} D_{-d/2}\left(\frac{\sqrt{\tau}}{\sqrt{2}\ell}\right), \quad (5)$$

with D_ν the parabolic-cylinder function [18]. For $d = 4$ this reduces to

$$P(\tau) \propto \frac{1}{\tau} \left[1 - \sqrt{\pi} s e^{s^2} \text{erfc}(s) \right], \quad s = \frac{\sqrt{\tau}}{2\ell}, \quad (6)$$

so $d_s \rightarrow 2$ for $\tau \ll \ell^2$ and $d_s \rightarrow 4$ for $\tau \gg \ell^2$. More generally, $d_s = 2d/\alpha$ in a regime where $E \propto k^\alpha$.

Proof. Set $u = k^2$ and use [18, 3.462.1]. For $d = 4$, evaluate $\int_0^\infty u e^{-\beta u^2 - \gamma u} du$ directly, with $\beta = \tau \ell^2$ and $\gamma = \tau$. The limits follow from $\sqrt{\pi} s e^{s^2} \text{erfc}(s) = 1 - \frac{1}{2s^2} + O(s^{-4})$ as $s \rightarrow \infty$, and from $\rightarrow 0$ as $s \rightarrow 0$. For a pure power law $E \propto k^\alpha$, Theorem 1 gives $\langle E \rangle_\tau = d/(\alpha\tau)$. $\square$

Remark (Ghost pole and the meaning of P). In Lorentzian signature the transverse-traceless propagator of quadratic gravity decomposes as $\frac{1}{k^2(1+\ell^2 k^2)} = \frac{1}{k^2} - \frac{1}{k^2 + \ell^{-2}}$. The second term has negative residue and describes a massive spin-2 ghost [15]. Proposition 2.2 uses $E = k^2(1 + \ell^2 k^2)$ only as an effective Euclidean dispersion that models the ultraviolet scaling of the spectral density, and it says nothing about unitarity. Moreover, the heat kernel of a fourth-order operator is not positive in position space, so $P(\tau)$ is then a spectral trace rather than a return probability. This is one reason why different definitions of diffusion can lead to different spectral dimensions [8].

Combining (6) with a momentum cutoff $|k| \leq \pi/a$ gives three regimes: $d_s \rightarrow 0$ for $\tau \ll a^4/\ell^2$, $d_s \approx 2$ in between, and $d_s \rightarrow 4$ for $\tau \gg \ell^2$. Numerically, with $\ell = 10a$, the cutoff result agrees with (6) to 10^{-4} for $\tau \geq 10^{-2}a^2$. It equals 0.001 at $\tau = 10^{-7}a^2$, averages 2.03 on $10^{-2} \leq \tau/a^2 \leq 1$, and reaches 3.999 at $\tau = 10^6 a^2$. We also checked the parabolic-cylinder identity (5) numerically to 10^{-11} (Fig. 1, bottom right).

3 Fractal graphs and dimensional splitting

3.1 Walk dimension

Let d_H be the Hausdorff (volume-growth) dimension and d_w the walk dimension, defined by $\langle r^2 \rangle \sim \tau^{2/d_w}$. Then [11, 12]

$$d_s = \frac{2d_H}{d_w}. \quad (7)$$

Dimensional splitting, $d_s < d_H$, is thus equivalent to subdiffusion, $d_w > 2$. For a dispersion $E \propto k^\alpha$ on a space with ordinary volume growth $d_H = d$, the diffusion is anomalous with $d_w = \alpha$, and (7) gives $d_s = 2d/\alpha$, consistent with Proposition 2.2. This makes precise how quantum-gravity proposals can combine $d_H \approx 4$ with $d_s \approx 2$. The splitting reflects an anomalous scaling of the kinetic operator ($\alpha = 4$) at unchanged volume scaling. It does not require the geometry to be fractal in the sense of volume growth.

3.2 Sierpinski gasket

For the Sierpinski gasket, $d_H = \ln 3 / \ln 2$ and $d_w = \ln 5 / \ln 2$, so $d_s = 2 \ln 3 / \ln 5 \simeq 1.3652$ [12]. Spectral decimation rescales eigenvalues by 5 under the geometric scaling by 2. As a result $P(\tau) = \tau^{-d_s/2} G(\ln \tau)$ with G periodic of period $\ln 5$, and $d_s(\tau)$ oscillates log-periodically. This is the heat-kernel counterpart of the complex dimensions studied in [13, 19].

Analysis on fractals. The graph Laplacians used here approximate the Laplacian of Kigami’s analysis on fractals [24, 25]. The renormalised graph energies converge to a self-similar Dirichlet form, and the graph Laplacians, rescaled by 5^m up to a normalisation constant, converge to the Laplacian with respect to the self-similar measure. Its spectrum is obtained by spectral decimation [26]. Kigami and Lapidus proved Weyl-type asymptotics $N(\lambda) \asymp \lambda^{d_s/2}$ with $d_s = 2 \ln 3 / \ln 5$, using renewal theory. In the arithmetic (“lattice”) self-similar case, where the logarithms of the contraction ratios are commensurable, the prefactor is periodic in $\ln \lambda$ and does not converge [27]. Here “lattice” is the renewal-theoretic term and is unrelated to the spatial lattices of Section 2. The log-periodic heat kernel is the Laplace-transform counterpart of this statement.

Numerical test. We diagonalised the graph Laplacian of the level-7 gasket ($N = 3282$ vertices). The finite-size scale is $1/\lambda_1 \simeq 4.3 \times 10^3$. Averaged over exactly three periods, d_s is 1.387, 1.357 and 1.347 for $\tau \in [1, 125]$, $[2, 250]$ and $[3, 375]$, bracketing $2 \ln 3 / \ln 5$. The drift reflects the ultraviolet transient on the left and the onset of finite-size effects on the right. After linear detrending, the dominant oscillation frequency is 1.02 cycles per unit of $\log_5 \tau$, against the predicted value 1 (Fig. 2, left).

3.3 Critical phase transitions and effective functionals

In CDT, macroscopic four-dimensional geometry emerges in one phase of the path integral, and a second-order transition between phases has been reported [16], which allows a continuum limit to be considered. A natural effective description of the near-critical regime, consistent with the scale dependence of d_s , is a higher-curvature functional of the form

$$\Gamma_k[g] = \int d^4x \sqrt{g} \left[\frac{1}{16\pi G_k} (R - 2\Lambda_k) + \alpha_k R^2 + \beta_k C_{\mu\nu\rho\sigma} C^{\mu\nu\rho\sigma} \right], \quad (8)$$

with scale-dependent couplings. On flat space its transverse-traceless kinetic operator has the form $k^2(1 + \ell^2 k^2)$ with $\ell^2 \propto \beta G$ [15], which is the dispersion of Proposition 2.2. We stress that (8) is a truncation motivated by these properties. We do not derive it from the CDT transition, whose universality class remains open.

Remark (Isotropy versus Lifshitz anisotropy). CDT imposes a preferred time foliation. It has therefore been proposed that its continuum limit near the second-order transition is described by an anisotropic Hořava–Lifshitz theory [3, 23]. In such a theory only spatial derivatives are of higher order, with dynamical exponent $z = 3$ in the ultraviolet, while time derivatives remain of second order, which avoids Ostrogradsky ghosts. The isotropic functional (8) should therefore be read as a toy truncation. The anisotropic dispersion $\omega^2 \sim k^2 + k^6/M^4$ gives $d_s = 1 + 3/z = 2$ in the ultraviolet [3], the same value as the isotropic k^4 case, so the spectral dimension alone does not distinguish the two.

4 Noncommutative spectral triples: from lattice to continuum

4.1 Norm-resolvent convergence of the lattice Hodge–Dirac operator

Let $T^d = (\mathbb{R}/2\pi\mathbb{Z})^d$ and $a = 2\pi/N$. The cubical lattice of spacing a carries discrete differential forms, the coboundary d_a and its adjoint, and the Hodge–Dirac operator $D_a = d_a + d_a^*$ on $\mathcal{H}_a = \bigoplus_p \Lambda_a^p$. We use the spectral triple $(C(\text{lattice}), \mathcal{H}_a, D_a)$. The continuum counterpart is $D = d + d^*$ on $L^2\Lambda^\bullet(T^d)$. Let J_a identify the lattice Fourier mode k in the Brillouin zone $B_a = (-\pi/a, \pi/a]^d$ with the corresponding continuum mode and form component.

Theorem 2 (Norm-resolvent convergence). *For every $z \in \mathbb{C} \setminus \mathbb{R}$ there is $C(z)$ such that*

$$\|J_a(D_a - z)^{-1}J_a^* - (D - z)^{-1}\| \leq C(z)a. \quad (9)$$

Proof. By the Künneth structure of the cubical complex of T^d , D_a^2 acts on each form component as the scalar lattice Laplacian. Hence D_a has eigenvalues $\pm s_a(k)$, with $s_a(k)^2 = \sum_i \frac{4}{a^2} \sin^2 \frac{k_i a}{2}$ for $k \in B_a$, while D has eigenvalues $\pm |k|$ with the same multiplicities. For $|k_i a| \leq \pi$ we have $\frac{2}{\pi}|k_i| \leq \frac{2}{a}|\sin \frac{k_i a}{2}| \leq |k_i|$ and $0 \leq |k_i| - \frac{2}{a}|\sin \frac{k_i a}{2}| \leq a^2|k_i|^3/24$. Therefore $\frac{2}{\pi}|k| \leq s_a(k) \leq |k|$ and $|k| - s_a(k) = \frac{|k|^2 - s_a^2}{|k| + s_a} \leq a^2|k|^3/12$. Take $z = iy$ for simplicity. For a mode $k \neq 0$ in B_a (the zero mode contributes nothing, since $s_a(0) = |0| = 0$),

$$\left| \frac{1}{\pm s_a - z} - \frac{1}{\pm |k| - z} \right| = \frac{||k| - s_a|}{|s_a \mp z| ||k| \mp z|} \leq \frac{a^2|k|^3/12}{s_a|k|} \leq \frac{\pi a^2|k|}{24} \leq \frac{\pi^2 \sqrt{d}}{24} a, \quad (10)$$

since $|k| \leq \pi\sqrt{d}/a$ in B_a . For modes outside B_a , which lie in the kernel of J_a^* , the continuum resolvent is bounded by $1/|k| \leq a/\pi$. Both operators are diagonal in the Fourier basis, so the norm is the supremum of these bounds. General z follows from the resolvent identity. $\square$

Counterexample: fermion doubling. The naive Dirac operator $\sum_\mu \gamma^\mu \nabla_\mu^{\text{sym}}$ has eigenvalues $\pm (\sum_\mu \sin^2(k_\mu a)/a^2)^{1/2}$, which vanish at every corner $k_\mu \in \{0, \pi/a\}$ of the Brillouin zone. At $k = (\pi/a, 0, \dots)$ the lattice resolvent equals $-1/z$, whereas the continuum resolvent at $|k| = \pi/a$ is $O(a)$. The difference therefore stays of order one, and there is no norm-resolvent convergence. This is the Nielsen–Ninomiya obstruction [14]. Numerically, on the circle at $z = i$, the Hodge–Dirac error decreases as 1.4×10^{-1} , 3.2×10^{-2} , 7.9×10^{-3} , 2.0×10^{-3} , 4.9×10^{-4} for $N = 16, \dots, 4096$, i.e. as $O(a)$. The naive error stays at 1.00 (Fig. 2, middle).

4.2 Connes distance on graphs

The Connes distance is $d_C(p, q) = \sup\{|f(p) - f(q)| : \|[D, f]\| \leq 1\}$ [9, 10]. For a graph with edges oriented and weighted by $1/a$, take the incidence Dirac operator $D = \begin{pmatrix} 0 & \partial^* \\ \partial & 0 \end{pmatrix}$ on $\ell^2(V) \oplus \ell^2(E)$, with $(\partial g)(e) = (g(e^+) - g(e^-))/a$. Functions act on edges by $f(e) = f(e^-)$.

Proposition 4.1 (Connes distance bounds). *Let $\deg^+$ be the maximal in-degree. Then $\|[D, f]\|^2 = \max_v \sum_{e:e^+=v} |\nabla_e f|^2$ with $\nabla_e f = (f(e^+) - f(e^-))/a$, and*

$$\frac{a d_{\text{hop}}(p, q)}{\sqrt{\deg^+}} \leq d_C(p, q) \leq a d_{\text{hop}}(p, q). \quad (11)$$

In one dimension, with consistent orientation, $d_C(p, q) = |p - q|$ exactly.

Proof. $[\partial, f]g(e) = \nabla_e f g(e^+)$ is a weighted map from vertices to edges, and its norm is $\max_v (\sum_{e:e^+=v} |\nabla_e f|^2)^{1/2}$. The same holds for $[\partial^*, f]$, which is minus its adjoint. Hence $\|[D, f]\| \leq 1$ implies $|\nabla_e f| \leq 1$ on every edge, so $d_C \leq a d_{\text{hop}}$. Conversely, $f = a d_{\text{hop}}(p, \cdot)/\sqrt{\deg^+}$ is admissible. In one dimension $\deg^+ = 1$. $\square$

On $a\mathbb{Z}^d$, $a d_{\text{hop}}$ converges to the ℓ^1 distance, not to the Euclidean distance; the ratio is $\sqrt{2}$ along a diagonal of $\mathbb{Z}^2$. The hop metric therefore does not deform into the Riemannian geodesic distance. The Connes distance behaves differently.

Proposition 4.2 (Euclidean lower bound on $a\mathbb{Z}^d$). *For the incidence Dirac operator on $a\mathbb{Z}^d$, with edges oriented along $+e_i$, one has $d_C(p, q) \geq |p - q|_2$ for all lattice points p, q and every $a > 0$.*

Proof. Let $\mathbf{n} = (p - q)/|p - q|_2$ and $f(x) = \mathbf{n} \cdot x$. On every edge along e_i , $\nabla_e f = n_i$. Each vertex has exactly one incoming edge in each direction, so by Proposition 4.1, $\|[D, f]\|^2 = \sum_i n_i^2 = 1$. Hence f is admissible and $d_C(p, q) \geq f(p) - f(q) = |p - q|_2$. If the algebra is required to consist of bounded functions, replace f by its clamp $\min\{\max\{f, f(q)\}, f(p)\}$. Clamping does not increase any edge difference, so the constraint still holds, and the values at p and q are unchanged. $\square$

The supremum defining d_C allows a different test function for each pair of points, and a linear function already gives the Euclidean distance. On $\mathbb{Z}^2$, with edges oriented along $+x$ and $+y$, the upwind recursion $f(v) = \frac{A+B}{2} + \frac{1}{2}\sqrt{2 - (A - B)^2}$, where A and B are the values at the two predecessors, defines an admissible f . It is a monotone discretisation of the eikonal equation $|\nabla f| = 1$ and gives another lower bound on d_C . Relative to the Euclidean distance, this bound equals 1.0186, 1.0063 and 1.0020 at 45° for grids of 50, 200 and 800 steps, and exactly 1 along the axes (Fig. 2, right). At finite resolution it is therefore *strictly* larger than the bound of Proposition 4.2, so $d_C > |p - q|_2$ off the axes, and the excess decreases as the grid is refined.

Conjecture 1. *For the incidence Dirac operator on $a\mathbb{Z}^d$, $d_C(p, q) \rightarrow |p - q|_2$ as $a \rightarrow 0$.*

By Proposition 4.2 the conjecture is equivalent to $\limsup_{a \rightarrow 0} d_C(p, q) \leq |p - q|_2$. An upper bound sharper than ℓ^1 is missing. A natural route is to extend admissible lattice functions to Lipschitz functions on $\mathbb{R}^d$, by multilinear interpolation or mollification, and to use the continuum bound. The difficulty is that admissibility constrains gradients vertex by vertex. Near a corner of a lattice cube, the edges that determine the different components of the interpolated gradient end at *different* vertices, so each component is only bounded by 1 and the interpolant can have $|\nabla F|$ up to $\sqrt{d}$. This reproduces only the ℓ^1 -type bound. The same vertex-wise decoupling is what allows $d_C > |p - q|_2$ at finite resolution, so an improvement must show that such local excesses do not accumulate along paths as $a \rightarrow 0$.

5 Discussion and limitations

1. The identity $d_s = 2\tau\langle\lambda\rangle_\tau$ shows that the spectral dimension is a property of the spectral density alone. It is insensitive to eigenvectors, and hence to geometric information beyond the spectrum. Two geometries with the same spectrum have the same running d_s .
2. The ultraviolet zero and the infrared drop in Corollary 2.1 are generic artefacts. Reported ultraviolet values of d_s should be read from plateaus well above the cutoff scale and below the finite-size scale.
3. Theorem 2 concerns flat tori. Extending norm-resolvent convergence to curved manifolds and random triangulations, as in CDT, and controlling fermion doubling there, are open problems.
4. For the Connes distance on $a\mathbb{Z}^d$ we prove $d_C \geq |p - q|_2$. The conjectured Euclidean limit requires a matching upper bound.
5. The functional (8) is a truncation, not a derivation from the CDT transition. It is isotropic, whereas CDT has a preferred foliation, and its higher-derivative pole is a ghost. Accordingly, Proposition 2.2 is used only as an effective spectral model.




Figure 1: Top left: integrated spectral densities of the 64×64 periodic lattice and the level-7 Sierpinski gasket. Top right: return probabilities. Bottom left: running spectral dimension. The lattice (exact diagonalisation) coincides with the infinite-lattice formula (3) until finite-size effects set in near $\tau \sim 10^2$; note the overshoot above $d = 2$. The gasket oscillates around $2 \ln 3 / \ln 5$. Bottom right: $d = 4$ with $E = k^2(1 + \ell^2 k^2)$, $\ell = 10a$, and cutoff $|k| \leq \pi/a$ (green), the closed form (6) (dotted), and $E = k^2$ with the same cutoff (grey).

6. *Connections within the series.* Log-periodic oscillations of d_s on self-similar graphs are the heat-kernel counterpart of the complex dimensions of [19]. The equipartition form of d_s is an instance of the thermodynamic formalism used for entanglement spectra in [20]. Its heat-capacity counterpart, $C = \tau^2 \text{Var}(\lambda) = \frac{1}{2} d_s - \frac{1}{2} \frac{d d_s}{d \ln \tau}$, turns the overshoot of d_s on $\mathbb{Z}^d$ into a Schottky-type anomaly [22]. Whether a computable graph can have an uncomputable spectral dimension, in analogy with the uncomputable fractal dimensions of [21], is left open.

Data and code availability

All numbers and figures are produced by the Python script `quantum_geometry_crossover.py`, available at <https://github.com/Ruqing1963/quantum-geometry-crossover> and archived on Zenodo under DOI [10.5281/zenodo.23195881](https://doi.org/10.5281/zenodo.23195881).




Figure 2: Left: log-periodic oscillation of d_s on the Sierpinski gasket (period $\ln 5$). Middle: norm-resolvent error at $z = i$ on the circle for the lattice Hodge-Dirac operator (convergent, $O(a)$) and the naive symmetric-difference operator (non-convergent, fermion doubling). Right: upwind lower bound on the Connes distance on the square lattice relative to the Euclidean distance, compared with the ℓ^1 distance; the linear test function of Proposition 4.2 gives exactly 1 (grey line).

A Numerical methods

- *Graphs.* The periodic 64×64 square lattice and the level-7 Sierpinski gasket (built recursively on integer coordinates; $N = (3^8 + 3)/2 = 3282$) are diagonalised exactly with LAPACK (`numpy.linalg.eigvalsh`). The lattice eigenvalues agree with the analytic spectrum to 10^{-13} .
- *Heat kernel.* $P(\tau) = \frac{1}{N} \sum_k e^{-\tau \lambda_k}$ on 321 logarithmically spaced points in $[10^{-4}, 10^4]$. d_s is computed both by centred differences in $\ln \tau$ and from (2); the two agree to 1.5×10^{-3} .
- *Modified dispersion.* Radial integrals with cutoff π/a are evaluated by adaptive quadrature (relative tolerance 10^{-11} , no absolute tolerance), split at $k = \tau^{-1/2}$, on $\tau \in [10^{-7}, 10^6]$. Tails with $\tau E > 700$ are omitted, since they underflow.
- *Dirac operators.* Resolvent differences are evaluated mode by mode on the circle, including the bound $1/|k|$ for continuum modes outside the Brillouin zone.
- *Connes distance.* The upwind recursion is checked to satisfy the admissibility constraint at every vertex; values are reported at the lattice point on each ray at the edge of the grid.

References

- [1] J. Ambjørn, J. Jurkiewicz, R. Loll, “Spectral dimension of the universe”, Phys. Rev. Lett. 95 (2005) 171301, arXiv:hep-th/0505113.
- [2] O. Lauscher, M. Reuter, “Fractal spacetime structure in asymptotically safe gravity”, JHEP 10 (2005) 050, arXiv:hep-th/0508202.
- [3] P. Hořava, “Spectral dimension of the universe in quantum gravity at a Lifshitz point”, Phys. Rev. Lett. 102 (2009) 161301, arXiv:0902.3657.
- [4] L. Modesto, “Fractal structure of loop quantum gravity”, Class. Quant. Grav. 26 (2009) 242002, arXiv:0812.2214.
- [5] D. Benedetti, “Fractal properties of quantum spacetime”, Phys. Rev. Lett. 102 (2009) 111303, arXiv:0811.1396.

- [6] G. Calcagni, “Fractal universe and quantum gravity”, *Phys. Rev. Lett.* 104 (2010) 251301, arXiv:0912.3142.
- [7] S. Carlip, “Dimension and dimensional reduction in quantum gravity”, *Class. Quant. Grav.* 34 (2017) 193001, arXiv:1705.05417.
- [8] G. Calcagni, A. Eichhorn, F. Saueressig, “Probing the quantum nature of spacetime by diffusion”, *Phys. Rev. D* 87 (2013) 124028, arXiv:1304.7247.
- [9] A. Connes, *Noncommutative Geometry*, Academic Press, San Diego (1994).
- [10] A. Connes, “Gravity coupled with matter and the foundation of non-commutative geometry”, *Commun. Math. Phys.* 182 (1996) 155, arXiv:hep-th/9603053.
- [11] S. Alexander, R. Orbach, “Density of states on fractals: fractons”, *J. Physique Lett.* 43 (1982) 625, doi:10.1051/jphyslet:019820043017062500.
- [12] R. Rammal, G. Toulouse, “Random walks on fractal structures and percolation clusters”, *J. Physique Lett.* 44 (1983) 13, doi:10.1051/jphyslet:0198300440101300.
- [13] E. Akkermans, G. V. Dunne, A. Teplyaev, “Physical consequences of complex dimensions of fractals”, *EPL* 88 (2009) 40007, arXiv:0903.3681.
- [14] H. B. Nielsen, M. Ninomiya, “Absence of neutrinos on a lattice (I). Proof by homotopy theory”, *Nucl. Phys. B* 185 (1981) 20.
- [15] K. S. Stelle, “Renormalization of higher derivative quantum gravity”, *Phys. Rev. D* 16 (1977) 953.
- [16] J. Ambjørn, S. Jordan, J. Jurkiewicz, R. Loll, “A second-order phase transition in CDT”, *Phys. Rev. Lett.* 107 (2011) 211303, arXiv:1108.3932.
- [17] NIST Digital Library of Mathematical Functions, §10.40, <https://dlmf.nist.gov/10.40>.
- [18] I. S. Gradshteyn, I. M. Ryzhik, *Table of Integrals, Series, and Products*, 7th ed., Academic Press (2007), formula 3.462.1.
- [19] R. Chen, “Holographic entanglement of fractal regions: Complex dimensions, non-local counterterms and cosmic-brane multifractality” (2026), doi:10.5281/zenodo.23189575.
- [20] R. Chen, “Multifractal large deviations in quantum entanglement: Ideal Fermi-gas mapping, Markov tensor cascades, and quartic Casimir symmetry” (2026), doi:10.5281/zenodo.23193512.
- [21] R. Chen, “Uncomputable fractal dimensions, Kolmogorov complexity cascades, and the Chaitin barrier in physical systems” (2026), doi:10.5281/zenodo.23194419.
- [22] R. Chen, “Thermodynamics of scale: Entropic c -functions, data processing, spectral thermodynamics, fluctuation relations, and the arrow of time” (2026), doi:10.5281/zenodo.23199122.
- [23] J. Ambjørn, A. Görlich, S. Jordan, J. Jurkiewicz, R. Loll, “CDT meets Hořava–Lifshitz gravity”, *Phys. Lett. B* 690 (2010) 413–419, arXiv:1002.3298.
- [24] J. Kigami, *Analysis on Fractals*, Cambridge Tracts in Mathematics 143, Cambridge University Press (2001).
- [25] R. S. Strichartz, *Differential Equations on Fractals: A Tutorial*, Princeton University Press (2006).

- [26] M. Fukushima, T. Shima, “On a spectral analysis for the Sierpiński gasket”, *Potential Anal.* 1 (1992) 1–35.
- [27] J. Kigami, M. L. Lapidus, “Weyl’s problem for the spectral distribution of Laplacians on p.c.f. self-similar fractals”, *Commun. Math. Phys.* 158 (1993) 93–125.