

(a) the bottom of $\bigcup_{0 \le t < L} t\mathcal{F}$
between the arcs and $y = 1$: $\Psi_0(c) - e^{-c}$
below the arcs: outside $\bigcup_t t\mathcal{F}$
$y = \sqrt{1 - \{x\}^2}$ (exact bottom)
$y = 1$ (Λ_1, lower bound)
$y = \sqrt{3}/2$ ($\Lambda_{\sqrt{3}/2}$, upper bound)
y
x (width coordinate of the cusp; period 1 of the scallop, period Q of $G|_2\varrho$)
(b) the anatomy of $B \approx 0.48$, p from 101 to 400009 (fits over $p \geqq 10^4$)
$\|G_{1,p}\|^2 - \frac{1}{\pi}(\log p)^2$: $b = +0.4817$
C_1': free part of the main cusps: $b = +0.3829$
C_2: Siegel term and $k \geq p$: $b = +0.0566$
C_4: arcs, main cusps: $b = +0.0407$
arcs on the free part: $b = +0.0288$
C_3 + rest (other cusps, correlations): $b = +0.0015$
proved: $b = 0.38301$ (Lemma 6.10 of Part II)
$b = 2\kappa/\pi = 0.02879$ (Mellin, Section 2)
contribution to $\|G_{1,p}\|^2 - \frac{1}{\pi}(\log p)^2$
$\log p$