SUMS OF TWO SQUARES AND HEEGNER NORMS WITH FEW PRIME FACTORS, IV: THE SCALLOPED PARSEVAL FORMULA, THE HEEGNER CLASSIFICATION, AND EISENSTEIN DISPERSION

RUQING CHEN

ABSTRACT. Parts I–III studied the cusp form $G_{m,m'}$ attached to $\theta^2(mz)\theta^2(m'z)$, whose Fourier coefficients are the remainders of the sieve for $r(n)r(N-n)$. Three results. (1) The Petersson norm of G is given *exactly* by a Parseval sum over the cusps whose lower boundary is the scalloped curve $y=\sqrt{1-\{x\}^2}$ of Waibel's fundamental domain, with correlation terms $a(k)\overline{a(k+jQ)}$ at distances multiple of the width. It gives the coefficient of $\log p$ in $\|G_{1,p}\|^2=\frac{1}{\pi}(\log p)^2+B\log p+O(1)$ in closed form: $B=0.48204$ is the sum of 0.383 from the free part of the two main cusps, $2\kappa/\pi=0.0288$ from the region between the arcs and the line $y=1$, and 0.0702 from the coefficients $k\geq p$, a sum of exponential integrals $E_1(\pi\beta)$, $E_1(4\pi\beta)$ weighted by $r(\beta)^2$; the Siegel main term contributes nothing to B. (2) For the seven odd discriminants of class number one, $D\in\{3,7,11,19,43,67,163\}$, $\Theta_{1,p}^{(D)}=\theta_D(z)\theta_D(pz)$ has the square-free level Dp with four Atkin–Lehner cusps. We prove, for all seven D by one argument, the Siegel main term with closed-form densities at D and p, the Atkin–Lehner signs, the necessary condition $a_\varphi(D)=\chi_{-D}(p)$, $a_\varphi(p)=-\chi_{-D}(p)$ for a newform to occur, with the non-vanishing of $L(1,\varphi)L(1,\varphi\otimes\chi_{-D})$ as the remaining criterion, the Rankin–Selberg formula with the constant $C_D=w_D^2\sqrt{D}/(8\pi^2)$, and the asymptotic $\|G_{1,p}^{(D)}\|^2=\frac{6}{\pi(D+1)}(\log p)^2+B_D\log p+O(1)$ with B_D in closed form; the leading coefficient is $2/\operatorname{vol}(\Gamma_0(D)\backslash\mathbb{H})$, and the corresponding lower bound holds for $D=8$ as well. (3) For the Eisenstein integers the Fourier transform of the circle $x^2+xy+y^2\equiv a$ modulo any c prime to 3 is a classical Kloosterman sum with the hexagonal dual norm, the ramified prime 3 takes over the role that 2 plays for the Gaussian integers, and the obstruction to a level beyond $1/2$ is unchanged. We also correct an error of Parts II–III: the Kloosterman sums of the dispersion form are $S(\overline{16m}N,|h|^2;m'k)$, not $S(16N,m|h|^2;m'k)$.

1. INTRODUCTION AND STATEMENT OF RESULTS

1.1. **Setting.** Parts I–III [3, 4, 5] prove the level of distribution $1/2$ for the weighted sequence $r(n)r(N-n)$, r the number of representations as a sum of two squares, and with it $\Omega(n(N-n))\leq 7$, $\max(\Omega(n),\Omega(N-n))\leq 5$ for every large $N\equiv 2 \pmod 4$, through the cusp forms $G_{m,m'}\in S_2(\Gamma_0(4mm'))$ with $\theta^2(mz)\theta^2(m'z)=E_{m,m'}+G_{m,m'}$: the Fourier coefficient $a_{G_{m,m'}}(N)=T(m,m';N)-M(m,m';N)$ is the remainder of the sieve at the modulus pair (m,m'), and the uniform bound $|a_G(N)|\ll N^{1/2+\varepsilon}\|G\|$ (Part I, Theorem 2.3) with $\|G_{m,m'}\|^2\asymp(\log mm')^2$ (Part II, Theorems 1.5 and 6.14) gives the level. Part III located the obstacles to a level beyond $1/2$ and to the pair $(7,4)$. The present part answers three questions that arose from it: how large exactly is the norm (the term $B\log p$ which Part II, Remark 6.16, could only estimate); what happens for the other imaginary quadratic fields of class number one; and whether the Eisenstein integers, with their square-free level, change the dispersion problem.

1.2. **Results.** Let $G\in S_2(\Gamma_0(L))$. Waibel's formula [14, (22)] (Part I, Lemma 2.8) writes $\|G\|^2$ as a sum over the cusp classes of integrals of $|G|_2\varrho|^2$ over the union of the L translates $t\mathcal{F}$,

Date: November 2, 2026.

$0 \le t < L$, of the standard fundamental domain, whose lower boundary is the *scalloped* curve $y = s(x) = \sqrt{1-\{x\}^2}$ ($\{x\}$ the distance to the nearest integer), not a horizontal line. Part II bounded these integrals by the strips $y \ge 1$ and $y \ge \sqrt{3}/2$: with Λ_h the sum of the integrals over $y \ge h$, $\Lambda_1 \le \|G\|^2 \le \Lambda_{\sqrt{3}/2}$ (Part II, Lemma 6.11). Parseval over the exact region gives an identity. Throughout, $S(Y) = \sum_{k\ge1} r(k)^2 k^{-1} e^{-k/Y}$, $E_1(x) = \int_x^\infty e^{-t}t^{-1}\mathrm{d}t$ is the exponential integral, and $\kappa = \int_{-1/2}^{1/2} \log(1-x^2)^{-1/2}\mathrm{d}x = 0.045231$.

Theorem 1.1 (Scalloped Parseval)**.** *For $G \in S_2(\Gamma_0(L))$ with $G|_{2\varrho} = \sum_{k\ge1} b_d(k)e(kz/Q_d)$ at a cusp of the class $d \mid L$ and width Q_d (Section 2), and $\Psi_j(c) = \int_{-1/2}^{1/2} \cos(2\pi jx)e^{-c\sqrt{1-x^2}}\mathrm{d}x$,*

$$\|G\|^2 = \sum_{d\mid L} w_d \sum_{k\ge1}\sum_j b_d(k)\overline{b_d(k+jQ_d)}\,\frac{\Psi_j(c_{k,j})}{c_{k,j}}, \qquad c_{k,j} = \frac{2\pi(2k+jQ_d)}{Q_d},$$

with explicit weights w_d (Proposition 2.1). For $G = G_{1,p}$ the formula, evaluated with $k \le 14p$, reproduces the Rankin–Selberg estimates of Part II where these have converged (7.42858 against 7.42871 at $p = 101$), lies inside the sandwich $\Lambda_1 \le \|G\|^2 \le \Lambda_{\sqrt{3}/2}$ at every p, and gives, over $p \ge 10^4$,

$$\|G_{1,p}\|^2 = \frac{1}{\pi}(\log p)^2 + B\log p + O(1), \qquad B = 0.48204\ldots,$$

where B is the explicit constant of Theorem 2.4: 0.383 is the free part $\frac{1}{4\pi}(S(p/4\pi) + S(p/\pi))$ of the two main cusps (Part II, Lemma 6.3), 0.0288 $= 2\kappa/\pi$ the arcs acting on that free part (Proposition 2.2), 0.0569 and 0.0119 the coefficients $k \ge p$ of the two main cusps at height 1 and through the arcs, $\frac{1}{\pi}\sum_\beta r(\beta)^2(E_1(\pi\beta) + E_1(4\pi\beta))$ and its arc version, 0.0019 the cusp of denominator 2 and -0.0005 the correlation terms; the Siegel main term and all cross terms are $O(1)$. The fit over $10^4 \le p \le 4\cdot10^5$ gives 0.4817, the local slopes at the largest primes 0.4819 and 0.4820.

The value $B \approx 0.46$ of Part II, Remark 6.16, was a fit to Rankin–Selberg estimates that had not converged; the fit of the exact formula gives 0.48, and Theorem 2.4 the constant itself.

What the identity adds to the sandwich is structural, not only a matter of precision. A horizontal cut-off $y \ge h$ is invariant under $x \mapsto x+1$, so Parseval over it keeps the diagonal alone, $\sum_k |b_d(k)|^2 e^{-4\pi hk/Q_d}/k$; the scalloped boundary has period 1 in x while the expansion at a cusp of width Q_d has period Q_d, so the integral over the union of the Q_d translates couples $b_d(k)$ with $b_d(k+jQ_d)$, and the result is the diagonal with the weights $\Psi_0(c)/c$ in place of e^{-c}/c together with the correlation terms $j \ne 0$ with the weights $\Psi_j(c)/c$. Both are visible in B: the arcs give $b_2 + b_4 = 0.0407$ and part of b_5, the correlations give $b_6 = -0.0005$ (Theorem 2.4), and neither can be seen from $\Lambda_1 \le \|G\|^2 \le \Lambda_{\sqrt{3}/2}$, whose two sides differ by $0.134\log p$ (Part II, Section 6.7).

Let now $D \in \{3,7,11,19,43,67,163\}$ be an odd discriminant of class number one, $\chi = \chi_{-D}$, w_D the number of units, $N_D(x,y) = x^2+xy+\frac{D+1}{4}y^2$, $r_D(n) = w_D\sum_{d\mid n}\chi(d)$, $\theta_D = \sum r_D(n)e(nz)$ and, for a prime $p \ne D$,

$$\Theta_{1,p}^{(D)} = \theta_D(z)\theta_D(pz) = E + G_{1,p}^{(D)} \in M_2(\Gamma_0(Dp)),$$
$$a_G(k) = T(k) - M(k) = \sum_{\alpha+p\beta=k} r_D(\alpha)r_D(\beta) - M(k).$$

Theorem 1.2 (The odd Heegner discriminants). (i) $M(k) = \frac{4\pi^2 k}{Dp}\prod_\ell \beta_\ell(k)$ *with the standard densities at* $\ell \nmid Dp$ *and, with* $v = v_\ell(k)$ *and* $\varepsilon = -\chi_{-D}(p)$,

$$\beta_p(k) = (1+\chi(p)) - \chi(p)\Big(1+\frac{1}{p}\Big)p^{-v}, \qquad \beta_D(k) = (1+\varepsilon) - \varepsilon\Big(1+\frac{1}{D}\Big)D^{-v}.$$

(ii) $\Theta|_2 W_p = \chi(p)\Theta$, $\Theta|_2 W_D = -\chi(p)\Theta$, $\Theta|_2 W_{Dp} = -\Theta$.

(iii) *A newform* φ *of level* Dp *occurs in* $G_{1,p}^{(D)}$ *only if* $a_\varphi(D) = \chi_{-D}(p)$ *and* $a_\varphi(p) = -\chi_{-D}(p)$ *(root number* $+1$*); an oldform from level* p *only if* $a_f(p) = -\chi_{-D}(p)$*; an oldform from level* D *only if* $\chi_{-D}(p) = +1$*. These conditions are necessary; a newform* φ *that satisfies them occurs if and only if* $L(1,\varphi)L(1,\varphi\otimes\chi_{-D}) \neq 0$ *(Proposition 3.4). Unlike* $D = 4$*, where* $\eta_4 = -1$ *for every newform of level* $4n$*, neither sign of* W_D *is excluded.*

(iv) $\langle \Theta_{1,p}^{(D)}, \varphi\rangle = C_D\,\chi(p)\,\eta_p(\varphi)\,L(1,\varphi)L(1,\varphi\otimes\chi_{-D})$ *with* $C_D = \dfrac{w_D}{4\pi L(1,\chi_{-D})} = \dfrac{w_D^2\sqrt{D}}{8\pi^2}$.

(v) $\|G_{1,p}^{(D)}\|^2 = \dfrac{6}{\pi(D+1)}(\log p)^2 + B_D\log p + O(1)$ *with* B_D *in closed form (Theorem 3.8):* $B_3 = 0.8211$, $B_7 = 0.3289$, $B_{11} = 0.2941$, $B_{19} = 0.3225$, $B_{43} = 0.3637$, $B_{67} = 0.3632$, $B_{163} = 0.3216$. *For all nine discriminants, including* $D = 4, 8$*, the two wide Atkin–Lehner cusps give, with* $\psi(D) = D\prod_{\ell\mid D}(1+1/\ell)$,

$$\|G_{1,p}^{(D)}\|^2 \;\geq\; \frac{6}{\pi\,\psi(D)}(\log p)^2 + O(\log p) = \frac{2}{\mathrm{vol}(\Gamma_0(D)\backslash\mathbb{H})}(\log p)^2 + O(\log p),$$

with equality up to $O(\log p)$ *for* $D = 4$ *(Part II, Theorem 6.14) and for the odd* D*; for* $D = 8$ *only the lower bound is proved here.*

The constants are those of the analytic class number formula with $h = 1$, $w_D^2 L(1,\chi_{-D})^2 = 4\pi^2/D$, which makes them independent of w_D; the remaining factor $\prod_{\ell\mid D}(1+1/\ell)^{-1}$ is the Euler factor at the ramified primes, where $\sigma_\chi(\ell^j) = 1$. Section 4 verifies (i)–(iv) for $D \leq 43$ (91 newform components, Rankin–Selberg ratios in $[0.9936, 1.0048]$) and (v) for all seven odd D by the exact Parseval formula: the local slopes of $\|G\|^2 - \frac{6}{\pi(D+1)}(\log p)^2$ at the largest primes computed are 0.8210, 0.3289, 0.2943, 0.3214, 0.3661, 0.3648, 0.3128 against $B_D = 0.8211$, 0.3289, 0.2941, 0.3225, 0.3637, 0.3632, 0.3216.

Theorem 1.3 (Eisenstein dispersion). *For* $(c,3) = 1$*,* c *odd or even,* $(a,c) = 1$ *and* $h \in \mathbb{Z}^2$*, with* $|h|_3^2 = h_1^2 - h_1h_2 + h_2^2$ *the hexagonal dual norm,*

$$\sum_{\substack{x,y \bmod c\\ x^2+xy+y^2\equiv a\ (c)}} e\Big(\frac{h_1x+h_2y}{c}\Big) = \sum_{g\mid(c,h)} g\,\chi_{-3}\Big(\Big(\frac{c}{g}\Big)^*\Big)S\Big(a,\ \overline{3}\Big|\frac{h}{g}\Big|_3^2;\ \frac{c}{g}\Big).$$

The remainder $a_{G^{(3)}_{m,m'}}(N) - a_{G^{(3)}_{m,3m'}}(N)$ *of the sieve for* $r_3(n)r_3(N-n)$ *has as its leading term a trilinear form in the sums* $S(\overline{27m}N, |h|_3^2; m'k)$ *over all moduli* $m'k$ *prime to* 3*, with a dual condition* $3 \mid |h|_3^2$ *in place of* $2 \mid |h|^2$*; its bilinear form in* (m',k) *and the Cauchy–Schwarz and Poisson sizes are those of Part III, and the obstruction to a level beyond* $1/2$ *is the same.*

1.3. **A correction to Parts II–III.** Throughout, a bar denotes the inverse modulo the modulus of the Kloosterman sum in which it occurs: $\bar m = m^{-1}$, $\overline{16} = 16^{-1}$, $\overline{16m} = (16m)^{-1} \bmod c$. The dispersion form of Part III, Proposition 3.2, arises from Poisson summation on the circle $x^2 + y^2 \equiv N\bar m \pmod c$, $c = m'k$, whose Fourier transform at h is, by completing the square (Part III, Lemma 3.1, c odd), the Kloosterman sum $S(N\bar m, \overline{16}|h|^2; c)$ up to the factor $\chi_{-4}(c)c$: the first argument is the residue class of the circle, the second comes from $\frac14|h|^2$ with the 4 of $|h|^2 = (h_1^2+h_2^2)$ and the 4 of $x^2+y^2 = \frac12 x^{\mathsf T}(2I)x$. Part III, Section 3.2 (and Part II,

Section 8.1) then rewrote $S(N\bar{m},\overline{16}|h|^2;c)$ as $S(16N,m|h|^2;c)$. This is false: for $(a,c)=1$ the substitution $t\mapsto \bar{a}t$ in $S(a,b;c)=\sum^*_{t \bmod c} e((at+b\bar{t})/c)$ gives $S(a,b;c)=S(1,ab;c)$, so a Kloosterman sum depends on its two arguments only through their product modulo the modulus, and the two products, $\overline{16m}N|h|^2$ and $16mN|h|^2$, differ by the factor $16m\cdot 16m=(16m)^2$. The correct form is $S(\overline{16m}N,|h|^2;c)$, the inverse of $16m$ taken modulo the modulus, and it cannot be removed. Explicitly, the products of the two arguments modulo c are $N\bar{m}\cdot\overline{16}|h|^2=\overline{16m}N|h|^2$ for the original sum and for its correct form, but $16N\cdot m|h|^2=16mN|h|^2$ for the rewritten one, and $\overline{16m}\not\equiv 16m$ in general (the script `kloosterman_argument_audit.py` checks the equality and the inequality numerically for random N,m,h and odd moduli c, together with the corresponding identities of Section 5 for c odd and even). The Weil, Cauchy–Schwarz and Poisson computations of Part III, which only open the sums, are unaffected; the Kuznetsov heuristic of its Section 3.5(b), which needs fixed integer arguments, is withdrawn (the integer representatives $(N+tc)/(16m)$ of the first argument vary with c); the correlation experiment of its Section 4.4 has been rerun with the correct sums, with the same conclusion. The corrected versions of Parts II and III are the ones cited here. Table 1 lists what depends on the identity and what does not.

Statement of Parts II–III	Uses the identity?	Status
Part II, Sections 1–7 and Part III, Sections 5–8: uniform bound $\lvert a_G(N)\rvert \ll N^{1/2+\varepsilon}\lVert G\rVert$, mean square, norm, level 1/2, $(7,5)$, dimension of the sieve, short intervals, spectral decomposition	no (theta-series side)	unchanged
Part II, Proposition 8.2 and Part III, Proposition 3.2: the dispersion form and the trilinear form (5) of Part III	yes, in the first argument	corrected to $S(\overline{16m}N,\lvert h\rvert^2;m'k)$
Part III, Section 3.3: Weil bound (level 1/4), theta bound (level 1/2), cancellation over m' (level 2/3)	opens the sums only	unchanged
Part III, Section 3.4(a)–(c): the three obstacles	yes	sharpened: the fixed argument carries a unit that depends on the modulus
Part III, Section 3.5(a): substitution of Drappeau's bound [8]	formally; N enters only as N_0	unchanged (both assignments stop at 1/2)
Part III, Section 3.5(b): Kuznetsov route [6] with the arguments $16N$, mj	yes	withdrawn
Part III, Sections 4.1–4.3: Cauchy–Schwarz (A1), (A2), Poisson in h	the unit sits in λ, invisible to (λ,q) and $\lvert G(\lambda;q)\rvert$	unchanged
Part III, Section 4.4: correlation experiment	yes (the sums computed)	rerun; same conclusion
Part III, Section 4.5: bilinear form (9) in (m',k)	yes	corrected arguments $\overline{16m}N\bar{k}$, $\overline{16m}N\overline{m'}$
Part II, Section 8.2: signs of $a_{G_{d,e}}(N)$	no ($a_G(N)$ computed directly)	unchanged

TABLE 1. The dependence of Parts II–III on the identity $S(N\bar{m},\overline{16}|h|^2;c)=S(\overline{16m}N,|h|^2;c)$ and its incorrect form $S(16N,m|h|^2;c)$.

2. The scalloped Parseval formula and the anatomy of B

Throughout, $G\in S_2(\Gamma_0(L))$, and for a class of pairs (a,c) modulo L with $(c,L)=d$ let $\varrho=\varrho_{a,c}\in \mathrm{SL}_2(\mathbb{Z})$ have first column (a,c), $\hat{d}\in\{d,d/2\}$ as in Part I, Lemma 2.9, $Q=L/\hat{d}$, and

$G|_2\varrho = \sum_{k\ge1} b(k)e(kz/Q)$. Part I, Lemma 2.8 is

$$\|G\|^2 = \frac{\mu}{\tilde\mu}\sum_{c\ (L)}\sum_{\substack{a\ (L)\\(a,c,L)=1}} I(\varrho_{a,c}), \qquad I(\varrho)=\int_{\bigcup_{0\le t<L} t\mathcal F}|G|_2\varrho|^2\,\mathrm dx\,\mathrm dy, \tag{1}$$

and $\bigcup_{0\le t<L} t\mathcal F = \{-\frac12 \le x \le L-\frac12,\ y\ge s(x)\}$ with $s(x)=\sqrt{1-\{x\}^2}$ of period 1.

Proposition 2.1. *For $j\in\mathbb Z$ and $c>0$ let $\Psi_j(c)=\int_{-1/2}^{1/2} e(-jx)\,e^{-c\sqrt{1-x^2}}\,\mathrm dx$ (real, $\Psi_{-j}=\Psi_j$). Then*

$$I(\varrho) = L\sum_{k\ge1}\sum_{j\ge(1-k)/Q} b(k)\overline{b(k+jQ)}\,\frac{\Psi_j(c_{k,j})}{c_{k,j}}, \qquad c_{k,j}=\frac{2\pi(2k+jQ)}{Q},$$

(the range of j is where $k+jQ\ge1$; a cusp form has $b(k')=0$ for $k'\le0$), and therefore $\|G\|^2=\sum_{d\mid L} w_d\sum_{k,j} b_d(k)\overline{b_d(k+jQ_d)}\Psi_j(c_{k,j})/c_{k,j}$ with $w_d=Q_d$ if $\hat d=d$ and $w_d=Q_d/4$ if $\hat d=d/2$. The terms $j=0$ with $\Psi_0(c)$ replaced by e^{-c} are the height-1 sums $\Lambda_1(d)$ of Part II; $\Psi_0(c)-e^{-c}>0$ integrates over the region between the arcs and the line $y=1$; the terms $j\ne0$ are correlations of coefficients at distance a multiple of the width, which a flat bottom annihilates. Since $c_{k,j}\ge 2\pi j$ and $\Psi_j(c)\le e^{-c\sqrt3/2}$, only $j\le7$ contribute above 10^{-16}.

Proof. $|G|_2\varrho|^2=\sum_{k,k'} b(k)\overline{b(k')}e((k-k')x/Q)e^{-2\pi(k+k')y/Q}$. Integrating in y from $s(x)$ to ∞ gives $e^{-c\,s(x)}/c$ with $c=2\pi(k+k')/Q$, and since s has period 1 and the x-range consists of L consecutive unit periods,

$$\int_{-1/2}^{L-1/2} e\Big(\frac{(k-k')x}{Q}\Big)e^{-c\,s(x)}\mathrm dx = \Big(\sum_{t=0}^{L-1} e\Big(\frac{(k-k')t}{Q}\Big)\Big)\int_{-1/2}^{1/2} e\Big(\frac{(k-k')x}{Q}\Big)e^{-c\,s(x)}\mathrm dx,$$

where the geometric sum is L if $Q\mid k-k'$ and 0 otherwise. With $k'=k+jQ$ the inner integral is $\Psi_j(c)$. The count of pairs in a class and $\mu/\tilde\mu = 1/(L\varphi(L))$ are as in Part II, Lemmas 6.2 and 6.12. □

For $G=G_{1,p}$, $L=4p$, the six classes are $d\in\{1,4,2\}$ (widths $4p,p,4p$; $b=a_{G_{1,p}}/Q$ at the Atkin–Lehner cusps, $b=\frac1p(T^{\mathrm{odd}}-M^{\mathrm{odd}})$ at the cusp of denominator 2) and $d\in\{p,4p,2p\}$ (widths $4,1,4$). The script `subleading_log_anatomy.py` evaluates the formula with $k\le14p$ and $j\le8$, Ψ_j by Gauss–Legendre quadrature on a logarithmic grid in c with cubic splines.

Proposition 2.2 (The arc term)**.** *At a cusp of width Q put $Y=Q/4\pi$ and $\kappa=\int_{-1/2}^{1/2}\log(1-x^2)^{-1/2}\mathrm dx = 0.045231$. Then*

$$\sum_{k\ge1}\frac{r(k)^2}{k}\Psi_0(k/Y) = 2(\log Y)^2 + (c_1+4\kappa)\log Y + O(1),$$

with $c_1=9.758$ the constant of Part II, Lemma 6.3; so the arcs add $4\kappa\log Y$ to the free part at each of the two main cusps, i.e. $\frac{2\kappa}{\pi}\log p = 0.028795\log p$ in total.

Proof. For $\operatorname{Re} w>0$, $\int_0^\infty \Psi_0(c)c^{w-1}\mathrm dc = \Gamma(w)\int_{-1/2}^{1/2}(1-x^2)^{-w/2}\mathrm dx = \Gamma(w)\,\Phi(w)$ with Φ entire, bounded on vertical strips, $\Phi(w)=1+\kappa w+O(w^2)$. With $D(1+w)=\sum_k r(k)^2k^{-1-w}=4w^{-2}(1+a_1w+a_2w^2+\dots)$ near $w=0$ (Part II, Lemma 6.3; $a_1=2\gamma+2\frac{L'}{L}(1,\chi_{-4})-2\frac{\zeta'}{\zeta}(2)+\frac13\log2$), Mellin inversion gives

$$\sum_{k\ge1}\frac{r(k)^2}{k}\Psi_0(k/Y) = \frac{1}{2\pi i}\int_{(2)}\Gamma(w)\Phi(w)D(1+w)Y^w\,\mathrm dw.$$

The integrand is holomorphic on $\operatorname{Re} w \geq -\frac{1}{2}$ except for the triple pole at $w=0$ ($L(1+w,\chi_{-4})^2$ is entire, $1/\zeta(2+2w)$ is holomorphic and of polynomial growth there, $\Gamma(w)$ decays exponentially), so moving the line to $\operatorname{Re} w = -\frac{1}{2}$ leaves the residue plus $O(Y^{-1/2})$. With $\Gamma(w) = w^{-1} - \gamma + O(w)$ the integrand is $4w^{-3}\big(1+(a_1+\kappa-\gamma)w+O(w^2)\big)\sum_{n\geq 0}(\log Y)^n w^n/n!$, whose residue is $2(\log Y)^2 + 4(a_1+\kappa-\gamma)\log Y + O(1)$: the double pole of $D(1+w)$ against the simple pole of $\Gamma(w)$ gives $(\log Y)^2$, and the first-order terms a_1, κ, $-\gamma$ of the three factors give the coefficient of $\log Y$. Without κ this is Part II, Lemma 6.3 with $c_1 = 4(a_1-\gamma)$; the arcs add $4\kappa\log Y$. The antiderivative of $\log(1-x^2)$ is $x\log(1-x^2) - 2x + \log\frac{1+x}{1-x}$, whence the value of κ. (The first-order approximation $\Psi_0(c) - e^{-c} \approx c\kappa_1 e^{-c}$, $\kappa_1 = \int(1-\sqrt{1-x^2})\mathrm{d}x = 0.04339$, gives $4\kappa_1$ instead: the higher orders in c also contribute to the $\log Y$ term.) □

2.1. **The anatomy of B.** Write $\|G_{1,p}\|^2 - \frac{1}{\pi}(\log p)^2 = C_1' + C_2 + C_3 + C_4$ with $C_1 = \frac{1}{4\pi}(S(p/4\pi) + S(p/\pi))$ the free part of the two main cusps, $C_1' = C_1 - \frac{1}{\pi}(\log p)^2 = 0.38301\log p + O(1)$, $C_2 = \Lambda_1(\text{main cusps}) - C_1$ (the correction $r(k)^2 \to |a(k)|^2$: the Siegel main term for $k<p$, the terms with $\beta \geq 1$ for $k \geq p$), $C_3 = \Lambda_1$ of the other four cusps, and $C_4 = \|G\|^2 - \Lambda_1$ (the part below height 1). Table 2 and Figure 1 give the result of the exact formula for twelve primes from 101 to 400009 (a few seconds per prime).

p	C_1'	C_2	C_3	C_4 arc, main	free part	C_4 corr.	total, $/\log p$
101	1.128	−0.815	0.154	0.164	0.151	−0.001	0.649, 0.141
1009	1.999	−0.711	0.162	0.259	0.218	−0.002	1.727, 0.250
10007	2.876	−0.582	0.166	0.352	0.284	−0.003	2.830, 0.307
100003	3.758	−0.453	0.169	0.446	0.350	−0.004	3.939, 0.342
400009	4.289	−0.374	0.171	0.502	0.390	−0.005	4.607, 0.357
fit b ($p \geq 10^4$)	0.3829	0.0566	0.0020	0.0407	0.0288	−0.0005	0.4817
local slope, $p \geq 10^5$	0.3830	0.0568	0.0019	0.0407	0.0288	−0.0005	0.4819
proved (Theorem 2.4)	0.3830	0.0569	0.0019	0.0407	0.0288	−0.0005	0.4820

TABLE 2. The parts of $\|G_{1,p}\|^2 - \frac{1}{\pi}(\log p)^2$; C_3 includes the arc terms of the four narrow cusps. The fit of the total is the three-parameter fit of the exact norm, with $a = 0.31845$ for $(\log p)^2$ ($1/\pi = 0.31831$) and $c = -1.607$; the local slopes are $(y(p_2) - y(p_1))/\log(p_2/p_1)$ over the last three primes. The part below height 1 is $0.041\log p$, against the bound $0.134\log p$ of Part II, Section 6.7. Scripts `subleading_log_anatomy.py`, `closed_form_B.py`; floating point.

So $B \approx 0.48$: 0.383 and 0.029 are proved by Mellin inversion, and the remaining 0.07 is the Siegel and tail corrections at height 1 (0.057) and through the arcs (0.012), plus the other cusps (0.002) and the correlations (−0.0005). These are not Mellin transforms of Dirichlet series, and the next subsection evaluates them by a different mechanism.

2.2. **The closed form of B.** For $k \geq p$ the coefficient is $a(k) = \sum_{\beta\geq 0} r(\beta)r(k-p\beta) - M(k)$ with $r(0)=1$: the term $\beta=0$ is the free part $r(k)$, and $T'(k) = \sum_{\beta\geq 1} r(\beta)r(k-p\beta)$ is the tail. In $|a(k)|^2$ the squares $r(k-p\beta)^2$, summed over k, produce a logarithm by $\sum_{a\leq X} r(a)^2 \sim 4X\log X$; every other product is a shifted convolution of r, or involves the Siegel main term, and is bounded by the following elementary estimates.

Lemma 2.3. *Uniformly for $X \geq 1$ and integers $h \geq 0$:*

(i) *for $h \geq 1$, $\sum_{a\leq X} r(a)r(a+h) \ll (X+h)\,\sigma_{-1}(h) + \sqrt{X+h}\,d(h)$;*
(ii) *$\sum_{a\leq X} r(a)\sigma_{-1}(a+h) \ll X+h$ and $\sum_{a\leq X}\sigma_{-1}(a)\sigma_{-1}(a+h) \ll X+h$;*




FIGURE 1. (a) The lower boundary of $\bigcup_{0\le t<L} t\mathcal{F}$ in the width coordinate of a cusp: the arcs $y=\sqrt{1-\{x\}^2}$, the line $y=1$ of the lower bound Λ_1 and the line $y=\sqrt{3}/2$ of the upper bound $\Lambda_{\sqrt{3}/2}$ of Part II; the shaded region between the arcs and $y=1$ is what $\Psi_0(c)-e^{-c}$ integrates. (b) The parts of $\|G_{1,p}\|^2-\frac{1}{\pi}(\log p)^2$ against $\log p$ for the twelve primes of Table 2, with their least-squares lines over $p\ge10^4$; the proved slopes of Theorem 2.4 are listed. Script `plot_scalloped_anatomy.py`; floating point.

(iii) $\sum_{a\le X} r(a)^2 = 4X\log X + O(X)$ *and* $\sum_{a\le X,\ a\equiv 2\,(4)} r(a)^2 = X\log X + O(X)$.

Proof. (i) The sum counts $(x,y,z,w)\in\mathbb{Z}^4$ with $x^2+y^2\le X$ and $z^2+w^2-x^2-y^2=h$. The substitution $u_1=z-x$, $v_1=z+x$, $u_2=w-y$, $v_2=w+y$ is injective, gives $u_1v_1+u_2v_2=h$ and keeps $|u_i|,|v_i|\le U:=2\sqrt{X+h}$. If $u_1=0$, then $u_2v_2=h$ has at most $2d(h)$ solutions and v_1 at most $2U+1$ values; symmetrically for $u_2=0$; together $\ll Ud(h)$. If $u_1u_2\ne0$ and $g=(u_1,u_2)$, there is no solution unless $g\mid h$, and then (v_1,v_2) runs over a coset of $\mathbb{Z}\,(u_2/g,-u_1/g)$, which has at most $1+2Ug/\max(|u_1|,|u_2|)$ points in the box. Summing over (u_1,u_2), the 1 gives $\ll U^2$, and with $u_i=gu_i'$, $(u_1',u_2')=1$, the second term gives $2U\sum_{g|h}\sum_{|u_i'|\le U/g} 1/\max(|u_1'|,|u_2'|)\ll U\sum_{g|h}U/g=U^2\sigma_{-1}(h)$, as at most $8m$ pairs have $\max(|u_1'|,|u_2'|)=m$.

(ii) Write $\sigma_{-1}(a+h)=\sum_{e|a+h}1/e$ and split at $E=\log^2(2X+2h)$. The divisors $e>E$ contribute at most $E^{-1}\sum_{a\le X} r(a)d(a+h)\le E^{-1}\big(\sum_{a\le X}r(a)^2\big)^{1/2}\big(\sum_{n\le X+h}d(n)^2\big)^{1/2}\ll E^{-1}(X+h)\log^2(2X+2h)\ll X+h$. For $e\le E$, $\sum_{a\le X,\ e|a+h} r(a)$ counts the lattice points of the disc $x^2+y^2\le X$ in the $N_2(-h;e)$ residue classes modulo e with $x^2+y^2\equiv -h$, so it is $N_2(-h;e)\big(\pi X/e^2+O(\sqrt{X}/e+1)\big)$ with $N_2(t;e)\le e\,d(e)^2$ (Part II, proof of Lemma 6.4), and the sum over $e\le E$ is $\ll X\sum_e d(e)^2e^{-2}+\sqrt{X}\sum_{e\le E}d(e)^2e^{-1}+\sum_{e\le E}d(e)^2\ll X+h$. The second bound is the same argument with $r(a)$ replaced by $\sigma_{-1}(a)=\sum_{f|a}1/f$: for $e,f\le E$ the number of $a\le X$ with $f\mid a$, $e\mid a+h$ is at most $X(e,f)/(ef)+1$ and $\sum_{e,f}(e,f)(ef)^{-2}$ converges, while $e>E$ or $f>E$ contribute $\ll E^{-1}\sum_{a\le X}\big(d(a)\sigma_{-1}(a+h)+\sigma_{-1}(a)d(a+h)\big)\ll E^{-1}(X+h)\log^2(2X+2h)$ by Cauchy–Schwarz.

(iii) The first statement is Part II, Lemma 6.3. For the second, $r(2m)=r(m)$ gives $\sum_{a\le X,\ a\text{ odd}} r(a)^2=R(X)-R(X/2)$ with $R(X)=\sum_{a\le X}r(a)^2$, and $a\equiv2\pmod4$ means $a=2m$ with m odd, so the sum is $R(X/2)-R(X/4)=X\log X+O(X)$. □

Theorem 2.4 (The closed form of B)**.** *Let* $E_1(x)=\int_x^\infty e^{-t}\,\mathrm{d}t/t$, $s=\sqrt{1-x^2}$, $I_j(v_0;\lambda)=\int_{v_0}^\infty \Psi_j(\lambda v)\,\mathrm{d}v/(\lambda v)$, *and* $r(0)=1$. *Then*

$$\|G_{1,p}\|^2=\frac{1}{\pi}(\log p)^2+B\log p+O(1),\qquad B=b_1+b_2+b_3+b_4+b_5+b_6=0.48204\ldots,$$

$$b_1=\frac{c_1-2\log(4\pi^2)}{2\pi}=0.383005,\qquad b_2=\frac{2\kappa}{\pi}=0.028794,$$

$$b_3=\frac{1}{\pi}\sum_{\beta\ge1}r(\beta)^2\big(E_1(\pi\beta)+E_1(4\pi\beta)\big)=0.056874,$$

$$b_4=\frac{1}{\pi}\sum_{\beta\ge1}r(\beta)^2\int_{-1/2}^{1/2}\big(E_1(\pi\beta s)+E_1(4\pi\beta s)-E_1(\pi\beta)-E_1(4\pi\beta)\big)\mathrm{d}x=0.011936,$$

$$b_5=\frac{1}{\pi}\sum_{\beta\equiv2\,(4)}r(\beta)^2\int_{-1/2}^{1/2}E_1(\pi\beta s)\,\mathrm{d}x=0.001893,$$

$$b_6=\sum_{j\ge1}\Big[4\sum_{\beta\ge0}r(\beta)r(\beta+j)\,I_j(2\beta+j;2\pi)+2\sum_{\beta\ge0}r(\beta)r(\beta+4j)\,I_j(\beta+2j;\pi)$$
$$+2\sum_{\beta\equiv2\,(4)}r(\beta)r(\beta+4j)\,I_j(\beta+2j;\pi)\Big]=-0.000460.$$

Here b_1 *is the free part of the two main cusps above height* 1 *and* b_2 *the arcs on it (Section 2);* b_3 *and* b_4 *are the coefficients* $k\ge p$ *of the two main cusps above height* 1 *and through the arcs;* b_5 *is the cusp of denominator* 2 *(width* $4p$*) with its arcs;* b_6 *the correlation terms. The Siegel main term, all cross terms and the three narrow cusps contribute* $O(1)$*. All series converge absolutely:* $r(\beta)\le 4d(\beta)\le 4\beta$, $E_1(x)\le e^{-x}$ *for* $x\ge1$, $|\Psi_j(c)|\le e^{-\sqrt{3}c/2}$ *and hence* $|I_j(v_0;\lambda)|\le e^{-\sqrt{3}\lambda v_0/2}$ *for* $\lambda v_0\ge 2$*, so that the terms of* b_3,b_4,b_5 *are* $\ll\beta^2e^{-\sqrt{3}\pi\beta/2}$ *and those of* b_6 *are* $\ll(\beta+j)^2e^{-\sqrt{3}\pi(2\beta+j)}$ *in the first sum* ($\lambda v_0=2\pi(2\beta+j)$) *and* $\ll(\beta+4j)^2e^{-\sqrt{3}\pi(\beta+2j)/2}$ *in the second and third* ($\lambda v_0=\pi(\beta+2j)$)*; the interchanges of sums and integrals in the proof are justified by this domination. The numbers are evaluations of these expressions (*`closed_form_B.py`*); the fits and local slopes quoted in Theorem 1.1 and Section 4 check them and play no part in the proof.*

Proof. We go through the six classes of Proposition 2.1; $O(1)$ means bounded by a constant independent of p, and $M(k)\ll k\sigma_{-1}(k)/p$ uniformly in k and p by Part II, Lemma 6.4 (the same for M^{odd}).

The dominating estimate. Every weight that occurs below has the form $F(t)=\phi(t/Y)/t$, up to a constant factor and a shift of t by $jQ/2\ge0$, with $Y=Q/4\pi\asymp p$ and $\phi\in\{e^{-u},\ \Psi_0(u)-e^{-u},\ \Psi_j(u)\}$; all of these satisfy $|\phi(u)|,|\phi'(u)|\le 2e^{-\sqrt{3}u/2}$ (for Ψ_j: $|\Psi_j'(u)|\le\int\sqrt{1-x^2}\,e^{-u\sqrt{1-x^2}}\mathrm{d}x\le e^{-\sqrt{3}u/2}$, as $\sqrt{1-x^2}\ge\sqrt{3}/2$ on $[-\frac12,\frac12]$), so that $|F'(t)|\le 2e^{-\sqrt{3}t/(2Y)}(1/(Yt)+1/t^2)$. Let $h\ge1$, let $c(a)$ be a sequence with $|c(0)|\ll d(h)$ and $\sum_{1\le a\le X}|c(a)|\ll(X+h)\sigma_{-1}(h)+\sqrt{X+h}\,d(h)$ for all $X\ge1$, and let $h'\ge\max(h,1)$. Then partial summation gives

$$\sum_{a\ge0}c(a)F(a+h')\ll\frac{d(h)}{h'}e^{-\sqrt{3}h'/(2Y)}+\int_{h'}^\infty\big(t\,\sigma_{-1}(h)+\sqrt{t}\,d(h)\big)e^{-\sqrt{3}t/(2Y)}\Big(\frac{1}{Yt}+\frac{1}{t^2}\Big)\mathrm{d}t$$
$$\ll d(h)\Big(1+\frac{Y}{h'}\Big)e^{-\sqrt{3}h'/(2Y)}\tag{2}$$

with an absolute implied constant: with $\lambda=\sqrt{3}/(2Y)$, $\int_{h'}^{\infty}e^{-\lambda t}(1/Y+1/t)\mathrm{d}t=e^{-\lambda h'}/(\lambda Y)+E_1(\lambda h')\le e^{-\lambda h'}\big(2/\sqrt{3}+\log(1+1/(\lambda h'))\big)$ by $E_1(x)\le e^{-x}\log(1+1/x)$, $\int_{h'}^{\infty}e^{-\lambda t}(t^{-1/2}/Y+t^{-3/2})\mathrm{d}t\le h'^{-1/2}e^{-\lambda h'}\big(1/(\lambda Y)+1/(\lambda h')\big)$, and $\sigma_{-1}(h)\le d(h)$. By Lemma 2.3(i) and (ii) the hypothesis holds for $c(a)=r(a)r(a+h)$ and for $c(a)=r(a)\sigma_{-1}(a+h)$. All $O(1)$ terms below are sums over $\beta,\beta'\ge1$ (and $j\ge1$) of bounds (2) with $h'=p\max(\beta,\beta')$ and $Y\le p/\pi$, so that $Y/h'\le1/\pi$ and $h'/Y\ge\pi\max(\beta,\beta')$; and the resulting series converge uniformly in p, because $d(pn)\le2d(n)\le2n$ for a prime p and $n\ge1$, together with $r(n)\le4d(n)\le4n$, gives

$$\sum_{\beta\ne\beta'}r(\beta)r(\beta')\,d(p|\beta-\beta'|)\,e^{-\sqrt{3}\pi\max(\beta,\beta')/2}\le32\sum_{\beta,\beta'\ge1}\beta\beta'\max(\beta,\beta')\,e^{-\sqrt{3}\pi\max(\beta,\beta')/2}<\infty,$$

and likewise on the diagonal $\beta=\beta'$ with $d(h)$ replaced by 1. This is the uniformity in p; the same argument with $e^{-\sqrt{3}\pi\max(\beta,\beta')/2}$ replaced by $e^{-c\max(\beta,\beta')}$ for a fixed $c>0$ is what Section 3.6 uses.

Main cusps, height 1. At the width $Q\in\{p,4p\}$ the class is $\frac{1}{4\pi}\sum_k|a(k)|^2e^{-k/Y}/k$, $Y=Q/4\pi$, so $p\beta/Y=4\pi p\beta/Q\in\{4\pi\beta,\pi\beta\}\ge\pi\beta$. With $a=r+T'-M$, $|a|^2-r^2=2rT'+T'^2-2rM-2T'M+M^2$. By partial summation and Lemma 2.3(ii) with $h=0$, $\sum_k r(k)M(k)e^{-k/Y}/k\ll p^{-1}\sum_k r(k)\sigma_{-1}(k)e^{-k/Y}\ll Y/p\ll1$ and $\sum_k M(k)^2e^{-k/Y}/k\ll p^{-2}\sum_k k\sigma_{-1}(k)^2e^{-k/Y}\ll Y^2/p^2\ll1$. Next, $\sum_k r(k)T'(k)e^{-k/Y}/k=\sum_{\beta\ge1}r(\beta)\sum_{a\ge0}r(a)r(a+p\beta)F(a+p\beta)$ with $F(t)=e^{-t/Y}/t$, and (2) with $h=h'=p\beta$ bounds the inner sum by $d(p\beta)e^{-\sqrt{3}\pi\beta/2}$, which sums to $O(1)$. Similarly $\sum_k T'(k)M(k)e^{-k/Y}/k\ll p^{-1}\sum_\beta r(\beta)\sum_a r(a)\sigma_{-1}(a+p\beta)e^{-(a+p\beta)/Y}\ll p^{-1}\sum_\beta r(\beta)(Y+p\beta)e^{-\pi\beta}\ll1$ by (ii) and partial summation. In $T'(k)^2=\sum_{\beta,\beta'\ge1}r(\beta)r(\beta')r(k-p\beta)r(k-p\beta')$ the terms $\beta<\beta'$ are, with $k=a+p\beta'$, the sums $\sum_{a\ge0}r(a)r(a+p(\beta'-\beta))F(a+p\beta')$, and (2) with $h=p(\beta'-\beta)$, $h'=p\beta'$ bounds them by $d(p(\beta'-\beta))e^{-\sqrt{3}\pi\beta'/2}$: again $O(1)$ in total. The diagonal $\beta=\beta'$ is

$$D_Q=\sum_{\beta\ge1}r(\beta)^2\sum_{a\ge0}r(a)^2\,\frac{e^{-(a+p\beta)/Y}}{a+p\beta},$$

and by (iii) and partial summation, $\sum_{a\ge1}r(a)^2F(a+h)=\int_1^\infty(4X\log X+O(X))(-F'(X+h))\,\mathrm{d}X$ for $F(t)=e^{-t/Y}/t$, where $-F'(t)=e^{-t/Y}(1/(Yt)+1/t^2)>0$; the $O(X)$ part is $\ll e^{-h/Y}+E_1(h/Y)\ll e^{-h/2Y}$, and the main part is, after an integration by parts and $X=Yu$,

$$4\int_1^\infty(\log X+1)\frac{e^{-(X+h)/Y}}{X+h}\,\mathrm{d}X=4\log Y\,E_1(h/Y)+O(e^{-h/2Y}).$$

With $h=p\beta$, $\log Y=\log p+O(1)$ and $h/Y\ge\pi\beta$, this gives $D_Q=4\log p\sum_\beta r(\beta)^2E_1(4\pi p\beta/Q)+O(1)$, and $\frac{1}{4\pi}(D_p+D_{4p})=b_3\log p+O(1)$.

Main cusps, arcs. The terms $j=0$ with $\Psi_0(c)-e^{-c}$ in place of e^{-c} are treated identically: the weight is $F(t)=(\Psi_0(t/Y)-e^{-t/Y})/t$, to which (2) applies as it stands, and in the diagonal $\int_0^\infty\Psi_0(u+v_0)\,\mathrm{d}u/(u+v_0)=\int_{-1/2}^{1/2}E_1(sv_0)\,\mathrm{d}x$ by Fubini, since $\int_0^\infty e^{-s(u+v_0)}\mathrm{d}u/(u+v_0)=E_1(sv_0)$. Subtracting $E_1(v_0)$ gives b_4.

Main cusps, correlations. For $j\ge1$ the term is $\frac{2}{Q}\sum_k a(k)a(k+jQ)\Psi_j(c)/c$ with $c=2\pi(2k+jQ)/Q=(k+jQ/2)/Y$, i.e. the weight $\frac{1}{2\pi}F(k+jQ/2)$ with $F(t)=\Psi_j(t/Y)/t$. Expanding both factors, every product containing M or two different arguments of r is bounded by (2) (the shift $jQ/2$ only increases h', and $jQ/(2Y)=2\pi j$), giving $O(e^{-\sqrt{3}\pi j})$ with geometric decay in β,β'. The squares $r(k-p\beta)^2$ occur exactly when $\beta'-\beta=jQ/p$: $\beta'=\beta+j$ for $Q=p$ and $\beta'=\beta+4j$ for $Q=4p$, now with $\beta\ge0$, the term $\beta=0$ being the free part $r(k)$ paired with the term $\beta'=jQ/p$ of the tail. With $k=a+p\beta$, $a=pu$, partial summation with (iii) gives

$\sum_a r(a)^2\Psi_j(c)/c = 4p\log p\int_0^\infty \Psi_j(c(u))\,\mathrm{d}u/c(u) + O(p)$, where $c(u) = 2\pi(2u+2\beta+j)$ for $Q = p$ and $c(u) = \pi(u+\beta+2j)$ for $Q = 4p$; the integrals are $\frac{1}{2}I_j(2\beta+j;2\pi)$ and $I_j(\beta+2j;\pi)$, and with the factors $2/Q$ the two widths give $4\log p\sum_\beta r(\beta)r(\beta+j)I_j(2\beta+j;2\pi)$ and $2\log p\sum_\beta r(\beta)r(\beta+4j)I_j(\beta+2j;\pi)$: the first two sums of b_6.

The cusp of denominator 2. Its width is $4p$, $w_d = p$ and $b = \frac{1}{p}(T^{\mathrm{odd}} - M^{\mathrm{odd}})$ (Part II, Lemma 6.12), where $T^{\mathrm{odd}}(k) = \sum_{\beta\geq 1} r^{\mathrm{o}}(\beta)r^{\mathrm{o}}(k-p\beta)$ with $r^{\mathrm{o}}(a) = r(a)$ for $a \equiv 2 \pmod 4$ and 0 otherwise (odd x, y force $x^2+y^2 \equiv 2 \pmod 8$, and conversely). The class is $\frac{1}{\pi}\sum_k (T^{\mathrm{odd}} - M^{\mathrm{odd}})^2(k)e^{-\pi k/p}/k$ (height 1) and the same with $\Psi_0 - e^{-c}$; there is no free part, the M^{odd} terms and the off-diagonal terms are $O(1)$ as above, and the diagonal gives, by the second part of (iii) (coefficient 1 in place of 4) and $Y = p/\pi$, $\frac{1}{\pi}\log p\sum_{\beta\equiv 2(4)} r(\beta)^2\int E_1(\pi\beta s)\mathrm{d}x + O(1) = b_5\log p + O(1)$, height 1 and arcs together. Its correlations are the third sum of b_6, by the computation for $Q = 4p$.

The narrow cusps. At the widths 4 and 1 the weights are $e^{-\pi k}$ and $e^{-4\pi k}$; the coefficients are $a(k) = r(k) - M(k)$ for $k < p$ with $M(k) \ll k\sigma_{-1}(k)/p$, so these classes are $\frac{1}{4\pi}\sum_k r(k)^2 e^{-4\pi k/Q}/k + O(1/p)$, and the cusp of denominator $2p$ has $T^{\mathrm{odd}}(1,p;k) = 0$ for $k < 2(p+1)$ (Part II, Lemma 6.12), hence is $O(1/p)$ as well; the same holds for their arc and correlation terms.

Adding b_1 (Lemma 6.3 of Part II, as in Section 2.1) and b_2 (Proposition 2.2) gives the theorem. The script `closed_form_B.py` evaluates $b_1,\dots,b_6$ (E_1 from `scipy`, the x- and v-integrals by quadrature, $\beta \leq 40$, $j \leq 8$) and compares them with the local slopes of the exact anatomy over $p = 2\cdot 10^5$, $4\cdot 10^5$: 0.05687/0.05676 (C_2), 0.00133/0.00134 (cusp of denominator 2 at height 1), 0.04073/0.04073 (arcs, main cusps), 0.00057/0.00058 (arcs, other cusps), $-0.00046/-0.00046$ (correlations) and 0.48204/0.48188 for the total; and the part of C_2 that remains after the diagonal is removed is -1.1691 ± 0.0001 at height 1 and -0.05762 through the arcs for all $10^4 \leq p \leq 4\cdot 10^5$. □

Remark 2.5. (a) The Siegel main term does not enter B: the sums $\sum r(k)\beta(k)$ and $\sum \beta(k)^2$ grow linearly, without a logarithm, which is Lemma 2.3(ii) and sharpens Part II, Lemma 6.4, by the factor $\log 2Y$ there. The whole of $B - b_1 - b_2 = 0.0702$ comes from the squares $r(k-p\beta)^2$, $\beta \geq 1$, i.e. from the representations $k = \alpha + p\beta$ of the coefficients $k \geq p$, through the exponential integral $E_1(4\pi p\beta/Q)$ of the position $p\beta/Y$ of the shifted copy of $\sum r(a)^2$ in the weight $e^{-k/Y}$; the terms $\beta = 1, 2$ carry all but 10^{-6} of it. (b) The same mechanism applies to the odd Heegner discriminants of Section 3, where $\sum_{a\leq X} r_D(a)^2 = \frac{24}{D+1}X\log X + O(X)$: at the two wide cusps (widths Dp and p, so $4\pi p\beta/Q \in \{4\pi\beta/D, 4\pi\beta\}$) the tail adds $\frac{6}{\pi(D+1)}\sum_{\beta\geq 1} r_D(\beta)^2\big(E_1(4\pi\beta/D) + E_1(4\pi\beta)\big)\log p$, which for $D = 4$ is b_3. Section 3.6 carries this out: Lemma 3.7 is the analogue of Lemma 2.3 for the form N_D, and Theorem 3.8 the closed form of B_D.

3. The theta series of the odd Heegner discriminants

3.1. **Setting.** Let D be one of the seven odd Heegner discriminants (the nine imaginary quadratic fields of class number one have $D \in \{3,4,7,8,11,19,43,67,163\}$, Heegner–Stark [13]), $K = \mathbb{Q}(\sqrt{-D})$, $\chi = \chi_{-D}$ the Kronecker symbol, w_D the number of units ($w_3 = 6$, $w_D = 2$ otherwise), $N_D(x,y) = x^2 + xy + \frac{D+1}{4}y^2$ (Gram determinant D), $r_D(n) = \#\{N_D = n\} = w_D\sum_{d|n}\chi(d)$ for $n \geq 1$ (class number one), and $\theta_D(z) = \sum_{n\geq 0} r_D(n)e(nz) \in M_1(\Gamma_0(D),\chi)$. The Eisenstein series $E(z,s) = \sum_{\Gamma_\infty\backslash\Gamma_0(D)} \chi(d_\gamma)j(\gamma,z)^{-1}\,\mathrm{Im}(\gamma z)^s$ has at $s = 0$ the expansion $1 + \frac{2}{L(0,\chi)}\sum\sigma_\chi(n)e(nz)$ with $L(0,\chi) = 2h/w_D = 2/w_D$ (Hecke; see [7, Chapter 4] for the weight-one Eisenstein series), so $E(z,0) = \theta_D$ without any dimension argument. For a prime $p \neq D$, $\Theta_{1,p} = \theta_D(z)\theta_D(pz) = E + G \in M_2(\Gamma_0(Dp))$, $T(k) = \sum_{\alpha+p\beta=k} r_D(\alpha)r_D(\beta)$, $a_G(k) =$

$T(k) - M(k)$. The level Dp is square-free: $\Gamma_0(Dp)$ has the four cusps $W_Q\infty$, $Q \in \{1, D, p, Dp\}$, of widths $Dp, p, D, 1$, and no cusp of denominator 2δ. We write $|_k$ for the weight-k slash operator normalised by $(\det\gamma)^{k/2}$, so that scalars act trivially and $f(mz) = m^{-k/2}(f|_k B_m)(z)$, $B_m = \left(\begin{smallmatrix} m & 0 \\ 0 & 1 \end{smallmatrix}\right)$.

3.2. **The Siegel main term.**

Theorem 3.1. *$M(k) = \dfrac{4\pi^2 k}{Dp}\prod_\ell \beta_\ell(k)$, where $\beta_\ell(k) = (1-\ell^{-2})\sigma_{-1}(\ell^{v_\ell(k)})$ for every prime $\ell \nmid Dp$, including $\ell = 2$, and, with $v = v_\ell(k)$,*

$$\beta_p(k) = (1+\chi(p)) - \chi(p)\Big(1+\frac{1}{p}\Big)p^{-v}, \qquad \beta_D(k) = (1+\varepsilon) - \varepsilon\Big(1+\frac{1}{D}\Big)D^{-v},$$

where $\varepsilon = \left(\frac{-p}{D}\right) = -\chi_{-D}(p)$. Explicitly, with $v = v_\ell(k)$:

	$\ell \nmid Dp$	$\ell = p$ *(sign $\chi(p)$)*	$\ell = D$ *(sign $\varepsilon = -\chi(p)$)*
$v = 0$	$1-\ell^{-2}$	$1-\chi(p)/p$	$1-\varepsilon/D$
$v \ge 1$, *sign* $+1$	$(1-\ell^{-2})\sigma_{-1}(\ell^v)$	$2-(1+1/p)p^{-v}$	$2-(1+1/D)D^{-v}$
$v \ge 1$, *sign* -1	$(1-\ell^{-2})\sigma_{-1}(\ell^v)$	$(1+1/p)p^{-v}$	$(1+1/D)D^{-v}$

The rows $v \ge 1$ follow from the recursions $\beta_p(k) = (1+\chi(p))(1-1/p) + \beta_p(k/p)/p$ and $\beta_D(k) = (1+\varepsilon)(1-1/D) + \beta_D(k/D)/D$ of the proof, iterated v times from the row $v = 0$. In particular $\beta_\ell(k) \le 4\sigma_{-1}(\ell^v)$ at every prime, so $M(k) \ll k\sigma_{-1}(k)/(Dp)$ uniformly in k, p and D.

Proof. The quaternary form $Q = N_D(x,y) + pN_D(z,w)$ has Gram determinant D^2p^2, and Siegel's theorem [12] (see [10, Chapter 6] for the local densities) gives $r(\operatorname{gen} Q, k) = 4\pi^2 k\prod_\ell \beta_\ell(k)/\sqrt{D^2p^2}$ with $\beta_\ell(k) = \lim_a \ell^{-3a}\#\{x \bmod \ell^a : Q(x) \equiv k\}$; the genus theta series is the Eisenstein part of $\Theta_{1,p}$. For $\ell \nmid Dp$ the form is unimodular over $\mathbb{Z}_\ell$ with square discriminant, which gives the standard density; this includes $\ell = 2$ because N_D is even unimodular over $\mathbb{Z}_2$, $pN_D \cong N_D$ there (p a unit; the Arf invariant is insensitive to unit scaling), and $N_D \oplus N_D$ has Arf invariant 0, i.e. is the sum of two hyperbolic planes, whatever the class of N_D modulo 2.

At $\ell = p$: the solutions with $(x,y) \not\equiv (0,0) \pmod p$ are non-singular in (x,y), so each solution modulo p lifts to p^{a-1} solutions modulo p^a; modulo p the equation $N_D(x,y) \equiv t$ has $p - \chi(p)$ solutions for $p \nmid t$ and $1+(p-1)(1+\chi(p))$ for $p \mid t$. With (z,w) free this gives $\beta_p(k) = 1-\chi(p)/p$ for $p \nmid k$ and the contribution $(1+\chi(p))(1-1/p)$ for $p \mid k$. The solutions with $x \equiv y \equiv 0 \pmod p$, $x = px'$, $y = py'$, satisfy $pN_D(x',y') + N_D(z,w) \equiv k/p \pmod{p^{a-1}}$, the same form with the blocks interchanged; counting (x',y') modulo p^{a-1} and the p^2 lifts of (z,w) gives $\beta_p(k/p)/p$. Hence $\beta_p(k) = (1+\chi(p))(1-1/p) + \beta_p(k/p)/p$ for $p \mid k$, which iterates to the closed form.

At $\ell = D$: since 2 is invertible, $N_D(x,y) = (x+y/2)^2 + \frac{D}{4}y^2 \cong u^2 + Dv^2$ over $\mathbb{Z}_D$, so $Q \cong u^2 + ps^2 + D(v^2+pt^2)$. The solutions with $(u,s) \not\equiv 0 \pmod D$ are non-singular in (u,s); modulo D the binary form u^2+ps^2, of discriminant $-4p$, represents $t \not\equiv 0$ exactly $D-\varepsilon$ times and $t \equiv 0$ exactly $1+(D-1)(1+\varepsilon)$ times, $\varepsilon = (-p/D)$. The solutions with $u \equiv s \equiv 0$ reduce to $D(u'^2+ps'^2) + (v^2+pt^2) \equiv k/D \pmod{D^{a-1}}$, the same form, with the factor $D^2/D^3 = 1/D$. Hence $\beta_D(k) = 1-\varepsilon/D$ for $D \nmid k$ and $\beta_D(k) = (1+\varepsilon)(1-1/D) + \beta_D(k/D)/D$ for $D \mid k$. Finally $(-p/D) = (-1/D)(p/D) = -(p/D) = -(-D/p) = -\chi_{-D}(p)$ by quadratic reciprocity, $D \equiv 3 \pmod 4$. □

3.3. **Atkin–Lehner signs and the spectral support.**

Theorem 3.2. *$\Theta_{1,p}|_2 W_p = \chi(p)\Theta_{1,p}$, $\Theta_{1,p}|_2 W_D = -\chi(p)\Theta_{1,p}$ and $\Theta_{1,p}|_2 W_{Dp} = -\Theta_{1,p}$, for any choice of the Atkin–Lehner matrices [1] of level Dp.*

Proof. Write $W_p = \left(\begin{smallmatrix} pa & b \\ Dpc & pd \end{smallmatrix}\right)$, $pad - Dbc = 1$. Then $W_p = \gamma B_p$ with $\gamma = \left(\begin{smallmatrix} a & b \\ Dc & pd \end{smallmatrix}\right) \in \Gamma_0(D)$, so $\theta_D|_1 W_p = \chi(pd)p^{1/2}\theta_D(pz)$; and $B_pW_p = p\left(\begin{smallmatrix} pa & b \\ Dc & d \end{smallmatrix}\right)$ with the second factor in $\Gamma_0(D)$, so $\theta_D(p\,\cdot)|_1 W_p = p^{-1/2}\chi(d)\theta_D$. The product is $\chi(p)\Theta_{1,p}$.

For W_D: the dual of the lattice $(\mathcal{O}_K, N_D)$ is $\mathcal{O}_K/\sqrt{-D}$, so Poisson summation gives $\theta_D(-1/z) = D^{-1/2}(-iz)\theta_D(z/D)$ and $\theta_D(-1/(Dz)) = -i\sqrt{D}z\,\theta_D(z)$, i.e. $\theta_D|_1 W_0 = -i\theta_D$ for $W_0 = \left(\begin{smallmatrix} 0 & -1 \\ D & 0 \end{smallmatrix}\right)$. Every matrix $W = \left(\begin{smallmatrix} Da' & b' \\ Dc' & Dd' \end{smallmatrix}\right)$ of determinant D is $W_0\gamma'$ with $\gamma' = \left(\begin{smallmatrix} c' & d' \\ -Da' & -b' \end{smallmatrix}\right) \in \Gamma_0(D)$, whence $\theta_D|_1 W = -i\chi(-b')\theta_D$. Now $W_D = \left(\begin{smallmatrix} Da & b \\ Dpc & Dd \end{smallmatrix}\right)$, $Dad - pbc = 1$, is such a matrix with $b' = b$, and $B_pW_D = \left(\begin{smallmatrix} Da & pb \\ Dc & Dd \end{smallmatrix}\right)B_p$ exhibits $\theta_D(p\,\cdot)|_1 W_D = p^{-1/2}(\theta_D|_1 W')|_1 B_p$ with W' such a matrix with $b' = pb$. Therefore $\Theta_{1,p}|_2 W_D = (-i)^2\chi(-b)\chi(-pb)\Theta_{1,p} = -\chi(p)\Theta_{1,p}$, using $(b, D) = 1$; and $W_{Dp} \equiv W_D W_p$ modulo $\Gamma_0(Dp)$ gives -1. □

Theorem 3.3 (Spectral support)**.** *Let φ be a newform of weight 2 and level Dp. If $\langle\Theta_{1,p}, \varphi\rangle \neq 0$ then $\eta_p(\varphi) = \chi(p)$, $\eta_D(\varphi) = -\chi(p)$, i.e. $a_\varphi(D) = \chi_{-D}(p)$, $a_\varphi(p) = -\chi_{-D}(p)$, and the root number of φ is $+1$. For a newform f of level p and $d \mid D$, $\langle\Theta_{1,p}, f(d\,\cdot)\rangle \neq 0$ requires $\eta_p(f) = \chi(p)$, i.e. $a_f(p) = -\chi_{-D}(p)$. For a newform f of level D and $d \mid p$, $\langle\Theta_{1,p}, f(d\,\cdot)\rangle = C_D(1 + \chi(p))L(1, f)L(1, f \otimes \chi)/d$ vanishes when $\chi_{-D}(p) = -1$.*

Proof. $\langle\Theta, \varphi\rangle = \langle\Theta|W, \varphi|W\rangle$ for $W = W_p, W_D$ and Theorem 3.2 give $\chi(p)\eta_p(\varphi) = 1$ and $-\chi(p)\eta_D(\varphi) = 1$; $a_\varphi(\ell) = -\eta_\ell(\varphi)$ for $\ell \parallel Dp$, and the root number is $-\eta_D\eta_p$. For f of level p, Part II, Lemma 6.8 (with $Q = p$) gives $f(d\,\cdot)|_2 W_p = \eta_p(f)f(d\,\cdot)$, and the same argument applies; W_D exchanges $f(z)$ and $f(Dz)$ up to a factor and gives no further condition. For f of level D the formula is the oldform computation in the proof of Part II, Theorem 6.7(ii), with D in the role of 4 and p in the role of n: the brackets there give $\sum_{Q_2|d}\chi(Q_2)\prod_{\ell|p/d}(1 + \chi(\ell)) = 1 + \chi(p)$ in both cases $d = 1, p$. □

Theorem 3.3 gives necessary conditions. What is proved about the actual support, and what is only observed, is the following.

Proposition 3.4. *Let $\mathcal{S}(D, p)$ be the set of newforms φ of weight 2 and level Dp with $\eta_p(\varphi) = \chi(p)$ and $\eta_D(\varphi) = -\chi(p)$.*

(i) *The newform part of $G_{1,p}^{(D)}$ lies in the span of $\mathcal{S}(D, p)$ (Theorem 3.3), and for $\varphi \in \mathcal{S}(D, p)$ one has $\langle\Theta_{1,p}, \varphi\rangle \neq 0$ if and only if $L(1, \varphi)L(1, \varphi \otimes \chi_{-D}) \neq 0$ (Theorem 3.5 below). Both central values have root number $+1$ in this situation, and their non-vanishing is not addressed here.*

(ii) *In contrast with $D = 4$, where every newform of level $4n$ has $\eta_4 = -1$ (Part II, Proposition 6.17), so that the newform support is empty whenever $\chi_{-4}(n) = -1$, neither sign of W_D is excluded at the square-free level Dp: the local component at D is the Steinberg representation or its unramified quadratic twist, and both occur. $\mathcal{S}(D, p)$ has dimension $\frac{1}{4}(\dim S^{\mathrm{new}} + \chi(p)\operatorname{tr} W_p - \chi(p)\operatorname{tr} W_D - \operatorname{tr} W_{Dp})$, the traces taken on $S_2^{\mathrm{new}}(\Gamma_0(Dp))$; these traces are sums of class numbers of size $O_\varepsilon((Dp)^{1/2+\varepsilon})$ by the trace formulae for the Atkin–Lehner involutions (Yamauchi [15] for the Fricke involution; Martin [11] for the eigenspaces with prescribed signs at square-free level), so $\#\mathcal{S}(D, p)$ is asymptotically a quarter of the new subspace, half of the forms with root number $+1$. Nothing in this paper depends on this count.*

(iii) Observed *(Section 4): for $D \leq 43$ and the primes tested, every newform component of $G_{1,p}^{(D)}$ found by the Hecke decomposition lies in $\mathcal{S}(D, p)$ (91 components). The converse, whether every $\varphi \in \mathcal{S}(D, p)$ occurs, i.e. whether $L(1, \varphi)L(1, \varphi \otimes \chi_{-D}) \neq 0$ for all of them, has not been tested: the decomposition of G finds only the forms that do occur.*

The oldforms from level D exist for $D \in \{11, 19, 43, 67, 163\}$ (genus $1, 1, 3, 5, 13$) and not for $D = 3, 7$; at the level 22 ($\chi_{-11}(2) = -1$) the cuspidal part vanishes identically, $\theta_{11}(z)\theta_{11}(2z) = E$, while at the level 55 ($\chi_{-11}(5) = +1$) the component of f_{11} is $f_{11}(z) + 5f_{11}(5z)$ up to a scalar and carries 25% of the norm.

3.4. **The Rankin–Selberg formula.**

Theorem 3.5. *For a newform φ of weight 2 and level Dp,*

$$\langle \Theta_{1,p}, \varphi \rangle = C_D\, \chi(p)\, \eta_p(\varphi)\, L(1,\varphi)\, L(1, \varphi \otimes \chi_{-D}), \qquad C_D = \frac{w_D}{4\pi L(1,\chi_{-D})} = \frac{w_D^2\sqrt{D}}{8\pi^2};$$

$C_3 = \frac{9\sqrt{3}}{2\pi^2} = 0.7897$, $C_D = \frac{\sqrt{D}}{2\pi^2}$ *for* $D \geq 7$, *against* $C_4 = \frac{4}{\pi^2}$ *and* $C_8 = \frac{\sqrt{2}}{\pi^2}$.

Proof. The proof of Part II, Theorem 6.7(i) carries over: $\theta_D = E(z, 0)$; the cusps of $\Gamma_0(Dp)$ over ∞ of $\Gamma_0(D)$ are $W_Q\infty$, $Q \mid p$; unfolding gives $I(s) = \sum_{Q|p} \chi(Q)\Gamma(s+1)(4\pi)^{-s-1} \sum_j r_D(j) c_Q((p/Q)j)((p/Q)j)^{-s-1}$, with $\chi(a)\chi(d) = \chi(Q)$ from $Qad \equiv 1 \pmod D$ and $c_Q(k) = \eta_Q(\varphi) a_\varphi(k)$; the Dirichlet series $\sum_j r_D(j) a_\varphi(j) j^{-w} = w_D \sum_j \sigma_\chi(j) a_\varphi(j) j^{-w}$ is $w_D L(w,\varphi)L(w,\varphi\otimes\chi)/(L(2w-1,\chi)\prod_{\ell|p}(1-\chi(\ell)\ell^{1-2w}))$. Here $L(w,\varphi) = \sum_n a_\varphi(n) n^{-w}$ with $a_\varphi(1) = 1$, so that the central point is $w = 1$, $L(w, \varphi\otimes\chi) = \sum_n \chi(n) a_\varphi(n) n^{-w}$, and the Euler factors are, with $x = \ell^{-w}$: for $\ell \nmid Dp$ the Rankin–Selberg factor $\sum_j \sigma_\chi(\ell^j) a_\varphi(\ell^j) x^j = (1 - \chi(\ell)\ell x^2) L_\ell(w,\varphi) L_\ell(w, \varphi\otimes\chi)$ (Part II, Theorem 6.7, with $\alpha\beta = \ell$).

The local factors at p and D. For $\ell = p$, where $a_\varphi(p^j) = a_\varphi(p)^j$, the factor $\sum_j \sigma_\chi(p^j) a_\varphi(p)^j x^j$ equals $(1 - a_\varphi(p)x)^{-1}(1 - \chi(p)a_\varphi(p)x)^{-1} = L_p L_p$, without the numerator, which is why the factor $\ell = p$ of $1/L(2w-1,\chi)$ is divided out; and for $\ell = D$, where $\chi(D) = 0$, $\sigma_\chi(D^j) = 1$ and $a_\varphi(D^j) = a_\varphi(D)^j$, the factor $\sum_j a_\varphi(D)^j x^j = (1 - a_\varphi(D)x)^{-1} = L_D(w,\varphi) L_D(w, \varphi\otimes\chi)$, the twisted factor at D being 1. Both are the local factors of $L(w,\varphi)L(w,\varphi\otimes\chi)$ at these primes, so the product over all ℓ is $L(w,\varphi)L(w,\varphi\otimes\chi)$ divided by $\prod_{\ell\nmid Dp}(1-\chi(\ell)\ell^{1-2w})^{-1} = L(2w-1,\chi)\prod_{\ell|p}(1-\chi(\ell)\ell^{1-2w})$, as displayed. At $s = 0$ the constant is $w_D/(4\pi L(1,\chi))$, and $\sum_{Q|p}\chi(Q)\eta_Q(\varphi)a_\varphi(p/Q)/(p/Q) = \chi(p)\eta_p(\varphi)(1-\chi(p)/p)$ cancels the local factor. The class number formula $L(1,\chi_{-D}) = 2\pi h/(w_D\sqrt{D})$ with $h = 1$ gives the second expression. □

3.5. **The norm.**

Theorem 3.6. *Let $D_D(s) = \sum_k r_D(k)^2 k^{-s}$. Then $D_D(s) = w_D^2\, \zeta(s)^2 L(s,\chi)^2 / \big((1 + D^{-s})\zeta(2s)\big)$ and*

$$S_D(Y) := \sum_{k\geq 1} \frac{r_D(k)^2}{k} e^{-k/Y} = \frac{12}{D+1}(\log Y)^2 + c_1^{(D)} \log Y + c_2^{(D)} + O(Y^{-1/2}),$$

$$c_1^{(D)} = \frac{24}{D+1}\Big(\gamma + 2\frac{L'}{L}(1,\chi) - 2\frac{\zeta'}{\zeta}(2) + \frac{\log D}{D+1}\Big),$$

$$\frac{L'}{L}(1,\chi_{-D}) = \gamma + \log 2 + \log\pi - \frac{w_D}{2}\sum_{a=1}^{D-1} \chi(a) \log\Gamma\Big(\frac{a}{D}\Big).$$

Consequently the regions above height 1 at the two wide cusps give

$$\|G_{1,p}^{(D)}\|^2 \;\geq\; \frac{1}{4\pi}\Big(S_D\Big(\frac{Dp}{4\pi}\Big) + S_D\Big(\frac{p}{4\pi}\Big)\Big) + O(1) = \frac{6}{\pi(D+1)}(\log p)^2 + b_D \log p + O(1),$$

$$b_D = \frac{6}{\pi(D+1)} \log\frac{D}{16\pi^2} + \frac{c_1^{(D)}}{2\pi},$$

with $b_3 = 0.713$, $b_7 = 0.207$, $b_{11} = 0.133$, $b_{19} = 0.121$, $b_{43} = 0.143$, $b_{67} = 0.150$, $b_{163} = 0.142$.

Proof. $r_D/w_D = \sigma_\chi$ is multiplicative with $\sigma_\chi(\ell^j) = j+1$, $\frac12(1+(-1)^j)$, 1 for $\chi(\ell) = 1, -1, 0$, so the Euler factor of $\sum \sigma_\chi(k)^2 k^{-s}$ at $\ell \neq D$ is that of $\zeta^2 L^2/\zeta(2s)$, and at $\ell = D$ it is $(1-x)^{-1}$ against $(1+x)/(1-x)$, whence the factor $(1+D^{-s})^{-1}$. At $s=1$: $w_D^2 L(1,\chi)^2 \cdot \frac{D}{D+1} \cdot \frac{6}{\pi^2} = \frac{24}{D+1}$ by the class number formula. Mellin inversion, $S_D(Y) = \frac{1}{2\pi i}\int_{(2)} \Gamma(s) D_D(1+s) Y^s \, \mathrm{d}s$, with the contour moved to $\operatorname{Re} s = -\frac12$ across the double pole at $s=0$ ($L(1+s,\chi)^2$ is entire, $1/\zeta(2+2s)$ is holomorphic and of polynomial growth for $\operatorname{Re} s \geq -\frac12$, and $\Gamma(s)$ decays exponentially), gives $S_D(Y)$ with the residue $\frac{12}{D+1}(\log Y)^2 + c_1^{(D)} \log Y + c_2^{(D)}$, exactly as in Part II, Lemma 6.3, where $c_1^{(D)} = \frac{24}{D+1}(\gamma + a_1)$ with a_1 the coefficient of s in $D_D(1+s) \cdot \frac{D+1}{24} s^2 = 1 + a_1 s + \dots$, i.e. $a_1 = 2\gamma + 2\frac{L'}{L}(1,\chi) - 2\frac{\zeta'}{\zeta}(2) + \frac{\log D}{D+1}$. The value of $\frac{L'}{L}(1,\chi)$ follows from the functional equation $\Lambda(s) = (D/\pi)^{s/2}\Gamma(\frac{s+1}{2}) L(s,\chi) = \Lambda(1-s)$ and $L(s,\chi) = D^{-s}\sum_a \chi(a)\zeta(s, a/D)$ with $\zeta'(0,a) = \log\Gamma(a) - \frac12 \log 2\pi$: $\frac{L'}{L}(1) = -\log\frac{D}{\pi} - \frac12\psi(\frac12) - \frac12\psi(1) - \frac{L'}{L}(0)$ and $\frac{L'}{L}(0) = -\log D + \frac{w_D}{2}\sum_a \chi(a)\log\Gamma(a/D)$, since $L(0,\chi) = 2/w_D$. The lower bound for the norm is Part II, Lemma 6.2 at the cusps $W_1\infty$ and $W_D\infty$ (widths Dp and p), which carry $\pm G_{1,p}$ by Theorem 3.2: the regions $y \geq 1$ there contribute $\frac{1}{4\pi}\sum_k |a(k)|^2 k^{-1} e^{-4\pi k/Q}$, $Q \in \{Dp, p\}$, and the difference between $|a(k)|^2$ and $r_D(k)^2$ (the Siegel main term and the coefficients $k \geq p$) contributes $O(1)$ plus the non-negative tail term $b_3^{(D)} \log p$ by Lemma 3.7 and the proof of Theorem 3.8 below. □

3.6. **The asymptotic formula for the odd discriminants.** The level Dp is square-free, so Proposition 2.1 has four classes, all Atkin–Lehner, of widths $Q \in \{Dp, p, D, 1\}$, each with $w_d = Q$ and $b = \pm a_G/Q$ (Theorem 3.2 and the computation of Part II, Lemma 6.2, whose pair count $\varphi(L)Q$ holds for every square-free L): the class of width Q is $\frac{1}{Q}\sum_{k,j} a(k)a(k+jQ)\Psi_j(c_{k,j})/c_{k,j}$. The proof of Theorem 2.4 transfers once Lemma 2.3 is available for r_D.

Lemma 3.7. *Let D be one of the odd discriminants and $r_D(0) = 1$. Uniformly for $X \geq 1$ and $h \geq 0$, with constants depending on D:*

(i) *for $h \geq 1$, $\sum_{a\leq X} r_D(a) r_D(a+h) \ll (X+h)\,\sigma_{-1}(h) + \sqrt{X+h}\,d(h)$;*
(ii) $\sum_{a\leq X} r_D(a)\sigma_{-1}(a+h) \ll X + h$;
(iii) $\sum_{a\leq X} r_D(a)^2 = \dfrac{24}{D+1} X \log X + O(X)$.

Proof. (i) Since $4N_D(x,y) = (2x+y)^2 + Dy^2$, the sum counts $(x,y,z,w) \in \mathbb{Z}^4$ with $N_D(x,y) \leq X$ and $N_D(z,w) - N_D(x,y) = h$, and the substitution $u_1 = (2z+w) - (2x+y)$, $v_1 = (2z+w) + (2x+y)$, $u_2 = w - y$, $v_2 = w + y$ is injective, gives $u_1 v_1 + D u_2 v_2 = 4h$ and keeps $|u_i|, |v_i| \leq U := 4\sqrt{X+h}$ (as $|2x+y| \leq 2\sqrt{N_D(x,y)}$ and $|y| \leq 2\sqrt{N_D(x,y)/D}$). The count is that of Lemma 2.3(i) for the equation $u_1 v_1 + D u_2 v_2 = 4h$: the terms with $u_1 = 0$ or $u_2 = 0$ give $\ll U d(4h)$; for $u_1 u_2 \neq 0$ and $g = (u_1, Du_2)$ there is no solution unless $g \mid 4h$, and then (v_1, v_2) lies on a coset of $\mathbb{Z}\,(Du_2/g, -u_1/g)$, which has at most $1 + 2Ug/\max(|u_1|, D|u_2|)$ points in the box. The 1 gives $\ll U^2$; with $u_1 = g u_1'$, $Du_2 = g u_2'$, $(u_1', u_2') = 1$, the second term gives $2U \sum_{g|4h} \sum_{|u_i'| \leq DU/g} 1/\max(|u_1'|, |u_2'|) \ll U \sum_{g|4h} DU/g \ll_D U^2 \sigma_{-1}(h)$.

(ii) As in Lemma 2.3(ii), with the number $N_D(t;e)$ of solutions of $N_D(x,y) \equiv t \pmod e$ in place of $N_2(t;e)$: $N_D(t;e)$ is multiplicative in e, and at each prime power the count for a binary form whose Gram matrix has determinant D (non-degenerate modulo every $\ell \neq D$, including $\ell = 2$; equal to $u^2 + Dv^2$ at $\ell = D$) is $N_D(t;\ell^a) \leq C\,\ell^a(a+1)$ with an absolute C, so $N_D(t;e) \leq C^{\omega(e)} e\, d(e) \ll_\varepsilon e^{1+\varepsilon}$, which is all the argument of Lemma 2.3(ii) needs (the lattice points of the ellipse $N_D \leq X$ in a residue class modulo e number $\frac{2\pi}{\sqrt{D}} X/e^2 + O(\sqrt{X}/e + 1)$).

(iii) The Dirichlet series $D_D(s)$ of Theorem 3.6 has a double pole at $s=1$ with leading coefficient $\frac{24}{D+1}$ and is otherwise holomorphic, of polynomial growth, on $\operatorname{Re} s \ge \frac12+\delta$; Perron's formula gives the assertion. □

Theorem 3.8 (The closed form of B_D)**.** *Let $D \in \{3,7,11,19,43,67,163\}$, $A_D = \frac{24}{D+1}$, and E_1, s, I_j as in Theorem 2.4. Then*

$$\|G_{1,p}^{(D)}\|^2 = \frac{6}{\pi(D+1)}(\log p)^2 + B_D \log p + O(1), \qquad B_D = b_1^{(D)} + b_2^{(D)} + b_3^{(D)} + b_4^{(D)} + b_6^{(D)},$$

$$b_1^{(D)} = b_D = \frac{6}{\pi(D+1)}\log\frac{D}{16\pi^2} + \frac{c_1^{(D)}}{2\pi}, \qquad b_2^{(D)} = \frac{A_D\kappa}{2\pi} = \frac{12\kappa}{\pi(D+1)},$$

$$b_3^{(D)} = \frac{6}{\pi(D+1)}\sum_{\beta\ge1} r_D(\beta)^2\big(E_1(4\pi\beta/D) + E_1(4\pi\beta)\big),$$

$$b_4^{(D)} = \frac{6}{\pi(D+1)}\sum_{\beta\ge1} r_D(\beta)^2\int_{-1/2}^{1/2}\big(E_1(4\pi\beta s/D) + E_1(4\pi\beta s) - E_1(4\pi\beta/D) - E_1(4\pi\beta)\big)\mathrm{d}x,$$

$$b_6^{(D)} = \sum_{j\ge1}\Big[A_D\sum_{\beta\ge0} r_D(\beta)r_D(\beta+j)\,I_j(2\beta+j;2\pi) + \frac{2A_D}{D}\sum_{\beta\ge0} r_D(\beta)r_D(\beta+Dj)\,I_j\Big(\beta+\frac{Dj}{2};\frac{4\pi}{D}\Big)\Big],$$

with $r_D(0)=1$. Numerically:

D	$b_1^{(D)}$	$b_2^{(D)}$	$b_3^{(D)}$	$b_4^{(D)}$	$b_6^{(D)}$	B_D	*local slope, Table 4*
3	0.71298	0.04319	0.05173	0.01411	−0.00091	0.82111	0.8210
7	0.20702	0.02160	0.08684	0.01369	−0.00026	0.32888	0.3289
11	0.13256	0.01440	0.13249	0.01482	−0.00017	0.29411	0.2943
19	0.12142	0.00864	0.17814	0.01439	−0.00013	0.32246	0.3214
43	0.14334	0.00393	0.20469	0.01182	−0.00009	0.36369	0.3661
67	0.14981	0.00254	0.20078	0.01018	−0.00007	0.36324	0.3648
163	0.14209	0.00105	0.17136	0.00717	−0.00005	0.32163	0.3128

Proof. The proof of Theorem 2.4 applies with r_D in place of r, Lemma 3.7 in place of Lemma 2.3, the bound $M(k) \ll k\sigma_{-1}(k)/p$ of Theorem 3.1, and the four classes listed above. The dominating estimate (2) is unchanged (it uses only the hypothesis on $\sum|c(a)|$, which (i), (ii) supply, and $|c(0)| = r_D(h) \ll d(h)$). At the two wide cusps, $Y = Q/4\pi \le Dp/4\pi$ and $h' = p\max(\beta,\beta') \ge p$, so the factor $1+Y/h'$ of (2) is at most $1+D/(4\pi)$ and the exponential is $e^{-\sqrt3 h'/(2Y)} \le e^{-2\sqrt3\pi\max(\beta,\beta')/D}$; the series over β,β' and j therefore converge as in the proof of Theorem 2.4, with $e^{-\sqrt3\pi\max/2}$ replaced by $e^{-2\sqrt3\pi\max/D}$, uniformly in p, with constants depending on D. The free part is Theorem 3.6 ($b_1^{(D)}$), and its arcs are Proposition 2.2 with the residue coefficient A_D in place of 4, i.e. $A_D\kappa\log Y$ at each wide cusp ($b_2^{(D)}$). In the diagonal of the tail, partial summation with (iii) replaces $4\log Y\,E_1(h/Y)$ by $A_D\log Y\,E_1(h/Y)$ with $h/Y = 4\pi\beta/D$ and $4\pi\beta$, whence $b_3^{(D)}$ and, through the arcs, $b_4^{(D)}$. In the correlations, $\beta' = \beta + jQ/p$ is $\beta+j$ at the width p and $\beta+Dj$ at the width Dp, where $c_{k,j} = \frac{4\pi}{D}(u+\beta+\frac{Dj}{2})$ for $k = pu+p\beta$, and the weights $\frac{2}{Q}$ give the factors A_D and $\frac{2A_D}{D}$ of $b_6^{(D)}$ (for $D=4$ these are the first two sums of b_6 in Theorem 2.4). The narrow cusps, widths D and 1, carry the weights $e^{-4\pi k/D}$, $e^{-4\pi k}$ and the coefficients $a(k) = r_D(k) - M(k)$ for $k<p$: they contribute $\frac{1}{4\pi}\sum_k r_D(k)^2k^{-1}e^{-4\pi k/Q} + O(1/p)$,

a constant, and the same for their arc and correlation terms. There is no cusp of denominator 2, so there is no analogue of b_5.

The script `heegner_closed_form_B.py` evaluates the closed forms and the exact Parseval formula for the seven odd D (Table 4); for $D=4$ its generic expressions reproduce b_3, b_4 and the main-cusp part of b_6 of Theorem 2.4 to six decimals. The local slopes of each component of the exact norm agree with $b_3^{(D)}$, $b_2^{(D)}+b_4^{(D)}$ and $b_6^{(D)}$ to three or four decimals for every D (e.g. $D=11$: 0.1327, 0.02922, -0.00017 against 0.13249, 0.02922, -0.00017), and the narrow cusps have slope 0. □

The tail term $b_3^{(D)}$ grows with D relative to the rest: $E_1(4\pi/D)$ is 0.0030 for $D=3$ but 2.06 for $D=163$, because the shifted copy of $\sum r_D(a)^2$ at the position $p\beta$ sits well inside the weight $e^{-4\pi k/(Dp)}$ of the widest cusp. For $D\geq 19$ the tail exceeds the free part $b_1^{(D)}$, and for $D=163$ the whole of $B_D=0.32$ is 28 times the leading coefficient $6/(164\pi)=0.0116$.

3.7. **The even discriminants, and the volume.** For $D=4$ ($\theta_4=\theta^2$, $w_4=4$) and $D=8$ ($\theta_8(z)=\sum e((x^2+2y^2)z)$, $w_8=2$) the theta series are again $E_1(z,\chi_{-D})$, and the level Dp has, besides the Atkin–Lehner cusps, the cusps of denominator 2δ ($D=4$) and $2\delta, 4\delta$ ($D=8$) of Part II, Section 6.6. The two ingredients that fix the constants are the same, $r_D=w_D\sigma_\chi$ and $w_D^2L(1,\chi_{-D})^2=4\pi^2/D$; hence $C_D=w_D^2\sqrt{D}/(8\pi^2)$ for all nine discriminants ($C_8=\sqrt{2}/\pi^2$ with the same proof, once the cusps of $\Gamma_0(8p)$ over ∞ of $\Gamma_0(8)$ are listed, which we have not written out), and the residue of D_D at $s=1$ is $\frac{4\pi^2}{D}\cdot\frac{6}{\pi^2}\prod_{\ell\mid D}(1+\ell^{-1})^{-1}=24/\psi(D)$, the Euler factor at each ramified prime being $(1+\ell^{-s})^{-1}$. The two wide Atkin–Lehner cusps (widths Dp and p, carrying $\pm G_{1,p}$ by the computation of Theorem 3.2, using $\theta_D(-1/(Dz))=-i\sqrt{D}z\,\theta_D(z)$ for $D=8$ as well) therefore give, uniformly for the nine discriminants,

$$\|G_{1,p}^{(D)}\|^2 \;\geq\; \frac{6}{\pi\,\psi(D)}(\log p)^2+O(\log p)=\frac{2}{\mathrm{vol}(\Gamma_0(D)\backslash\mathbb{H})}(\log p)^2+O(\log p),$$

since $\mathrm{vol}(\Gamma_0(D)\backslash\mathbb{H})=\frac{\pi}{3}\psi(D)$: $\frac{1}{\pi}$ for $D=4$ (Part II, Theorem 6.14, where it is the asymptotic), $\frac{1}{2\pi}$ for $D=8$, $\frac{6}{\pi(D+1)}$ for the odd D (Theorem 3.8, where it is the asymptotic). The growth constant is the reciprocal volume of the modular curve of the *discriminant*, not of the level: in the Rankin–Selberg normalisation $\sum_{k\leq Y}|a(k)|^2/k\sim 16\pi^2\|G\|^2Y/\mathrm{vol}(\Gamma_0(Dp)\backslash\mathbb{H})$, the index $[\Gamma_0(D):\Gamma_0(Dp)]=p+1$ cancels against the sparseness $1/p$ of the sequence $p\beta$. The value $\sqrt{D}/(2\pi)$ that one might guess from the discriminant alone is wrong already at $D=3$ (0.276 against $6/(4\pi)=0.4775$, fitted 0.4776). For $D=8$ the asymptotic formula would need the cusps of denominator 2δ and 4δ of $\Gamma_0(8p)$ (Part II, Section 6.6, for $4p$), which we have not written out; Theorem 1.2(v) states the lower bound for $D=8$ and the asymptotic for the eight other discriminants.

4. Numerical verification

The script `heegner_theta_unified.py` (self-contained: Kronecker symbol, r_D, T, M, counts of local densities, Atkin–Lehner check at a base point, Hecke decomposition by T_2 then T_3,T_5,T_7,T_{13} where eigenvalues coincide, central values by the smoothed functional equation, norms by Rankin–Selberg and by the cusp sums) was run for each odd D; the norms of Table 4 (the exact identity, truncated as stated there) are from `heegner_closed_form_B.py`, which evaluates the scalloped Parseval formula over the four Atkin–Lehner cusps with the coefficients $k\leq 3Dp$ (the first script truncates at $k\leq 12p$, which at the cusp of width Dp leaves out weights up to $e^{-48\pi/D}$, i.e. 0.03, 0.11, 0.40 for $D=43,67,163$; its norm values for these D are superseded). The checks, uniformly: the closed-form densities agree with direct counts modulo

D^a and p^a ($p \le 13$) to $2 \cdot 10^{-16}$; at the level 6, $T = M$ for all $k \le 2^{16}$ (no cusp forms); at the level 14, $T - M = \frac{2}{3} f_{14}$, at the level 15, $T - M = \frac{9}{2} f_{15}$, at the level 22, $T - M = 0$; the three Atkin–Lehner signs hold to 10^{-15} for $p \le 13$.

D	w_D	$L(1,\chi)$	$\frac{L'}{L}(1,\chi)$	C_D	$\frac{12}{D+1}$	$c_1^{(D)}$	$c_2^{(D)}$	$\frac{6}{\pi(D+1)}$	B_D	newforms, RS ratios
3	6	0.604600	0.368282	0.789720	3	16.370	13.059	0.4775	0.8211	21: [0.9944, 1.0048]
7	2	1.187410	0.015636	0.134035	1.5	5.975	1.450	0.2387	0.3289	14: [0.9966, 1.0047]
11	2	0.947226	−0.084218	0.168022	1	3.497	0.492	0.1592	0.2941	20: [0.9952, 1.0029]
19	2	0.720731	−0.084914	0.220824	0.6	2.033	0.698	0.0955	0.3225	23: [0.9965, 1.0047]
43	2	0.479088	0.249482	0.332204	0.2727	1.255	1.292	0.0434	0.3637	13: [0.9936, 1.0038]
67	2	0.383746	0.658364	0.414695	0.1765	1.093	1.505	0.0281	0.3632	(not resolved)
163	2	0.246050	2.168327	0.646797	0.0732	0.891	1.868	0.0116	0.3216	(not resolved)

TABLE 3. The constants of Theorems 3.5, 3.6 and 3.8 ($c_2^{(D)}$ numerically, from $S_D(Y)$ at $Y = 10^5$), and the newform components of $G_{1,p}^{(D)}$ resolved by the Hecke decomposition for $p \le 37$ ($D = 3$) and $p \le 19$ (otherwise), all of which obey Theorem 3.3, with the range of the ratios of the two sides of Theorem 3.5 (the twist has level $D^2 p$ and root number $+1$ in every case). $L(1,\chi)$ and $\frac{L'}{L}(1,\chi)$ agree with mpmath for $D \le 19$. For $D = 67, 163$ the dimension of $S_2(\Gamma_0(Dp))$ (≥ 33, ≥ 81) is beyond the Prony recurrence of order ≤ 11 that resolves the components, and the support is not tested; a test would need the full newform space of level Dp (modular symbols), which we have not done. Files `results/heegner_theta_unified_*.txt`; floating point.

D	$p = 101$ (RS)	1009	10007	$p_{\max}$	$\|G\|^2$ at $p_{\max}$	fitted b	B_D
3	11.890 (11.893)	26.368	45.917	400009	87.879	0.8203	0.8211
7	5.440 (5.437)	12.490	22.076	200003	38.375	0.3285	0.3289
11	4.089 (4.092)	8.971	15.532	200003	26.621	0.2935	0.2941
19	3.399 (3.402)	6.663	10.948	100003	16.243	0.3245	0.3225
43	3.251 (3.241)	5.219	7.651	40009	9.347	0.3615	0.3637
67	3.292 (3.342)	4.906	6.760	40009	8.028	0.3564	0.3632
163	3.643 (3.487)	4.678	5.841	10007	5.841	0.3192	0.3216

TABLE 4. The norm $\|G_{1,p}^{(D)}\|^2$ evaluated by the exact scalloped Parseval identity over the four Atkin–Lehner cusps of $\Gamma_0(Dp)$, truncated at the coefficients $k \le 3Dp$ and the correlations $j \le 8$ (truncation errors below 10^{-12}, see "Truncation and rounding" below; the widest cusp has the weight $e^{-4\pi k/(Dp)}$; $p_{\max}$ is limited by memory, $3Dp \le 8 \cdot 10^6$). The sandwich $\Lambda_1 \le \|G\|^2 \le \Lambda_{\sqrt{3}/2}$ of Part II, Lemma 6.11 holds at every p. In brackets at $p = 101$, the Rankin–Selberg estimate with the window 2^{21}: within 0.004 for $D \le 19$, not converged at the levels $67 \cdot 101$ and $163 \cdot 101$ ($X/L \le 310$). The fitted b is the slope of $\|G\|^2 - \frac{6}{\pi(D+1)}(\log p)^2$ in $\log p$ over $p \ge 1009$, against the closed form B_D of Theorem 3.8; the local slopes at the last two primes are in that theorem. Script `heegner_closed_form_B.py`, file `results/heegner_closed_form_B.txt`; floating point.

Four different quantities. Table 4 and Theorem 3.8 involve (a) the proved constant B_D; (b) the exact Parseval formula, truncated at $k \le 3Dp$ and $j \le 8$; (c) the slope of a linear fit of




FIGURE 2. (a) $\|G_{1,p}^{(D)}\|^2/(\log p)^2$ against p for $D = 4$ (Table 2) and the seven odd discriminants (Table 4), with the limits $2/\operatorname{vol}(\Gamma_0(D)\backslash\mathbb{H}) = 6/(\pi\psi(D))$ dotted; the approach is slow, $\frac{6}{\pi\psi(D)} + B_D/\log p$, and for $D \geq 43$ the second term still dominates. (b) The coefficient of $(\log p)^2$ fitted with the proved B_D (two parameters, $\|G\|^2 - B_D \log p = a(\log p)^2 + c$ over $p \geq 1009$) against D, with the curve $6/(\pi\psi(D))$ through the primes: 0.4774, 0.3183, 0.2387, 0.1591, 0.0956, 0.0433, 0.0277, 0.0115 for $D = 3, 4, 7, 11, 19, 43, 67, 163$ against 0.4775, 0.3183, 0.2387, 0.1592, 0.0955, 0.0434, 0.0281, 0.0116; a factor 41 between $D = 3$ and $D = 163$. $D = 8$ has not been computed. Script `plot_heegner_volume_scaling.py`; floating point.




FIGURE 3. The level 77 is $7 \cdot 11$ in two ways. The three newforms of level 77 with $a(7) = +1$, $a(11) = -1$ (and $a(2) = -\sqrt{5}, 0, \sqrt{5}$) are exactly the support of both $G_{1,11}^{(7)}$ ($\chi_{-7}(11) = +1$: $a(7) = \chi_{-7}(11)$, $a(11) = -\chi_{-7}(11)$) and $G_{1,7}^{(11)}$ ($\chi_{-11}(7) = -1$: $a(11) = \chi_{-11}(7)$, $a(7) = -\chi_{-11}(7)$). (a) Their shares of the two norms, $0.001, 0.019, 0.981$ against $0.502, 0.495, 0.005$. (b) The central values that decide them through Theorem 3.5: the same $L(1, \varphi)$, but the twists by χ_{-7} (level 539) and by χ_{-11} (level 847) favour opposite forms. Script `plot_level77_duality.py`; floating point.

$\|G\|^2 - \frac{6}{\pi(D+1)}(\log p)^2$ over the computed primes; (d) the local slope between two consecutive primes. Only (a) is a theorem; (c) and (d) estimate it with an error that comes from the $O(1)$ term of the asymptotic formula, which is not a single constant: it depends on p through $\chi_{-D}(p)$ (the densities β_p, β_D of Theorem 3.1 and the support of Theorem 3.3), so that the local slope between two primes with different $\chi_{-D}(p)$ carries the difference of two constants divided by $\log(p_2/p_1)$. The data show exactly this. For $D = 67$ the primes $1009, 4001, 10007, 20011, 40009$ have $\chi_{-67}(p) = +, -, +, -, +$, and the local slopes alternate, 0.349, 0.364, 0.352, 0.365, around $B_{67} = 0.3632$. For $D = 163$ the range is the shortest ($p \le 10007$, since $K = 3Dp$ is limited to $8 \cdot 10^6$ by memory): the pair $1009, 4001$ with equal $\chi_{-163}(p) = -1$ gives 0.3229 against $B_{163} = 0.3216$, and the last pair $4001, 10007$, where χ changes sign, gives 0.3128; this is the 0.009 of Theorem 3.8's table, and it is the same effect, not a discrepancy with the theorem. For $D = 43$ the pair $10007, 20011$ with equal $\chi = +1$ gives 0.3628 against 0.3637. Likewise the three-parameter fits of the leading coefficient are unreliable for $D \ge 43$ over these ranges (0.045, 0.030, 0.0073 against 0.0434, 0.0281, 0.0116), because $B_D \log p$ is larger than $\frac{6}{\pi(D+1)}(\log p)^2$ there, while the two-parameter fits with B_D fixed (Figure 2) give 0.0433, 0.0277, 0.0115. A run to $p = 10^5$ for $D = 163$ would need $K = 5 \cdot 10^7$; we have not done it.

The direct test of this explanation is to group the primes by the sign of $\chi_{-D}(p)$ and to take the local slopes within each group (`chi_grouped_slopes.py`, from the data of Table 4); Table 5 gives the last slope of each group. For $D \le 43$ every grouped slope lies within 0.0012 of B_D, for both signs where both groups contain two computed primes ($D = 3, 7, 19, 43$) and for the sign -1 alone for $D = 11$, whose computed primes up to 10^5 all have $\chi_{-11}(p) = -1$ (against 0.003 for $D = 19$ and 0.0024 for $D = 43$ between consecutive primes of different sign); for $D = 163$ the only pair available, $1009, 4001$ with $\chi = -1$, gives 0.3229. For $D = 67$ the two groups agree with each other, 0.3584 and 0.3589, but lie 0.005 below $B_{67} = 0.3632$: the ranges are a factor 4 to 5 in p, so a drift of 0.007 in the $O(1)$ term over the range, which the proof allows, accounts for it, and the data for $D = 67$ do not determine B_{67} beyond ± 0.005. This is a numerical check of the explanation, not a condition of the theorem.

D	$\chi_{-D}(p) = +1$: last pair, slope		$\chi_{-D}(p) = -1$: last pair, slope		B_D
3	$(100003, 400009)$	0.8211	$(10007, 200003)$	0.8206	0.8211
7	$(40009, 100003)$	0.3289	$(20011, 200003)$	0.3287	0.3289
11	(one prime)		$(40009, 100003)$	0.2936	0.2941
19	$(20011, 100003)$	0.3221	$(10007, 40009)$	0.3213	0.3225
43	$(10007, 20011)$	0.3628	$(4001, 40009)$	0.3644	0.3637
67	$(10007, 40009)$	0.3584	$(4001, 20011)$	0.3589	0.3632
163	(one prime)		$(1009, 4001)$	0.3229	0.3216

TABLE 5. Local slopes of $\|G_{1,p}^{(D)}\|^2 - \frac{6}{\pi(D+1)}(\log p)^2$ in $\log p$ between the last two computed primes with the same $\chi_{-D}(p)$, by sign, against B_D (Theorem 3.8). The primes are those of Table 4 (101, 1009, 4001, 10007, 20011, 40009, 100003, 200003, 400009 as far as computed). Script `chi_grouped_slopes.py`, file `results/chi_grouped_slopes.txt`; floating point.

Truncation and rounding. In Table 4 the sums over k stop at $K = 3Dp$, where the smallest weight, at the cusp of width Dp, is $e^{-4\pi K/(Dp)} = e^{-12\pi} = 4 \cdot 10^{-17}$; since $|a(k)| \le T(k) + M(k) \ll_\varepsilon k^{1+\varepsilon}/p$, the omitted tail $\sum_{k>K} |a(k)|^2 k^{-1} e^{-4\pi k/(Dp)}$ is $\ll_\varepsilon D^2 K^{2\varepsilon} e^{-12\pi}$, below 10^{-12} for every entry. The correlation terms with $j \ge 9$ are bounded by $|\Psi_j(c)| \le e^{-\sqrt{3}c/2} \le e^{-9\sqrt{3}\pi}$, below 10^{-21} times the height-1 sum. The functions Ψ_j are computed by 240-point Gauss–Legendre

quadrature in x (the integrand is analytic on $[-\frac12,\frac12]$) and cubic splines on a geometric grid of 8000 points in $c\in[10^{-9},70]$; against adaptive quadrature at 360 random (j,c) the splines are accurate to $2.5\cdot10^{-13}$ absolutely and to 10^{-9} relatively wherever $|\Psi_j|>10^{-6}$. The closed forms use E_1 from `scipy` (10^{-15}), 200-point Gauss–Legendre quadrature for the x-integrals of $b_4^{(D)}$, adaptive quadrature (10^{-10}) for I_j, and $\beta\le 4D+40$, where $E_1(4\pi\beta/D)<e^{-16\pi}$. The sums over k have up to $8\cdot10^6$ non-negative terms in double precision (relative rounding below 10^{-12}). So the entries of Table 4 are correct to the digits shown, and the comparison with B_D is limited by the $O(1)$ term, not by the computation.

The two curiosities of the data. At the level 77 the same three newforms carry $G_{1,11}^{(7)}$ and $G_{1,7}^{(11)}$ with shares $(0.001,0.019,0.981)$ and $(0.502,0.495,0.005)$ (Figure 3): the support conditions of Theorem 3.3 coincide for the two pairs (D,p), and the two twisted central values distribute the norm in opposite ways. And at $D=163$ the term $B_D\log p$ with $B_{163}=0.3216$ (Theorem 3.8; 0.171 of it is the tail $b_3^{(163)}$) exceeds the leading term $0.0116(\log p)^2$ for $\log p<27.6$, i.e. $p<10^{12}$: the suppression of the norm by the factor 41 is, in the range of any computation, a suppression of $(\log p)^2$ only, and at $p\le10^4$ the norms of the seven odd discriminants are all between 3 and 46.

5. Sieve limits and the Eisenstein dispersion form

5.1. **Does a large discriminant help the sieve?** For the sequence $r_D(n)r_D(N-n)$, the norm $\|G_{m,m'}^{(D)}\|^2\sim\frac{6}{\pi\psi(D)}(\log mm')^2$ enters the remainder through the uniform bound $|a_G(N)|\ll N^{1/2+\varepsilon}\|G\|$ (Part I, Theorem 2.3, whose proof is independent of D), and it decreases like $1/D$. This does not help. The main term decreases as well: the mean value of r_D is $w_DL(1,\chi_{-D})=2\pi/\sqrt{D}$, so $X_N=\sum_n r_D(n)r_D(N-n)\asymp\frac{4\pi^2}{D}N\mathfrak{S}_D(N)$ is of size N/D, while the remainder for one pair (d,e) of sieve moduli is $\ll N^{1/2+\varepsilon}(\log de)/\sqrt{\psi(D)}$; the ratio remainder/main is $\asymp\sqrt{D}\,N^{-1/2}\log de$ for prime D, larger by $\sqrt{D}$, not smaller. And the level of distribution is an exponent while D is fixed: the proof of the level $1/2$ (Part I, Theorem 3.2) loses only constants and N^ε in D, so the level is $1/2$ for every D, and the sieve dimension is 2 for every D, since $\sum_{p\le z}\omega_D(p)/p=2\log\log z+O(1)$ with $\omega_D(p)=2(1+\chi_{-D}(p))$ on average. What a large discriminant changes is the constant in the final count, $\asymp N/(D(\log N)^2)$, and, for the odd D, the shape of the fundamental domain, which simplifies the norm and has no bearing on the dispersion problem. All of this is for fixed D, with constants depending on D; nothing is claimed uniformly as $D\to\infty$, and the discriminants of class number one are finitely many in any case.

5.2. **The circle modulo c for the Eisenstein integers.**

Lemma 5.1. *Let $(c,3)=1$, c odd or even, $(a,c)=1$, $h\in\mathbb{Z}^2$, $|h|_3^2=h_1^2-h_1h_2+h_2^2$, c'^* the product of the primes dividing c' to an odd power, and $\chi=\chi_{-3}$ ($\chi(2)=-1$). Then*

$$F_c(a;h):=\sum_{\substack{x,y \bmod c\\ N_3(x,y)\equiv a\ (c)}} e\Big(\frac{h_1x+h_2y}{c}\Big)=\sum_{g\mid(c,h_1,h_2)} g\,\chi\Big(\Big(\frac{c}{g}\Big)^*\Big)S\Big(a,\ \overline{3}\,\Big|\frac{h}{g}\Big|_3^2;\ \frac{c}{g}\Big).$$

Proof. Write $N_3(x)=\frac12x^{\mathsf T}Ax$ for $x=(x,y)$ with the Gram matrix $A=\left(\begin{smallmatrix}2&1\\1&2\end{smallmatrix}\right)$, $\det A=3$, and put $G_A(t;c)=\sum_{x \bmod c}e(tN_3(x)/c)$ for $(t,c)=1$. Three steps; the first two are the Gauss sums of the hexagonal lattice, for which see also [2, Chapter 1] and [9, Section 3.5].

Step 1: $G_A(t;c)=\chi(c)\,c$ for every c prime to 3 and every t prime to c. For $(c_1,c_2)=1$ write $x=c_2x_1+c_1x_2$; then $N_3(x)\equiv c_2^2N_3(x_1)+c_1^2N_3(x_2)\pmod{c_1c_2}$, the cross term $c_1c_2\,x_1^{\mathsf T}Ax_2$ being divisible by c_1c_2, so $G_A(t;c_1c_2)=G_A(tc_2;c_1)\,G_A(tc_1;c_2)$ and it suffices to treat $c=\ell^\nu$. For $\nu\ge2$ write $x=x_0+\ell^{\nu-1}x_1$ with x_0 modulo $\ell^{\nu-1}$ and x_1 modulo ℓ: $N_3(x)\equiv N_3(x_0)+\ell^{\nu-1}x_0^{\mathsf T}Ax_1$

(mod ℓ^ν), the sum over x_1 is $\ell^2[Ax_0 \equiv 0\ (\ell)] = \ell^2[x_0 \equiv 0\ (\ell)]$ because A is invertible modulo $\ell \neq 3$, and with $x_0 = \ell x_0'$ this gives $G_A(t;\ell^\nu) = \ell^2 G_A(t;\ell^{\nu-2})$. Hence $G_A(t;\ell^\nu) = \ell^\nu$ for even ν and $\ell^{\nu-1}G_A(t;\ell)$ for odd ν. For $\nu = 1$ and $\ell \neq 2$, $4N_3 = (2x+y)^2 + 3y^2$ and the substitution $(2x+y, y) \mapsto (x, y)$ (a bijection modulo ℓ) give $G_A(t;\ell) = G(\bar4 t;\ell)G(3\bar4 t;\ell)$ with $G(a;\ell) = \sum_x e(ax^2/\ell) = \left(\frac{a}{\ell}\right)\varepsilon_\ell\sqrt{\ell}$, $\varepsilon_\ell^2 = \left(\frac{-1}{\ell}\right)$, so $G_A(t;\ell) = \left(\frac{3}{\ell}\right)\left(\frac{-1}{\ell}\right)\ell = \left(\frac{-3}{\ell}\right)\ell = \chi(\ell)\ell$; and directly $G_A(t;2) = 1-1-1-1 = -2 = \chi(2)\cdot 2$. So $G_A(t;\ell^\nu) = \chi(\ell)^\nu\ell^\nu$ in every case, independently of t, and the multiplicativity gives $G_A(t;c) = \chi(c)c$; this equals $\chi(c^*)c$ because $\chi(\ell)^2 = 1$ for $\ell \neq 3$.

Step 2: completing the square. For $(t,c) = 1$ and $h \in \mathbb{Z}^2$ the vector $v = \bar t\, A^{-1}h$ is defined modulo c, since $A^{-1} = \frac13\left(\begin{smallmatrix} 2 & -1 \\ -1 & 2\end{smallmatrix}\right)$ and 3 is invertible modulo c; and $tN_3(x+v) = tN_3(x) + t\,x^{\mathsf T}Av + tN_3(v) = tN_3(x) + h\cdot x + \bar t\,\frac12 h^{\mathsf T}A^{-1}h$, where $\frac12 h^{\mathsf T}A^{-1}h = \frac13(h_1^2 - h_1h_2 + h_2^2) = \frac13|h|_3^2$. Both sides are polynomials with integer coefficients in x and v ($N_3(x) = x^2+xy+y^2$), so no division by 2 occurs, and $\sum_{x \bmod c} e\big((tN_3(x) + h\cdot x)/c\big) = e(-\bar t\,\bar3\,|h|_3^2/c)\,G_A(t;c)$.

Step 3: the sum over t. $F_c(a;h) = \frac1c\sum_{t \bmod c} e(-ta/c)\sum_{x \bmod c} e((tN_3(x)+h\cdot x)/c)$. For $(t,c) = g$ write $t = gt'$, $c = gc'$ and $x = x_0 + c'x_1$ with x_0 modulo c', x_1 modulo g: $tN_3(x)/c \equiv t'N_3(x_0)/c'$ and $h\cdot x/c = h\cdot x_0/c + h\cdot x_1/g$ modulo 1, so the sum over x_1 is $g^2[g \mid h]$, and for $g \mid h$ the sum over x_0 is $\sum_{x_0 \bmod c'} e((t'N_3(x_0) + (h/g)\cdot x_0)/c') = e(-\bar t'\,\bar3\,|h/g|_3^2/c')\,\chi(c')c'$ by Steps 1 and 2. Collecting the t' prime to c',

$$F_c(a;h) = \sum_{g\mid(c,h)} \frac{g^2c'}{c}\,\chi(c') \sideset{}{^*}\sum_{t' \bmod c'} e\Big(\frac{-t'a - \bar t'\,\bar3\,|h/g|_3^2}{c'}\Big) = \sum_{g\mid(c,h)} g\,\chi\Big(\frac{c}{g}\Big) S\Big(a,\ \bar3\Big|\frac{h}{g}\Big|_3^2;\ \frac{c}{g}\Big),$$

by $S(-m,-n;c') = S(m,n;c')$ ($t' \mapsto -t'$); for $c' = 1$ both G_A and S are 1. The script `eisenstein_circle_check.py` verifies the identity for all a, h and all $c \le 40$ prime to 3, odd and even (285 902 cases). □

The hypotheses. The proof uses $(c,3) = 1$ exactly twice, for the invertibility of A modulo every $\ell \mid c$ (Step 1) and of 3 (Step 2); it uses $(a,c) = 1$ nowhere, so the identity holds for every a with $S(a,b;c')$ defined by the same sum (the script checks all a), and it holds for $c = 1$ ($F_1 = 1$, $g = 1$). Even moduli need no separate argument: for $\ell = 2$ the reduction $G_A(t;2^\nu) = 4G_A(t;2^{\nu-2})$ uses only $\det A = 3 \not\equiv 0 \pmod 2$, and the base case $G_A(t;2) = -2$ is a direct count, so no Gauss sum of odd modulus enters, and 2^ν with ν large is covered by the induction. This is where the Eisenstein case differs from the Gaussian one: for $x^2+y^2 = \frac12 x^{\mathsf T}(2I)x$ the Gram matrix has $\det = 4$, Step 1 fails at $\ell = 2$, and Part III, Lemma 3.1 needs c odd. The condition $g \mid (c,h)$ means $g \mid \gcd(c,h_1,h_2)$, in Theorem 1.3 and here.

5.3. **The dispersion form.** Let m, m' be coprime square-free integers prime to $3N$. Since the ideal $(\sqrt{-3})$ is principal of norm 3, $r_3(3\beta') = r_3(\beta')$, and the terms of $T(m,m';N)$ with $3 \mid \beta$ are $T(m,3m';N)$: with $T^\flat(m,m';N) = \sum_{3\nmid\beta} r_3(\alpha)r_3(\beta)$,

$$a_{G_{m,m'}}(N) - a_{G_{m,3m'}}(N) = T^\flat(m,m';N) - M^\flat(m,m';N),$$

the analogue of $a_G(N) - a_G(N/2) = T^{\mathrm{odd}} - M^{\mathrm{odd}}$ of Part III. In $T^\flat$, $r_3(\beta) \neq 0$ forces $\beta \equiv 1 \pmod 3$, so $\chi(\beta/k) = \chi(k)$ and $r_3(\beta) = 12\sum_{k\mid\beta,\,k<\sqrt\beta}\chi(k) + 6\chi(\sqrt\beta)[\beta = \square]$, the divisors k being prime to 3 and to m but possibly even. Hence

$$T^\flat(m,m';N) = 12 \sum_{\substack{k<\sqrt{N/m'} \\ 3\nmid k}} \chi(k)\,\mathcal{N}_k + 6\sum_j \chi(j)\,r_3^\flat\Big(\frac{N-m'j^2}{m}\Big),$$

$$\mathcal{N}_k = \#\Big\{(x,y):\ N_3(x,y) \equiv N\bar m\ (m'k),\ mN_3(x,y) \not\equiv N\ (3),\ 0 < N_3(x,y) < Y_k\Big\},$$

$Y_k = (N - m'k^2)/m$. The condition modulo 3 is the analogue of the parity condition $x + y$ odd of Part III: Poisson summation on the classes modulo $3c$, $c = m'k$, gives

$$\mathcal{N}_k = \frac{Y_k}{(3c)^2} \sum_{h \in \mathbb{Z}^2} \hat{w}\Big(\frac{h\sqrt{Y_k}}{3c}\Big)\, \epsilon_3(h)\, F_c(N\bar{m};\ \bar{3}h),$$

where $\bar{3}h$ is the Chinese-remainder twist and $\epsilon_3(h)$ the Fourier transform of the admissible classes modulo 3: as $N_3(x,y) \equiv (x-y)^2 \pmod 3$, the condition is $x \equiv y$ if $N\bar{m} \equiv 1$, $x \not\equiv y$ if $N\bar{m} \equiv 0$, and empty if $N\bar{m} \equiv 2 \pmod 3$, so that $\epsilon_3(h) = 3[h_1 + h_2 \equiv 0\ (3)]$, $9[h \equiv 0\ (3)] - 3[h_1 + h_2 \equiv 0\ (3)]$, 9, respectively. Since $|h|_3^2 \equiv (h_1 + h_2)^2 \pmod 3$, the condition $h_1 + h_2 \equiv 0 \pmod 3$ is $3 \mid |h|_3^2$: exactly the condition $h_1 \equiv h_2 \pmod 2$, i.e. $2 \mid |h|^2$, of Part III, with the ramified prime of the field in the role of 2. By Lemma 5.1 applied to the vector $\bar{3}h$, whose norm is $|\bar{3}h|_3^2 \equiv \bar{9}|h|_3^2 \pmod c$, the leading term $g = 1$ of $F_c(N\bar{m}; \bar{3}h)$ is $\chi((m'k)^*)S(N\bar{m}, \bar{3}\cdot\bar{9}|h|_3^2; m'k) = \chi((m'k)^*)S(N\bar{m}, \overline{27}|h|_3^2; m'k) = \chi((m'k)^*)S(\overline{27m}N, |h|_3^2; m'k)$, the last step because a Kloosterman sum depends on its two arguments only through their product, here $N\bar{m}\cdot\overline{27}|h|_3^2 = \overline{27m}N|h|_3^2$ modulo $m'k$, with $\overline{27m} = (27m)^{-1} \bmod m'k$ (Section 1.3; checked numerically in `kloosterman_argument_audit.py`); the parameter rule is the same as in the Gaussian case, with $\bar{3}\cdot\bar{9} = \overline{27}$ in place of $\overline{16}$, the $\bar{3}$ from A^{-1} in Step 2 of Lemma 5.1 and the $\bar{9}$ from the twist $\bar{3}h$; and $\chi(k)\chi((m'k)^*) = \chi(m')$ for square-free m'.

Proposition 5.2. *The leading term ($g = 1$) of $a_{G_{m,m'}}(N) - a_{G_{m,3m'}}(N)$ is*

$$12\,\chi(m') \sum_{\substack{k < \sqrt{N/m'} \\ 3\nmid k}} \frac{Y_k}{(3m'k)^2} \sum_{h \neq 0} \hat{w}\Big(\frac{h\sqrt{Y_k}}{3m'k}\Big)\, \epsilon_3(h)\, S\big(\overline{27m}\,N,\ |h|_3^2;\ m'k\big),$$

and the remainder sum of the sieve at the level N^θ has as its leading term the trilinear form $\sum_{m'} \Lambda'_{m'} \sum_{c \equiv 0\,(m')} \sum_m \Lambda_m \sum_j b_{m,j}(c) S(\overline{27m}N, j; c)$ over the moduli $c = m'k$ prime to 3, odd or even, with $j = |h|_3^2 \ll mm'$ a norm from $\mathbb{Z}[\zeta_3]$, whose m-th piece is, by the Chinese remainder theorem, the bilinear form in the two moduli

$$\mathfrak{B}_m^{(3)} = \sum_{m'} \Lambda'_{m'} \sum_{k \le K} \sum_h w_{k,m'}(h)\, S\big(\overline{27m}N\bar{k},\ |h|_3^2\bar{k};\ m'\big)\, S\big(\overline{27m}N\overline{m'},\ |h|_3^2\overline{m'};\ k\big).$$

What is an identity and what is a bound. Behind Proposition 5.2 is the exact decomposition, as in Part III, Proposition 3.2,

$$a_{G_{m,m'}}(N) - a_{G_{m,3m'}}(N) = R_1 + R_{g>1} + \Sigma_\square + \Sigma_\partial,$$

where R_1 is the displayed leading term (the terms $g = 1$, $h \neq 0$ of Lemma 5.1; the terms $h = 0$, summed over k, are the Siegel main term $M^\flat$), $R_{g>1}$ collects the terms $g \mid (m'k, h)$ with $g > 1$, sums of the same shape with the modulus $m'k/g$ and the dual variable h/g, $\Sigma_\square = 6\sum_j \chi(j) r_3^\flat((N - m'j^2)/m) \ll N^{1/2+\varepsilon}m'^{-1/2}$ is the ternary term, and Σ_∂ the error of the smooth cutoff at Y_k. This identity, Lemma 5.1 and the shape of the bilinear form are what is proved here. The bounds for R_1 that are asserted are exactly those of Part III, which only open the Kloosterman sums or use their Weil mean square: the Weil bound $|S(a, |h|_3^2; c)| \le d(c)c^{1/2}(|h|_3^2, c)^{1/2}$ gives the level 1/4, the theta-function bound (Part I, Theorem 2.3, whose proof applies to θ_3 unchanged) the level 1/2, and the Cauchy–Schwarz arrangements (A1), (A2) and the Poisson step in h of Part III, Section 4, transfer verbatim because they depend only on the lengths of the variables ($|h| \ll m'k\sqrt{m/N}$, $H \le \sqrt{mm'}$) and on $\sum_h |S|^2$. No bound beyond the level 1/2 is proved or claimed, for R_1 or for the whole remainder, and $R_{g>1}$, $\Sigma_\square$, Σ_∂ are not estimated here, exactly as in Part III, Remark 3.3(iv).

Comparison with the Gaussian case. Nothing in the structure changes. (i) The fixed argument is the single integer N, carried by a unit ($\overline{27m}$ in place of $\overline{16m}$) modulo the modulus, so the obstacle of Part III, Section 3.4, the absence of an average over it and the inapplicability of the Kuznetsov formula and of the bounds of [6, 8], is the same. (ii) The sizes are the same: $|h| \ll m'k\sqrt{m/N}$, the h-box has side $H \le \sqrt{mm'}$, below $\sqrt{q}$ for every modulus $q = [c_1, c_2]$ of a Poisson step, and the Cauchy–Schwarz arrangements (A1), (A2) of Part III, Section 4 give the same diagonals ($E^{1/4}N^{3/4}$ and $\sqrt{EN}$); there is no tighter grouping, because the groupings depend only on the lengths of the variables and on the Weil mean square of Kloosterman sums over the dual lattice, which is the same for the hexagonal lattice. (iii) The moduli $m'k$ range over all k prime to 3, even k included: more moduli, no gain. (iv) The 2-adic "impurity" of Part III has moved to 3: the condition $3 \nmid \beta$ produces $\epsilon_3(h)$ and $3 \mid |h|_3^2$, as $2 \nmid \beta$ produced $h_1 \equiv h_2 \pmod 2$. This is not an accident: the ramified prime is where the ideal of norm ℓ is principal, and removing it from the divisor sum of $r_D(\beta)$ is what makes the divisors $k < \sqrt{\beta}$ pair with β/k under χ. The clean level Dp matters for the holomorphic side (Section 3), not for the dispersion form; a level beyond $1/2$ needs the bilinear saving over (m', k) of Part III, Section 4.5, in the Eisenstein setting as in the Gaussian one.

6. Outlook

In summary: the Petersson norm of $G_{1,p}$ is an exact Parseval sum over the scalloped fundamental domain, and its second coefficient $B = 0.48204$ is a closed form of six pieces (Theorems 1.1 and 2.4); for the seven odd discriminants of class number one the Siegel main term, the Atkin–Lehner signs, the necessary support conditions, the Rankin–Selberg constant C_D and the asymptotic with the closed form B_D are proved for all seven odd D by one argument, with constants depending on D (Theorem 1.2), with the actual occurrence of a newform decided by $L(1,\varphi)L(1,\varphi\otimes\chi_{-D}) \neq 0$ (Proposition 3.4), the lower bound $\frac{2}{\mathrm{vol}}(\log p)^2$ for all nine discriminants and only the lower bound for $D = 8$; for the Eisenstein integers the circle modulo any c prime to 3 has a Kloosterman Fourier transform (Lemma 5.1), the dispersion form is that of Part III with $\overline{27m}$ for $\overline{16m}$ and 3 for 2 (Proposition 5.2), and no level beyond $1/2$ is proved for any discriminant. The problem of Parts I–IV, $N = n + (N - n)$ with both summands norms from an imaginary quadratic field of class number one and few prime factors, is a binary additive problem of the Goldbach type with two sparse sequences, and it shares with Goldbach's problem the shape of its obstacles: a level of distribution $1/2$ obtained from the modular side (here from a theta series, there from the Bombieri–Vinogradov theorem), a sieve whose dimension (here 2) fixes the attainable pair (k, k') at that level, and a remainder beyond $1/2$ that is a bilinear form in Kloosterman sums with one argument fixed. What the present part adds to this picture is structural. The spectral support of the theta series is governed by the Atkin–Lehner signs at the ramified prime and at p (Theorem 3.3): the remainder sees only the newforms whose root number is $+1$ and whose signs match $\chi_{-D}(p)$, so the parity of the root number, which decides the vanishing of $L(1,\varphi)$, enters the sieve remainder at the level of individual forms; the Rankin–Selberg formula (Theorem 3.5) writes the remainder as a sum of products of two central values, a twist by χ_{-D} against the trivial one, so that the automorphic lift to GL_2 of the two quadratic fields (the one of the discriminant and the one of the twist) is where the arithmetic of N is encoded; and the norm, which measures the size of the remainder on average, is the reciprocal volume of $X_0(D)$ (Theorem 1.2(v)), the modular curve of the discriminant, which is rational for the four small discriminants $D \in \{3, 4, 7, 8\}$ and has genus $1, 1, 3, 5, 13$ for $D = 11, 19, 43, 67, 163$; the second coefficient B_D (Theorems 2.4 and 3.8) has, for all eight discriminants treated, the same five pieces, the free part, the arcs, the tail of exponential integrals at the two wide cusps, its arcs, and the correlations, so that a uniform formula for B_D over imaginary quadratic fields, with the

class group entering through the theta series of the classes, seems within reach, and would be the natural continuation. We do not claim that any of this moves the Goldbach problem; the honest content is that the two problems have the same anatomy, and that in the theta-series setting every piece of that anatomy (main term, support, central values, norm) has been made explicit. The question that remains, here as there, is the bilinear form of Part III, Section 4.5.

Data and code

The scripts `subleading_log_anatomy.py` and `closed_form_B.py` (Section 2; they import the tables and the Siegel main term of Part II), `heegner_theta_unified.py` and `heegner_closed_form_B.py` (Sections 3–4), `eisenstein_circle_check.py` (Lemma 5.1), `kloosterman_argument_audit.py` (Sections 1.3 and 5) and the plotting scripts `plot_scalloped_anatomy.py`, `plot_heegner_volume_scaling.py`, `plot_level77_duality.py`, with their outputs in `results/` and the working notes in `notes/`, are in the repository https://github.com/Ruqing1963/two-squares-part4, archived with this manuscript at Zenodo, doi:10.5281/zenodo.23866356 (version 1.0).
Numerical methodology. Python 3.12.4 with numpy 1.26.4, scipy 1.13.1, matplotlib 3.8.4, mpmath 1.3.0 and sympy 1.13.1; IEEE double precision throughout, except for the L-values checked with mpmath. Table 2 is `subleading_log_anatomy.py` (output `results/subleading_log_anatomy.txt`), the constants of Theorem 2.4 are `closed_form_B.py`, Table 3 is `heegner_theta_unified.py` (one D per run for $D \geq 19$, outputs `results/heegner_theta_unified_*.txt`), Table 4 and the constants of Theorem 3.8 are `heegner_closed_form_B.py` (about ten minutes and 1 GB), Lemma 5.1 is `eisenstein_circle_check.py`, and Figures 1–3 are the three plotting scripts reading these outputs; the run commands are in the README. The truncation and rounding errors of the norm computations are estimated in Section 4 ("Truncation and rounding"): tails below 10^{-12}, Ψ_j to $2.5 \cdot 10^{-13}$, E_1 to 10^{-15}. All computations are in floating point and are evidence, not proof; the theorems do not depend on them.

References

[1] A. O. L. Atkin and J. Lehner, Hecke operators on $\Gamma_0(m)$, Math. Ann. 185 (1970), 134–160.
[2] B. C. Berndt, R. J. Evans and K. S. Williams, Gauss and Jacobi Sums, Wiley, New York, 1998.
[3] R. Chen, Sums of two squares with few prime factors via theta series of quaternary forms, preprint (2026), doi:10.5281/zenodo.23826176.
[4] R. Chen, Sums of two squares with few prime factors, II: the mean square of the cuspidal error of $\theta^2(mz)\theta^2(m'z)$, almost all N, and the size of the Petersson norm, preprint, version 3 (2026), doi:10.5281/zenodo.23866064 (version 2, doi:10.5281/zenodo.23826297, precedes the correction of Section 1.3).
[5] R. Chen, Sums of two squares with few prime factors, III: the dispersion obstacle, sieve barriers at the level 1/2, and short-interval mean squares, preprint (2026), doi:10.5281/zenodo.23826860.
[6] J.-M. Deshouillers and H. Iwaniec, Kloosterman sums and Fourier coefficients of cusp forms, Invent. Math. 70 (1982), 219–288.
[7] F. Diamond and J. Shurman, A First Course in Modular Forms, Springer, 2005.
[8] S. Drappeau, Sums of Kloosterman sums in arithmetic progressions, and the error term in the dispersion method, Proc. London Math. Soc. 114 (2017), 684–732.
[9] H. Iwaniec and E. Kowalski, Analytic Number Theory, AMS Colloquium Publications 53, 2004.
[10] Y. Kitaoka, Arithmetic of Quadratic Forms, Cambridge Tracts in Mathematics 106, Cambridge University Press, 1993.
[11] K. Martin, Refined dimensions of cusp forms, and equidistribution and bias of signs, J. Number Theory 188 (2018), 1–17; arXiv:1609.05386.
[12] C. L. Siegel, Über die analytische Theorie der quadratischen Formen, Ann. of Math. 36 (1935), 527–606.
[13] H. M. Stark, A complete determination of the complex quadratic fields of class-number one, Michigan Math. J. 14 (1967), 1–27.

[14] F. Waibel, Uniform bounds for norms of theta series and arithmetic applications, Math. Proc. Cambridge Philos. Soc. 173 (2022), 669–691; arXiv:2012.04311 (equation numbers refer to the arXiv version).
[15] M. Yamauchi, On the traces of Hecke operators for a normalizer of $\Gamma_0(N)$, J. Math. Kyoto Univ. 13 (1973), 403–411.

GUT Geoservice Inc., Montreal, Canada
Email address: ruqing@hotmail.com