

Riverside Family Medicine

After-Visit Clinical Note

Patient: Alex Rivera DOB: 03/14/1985 (Age 41) Sex: M
MRN: 00482913 Date of Service: 2026-06-12 Provider: Sarah Lee, MD

Chief Complaint

Fatigue, increased thirst, and intermittent blurred vision for approximately 6 weeks.

History of Present Illness

41-year-old male presents with a 6-week history of fatigue, polydipsia, polyuria, and intermittent blurred vision. Reports increased nocturia (2-3x/night) and unintentional 6 lb weight loss. No chest pain, no focal neuro deficits. Family history significant for type 2 diabetes (father).

Vitals

BP 138/86 mmHg HR 78 Temp 98.4F BMI 31.2 kg/m2

Assessment

1. Type 2 Diabetes Mellitus, newly diagnosed (E11.9). HbA1c 7.8%.
2. Essential hypertension (I10).
3. Mixed hyperlipidemia (E78.2).

Plan

- Initiate metformin 500 mg PO BID with meals; titrate to 1000 mg BID over 4 weeks as tolerated.
- Medical nutrition therapy; ≥ 150 min/week moderate-intensity physical activity.
- Referral to ophthalmology for dilated diabetic retinopathy screening.
- Order urine albumin-to-creatinine ratio (microalbuminuria screen).
- Repeat HbA1c in 3 months. Continue lisinopril 10 mg daily.
- Follow-up appointment in 3 months, or sooner as needed.

Patient Instructions

You have been diagnosed with type 2 diabetes. This means your blood sugar (glucose) is higher than the normal range. Your HbA1c, a 3-month average of your blood sugar, is 7.8% (a value of 6.5% or higher indicates diabetes).

Start metformin as directed with food to reduce stomach upset. Pick up your prescription at your pharmacy. Take your medication daily.

When to Seek Care

Symptoms of low blood sugar: shakiness, sweating, confusion, rapid heartbeat.

Persistent vomiting, very high blood sugar, or vision changes that worsen.

Electronically signed by Sarah Lee, MD on 2026-06-12 18:42

