

Math, Multivariable Calculus

RustColeone

May 22, 2020

Contents

1	Vectors	3
1.1	Vector Calculations	3
2	Vector functions	3
2.1	Vector Valued functions in multiple variables	3
2.2	Limits and continuity	4
2.3	Partial Derivatives	6
2.4	Properties of Derivatives DF	7
2.5	Directional derivatives	9
2.6	Partial derivative along a direction	10
2.7	Summary	12
2.8	Gradients and level sets	12
2.9	Parametric Curves	14
2.10	Acceleration	15
2.11	Curvature	16
2.12	Higher order derivatives and Taylor expansions	19
2.13	Taylor Expansions	20
2.14	Vector Field and Flow	27
2.15	Integration along a path	29
2.16	Gradient Vector Field	31
2.17	Double Integrals	33

1 Vectors

1.1 Vector Calculations

2 Vector functions

2.1 Vector Valued functions in multiple variables

Calculus of single variable is a study of $y = f(x)$

$$y = f(x), \text{ where } (x \in \mathbb{R}) \rightarrow (f(x) \in \mathbb{R}) \quad (1)$$

(2)

However, when we are dealing with multi-variable calculus, we have: an n variable vector valued function (if $m > 1$)

$$F = \begin{cases} (x_1, \dots, x_n) & \rightarrow F(x_1, \dots, x_n) \\ \in & \in \\ \mathbb{R}^n & \longrightarrow \mathbb{R}^m \end{cases} \quad (3)$$

where $F(x_1, \dots, x_n) = (F_1(x_1, \dots, x_n), F_2(x_1, \dots, x_n) \dots F_m(x_1, \dots, x_n))$


The inputs and outputs are Vectors, for example

$$d(x, y) = \sqrt{x^2 + y^2}, (2 \text{ variables}) \quad (4)$$

$$(x, y) \mathbb{R}^2 \rightarrow (\sqrt{x^2 + y^2}) \mathbb{R} \quad (5)$$

$$\text{distance} = \sqrt{x^2 + y^2} \quad (6)$$



Graph for $f(x, y) = \sqrt{(x^2 + y^2)}$





The graph should look like a cone

Similarly, we can have a function that takes all the x , y and z can calculate that distance. That would however be a four dimensional plot, which we cannot plot.

For a similar function, $z = x^2 + y^2$



For another function $f(z) = (\cos z, \sin z)$




2.2 Limits and continuity

Definition:

$$\lim_{(x,y) \rightarrow (a,b)} f(x,y) = L \quad (7)$$

if $f(x, y)$ goes to L as (x, y) approaches (a, b)



Similarly, for $(x, y) \rightarrow (a, b)$ we need to look at all possible paths. If you can find 2 oaths to approach (a, b) such limits along these 2 paths are different, then the function is not continuous. For example:

$$\lim_{(x,y) \rightarrow (0,0)} \frac{2xy}{x^2 + y^2} = \text{DNE}(\text{do not exists}) \text{ since:} \quad (8)$$

$$(x, 0) \rightarrow (0, 0), \lim = \lim_{x \rightarrow 0} \frac{0}{x^2} = 0 \quad (9)$$

$$(x, x) \rightarrow (0, 0), \lim = \lim_{y \rightarrow 0} \frac{2x * x}{x^2 + x^2} = 1 \quad (10)$$

$1 \neq 0$ hence D.N.E.

Remember to take a look at all directions

Definition: $f(x, y)$ continuous at (a, b) if:

1. $f(x, y)$ defined at (a, b)
2. $\lim_{(x,y) \rightarrow (a,b)} f(x, y) = f(a, b)$

How to tell if a function is continuous? $f(x, y, z) = e^{x-z} \sin(x + yz^2)$

1. (a) $f(x, y, z) = x$
 (b) $f(x, y, z) = y$
 (c) $f(x, y, z) = \text{zallcontinuous}$
2. $f(x, y, z) = x - z$
3. $f(x, y, z) = e^{x-z}(\text{composition of contin. func is contin.})$
4. $g(x, y, z) = \sin(x + yz^2)$ continuous similarly

Test: $\lim_{(x,y,z) \rightarrow (0,1,-1)} = e^1 \sin(1)$, hence continuous. A special case is when you have $\frac{f(x,y)}{g(x,y)}$, it will be continuous at (a, b) as long as $g(a, b) \neq 0$

For a vector function $F(\vec{v}) = (F_1(\vec{v}), F_2(\vec{v}) \dots F_m(\vec{v}))$, where \vec{v} is a vector or $\mathbb{R}^n, (x_1, x_2 \dots x_n)$, is a continuous function at vector \vec{n} if all its component functions are continuous at \vec{n}

Example:

$$f(x, y, z) = (e^{x+y}, z, \cos(x^2 - z)) \text{ continuous everywhere} \quad (11)$$

$$\text{since } \begin{cases} e^{x+y} \text{ continuous} \\ z \text{ continuous} \\ \cos(x^2 - z) \text{ continuous} \end{cases} \quad (12)$$

2.3 Partial Derivatives

The derivative of a function is the rate of change along the x axis. The partial derivatives of $F(x, y)$ would be the rate of change of F as one of the axis is fixed (along one axis).

$$\frac{dF(x, y)}{dx} = F_x(x, y) = \lim_{\Delta x \rightarrow 0} \frac{F(x + \Delta x, y) - F(x, y)}{\Delta x} \quad (13)$$

$$\frac{dF(x, y)}{dy} = F_y(x, y) = \lim_{\Delta y \rightarrow 0} \frac{F(x, y + \Delta y) - F(x, y)}{\Delta y} \quad (14)$$

In order to compute the derivative, we treat all other variables as constants.

Example:

$$f(x, y, z) = xe^{xy} - \sin(y^2 + z^2) \quad (15)$$

$$\frac{df}{dx} = e^{xy} + yxe^{xy} - 0 = e^{xy} + yxe^{xy} \quad (16)$$

$$\frac{df}{dy} = x^2e^{xy} - \cos(y^2 + z^2)(2y) \frac{df}{dz} = 0 - \cos(y^2 + z^2)(2z) \quad (17)$$

Take a point (a, b) for example

1. $\frac{df}{dx}$ = gradient of tangent at that point along x axis
2. $\frac{df}{dy}$ = gradient of tangent at that point along y axis
3. These two direction can span the tangent plane of f at that point
 - (a) Tangent Vector
 - (b) along x direction = $(1, 0, \frac{df}{dx}(a, b)) = \vec{v}$
 - (c) along y direction = $(0, 1, \frac{df}{dy}(a, b)) = \vec{w}$
 - (d) Using these vectors we can find the plane's equation

Example, find the tangent plane of $f(x, y)$ at $(1, 1)$:

$$f(x, y) = 1 - x^2 - 2y^2 \quad (18)$$

$$f(1, 1) = z = -2, \text{ point at } (1, 1, -2) \quad (19)$$

$$\frac{df}{dx} = (1, 0, -2x)|_{(1,1)} = (1, 0, -2) \quad (20)$$

$$\frac{df}{dy} = (1, 0, -4y)|_{(1,1)} = (0, 1, -4) \quad (21)$$

$$(1, 0, -2) \times (0, 1, -4) = (2, 4, 1) \quad (22)$$

$$2x + 4y + z = 4 \text{ a linear approximation of } f(x, y) \text{ at } (1, 1) \quad (23)$$

Definition of a vector function:

$$F(x_1, x_2, \dots, x_n) = F(\vec{x}) = (F_1(\vec{x}) \dots F_m(\vec{x})) \in \mathbb{R}^m \quad (24)$$

Where each F_i is a component function. The derivative the size $(m \times n)$ matrix of F (Jacobian matrix) denoted by $DF(\vec{x})$

$$DF(\vec{x}) = \begin{pmatrix} \frac{dF_1}{dx_1} & \frac{dF_1}{dx_2} & \dots & \frac{dF_1}{dx_n} \\ \frac{dF_2}{dx_1} & \frac{dF_2}{dx_2} & \dots & \frac{dF_2}{dx_n} \\ \dots & \dots & \dots & \dots \\ \frac{dF_m}{dx_1} & \frac{dF_m}{dx_2} & \dots & \frac{dF_m}{dx_n} \end{pmatrix} \quad (25)$$

The function F transforms $\mathbb{R}^n \rightarrow \mathbb{R}^m$, $DF: \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a linear map, a linear approximation of F .

Example:

$$F(x, y, z) = (e^{x+yz}, x^2 + 1, \sin(y + z), 4y) \quad (26)$$

$$DF(x, y, z) = \begin{pmatrix} e^{x+yz} & ze^{x+yz} & ye^{x+yz} \\ 2x & 0 & 0 \\ 0 & \cos(y + z) & \cos(y + z) \\ 0 & 4 & 0 \end{pmatrix} DF(1, 1, 2) = \quad (27)$$

$$DF(1, 1, 2) = \begin{pmatrix} e^3 & 2e^3 & e^3 \\ 2 & 0 & 0 \\ 0 & \cos(3) & \cos(3) \\ 0 & 4 & 0 \end{pmatrix} \quad (28)$$

Matrix of numbers, a linear map

$$F(x, y, z) = (e^{x+yz}, x^2 + 1, \sin(y + z), 4y) \quad (29)$$

2.4 Properties of Derivatives DF

Given two vector function F and $G: \mathbb{R}^n \rightarrow \mathbb{R}^m$ both differentiable at some point \vec{x}

$$D(f \pm G) = D(F) \pm D(G) \quad (30)$$

$$C \in \mathbb{R}, D(cF) = cD(F) \quad (31)$$

suppose $f, g : \mathbb{R}^n \rightarrow \mathbb{R}$ a function in n variables, product rule still holds

$$D(f \cdot g) = D(f) \cdot g + f \cdot D(g) \quad (32)$$

where $D(f \cdot g)$ is a n^{th} dimension vector, $D(f \text{ or } g)$ is a vector function and f or g is a scalar function

$$D\left(\frac{f}{g}\right) = \frac{g \cdot D(f) - f \cdot D(g)}{g^2}, g(\vec{x}) \neq 0 \quad (33)$$

Given two vector function $\vec{v}(t)$ and $\vec{w}(t) : \mathbb{R} \rightarrow \mathbb{R}^n$ so $t \rightarrow \vec{v}(t)$

$$f(t) = \vec{v}(t) \cdot \vec{w}(t) \text{ scalar function} \quad (34)$$

$$f'(t) = \vec{v}(t)' \cdot \vec{w} + \vec{w}(t)' \cdot \vec{v} \quad (35)$$

$$u(t) = \vec{v}(t) \times \vec{w}(t) \text{ vector function} \quad (36)$$

$$u'(t) = \vec{v}(t)' \times \vec{w}(t) + \vec{w}(t)' \times \vec{v}(t) \quad (37)$$

$$\vec{v}(t) = \langle v_1(t), v_2(t), \dots, v_n(t) \rangle \quad (38)$$

Chain rule, suppose $f : \mathbb{R}^m \rightarrow \mathbb{R}^n$, $g : \mathbb{R}^n \rightarrow \mathbb{R}^p$, $g \circ f : \mathbb{R}^m \rightarrow \mathbb{R}^p$, $\vec{x} \in \mathbb{R}^m$

$$D(g \circ f)(\vec{x}) (\text{size } p \times m) = DG(F(\vec{x})) (\text{size } p \times n) \cdot DF\vec{x} (\text{size } n \times m) \quad (39)$$

$$(g \circ f)'(x) = g'(f(x)) \cdot f'(x) \quad (40)$$

Example:

$$f(x, y) = (x^3 + y, e^{xy}, 2 + xy) : \mathbb{R}^2 \rightarrow \mathbb{R}^3 \quad (41)$$

$$g(y, v, w) = (y^2 + v, uv + w^3) : \mathbb{R}^3 \rightarrow \mathbb{R}^2 \quad (42)$$

$$g(f(x, y)) = g \circ f = ((x^3 + y)^2 + e^{xy}, (x^3 + y)e^{xy} + (2 + xy)^3) \quad (43)$$

$$D(g \circ f) = \begin{pmatrix} 2(x^3 + y)(3x^2) + ye^{xy} & 2(x^3 + y) + xe^{xy} \\ 3x^2e^{xy} + (x^3 + y)ye^{xy} & e^{xy} + (x^3 + y)xe^{xy} \\ +3(2 + xy)^2y & +3(2 + xy)^2x \end{pmatrix} \quad (44)$$

$$DG = \begin{pmatrix} 2u & 1 & 0 \\ v & u & 3w^2 \end{pmatrix} \quad (45)$$

$$(g \circ f)'(x) = g'(f(x)) \cdot f'(x) \quad (46)$$

$$DG = \begin{pmatrix} 2(x^3 + y) & 1 & 0 \\ e^{xy} & x^3 + y & 3(2 + xy)^2 \end{pmatrix} \times \begin{pmatrix} 3x & 1 \\ ye^{xy} & xe^{xy} \\ y & x \end{pmatrix} \quad (47)$$

Check that the equality holds.

2.5 Directional derivatives

Recall the Jacobian matrix, in the following context, the function F is not necessarily defined on the entire \mathbb{R}^n , but rather a subset $\subseteq \mathbb{R}^n$

$$F(x_1, x_2, \dots, x_n) = F(\vec{x}) = (F_1(\vec{x}) \dots F_m(\vec{x})) \in \mathbb{R}^m \quad (48)$$

Where each F_i is a component function. The derivative the size $(m \times n)$ matrix of F (Jacobian matrix) denoted by $DF(\vec{x})$

$$DF(\vec{x}) = \begin{pmatrix} \frac{dF_1}{dx_1} & \frac{dF_1}{dx_2} & \dots & \frac{dF_1}{dx_n} \\ \frac{dF_2}{dx_1} & \frac{dF_2}{dx_2} & \dots & \frac{dF_2}{dx_n} \\ \dots & \dots & \dots & \dots \\ \frac{dF_m}{dx_1} & \frac{dF_m}{dx_2} & \dots & \frac{dF_m}{dx_n} \end{pmatrix} \quad (49)$$

Examples: 1)

$$f(x, y, z) = x^2y + \ln z : \mathbb{R}^3 \rightarrow \mathbb{R}^1 \quad (50)$$

$$\begin{aligned} Df &= \left(\frac{df}{dx}, \frac{df}{dy}, \frac{df}{dz} \right) \\ &= (2x, 1, \frac{1}{z}) \end{aligned} \quad (51)$$

2)

$$\vec{v}(t) = (\cos(t), \sin(t)); \text{ vector function} \quad (52)$$

$$D\vec{v}(t) = \begin{pmatrix} -\sin(t) \\ \cos(t) \end{pmatrix} = (-\sin(t), \cos(t))^T \quad (53)$$

$$D\vec{v}(t) \cdot \vec{v}(t) = -\sin(t) \cos(t) + \cos(t) \sin(t) = 0 \quad (54)$$

$$D\vec{v}(t) \perp \vec{v}(t); \text{ hence a tangent vector} \quad (55)$$

3)

$$g(t) = e^t \text{ scalar function} \quad (56)$$

$$g(t) \cdot \vec{v}(t) = e^t(\cos(t), \sin(t)) = (e^t \cos(t), e^t \sin(t)) \quad (57)$$

$$D(g(t) \cdot \vec{v}(t)) = \begin{bmatrix} e^t \cos(t) - e^t \sin(t) \\ e^t \cos(t) + e^t \sin(t) \end{bmatrix} \quad (58)$$

4)

$$\vec{v}(t) = (t, \sin(t), \cos(t)), \vec{w}(t) = (3t, 0, 2) \quad (59)$$

$$\begin{aligned} f(t) &= \vec{v}(t) \cdot \vec{w}(t) \\ &= 3t^2 + 0 + 2 \cos(t); \text{ scalar function} \end{aligned} \quad (60)$$

$$D(f(t)) = 6t - 2 \sin(t) \quad (61)$$

$$\vec{w} \cdot D\vec{v} + \vec{v} \cdot D\vec{w} = \begin{bmatrix} 3t \\ 0 \\ 2 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ +\cos(t) \\ -\sin(t) \end{bmatrix} + \begin{bmatrix} t \\ \sin(t) \\ \cos(t) \end{bmatrix} \cdot \begin{bmatrix} 3 \\ 0 \\ 0 \end{bmatrix} \quad (62)$$

$$= 6t - 2 \sin(t) \quad (63)$$

2.6 Partial derivative along a direction

Definition: Given a function $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ is differentiable at (a, b) , given a unit vector $\vec{u} = (u_1, u_2) \in \mathbb{R}^2$ then the directional derivatives of F along \vec{u} is:

$$D_{\vec{u}}F(a, b) = \left[\frac{d}{dt} f((a, b) + t\vec{u}) \right]_{t=0} \quad (64)$$

$$= \lim_{h \rightarrow 0} \frac{f(a + hu_1, b + hu_2) - f(a, b)}{h} \quad (65)$$

is the rate of change of F along the direction of \vec{u}

Two special cases would be when u are along the typical x or y direction, which would mean the class derivative along x -axis $\frac{df}{dx}$ or the y -axis $\frac{df}{dy}$

Theorem $\vec{u} = (u_1, u_2)$ unit vector,

$$D_{\vec{u}}f(x, y) = \left(\frac{df}{dx}, \frac{df}{dy} \right) \cdot \vec{u} = (u_1, u_2) \quad (66)$$

$$\text{since } \vec{u} = u_1(1, 0) + u_2(0, 1) \quad (67)$$

$$D_{\vec{u}}f = u_1 \frac{df}{dx} + \frac{df}{dy} \quad (68)$$

Proof:

$$D_{\vec{u}}f = \lim_{h \rightarrow 0} \frac{F(a + hu_1, b + hu_2) - F(a, b)}{h} \quad (69)$$

$$= \lim_{h \rightarrow 0} \frac{(F(a + hu_1, b + hu_2) - F(a, b + hu_2))}{h} \quad (70)$$

$$+ \lim_{h \rightarrow 0} \frac{F(a, b + hu_2) - F(a, b)}{h} \quad (71)$$

$$= \lim_{u_1 h \rightarrow 0} \frac{[F(a + hu_1, b + hu_2) - F(a, b + hu_2)] \cdot u_1}{hu_1} \quad (72)$$

$$+ \lim_{h \rightarrow 0} \frac{F(a, b + hu_2) - F(a, b)}{h} \quad (73)$$

$$= \frac{df}{dx}(a, b + 0) \cdot u_1 + \frac{df}{dy} \cdot u_s \quad (74)$$

Example: Find the directional Deriv of $f(x, y) = x^2 + 3xy$ along the direction

of $(3, 4)$ at the point $p = (2, -1)$

$$\vec{u} = \frac{(3, 4)}{\sqrt{(3^2 + 4^2)}} = \left(\frac{3}{5}, \frac{4}{5}\right) \quad (75)$$

$$D_{\vec{u}}f(2, -1) = (2x + 3y, 3x)|_{(2, -1)} \cdot \left(\frac{3}{5}, \frac{4}{5}\right) \quad (76)$$

$$= (1, 6) \cdot \left(\frac{3}{5}, \frac{4}{5}\right) \quad (77)$$

$$= \frac{27}{5} \quad (78)$$

$$D_{\vec{u}}f = \nabla f \cdot \vec{u} = \left(\frac{df}{dx}, df dy\right) \cdot \vec{u} \quad (79)$$

Definition: Given $F : \mathbb{R}^n \rightarrow \mathbb{R}$, then define the gradient.

$$\nabla f = D_{\vec{u}}f = \left(\frac{df}{x_1}, \frac{df}{x_2}, \dots, \frac{df}{x_n}\right) \quad (80)$$

Hence the directional derivative of a function along a direction is:

$$D_{\vec{u}}f = \nabla f \cdot \vec{u} \quad (81)$$

The consequence of this:

1)

$$D_{\vec{u}}f \text{ max when } \vec{u} // \nabla f = \left(\frac{df}{dx}, \frac{df}{dy}\right) \quad (82)$$

since

$$\nabla f \cdot \vec{u} = |\nabla f| |\vec{u}| \cos(\theta) \quad (83)$$


$$= |\nabla f| \cdot 1 \cdot \cos(\theta) \theta = \text{ang}(\nabla f, \vec{u}) \quad (84)$$

When maximized, $//, \theta = 0$

$$\nabla f \cdot \vec{u} = |\nabla f| = \sqrt{\frac{df^2}{dx} + \frac{df^2}{dy}} \quad (85)$$

2)

$$\text{When } \vec{u} \perp \nabla f, D_{\vec{u}}f = 0 \quad (86)$$



Example: $f(x, y) = e^{-(x^2 + y^2)}$, find the direction along this function which has the largest rate of increase/decrease at point $(1, 1)$

Since at maximum, $\vec{u} \parallel \nabla f$

$$\nabla f = (-2xe^{-(x^2+y^2)}, -2ye^{-(x^2+y^2)})|_{(1,1)} \quad (87)$$

$$= (-2e^{-2}, -2e^{-2}) \quad (88)$$

$$\therefore \vec{u} \parallel \nabla f \parallel (1, 1) \quad (89)$$

$$\therefore \vec{u} = \left(-\frac{1}{\sqrt{2}}, -\frac{1}{\sqrt{2}}\right) \quad (90)$$

Largest rate of increase at $\left(\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right)$

Largest rate of decrease at $-\left(\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right)$

The gradient ∇f tells us the direction along which F changes the most.

2.7 Summary

Given a function $f(x, y)$	n-th dim analogue
Define Partial derivatives (deriv) $\frac{df}{dx}, \frac{df}{dy}$ or (f_x, f_y)	$f(x_1, x_2, \dots, x_n)$
Define gradient $\frac{df}{dx}, \frac{df}{dy}$ or (f_x, f_y)	$f : \mathbb{R}^n \rightarrow \mathbb{R}$
Directional deriv: given unit vector $\vec{u} = (u_1, u_2)$	$\frac{df}{dx_1} \dots \frac{df}{dx_n}$
$D_{\vec{u}}f = \lim_{h \rightarrow 0} \frac{f((a,b)+h\vec{u}) - f(a,b)}{h}$	$\nabla f = (\uparrow)$
if $(\vec{u} = (1, 0))$, $D_{\vec{u}}f = \frac{df}{dx}$	$\vec{u}f \in \mathbb{R}^n$
if $(\vec{u} = (0, 1))$, $D_{\vec{u}}f = \frac{df}{dy}$	$D_{\vec{u}}f = ?$
Theorem: $D_{\vec{u}} = \nabla f \cdot \vec{u} = u_1 \frac{df}{dx} + u_2 \frac{df}{dy}$	generalizations?
Remark: We can define $D_{\vec{u}}$ for any $\vec{v} \in \mathbb{R}^2$ in the exact same way $D_{\vec{u}} = \vec{v} D_{\frac{\vec{v}}{ \vec{v} }} f$	

To each unit vector \vec{u} , we have a tangent vector associated to it:

$$(u_1, u_2, D_{\vec{u}}f) = u_1(1, 0, \frac{df}{dx}) + u_2(0, 1, \frac{df}{dy}) \quad (91)$$

Two special cases: $(1, 0, \frac{df}{dx})(0, 1, \frac{df}{dy})$

The two tangent vector span a tangent plane at $(a, b, f(a, b))$

2.8 Gradients and level sets

Recall

$$\nabla f = \left(\frac{df}{dx}, \frac{df}{dy}\right) \quad (92)$$

$$\therefore D_{\vec{u}} = \nabla f \cdot \vec{u} = |\nabla f| |\vec{u}| \cos(\theta) \quad (93)$$

$D_{\vec{u}}$ is the rate of change along \vec{u}

if $u \parallel \nabla f$, ($\theta = 0$) then $D_{\vec{u}}F$ max


$$|D_{\vec{u}}F| = \sqrt{\left(\frac{df}{dx}\right)^2 + \left(\frac{df}{dy}\right)^2} \quad (94)$$

In other words, along the direction of ∇F , the function F changes the most.

Definition: A level set of $F(x,y)$ is:

$$(x, y) \in \mathbb{R}^2 : f(x, y) = c \quad (95)$$

Example:



$$f(x, y) = x^2 + \frac{y^2}{4} \quad (96)$$

$$\{f(x, y) = 1\} \text{ (Level set)} \quad (97)$$

$$= \{(x, y) | x^2 + \frac{y^2}{4} = 1\} \quad (98)$$

Theorem: Given $f(x, y)$ differentiable, then the gradient $\nabla f \perp$ the level set.

We can prove it by using the chain rule: let $\gamma(t)$ be a parametric function of my level set $\{f(x, y) = 1\}$, $f(\gamma(t)) = 1$ and differentiate both sides.

$$0 = D(f(\gamma(t))) = Df(\gamma(t))D(\gamma(t)) \quad (99)$$

$$= \nabla F(\gamma(t))_{1 \times 2} \cdot (\gamma'_1(t), \gamma'_2(t))_{2 \times 1} \quad (100)$$

$$\text{Hence } \Rightarrow \nabla F(\gamma(t))_{1 \times 2} \perp (\gamma'_1(t), \gamma'_2(t))_{2 \times 1} \quad (101)$$

Example: Look at a surface $\sin(xy) - 2\cos(yz) = 0$ Find the tangent plane of this surface at $(\frac{\pi}{2}, 1, \frac{\pi}{3})$. The idea is that consider the surface as a level set of $F(x, y, z) = \sin(xy) - 2\cos(yz)$, $\{F = 0\}$

Using the Theorem: ∇F is hence perpendicular to this surface, giving us the normal direction of our tangent plane.

$$\nabla F = \left(\frac{df}{dx}, \frac{df}{dy}, \frac{df}{dz}\right) = (y \cos(xy), x \cos(xy) + 2z \sin(yz), 2y \sin(tz)) \quad (102)$$

$$= \left(\cos\left(\frac{\pi}{2}\right), \frac{\pi}{2} \cos\left(\frac{\pi}{2}\right) + \frac{2\pi}{3} \sin\left(\frac{\pi}{3}\right), 2 \sin\left(\frac{\pi}{3}\right)\right) \quad (103)$$

$$= \left(0, \frac{\sqrt{3}\pi}{3}, \sqrt{3}\right) \quad (104)$$

$$0 = 0 \cdot \left(x - \frac{\pi}{2}\right) + \frac{\sqrt{3}\pi}{3}(y - 1) + \sqrt{3}\left(z - \frac{\pi}{3}\right) \quad (105)$$

Where the last equation above is the equation of the tangent plane.

2.9 Parametric Curves

... are vector functions in 1 variable.

Definition: A parametric curve in \mathbb{R}^n is a vector valued function where $\vec{c}(t) : \mathbb{R}^1 \rightarrow \mathbb{R}^n$

All the component functions of $\vec{c}(t)$ are functions in 1 variable.

Example:

1) Draws a Circle

$$\vec{c}(t) = (\cos(t), \sin(t)), t \in [0, 2\pi] \quad (106)$$

2) Draws a ellipse at plane $z = 3$

$$\vec{c}(t) = (a \cos(t), b \sin(t), 3), t \in [0, 2\pi] \quad (107)$$

3) Ellipse spiraling up

$$\vec{c}(t) = (a \cos(t), b \sin(t), t), t \in [0, 2\pi] \quad (108)$$

Definition: Orientation of $\vec{c}(t)$, the direction corresponding to increasing t-value is called the positive orientation of $\vec{c}(t)$. The opposite direction is the negative orientation.

Example: Positive orientation = Clockwise

$$\vec{c}(t) = (\cos(t), -\sin(t), t), t \in [0, 2\pi] \quad (109)$$

Derivative of $\vec{c}(t)$

If $\vec{c}(t) = (x(t), y(t), z(t))$, $\vec{c}'(t) = (x'(t), y'(t), z'(t))$

1. $\vec{c}(t)$ is a vector function
2. $\vec{c}'(t)$ is a tangent to curve $\vec{c}(t)$

Terminology: $\vec{c}'(t)$ is the velocity of $\vec{c}(t)$, and $|\vec{c}'(t)|$ is the speed of $\vec{c}(t)$. The speed of the curve $\vec{c}(t) = (\cos(t), \sin(t))$ for example, is 1.

What is the length of curve $\vec{c}(t) = (\cos(t), \sin(t), t)$ between (0 to 2π) Given $\vec{c}(t)$ differentiable between range a and b, length of the curve would be

$$l = \int_a^b |\vec{c}'(t)| dt \quad (110)$$

$$|\vec{c}'(t)| = \sqrt{\sin(t)^2 + \cos(t)^2 + 1} = \sqrt{2} \quad (111)$$

$$l = \int_0^{2\pi} \sqrt{2} dt \quad (112)$$

Think of length as a function in t

$$l(t) = \int_a^t |\vec{c}'(x)| dx \quad (113)$$

If curve $\vec{c}(t)$ differentiable and $\vec{c}'(t) \neq 0$,

1. $|\vec{c}'(t)| > 0$
2. $l(t)$ strictly increasing.
3. l is one to one function so has inverse.

$$l[a, b] \rightarrow \mathbb{R}^1, l \text{ has an inverse.} \quad (114)$$

$$t \rightarrow l(t), \text{ think of our parameter } t \quad (115)$$

$$(116)$$

t can now be expressed as a function in terms of length, $\vec{c}(t) = \vec{c}(l(t)) = \vec{c}(l)$

Proof:

In our last example of calculating length, we can find an inverse for $l(t) = \sqrt{2}t$ resulting $t = \frac{l}{\sqrt{2}}$, by substitution we can reparameterize $\vec{c}(t)$ into $\vec{c}(l)$

$$\vec{c}(l) = \vec{c}(l(t)) \quad (117)$$

$$\frac{d\vec{c}(l)}{dl} = \frac{\vec{c}'(t(l))}{dt} \cdot \frac{dt}{dl} \quad (118)$$

$$= \vec{c}'(t) \cdot \frac{1}{|\vec{c}'(t)|} \quad (119)$$

$$|\frac{d\vec{c}(l)}{dl}| = |\vec{c}'(t)| \cdot \frac{1}{|\vec{c}'(t)|} = 1 \quad (120)$$

2.10 Acceleration

Acceleration is the second derivatives of length, hence the definition of second derivative can be expressed as below:

Definition: Given $\vec{c}(t)$, $\vec{a}(t) = \vec{c}''(t)$ the acceleration of \vec{c} .

Example:

1) Ellipse

$$\vec{c}_1(t) = (\cos(t), 2 \sin(t)), t \in [0, 2\pi] \quad (121)$$

$$\vec{c}'_1(t) = (-\sin(t), 2 \cos(t)) \quad (122)$$

$$\vec{c}''_1(t) = (-\cos(t), -2 \sin(t)) \quad (123)$$

2) Different para of the same Ellipse

$$\vec{c}_2(t) = (\cos(t^2), 2 \sin(t^2)), t \in [0, \sqrt{2\pi}] \quad (124)$$

$$\vec{c}'_2(t) = 2t(-\sin(t^2), 2 \cos(t^2)) \quad (125)$$

$$\vec{c}''_2(t) = (-2 \cos(t^2) - 4t^2 \cos(t^2), 4 \cos(t^2) - 8t^2 \sin(t^2)) \quad (126)$$

Facts:

1. We can have different parameterations of a curve
2. Velocity, acceleration depend on the para we choose in general
3. For different parameterization \vec{c}_1, \vec{c}_2 of the same curve, $\vec{c}_1'(t) // \vec{c}_2'(t)$
4. "Unit Velocity" $\frac{\vec{c}_1'(t)}{|\vec{c}_1'(t)|} = \pm \frac{\vec{c}_2'(t)}{|\vec{c}_2'(t)|}$ are equal, the \pm depends on the orientations of the \vec{c}_1 and \vec{c}_2 .

Definition: Given $\vec{c}(t)$, assume $|\vec{c}'(t)| \neq 0$, defined the unit tangent vector $\vec{T}(t) = \frac{\vec{c}'(t)}{|\vec{c}'(t)|}$. If $\vec{c}_1(t)$ and $\vec{c}_2(t)$ are two different parameter of a curve with the same orientation, their unit tangent vector equals.

2.11 Curvature

Idea: Curvature is a measure of how sharply a curve is bending at a point, it is the "rate of change" of directions along a curve. So a straight line has no curvature, and a slightly curved line has a small curvature.

Definition: Quantitatively, if $\vec{c}(t)$ is parametered by length, (namely, $|\vec{c}'(t)| = 1$), then we define the curvature at $l = l_0$ to be $k(l_0) = |\vec{c}''(l_0)| = \left| \frac{d^2 \vec{c}(l)}{dl^2} \right|_{l=l_0}$

Remark: IF $\vec{c}(t)$ not parametered by length, then $|\vec{c}''(t)|$ is not the curvature.

Explanation: a curve is like a highway and a parameterized function $\vec{c}(t)$ is like someone driving along this highway giving his position vector at a given time. Parameterized by length would be driving at a constant/normalized speed. A curvature is how sharp a turn is, measured by looking at how quickly the driver need to turn their steering wheel.

Generally, $|\vec{c}'(l)| = 1 \Rightarrow \vec{c}'(l)$ unit tangent, so $\vec{c}''(t)$ is curvature.

Example:

1) Find parameter by length

$$\vec{c}(t) = \vec{a} + t\vec{v} \quad (127)$$

$$\text{Constant vectors: } \vec{a} = (1, 2, 3), \vec{v} = (0, 2, 1)$$

$$\text{Length function } l(t) = \int_0^t |\vec{c}'(x)| dx \quad (128)$$

$$= \int_0^t |\vec{v}| dx, \quad \vec{v} = (0, 2, 1) \quad (129)$$

$$l = l(t) = \sqrt{5}t \quad (130)$$

$$t = t(l) = \frac{l}{\sqrt{5}} \quad (131)$$

Curvature would be zero since it is a line, its second derivative would be a zero.

2) Find the curvature for a circle

$$\vec{c}(t) = (r \cos(t), r \sin(t)), r > 0 \text{ circle } x^2 + y^2 = r^2 \quad (132)$$

$$\text{length function } l(t) = \int_0^t |\vec{c}'(x)| dx = rt \quad (133)$$

$$t(l) = \frac{l}{r} \quad (134)$$

$$\vec{c}(l) = (r \cos(\frac{l}{r}), r \sin(\frac{l}{r})) \quad (135)$$

$$\vec{c}''(l) = (-\frac{\cos(l/r)}{r}, -\frac{\sin(l/r)}{r}) \quad (136)$$

$$k(l) = |\vec{c}''(l)| \quad (137)$$

$$= \sqrt{(-\frac{\cos(l/r)}{r})^2 + (-\frac{\sin(l/r)}{r})^2} = \frac{1}{r} \quad (138)$$

Curvature for circle of radius $r = \frac{1}{r}$

Second Definition of Curvature:

If $\vec{c}(t)$ any parametric curve, $\vec{c}'(t) \neq 0$, recall the unit tangent vector $\vec{T} = \frac{\vec{c}'(t)}{|\vec{c}'(t)|}$,

then define curvature $k(t) = \frac{|\vec{T}'(t)|}{|\vec{c}'(t)|}$

Theorem: Definition 1 and 2 of curvature are the same.

$$\vec{c}'(t) = \frac{d\vec{c}(t(l))}{dt} \quad (139)$$

$$= \frac{d\vec{c}(t)}{dt} \cdot \frac{dt}{dl} \quad (140)$$

$$= \frac{\vec{c}'(t)}{|\vec{c}'(t)|} = \vec{T} \quad (141)$$

$$k(l(t)) = |\vec{c}''(t)| = \left| \frac{d^2\vec{c}}{dl^2} \right| = \left| \frac{d\vec{T}(t(l))}{dl} \right| \quad (142)$$

$$= \frac{d\vec{c}'(t)}{dt} \cdot \frac{dt}{dl} \quad (143)$$

$$= |\vec{T}'(t)| \cdot \frac{1}{|\vec{c}'(t)|} = k(t) \quad (144)$$

Using the second definition to find the curvature of the parabola $y = x^2$

$$\vec{c} = (t, t^2) \quad (145)$$

$$\vec{c}'(t) = (1, 2t) \quad (146)$$

$$\vec{T}(t) = \frac{(1, 2t)}{\sqrt{1 + 4t^2}} \quad (147)$$

$$\vec{T}'(t) = \frac{(0, 2)}{\sqrt{1 + 4t^2}} + (1, 2t) \cdot ((a + 4t^2)^{-\frac{1}{2}})' \quad (148)$$

$$= \frac{(-4t, 2)}{(1 + 4t^2)^{\frac{3}{2}}} \quad (149)$$

$$|T'(t)| = \frac{\sqrt{4 + 16t^2}}{(1 + 4t^2)^{\frac{3}{2}}} \quad (150)$$

$$= \frac{2}{1 + 4t^2} \quad (151)$$

$$K(t) = |T'(t)| \cdot \frac{1}{|c'(t)|} \quad (152)$$

$$= \frac{2}{(1 + 4t^2)} \cdot \frac{1}{\sqrt{1 + 4t^2}} \quad (153)$$

$$= 2 \cdot (1 + 4t^2)^{-\frac{3}{2}} \quad (154)$$

As x approaches ∞ , the curvature approaches 0, $y = x^2$ became more like a straight line.

Osculating Plane and Circle

1. Derivative — Tangent line
2. Curvature — Osculating Circle

Observations:

1) If $\vec{c}(l)$ parameterized by length, then

$$\vec{c}'(l) \perp \vec{c}''(l) \quad (155)$$

$$\because (\vec{c}'(l) \cdot \vec{c}'(l))' = (|\vec{c}'(l)|^2)' = (1)' \quad (156)$$

$$\vec{c}'(l) \cdot \vec{c}''(l) + \vec{c}'(l) \cdot \vec{c}''(l) = 0 \quad (157)$$

$$\Rightarrow \vec{c}'(l) \cdot \vec{c}''(l) = 0 \quad (158)$$

so second derivative is the normal and the first derivative is the tangent.

2) If $\vec{c}(t)$ is any parameterized curve, its unit tangent vector \vec{T} and \vec{T}' is

$$\vec{T}(t) = \frac{\vec{c}'(t)}{|\vec{c}'(t)|} \quad (159)$$

$$\because (\vec{T}(t) \cdot \vec{T}(t))' = (|\vec{T}(t)|^2)' = (1)' \quad (160)$$

$$\Rightarrow \vec{T}(t) \cdot \vec{T}'(t) = 0 \quad (161)$$

$$\therefore \vec{T}(t) \perp \vec{T}'(t) \quad (162)$$

Definition: Osculating Plane: Given $\vec{c}(t)$, $\vec{c}'(t) \neq 0$, then the osculating plane is the plane spanned by $\vec{T}(t)$ and $\vec{T}'(t)$

Recall that we computed for a circle with radius r its curvature $k = \frac{1}{r}$, therefore, if have a $\vec{c}(t)$ with $k(t_0) = k$, then at this point, the curve $\vec{c}(t)$ is "modeled" by a circle of radius $\frac{1}{k}$

Definition: Osculating Circle: Given $\vec{c}(t)$, $\vec{c}'(t) \neq 0$, then the osculating circle of \vec{c} at $t = t_0$ is a circle on the osculating plane at $t = t_0$ and has radius $\frac{1}{k(t_0)}$, and the tangent to the curve. (The best approximation of the curve at that point by the circle)

Example: Find the equation of the osculating plane at $\vec{c}(\pi) = (-a, 0, b\pi)$ of $\vec{c}(t) = (a \cos(t), a \sin(t), bt)$, $t \geq 0$; $a, b > 0$,

$$\vec{T} = \frac{\vec{c}'(t)}{|\vec{c}'(t)|} = \frac{(-a \sin(t), a \cos(t), b)}{\sqrt{a^2 + b^2}} \quad (163)$$

$$\vec{T}' = \frac{(-a \cos(t), -a \sin(t), 0)}{\sqrt{a^2 + b^2}} \quad (164)$$

$$\text{Osc.Plane} = \vec{T} \times \vec{T}' \quad (165)$$

$$= \frac{1}{a^2 + b^2} (ab \sin(t), -ab \cos(t), a^2)|_{t=\pi} \quad (166)$$

$$= \frac{1}{a^2 + b^2} (0, ab, a^2) \quad (167)$$

calculating plane at given point

$$a^2 b \pi = 0x + aby + a^2 z \quad (168)$$

2.12 Higher order derivatives and Taylor expansions

Recall partial derivatives of $F(x, y)$ by x and y denoted by F_x and F_y , second order derivative $= F_{xx}$ and F_{yy} , first x then y F_{xy} and first y then x F_{yx} , vise versa.

Theorem: If $F(x, y)$ differentiable and F_{xy} , F_{yx} continuous, $F_{xy} = F_{yx}$, hence the order can be switched

Example: $f(x, y, z) = x^2 e^{y^2 - x} + x^2 y z^2 - \cos(x^2 + y^2)$, $f_{xzzz} = ?$ switch order, try finding f_{zzxy} instead

$$df/dz = f_z = 2zx^2y \Rightarrow f_{zz} = 2x^2y \Rightarrow f_{zzxy} = 4x \quad (169)$$

2.13 Taylor Expansions

One Variable: $y = f(x)$, $f'(x)$ =, tangent line of graph of the function, its linear approximation at point $x = x_0$, equation of tangent line is $y = f(x_0) + f'(x_0)(x - x_0)$. Higher order $f'(x) \rightarrow$ higher degree approximation.

$$f(x) - f(x_0) = \int_{x_0}^x f'(t) dt \quad (170)$$

$$\int u'v = uv - \int v'u, \quad u = t, v = f'(t) \quad (171)$$

$$= tf'(t)|_{x_0}^x - \int_{x_0}^x t df'(t) \text{ by parts} \quad (172)$$

$$= [xf'(x) - x_0 f'(x_0)] - \int_{x_0}^x t f''(t) dt \quad (173)$$

$$= (xf'(x) - x_0 f'(x_0)) + (x_0 f'(x_0) - x_0 f'(x_0)) - \int_{x_0}^x t f''(t) dt \quad (174)$$

$$= (x - x_0)f'(x_0) + x(f'(x) - f'(x_0)) - x \int_{x_0}^x t f''(t) dt \quad (175)$$

$$= f'(x_0)(x - x_0) + x \int_{x_0}^x f''(t) dt - \int_{x_0}^x t f''(t) dt \quad (176)$$

$$f(x) = f(x_0) + f'(x_0)(x - x_0) + \int_{x_0}^x (x - t) f''(t) dt \quad (177)$$

$$(178)$$

This is the 1 st Taylor expansion where $f(x_0) + f'(x_0)(x - x_0)$ is the 1st order approximation, a tangent line, and $\int_{x_0}^x (x - t) f''(t) dt$ is the 1st order remainder. error/diff. to linear approx.

We can continue to get a higher order Taylor expansion using integration by parts by setting $v = f''(t)$ and $u' = (x - t)$ which would yield:

$$\begin{aligned} f(x) = & f(x_0) + f'(x_0)(x - x_0) \\ & + f''(x_0) \cdot \frac{(x - x_0)^2}{2} \\ & + \int_{x_0}^x (x - t) f'''(t) \cdot \frac{(x - t)^2}{2} dt \end{aligned} \quad (179)$$

Theorem: if $f(x)$ differentiable, then $\frac{d^n(f(x))}{dx^n}$ exists for all n . Hence the n -th order of the Taylor expansion would be:

$$f(x) = f(x_0) + \frac{f'(x_0)(x-x_0)}{1!} + \frac{f''(x_0)(x-x_0)^2}{2!} \dots + f^{(n)}(x_0) \frac{(x-x_0)^n}{n!} + \int_{x_0}^x (x-t) f^{(n+1)}(t) \cdot \frac{(x-t)^n}{n!} dt \quad (180)$$

Example:

Find the 2nd order Taylor expansion at $x = 0$ of $f(x) = x + e^{-2x}$ Use that to approximate the value $f(0.075) = 0.075 + e^{-0.15}$ and estimate the error.

$$\begin{aligned} f'(x) &= (1 - 2e^{2x}), f''(x) = 4e^{-2x} \\ f(x) &= f(0) + f'(0)(x-0) \\ &\quad + \frac{f''(0)(x-0)^2}{2} \\ &\quad + \int_0^x f'''(t) \frac{(x-t)^2}{2} dt \\ f(0.075) &\approx 1 - (0.075) + 2(0.075)^2 \end{aligned} \quad (181)$$

Error given by the remainder $R(x)$, a bound for the error

$$\text{error} \leq |R(x)| \leq \left| \int_0^x -8e^{2t} \cdot \frac{(x-t)^2}{2} dt \right| \quad (182)$$

$$\leq 4x^2(x-0) = 4x^3 = (4 \cdot (0.075)^3) \quad (183)$$

For 2 dimension, the second order expansion of $F(x, y)$ at (x_0, y_0) is:

$$\begin{aligned} F(x, y) &= F(x_0, y_0) \\ &\quad + \frac{1}{1!} \left[\frac{\sigma F}{\sigma x}(x_0, y_0)(x-x_0) + \frac{\sigma F}{\sigma y}(x_0, y_0)(y-y_0) \right] \\ &\quad + \frac{1}{2!} \left[\frac{\sigma^2 F}{\sigma x^2}(x_0, y_0)(x-x_0)^2 + \frac{\sigma^2 F}{\sigma y^2}(x_0, y_0)(y-y_0)^2 \right. \\ &\quad \left. + 2 \frac{\sigma^2 F}{\sigma x \sigma y} \right] + R_2(x, y) \end{aligned} \quad (184)$$

Example: Find the second Taylor expansion of $f(x, y) = \sin(x+y) + \cos(x-3y)$

at (0,0)

$$f_x = \cos(x+y) - \sin(x-3y) \quad (185)$$

$$f_y = \cos(x+y) + 3\sin(x-3y) \quad (186)$$

$$f_{xx} = -\sin(x+y) - \cos(x-3y) \quad (187)$$

$$f_{xy} = -\sin(x+y) + 3\cos(x-3y) \quad (188)$$

$$f_{yy} = -\sin(x+y) - 9\cos(x-3y) \quad (189)$$

$$f(x,y) = f(0,0) + f_x(0,0)x + f_y(0,0)y \quad (190)$$

$$+ \frac{1}{2} [f_{xx}x^2 + f_{yy}y^2 + 2f_{xy}xy] + R_2(x,y) \quad (191)$$

$$= 1 + x + y + \frac{1}{2}(-x^2 - 9y^2 + 6xy) + R_2(x,y) \quad (192)$$

Second Approximation, Taylor Polynomial.

For extreme values of $F(x,y)$: Recall that given $y = f(x)$, differentiable, n -th Taylor expansion at $x = x_0$

$$f(x) = f(x_0) + f'(x_0)(x - x_0) + f''(x_0) \cdot \frac{(x - x_0)^2}{2} \dots \quad (193)$$

$$+ f^{(n)}(x_0) \frac{(x - x_0)^n}{n!} \text{ Taylor polynomial} \quad (194)$$

$$+ \int_{x_0}^x f^{(n+1)}(t) \frac{(x - t)^n}{n!} dt \text{ } R_n(x) \text{ remainder} \quad (195)$$

Taylor Expansion for $F(x,y)$ at (x_0, y_0)

$$F(x,y) = F(x_0, y_0) + F_x(x_0, y_0)x + F_y(x_0, y_0)y \quad (196)$$

$$+ F_{xx}(x_0, y_0) \frac{x^2}{2} + F_{xy}(x_0, y_0) \frac{xy}{2} \quad (197)$$

$$+ F_{yx}(x_0, y_0) \frac{yx}{2} + F_{yy}(x_0, y_0) \frac{y^2}{2} \quad (198)$$

$$+ R_2(x,y) \quad (199)$$

Recall: for $y = f(x)$, critical points, $f'(x_0) = 0$ are candidates for local max and local min.

1. $f''(x_0) > 0$ local min
2. $f''(x_0) < 0$ local max
3. $f''(x_0) = 0$ inconclusive

if x_0 is critical point, first derivative term $f'(x_0) = 0$ removed from polynomial. The two variable are $F(x,y)$. Definition $F(x,y)$ differentiable, then say (x_0, y_0) is a critical point if $\nabla F(x_0, y_0) = 0$.

Recall equation of tangent plane

$$z = F(x_0, y_0) + F_x(x_0, y_0)(x - x_0) + F_y(y - y_0) \quad (200)$$

Tangent plane is horizontal. Definition: for $F(x, y)$, define the Hessian matrix.

$$\begin{pmatrix} F_{xx} & F_{xy} \\ F_{yx} & F_{yy} \end{pmatrix} (x_0, y_0) \quad (201)$$

Define the discriminant, (x_0, y_0) is critical point.

$$D(x_0, y_0) = \det \begin{pmatrix} F_{xx}(x_0, y_0) & F_{xy}(x_0, y_0) \\ F_{yx}(x_0, y_0) & F_{yy}(x_0, y_0) \end{pmatrix} (x_0, y_0) = F_{xx}F_{yy} - F_{xy}F_{yx} \quad (202)$$

Theorem (2nd derivative test): Suppose (x_0, y_0) is a critical point, then

1. If $D(x_0, y_0) > 0$ and $F_{xx} > 0$ or $(F_{yy} > 0)$, then (x_0, y_0) is a local minimum
2. If $D(x_0, y_0) > 0$ and $F_{xx} < 0$ or $(F_{yy} < 0)$, then (x_0, y_0) is a local maximum
3. if $D(x_0, y_0) < 0$, then (x_0, y_0) is a saddle point.
4. if $D(x_0, y_0) = 0$, then it is inconclusive.

Find and classify all critical points of $F(x, y) = x^3 - 3x^2 - 3y^2 + 3xy^2$

$$F_x = 3x^2 - 6x + 3y^2 = 0 \quad (203)$$

$$F_y = -6y + 6xy = 0 \quad (204)$$

$$\Rightarrow y = 0 \text{ or } x = 1 \text{ giving two critical point} \quad (205)$$

$$y = 0, \Rightarrow (0, 0), (2, 0) \quad (206)$$

$$x = 1, \Rightarrow (1, 1), (1, -1) \quad (207)$$

Use Second derivative to test for the four critical points

$$D(x, y) = F_{xx}F_{yy} - F_{xy}^2 \quad (208)$$

$$D(1, 1) = \Rightarrow \text{ is a saddle point} \quad (209)$$

$$D(1, -1) = \Rightarrow \text{ is a saddle point} \quad (210)$$

$$D(0, 0) = \Rightarrow \text{ is local max} \quad (211)$$

$$D(2, 0) = \Rightarrow \text{ is local min} \quad (212)$$

Why is this true?

From the second Taylor expansion, at a critical point, the dominant part:

$$F_{xx}(x_0, y_0)(x - x_0)^2 + 2F_{xy}(x_0, y_0)(x - x_0)(y - y_0) + F_{yy}(y - y_0)^2 \quad (213)$$

Homogeneous degree 2 polynomial in 2 variables:

$$P(x, y) = ax^2 + 2Cxy + by^2 \quad (214)$$

$$= a(x^2 + \frac{2c}{a}xy + \frac{b}{a}y^2) \quad (215)$$

$$= a((x - \frac{c}{a}y)^2 + \frac{ab - c^2}{a^2}y^2) \quad (216)$$

$$= a((x - \frac{c}{a}y)^2 + \frac{ab - c^2}{a^2}y^2) \quad (217)$$

$$\Delta = ab - c^2 > 0, \text{ then } \begin{cases} x' = x - \frac{c}{a}y \\ y' = x - \frac{\sqrt{ab-c^2}}{a}y \end{cases} \quad (218)$$

$$= a((x')^2 + (y')^2) \quad (219)$$

$$\Rightarrow a > 0 \text{ local min, } a < 0 \text{ local max} \quad (220)$$

$$\text{If } \Delta < 0 \begin{cases} x' = x - \frac{c}{a}y \\ y' = x - \frac{\sqrt{c^2-ab}}{a}y \end{cases} \quad (221)$$

$$\text{then } P(x, y) = P(x', y') = a((x')^2 - (y')^2) \quad (222)$$

$$a > 0 \text{ saddle point} \quad (223)$$

Apply this to $F_{xx}(x_0, y_0)(x - x_0)^2 + 2F_{xy}(x_0, y_0)(x - x_0)(y - y_0) + F_{yy}(y - y_0)^2$

$$a = F_{xx}(x_0, y_0)(x - x_0)^2 \quad (224)$$

$$b = F_{yy}(x_0, y_0)(x - x_0)^2 \quad (225)$$

$$2c = 2F_{xy}(x_0, y_0)(x - x_0)^2 \quad (226)$$

$$ab - c^2 = D(x, y) \quad (227)$$

Global Extreme Values for F(x,y).

Recall local max/min:

1. Find critical points $\nabla F(x_0, y_0) = \vec{0}$
2. Use second derivatives to test the critical points.

For Global Extreme values:

Example:

1) Find global min and max of $f(x, y) = x^3 - 3xy + 3y^2$ defined on the rectangle $D = \{(x, y) | 0 \leq x \leq 1, 0 \leq y \leq 2\}$

First, find local critical point

$$f_x = 3x^2 - 3y = 0 \quad (228)$$

$$f_y = -3x + 6y = 0 \quad (229)$$

$$\Rightarrow x = 2y \quad (230)$$

$$\text{plug in to } f_x : 0 = 3(4y^2 - y) \quad (231)$$

$$0 = y(4y - 1) \quad (232)$$

$$\Rightarrow y = 0 \text{ or } y = \frac{1}{4} \quad (233)$$

$$\therefore (0, 0) \text{ \& } (0.5, 0.25) \text{ two critical point} \quad (234)$$

$$\text{Hessian Matrix } D(x, y) = \det \begin{bmatrix} f_{xx} & f_{xy} \\ f_{xy} & f_{yy} \end{bmatrix} \quad (235)$$

$$= f_{xx} - f_{xy}^2 = 36x - 9 = 9(4x - 1) \quad (236)$$

$$\Rightarrow D(0, 0) = -9 < 0, \text{ saddle, neither max/min} \quad (237)$$

$$\Rightarrow D(0.5, 0.25) = 9 > 0, \quad f_{xx}(0.5, 0.25) = 3 > 0 \quad (238)$$

$$(239)$$

Need to compare with boundary values.

$$\text{Side 1: } \{x = 0, 0 \leq y \leq 2\} \quad (240)$$

$$f(x, y) = x^3 - 3xy + 3y^2 |_{x=0} \quad (241)$$

$$f(0, y) = 3y^2 \Rightarrow \max 12 \min 0 \quad (242)$$

$$\text{Side 2: } \{y = 0, 0 \leq x \leq 1\} \quad (243)$$

$$f(x, 0) = x^3, \max 1 \min 0 \quad (244)$$

$$\text{Side 3: } \{x = 1, 0 \leq y \leq 2\} \quad (245)$$

$$f(1, y) = 1 - 3y + 3y^2, \max 7 \min 0.25 \quad (246)$$

$$\text{Side 4: } \{y = 2, 0 \leq x \leq 1\} \quad (247)$$

$$f(x, 2) = x^3 - 6x + 12, \max 12 \min 7 \quad (248)$$

Hence global max 12 at (0,2) and -1/16 at (0.5,0.25)

2) Find for the function $f(x, y) = 2x^2 + 2y^2 - x + y$ on disk $D = \{(x, y) : x^2 + y^2 \leq$

1} its global Max and min/min

$$f_x = 4x - 1, f_y = 4y + 1, \quad (249)$$

$$f_x = 0 \text{ and } f_y = 0 \rightarrow (x = 1/4, y = -1/4) \quad (250)$$

$$f_{xx} = 4, f_{yy} = 4, f_{xy} = 0 \quad (251)$$

$$D(x, y) = 16 - 0^2 = 16 > 0, f_{xx} = 4 > 0 \quad (252)$$

$$\text{see } (x = 1/4, y = -1/4) \text{ is local min} \quad (253)$$

Parameterize the bounding (unit circle)

$$\vec{c}(t) = (\cos(t), \sin(t)), t \in [0, 2\pi] \quad (254)$$

$$\text{plugin } f(\cos(t), \sin(t)) = 2 - \cos(t) + \sin(t), t \in [0, 2\pi] \quad (255)$$

$$f'(t) = \sin(t) + \cos(t) = 0 \quad (256)$$

$$-1 = \tan(t) \quad (257)$$

$$\therefore \text{critical points } t = \frac{3\pi}{4}, t = \frac{7\pi}{4} \quad (258)$$

$$f(t) = 2 - \cos(t) + \sin(t) \text{ at } 0 \& 2\pi = 1 \& 1 \quad (259)$$

$$f\left(\frac{3\pi}{4}\right) = 2 - \cos\left(\frac{3\pi}{4}\right) + \sin\left(\frac{3\pi}{4}\right) = 2 + \sqrt{2} \quad (260)$$

$$f\left(\frac{7\pi}{4}\right) = 2 - \sqrt{2} \text{ min on boundary} \quad (261)$$

Recall have local min at (0.25, -0.25)

Global max at $(-\frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2})$, $2 + \sqrt{2}$

Global max at $(\frac{1}{4}, -\frac{1}{4})$, $-\frac{1}{4}$

Optimization with constraint — Lagrangian multiplier.

Given $f(x, y)$, find max/min with a constraint $g(x, y) = k$, $k = \text{a fixed number}$.

The langrangian multiplier method would be:

$$\text{Step 1 } \begin{cases} \nabla f = \lambda \nabla g, \lambda = \text{constant} \\ g(x, y) = k \end{cases} \quad (262)$$

solve this system of equation to get (x, y)

Step 2, plug in (x, y) into $f(x, y)$

Example: $f(x, y) = 4xy$, find max/min with condition $x^2 + y^2 = 4$

$$g(x, y) = x^2 + y^2, g(x, y) = 4 \quad (263)$$

$$f \& g = \begin{cases} \nabla f = & (2y, 2x) = \lambda \cdot \nabla g = (2x, 2y) \\ g(x, y) = & 4 = x^2 + y^2 \end{cases} \quad (264)$$

$$\therefore 2y = \lambda 2x \rightarrow \lambda = \frac{y}{x} \quad (265)$$

$$2x = \lambda 2y \rightarrow \lambda = \frac{x}{y} \quad (266)$$

$$\therefore x^2 = y^2 \& g(x, y) : x^2 + y^2 = 4, x^2 = y^2 = 2 \quad (267)$$

$$x, y = \pm \sqrt{2}, 4 \text{ solutions, max } 8, \text{ min } -8 \quad (268)$$

Why does it work? at max $\nabla g // \nabla f$

2.14 Vector Field and Flow

Definition: A vector field $\vec{F} : \mathbb{R}^m \rightarrow \mathbb{R}^m$

1. Example 1: $\vec{F}(x, y) = (x, y), \mathbb{R}^2 \rightarrow \mathbb{R}^2$
2. Example 2: $\vec{F}(x, y) = (-y, x), \mathbb{R}^2 \rightarrow \mathbb{R}^2$
3. Example 3: $\vec{F}(x, y, z) = (x^3, xy, e^{xyz}), \mathbb{R}^3 \rightarrow \mathbb{R}^3$

Definition: (Flow/Integral Curve) Given $\vec{F} : \mathbb{R}^m \rightarrow \mathbb{R}^m$, a flow of \vec{F} is curve $\vec{C}(t) : \mathbb{R}^1 \rightarrow \mathbb{R}^m$ such that $\vec{C}'(t) = \vec{F}(\vec{C}(t))$

Example:

$$\vec{F}(x, y) = (-y, x) \quad (269)$$

Verify that the curve $\vec{C}(t) = (\cos(t), \sin(t))$, is a flow/integral curve of \vec{F} by checking if $\vec{C}'(t) = \vec{F}(\vec{C}(t))$

$$\vec{C}'(t) = (-\sin(t), \cos(t)) \quad (270)$$

$$\vec{C}'(t) \equiv \vec{F}(\vec{C}(t)) \quad (271)$$

Remark: Finding flows of a vector field is the focus of differential equations.

An important class of example of vector field is given by ∇F where $F : \mathbb{R}^m \rightarrow \mathbb{R}^1$, $\nabla F = (\frac{\sigma F}{\sigma x}, \frac{\sigma F}{\sigma y}, \frac{\sigma F}{\sigma z}) : \mathbb{R}^3 \rightarrow \mathbb{R}^3$

Looking at ∇F tells us something about F

1. ∇F is the direction along which F changes most dramatically.
2. $\nabla F \perp$ level sets ($F(x, y, z) = c$).

Definition (Divergence and Curl): Given $\vec{F} : \mathbb{R}^3 \rightarrow \mathbb{R}^3$, $\vec{F} = (F_1(x, y, z), F_2(x, y, z), F_3(x, y, z))$. Define the divergence of \vec{F}

1. $\text{div } \vec{F} = (\frac{\sigma F}{\sigma x}, \frac{\sigma F}{\sigma y}, \frac{\sigma F}{\sigma z})$
2. $\text{div } \vec{F} = \mathbb{R}^3 \rightarrow \mathbb{R}^1$ a scalar function

Define the curl of \vec{F}

1. $\text{Curl } \vec{F} = \begin{bmatrix} \sigma/\sigma x \\ \sigma/\sigma y \\ \sigma/\sigma z \end{bmatrix} \times \begin{bmatrix} F_1 \\ F_2 \\ F_3 \end{bmatrix}$

2. $\text{div } \vec{F} = \mathbb{R}^3 \rightarrow \mathbb{R}^3$ a vector function

Easier way to memorize these 2 notations: ∇ is a vector of differential operators $\langle \frac{\sigma}{\sigma x}, \frac{\sigma}{\sigma y}, \frac{\sigma}{\sigma z} \rangle$

Hence ∇F means the gradient of F. So $\text{div } \vec{F}$ is the dot product between ∇ and \vec{F} : $\text{div } \vec{F} = \nabla \cdot \vec{F}$

Hence ∇F means the gradient of F. So $\text{curl } \vec{F}$ is the cross product between ∇ and \vec{F} : $\text{curl } \vec{F} = \nabla \times \vec{F}$

Similar for two dimension curves, except for the curl, we add a zero to the third dimension making the original function a function plot on a plane in the three dimensional plot. $\text{Curl } \vec{F} = \text{curl}(F_1, F_2, 0)$, and the result will be $(0, 0, \frac{\sigma F_2}{\sigma x} - \frac{\sigma F_1}{\sigma y})$, and the scalar curl for $\vec{F} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is defined as $\frac{\sigma F_2}{\sigma x} - \frac{\sigma F_1}{\sigma y}$, a scalar function.

Meanings of divergence and curl:

Divergence of \vec{F} at (x_0, y_0) is measuring the change of density at this point.
Change of volume of flows

$$= F_1(x + \Delta x, y)\Delta y - F_1(x, y)\Delta x \quad (272)$$

$$+ \frac{F_2(x, y + \Delta y)\Delta x - F_2(x, y)\Delta y}{\Delta x \Delta y} \quad (273)$$

$$\approx \frac{\sigma F_1}{\sigma x} + \frac{\sigma F_2}{\sigma y} \text{div } \vec{F} \quad (274)$$

Curl of $\vec{F} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$, scalar curl $= \frac{\sigma F_2}{\sigma x} - \frac{\sigma F_1}{\sigma y}$. is the measurement of the tendency of spin at a point. The Jacobian matrix $D\vec{F}$ capture the instant change of \vec{F} In particular, $D\vec{F}$ capture the infinitely minimal tendency of spin at the point.

A bit about Linear Algebra

If The Jacobian matrix $D\vec{F}$ is symmetric, i.e. $D\vec{F} = D\vec{F}^T$, i.e. $\frac{\sigma F_2}{\sigma x} = \frac{\sigma F_1}{\sigma y}$.

Then $D\vec{F}$ introduces no rotation. Curl measures how far away $D\vec{F}$ from being symmetric, the more asymmetric, the more $D\vec{F}$ rotates a basis.

Curl in third dimension:

1. $\text{Curl } \vec{F}$ = the axis of rotation of $D\vec{F}$.
2. $|\text{Curl } \vec{F}|$ = rate of rotation of $D\vec{F}$ the instant rate of rotation of \vec{F} at the point.

Relationship between ∇ , div , $curl$

$$\mathbb{R}^1 - (\nabla) \rightarrow \mathbb{R}^3 - (curl) \rightarrow \mathbb{R}^3 - (div) \rightarrow \mathbb{R}^1 \quad (275)$$

$\mathbb{R}^1 - (\nabla) \rightarrow \mathbb{R}^3$ are both tangent of functions.

$$curl(\nabla f) = \vec{0} \quad (276)$$

$$div(curl \vec{F}) = 0 \quad (277)$$

2.15 Integration along a path

Definition: A path (parametric curve) $\vec{c}: [a, b] \rightarrow \mathbb{R}^m$ is C' if $\vec{C}(t)$ exists and is continuous

Definition: A path $\vec{c}: [a, b] \rightarrow \mathbb{R}^m$ is piece-wise C' if can divide $[a, b]$ into finitely many sub-integrals. $[a_1 = a, a_2], [a_2, a_3] \dots, [a_{n-1}, a_n = b]$ such that \vec{C} on $[a_i, a_{i+1}]$ is C' .

Convention: From now on, we assume a path $\vec{C}(t)$ is always piece-wise C' .

Definition: If $\vec{C}(t)$ is a piece-wise C' path, we said that

1. $\vec{C}(t)$ is simple if it does not intersect itself or equivalently, $\vec{c}: [a, b] \rightarrow \mathbb{R}^m$ is an 1 to 1.
2. $\vec{C}(t)$ is simple closed path/curve if there exists $\vec{C}(a) = \vec{C}(b)$ and does not intersect itself except for the 2 end points.

Definition: for $\vec{C}: [a, b] \rightarrow \mathbb{R}^m$, $f(x_1 \dots x_m): \mathbb{R}^m \rightarrow \mathbb{R}^1$ then integral of f along \vec{C} is denoted by:

$$\int_C f ds = \int_a^b f(\vec{C}(t)) \cdot |\vec{C}'(t)| dt \quad (278)$$

Example: Finding the integral of $f(x, y, z) = xyz$ along the path $\vec{C}(t) = (-\sin(t), \sqrt{2}\cos(t), \sin(t))$, $t \in [0, \pi/2]$

$$\int_c f ds = \int_0^{\pi/2} f(\vec{C}(t)) \cdot |\vec{C}'(t)| dt \quad (279)$$

$$= \dots \quad (280)$$

$$= -2 \int_0^{\pi/2} \sin^2(t) \cos(t) dt \quad (281)$$

$$= -\frac{2}{3} \quad (282)$$

Theorem: Path integral of a function does not depend on the parametrization of the path.

Namely, $\vec{C}_1(t)$ and $\vec{C}_2(t)$ are 2 different parameter of the same path, $\int_{C_1} f ds = \int_{C_2} f ds$

The integration of vector field along a path is similar:

Definition: For a path $\vec{C}: [a, b] \rightarrow \mathbb{R}^m$ and a vector function $\vec{F}: \mathbb{R}^m \rightarrow \mathbb{R}^m$ the integral of \vec{F} along \vec{C} is

$$\int_{\vec{C}} \vec{F} d\vec{s} = \int_a^b \vec{F}(\vec{C}(t)) \cdot \vec{C}'(t) dt \quad (283)$$

This can be used in occasion like calculating the work done by a force field.

Example: $\vec{F}(x, y, z) = (-y, x, 1)$ force field, and there is a particle moving along a three dimension path $\vec{C}(t)$ and similarly, when the path is perpendicular, work done(hence the integral) would also be zero.

1) for $\vec{C}(t) = (t, t, t)$, $t \in [0, 1]$

$$\int_{\vec{C}} \vec{F} d\vec{s} = \int_0^1 (-t, t, 1) \cdot (1, 1, 1) dt \quad (284)$$

$$= 1 \text{ (J)} \quad (285)$$

2) for $\vec{C}(t) = (\cos(t), \sin(t), t)$, $t \in [0, 2\pi]$

$$\int_{\vec{C}} \vec{F} d\vec{s} = \int_0^{2\pi} \pi(-\sin(t), \cos(t), 1) \cdot (-\sin(t), \cos(t), 1) dt \quad (286)$$

$$= 4\pi \text{ (J)} \quad (287)$$

Properties of $\int_{\vec{C}} \vec{F} d\vec{s}$

$$\int_{\vec{C}} \vec{F} d\vec{s} = \int_a^b \vec{F}(\vec{C}(t)) \cdot \vec{C}'(t) dt \quad (288)$$

$$= \int_a^b |\vec{F}(\vec{C}(t))| \cdot |\vec{C}'(t)| \cos(\theta t) dt \quad (289)$$

The angle $\theta(t)$ is the angle formed by $\vec{F}(\vec{C}(t))$, and $\vec{C}'(t)$ is always < 90 degree. If $\vec{F}(\vec{C}(t)) \perp \vec{C}'(t)$

$$\int_{\vec{C}} \vec{F} d\vec{s} = 0 \quad (290)$$

If $\vec{C}_1(t)$ and $\vec{C}_2(t)$ are different parametrization of the same path with the same orientation, then

$$\int_{\vec{C}_1} \vec{F} d\vec{s} = \int_{\vec{C}_2} \vec{F} d\vec{s} \quad (291)$$

If opposite orientation

$$\int_{\vec{C}_1} \vec{F} d\vec{s} = - \int_{\vec{C}_2} \vec{F} d\vec{s} \quad (292)$$

Definition of Circulation: $\vec{C} : [a, b] \rightarrow \mathbb{R}^m$ a simple closed path, for a vector field F , the integration of this vector field along this simple closed path is called the circulation of \vec{F} along C .

For example, along a magnetic field, and such that C is a closed curve, the Ampere's law states that: $\int_C \vec{B} d\vec{s} = \mu I$ where I is the current.

Example: Take for example, the integration of the vector field $F_1(\vec{x}, y) = (-y, x)$ a simple closed path $\vec{C}(t) = (\cos(t), \sin(t))$, $t \in [0, 2\pi]$, the circulation would be

$$\vec{F}_1 = \int_{\vec{C}} \vec{F}_1 d\vec{s} \quad (293)$$

$$= \int_0^{2\pi} \begin{bmatrix} -\sin(t) \\ \cos(t) \end{bmatrix} \cdot \begin{bmatrix} -\sin(t) \\ \cos(t) \end{bmatrix} dt = 2\pi \quad (294)$$

2.16 Gradient Vector Field

Recall the two different kinds of integrals along a path:

$$\int_{\vec{C}} \vec{F} d\vec{s} = \int_a^b \vec{F}(\vec{C}(t)) \cdot \vec{C}'(t) dt, \quad \int_C f ds = \int_a^b f(\vec{C}(t)) |\vec{C}'(t)| dt$$

Definition: $\vec{F} : \mathbb{R}^m \rightarrow \mathbb{R}^m$ is called a gradient (conservative) vector field if there exists a function $f : \mathbb{R}^m \rightarrow \mathbb{R}$ such that $\vec{F} = \nabla f$ There are vector fields that are not gradient, recall that $Curl(\nabla f) = 0$:

$$Curl(\nabla f) = Curl\left(\frac{\sigma f}{\sigma x}, \frac{\sigma f}{\sigma y}\right) \quad (295)$$

$$= \frac{\sigma(\sigma f / \sigma y)}{\sigma x} - \frac{\sigma(\sigma f / \sigma x)}{\sigma y} \quad (296)$$

$$= 0 \quad (297)$$

Hence the curl test: if $\vec{F} : \mathbb{R}^m \rightarrow \mathbb{R}^m$, $\text{curl} \vec{F} \neq 0$, then \vec{F} cannot be a gradient curve.

Theorem:: Suppose $\vec{F} = \nabla f$, $\vec{C} : [a, b] \rightarrow \mathbb{R}^m$ a path, then

$$\int_{\vec{C}} \vec{F} d\vec{s} = \int_{\vec{C}} \nabla f d\vec{s} = f(\vec{C}(b)) - f(\vec{C}(a)) \quad (298)$$

Example: find the integration of a vector field $F = (2x, 2e^z, 2ye^z)$ along a curve abc $\vec{C}(a) = [0, 0, 0]$, $\vec{C}(b) = [0, -1, 1]$, $\vec{C}(c) = [1, -1, 0]$

$$\text{Observed that: } \vec{F} = \nabla(x^2 + 2ye^z) = \nabla f \quad (299)$$

Hence from theorem:

$$\int_{\vec{C}} \vec{F} d\vec{s} = f(1, -1, 0) - f(0, 0, 0) = -1 \quad (300)$$

Observations, for two different closed path with the same end points:

$$\int_{\vec{C}_1} \nabla f ds = \int_{\vec{C}_2} \nabla f ds \quad (301)$$

equivalently, for a simple closed curve:

$$\int_{\vec{C}} \vec{F} d\vec{s} = \int_{\vec{C}} \nabla f d\vec{s} = f(\vec{C}(b)) - f(\vec{C}(a)) = 0 \quad (302)$$

Theorem: Hence given a vector field \vec{F} , the following statements about it are equivalent:

1. $\vec{F} = \nabla f$
2. $\int_{\vec{C}_1} \nabla f ds = 0$
3. For any two paths c_1 c_2 with the same end points, $\int_{\vec{C}_1} \vec{F} ds = \int_{\vec{C}_2} \vec{F} ds$

The curl of $\nabla f = 0$ is a necessary condition for gradient vector field, but is this true such that when $\text{curl} \vec{F} = 0$, $\vec{F} = \nabla f$? no

Example: $\vec{F} = (\frac{-y}{x^2+y^2}, \frac{x}{x^2+y^2}, z)$ (its $\text{Curl} = 0$) defined on $u = \mathbb{R}^3$ (is not simple connected), but if we take a simple closed curve $\vec{C}(t) = (\cos(t), \sin(t), 0)$ $t \in [0, 2\pi]$, and $\int_{\vec{C}} \vec{F} d\vec{s} \neq 0$, hence not a gradient vector field by theorem.

Definition: Simply connected domains, n belongs to \mathbb{R}^n is said to be simply connected if any closed path can be deformed continuously to a point, u is simply connected. For example, a disk is simply connected and a ring is not as it cannot be shrunk to a simple point.

Theorem: if $\vec{F} : \mathbb{R}^m \rightarrow \mathbb{R}^m$ vector field defined on a simple connected domain and $\text{curl} \vec{F} = 0$, then \vec{F} is a gradient vector field.

Proof: supposed that $u = \text{Disk}$ and \vec{F} defined on u , $\text{curl} \vec{F} = 0$, then we claim there exists a function $f(x,y)$ defined on u such that $\nabla f = \vec{F}$

$$\vec{F} = (F_1(x, y), F_2(x, y)), (a, b) \in u \quad (303)$$

$$f(a, b) = \int_{\vec{C}_{a,b}} \vec{F} d\vec{s}, \text{ where } C_{a,b} \text{ is a line } (0, 0) - - - (a, b) \quad (304)$$

$$= \int_0^1 (F_1(ta, tb), F_2(ta, tb)) \cdot (a, b) dt \left\{ C_{a,b}(t) = (ta, tb) t \in [0, 1] \right. \quad (305)$$

$$= f(a, b) \quad (306)$$

$$(307)$$

Claim: $\nabla F = \vec{F}$, check for $\frac{\sigma f}{\sigma a} F_1$:

$$\frac{\sigma f(a, b)}{\sigma a} = \frac{\sigma}{\sigma a} \int_0^1 (F_1(ta, tb), F_2(ta, tb)) \cdot (a, b) dt \quad (308)$$

$$= \int_0^1 (F_1(ta, tb) + a \frac{\sigma F_1}{\sigma x} \cdot \frac{\sigma x}{\sigma a} + b \frac{\sigma F_2}{\sigma x} \cdot \frac{\sigma x}{\sigma a}) dt, \quad x = ta, \quad \frac{\sigma x}{\sigma a} = t \quad (309)$$

$$Curl \vec{F} = \frac{\sigma F_2}{\sigma x} - \frac{\sigma F_1}{\sigma y} = 0 \quad (310)$$

$$= \int_0^1 (F_1(ta, tb) + at \frac{\sigma F_1}{\sigma x} + bt \frac{\sigma F_1}{\sigma x}) dt \quad (311)$$

$$x = at, \quad \frac{\sigma x}{\sigma t} = a, \quad y = bt, \quad \frac{\sigma y}{\sigma t} = b \quad (312)$$

$$= \int_0^1 (F_1(ta, tb) + t \frac{\sigma F_1}{\sigma x} \cdot \frac{\sigma x}{\sigma t} + t \frac{\sigma F_1}{\sigma y} \cdot \frac{\sigma y}{\sigma t}) dt \quad (313)$$

$$= \int_0^1 (F_1(a, b) + t \frac{df}{dt}) dt \quad (314)$$

$$= \int_0^1 \frac{d}{dt} (tF_1(ta, tb)) dt \quad (315)$$

$$= (tF_1(ta, tb)) \quad (316)$$

$$\frac{\sigma f}{\sigma a} = F_1(a, b) \quad (317)$$

$$(318)$$

Rest can be proved similarly.

Next for arbitrary domain u, assume that $curl \vec{F} = 0$ and we want to show that the integral for any two path with the same end points $\int_{\vec{C}_1} \vec{F} d\vec{s} = \int_{\vec{C}_2} \vec{F} d\vec{s}$

Idea: deform C_1 to C_2 such that any 2 adjacent curves can be covered by a series of small disks. Because for each disk, we shown $\int F$ independent of paths.

Then

$$\int_{c_1} \vec{F} ds = \sum_i \int_{P_i} \vec{F} ds = \sum_i \int_{Q_i} \vec{F} ds = \int_{C_3} \vec{F} ds \quad (319)$$

Similarly:

$$\int_{c_1} \vec{F} ds = \int_{C_3} \vec{F} ds = \int_{C_n} \vec{F} ds = \int_{C_2} \vec{F} ds \quad (320)$$

so $\int \vec{F}$ is indenependent of choice of paths, by theorem 1, \vec{F} is a gradient vector field.

2.17 Double Integrals

Green's Theorem: the circulation of \vec{F} along a closed curve = Double integral of $\text{curl}(\vec{F})$

$D \subseteq \mathbb{R}^2$ is a domain, we say D is closed if it contains its boundary in D . If there exists a disk $B(r)$ of radius r such that $D \subset B(r)$.

Definition: $f(x,y)$ is continuous function defined on a bounded and closed domain D , the double integral denoted by $vol = \iint_D f(x,y) dA$ is supposed to represent the volume of the solid bounded by the function. The idea is to use the Riemann sum, $R_n = \sum_{i=1}^n f(x_i, y_i) \Delta x \cdot \Delta y$, as $\lim_{n \rightarrow \infty} R_n$. We can compute these integrals as normal integrals one by one.

Example:

1) For a nice domain $D = [a, b] \times [c, d]$, $\iint_R 6x^2 y dA = ?$ By Fubini's theorem:

$$\iint_R 6x^2 y dA = \int_0^4 \left(\int_{-1}^1 6x^2 y dx \right) dy \quad (321)$$

$$= \int_0^4 \left. \frac{6yx^3}{3} \right|_{-1}^1 dy = \int_0^4 4y dy \quad (322)$$

$$= 2y^2 \Big|_0^4 = 32 \quad (323)$$

$$\text{Alternate method:} = \left(\int_{-1}^1 6x^2 dx \right) \cdot \left(\int_0^4 y dy \right) \quad (324)$$

$$= 2x^3 \Big|_{-1}^1 \cdot \frac{y^2}{2} \Big|_0^4 \quad (325)$$

$$= 32 \quad (326)$$

Remark: if $f(x, y) = g(x) \times h(y)$, $\iint f(x, y) = \left(\int g(x) dx \right) \cdot \left(\int h(y) dy \right)$

2) For a domain $x \in [a, b]$, $y \in [p(x), q(x)]$, evaluate $\iint_D e^{2x+y} dA$, D is the region bounded by the line $y = 2x$ and $y = x$, $x = 1$ and $x = 2$.

$$\int_1^2 \int_x^{2x} e^{2x+y} dy dx \quad (327)$$

$$= \int_1^2 (e^{2x} \cdot e^y \Big|_x^{2x}) dx \quad (328)$$

$$= \int_1^2 e^{4x} - e^{3x} dx \quad (329)$$

$$= \frac{e^{4x}}{4} - \frac{e^{3x}}{3} \Big|_1^2 \quad (330)$$

3) Type 2 elementary region $y \in [c, d]$, $x \in [p(y), q(y)]$, evaluate $\iint_D 2y dA$, D is the region bounded by the line $y = x - 6$ and $y^2 = x$. and by finding

boundaries, $y \in [-2, 3]$, $x \in [y^2, y + 6]$

$$\int_{-2}^3 \int_{y^2}^{y+6} 2y dx dy \quad (331)$$

$$= \int_{-2}^3 2y(y + 6 - y^2) dy \quad (332)$$

$$= -\frac{y^4}{2} + \frac{2y^3}{3} + 6y^2 \Big|_{-2}^3 \quad (333)$$

$$= \dots \quad (334)$$

4) We can change the order of integration by Fubini's theorem, for example, evaluate $\int_0^1 \int_x^{\sqrt[3]{x}} e^{x/y} dy dx$, we don't know the antiderivative of $e^{x/y}$ with respect to y , however, we can change the order

$$\int_0^1 \int_x^{\sqrt[3]{x}} e^{x/y} dy dx = \int_0^1 \int_{y^3}^y e^{x/y} dx dy \quad (335)$$

$$= \int_0^1 y e^{x/y} \Big|_{y^3}^y dy \quad (336)$$

$$= \int_0^1 y e^{x/y} \Big|_{y^3}^y dy \quad (337)$$

$$= \dots \quad (338)$$

$$= \frac{1}{2} \quad (339)$$

Some properties of double integrals.

1. Linearity

2. $D = D_1 \sqcup D_2$, then $\iint_D f dA = \iint_{D_1} f dA + \iint_{D_2} f dA$

3. $m \leq f(x, y) \leq M$, then $m \cdot \text{Area}(D) \leq \iint_D f(x, y) dA \leq M \cdot \text{Area}(D)$

Change of Variables(substitution) in two dimension, $(x, y) \in \mathbb{R}^2 \rightarrow (u, v) \in \mathbb{R}^2$, $(T(x, y) = (u(x, y), v(x, y)))$:

$$\iint_D f(x, y) dx dy \quad (340)$$

$$= \iint_{T(D)} f(x(u, v), y(u, v)) \left| \begin{array}{cc} \frac{\sigma x}{\sigma u} & \frac{\sigma x}{\sigma v} \\ \frac{\sigma y}{\sigma u} & \frac{\sigma y}{\sigma v} \end{array} \right| du dv \quad (341)$$

Example: Find $\iint_D (3x + 6y)^2 dA$, where D is the region enclosed by the four

lines $x \pm 2y = \pm 2$. By substitution, $u = x+2y$, $v = x-2y$.

$$\iint_D (3x + 6y)^2 dA \quad (342)$$

$$= \iint_{T(D)} \left(3\frac{u+v}{2} + 6\frac{u-v}{4} \right)^2 \left| \begin{array}{cc} \frac{1}{2} & \frac{1}{2} \\ \frac{1}{4} & -\frac{1}{4} \end{array} \right| dudv \quad (343)$$

$$= \int_{-2}^2 \int_{-2}^2 \frac{(3u)^2}{4} dudv \quad (344)$$

$$= 48 \quad (345)$$

Next, A special case of change of variables: by polar coordinate, $x = r \cos(\theta)$, $y = r \sin(\theta)$, $r = \sqrt{x^2 + y^2}$, $\tan(\theta) = \frac{y}{x}$

Example: Find the area of the unit disk D:

1) $f = 1$

$$Area(D) = \iint_D 1 dA \quad (346)$$

$$= \int_{-1}^1 \int_{-\sqrt{1-x^2}}^{\sqrt{1-x^2}} 1 dy dx \quad (347)$$

$$= \int_0^{2\pi} \int_0^1 1 \left| \begin{array}{cc} \frac{\sigma x}{\sigma u} & \frac{\sigma x}{\sigma v} \\ \frac{\sigma y}{\sigma u} & \frac{\sigma y}{\sigma v} \end{array} \right| dr d\theta \quad (348)$$

$$= \int_0^{2\pi} \int_0^1 1 \times r dr d\theta \quad (349)$$

$$= \dots \quad (350)$$

$$= \pi \quad (351)$$

In fact, $dx dy = r dr d\theta$
2) $e^{x^2+y^2}$

$$Area(D) = \iint_D e^{x^2+y^2} dA \quad (352)$$

$$= \int_0^{2\pi} \int_0^1 e^{r^2} r dr d\theta \quad (353)$$

$$= \frac{1}{2} e^{r^2} \Big|_0^1 (2\pi - 0) \quad (354)$$

$$= (e - 1)\pi \quad (355)$$