

Calibrating gyrochronology using Galactic kinematics

RUTH ANGUS,^{1,2,3} YUXI (LUCY) LU,^{3,1} DAN FOREMAN-MACKEY,² ADRIAN M. PRICE-WHELAN,² JASON CURTIS,¹ AND
EMILY CUNNINGHAM²

¹*Department of Astrophysics, American Museum of Natural History, 200 Central Park West, Manhattan, NY, USA*

²*Center for Computational Astrophysics, Flatiron Institute, 162 5th Avenue, Manhattan, NY, USA*

³*Department of Astronomy, Columbia University, Manhattan, NY, USA*

ABSTRACT

Gyrochronology, the method of inferring the age of a star from its rotation period, could provide ages for billions of stars over the coming decade of time-domain astronomy. However, the gyrochronology relations remain poorly calibrated due to a lack of precise ages for old, cool main-sequence stars. Now however, with proper motion measurements from Gaia, Galactic kinematics can be used as an age proxy, and the magnetic and rotational evolution of stars can be examined in detail. We demonstrate that kinematic ages, inferred from the velocity dispersions of groups of stars, beautifully illustrate the time and mass-dependence of the gyrochronology relations. We use the kinematic ages of field stars, plus benchmark clusters and asteroseismic stars, to calibrate a new empirical Gaussian process gyrochronology relation, that fully captures the complex rotational evolution of cool dwarfs over a range of masses and ages. We use cross validation to demonstrate that this relation accurately predicts ages for FGKM dwarfs.

Keywords: Stellar Rotation — Stellar Evolution — Stellar Activity — Stellar Magnetic Fields — Low Mass Stars — Solar Analogs — Milky Way Dynamics

1. INTRODUCTION

Low mass dwarfs are the most common stars in the Milky Way, and their ages could reveal the evolution of planetary systems and stellar populations in the Galaxy. However, the ages of GKM stars are difficult to measure because their luminosities and temperatures evolve slowly on the main sequence. Fortunately, rotation-dating, or ‘gyrochronology’ provides a promising means to measure precise ages for these cool dwarfs. The rotation periods of these stars evolve relatively rapidly, and a fully calibrated gyrochronology model that captures the time and mass-dependence of stellar spin down could provide ages that are precise to within 20% for millions of Milky Way stars in the time-domain era (Epstein & Pinsonneault 2014; Najita et al. 2016; Angus et al. 2019).

With the thousands of new photometric rotation period measurements provided by specialized ground and space-based missions (particularly *Kepler/K2* and *TESS* Borucki et al. 2010; Howell et al. 2014; Ricker et al. 2015), we are making progress towards the ultimate goal for rotation-dating: a fully calibrated gyrochronology relation, applicable to GKM main-sequence stars of all ages. A lack of low-mass and old calibration stars has previously limited the mass and age coverage of gyrochronology relations, which can only be calibrated using stars with precise age and rotation period measurements. Historically, the calibration sample has been limited to open clusters and asteroseismic stars because cluster members and Solar-like oscillators can be precisely dated via main-sequence turn off and asteroseismology. Open clusters provide good mass coverage for young stars: rotation periods have been measured for F through mid M dwarfs for stars in clusters with precisely measured ages up to around 700 Myr. Asteroseismic stars provide reasonable age coverage for hot stars: seismic masses and photometric surface rotation periods have been measured for F, G and early K dwarfs for stars as old as 10 Gyr. However, neither asteroseismology nor cluster analysis can provide rotation periods and ages for old, late K and M dwarfs. In addition, cluster and asteroseismic stars generally provide sparse coverage of the rotation period-effective temperature plane, and cannot reveal the detailed evolution of stellar rotation rates. As a result, most empirical gyrochronology relations are only reliable for G dwarfs up to Solar age, K dwarfs up to 2-3 Gyr, and early M dwarfs up to < 1 Gyr. For this reason, the rotational evolution of cool dwarfs is not well understood. However, as we showed in ?, *kinematic* ages can be used to turn field stars, observed by *Kepler*, into a calibration sample that provides good mass and age coverage. Although the *Kepler* sample does not include late

M dwarfs, it can still be used to extend gyrochronology relations to much older ages for late K and early M dwarfs. By adding the ages and rotation periods of thousands of field stars to the open cluster and asteroseismic calibration sample, we can calibrate a gyrochronology relation that is applicable to FGK and early M dwarfs between the ages of ~ 500 Myr and 8 Gyr.

1.1. *Using kinematics as an age proxy*

In ? we demonstrated that kinematics can be used to explore the evolution of stellar rotation. In this paper we take the next step and use kinematics to calibrate a new gyrochronology relation. This paper is laid out as follows. In section ?? we describe the calibration of a new AVR, and how we used it to calculate ages for over 6000 stars with measured rotation periods. In this section we also compare these kinematic ages with literature age measurements. In section ?? we describe how we use these kinematic ages to calibrate a new empirical gyrochronology relation.

2. METHOD

2.1. Data

This study focuses on stellar rotation in the original *Kepler* field, partly because *Kepler* provides the largest sample of published rotation periods, and partly because its low Galactic latitude allows us to marginalize over missing RV measurements to approximate v_z . We combined three rotation period catalogs constructed from original *Kepler* data: McQuillan et al. (2014), ? and García et al. (2014).

2.2. Inferring 3D velocities (marginalizing over missing RV measurements)

It has been demonstrated that the dispersion in vertical velocity, v_z for a group of stars increases with the age of that group (citations). However, velocities in Galactocentric coordinates, v_x , v_y and v_z , can only be calculated with full 6-D position and velocity information, *i.e.* proper motions, position and radial velocity. In ? we introduced the idea that kinematic ages could be used to calibrate gyrochronology and showed, in the appendix of that paper, that velocity, v_b in the Galactic frame, which can be calculated without an RV measurement, can be used as an approximation to v_z for *Kepler* stars. This is because the *Kepler* field of view lies at relatively low Galactic latitudes, ($\sim 5 - 20^\circ$), so the z -direction is similar to the b -direction for *Kepler* stars. However, v_b is only a close approximation to v_z at extremely low latitudes, and even in the *Kepler* field, kinematic ages calculated with v_b instead of v_z are systematically larger because of extra noise introduced by the imperfect translation between v_b and v_z . In order to calculate accurate vertical velocities and therefore ages, the appropriate approach is to *infer* v_z by marginalizing over missing RV measurements.

Three-dimensional velocities in galactocentric coordinates: v_x , v_y , and v_z can only be directly computed via a transformation from 3D velocities in another coordinate system, like the equatorial coordinates provided by *Gaia*: μ_α , μ_δ , and RV. For stars with no measured RV in *Gaia* DR2, v_x , v_y , and v_z can still be inferred from positions and proper motions alone, by marginalizing over missing RV measurements. For each star in our sample, we inferred v_x , v_y , and v_z from the 3D positions and proper motions provided in the *Gaia* DR2 catalog (Brown et al. 2011). We also simultaneously inferred distance, instead of using $1/\pi$, to model velocities (?).

Using Bayes rule, the posterior probability of the parameters given the data can be written:

$$p(v_{xyz}, D | \mu_\alpha, \mu_\delta, \alpha, \delta, \pi) = p(\mu_\alpha, \mu_\delta, \alpha, \delta, \pi | v_{xyz}, D) p(v_{xyz}) p(D), \quad (1)$$

where D is distance, α is Right Ascension (RA), δ is declination (dec), π is parallax, μ_α is proper motion in RA, and μ_δ is proper motion in dec. The prior over $\log(\text{distance})$ and velocities was a multivariate Gaussian with mean and covariance determined from the distance and velocity distributions of *Kepler* targets with RV measurements.

The posterior PDF was explored using **emcee** (?), an affine-invariant, ensemble MCMC sampler.

Initialization.

We found that 10,000 samples with 16 walkers was sufficient to calculate a converged autocorrelation time, and produce 150-500 independent samples per parameter.

Over 3000 stars in the McQuillan et al. (2014) sample do have RV measurements and provide an opportunity to test this method of inferring velocities. Figure ?? shows the velocities of these 3000 stars, calculated using RV measurements, compared with their inferred velocities.

2.3. Calculating velocity dispersions


A kinematic age can be calculated from the velocity dispersion (*e.g.* the standard deviation or Median Absolute Deviation, MAD, of velocities) of a group of stars. Many empirical Age-Velocity dispersion Relations (AVRs) have been calibrated using stars with measured proper motions, radial velocities, and ages (*e.g.* ??). The major assumption underlying kinematic ages is that the stars used to calculate a velocity dispersion are *all the same age*. So, in order to calculate kinematic ages from velocity dispersions for *Kepler* stars, it is necessary to group them by age.

2.4. Converting velocity dispersion to age with an AVR

2.5. Comparing kinematic ages with asteroseismic and cluster ages

2.6. A Gaussian process gyrochronology relation

Figure 1. Vertical velocities calculated with full 6D information vs vertical velocities inferred without RV, for all 3000 [McQuillan et al. \(2014\)](#) stars with *Gaia* RV measurements.



3. RESULTS

4. DISCUSSION

5. CONCLUSION

This work was partly developed at the 2019 KITP conference ‘Better stars, better planets’. Parts of this project are based on ideas explored at the Gaia sprints at the Flatiron Institute in New York City, 2016 and MPIA, Heidelberg, 2017. This work made use of the *gaia-kepler.fun* crossmatch database created by Megan Bedell.

Some of the data presented in this paper were obtained from the Mikulski Archive for Space Telescopes (MAST). STScI is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS5-26555. Support for MAST for non-HST data is provided by the NASA Office of Space Science via grant NNX09AF08G and by other grants and contracts. This paper includes data collected by the Kepler mission. Funding for the *Kepler* mission is provided by the NASA Science Mission directorate.

This work has made use of data from the European Space Agency (ESA) mission *Gaia* (<https://www.cosmos.esa.int/gaia>), processed by the *Gaia* Data Processing and Analysis Consortium (DPAC, <https://www.cosmos.esa.int/web/gaia/dpac/consortium>). Funding for the DPAC has been provided by national institutions, in particular the institutions participating in the *Gaia* Multilateral Agreement.

REFERENCES

- Angus, R., Morton, T. D., & Foreman-Mackey *et al*, D. 2019, AJ, 158, 173, doi: [10.3847/1538-3881/ab3c53](https://doi.org/10.3847/1538-3881/ab3c53)
- Borucki, W. J., Koch, D., & Basri *et al*, G. 2010, Science, 327, 977, doi: [10.1126/science.1185402](https://doi.org/10.1126/science.1185402)
- Brown, T. M., Latham, D. W., & Everett *et al*, M. E. 2011, AJ, 142, 112, doi: [10.1088/0004-6256/142/4/112](https://doi.org/10.1088/0004-6256/142/4/112)
- Epstein, C. R., & Pinsonneault, M. H. 2014, ApJ, 780, 159, doi: [10.1088/0004-637X/780/2/159](https://doi.org/10.1088/0004-637X/780/2/159)
- García, R. A., Ceillier, T., & Salabert *et al*, D. 2014, A&A, 572, A34, doi: [10.1051/0004-6361/201423888](https://doi.org/10.1051/0004-6361/201423888)
- Howell, S. B., Sobeck, C., & Haas *et al*, M. 2014, PASP, 126, 398, doi: [10.1086/676406](https://doi.org/10.1086/676406)
- McQuillan, A., Mazeh, T., & Aigrain, S. 2014, ApJS, 211, 24, doi: [10.1088/0067-0049/211/2/24](https://doi.org/10.1088/0067-0049/211/2/24)
- Najita, J., Willman, B., Finkbeiner, D. P., et al. 2016, arXiv e-prints, arXiv:1610.01661. <https://arxiv.org/abs/1610.01661>
- Ricker, G. R., Winn, J. N., & Vanderspek *et al*, R. 2015, Journal of Astronomical Telescopes, Instruments, and Systems, 1, 014003, doi: [10.1117/1.JATIS.1.1.014003](https://doi.org/10.1117/1.JATIS.1.1.014003)