

Mapping landscape suitability for forest thinning to reduce evapotranspiration and enhance groundwater recharge in Arizona

Ryan E Lima^{a,*}, Neha Gupta^b, Travis Zalesky^b, Temuulen Tsagaan Sankey^a,
Abraham E Springer^a, Katherine Jacobs^b

^a*Northern Arizona University,*

^b*University of Arizona,*

Abstract

Literature on the relationship between forest thinning and water yield was used to develop suitability criteria to map where forest treatment is most likely to enhance groundwater recharge across the Coconino National Forest in Arizona. Recharge in the region is ephemeral and focused in periods of snowmelt and locations of enhanced permeability when soil moisture exceeds threshold levels. Our approach combines thematic maps of criteria such as average precipitation, snow dominance, slope, aspect, landscape morphology, forest basal area, canopy cover, lithology and hydrologic soil type into a GIS-Multi-Criteria Decision Making model. Pairwise comparisons were made between criteria, and Analytic Hierarchy Process was used as a weighting method.

Keywords: suitability mapping, Forest thinning, Water yield, groundwater recharge, GIS-MCDM, AHP

1. Introduction

Warming associated with anthropocentric climate change has increased the number of extreme hydroclimate events in the Colorado River Basin. Droughts, heatwaves, and floods have more than doubled in frequency since 2010 [1]. Since 2000, the Colorado River Basin has been in the midst of a historic drought [2, 3]. During that time, streamflow in the Colorado River has declined by 19% relative to the 1906-1999 average [4, 5]. Rapid population growth in the Southwest and Arizona, in particular, is increasing the demands on already strained water supplies in the State. Reductions in streamflow have increased

*Corresponding author

Email address: ryan.lima@nau.edu (Ryan E Lima)

reliance on groundwater pumping, while groundwater levels have declined for decades in many groundwater basins across Arizona [6].

Concurrently, the risk of catastrophic wildfires is increasing in Western forests—an emerging driver of runoff change that will increase the impact on the water supply [3]. Forest structure in Northern Arizona and New Mexico has changed significantly post-Euro-American settlement. Many forests are overstocked relative to pre-settlement conditions due to grazing, logging, and wildfire exclusion [7, 8]. These changes have increased the risk of catastrophic wildfire [9]. Rising temperatures and related droughts have contributed to extensive tree mortality from wildfire, disease, and insect infestation [10]. Warming temperatures have tripled the frequency and quadrupled the size of wildfires since 2000 [[11]].

Landscape-scale forest restoration efforts have been planned or implemented across much of Arizona. For example, the Four Forest Restoration Initiative (4FRI) includes plans for restoration across over 1 million hectares of Arizona’s forests [12]. The primary goal of restoration efforts is to reduce wildfire risk [9, 8]. However, numerous studies have linked forest treatments to increased water yields in semi-arid forests and have emphasized the role of forest restoration in improving hydrologic services and increasing water availability [13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24]. Forest treatments such as thinning and burning can significantly impact the hydrologic cycle of forests [25]. For example, forest thinning in Arizona has been associated with increased snow cover days [26, 27, 28], greater soil moisture [29, 30], and greater forest canopy moisture [31].

While the connection between forest treatment and water yield is well documented, the response of forests to treatments is complex and non-linear and differs across forest types, with treatment level, and along aspect and elevational gradients [25, 32, 17, 24, 33]. Regardless of the potential for increased water yield, the enhancement of groundwater recharge rarely, if ever, ranks among the primary motivations for forest treatment even among projects with the stated goal of improving watershed health [34, 35, 9, 8, 36]. This study examines forest restoration through the lens of groundwater recharge enhancement and aims to identify potential recharge zones. We map suitability for forest thinning with the goal of enhancing recharge. Suitability maps like these may complement (or supplement) existing frameworks for prioritizing landscape-scale forest management.

Suitability mapping, and particularly GIS-based Multi-criteria decision making (GIS-MCDM), is widely used to map potential recharge zones and areas suitable for Managed Aquifer Recharge (MAR), but to our knowledge, it has not yet been implemented to map incidental recharge, or recharge enhancement potential, from forest thinning [37, 38]. Pairwise comparisons were made between criteria including forest basal area, canopy cover, average precipitation, snow dominance, slope, aspect, landscape morphology, forest density, lithology, and hydrologic soil type, and the Analytic Hierarchy Process (AHP) was used as a weighting method.

1.1. Literature Reviewed

- How forest thinning may enhance recharge
 - Reduced canopy interception, increased snow retention
 - decreases ET - TEKI fort valley research
 - Increases Soil Moisture
 - Precipitation thresholds in thinning literature
- Caveats
 - Low Elevations - less water yield in low elevation forests in the Salt-Verde System [32]
 - Sunny Aspects - increased sublimation with decreased canopy cover [39]

2. Methods

We used the ArcGIS Pro Suitability Modeler within the Spatial Analysis toolbox to create a weighted suitability model. The first step involved screening out unsuitable areas including non-forested land covers, wilderness areas, and areas with maximum annual precipitation below 500 *mm*. After this initial screening process thematic maps were created using criteria known to affect water yield in thinned forests.

2.1. Study Area

The Coconino National Forest is located in Northern Arizona near the cities of Flagstaff and Sedona. It ranges in elevation from 790 *m* near the Verde River to 3,851 *m* at the summit of Humphreys Peak. It is located within the largest contiguous ponderosa pine (*Pinus ponderosa*) forest in North America. Ponderosa pine covers roughly 40% of the national forest an area of about 340,000 hectares, primarily between 2000 - 2400 *m* in elevation.

- The Coconino National Forest also spans the Mogollon Rim, which has been identified as an important recharge area for the regional groundwater aquifers
- Of the estimated 1,740,000 acre-feet of precipitation that falls on the Mogollon Rim annually, about 8 percent is estimated to recharge the regional aquifers [40].

2.2. Initial Suitability Screen

Areas were considered unsuitable for the following reasons:

1. Areas with Max annual precipitation 1991 - 2020 < 500mm [source: AORC Max Annual Precipitation 1991 - 2020]
2. All Landcover types other than 410 Deciduous Forest, 42 - Evergreen Forest, & 43 - Mixed Forest [source: NLCD Landcover 2021]
3. Forest Vegetation types that include attributes EVT_LF = ["Water", "Shrub", "Sparse", "Herb", "Developed", "Barren", "Agriculture"] and EVT_PHYS = ["Developed", "Riparian", "Agriculture"] and including keywords in EVT_NAME = ["Madrean", "Savanna"]. [Source: LF2023_240_EVT] Landfire
4. Wilderness Areas

2.3. Suitability Criteria

2.3.1. Topographic Relative Moisture Index

Topographic Relative Moisture Index (TRMI) incorporates several topographic parameters that influence moisture dynamics including slope gradient, aspect, relative elevation (or topographic position) and landscape convexity or concavity [41]. TRMI data comes from the Southwest Regional Gap Analysis (SWReGAP) where terrain features are separated into 10 categories ranked in terms of soil moisture. Del Campo and others [42] examined below ground hydrological processes in thinned semi-arid watersheds and found that sites with high antecedent soil moisture had the highest response, with drainage to deeper soil layers increasing by 50mm/year relative to control sites.

| Class Name | Description | Suitability Value |
|-------------------------------------|---|-------------------|
| Valley Flats | These areas are typically the lowest points in a landscape where water naturally accumulates, making them the moistest. | 10 |
| Very Moist Steep Slopes | Despite the slope, these areas (often with north-facing or shaded aspects in the Northern Hemisphere) retain moisture due to cool temperatures and reduced evaporation. | 9 |
| Toe Slopes | These are the lower parts of slopes where water tends to collect after flowing downhill, making them moist but slightly less so than valley flats. | 8 |
| Cool Aspect Scarps, Cliffs, Canyons | North-facing scarps and shaded canyons are cooler and retain moisture better than exposed slopes, particularly in arid regions. | 7 |

| Class Name | Description | Suitability Value |
|--|--|-------------------|
| Nearly Level Plateaus or Terraces | These flat areas may retain moderate amounts of moisture but are generally more exposed to sunlight and wind than valleys or steep moist slopes. | 6 |
| Gently Sloping Ridges | These ridges are not as steep, so water may infiltrate rather than run off completely, but they are still relatively dry due to elevation. | 5 |
| Moderately Moist Steep Slopes | These are typically slopes with intermediate aspects or conditions, retaining some moisture but less than moist slopes. | 4 |
| Moderately Dry Slopes | These areas have greater runoff and less water retention, often due to steeper gradients and/or sun exposure. | 3 |
| Very Dry Steep Slopes | Steep slopes with sun exposure (e.g., south-facing in the Northern Hemisphere) promote high runoff and evaporation, leaving them very dry. | 2 |
| Hot Aspect Scarps, Cliffs, and Canyons | South-facing cliffs and canyons with maximum sun exposure (in the Northern Hemisphere) are the driest due to high evaporation and limited water retention. | 1 |

2.3.2. Basal Area

We extracted basal area estimates from the TreeMap 2016 [43] conus dataset.

2.3.3. Snowfall Dominance

Ask Patrick Broxton how it was calculated.

2.3.4. Mean Annual Precipitation

Utilized AORC Retrospective forcing data for average water year (WY) precipitation 1991 - 2020

2.3.5. Subsurface Infiltration Capacity

We utilized the Global Hydrological Maps of Permeability and Porosity (GLHYMPS) to estimate subsurface infiltration capacity [44]

2.3.6. Soil Hydrologic Type

Data from gNATSGO was used to estimate the soil's ability to infiltrate water, we used the Hydrologic Soil Type.

2.3.7. Canopy Cover

Canopy Cover was obtained from the 2021 National Land Cover Dataset , which provides an estimate of Total Canopy Cover

2.4. Weighting

| Criteria | Weight (%) |
|----------------------------|------------|
| Subsurface Infiltration | 10.12 |
| Canopy Cover | 9.82 |
| Basal Area | 15.06 |
| Topographic Moisture Index | 29.48 |
| Mean annual Precipitation | 10.73 |
| Snowfall Fraction | 14.77 |

3. Results

We identified 30 forest patches of between 50 and 60 Hectares that we consider suitable for thinning for the purposes of enhancing groundwater recharge. The patches have mean suitability values ranging from 5.8 to 6.25




Figure 1: Figure x: Suitability Map for Thinning to Enhance Groundwater Recharge

4. Discussion

4.1. Acknowledgments

Phasellus interdum tincidunt ex, a euismod massa pulvinar at. Ut fringilla ut nisi nec volutpat. Morbi imperdiet congue tincidunt. Vivamus eget rutrum purus. Etiam et pretium justo. Donec et egestas sem. Donec molestie ex sit amet viverra egestas. Nullam justo nulla, fringilla at iaculis in, posuere non mauris. Ut eget imperdiet elit.

4.2. Open research

Phasellus interdum tincidunt ex, a euismod massa pulvinar at. Ut fringilla ut nisi nec volutpat. Morbi imperdiet congue tincidunt. Vivamus eget rutrum purus. Etiam et pretium justo. Donec et egestas sem. Donec molestie ex sit amet viverra egestas. Nullam justo nulla, fringilla at iaculis in, posuere non mauris. Ut eget imperdiet elit.

References

- [1] K. E. Bennett, C. Talsma, R. Boero, [Concurrent Changes in Extreme Hydroclimate Events in the Colorado River Basin](#), *Water* 13 (7) (2021) 978. doi:10.3390/w13070978.
URL <https://www.mdpi.com/2073-4441/13/7/978>
- [2] D. M. Meko, C. A. Woodhouse, A. G. Winitzky, [Tree-Ring Perspectives on the Colorado River: Looking Back and Moving Forward](#), *JAWRA Journal of the American Water Resources Association* 58 (5) (2022) 604–621. doi:10.1111/1752-1688.12989.
URL <https://onlinelibrary.wiley.com/doi/10.1111/1752-1688.12989>
- [3] A. P. Williams, B. I. Cook, J. E. Smerdon, [Rapid intensification of the emerging southwestern North American megadrought in 2020–2021](#), *Nature Climate Change* 12 (3) (2022) 232–234. doi:10.1038/s41558-022-01290-z.
URL <https://www.nature.com/articles/s41558-022-01290-z>
- [4] D. Hogan, J. D. Lundquist, [Recent Upper Colorado River Streamflow Declines Driven by Loss of Spring Precipitation](#), *Geophysical Research Letters* 51 (16) (2024) e2024GL109826. doi:10.1029/2024GL109826.
URL <https://agupubs.onlinelibrary.wiley.com/doi/10.1029/2024GL109826>
- [5] B. Udall, J. Overpeck, [The twenty-first century Colorado River hot drought and implications for the future](#), *Water Resources Research* 53 (3) (2017) 2404–2418. doi:10.1002/2016WR019638.
URL <https://agupubs.onlinelibrary.wiley.com/doi/10.1002/2016WR019638>
- [6] D. E. Tadych, M. Ford, B. G. Colby, L. E. Condon, [Historical patterns of well drilling and groundwater depth in Arizona considering groundwater regulation and surface water access](#), *JAWRA Journal of the American Water Resources Association* (2024) 1752–1688.13234doi:10.1111/1752-1688.13234.
URL <https://onlinelibrary.wiley.com/doi/10.1111/1752-1688.13234>
- [7] W. Covington, M. M. Moore, *Southwestern ponderosa pine forest structure: changes since Euro-American settlement*, *Journal of forestry*. 92 (1994) 39–47, place: Bethesda, MD : Publisher: Society of American Foresters,.

- [8] P. Friederici, Ecological Restoration of Southwestern Ponderosa Pine Forests, no. v.2 in Science Practice Ecological Restoration, Island Press, Chicago, 2013.
- [9] C. D. Allen, M. Savage, D. A. Falk, K. F. Suckling, T. W. Swetnam, T. Schulke, P. B. Stacey, P. Morgan, M. Hoffman, J. T. Klingel, [Ecological restoration of southwestern ponderosa pine ecosystems: a broad perspective](#), Ecological Applications 12 (5) (2002) 1418–1433. doi:10.1890/1051-0761(2002)012[1418:EROSPP]2.0.CO;2.
URL [http://doi.wiley.com/10.1890/1051-0761\(2002\)012\[1418:EROSPP\]2.0.CO;2](http://doi.wiley.com/10.1890/1051-0761(2002)012[1418:EROSPP]2.0.CO;2)
- [10] L. T. Berner, B. E. Law, A. J. H. Meddens, J. A. Hicke, [Tree mortality from fires, bark beetles, and timber harvest during a hot and dry decade in the western United States \(2003–2012\)](#), Environmental Research Letters 12 (6) (2017) 065005. doi:10.1088/1748-9326/aa6f94.
URL <https://iopscience.iop.org/article/10.1088/1748-9326/aa6f94>
- [11] V. Iglesias, J. K. Balch, W. R. Travis, [U.s. fires became larger, more frequent, and more widespread in the 2000s](#), Science Advances 8 (11) (03 2022). doi:10.1126/sciadv.abc0020.
URL <http://dx.doi.org/10.1126/sciadv.abc0020>
- [12] C. A. Schultz, T. Jedd, R. D. Beam, [The Collaborative Forest Landscape Restoration Program: A History and Overview of the First Projects](#), Journal of Forestry 110 (7) (2012) 381–391. doi:10.5849/jof.11-082.
URL <https://academic.oup.com/jof/article/110/7/381-391/4599006>
- [13] J. Bosch, J. Hewlett, [A review of catchment experiments to determine the effect of vegetation changes on water yield and evapotranspiration](#), Journal of Hydrology 55 (1-4) (1982) 3–23. doi:10.1016/0022-1694(82)90117-2.
URL <https://linkinghub.elsevier.com/retrieve/pii/0022169482901172>
- [14] M. B. Baker, [Effects of Ponderosa Pine Treatments on Water Yield in Arizona](#), Water Resources Research 22 (1) (1986) 67–73. doi:10.1029/WR022i001p00067.
URL <https://agupubs.onlinelibrary.wiley.com/doi/10.1029/WR022i001p00067>
- [15] G. J. Gottfried, [Moderate timber harvesting increases water yields from an Arizona Mixed Conifer Watershed](#), JAWRA Journal of the American Water Resources Association 27 (3) (1991) 537–546, publisher: John Wiley & Sons, Ltd.
URL <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1752-1688.1991.tb01454.x>
- [16] B. D. Smerdon, T. Redding, J. Beckers, [An overview of the effects of forest management on groundwater hydrology](#), Journal of Ecosystems and Management (Mar. 2009). doi:10.22230/jem.2009v10n1a409.
URL <https://jem-online.org/index.php/jem/article/view/409>

- [17] C. B. Zou, P. F. Ffolliott, M. Wine, [Streamflow responses to vegetation manipulations along a gradient of precipitation in the Colorado River Basin](#), *Forest Ecology and Management* 259 (7) (2010) 1268–1276. doi:10.1016/j.foreco.2009.08.005.
URL <https://linkinghub.elsevier.com/retrieve/pii/S0378112709005350>
- [18] C. J. W. Wyatt, [Estimating groundwater yield following forest restoration along the Mogollon Rim](#), Master’s thesis, Northern Arizona University, Flagstaff, AZ (2013).
URL <https://cdm17192.contentdm.oclc.org/digital/collection/p17192coll1/id/476/rec/3>
- [19] H. A. Moreno, H. V. Gupta, D. D. White, D. A. Sampson, [Modeling the distributed effects of forest thinning on the long-term water balance and stream flow extremes for a semi-arid basin in the southwestern US](#) (Oct. 2015). doi:10.5194/hessd-12-10827-2015.
URL <https://hess.copernicus.org/preprints/12/10827/2015/hessd-12-10827-2015.pdf>
- [20] S. Simonit, J. P. Connors, J. Yoo, A. Kinzig, C. Perrings, [The Impact of Forest Thinning on the Reliability of Water Supply in Central Arizona](#), *PLOS ONE* 10 (4) (2015) e0121596. doi:10.1371/journal.pone.0121596.
URL <https://dx.plos.org/10.1371/journal.pone.0121596>
- [21] C. Wyatt, F. C. O’Donnell, Abraham E. Springer, [Semi-Arid Aquifer Responses to Forest Restoration Treatments and Climate Change](#), *Ground Water* (2015). doi:10.1111/gwat.12184.
- [22] F. C. O’Donnell, W. T. Flatley, A. E. Springer, P. Z. Fulé, [Forest restoration as a strategy to mitigate climate impacts on wildfire, vegetation, and water in semiarid forests](#), *Ecological Applications* 28 (6) (2018) 1459–1472. doi:10.1002/eap.1746.
URL <https://esajournals.onlinelibrary.wiley.com/doi/10.1002/eap.1746>
- [23] E. R. Schenk, F. O’Donnell, A. E. Springer, L. E. Stevens, [The impacts of tree stand thinning on groundwater recharge in aridland forests](#), *Ecological Engineering* 145 (2020) 105701. doi:10.1016/j.ecoleng.2019.105701.
URL <https://linkinghub.elsevier.com/retrieve/pii/S0925857419304252>
- [24] A. R. Hibbert, [Managing vegetation to increase flow in the colorado river basin](#), Tech. rep. (1979).
- [25] A. D. Del Campo, K. Otsuki, Y. Serengil, J. A. Blanco, R. Yousefpour, X. Wei, [A global synthesis on the effects of thinning on hydrological processes: Implications for forest management](#), *Forest Ecology and Management* 519 (2022) 120324. doi:10.1016/j.foreco.2022.120324.
URL <https://linkinghub.elsevier.com/retrieve/pii/S0378112722003188>

- [26] T. Sankey, J. Donald, J. McVay, M. Ashley, F. O'Donnell, S. M. Lopez, A. Springer, [Multi-scale analysis of snow dynamics at the southern margin of the North American continental snow distribution](#), *Remote Sensing of Environment* 169 (2015) 307–319. doi:10.1016/j.rse.2015.08.028. URL <https://linkinghub.elsevier.com/retrieve/pii/S0034425715301140>
- [27] A. Belmonte, T. Sankey, J. Biederman, J. Bradford, S. Goetz, T. Kolb, [UAV-Based Estimate of Snow Cover Dynamics: Optimizing Semi-Arid Forest Structure for Snow Persistence](#), *Remote Sensing* 13 (5) (2021) 1036. doi:10.3390/rs13051036. URL <https://www.mdpi.com/2072-4292/13/5/1036>
- [28] J. Donager, T. T. Sankey, A. J. Sánchez Meador, J. B. Sankey, A. Springer, [Integrating airborne and mobile lidar data with UAV photogrammetry for rapid assessment of changing forest snow depth and cover](#), *Science of Remote Sensing* 4 (2021) 100029. doi:10.1016/j.srs.2021.100029. URL <https://linkinghub.elsevier.com/retrieve/pii/S266601722100016X>
- [29] A. Belmonte, T. Ts. Sankey, J. Biederman, J. B. Bradford, T. Kolb, [Soil moisture response to seasonal drought conditions and post-thinning forest structure](#), *Ecohydrology* 15 (5) (2022) e2406. doi:10.1002/eco.2406. URL <https://onlinelibrary.wiley.com/doi/10.1002/eco.2406>
- [30] T. Sankey, J. Tatum, [Thinning increases forest resiliency during unprecedented drought](#), *Scientific Reports* 12 (1) (2022) 9041. doi:10.1038/s41598-022-12982-z. URL <https://www.nature.com/articles/s41598-022-12982-z>
- [31] T. Sankey, A. Belmonte, R. Massey, J. Leonard, [Regional-scale forest restoration effects on ecosystem resiliency to drought: a synthesis of vegetation and moisture trends on Google Earth Engine](#), *Remote Sensing in Ecology and Conservation* 7 (2) (2021) 259–274. doi:10.1002/rse2.186. URL <https://zslpublications.onlinelibrary.wiley.com/doi/10.1002/rse2.186>
- [32] J. A. Biederman, M. D. Robles, R. L. Scott, J. F. Knowles, [Streamflow Response to Wildfire Differs With Season and Elevation in Adjacent Headwaters of the Lower Colorado River Basin](#), *Water Resources Research* 58 (3) (2022) e2021WR030687. doi:10.1029/2021WR030687. URL <https://agupubs.onlinelibrary.wiley.com/doi/10.1029/2021WR030687>
- [33] R. Moore, S. Wondzell, [Physical hydrology and the effects of forest harvesting in the pacific northwest: A review](#), *Journal of the American Water Resources Association* 41 (4) (2005) 763–784. doi:10.1111/j.1752-1688.2005.tb04463.x. URL <http://doi.wiley.com/10.1111/j.1752-1688.2005.tb04463.x>
- [34] J. A. Stanturf, B. J. Palik, R. K. Dumroese, [Contemporary forest restoration: A review emphasizing function](#), *Forest Ecology and Management* 331

- (2014) 292–323. doi:10.1016/j.foreco.2014.07.029.
URL <https://linkinghub.elsevier.com/retrieve/pii/S0378112714004654>
- [35] S. Filoso, M. Bezerra, K. C. B. Weiss, M. A. Palmer, *Impacts of forest restoration on water yield: A systematic review*, PLOS ONE 12 (8) (2017) e0183210. doi:10.1371/journal.pone.0183210.
URL <https://dx.plos.org/10.1371/journal.pone.0183210>
- [36] F. C. O'Donnell, *The influence of restoration treatments on hydrologic output in fire-adapted forests of the southwest*, Tech. rep., Flagstaff, AZ (11 2016).
URL <https://cdm17192.contentdm.oclc.org/digital/collection/p17192coll1/id/662/rec/1>
- [37] S. Fathi, J. Hagen, A. Matanó, G. E. H. Nogueira, *Review of GIS Multi-Criteria Decision Analysis for Managed Aquifer Recharge in Semi-Arid Regions*, Springer International Publishing, Cham, 2021, pp. 19–52, dOI: 10.1007/978-3-030-68124-1_2. doi:10.1007/978-3-030-68124-1_2.
URL https://link.springer.com/10.1007/978-3-030-68124-1_2
- [38] P. Raja Shekar, A. Mathew, *Assessing groundwater potential zones and artificial recharge sites in the monsoon-fed murredu river basin, india: An integrated approach using gis, ahp, and fuzzy-ahp*, Groundwater for Sustainable Development 23 (2023) 100994. doi:10.1016/j.gsd.2023.100994.
URL <https://linkinghub.elsevier.com/retrieve/pii/S2352801X23000942>
- [39] J. A. Biederman, A. J. Somor, A. A. Harpold, E. D. Gutmann, D. D. Breshears, P. A. Troch, D. J. Gochis, R. L. Scott, A. J. H. Meddens, P. D. Brooks, *Recent tree die-off has little effect on streamflow in contrast to expected increases from historical studies*, Water Resources Research 51 (12) (2015) 9775–9789. doi:10.1002/2015WR017401.
URL <https://agupubs.onlinelibrary.wiley.com/doi/10.1002/2015WR017401>
- [40] J. T. Parker, W. C. Steinkampf, M. E. Flynn, *Hydrogeology of the mogolon highlands, central arizona*, Tech. rep., Reston, VA, series: Scientific Investigations Report (2005).
URL <https://pubs.usgs.gov/sir/2004/5294/pdf/Parker%20SIR%202004-5294%20WEB.pdf>
- [41] A. J. Parker, *The topographic relative moisture index: An approach to soil-moisture assessment in mountain terrain*, Physical Geography 3 (2) (1982) 160–168. doi:10.1080/02723646.1982.10642224.
URL <https://www.tandfonline.com/doi/full/10.1080/02723646.1982.10642224>
- [42] A. D. Del Campo, M. González-Sanchis, A. J. Molina, A. García-Prats, C. J. Ceacero, I. Bautista, *Effectiveness of water-oriented thinning in two semiarid forests: The redistribution of increased net rainfall into soil water*,

- drainage and runoff, *Forest Ecology and Management* 438 (2019) 163–175.
[doi:10.1016/j.foreco.2019.02.020](https://doi.org/10.1016/j.foreco.2019.02.020).
 URL <https://linkinghub.elsevier.com/retrieve/pii/S0378112718323533>
- [43] K. L. Riley, I. C. Grenfell, J. D. Shaw, M. A. Finney, *Treemap 2016 dataset generates conus-wide maps of forest characteristics including live basal area, aboveground carbon, and number of trees per acre*, *Journal of Forestry* 120 (6) (2022) 607–632. [doi:10.1093/jofore/fvac022](https://doi.org/10.1093/jofore/fvac022).
 URL <https://academic.oup.com/jof/article/120/6/607/6701541>
- [44] T. Gleeson, N. Moosdorf, J. Hartmann, L. P. H. Van Beek, *A glimpse beneath earth’s surface: Global hydrogeology maps (glhymps) of permeability and porosity*, *Geophysical Research Letters* 41 (11) (2014) 3891–3898.
[doi:10.1002/2014GL059856](https://doi.org/10.1002/2014GL059856).
 URL <http://doi.wiley.com/10.1002/2014GL059856>