

# On Rowland's Sequence

Serafín Ruiz Cabello

Young Workshop on Arithmetics and Combinatorics  
ICMAT

June 21, 2011

Joint work with Fernando Chamizo and Dulcinea Raboso

## 1 Introduction

- Rowland's Sequence
- Auxiliary Sequences
- Generalization

## 2 Conjectures

- Conjectures
- Relation between conjectures

## 3 Primes

- Results
- Rowland's Chains

## 1 Introduction

- Rowland's Sequence
- Auxiliary Sequences
- Generalization

## 2 Conjectures

- Conjectures
- Relation between conjectures

## 3 Primes

- Results
- Rowland's Chains

# Rowland's Sequence

Rowland's Sequence

2008

$$\begin{cases} a_1 = 7 \\ a_k = a_{k-1} + \gcd(k, a_{k-1}); & k > 1 \end{cases}$$

# Rowland's Sequence

## Rowland's Sequence

2008

$$\begin{cases} a_1 = 7 \\ a_k = a_{k-1} + \gcd(k, a_{k-1}); \quad k > 1 \end{cases}$$

## First terms

$k$	1	2	3	4	5	6	7	8	9	10	11	...
$a_k$	7	8	9	10	15	18	19	20	21	22	33	...
												...

# Rowland's Sequence

## Rowland's Sequence

2008

$$\begin{cases} a_1 = 7 \\ a_k = a_{k-1} + \gcd(k, a_{k-1}); \quad k > 1 \end{cases}$$

## First terms

$k$	1	2	3	4	5	6	7	8	9	10	11	...
$a_k$	7	8	9	10	15	18	19	20	21	22	33	...
$a_k - a_{k-1}$		1	1	1	5	3	1	1	1	1	11	...

# Rowland's Sequence

## Rowland's Sequence

2008

$$\begin{cases} a_1 = 7 \\ a_k = a_{k-1} + \gcd(k, a_{k-1}); \quad k > 1 \end{cases}$$

## First terms

$k$	1	2	3	4	5	6	7	8	9	10	11	...
$a_k$	7	8	9	10	15	18	19	20	21	22	33	...
$a_k - a_{k-1}$		1	1	1	5	3	1	1	1	1	11	...

The sequence  $\{a_k - a_{k-1}\}_{k>1}$  has a surprising property:

$$\{a_k - a_{k-1}\} =$$

1, 1, 1, 5, 3, 1, 1, 1, 1, 11, 3, 1, 1, 1, 1, 1, 1, 1, 1, 1, 23, 3, 1,  
1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 47, 3, 1, 5, 3, 1, 1,  
1,  
1, 101, 3, 1, 1, 7,  
1, 1, 1, 1, 11, 3, 1, 1, 1, 1, 1, 13, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1,  
1,  
1,  
1,  
1,  
1,  
1,  
1, ...



[illegible]

Every term distinct than 1 is a prime number.

# Rowland's Sequence

## Rowland's Sequence

2008

$$\begin{cases} a_1 = 7 \\ a_k = a_{k-1} + \gcd(k, a_{k-1}); \quad k > 1 \end{cases}$$

## First terms

$k$	1	2	3	4	5	6	7	8	9	10	11	...
$a_k$	7	8	9	10	15	18	19	20	21	22	33	...
$a_k - a_{k-1}$		1	1	1	5	3	1	1	1	1	11	...

# Rowland's Sequence

## Rowland's Sequence

2008

$$\begin{cases} a_1 = 7 \\ a_k = a_{k-1} + \gcd(k, a_{k-1}); \quad k > 1 \end{cases}$$

## First terms

$k$	1	2	3	4	5	6	7	8	9	10	11	...
$a_k$	7	8	9	10	15	18	19	20	21	22	33	...
$a_k - a_{k-1}$		1	1	1	5	3	1	1	1	1	11	...

## Theorem

$a_k - a_{k-1}$  is 1 or prime for  $k > 1$ .

# Auxiliary sequences

We have an elementary proof for this theorem.

# Auxiliary sequences

We have an elementary proof for this theorem.

## Auxiliary sequences

$$\begin{cases} c_1^* = 5 \\ c_n^* = c_{n-1}^* + \text{lpf}(c_{n-1}^*) - 1, n > 1 \end{cases} ; \quad r_n^* = \frac{c_n^* + 1}{2}, n \geq 1.$$

# Auxiliary sequences

We have an elementary proof for this theorem.

## Auxiliary sequences

$$\begin{cases} c_1^* = 5 \\ c_n^* = c_{n-1}^* + \text{lpf}(c_{n-1}^*) - 1, n > 1 \end{cases} ; \quad r_n^* = \frac{c_n^* + 1}{2}, n \geq 1.$$

Here,  $\text{lpf}(\cdot)$  is the least prime factor of an integer.

# Auxiliary sequences

We have an elementary proof for this theorem.

## Auxiliary sequences

$$\begin{cases} c_1^* = 5 \\ c_n^* = c_{n-1}^* + \text{lpf}(c_{n-1}^*) - 1, n > 1 \end{cases} ; \quad r_n^* = \frac{c_n^* + 1}{2}, n \geq 1.$$

Here,  $\text{lpf}(\cdot)$  is the least prime factor of an integer.

## Proposition

$$a_k - a_{k-1} = \begin{cases} \text{lpf}(c_{n-1}^*), & \text{if } k = r_n^* \text{ for some } n > 1. \\ 1, & \text{otherwise} \end{cases}$$

# Auxiliary sequences

We have an elementary proof for this theorem.

## Auxiliary sequences

$$\begin{cases} c_1^* = 5 \\ c_n^* = c_{n-1}^* + \text{lpf}(c_{n-1}^*) - 1, n > 1 \end{cases} ; \quad r_n^* = \frac{c_n^* + 1}{2}, n \geq 1.$$

Here,  $\text{lpf}(\cdot)$  is the least prime factor of an integer.

## Proposition

$$a_k - a_{k-1} = \begin{cases} \text{lpf}(c_{n-1}^*), & \text{if } k = r_n^* \text{ for some } n > 1. \\ 1, & \text{otherwise} \end{cases}$$

Using this proof, we obtain another amazing result:



# Auxiliary sequences

We have an elementary proof for this theorem.

## Auxiliary sequences

$$\begin{cases} c_1^* = 5 \\ c_n^* = c_{n-1}^* + \text{lpf}(c_{n-1}^*) - 1, n > 1 \end{cases} ; \quad r_n^* = \frac{c_n^* + 1}{2}, n \geq 1.$$

Here,  $\text{lpf}(\cdot)$  is the least prime factor of an integer.

## Proposition

$$a_k - a_{k-1} = \begin{cases} \text{lpf}(c_{n-1}^*), & \text{if } k = r_n^* \text{ for some } n > 1. \\ 1, & \text{otherwise} \end{cases}$$

Using this proof, we obtain another amazing result:

## Proposition

$\{a_k - a_{k-1}\}_{k>1}$  contains infinitely many primes.

# Generalized Rowland's Sequence

We want to study a more general sequence replacing  $a_1 = 7$  by any integer  $a_1 \geq 1$ .

# Generalized Rowland's Sequence

We want to study a more general sequence replacing  $a_1 = 7$  by any integer  $a_1 \geq 1$ .

## Generalized Rowland's Sequence

$$\begin{cases} a_1 \in \mathbb{Z}^+ \\ a_k = a_{k-1} + \gcd(k, a_{k-1}); \quad k > 1 \end{cases}$$

# Generalized Rowland's Sequence

We want to study a more general sequence replacing  $a_1 = 7$  by any integer  $a_1 \geq 1$ .

## Generalized Rowland's Sequence

$$\begin{cases} a_1 \in \mathbb{Z}^+ \\ a_k = a_{k-1} + \gcd(k, a_{k-1}); \quad k > 1 \end{cases}$$

- We can discard any even  $a_1$ , since  $a_1 = 2A$  and  $a_1 = 2A + 1$  give the same  $a_2$ .

# Generalized Rowland's Sequence

We want to study a more general sequence replacing  $a_1 = 7$  by any integer  $a_1 \geq 1$ .

## Generalized Rowland's Sequence

$$\begin{cases} a_1 \in \mathbb{Z}^+ \\ a_k = a_{k-1} + \gcd(k, a_{k-1}); \quad k > 1 \end{cases}$$

- We can discard any even  $a_1$ , since  $a_1 = 2A$  and  $a_1 = 2A + 1$  give the same  $a_2$ .
- Also,  $a_1 = 1$  and  $a_1 = 3$  lead to the sequences  $a_k = k$  and  $a_k = k + 2$ , respectively.

# Generalized Rowland's Sequence

We want to study a more general sequence replacing  $a_1 = 7$  by any integer  $a_1 \geq 1$ .

## Generalized Rowland's Sequence

$$\begin{cases} a_1 \in \mathbb{Z}^+, a_1 \text{ odd}, a_1 > 3 \\ a_k = a_{k-1} + \gcd(k, a_{k-1}); \quad k > 1 \end{cases}$$

- We can discard any even  $a_1$ , since  $a_1 = 2A$  and  $a_1 = 2A + 1$  give the same  $a_2$ .
- Also,  $a_1 = 1$  and  $a_1 = 3$  lead to the sequences  $a_k = k$  and  $a_k = k + 2$ , respectively.

# Generalized Rowland's Sequence

We want to study a more general sequence replacing  $a_1 = 7$  by any integer  $a_1 \geq 1$ .

## Generalized Rowland's Sequence

$$\begin{cases} a_1 \in \mathbb{Z}^+, a_1 \text{ odd}, a_1 > 3 \\ a_k = a_{k-1} + \gcd(k, a_{k-1}); \quad k > 1 \end{cases}$$

- We can discard any even  $a_1$ , since  $a_1 = 2A$  and  $a_1 = 2A + 1$  give the same  $a_2$ .
- Also,  $a_1 = 1$  and  $a_1 = 3$  lead to the sequences  $a_k = k$  and  $a_k = k + 2$ , respectively.
- Unfortunately, under these conditions,  $a_k - a_{k-1}$  is not necessarily prime. For instance, if  $a_1 = 533$ , then  $a_{18} - a_{17} = 9$ .

The proof of the theorem suggests to introduce a general recurrence:



The proof of the theorem suggests to introduce a general recurrence:

$$\begin{cases} r_1 = 1 \\ r_{n+1} = \min\{p + p\lfloor r_n/p \rfloor : p|c_n\}, \quad n \geq 1 \end{cases}$$
$$\begin{cases} c_1 = a_1 - 2 \\ c_{n+1} = c_n + \gcd(c_n, r_{n+1}) - 1, \quad n \geq 1 \end{cases}$$

Equivalently,  $r_{n+1} = \min\{k > r_n : (k, c_n) \neq 1\}$ .

The proof of the theorem suggests to introduce a general recurrence:

$$\begin{cases} r_1 = 1 \\ r_{n+1} = \min\{p + p\lfloor r_n/p \rfloor : p|c_n\}, \quad n \geq 1 \\ c_1 = a_1 - 2 \\ c_{n+1} = c_n + \gcd(c_n, r_{n+1}) - 1, \quad n \geq 1 \end{cases}$$

Equivalently,  $r_{n+1} = \min\{k > r_n : (k, c_n) \neq 1\}$ .

### Proposition

For any odd  $a_1 > 3$ , the sequence  $\{a_k\}$  satisfies  $a_k = c_n + k + 1$ , for  $r_n \leq k < r_{n+1}$ . Moreover,

$$a_k - a_{k-1} = \begin{cases} \gcd(c_{n-1}, r_n), & \text{if } k = r_n \text{ for some } n > 1. \\ 1, & \text{otherwise.} \end{cases}$$

## 1 Introduction

- Rowland's Sequence
- Auxiliary Sequences
- Generalization

## 2 Conjectures

- Conjectures
- Relation between conjectures

## 3 Primes

- Results
- Rowland's Chains

We have seen that for an arbitrary  $a_1 > 3$ , composite numbers may appear. Nevertheless, counterexamples are hard to find. Computational tests suggest the following:

We have seen that for an arbitrary  $a_1 > 3$ , composite numbers may appear. Nevertheless, counterexamples are hard to find. Computational tests suggest the following:

### Conjecture (A)

For any Generalized Rowland's Sequence, there exists a positive integer  $N$  such that  $a_k - a_{k-1}$  is 1 or prime for every  $k > N$ .

We have seen that for an arbitrary  $a_1 > 3$ , composite numbers may appear. Nevertheless, counterexamples are hard to find. Computational tests suggest the following:

### Conjecture (A)

For any Generalized Rowland's Sequence, there exists a positive integer  $N$  such that  $a_k - a_{k-1}$  is 1 or prime for every  $k > N$ .

The proof of the first theorem may be applied when  $c_n = 2r_n - 1$  for some  $n \in \mathbb{Z}^+$ . This implies  $c_l = 2r_l - 1$  for  $l > n$ .

We have seen that for an arbitrary  $a_1 > 3$ , composite numbers may appear. Nevertheless, counterexamples are hard to find. Computational tests suggest the following:

### Conjecture (A)

For any Generalized Rowland's Sequence, there exists a positive integer  $N$  such that  $a_k - a_{k-1}$  is 1 or prime for every  $k > N$ .

The proof of the first theorem may be applied when  $c_n = 2r_n - 1$  for some  $n \in \mathbb{Z}^+$ . This implies  $c_l = 2r_l - 1$  for  $l > n$ . For instance, if  $a_1 = 117$ ,

We have seen that for an arbitrary  $a_1 > 3$ , composite numbers may appear. Nevertheless, counterexamples are hard to find. Computational tests suggest the following:

### Conjecture (A)

For any Generalized Rowland's Sequence, there exists a positive integer  $N$  such that  $a_k - a_{k-1}$  is 1 or prime for every  $k > N$ .

The proof of the first theorem may be applied when  $c_n = 2r_n - 1$  for some  $n \in \mathbb{Z}^+$ . This implies  $c_l = 2r_l - 1$  for  $l > n$ . For instance, if  $a_1 = 117$ ,

$n$	1	2	3	4	5	6	7	8	9	10	...
$r_n$	1						...	...	...	...	...
$c_n$	115						...	...	...	...	...



We have seen that for an arbitrary  $a_1 > 3$ , composite numbers may appear. Nevertheless, counterexamples are hard to find. Computational tests suggest the following:

### Conjecture (A)

For any Generalized Rowland's Sequence, there exists a positive integer  $N$  such that  $a_k - a_{k-1}$  is 1 or prime for every  $k > N$ .

The proof of the first theorem may be applied when  $c_n = 2r_n - 1$  for some  $n \in \mathbb{Z}^+$ . This implies  $c_l = 2r_l - 1$  for  $l > n$ . For instance, if  $a_1 = 117$ ,

$n$	1	2	3	4	5	6	7	8	9	10	...
$r_n$	1	5	7	10	12		...	...	...	...	...
$c_n$	115	119	125	129	131		...	...	...	...	...

We have seen that for an arbitrary  $a_1 > 3$ , composite numbers may appear. Nevertheless, counterexamples are hard to find. Computational tests suggest the following:

### Conjecture (A)

For any Generalized Rowland's Sequence, there exists a positive integer  $N$  such that  $a_k - a_{k-1}$  is 1 or prime for every  $k > N$ .

The proof of the first theorem may be applied when  $c_n = 2r_n - 1$  for some  $n \in \mathbb{Z}^+$ . This implies  $c_l = 2r_l - 1$  for  $l > n$ . For instance, if  $a_1 = 117$ , then  $c_n = 2r_n - 1$  **for the first time when  $n = 6$** .

$n$	1	2	3	4	5	6	7	8	9	10	...
$r_n$	1	5	7	10	12	131	...	...	...	...	...
$c_n$	115	119	125	129	131	261	...	...	...	...	...

We have seen that for an arbitrary  $a_1 > 3$ , composite numbers may appear. Nevertheless, counterexamples are hard to find. Computational tests suggest the following:

### Conjecture (A)

For any Generalized Rowland's Sequence, there exists a positive integer  $N$  such that  $a_k - a_{k-1}$  is 1 or prime for every  $k > N$ .

The proof of the first theorem may be applied when  $c_n = 2r_n - 1$  for some  $n \in \mathbb{Z}^+$ . This implies  $c_l = 2r_l - 1$  for  $l > n$ . For instance, if  $a_1 = 117$ , then  $c_n = 2r_n - 1$  **for the first time when  $n = 6$** . And, for every  $l > 6$ ,

$n$	1	2	3	4	5	6	7	8	9	10	...
$r_n$	1	5	7	10	12	131	...	...	...	...	...
$c_n$	115	119	125	129	131	261	...	...	...	...	...

We have seen that for an arbitrary  $a_1 > 3$ , composite numbers appear. Nevertheless, counterexamples are hard to find. Computational tests suggest the following:

### Conjecture (A)

For any Generalized Rowland's Sequence, there exists a positive integer  $N$  such that  $a_k - a_{k-1}$  is 1 or prime for every  $k > N$ .

The proof of the first theorem may be applied when  $c_n = 2r_n - 1$  for some  $n \in \mathbb{Z}^+$ . This implies  $c_l = 2r_l - 1$  for  $l > n$ . For instance, if  $a_1 = 117$ , then  $c_n = 2r_n - 1$  **for the first time when  $n = 6$** . And, for every  $l > 6$ , **all pairs  $(r_l, c_l)$  verify  $c_l = 2r_l - 1$** .

$n$	1	2	3	4	5	6	7	8	9	10	...
$r_n$	1	5	7	10	12	131	132	263	264	272	...
$c_n$	115	119	125	129	131	261	263	525	527	543	...

We have seen that for an arbitrary  $a_1 > 3$ , composite numbers appear. Nevertheless, counterexamples are hard to find. Computational tests suggest the following:

### Conjecture (A)

For any Generalized Rowland's Sequence, there exists a positive integer  $N$  such that  $a_k - a_{k-1}$  is 1 or prime for every  $k > N$ .

In this example,  $c_5$  is the first prime term in the sequence  $\{c_n\}$ .

$n$	1	2	3	4	5	6	7	8	9	10	...
$r_n$	1	5	7	10	12	131	132	263	264	272	...
$c_n$	115	119	125	129	131	261	263	525	527	543	...

We have seen that for an arbitrary  $a_1 > 3$ , composite numbers appear. Nevertheless, counterexamples are hard to find. Computational tests suggest the following:

### Conjecture (A)

For any Generalized Rowland's Sequence, there exists a positive integer  $N$  such that  $a_k - a_{k-1}$  is 1 or prime for every  $k > N$ .

In this example,  $c_5$  is the first prime term in the sequence  $\{c_n\}$ .

Given  $n > 1$ ,  $r_n = c_{n-1}$  if and only if  $c_{n-1}$  is prime.

$n$	1	2	3	4	5	6	7	8	9	10	...
$r_n$	1	5	7	10	12	131	132	263	264	272	...
$c_n$	115	119	125	129	131	261	263	525	527	543	...

We have seen that for an arbitrary  $a_1 > 3$ , composite numbers appear. Nevertheless, counterexamples are hard to find. Computational tests suggest the following:

### Conjecture (A)

For any Generalized Rowland's Sequence, there exists a positive integer  $N$  such that  $a_k - a_{k-1}$  is 1 or prime for every  $k > N$ .

In this example,  $c_5$  is the first prime term in the sequence  $\{c_n\}$ .

Given  $n > 1$ ,  $r_n = c_{n-1}$  if and only if  $c_{n-1}$  is prime.  
If  $r_n = c_{n-1}$ , then  $c_n = 2r_n - 1$ .

$n$	1	2	3	4	5	6	7	8	9	10	...
$r_n$	1	5	7	10	12	131	132	263	264	272	...
$c_n$	115	119	125	129	131	261	263	525	527	543	...

$n$	1	2	3	4	5	6	7	8	9	10	...
$r_n$	1	5	7	10	12	131	132	263	264	272	...
$c_n$	115	119	125	129	131	261	263	525	527	543	...



$n$	1	2	3	4	5	6	7	8	9	10	...
$r_n$	1	5	6	11	12	23	24	47	48	50	...
$c_n$	5	9	11	21	23	45	47	93	95	99	...

$n$	1	2	3	4	5	6	7	8	9	10	...
$r_n$	1	3	5	6	41	42	83	84	167	168	...
$c_n$	33	35	39	41	81	83	165	167	333	335	...

$n$	1	2	3	4	5	6	7	8	9	10	11
$r_n$	1	7	11	17	18	20	21	29	30	35	587
$c_n$	511	517	527	543	545	549	551	579	581	587	1173

$n$	1	2	3	4	5	6	7	8	9	10	...
$r_n$	1	5	7	10	12	131	132	263	264	272	...
$c_n$	115	119	125	129	131	261	263	525	527	543	...

Conjecture (A) holds if  $c_n = 2r_n - 1$  for some  $n \in \mathbb{Z}^+$ , or if  $c_m$  is prime for some  $m \in \mathbb{Z}^+$ . Computations suggest that this happens for any initial condition  $a_1$ . Moreover, it seems that the minimal choices of  $m$  and  $n$  in these claims are always consecutive.

Conjecture (A) holds if  $c_n = 2r_n - 1$  for some  $n \in \mathbb{Z}^+$ , or if  $c_m$  is prime for some  $m \in \mathbb{Z}^+$ . Computations suggest that this happens for any initial condition  $a_1$ . Moreover, it seems that the minimal choices of  $m$  and  $n$  in these claims are always consecutive.

### Conjecture (B)

For an odd  $a_1 > 3$ , we define (writing  $\inf \emptyset = \infty$ )

$$\begin{aligned}n_0 &= \inf\{n \in \mathbb{Z}^+ : c_n = 2r_n - 1\}, \\m_0 &= \inf\{n \in \mathbb{Z}^+ : c_n \text{ is prime}\}.\end{aligned}$$

Then

$$(i) \, n_0 < \infty, \quad (ii) \, m_0 < \infty, \quad (iii) \, n_0 = m_0 + 1 < \infty$$

Conjecture (A) holds if  $c_n = 2r_n - 1$  for some  $n \in \mathbb{Z}^+$ , or if  $c_m$  is prime for some  $m \in \mathbb{Z}^+$ . Computations suggest that this happens for any initial condition  $a_1$ . Moreover, it seems that the minimal choices of  $m$  and  $n$  in these claims are always consecutive.

### Conjecture (B)

For an odd  $a_1 > 3$ , we define (writing  $\inf \emptyset = \infty$ )

$$\begin{aligned}n_0 &= \inf\{n \in \mathbb{Z}^+ : c_n = 2r_n - 1\}, \\m_0 &= \inf\{n \in \mathbb{Z}^+ : c_n \text{ is prime}\}.\end{aligned}$$

Then

$$(i) \, n_0 < \infty, \quad (ii) \, m_0 < \infty, \quad (iii) \, n_0 = m_0 + 1 < \infty$$

As we saw before, (ii) implies (i). Trivially, (iii) implies (ii) and (i).

# Relation between the conjectures

If Conjecture (B) is true, then so it is Conjecture (A):

$n$	1	2	3	4	5	6	7	8	9	10	...
$r_n$	1	5	7	10	12	131	132	263	264	272	...
$c_n$	115	119	125	129	131	261	263	525	527	543	...

# Relation between the conjectures

If Conjecture (B) is true, then so it is Conjecture (A):

## Proposition

Under (i), (ii) or (iii), Conjecture (A) is true. Moreover,  $\{a_k - a_{k-1}\}_{k \geq 1}$  contains infinitely many primes.

$n$	1	2	3	4	5	6	7	8	9	10	...
$r_n$	1	5	7	10	12	131	132	263	264	272	...
$c_n$	115	119	125	129	131	261	263	525	527	543	...

## Relation between the conjectures

If Conjecture (B) is true, then so it is Conjecture (A):

### Proposition

Under (i), (ii) or (iii), Conjecture (A) is true. Moreover,  $\{a_k - a_{k-1}\}_{k \geq 1}$  contains infinitely many primes.

### Proposition

For any odd  $a_1 > 3$ ,  $r_n \leq (c_n + 1)/2$ , for every  $n > 1$ . Moreover, the equality occurs if and only if  $\gcd(c_{n-1}, r_n)$  is prime  $p$  and  $p \lfloor r_{n-1}/p \rfloor = (c_{n-1} - p)/2$ .

$n$	1	2	3	4	5	6	7	8	9	10	...
$r_n$	1	5	7	10	12	131	132	263	264	272	...
$c_n$	115	119	125	129	131	261	263	525	527	543	...

A little more is needed for proving (iii).



A little more is needed for proving (iii).

### Proposition

If (i) holds and  $\gcd(c_{n_0-1}, r_{n_0}) > r_{n_0-1}$ , then (iii) is true.

A little more is needed for proving (iii).

### Proposition

If (i) holds and  $\gcd(c_{n_0-1}, r_{n_0}) > r_{n_0-1}$ , then (iii) is true.

### Proposition

Assume (i) and  $(2 + 1/2500)r_n < c_n + 1$ , for  $n < n_0$ . Then (iii) holds.

A little more is needed for proving (iii).

### Proposition

If (i) holds and  $\gcd(c_{n_0-1}, r_{n_0}) > r_{n_0-1}$ , then (iii) is true.

### Proposition

Assume (i) and  $(2 + 1/2500)r_n < c_n + 1$ , for  $n < n_0$ . Then (iii) holds.

### Proposition

Under (iii), there exists a prime  $p$  such that

$$\begin{aligned}\frac{p+1}{2} &= \inf\{k : a_k = 3k\}, \\ p &= \inf\{k : a_k = 3k, a_k - a_{k-1} > 1\}.\end{aligned}$$

## 1 Introduction

- Rowland's Sequence
- Auxiliary Sequences
- Generalization

## 2 Conjectures

- Conjectures
- Relation between conjectures

## 3 Primes

- Results
- Rowland's Chains

# Size of $m_0$

For a fixed  $a_1$ , usually  $m_0$  is *small*.

Size of  $m_0$ 

For a fixed  $a_1$ , usually  $m_0$  is *small*. In this example,  $m_0 = 5$ :

$n$	1	2	3	4	5	6	7	8	9	10	...
$r_n$	1	5	7	10	12	131	132	263	264	272	...
$c_n$	115	119	125	129	131	261	263	525	527	543	...

Size of  $m_0$ 

For a fixed  $a_1$ , usually  $m_0$  is *small*. In this example,  $m_0 = 5$ :

$n$	1	2	3	4	5	6	7	8	9	10	...
$r_n$	1	5	7	10	12	131	132	263	264	272	...
$c_n$	115	119	125	129	131	261	263	525	527	543	...

Anyway, given  $N$ , there exists  $a_1$  such that  $m_0 > N$ . Hence:

Size of  $m_0$ 

For a fixed  $a_1$ , usually  $m_0$  is *small*. In this example,  $m_0 = 5$ :

$n$	1	2	3	4	5	6	7	8	9	10	...
$r_n$	1	5	7	10	12	131	132	263	264	272	...
$c_n$	115	119	125	129	131	261	263	525	527	543	...

Anyway, given  $N$ , there exists  $a_1$  such that  $m_0 > N$ . Hence:

**Proposition**

$m_0$  can be arbitrarily large.



Size of  $m_0$ 

For a fixed  $a_1$ , usually  $m_0$  is *small*. In this example,  $m_0 = 5$ :

$n$	1	2	3	4	5	6	7	8	9	10	...
$r_n$	1	5	7	10	12	131	132	263	264	272	...
$c_n$	115	119	125	129	131	261	263	525	527	543	...

Anyway, given  $N$ , there exists  $a_1$  such that  $m_0 > N$ . Hence:

**Proposition**

$m_0$  can be arbitrarily large.

Given  $a_1$ , with  $m_0 < \infty$ ; and taking  $a'_1 = a_1 + c_{m_0}!$ , it can be proved that  $m'_0 > m_0$ .

# Rowland's Chains

We are interested on studying subsequences of primes appearing inside  $\{a_k - a_{k-1}\}$ .

# Rowland's Chains

We are interested on studying subsequences of primes appearing inside  $\{a_k - a_{k-1}\}$ . For instance, taking  $a_1 = 7$ ,

# Rowland's Chains

We are interested on studying subsequences of primes appearing inside  $\{a_k - a_{k-1}\}$ . For instance, taking  $a_1 = 7$ ,  $\{a_k - a_{k-1}\} =$

# Rowland's Chains

We are interested on studying subsequences of primes appearing inside  $\{a_k - a_{k-1}\}$ . For instance, taking  $a_1 = 7$ ,  $\{a_k - a_{k-1}\} =$   
1, 1, 1, 5, 3, 1, 1, 1, 1, 11, 3, 1, 1, 1, 1, 1, 1, 1, 1, 1, 23, 3, 1, 1,  
1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 47, 3, 1, 5, 3, 1, 1, 1,  
1,  
1,  
1, 1, 11, 3, 1, 1, 1, 1, 1, 13, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1,  
1,  
1,  
1,  
1,  
1, ...

# Rowland's Chains

We are interested on studying subsequences of primes appearing inside  $\{a_k - a_{k-1}\}$ . For instance, taking  $a_1 = 7$ ,  $\{a_k - a_{k-1}\} =$   
 $1, 1, 1, \textcolor{red}{5}, \textcolor{red}{3}, 1, 1, 1, 1, \textcolor{red}{11}, \textcolor{red}{3}, 1, 1, 1, 1, 1, 1, 1, 1, \textcolor{red}{23}, \textcolor{red}{3}, 1, 1,$   
 $1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, \textcolor{red}{47}, \textcolor{red}{3}, 1, \textcolor{red}{5}, \textcolor{red}{3}, 1, 1, 1,$   
 $1, 1,$   
 $1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, \textcolor{red}{101}, \textcolor{red}{3}, 1, 1, \textcolor{red}{7}, 1, 1,$   
 $1, 1, \textcolor{red}{11}, \textcolor{red}{3}, 1, 1, 1, 1, 1, \textcolor{red}{13}, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1,$   
 $1, 1,$   
 $1, 1,$   
 $1, 1,$   
 $1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, \textcolor{red}{233}, \textcolor{red}{3}, 1, 1, 1, 1, \dots$

Removing the ones, we obtain a sequence of primes:

# Rowland's Chains

[illegible]

Removing the ones, we obtain a sequence of primes:

5, 3, 11, 3, 23, 3, 47, 3, 5, 3, 101, 3, 7, 11, 3, 13, 233, 3.

### On Rowland's Sequence

This sequence is a **Rowland's Chain**.



# Rowland's Chains

[illegible]

In general, a **Rowland's Chain** is any subsequence of concatenated primes inside a sequence  $\{a_k - a_{k-1}\}_{k>n_0}$ , for any odd  $a_1 > 3$ .

We associate to the chain  $C_k = \{p_1, p_2, \dots, p_k\}$  the shifted partial sums

$$S(n) = \sum_{j < n} (p_j - 1), \quad S(1) = 0.$$

We associate to the chain  $C_k = \{p_1, p_2, \dots, p_k\}$  the shifted partial sums

$$S(n) = \sum_{j < n} (p_j - 1), \quad S(1) = 0.$$

The following is a characterization of Rowland's chains:

We associate to the chain  $C_k = \{p_1, p_2, \dots, p_k\}$  the shifted partial sums

$$S(n) = \sum_{j < n} (p_j - 1), \quad S(1) = 0.$$

The following is a characterization of Rowland's chains:

### Proposition

$C_k$  is a Rowland's Chain if and only if it verifies these conditions:

- $S(m) \equiv S(n) \pmod{p_n}$ , when  $p_n = p_m$ .
- $S(m) \not\equiv S(n) \pmod{p_n}$ , when  $p_n < p_m$ .
- For any prime  $q$ , the set  $\{S(j) \pmod{q} : p_j > q\}$  does not contain all residue classes modulo  $q$ .

We associate to the chain  $C_k = \{p_1, p_2, \dots, p_k\}$  the shifted partial sums

$$S(n) = \sum_{j < n} (p_j - 1), \quad S(1) = 0.$$

The following is a characterization of Rowland's chains:

### Proposition

$C_k$  is a Rowland's Chain if and only if it verifies these conditions:

- $S(m) \equiv S(n) \pmod{p_n}$ , when  $p_n = p_m$ .
- $S(m) \not\equiv S(n) \pmod{p_n}$ , when  $p_n < p_m$ .
- For any prime  $q$ , the set  $\{S(j) \pmod{q} : p_j > q\}$  does not contain all residue classes modulo  $q$ .

There are lots of restrictions. For instance,  $\{11, 5, p\}$  is not a Rowland's chain for any  $p > 3$ .

## Corollary

If  $p_1, p_2, \dots, p_k$  are distinct primes, then  
 $C_{2k} = \{p_1, p_2, \dots, p_k, p_1, p_2, \dots, p_k\}$  is not a Rowland's chain.

## Corollary

If  $p_1, p_2, \dots, p_k$  are distinct primes, then  $C_{2k} = \{p_1, p_2, \dots, p_k, p_1, p_2, \dots, p_k\}$  is not a Rowland's chain.

In spite of the last Corollary, large non-consecutive repetitions can be found. For instance,

$$C_{27} = \{3, 5, 3, 23, 3, 5, 3, 653, 3, 5, 3, 23, 3, 5, 3, 3603833, \\ 3, 5, 3, 23, 3, 5, 3, 653, 3, 5, 3\}$$

has length 27 but just 5 primes. This Chain corresponds to  $a_1 = 1550303031682205$ .

## References

- FERNANDO CHAMIZO, DULCINEA RABOSO AND SERAFÍN RUIZ-CABELLO. On Rowland's Sequence. *The Electronic Journal of Combinatorics*, 18(2): Article 10, 2011.  
[http://www.combinatorics.org/Volume\\_18/PDF/v18i2p10.pdf](http://www.combinatorics.org/Volume_18/PDF/v18i2p10.pdf)
- ERIC S. ROWLAND. A Natural Prime-generating Recurrence. *J. Integer Seq.*, 11(2): Article 08.2.8, 13, 2008.  
<http://www.cs.uwaterloo.ca/journals/JIS/VOL11/Rowland/rowland21.pdf>