

Modelling the Relationship between Climate Modes and Eastern Australia Atmospheric Carbon Monoxide

Ryan H. Peterson & Lakin Moser

Colorado School of Mines

May 8, 2024

Introduction

- The 2019/2020 Australian wildfire season was notably severe, with impacts on local ecosystems, lives, and property.
- Climate modes, especially El Niño Southern Oscillation (ENSO), have an effect on global and local climate conditions.
 - Specifically, the effect of southern hemisphere climate modes on eastern Australian wildfires.
- NSF NCAR project: Atmospheric impacts of the extreme Australian 2019/2020 wildfire season.
- Goal is to create a statistical model for the relationships between climate modes and atmospheric compositions during the fire season.
 - Building on earlier work for maritime south east Asia done by Daniels et al., 2022.

Background

- Using anomaly carbon monoxide (CO) as proxy for wildfire intensity Buchholz et al., 2018.
 - MOPITT total column CO.
 - Excluding 2019/2020 data.
- Exploring CO anomalies in north east Australia (NE Aus) and south east Australia (SE Aus).
- Using climate mode indices in below table.
 - Sea-surface temperature (SST) anomalies.

Climate Mode	Index	Source
El Niño-Southern Oscillation (ENSO)	Niño 3.4	NOAA, 2021b
Indian Ocean Dipole (IOD)	DMI	NOAA, 2021b
Tropical South Atlantic (TSA)	TS)	NOAA, 2021b
Antarctic Oscillation (AAO)	SAM	NOAA, 2021a

Table: Climate modes considered as predictors with index and source.

Motivation




Figure: Climate Mode indices and response regions.

Response Time Series




Figure: Response time series for response regions.

Predictor Time Series




Figure: Predictors time series for climate mode indices.

Vector Autoregression

- Consider the following time series for NE Aus $Y_{t,1}$ and SE Aus $Y_{t,2}$.
- And for climate mode indices; Niño 3.4 $X_{t,1}$, DMI $X_{t,2}$, TSA $X_{t,3}$, and SAM (AAO) $X_{t,4}$.
- Then a VAR(p) model for a given time series $Y_{t,j}$ for $j = 1, 2$, can be written as

$$Y_{t,j} = \alpha + \sum_{i=1}^p \Phi_i^T \mathbf{X}_{t-i} + w_t,$$

where $\Phi_i = [\phi_{i,j}, \phi_{i,1}, \phi_{i,2}, \phi_{i,3}, \phi_{i,4}]^T$ and $\mathbf{X}_{t-i} = [Y_{t-i,j}, X_{t-i,1}, X_{t-i,2}, X_{t-i,3}, X_{t-i,4}]^T$, with α as an intercept and $w_t \sim iid(0, \sigma_w^2)$.

Model Selection





Figure: Used AIC to inform choice of p .


VAR Models

- We choose a VAR(4) for both NE Aus and SE Aus.
 - Parameters estimated by OLS.
 - Includes trend and constant.
- Using the 'vars' package to fit the model.



PACF of Residuals for NE Aus





PACF of Residuals for SE Aus



Fitted Model



Residuals



Summary

- Using climate mode indices we are able create a time series model for CO anomalies in eastern Australia.
- Surprisingly, we did not need to distinguish wildfire season data.
- We also checked a VARMA(p,q) model but results were effectively the same.

- Further refinement of these models specifically using:
 - Regression with ARMA errors.
 - Regularization methods.
 - Dynamic regression models.
- Add in Outgoing Longwave Radiation (OLR) as a predictor time series over the response regions.
 - This is effectively a measurement of rainfall.

Acknowledgements



Joint work with NSF NCAR
Atmospheric Chemistry Observations
& Modeling

References


- Buchholz, R., Hammerling, D., Worden, H., Deeter, M., Emmons, L., Edwards, D., & Monks, S. (2018). Links between carbon monoxide and climate indices for the southern hemisphere and tropical fire regions. *Journal of Geophysical Research: Atmospheres*, 123(17), 9786–9800.
- Daniels, W. S., Buchholz, R. R., Worden, H. M., Ahamad, F., & Hammerling, D. M. (2022). Interpretable models capture the complex relationship between climate indices and fire season intensity in maritime southeast asia. *Journal of Geophysical Research: Atmospheres*, 127(17), e2022JD036774.
- NOAA. (2021a). Climate prediction center-teleconnections: Antarctic oscillation.
- NOAA. (2021b). Ocean observations panel for climate-state of the ocean climate.

Thank you! Questions?


Appendix 1

ccf 1


NE Aus & Nino




SE Aus & Nino



NE Aus & DMI




SE Aus & DMI




Appendix 2

ccf 2


NE Aus & TSA




SE Aus & TSA



NE Aus & SAM



SE Aus & SAM




Appendix 3

pacf residuals

What if we select p by $\min(AIC)$?

PACF of Residuals for NE Aus VAR(1)



PACF of Residuals for SE Aus VAR(2)

