

MAE 3013 Mechanical and Aerospace Engineering Analysis

Chapter 19 Numerics in General

School of Mechanical and Aerospace Engineering
Oklahoma State University

Newton-Raphson Method

Numerics

Sec. 19.1

Sec. 19.2

Example

Sec. 19.3

Example


Sec. 19.4

Example

Sec. 19.5

Example

- Consider the function shown below. We want to find the location where the function crosses the x -axis.



- Starting from point x_0 , a tangent line is drawn to the curve at the point $f(x_0)$. The location where this line intersects the x -axis will be our next approximation of the root.

Newton-Raphson Method, cont.

Numerics

- The slope of the line can be given by:

$$f'(x_0) = \frac{f(x_0)}{x_0 - x_1}$$

solving for x_1 yields:

$$x_1 = x_0 - \frac{f(x_0)}{f'(x_0)}$$

Sec. 19.1

Sec. 19.2

Example

Sec. 19.3

Example

Sec. 19.4

Example

Sec. 19.5

Example

Newton-Raphson Method, cont.

Numerics

- The slope of the line can be given by:

$$f'(x_0) = \frac{f(x_0)}{x_0 - x_1}$$

solving for x_1 yields:

$$x_1 = x_0 - \frac{f(x_0)}{f'(x_0)}$$

- This can be generalize to:

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$$

Sec. 19.1

Sec. 19.2

Example

Sec. 19.3

Example

Sec. 19.4

Example

Sec. 19.5

Example

Example

Numerics

Sec. 19.1

Sec. 19.2

Example

Sec. 19.3

Example

Sec. 19.4

Example

Sec. 19.5

Example

Use the same example as before: $x - 2 \sin x = 0$.

Secant Method

Numerics

Sec. 19.1

Sec. 19.2

Example

Sec. 19.3

Example

Sec. 19.4

Example

Sec. 19.5

Example

What if the derivative of the function is unknown? In the secant method the difference formula is used to approximate the derivative:

$$f'(x_n) \approx \frac{f(x_n) - f(x_{n-1})}{x_n - x_{n-1}}$$

This approximation is substituted into the Newton-Raphson formula to obtain:

$$x_{n+1} = x_n - f(x_n) \frac{x_n - x_{n-1}}{f(x_n) - f(x_{n-1})}$$

Note we need two starting values x_n and x_{n-1} .

Example

Numerics

Sec. 19.1

Sec. 19.2

Example

Sec. 19.3

Example

Sec. 19.4

Example

Sec. 19.5

Example

Using the same equation as before: $x - 2 \sin x = 0$

Newton's Method for a Set of Equations

Numerics

What if we have an N-equation, N-unknown set of non-linear equations? The Newton-Raphson Method can be extended from:

$$x_1 = x_0 - \frac{f(x_0)}{f'(x_0)}$$

to:
$$\bar{x}_{new} = \bar{x}_{old} - [J(\bar{x}_{old})]^{-1} * \overline{f(\bar{x}_{old})}$$

The Jacobian
$$J(\bar{x}_{old}) = \begin{bmatrix} \frac{\partial f_1}{\partial x_1} & \frac{\partial f_1}{\partial x_2} & \vdots \\ \frac{\partial f_2}{\partial x_1} & \frac{\partial f_2}{\partial x_2} & \vdots \\ \vdots & \vdots & \ddots \end{bmatrix}$$
 is a matrix of partial deris.

Notice that we need to take the inverse of a matrix or do we?

Newton's Method for a Set of Equations

Numerics

$$\bar{x}_{new} = \bar{x}_{old} - [J(\bar{x}_{old})]^{-1} * \overline{f(\bar{x}_{old})}$$

Define: $\Delta x = [J(\bar{x}_{old})]^{-1} * \overline{f(\bar{x}_{old})}$

Which can be rearranged as: $[J(\bar{x}_{old})]\Delta x = \overline{f(\bar{x}_{old})}$

Which resembles $Ax = b$, and can be solved $[J(\bar{x}_{old})]\Delta x = \overline{f(\bar{x}_{old})}$

Procedure: given a vector of functions, $\overline{f(\bar{x})}$ and an initial guess: \bar{x}

Form the Jacobian: $J(\bar{x}) = \begin{bmatrix} \frac{\partial f_1}{\partial x_1} & \frac{\partial f_1}{\partial x_2} \\ \frac{\partial f_2}{\partial x_1} & \frac{\partial f_2}{\partial x_2} \\ \vdots & \vdots \end{bmatrix}$

and use GE to solve: $[J(\bar{x}_{old})]\Delta x = \overline{f(\bar{x}_{old})}$ for: Δx

and finally: $\bar{x}_{new} = \bar{x}_{old} - \Delta x$

Sec. 19.1

Sec. 19.2

Example

Sec. 19.3

Example

Sec. 19.4

Example

Sec. 19.5

Example

Example

Numerics

Example:

$$\overline{f(x,y)} = \begin{bmatrix} \sin(x) - 3y \\ e^y + \cos(x) \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

