



OceanX benchmark | Data geography

Three regional clusters · six acquisition groups · CMOMS supplied separately



A Gulf of Mexico



GLORYS12 / CMEMS


2011–2017 · daily · temperature, salinity, sea level

98–80°W, 18–31°N · 0–2000 m

Analysis: 98–82°W, 18–30°N

Q13–Q18, Q29–Q30 | 8 tasks

B South China Sea



MODIS-Aqua + NOAA Blended Sea Winds


Chlorophyll: 2003–2017 · monthly

Wind: 2003–2012 · monthly

104–122°E, 1–25°N

Q19–Q22, Q28 | 5 tasks

C East China Sea



OISST + GLORYS12 + ERA5

SST: 1982–2023 · daily

GLORYS / ERA5: 1993–2011 + 2023 · May–Nov

119–129°E, 24–35°N; inner box: 120–128°E, 25–34°N

Q23–Q24 | 2 tasks

CMOMS South China Sea basin, shelf and Pearl River estuary · Q01–Q12, Q25–Q27 (15 tasks)

CMOMS extents follow supplied grids and masks; no exact boundary is inferred. The purple star locates the Pearl River estuary only.

Solid outline: acquisition domain Dashed outline: analysis box Dotted lines: Gulf split / Q17 subregion

Source: benchmark manifest 2026-09-08-accessible-v1 · Boundaries: Natural Earth · Required coverage, not download-completion status