

DATE: 2020-09-12

ANNOUNCEMENTS:

0.0.1 Curvature

Definition 1 (Curvature). .

- A measure of how quickly the curve changes direction at that point
- The magnitude of the rate of change of the unit tangent vector with respect to arc length.

$$\kappa = \left| \frac{d\bar{T}}{ds} \right|$$

Problem: don't want to use the arclength.

$$\bar{T}(t) \rightarrow \frac{d\bar{T}}{dt}$$

$$\frac{d\bar{T}}{dt} \cdot \frac{dt}{ds} \rightarrow$$

$$\left| \frac{d\bar{T}}{dt} \right|$$

$$\frac{\frac{d\bar{T}}{ds}}{\frac{ds}{dt}}$$

$$\implies \kappa = \frac{\frac{d\bar{T}}{dt}}{\frac{ds}{dt}}$$

Arclength so far function:

$$s = \int_a^t |\bar{r}'(u)| du$$

$$\frac{ds}{dt} = |\bar{r}'(t)|$$

$$\left| \frac{ds}{dt} \right| = |\bar{r}'(t)|$$

Definition 2 (Curvature).

$$\kappa = \frac{\left| \bar{T}'(t) \right|}{\left| \bar{r}'(t) \right|}$$

Example 1 (Curvature).

$$\bar{r}(t) = \langle 3t + 2, t, t - 7 \rangle$$

, find κ

Solution 1 (Curvature).

$$r'(t) = \langle 3, 1, 1 \rangle$$

$$|r'(t)| = \sqrt{9 + 1 + 1} = \sqrt{11}$$

$$\bar{T} = \frac{1}{\sqrt{11}} \langle 3, 1, 1 \rangle$$

$$\bar{T}' = \frac{1}{\sqrt{11}} \langle 0, 0, 0 \rangle$$

$$\bar{T}' = \langle 0, 0, 0 \rangle$$

$$|\bar{T}'| = 0$$

$$\kappa = \frac{0}{\sqrt{11}} = 0$$

Example 2 (Curvature). let $\bar{r}(t) = \langle \sqrt{2}t, e^t, e^{-t} \rangle$, Find κ

Solution 2 (Curvature).

$$r'(t) = \langle \sqrt{2}, e^t, -e^{-t} \rangle$$

$$|r'(t)| = \sqrt{2 + e^{2t} + e^{-2t}}$$

$$|r'(t)| = \sqrt{e^{2t} + 2 + e^{-2t}}$$

$$|r'(t)| = \sqrt{(e^t + e^{-t})^2}$$

$$|r'(t)| = e^t + e^{-t}$$

$$\bar{T}(t) = \frac{1}{e^t + e^{-t}} \langle \sqrt{2}, e^t, -e^{-t} \rangle$$

$$\bar{T}(t) = \frac{e^t}{e^{2t} + 1} \langle \sqrt{2}, e^t, -e^{-t} \rangle$$

$$\bar{T}(t) = \frac{1}{e^{2t} + 1} \langle \sqrt{2}e^t, e^{2t}, -1 \rangle$$

$$\bar{T}'(t) = \frac{1}{e^{2t} + 1} \langle \sqrt{2}e^t, 2e^{2t}, 0 \rangle + -1(e^{2t} + 1)^{-2} 2e^{2t} \langle \sqrt{2}e^t, e^{2t}, -1 \rangle$$

$$\bar{T}'(t) = \left\langle \frac{\sqrt{2}e^t}{e^{2t} + 1}, \frac{2e^{2t}}{e^{2t} + 1}, 0 \right\rangle + \left\langle \frac{-2\sqrt{2}e^{3t}}{(e^{2t} + 1)^2}, \frac{-2e^{4t}}{(e^{2t} + 1)^2}, \frac{2e^{2t}}{(e^{2t} + 1)^2} \right\rangle$$

$$\bar{T}'(t) = \left\langle \frac{\sqrt{2}e^{3t} + \sqrt{2}e^t - 2\sqrt{2}e^{3t}}{(e^{2t} + 1)^2}, \dots \right\rangle$$

$$\bar{T}'(t) = \left\langle \frac{-\sqrt{2}e^{3t} + \sqrt{2}e^t}{(e^{2t} + 1)^2}, \dots \right\rangle$$

$$\bar{T}'(t) = \left\langle \frac{-\sqrt{2}e^{3t} + \sqrt{2}e^t}{(e^{2t} + 1)^2}, \frac{2e^{2t}}{(e^{2t} + 1)^2}, \frac{2e^{2t}}{(e^{2t} + 1)^2} \right\rangle$$

$$\bar{T}'(t) = \frac{\sqrt{2}e^t}{(e^{2t} + 1)^2} \langle 1 - e^{2t}, \sqrt{2}e^t, \sqrt{2}e^t \rangle$$

$$|c\bar{V}| = |c| |\bar{v}|$$

$$\left| \bar{T}'(t) \right| = \frac{\sqrt{2}e^t}{(e^{2t} + 1)^2} \sqrt{1 - 2e^{2t} + e^{4t} + 2e^{2t} + 2e^{2t}}$$

$$\left| \bar{T}'(t) \right| = \frac{\sqrt{2}e^t}{(e^{2t} + 1)^2} \sqrt{e^{4t} + 2e^{2t} + 1}$$

$$\left| \bar{T}'(t) \right| = \frac{\sqrt{2}e^t}{(e^{2t} + 1)^2} \sqrt{(e^{2t} + 1)^2}$$

$$\left| \bar{T}'(t) \right| = \frac{\sqrt{2}e^t}{e^{2t} + 1}$$

$$\kappa = \frac{\frac{\sqrt{2}e^t}{e^{2t}+1}}{e^t + e^{-t}}$$

$$\kappa = \frac{\frac{\sqrt{2}e^{2t}}{e^{2t}+1}}{e^{2t} + 1}$$

$$\kappa = \frac{\sqrt{2}e^{2t}}{(e^{2t} + 1)^2}$$

•

This example worked, but is a contrived textbook version of the problem. A better formula is still needed..