

Chapter 1

Unit 1: Differentiation

DATE: 2020-06-01

ANNOUNCEMENTS:

1.1 A. What is a derivative?

- Geometric interpretation
- Physical interpretation
- Importance of derivatives to all measurements
 - Science
 - Enineering
 - Econ
 - PolSci

1.2 B. How to differentiate anything

Example: $e^{x \arctan(x)}$

1.2.1 Geometric interpretation:

Find the tangent line to $y = f(x)$ at $P = (x_0, y_0)$

The tangent line is defined by the equation $y - y_0 = m(x - x_0)$

Point: $y_0 = f(x_0)$

Slope: $m = f'(x_0)$

Definition 1 (Derivative). $f'(x_0)$, the derivative of f at x_0 , is the slope of the tangent line to $y = f(x)$ at the point P

Taken for granted the geometric interpretation, we know the line through a point, but we want to analytically describe the tangent in a way that a machine could reproduce.

First grasp this with language.

$$y - y_0 = m(x - x_0)$$

point

$$y_0 = f(x_0).$$

slope:

$$m = f'(x_0).$$

Definition 2 (Derivative). $f'(x_0)$, the derivative of f at x_0 , is the slope of the tangent line to $y = f(x)$ at P .

Definition 3 (Tangent Line). Limit of the secant lines PQ as $Q \rightarrow P$ (P fixed)

1.2.2 Finding the Slope

To find the slope between P and Q , we denote the horizontal distance as Δx , and the height as Δf

The slope of the secant line is $\frac{\Delta f}{\Delta x}$. The slope of the tangent line is

$$\boxed{m = \lim_{\Delta x \rightarrow 0} \frac{\Delta f}{\Delta x}} \quad (1.1)$$

Slope of the tangent line

So we redefine the points P and Q to make this formula more usable,

$$P = (x_0, f(x_0)).$$

$$Q = (x_0 + \Delta x, f(x_0 + \Delta x)).$$

$$\boxed{f'(x_0) = \lim_{\Delta x \rightarrow 0} \frac{f(x_0 + \Delta x) - f(x_0)}{\Delta x}} \quad (1.2)$$

Limit definition of derivative

Example 1 (Example 1). $f(x) = \frac{1}{x}$

Solution 1 (Example 1).

$$\begin{aligned} \frac{\Delta f}{\Delta x} &= \frac{\frac{1}{x_0 + \Delta x} - \frac{1}{x_0}}{\Delta x} \\ &= \frac{1}{\Delta x} \left(\frac{x_0 - (x_0 + \Delta x)}{(x_0 + \Delta x)x_0} \right) \\ &= \frac{-1}{(x_0 + \Delta x)x_0} \rightarrow (\Delta x \rightarrow 0) \rightarrow \\ &= \frac{-1}{x_0^2} \end{aligned}$$

Note: The calculus part of calculus is easy, but it is made hard by the putting it in the context of everything learned up to that point.

Example 2 (Geometry problem). *Find areas of triangles enclosed by the axes and tangent to $y = \frac{1}{x}$*

Solution 2 (Geometry problem). $y = \frac{1}{x}$

Pick a point (x_0, y_0) on this curve, and we must find the base and the height. Find the formula for the tangent line:

$$y - y_0 = \frac{-1}{x_0^2}(x - x_0).$$

Find the x -intercept:

$$\begin{aligned} & (y = 0) \\ 0 - \frac{1}{x_0} &= \frac{-1}{x_0^2}(x - x_0) \\ &= \frac{-x}{x_0^2} + \frac{1}{x_0} \\ \frac{x}{x_0^2} &= \frac{2}{x_0} \\ &= 2x_0 \end{aligned}$$