Hindawi Publishing Corporation International Journal of Hypertension Volume 2012, Article ID 594361, 12 pages doi:10.1155/2012/594361

Review Article

Protective Role of the ACE2/Ang-(1-9) Axis in Cardiovascular Remodeling

María Paz Ocaranza and Jorge E. Jalil

Cardiovascular Diseases Division, School of Medicine, Faculty of Medicine, Pontificia Universidad Catolica de Chile, Chile

Correspondence should be addressed to María Paz Ocaranza, mocaran@med.puc.cl

Received 1 August 2011; Revised 5 October 2011; Accepted 9 October 2011

Academic Editor: Anderson J. Ferreira

Copyright © 2012 M. P. Ocaranza and J. E. Jalil. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Despite reduction in cardiovascular (CV) events and end-organ damage with the current pharmacologic strategies, CV disease remains the primary cause of death in the world. Pharmacological therapies based on the renin angiotensin system (RAS) blockade are used extensively for the treatment of hypertension, heart failure, and CV remodeling but in spite of their success the prevalence of end-organ damage and residual risk remain still high. Novel approaches must be discovered for a more effective treatment of residual CV remodeling and risk. The ACE2/Ang-(1–9) axis is a new and important target to counterbalance the vasoconstrictive/proliferative RAS axis. Ang-(1–9) is hydrolyzed slower than Ang-(1–7) and is able to bind the Ang II type 2 receptor. We review here the current experimental evidence suggesting that activation of the ACE2/Ang-(1–9) axis protects the heart and vessels (and possibly the kidney) from adverse cardiovascular remodeling in hypertension as well as in heart failure.

1. Introduction

All epidemiological studies show that the risk of adverse cardiovascular (CV) outcomes, such as stroke, myocardial infarction (MI), heart failure (HF), and kidney disease [1], increase progressively with increasing blood pressure (BP). On the other hand, clinical trials demonstrate that lowering BP reduces such risks [1]. All antihypertensive medications lower BP, but specific drug classes display effects beyond BP reduction (pleiotropic effects) that might contribute to cardiovascular risk reduction.

Remodeling of the cardiovascular structure occurs in response, not only to changes in BP and flow, but also to modifications in the neurohormonal environment, in which the rennin-angiotensin-aldosterone system (RAAS) exerts a most predominant influence [2].

The RAAS is a major regulator of BP [3, 4]. In addition, the RAAS has a role in the vascular response to injury and inflammation [4]. Chronic RAAS activation, through both angiotensin (Ang) II and aldosterone, leads to hypertension and perpetuates a cascade of proinflammatory, prothrombotic, and atherogenic effects associated with endorgan damage [3, 4]. Based on these facts, several drugs have

been developed that work by (a) reduction of Ang II levels, (b) inhibition of the Ang II type 1 receptor (AT1R), (c) blockade of the aldosterone receptor, and (d) renin receptor blockade [5, 6]. During the last 25 years several clinical trials have shown the benefits with these drugs that inhibit the RAAS with regard to BP reduction, regression of cardiac hypertrophy, prevention of kidney damage and reduction of cardiovascular morbidity reduction in hypertensive patients. Besides, with most of these RAAS blockers, quality of life as well as survival has been significantly improved in patients with heart failure. Consequently, the RAAS is currently a main therapeutic target in hypertension treatment [3, 4]. Aggressive BP control improves outcomes in patients with CV disease, stroke, and nephropathy and might have beneficial effects beyond BP lowering [7].

Despite the reduction of CV events and end-organ damage with the current pharmacologic strategies, CV disease remains the primary cause of death in the world, and more than 94,000 Americans annually experience progression to end-stage renal disease (ESRD). As population ages, the proportion affected by end-organ damage is expected to grow [8]. Thus, it is most relevant to find new molecules

in order to prevent and reduce hypertension as well as pathologic CV and kidney remodeling and dysfunction. In this regard, activation of the new ACE2/Ang-(1–9) pathway seems to counterbalance the damage due to the RAAS system activation.

We review here the current experimental evidence suggesting that activation of the ACE2/Ang-(1–9) pathway protects the heart and vessels (and possibly the kidney) from adverse cardiovascular remodeling in hypertension as well as in heart failure.

2. Angiotensin-Converting Enzyme 2

The discovery of angiotensin-converting enzyme homologue, ACE2, added further complexity to the main axis of the RAAS, in which Ang II and its forming enzyme ACE play major roles [9, 10]. A growing body of evidence points to a possible promising role for this new member of the RAAS by opposing to the effects of the main axis [11, 12]. ACE2 has dramatically changed the direction of cardiovascular and renal research in view of the pivotal role of this enzyme in the regulation of the RAAS [12, 13].

ACE2 is the newest member of the RAAS and shares approximately 40% similarity with the somatic form of ACE [9, 10]. ACE2 is a membrane-bound carboxypeptidase and its cellular and tissue distribution is different from that of ACE. While ACE is expressed in the endothelium throughout the vasculature, ACE2 is distributed in tissues with the most abundant expression in heart, kidney, lung, small intestine, and testis [14]. ACE2 can be released into the circulation and urine by shedding [15]. Tumor necrosis factoralpha-converting enzyme (TACE/ADAM17) is the sheddase responsible for the ectodomain cleavage and shedding of ACE2 [16].

However, normal ACE2 enzymatic activity in plasma is very low, probably due to the presence of an endogenous inhibitor [17–19]. ACE2 is different from ACE in both substrate specificity and functions [9, 20, 21]. ACE2 can form (a) Ang-(1-7) through hydrolysis of Ang II and (b) Ang-(1-9) through hydrolysis of Ang I. This last reaction is negligibly slow and is several hundred times slower than Ang II hydrolysis by ACE2 to form Ang(1-7)—a vasodepressor peptide counterbalancing the vasopressor effect of Ang II [20, 21]. Ang-(1-7) can be subsequently converted to Ang-(1-5) by ACE [9, 20] or by neutral endopeptidases [9], while Ang-(1–9) may be converted to Ang-(1–7) by ACE [9]. There is little evidence proving the existence of alternative hydrolysis of Ang-(1-9) to Ang II in some tissues. Drummer et al. [22] proved that homogenates of rat kidney, and in a lesser extent of lung, convert Ang-(1–9) to Ang II due to an ACEindependent aminopeptidase and N-like carboxypeptidase. Singh et al. [23] confirmed that the pathway Ang I-Ang-(1-9)-Ang II really exists in glomeruli of streptozotocininduced diabetes mellitus rats. Moreover, in human heart tissue the main products of Ang I degradation are both Ang-(1-9) and Ang II generated by heart chymase, ACE and a poorly identified carboxypeptidase A [24]. Although the data proving the existence of alternative pathways of Ang II

production, in clinical practice we can still block only ACE or AT1R.

ACE2 does not act on bradykinin metabolism and its activity is not inhibited by classic ACE inhibitors (ACEIs) [9]. Thus it has been proposed that ACE2 activity may counterbalance the effects of ACE by preventing the accumulation of Ang II in tissues where both ACE2 and ACE are expressed [25, 26]. ACE2 has several biological substrates and it is considered a multifunctional enzyme. Acting as a monocarboxypeptidase, it cleaves several other non-RAAS peptides which have roles in maintaining cardiovascular homeostasis such as (des-Arg9)-bradykinin, a member of the kininogen-kinin system [13]. (des-Arg9)-Bradykinin is formed from bradkinin by the action of carboxypeptidases and is an agonist of the B1 receptor, which is induced after tissue injury [27]. Bradykinin, a vasodilator which acts through the B2 receptor, is produced from its precursor kiningen by kallikrein and is degraded by ACE [13]. While, degradation of bradykinin by ACE is known to be an important aspect of BP regulation, the significance of the degradation of (des-Arg9)-bradykinin by ACE2 remains to be established. In addition to (des-Arg9)-bradykinin, ACE2 is also able to degrade apelin-13, a peptide proposed to cause vasoconstriction and known to regulate fluid homeostasis, and other non-RAAS peptides such as kinetensin, dynorphin A and neurotensin [20].

For a long time, Ang-(1–7) was thought to be devoid of biological activity, in spite of early reports on biological effects [28]. The importance of Ang-(1–7) was emphasized by the discovery of ACE2. Ang-(1–7) has been shown to release vasopressin as effectively as Ang II from neurohypophyseal explants [28] and to have actions opposing those of Ang II, namely vasodilation, antitrophic effects and implications of vasodilation caused by bradykinin [29, 30].

Several experiments suggest an important interaction between Ang-(1-7) and prostaglandin-bradykinin-nitric oxide (NO) systems. Ang-(1-7) binds to the Mas receptor (G protein-coupled receptor) which mediates vasodilating and antiproliferative actions of this peptide [31]. The Mas receptor can hetero-oligomerize with the AT1 receptor and acts as a physiological antagonist of Ang II [32]. Studies revealed that Ang-(1-7) activated endothelial nitric oxide synthase and NO production via Akt-dependent pathways [33]. Furthermore, Tallant et al. [34] showed that the presence of an antisense probe directed against Mas abolished the Ang-(1-7)-induced inhibition of protein synthesis in cardiomyocytes. This study also revealed that Ang-(1-7) decreased serum-stimulated ERK1/ERK2 mitogen-activated protein kinase activity, a response that was blocked by D-Ala 7-Ang-(1-7), an antagonist of Mas receptor.

Ang II binds with high affinity to two different receptor subtypes—AT1R and AT2R—which are members of the seven-transmembrane-domain G-protein-coupled receptors (GPCR) superfamily, through Gq and Gi, respectively [35]. Whereas the AT1R mediates most of the recognized actions of Ang II, it appears that the AT2R opposes, in part, to the effects mediated by the AT1R. As the AT2R is expressed in adult tissues in smaller amounts than the AT1R, the actions and cell signaling of AT2R have been less well characterized

than those of AT1R [36–38]. Current knowledge suggests that AT2R stimulation mediates vasodilation, antigrowth, proapoptotic and antiinflammatory effects [39, 40]. Hence, the AT2R can modulate cardiovascular remodeling as well as progression of atherosclerosis.

AT2R stimulation activates the NO-cGMP-dependent pathway [41]. This occurs either directly or indirectly through bradykinin or by increased endothelial NOS activity or expression. AT2R activation is associated with phosphorylation of JNK, PTPs, $I\kappa B\alpha$ (inhibitor of NF- κB), and the transcription factor ATF2, and dephosphorylation of p38MAPK, ERK1/2, and STAT3, which are linked to antiproliferative and antiinflammatory effects and apoptosis [38, 42–44]. AT2R may induce relaxation by opening large-conductance Ca^{2+} -activated K^+ channels (BKCa) [45] and by negative regulation of the vascular Rho A/Rho kinase pathway. The AT2R also enhances the activity of tyrosine phosphatases and vanadate-sensitive phosphatases MKP1 (DUSP1), SHP1 (PTPN6) and PP2A [46, 47].

3. Ang-(1-9)

There is little information in the literature with respect to Ang-(1–9) probably because this peptide was initially thought to be active only after conversion to Ang-(1–7). Ang-(1–9) can be generated by several carboxypeptidase-type enzymes including ACE2 or cathepsin A [48, 49]. Ang-(1–9) is present in healthy volunteers, in patients or in animals treated with ACE inhibitors (ACEIs) or AT1 receptor blockers (ARBs) [50–52], and its circulating levels are increased by pathological conditions (i.e., early after MI) [51]. However, very little is currently known about Ang-(1–9) biological effects [50, 53].

Initial studies showed that incubation of Chinese hamster ovary cells (CHO) with Ang-(1–9) potentiated the release of arachidonic acid by [Hyp³Tyr(Me)⁸]BK, elevated [Ca²]i and also resensitized the B2 receptor desensitized by BK [48]. At the same time, Jackman et al. [54] showed in CHO cells and in human pulmonary endothelial cells that Ang-(1–9) was significantly more active than Ang-(1–7) enhancing the effect of an ACE-resistant bradykinin analogue on the B2 receptor and that Ang-(1–9) also augmented arachidonic acid and NO release by kinin [54].

Some studies have suggested that Ang-(1–9) may be an endogenous inhibitor of ACE. Donoghue et al. [9] proposed that Ang-(1–9) is a competitive inhibitor of ACE because it is by itselfan ACE substrate. Under conditions of ACE inhibition, such as after long-term administration of an ACEI in rats, Ang-(1–9) levels increased in plasma and kidney [50, 53]. This increase in Ang-(1–9) steady-state levels could be due to decreased catabolism of Ang-(1–9) by ACE. Conversely, the increased levels of Ang-(1–9) could be due to increased production by ACE2 as a result of increased availability of Ang I substrate. These results indicate that an alternate pathway of Ang I metabolism by ACE2 exists and that this pathway may be amplified in the presence of ACE inhibitors.

To determine whether Ang-(1–9) is active per se or it becomes active only after conversion to Ang-(1-7), Chen et al. [55] examined the metabolism of Ang I, Ang-(1–9) and Ang-(1–7) in stably transfected CHO cells that express human ACE and human bradykinin B2 receptors coupled to green fluorescent protein (B2GFP). They found that Ang-(1-9) was hydrolyzed 18 times slower than Ang I and 30% slower than Ang-(1-7). Ang-(1-9) inhibited ACE and it resensitized the desensitized B2GFP receptors, independently of ACE inhibition [55]. This is reflected by release of arachidonic acid through a mechanism involving cross-talk between ACE and B2 receptors. They concluded that Ang-(1–9) enhanced bradykinin activity, probably by acting as an endogenous allosteric modifier of the ACE and B2 receptor complex. Therefore, when ACE inhibitors block conversion of Ang I, other enzymes like ACE2 can still release Ang I metabolites like Ang-(1-9) and enhance the efficacy of ACEIs.

Recently, Flores-Muñoz et al. [56] using radioligand binding assays observed that Ang-(1-9) is able to bind the Ang II type 2 receptor (AT2R) (pKi = 6.28 ± 0.1). They demonstrated that Ang-(1-9) and not Ang II, affected hypertrophy through the AT2R, as PD123319 (an AT2 receptor blocker) did not alter Ang II-mediated growth but did block the effects of Ang-(1-9). Despite having ~100fold lower affinity than Ang II for the AT2R [57], the selective AT2R activity of Ang-(1-9) is not inconsistent with current pharmacological models of G protein-coupled receptor signalling and activation. Indeed, the concept of functional selectivity, where individual receptor ligands have the capacity to selectively stabilize conformations which lead to distinct signalling outcomes [57-59], is supported by a previous study in which the critical amino acids and the mode of binding of ligands at the AT1R and AT2R were investigated [60]. While agonist activation of the AT1R was particularly sensitive to peptide modifications that disrupted contact points between Ang II and its receptor, substitutions within Ang II were far better tolerated by the AT2R [60]. The AT2R exists in a relaxed conformation and Ang II therefore binds to multiple indistinct contact points [60]. Since Ang-(1-9) contains the entire Ang II sequence plus a C-terminal histidine, these observations indicate that this difference may stabilize the AT2R in a conformation able to counteract hypertrophic signalling in cardiomyocytes. Flores-Muñoz et al. [56] did not observe functional competition between Ang II and Ang-(1-9) at the AT2R and they concluded that that Ang-(1-9) is able to antagonize Ang II signalling in cardiomyocytes selectively via the AT2R, highlighting that Ang-(1–9), along with Ang-(1–7), makes up part of the counter-regulatory arm of the RAS. What remains to be determined is the downstream signalling effects from Ang-(1–9). Preliminary studies indicate that the classical pathways via PKC translocation and ERK1/2 activation [61-63] are not different between Ang II-, Ang-(1-7)- and Ang-(1-9) stimulated cells. Since the downstream signalling from the AT2R is unclear at present, future studies will be required to establish these mechanisms.

4. Role of the ACE2/Ang-(1-9) Axis in Hypertension

Crackower et al. [64] were the first to test ACE2 as the gene underlying the blood pressure locus on the X chromosome. They showed reduced expression of renal ACE2 in the salt-sensitive Sabra hypertensive rat compared with the normotensive rat. Both spontaneously hypertensive rats (SHR) and spontaneously hypertensive stroke-prone rats (SHRSP) rats showed reduced renal ACE2 protein levels compared with the normotensive Sabra and Wistar Kyoto (WKY) strains. Two other groups confirmed some of these findings showing lower renal ACE2 mRNA, protein, and activity in the SHR compared to WKY rats [65, 66]. However, other investigators were unable to detect any difference in renal ACE2 mRNA, protein, and activity between adult hypertensive rats and their normotensive controls [67].

Rentzsch et al. [68], assessed in SHRSP (that display reduced ACE2 mRNA and protein expression compared with control animals in the kidney) the role of ACE2 in the pathogenesis of hypertension. They generated transgenic rats on a SHRSP genetic background expressing the human ACE2 in vascular smooth muscle cells by the use of the SM22 promoter, called SHRSP-ACE2. In these transgenic rats, vascular smooth muscle cells (VSMC) expression of human ACE2 was confirmed by RNase protection, real-time RT-PCR, and ACE2 activity assays. Transgene ACE2 expression leads to significantly increased circulating levels of Ang-(1-7), a prominent product of ACE2. Mean arterial blood pressure was reduced in SHRSP-ACE2 compared to SHRSP rats, and the vasoconstrictive response to intraarterial administration of Ang II was attenuated. The latter effect was abolished by previous administration of an ACE2 inhibitor. To evaluate the endothelial function in vivo, endotheliumdependent and endothelium-independent agents such as acetylcholine and sodium nitroprusside, respectively, were applied to the descending thoracic aorta and blood pressure was monitored. Endothelial function turned out to be significantly improved in SHRSP-ACE2 rats compared to SHRSP. These data indicate that vascular ACE2 overexpression in SHRSP reduces hypertension probably by local Ang II degradation and by improving endothelial function [68].

A target gene therapy strategy holds significant potential to translate the available fundamental research of ACE2 into therapeutics. In fact, initial animal experiments have been extremely encouraging. For example, in SHR, viral-mediated ACE2 overexpression in the heart decreased high BP [69]. This strategy also preserved cardiac function, as well as left ventricular wall motion and contractility, and attenuated left ventricular wall thinning induced by myocardial infarction [70]. ACE2 overexpression in the rostral ventrolateral medulla causes significant decreases in BP and heart rate (HR) [71].

Compared with ACEIs and ARBs, the targeting of ACE2 has the following potential therapeutic advantages, first, it degradates both Ang I to generate Ang-(1–9) and Ang II to generate Ang-(1–7). Thus, targeting ACE2 would not only produce the antihypertrophic peptide Ang-(1–9) [52]

and the vasoprotective/antiproliferative peptide Ang-(1–7) [72–74], but would also influence the vasoconstrictive/proliferative effects of the ACE/Ang II/AT1R axis [75]. Second, it is a multifunctional enzyme with many biologically active substrates [9, 20]. Third, unlike ARB/ACEI therapy, ACE2 is an endogenous regulator of the RAS [75]. Fourth, it is a part of the vasodilatory/antiproliferative axis of the RAS [20] and fifth, although treatment with ACEIs or ARBs indirectly increases ACE2 expression, direct activation of this enzyme could result in a better outcome in cardiovascular diseases [68, 75]. Thus, the activation of the ACE2 axis may be a novel therapeutic strategy in hypertension.

So far, all attention has been focused on Ang-(1–7), that opposes the pressor, proliferative, profibrotic, and prothrombotic actions mediated by Ang II [76]. Experimental and clinical studies have demonstrated a role for the Ang-(1–7)/ACE2/Mas axis in the evolution of hypertension, the regulation of cardiovascular and renal function, and the progression of cardiovascular and renal disease including diabetic nephropathy [77]. Additional evidence suggests that a reduction in the expression and activity of this vasode-pressor component may be a critical factor in mediating the progression of cardiovascular and renal disease. These findings support a role for the Ang-(1–7)/ACE2/Mas axis and, in particular, on its putative role as an ACE-Ang II-AT1 receptor counter-regulatory axis within the RAS [76, 77].

Recently, the alternative angiotensin peptide, Ang-(1–9) has shown relevant biological functions. Ocaranza et al. [51] have observed increased ACE2 activity and Ang-(1–9) plasma levels in MI and sham rats treated with enalapril for 8 weeks while circulating Ang-(1–7) levels did not change in any phase after MI [51] (Figure 1). These findings support the hypothesis that, in this second arm of the RAS, ACE2 through Ang-(1–9) instead of Ang-(1–7), could act as a counterregulator of the first arm, where ACE catalyzes the formation of Ang II.

Besides, in experimental hypertension (DOCA salt model) and in normotensive sham animals, RhoA/Rho-kinase inhibition (a signaling pathway that participates in pathological cardiovascular and renal remodeling and also in blood pressure regulation) by fasudil reduced BP and increased vascular and plasma ACE2 enzymatic activity. At the same time, fasudil reduced Ang II and increased Ang-(1-9) plasma levels (Figure 2) [78]. No modifications were observed here in Ang-(1-7) levels despite increased ACE2 levels with RhoA/Rho-kinase inhibition [78]. Thus, RhoA/Rho-kinase inhibition, by increasing eNOS and/or by reducing both ACE and Ang II, does not activate the Ang-(1–7) pathway. This novel effect of RhoA/Rho-kinase inhibition on both ACE2 expression and Ang-(1-9) levels might additionally contribute to the antihypertensive effects of RhoA/Rhokinase inhibitors. Besides, these results strongly suggest that in this experimental model, hypertension is more dependent on ACE2 and Ang-(1-9) levels than on ACE and Ang II levels. Therefore, this second RAAS axis through ACE2 and Ang-(1-9) could be an important target for the treatment of hypertension.




FIGURE 1: Plasma levels of Ang-(1–7), Ang-(1–9) and Ang II in rats with myocardial infarction treated with the ACE inhibitor enalapril (8 weeks). Increased plasma levels of Ang-(1–9) were observed in rats with myocardial infarction treated with the ACE inhibitor enalapril. Myocardial infarction was induced by coronary artery ligation. Data are presented as mean \pm SEM ($n=12/{\rm group}$). AMI: acute myocardial infarction, E: enalapril. *P<0.05 compared to both Sham and untreated myocardial infarction groups; **P<0.05 compared to both Sham and enalapril-treated myocardial infarction groups. (adapted with permission from [51]).

5. Role of ACE2/Ang-(1-9) Axis in Vascular Remodeling

The vascular wall is continuously exposed to hemodynamic forces such as the luminal pressure and shear stress. Changes in these forces, either physiological or pathological, lead to functional and/or structural alterations of the vascular wall [79]. Acute changes in hemodynamic forces can modify vessel diameter. Chronic changes in hemodynamic forces result in structural alterations of the vessel wall, indicated by changes in wall diameter and thickness. In addition, changes in vascular structure are not solely determined by hemodynamic forces [80], but also by inflammatory responses and changes in extracellular matrix components [81]. Structural changes of the medial layer of the vascular wall during hypertension are termed "eutrophic remodeling" [82] and subsequently translate to other vascular pathologies. This involves an inward encroachment of the arterial wall thereby, reducing the diameter of the lumen [83].

Several RAAS components are involved in neointimal formation after vascular endothelial damage [84]. In particular, Rakugi et al. [85] observed that vascular endothelial damage results in the induction of vascular ACE. Their results suggested that inhibition of vascular ACE might be critical in the prevention of restenosis after balloon injury. Patients with previously untreated essential hypertension and eutrophic inward remodeling appears to respond to antihypertensive medication. Reduction in BP with drugs that block the RAAS such as ACEIS [86–88] or ARBs [86, 87, 89] and calcium channel antagonists [90] are able to reverse the eutrophic inward remodeling [88].

The protein and mRNA of ACE2 are expressed in human coronary arteries and arterioles and the vasa vasorum of most organs [9, 91]. Recently, ACE2 expression has also been




FIGURE 2: Plasma levels of Ang-(1–7), Ang-(1–9) and Ang II in DOCA salt hypertensive rats treated with the Rho kinase inhibitor fasudil. Increased plasma levels of Ang-(1–9) were observed in DOCA salt hypertensive rats treated with the Rho kinase inhibitor fasudil. Fasudil (100 mh/kg/day) by gavage was administered during 3 weeks, starting on the third week after DOCA administration. Data are presented as mean \pm SEM (n = 8-11/group). DOCA: deoxycorticosterone, F: fasudil. *P < 0.05 compared to both Sham and untreated DOCA groups (adapted with permission from [78]).

observed in the large conduit arteries (aorta and carotid) in the HR [92]. ACE2 localizes preferentially in endothelial cells and arterial smooth muscle cells (SMCs) [9, 91]. As for the role of ACE2 in vascular remodeling, the effect of ACE2 on neointima formation has not yet been studied, but Ang-(1-7) infusion after balloon-catheter injury of the rat carotid artery reduced neointima formation [93]. This effect was probably mediated by its inhibition of vascular SMC proliferation [94]. In hypertensive animal models, ACE2 mRNA and protein were associated with immunoreactive Ang-(1–7) in the large conduit arteries of SHRs. Treatment with an ARB induced a fivefold increase in ACE2 mRNA and was associated with a significant increase in aortic Ang-(1–7) protein expression. This effect was associated with a decrease in aortic medial thickness, suggesting that this may be a protective mechanism in the prevention of cardiovascular events during hypertension [94]. Igase et al. [95] showed that ACE2 protein is expressed not only in the media of the carotid artery but also in the neointima of the ballooninjured carotid artery in SHR. The increase in ACE2 protein expression in the neointima following exposure of the rats to an ARB compared to vehicle was associated with a reduction in neointima thickness. These results lead to the hypothesis that there is a strong correlation between the increase in ACE2 protein in the injured carotid artery of SHR and vascular remodeling during blockade of Ang II receptors [95].

There is known the prothrombotic effect of Ang II [96, 97] and the antithrombotic action of Ang-(1–7) [98] in renovascular hypertensive rats. Thus, in this context, the question arises whether Ang-(1–9) effects are similar to Ang II or to Ang-(1–7) in in vivo conditions. Kramkowski et al. [99] described that Ang-(1–9) enhances electrically

stimulated thrombosis in rats and that this effect was abolished by losartan—an antagonist of the AT1 receptor. The prothrombotic activity of Ang-(1–9) was accompanied by the enhancement of ex vivo platelet aggregation and in vitro Ang-(1–9) increased platelet aggregation. However, there are some points in this paper that should be clarified. First, thrombus formation was initiated by electrical stimulation producing arterial injury that is unrelated to a clinical situation. Second, the prothrombotic effect of Ang-(1–9) was much weaker, to the prothrombotic action of Ang II [96, 97]. Third, Ang-(1–9) slightly increased platelet aggregation in in vitro conditions.

On the contrary Ocaranza et al. [78] showed that by inhibiting the RhoA/Rho-kinase pathway with fasudil, gene expression and enzymatic ACE activity and plasma levels of Ang II were reduced (Figure 2) and whereas aortic gene expression and ACE2 activity were importantly increased. Simultaneously, plasma levels of Ang-(1–9) (Figure 2), mRNA eNOS levels increased and the aortic overexpression of the remodeling promotion proteins TGF- β 1, PAI-1, and MCP-1 as well as the increased aortic NADPH oxidase activity and O²⁻ production were reduced, as a consequence of direct RhoA/Rho-kinase inhibition [100]. This novel effect of RhoA/Rho-kinase inhibition on ACE2 gene expression, enzymatic activity, and Ang-(1-9) levels might additionally contribute to its benefits in hypertension, atherosclerosis, and in cardiovascular and renal pathologic remodeling. This is the first observation concerning a pharmacologic ACE2 and Ang-(1-9) levels activator, both in normotensive and in hypertensive animals, one of the most interesting findings of that study (Figure 2). Additionally, in experimental hypertension, direct RhoA/Rho-kinase inhibition also normalizes overexpression of genes that promote vascular remodeling. Interestingly, the observed changes in ACE/ACE2 and in Ang-(1–9) levels were present only during fasudil treatment both in sham and in the DOCA hypertensive rats [78]. Thus, vascular remodeling could be more dependent on the tissue ACE2/Ang-(1-9) axis than on Ang-(1-7) levels in normotensive as well as in hypertensive rats.

In vessels, new members of the RAS have been detected, including ACE2, Ang-(1-7) and Mas. Vascular ACE2 is functionally active and generates Ang-(1-7) from Ang II. Ang-(1-7) is found in the endothelium and vascular wall [101-103] and immunohistochemical staining shows abundant presence in aortic perivascular adventitial tissue [104, 105]. Ang-(1-7), by binding to receptor Mas on endothelial cells, opposes Ang II actions by mediating vasodilation, growthinhibition, antiinflammatory responses, antiarrhythmogenic and antithrombotic effects [33, 68] through NOS-derived NO production, activation of protein tyrosine phosphatases, reduced MAPK activation and inhibition of NADPH oxidase-derived generation of reactive oxygen species (ROS) [106, 107]. Overexpression of ACE2 in the vascular wall of SHR is associated with improved endothelial function and attenuated development of hypertension [68]. Ang-(1-7)-Mas can hetero-oligomerize with AT1R, thereby inhibiting Ang II actions. The ACE2/Ang-(1-7)-Mas axis is now considered as a counter-regulatory system to the ACE-Ang II-AT1R axis in the vasculature [107], although some

evidence indicates that Ang-(1–7) may also promote fibrosis and inflammation in certain conditions [108, 109].

6. Role of the ACE2/Ang-(1-9) Axis in Cardiac Remodeling

After myocardial injury or in response to chronically increased hemodynamic load, cardiac mass increases as a result of cardiomyocyte hypertrophy and ventricular wall thickening. Initially these changes are compensatory mechanisms which help to maintain ejection performance and heart function. With continued hemodynamic overload the heart becomes dilated and its walls thinner, resulting in a geometry that contributes to systolic dysfunction by increasing wall stress [110]. At the cellular level, cardiac myocytes increase in size (hypertrophy), rearrange within the myocardial matrix (cell slippage), and die, to be replaced by fibrous tissue, which include fibroblasts and collagen. These changes are collectively referred to as "remodeling" [111]. Cardiac remodeling has been consistently associated with an impaired prognosis in patients with hypertension, MI and chronic heart failure (CHF) [112].

Despite recent advances in our understanding of the ACE2/Ang-(1-7)/axis, the functional role of ACE2 in the heart is somewhat controversial. Crackower et al. [64] originally reported a progressive reduction in LV contractile function in ACE2-null mice without significant changes in fibrosis, left ventricular and cardiac myocyte hypertrophy, or in mean arterial pressure [64]. Interestingly, whereas plasma and tissue levels of Ang II were increased, a decrease in blood pressure was only observed in 6-month-old male ACE-/homozygote mice but not in age-matched females or 3month-old males. Conversely, Gurley et al. [113] reported that ACE2 deletion enhanced the susceptibility to Ang IIinduced hypertension but had no effect on cardiac structure or function [113]. Huentelman et al. [114] showed that the ACE2 overexpression protects the heart from Ang II-induced hypertrophy and fibrosis. More recently, in SHR hypertensive rats Díez-Freire et al. by using lentiviral-based ACE2 gene transfer, attenuated cardiac fibrosis and hypertrophy [70] and also improved LV and remodeling after experimental MI [115]. Finally, Yamamoto et al. [116] reported that ACE2 deletion exacerbated pressure overload-induced cardiac dysfunction and remodeling that was associated with increased intracardiac Ang II levels and AT1R activation. The reasons for these discrepancies seem to be: (a) the genetic background of the mice used for ACE2 gene deletion [113], (b) global versus tissue-specific ACE2 manipulation, or (c) the cardiac responses were monitored under basal or pathophysiological conditions.

In MI Ocaranza et al. [51] observed that (a) circulating and LV enzymatic activities of ACE2 were downregulated in the long-term phase of LV dysfunction in rats, (b) these effects were prevented by the conventional ACE inhibitor enalapril, (c) plasma Ang-(1–9) levels were significantly increased when MI rats or sham-operated rats were treated with enalapril for 8 weeks but circulating Ang-(1–7) levels did not change at that time (Figure 1) [51]. Based on these




FIGURE 3: Signaling events and cellular effects induced by Ang II via AT1R and opposing effects of Ang-(1–9) acting through AT2R. Proposed Ang-(1–9)-dependent mechanisms that antagonize the cardiovascular remodeling effects of Ang II. ACE2 can directly cleave Ang I to form Ang-(1–9). This peptide activates the AT2R to initiate signaling pathways that antagonize AT1R-mediated tyrosine kinase cascades. In this simplified scenario, Ang-(1–9) increases SHP-1 tyrosine phosphatase activity to inactivate src-dependent signaling. AT2R activation also acts other pathways such as NO-AKT. AT1R: Ang II type 1 receptor; AT2R: Ang II type 2 receptor; ERK1/2: extracellular signal-regulated kinase 1/2; JAK: Janus-activated kinase; MAPK: mitogen-activated protein kinase; p38: p38 MAPK; PKC: protein kinase C; STAT: signal transducer and activator of transcription; NO: nitric oxide; SHP-1: protein tyrosine phosphatase SH2 domain-containing phosphatase 1; MEK: mitogen/ERK kinase. Solid arrows indicates activation broken arrows indicates inactivation.

findings, it was proposed in this model of HF, that Ang-(1-9) rather than Ang-(1-7) acts as a counterregulator of Ang II [51].

Recently, in MI rats randomized to receive either vehicle, the ACEI enalapril, or the ARB candesartan for 8 weeks, Ocaranza et al. [52] observed that both drugs prevented LVH and increased plasma Ang-(1-9) levels by several folds. Ang-(1–9) levels correlated negatively with different LVH markers with or without adjustment for BP reduction. This effect was specific as neither Ang-(1-7), Ang II nor bradykinins were correlated with LVH. Chronic administration of Ang-(1–9) to MI rats by osmotic minipumps versus vehicle for two weeks decreased plasma Ang II levels, inhibited ACE activity and also prevented cardiac myocyte hypertrophy. Because there are in vitro evidences that the incubation of Ang-(1–9) with ACE generates Ang-(1–7) [9], and Ang-(1– 7) negatively regulates hypertrophy [34, 117], the authors used the Ang-(1-7) receptor blocker A779 to investigate whether Ang-(1-7) could mediate the effects of Ang-(1-9). Even though A779 was bioactive, with significant increase in circulating Ang-(1–7) levels by 2.7 fold, this compound did not modify the Ang-(1-9)-dependent suppression of cardiac myocytes hypertrophy induced by MI [52]. In in vitro experiments with cardiac myocytes incubated with norepinephrine (10 μ M) or with IGF-1 (10 nM), Ang-(1–9) also prevented hypertrophy and this effect was not modified by the coincubation with Ang-(1–9) and A779 [52].

To further understand the role of Ang-(1-9) compared to Ang-(1–7) in cardiomyocyte hypertrophy, Flores-Muñoz et al. [56] studied Ang-(1-9) effects in rat neonatal H9c2 and in rabbit left ventricular cardiomyocytes. Cardiomyocyte hypertrophy was stimulated with Ang II or vasopressin, significantly increasing cell size by approximately 1.2-fold as well as stimulating expression of the hypertrophy gene markers atrial natriuretic peptide, brain natriuretic peptide, β -myosin heavy chain and myosin light chain (2- to 5fold). Both Ang-(1-9) and Ang-(1-7) were able to block hypertrophy induced by either agonist. The effects of Ang-1-9) were not inhibited by captopril, supporting previous evidence that Ang-(1-9) acts independently of Ang-(1-7). The authors investigated receptor signalling via angiotensin type 1 and type 2 receptors (AT1R, AT2R) and Mas. The AT1R antagonist losartan blocked Ang II-induced, but not vasopressin-induced, hypertrophy. Losartan did not block the antihypertrophic effects of Ang-(1-9), or Ang-(1-7) on vasopressin-stimulated cardiomyocytes. The Mas antagonist A779 efficiently blocked the antihypertrophic effects of Ang-(1–7), without affecting Ang-(1–9). Furthermore, Ang-(1–7) activity was also inhibited in the presence of the bradykinin type 2 receptor antagonist HOE140, without affecting Ang-(1–9). Moreover, Flores-Muñoz et al. [56] observed that the AT2R antagonist PD123,319 abolished the antihypertrophic effects of Ang-(1-9), without affecting Ang-(1-7), suggesting Ang-(1–9) signals via the AT2R. Radioligand binding assays demonstrated that Ang-(1-9) was able to bind the AT2R (pKi = 6.28 ± 0.1). The data indicate that ACE2/Ang-(1-9) axis, acting as a counterregulator of Ang II, is an effective, and possibly direct novel anticardiac hypertrophy axis.

7. Conclusions

Pharmacological treatments based on the RAS blockade are used extensively for the treatment of hypertension and CV remodeling. However, in spite of their success in pharmacological blockade of the RAS, the prevalence of end-organ damage has risen steadily in the last several decades. These observations indicate that novel and innovative approaches must be used in an attempt to promote a more effective treatment for the residual CV remodeling. In this environment, the ACE2/Ang-(1–9) axis is an important target, that is critical in tipping the balance of vasoconstrictive/proliferative to vasodilatory/antiproliferative axis of the RAS. Conceptually, the ACE2/Ang-(1-9)/AT2 axis balances the adverse effects of the ACE-Ang II-AT1 receptor axis (Figure 3). Accumulating evidence suggests that ACE2 expression and Ang-(1-9) levels are altered in diastolic and systolic dysfunction and remodeling and the activation of the ACE2/Ang-(1-9) axis protects the heart and vessels from cardiovascular remodeling. In conclusion, the noncanonical RAS arm has new biological effector Ang-(1–9) to counterregulate the classical RAS.

Acknowledgments

This work was supported by Fondo Nacional de Desarrollo Científico y Tecnológico (Fondecyt) Grants 1100874 and 1085208.

References

- [1] A. V. Chobanian, G. L. Bakris, H. R. Black et al., "Seventh report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure," *Hypertension*, vol. 42, no. 6, pp. 1206–1252, 2003.
- [2] H. D. Intengan and E. L. Schiffrin, "Structure and mechanical properties of resistance arteries in hypertension: role of adhesion molecules and extracellular matrix determinants," *Hypertension*, vol. 36, no. 3, pp. 312–318, 2000.
- [3] A. W. J. H. Dielis, M. Smid, H. M. H. Spronk et al., "The prothrombotic paradox of hypertension: role of the reninangiotensin and kallikrein-kinin systems," *Hypertension*, vol. 46, no. 6, pp. 1236–1242, 2005.
- [4] C. M. Ferrario, "Role of angiotensin II in cardiovascular disease therapeutic implications of more than a century of research," *Journal of the Renin-Angiotensin-Aldosterone System*, vol. 7, no. 1, pp. 3–14, 2006.
- [5] M. R. Weir, "Effects of renin-angiotensin system inhibition end-organ protection: can we do better?" *Clinical Therapeutics*, vol. 29, no. 9, pp. 1803–1824, 2007.
- [6] A. H. J. Danser, "(Pro)renin receptors: are they biologically relevant?" Current Opinion in Nephrology and Hypertension, vol. 18, no. 1, pp. 74–78, 2009.
- [7] M. R. Weir, "Targeting mechanisms of hypertensive vascular disease with dual calcium channel and renin-angiotensin

- system blockade," *Journal of Human Hypertension*, vol. 21, no. 10, pp. 770–779, 2007.
- [8] P. M. Kearney, M. Whelton, K. Reynolds, P. Muntner, P. K. Whelton, and J. He, "Global burden of hypertension: analysis of worldwide data," *The Lancet*, vol. 365, no. 9455, pp. 217–223, 2005.
- [9] M. Donoghue, F. Hsieh, E. Baronas et al., "A novel angiotensin-converting enzyme-related carboxypeptidase (ACE2) converts angiotensin I to angiotensin 1-9," *Circulation Research*, vol. 87, no. 5, pp. E1–E9, 2000.
- [10] S. R. Tipnis, N. M. Hooper, R. Hyde, E. Karran, G. Christie, and A. J. Turner, "A human homolog of angiotensin-converting enzyme: cloning and functional expression as a captopril-insensitive carboxypeptidase," *Journal of Biological Chemistry*, vol. 275, no. 43, pp. 33238–33243, 2000.
- [11] U. Danilczyk, U. Eriksson, M. A. Crackower, and J. M. Penninger, "A story of two ACEs," *Journal of Molecular Medicine*, vol. 81, no. 4, pp. 227–234, 2003.
- [12] G. Y. Oudit, M. A. Crackower, P. H. Backx, and J. M. Penninger, "The role of ACE2 in cardiovascular physiology," *Trends in Cardiovascular Medicine*, vol. 13, no. 3, pp. 93–101, 2003.
- [13] A. J. Turner, S. R. Tipnis, J. L. Guy, G. I. Rice, and N. M. Hooper, "ACEH/ACE2 is a novel mammalian metallocarboxypeptidase and a homologue of angiotensin-converting enzyme insensitive to ACE inhibitors," *Canadian Journal of Physiology and Pharmacology*, vol. 80, no. 4, pp. 346–353, 2002.
- [14] F. Gembardt, A. Sterner-Kock, H. Imboden et al., "Organspecific distribution of ACE2 mRNA and correlating peptidase activity in rodents," *Peptides*, vol. 26, no. 7, pp. 1270– 1277, 2005.
- [15] S. Epelman, K. Shrestha, R. W. Troughton et al., "Soluble angiotensin-converting enzyme 2 in human heart failure: relation with myocardial function and clinical outcomes," *Journal of Cardiac Failure*, vol. 15, no. 7, pp. 565–571, 2009.
- [16] D. W. Lambert, M. Yarski, F. J. Warner et al., "Tumor necrosis factor-α convertase (ADAM17) mediates regulated ectodomain shedding of the severe-acute respiratory syndromecoronavirus (SARS-CoV) receptor, angiotensin-converting enzyme-2 (ACE2)," *Journal of Biological Chemistry*, vol. 280, no. 34, pp. 30113–30119, 2005.
- [17] K. M. Elased, T. S. Cunha, S. B. Gurley, T. M. Coffman, and M. Morris, "New mass spectrometric assay for angiotensin-converting enzyme 2 activity," *Hypertension*, vol. 47, no. 5, pp. 1010–1017, 2006.
- [18] G. I. Rice, A. L. Jones, P. J. Grant, A. M. Carter, A. J. Turner, and N. M. Hooper, "Circulating activities of angiotensin-converting enzyme, its homolog, angiotensin-converting enzyme 2, and neprilysin in a family study," *Hypertension*, vol. 48, no. 5, pp. 914–920, 2006.
- [19] R. A. Lew, F. J. Warner, I. Hanchapola et al., "Angiotensin-converting enzyme 2 catalytic activity in human plasma is masked by an endogenous inhibitor," *Experimental Physiology*, vol. 93, no. 5, pp. 685–693, 2008.
- [20] C. Vickers, P. Hales, V. Kaushik et al., "Hydrolysis of biological peptides by human angiotensin-converting enzymerelated carboxypeptidase," *Journal of Biological Chemistry*, vol. 277, no. 17, pp. 14838–14843, 2002.
- [21] G. I. Rice, D. A. Thomas, P. J. Grant, A. J. Turner, and N. M. Hooper, "Evaluation of angiotensin-converting enzyme (ACE), its homologue ACE2 and neprilysin in angiotensin peptide metabolism," *Biochemical Journal*, vol. 383, no. 1, pp. 45–51, 2004.

- [22] O. H. Drummer, S. Kourtis, and H. Johnson, "Formation of angiotensin II and other angiotensin peptides from desleu 10-angiotensin I in rat lung and kidney," *Biochemical Pharmacology*, vol. 37, no. 22, pp. 4327–4333, 1988.
- [23] R. Singh, A. K. Singh, and D. J. Leehey, "A novel mechanism for angiotensin II formation in streptozotocin-diabetic rat glomeruli," *American Journal of Physiology*, vol. 288, no. 6, pp. F1183–F1190, 2005.
- [24] J. O. Kokkonen, J. Saarinen, and P. T. Kovanen, "Regulation of local angiotensin II formation in the human heart in the presence of interstitial fluid: inhibition of chymase by protease inhibitors of interstitial fluid and of angiotensin-converting enzyme by ANG-(1-9) formed by heart carboxypeptidase Alike activity," *Circulation*, vol. 95, no. 6, pp. 1455–1463, 1997.
- [25] M. Ye, J. Wysocki, P. Naaz, M. R. Salabat, M. S. LaPointe, and D. Batlle, "Increased ACE 2 and decreased ACE protein in renal tubules from diabetic mice: a renoprotective combination?" *Hypertension*, vol. 43, no. 5, pp. 1120–1125, 2004.
- [26] G. Rivière, A. Michaud, C. Breton et al., "Angiotensin-converting enzyme 2 (ACE2) and ACE activities display tissue-specific sensitivity to undernutrition-programmed hypertension in the adult rat," *Hypertension*, vol. 46, no. 5, pp. 1169–1174, 2005.
- [27] M. Kakoki, R. W. McGarrah, H. S. Kim, and O. Smithies, "Bradykinin B1 and B2 receptors both have protective roles in renal ischemia/reperfusion injury," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 104, no. 18, pp. 7576–7581, 2007.
- [28] M. T. Schiavone, R. A. S. Santos, K. B. Brosnihan, M. C. Khosla, and C. M. Ferrario, "Release of vasopressin from the rat hypothalamo-neurohypophysial system by angiotensin-(1-7) heptapeptide," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 85, no. 11, pp. 4095–4098, 1988.
- [29] R. A. Souza Dos Santos, K. T. Passaglio, J. B. Pesquero, M. Bader, and A. C. Simões E Silva, "Interactions between angiotensin-(1-7), kinins, and angiotensin II in kidney and blood vessels," *Hypertension*, vol. 38, no. 3, pp. 660–664, 2001.
- [30] L. G. Maia, M. C. Ramos, L. Fernandes, M. H.C. De Carvalho, M. J. Campagnole-Santos, and R. A.S. Dos Santos, "Angiotensin-(1-7) antagonist A-779 attenuates the potentiation of bradykinin by captopril in rats," *Journal of Cardiovascular Pharmacology*, vol. 43, no. 5, pp. 685–691, 2004.
- [31] R. A. S. Santos, A. C. Simoes e Silva, C. Maric et al., "Angiotensin-(1-7) is an endogenous ligand for the G protein-coupled receptor Mas," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 100, no. 14, pp. 8258–8263, 2003.
- [32] E. Kostenis, G. Milligan, A. Christopoulos et al., "G-protein-coupled receptor Mas is a physiological antagonist of the angiotensin II type 1 receptor," *Circulation*, vol. 111, no. 14, pp. 1806–1813, 2005.
- [33] W. O. Sampaio, R. A. S. Dos Santos, R. Faria-Silva, L. T. Da Mata Machado, E. L. Schiffrin, and R. M. Touyz, "Angiotensin-(1-7) through receptor Mas mediates endothelial nitric oxide synthase activation via Akt-dependent pathways," *Hypertension*, vol. 49, no. 1, pp. 185–192, 2007.
- [34] E. A. Tallant, C. M. Ferrario, and P. E. Gallagher, "Angiotensin-(1-7) inhibits growth of cardiac myocytes through activation of the mas receptor," *American Journal of Physiology*, vol. 289, no. 4, pp. H1560–H1566, 2005.
- [35] E. Kaschina and T. Unger, "Angiotensin AT1/AT2 receptors: regulation, signalling and function," *Blood Pressure*, vol. 12, no. 2, pp. 70–88, 2003.

- [36] M. Horiuchi, M. Akishita, and V. J. Dzau, "Recent progress in angiotensin II type 2 receptor research in the cardiovascular system," *Hypertension*, vol. 33, no. 2, pp. 613–621, 1999.
- [37] C. Savoia, F. Tabet, G. Yao, E. L. Schiffrin, and R. M. Touyz, "Negative regulation of RhoA/Rho kinase by angiotensin II type 2 receptor in vascular smooth muscle cells: role in angiotensin II-induced vasodilation in stroke-prone spontaneously hypertensive rats," *Journal of Hypertension*, vol. 23, no. 5, pp. 1037–1045, 2005.
- [38] C. Hu, A. Dandapat, J. Chen et al., "Over-expression of angiotensin II type 2 receptor (agtr2) reduces atherogenesis and modulates LOX-1, endothelial nitric oxide synthase and heme-oxygenase-1 expression," *Atherosclerosis*, vol. 199, no. 2, pp. 288–294, 2008.
- [39] R. M. Touyz and E. L. Schiffrin, "Signal transduction mechanisms mediating the physiological and pathophysiological actions of angiotensin II in vascular smooth muscle cells," *Pharmacological Reviews*, vol. 52, no. 4, pp. 639–672, 2000.
- [40] D. Henrion, N. Kubis, and B. I. Lévy, "Physiological and pathophysiological functions of the AT2 subtype receptor of angiotensin II from large arteries to the microcirculation," *Hypertension*, vol. 38, no. 5, pp. 1150–1157, 2001.
- [41] P. M. Abadir, R. M. Carey, and H. M. Siragy, "Angiotensin AT2 receptors directly stimulate renal nitric oxide in Bradykinin B2-receptor-null mice," *Hypertension*, vol. 42, no. 4 I, pp. 600–604, 2003.
- [42] C. Savoia, T. Ebrahimian, Y. He, J. P. Gratton, E. L. Schiffrin, and R. M. Touyz, "Angiotensin II/AT2 receptor-induced vasodilation in stroke-prone spontaneously hypertensive rats involves nitric oxide and cGMP-dependent protein kinase," *Journal of Hypertension*, vol. 24, no. 12, pp. 2417–2422, 2006.
- [43] A. Dandapat, C. P. Hu, J. Chen et al., "Over-expression of angiotensin II type 2 receptor (agtr2) decreases collagen accumulation in atherosclerotic plaque," *Biochemical and Biophysical Research Communications*, vol. 366, no. 4, pp. 871– 877, 2008.
- [44] P. Brassard, F. Amiri, and E. L. Schiffrin, "Combined angiotensin II type 1 and type 2 receptor blockade on vascular remodeling and matrix metalloproteinases in resistance arteries," *Hypertension*, vol. 46, no. 3, pp. 598–606, 2005.
- [45] C. Dimitropoulou, R. E. White, L. Fuchs, H. Zhang, J. D. Catravas, and G. O. Carrier, "Angiotensin II relaxes microvessels via the AT2 receptor and Ca2+-activated K+ (BKCa) channels," *Hypertension*, vol. 37, no. 2 I, pp. 301–307, 2001.
- [46] M. Horiuchi, W. Hayashida, T. Kambe, T. Yamada, and V. J. Dzau, "Angiotensin type 2 receptor dephosphorylates Bcl-2 by activating mitogen-activated protein kinase phosphatase-1 and induces apoptosis," *Journal of Biological Chemistry*, vol. 272, no. 30, pp. 19022–19026, 1997.
- [47] K. Bedecs, N. Elbaz, M. Sutren et al., "Angiotensin II type 2 receptors mediate inhibition of mitogen-activated protein kinase cascade and functional activation of SHP-1 tyrosine phosphatase," *Biochemical Journal*, vol. 325, no. 2, pp. 449–454, 1997.
- [48] E. G. Erdös, H. L. Jackman, V. Brovkovych, F. Tan, and P. A. Deddish, "Products of angiotensin I hydrolysis by human cardiac enzymes potentiate bradykinin," *Journal of Molecular and Cellular Cardiology*, vol. 34, no. 12, pp. 1569–1576, 2002.
- [49] F. J. Warner, A. I. Smith, N. M. Hooper, and A. J. Turner, "Angiotensin-converting enzyme-2: a molecular and cellular perspective," *Cellular and Molecular Life Sciences*, vol. 61, no. 21, pp. 2704–2713, 2004.

- [50] H. Johnson, S. Kourtis, J. Waters, and O. H. Drummer, "Radioimmunoassay for immunoreactive [des-Leu10]-angiotensin I," *Peptides*, vol. 10, no. 3, pp. 489–492, 1989.
- [51] M. P. Ocaranza, I. Godoy, J. E. Jalil et al., "Enalapril attenuates downregulation of angiotensin-converting enzyme 2 in the late phase of ventricular dysfunction in myocardial infarcted rat," *Hypertension*, vol. 48, no. 4, pp. 572–578, 2006.
- [52] M. P. Ocaranza, S. Lavandero, J. E. Jalil et al., "Angiotensin-(1-9) regulates cardiac hypertrophy in vivo and in vitro," *Journal of Hypertension*, vol. 28, no. 5, pp. 1054–1064, 2010.
- [53] O. H. Drummer, S. Kourtis, and H. Johnson, "Effect of chronic enalapril treatment on enzymes responsible for the catabolism of angiotensin I and formation of angiotensin II," *Biochemical Pharmacology*, vol. 39, no. 3, pp. 513–518, 1990.
- [54] H. L. Jackman, M. G. Massad, M. Sekosan et al., "Angiotensin 1-9 and 1-7 release in human heart role of cathepsin A," *Hypertension*, vol. 39, no. 5, pp. 976–981, 2002.
- [55] Z. Chen, F. Tan, E. G. Erdös, and P. A. Deddish, "Hydrolysis of angiotensin peptides by human angiotensin I-converting enzyme and the resensitization of B2 kinin receptors," *Hypertension*, vol. 46, no. 6, pp. 1368–1373, 2005.
- [56] M. Flores-Muñoz, N. J. Smith, C. Haggerty, G. Milligan, and S. A. Nicklin, "Angiotensin1-9 antagonises pro-hypertrophic signalling in cardiomyocytes via the angiotensin type 2 receptor," *Journal of Physiology*, vol. 589, no. 4, pp. 939–951, 2011.
- [57] S. Galandrin, G. Oligny-Longpré, and M. Bouvier, "The evasive nature of drug efficacy: implications for drug discovery," *Trends in Pharmacological Sciences*, vol. 28, no. 8, pp. 423–430, 2007.
- [58] T. Kenakin, "Functional selectivity through protean and biased agonism: who steers the ship?" *Molecular Pharmacology*, vol. 72, no. 6, pp. 1393–1401, 2007.
- [59] N. J. Smith, K. A. Bennett, and G. Milligan, "When simple agonism is not enough: emerging modalities of GPCR ligands," *Molecular and Cellular Endocrinology*, vol. 331, no. 2, pp. 241–247, 2011.
- [60] S. I. Miura and S. S. Karnik, "Angiotensin II type 1 and type 2 receptors bind angiotensin II through different types of epitope recognition," *Journal of Hypertension*, vol. 17, no. 3, pp. 397–404, 1999.
- [61] Y. Zou, I. Komuro, T. Yamazaki et al., "Protein kinase C, but not tyrosine kinases or Ras, plays a critical role in angiotensin II-induced activation of Raf-1 kinase and extracellular signalregulated protein kinases in cardiac myocytes," *Journal of Biological Chemistry*, vol. 271, no. 52, pp. 33592–33597, 1996.
- [62] K. Vijayan, E. L. Szotek, J. L. Martin, and A. M. Samarel, "Protein kinase C-α-induced hypertrophy of neonatal rat ventricular myocytes," *American Journal of Physiology*, vol. 287, no. 6, pp. H2777–H2789, 2004.
- [63] J. Pan, U. S. Singh, T. Takahashi et al., "PKC mediates cyclic stretch-induced cardiac hypertrophy through Rho family GTPases and mitogen-activated protein kinases in cardiomyocytes," *Journal of Cellular Physiology*, vol. 202, no. 2, pp. 536–553, 2005.
- [64] M. A. Crackower, R. Sarao, G. Y. Oudit et al., "Angiotensin-converting enzyme 2 is an essential regulator of heart function," *Nature*, vol. 417, no. 6891, pp. 822–828, 2002.
- [65] J. C. Zhong, D. Y. Huang, Y. M. Yang et al., "Upregulation of angiotensin-converting enzyme 2 by all-trans retinoic acid in spontaneously hypertensive rats," *Hypertension*, vol. 44, no. 6, pp. 907–912, 2004.
- [66] C. Tikellis, M. E. Cooper, K. Bialkowski et al., "Developmental expression of ACE2 in the SHR kidney: a role in


- hypertension?" *Kidney International*, vol. 70, no. 1, pp. 34–41, 2006.
- [67] I. Hamming, R. Kreutz, J. Sluimer et al., "Renal angiotensinconverting enzyme 2 is unaltered in experimental hypertension," *Journal of the American Society of Nephrology*, vol. 16, article TH–PO294-TH-PO294, 2005.
- [68] B. Rentzsch, M. Todiras, R. Iliescu et al., "Transgenic angiotensin-converting enzyme 2 overexpression in vessels of SHRSP rats reduces blood pressure and improves endothelial function," *Hypertension*, vol. 52, no. 5, pp. 967–973, 2008.
- [69] C. Díez-Freire, J. Vázquez, M. F. Correa De Adjounian et al., "ACE2 gene transfer attenuates hypertension-linked pathophysiological changes in the SHR," *Physiological Genomics*, vol. 27, no. 1, pp. 12–19, 2006.
- [70] S. Der Sarkissian, J. L. Grobe, L. Yuan et al., "Cardiac overexpression of angiotensin converting enzyme 2 protects the heart from ischemia-induced pathophysiology," *Hypertension*, vol. 51, no. 3, pp. 712–718, 2008.
- [71] M. Yamazato, Y. Yamazato, C. Sun, C. Diez-Freire, and M. K. Raizada, "Overexpression of angiotensin-converting enzyme 2 in the rostral ventrolateral medulla causes long-term decrease in blood pressure in the spontaneously hypertensive rats," *Hypertension*, vol. 49, no. 4, pp. 926–931, 2007.
- [72] K. B. Brosnihan, P. Li, and C. M. Ferrario, "Angiotensin-(1-7) dilates canine coronary arteries through kinins and nitric oxide," *Hypertension*, vol. 27, no. 3, pp. 523–528, 1996.
- [73] S. Y. Osei, R. S. Ahima, R. K. Minkes, J. P. Weaver, M. C. Khosla, and P. J. Kadowitz, "Differential responses to angiotensin-(1-7) in the feline mesenteric and hindquarters vascular beds," *European Journal of Pharmacology*, vol. 234, no. 1, pp. 35–42, 1993.
- [74] I. Porsti, A. T. Bara, R. Busse, and M. Hecker, "Release of nitric oxide by angiotensin-(1-7) from porcine coronary endothelium: implications for a novel angiotensin receptor," *British Journal of Pharmacology*, vol. 111, no. 3, pp. 652–654, 1994.
- [75] J. Zhong, R. Basu, D. Guo et al., "Angiotensin-converting enzyme 2 suppresses pathological hypertrophy, myocardial fibrosis, and cardiac dysfunction," *Circulation*, vol. 122, no. 7, pp. 717–728, 2010.
- [76] R. A. S. Santos, A. J. Ferreira, and A. C. Simões E Silva, "Recent advances in the angiotensin-converting enzyme 2angiotensin(1-7)-Mas axis," *Experimental Physiology*, vol. 93, no. 5, pp. 519–527, 2008.
- [77] C. M. Ferrarlo, "ACE2: more of Ang-(1-7) or less Ang II?" *Current Opinion in Nephrology and Hypertension*, vol. 20, no. 1, pp. 1–6, 2011.
- [78] M. P. Ocaranza, P. Rivera, U. Novoa et al., "Rho kinase inhibition activates the homologous angiotensin-converting enzyme-angiotensin-(1-9) axis in experimental hypertension," *Journal of Hypertension*, vol. 29, no. 4, pp. 706–715, 2011
- [79] S. Heeneman, J. C. Sluimer, and M. J. A. P. Daemen, "Angiotensin-converting enzyme and vascular remodeling," *Circulation Research*, vol. 101, no. 5, pp. 441–454, 2007.
- [80] W. J. G. Hacking, E. Vanbavel, and J. A. E. Spaan, "Shear stress is not sufficient to control growth of vascular networks: Aa model study," *American Journal of Physiology*, vol. 270, no. 1, pp. H364–H375, 1996.
- [81] G. Pasterkamp, Z. S. Galis, and D. P. V. De Kleijn, "Expansive arterial remodeling: location, location," *Arterioscle-rosis, Thrombosis, and Vascular Biology*, vol. 24, no. 4, pp. 650–657, 2004.

- [82] P. I. Korner, J. A. Angus, and M. J. Mulvany, "Vascular remodeling," *Hypertension*, vol. 29, no. 4, pp. 1065–1066, 1997.
- [83] J. G. R. De Mey, P. M. Schiffers, R. H. P. Hilgers, and M. M. W. Sanders, "Toward functional genomics of flow-induced outward remodeling of resistance arteries," *American Journal* of *Physiology*, vol. 288, no. 3, pp. H1022–H1027, 2005.
- [84] N. Iwai, M. Izumi, T. Inagami, and M. Kinoshita, "Induction of renin in medial smooth muscle cells by balloon injury," *Hypertension*, vol. 29, no. 4, pp. 1044–1050, 1997.
- [85] H. Rakugi, D. S. Wang, V. J. Dzau, and R. E. Pratt, "Potential importance of tissue angiotensin-converting enzyme inhibition in preventing neointima formation," *Circulation*, vol. 90, no. 1, pp. 449–455, 1994.
- [86] D. Rizzoni, E. Porteri, G. Bettoni et al., "Effects of candesartan cilexetil and enalapril on structural alterations and endothelial function in small resistance arteries of spontaneously hypertensive rats," *Journal of Cardiovascular Phar*macology, vol. 32, no. 5, pp. 798–806, 1998.
- [87] D. Rizzoni, E. Porteri, A. Piccoli et al., "Effects of losartan and enalapril on small artery structure in hypertensive rats," *Hypertension*, vol. 32, no. 2, pp. 305–310, 1998.
- [88] N. K. Thybo, N. Stephens, A. Cooper, C. Aalkjaer, A. M. Heagerty, and M. J. Mulvany, "Effect of antihypertensive treatment on small arteries of patients with previously untreated essential hypertension," *Hypertension*, vol. 25, no. 4 I, pp. 474–481, 1995.
- [89] E. L. Schiffrin, J. B. Park, H. D. Intengan, and R. M. Touyz, "Correction of arterial structure and endothelial dysfunction in human essential hypertension by the angiotensin receptor antagonist losartan," *Circulation*, vol. 101, no. 14, pp. 1653– 1659, 2000.
- [90] E. Porteri, D. Rizzoni, A. Piccoli et al., "Effects of hypotensive and non-hypotensive doses of manidipine on structure, responses to endothelin-1 and ICAM-1 production in mesenteric small resistance arteries of spontaneously hypertensive rats," *Blood Pressure*, vol. 7, no. 5-6, pp. 324–330, 1998.
- [91] I. Hamming, W. Timens, M. L. C. Bulthuis, A. T. Lely, G. J. Navis, and H. van Goor, "Tissue distribution of ACE2 protein, the functional receptor for SARS coronavirus. A first step in understanding SARS pathogenesis," *Journal of Pathology*, vol. 203, no. 2, pp. 631–637, 2004.
- [92] M. Igase, W. B. Strawn, P. E. Gallagher, R. L. Geary, and C. M. Ferrario, "Angiotensin II at1 receptors regulate ACE2 and angiotensin-(1-7) expression in the aorta of spontaneously hypertensive rats," *American Journal of Physiology*, vol. 289, no. 3, pp. H1013–H1019, 2005.
- [93] W. B. Strawn, C. M. Ferrario, and E. A. Tallant, "Angiotensin-(1-7) reduces smooth muscle growth after vascular injury," *Hypertension*, vol. 33, no. 1, pp. 207–211, 1999.
- [94] E. J. Freeman, G. M. Chisolm, C. M. Ferrario, and E. A. Tallant, "Angiotensin-(1-7) inhibits vascular smooth muscle cell growth," *Hypertension*, vol. 28, no. 1, pp. 104–108, 1996.
- [95] M. Igase, K. Kohara, T. Nagai, T. Miki, and C. M. Ferrario, "Increased expression of angiotensin converting enzyme 2 in conjunction with reduction of neointima by angiotensin II type 1 receptor blockade," *Hypertension Research*, vol. 31, no. 3, pp. 553–559, 2008.
- [96] A. Mogielnicki, E. Chabielska, R. Pawlak, J. Szemraj, and W. Buczko, "Angiotensin II enhances thrombosis development in renovascular hypertensive rats," *Thrombosis and Haemostasis*, vol. 93, no. 6, pp. 1069–1076, 2005.
- [97] M. Kamińska, A. Mogielnicki, A. Stankiewicz et al., "Angiotensin II via AT1 receptor accelerates arterial thrombosis

- in renovascular hypertensive rats," *Journal of Physiology and Pharmacology*, vol. 56, no. 4, pp. 571–585, 2005.
- [98] I. Kucharewicz, R. Pawlak, T. Matys, D. Pawlak, and W. Buczko, "Antithrombotic effect of captopril and losartan is mediated by angiotensin-(1-7)," *Hypertension*, vol. 40, no. 5, pp. 774–779, 2002.
- [99] K. Kramkowski, A. Mogielnicki, A. Leszczynska, and W. Buczko, "Angiotensin-(1-9), the product of angiotensin I conversion in platelets, enhances arterial thrombosis in rats," *Journal of Physiology and Pharmacology*, vol. 61, no. 3, pp. 317–324, 2010.
- [100] P. Rivera, M. P. Ocaranza, S. Lavandero, and J. E. Jalil, "Rho kinase activation and gene expression related to vascular remodeling in normotensive rats with high angiotensin Iconverting enzyme levels," *Hypertension*, vol. 50, no. 4, pp. 792–798, 2007.
- [101] V. J. Dzau, K. Bernstein, D. Celermajer et al., "Pathophysiologic and therapeutic importance of tissue ACE: a consensus report," *Cardiovascular Drugs and Therapy*, vol. 16, no. 2, pp. 149–160, 2002.
- [102] D. M. R. Silva, H. R. Vianna, S. F. Cortes, M. J. Campagnole-Santos, R. A. S. Santos, and V. S. Lemos, "Evidence for a new angiotensin-(1-7) receptor subtype in the aorta of Sprague-Dawley rats," *Peptides*, vol. 28, no. 3, pp. 702–707, 2007.
- [103] D. Weiss, K. E. Bernstein, S. Fuchs, J. Adams, A. Synetos, and W. R. Taylor, "Vascular wall ACE is not required for atherogenesis in ApoE-/- mice," *Atherosclerosis*, vol. 209, no. 2, pp. 352–358, 2010.
- [104] R. M. K. W. Lee, M. Bader, N. Alenina, R. A. Santos, Y. J. Gao, and C. Lu, "Mas receptors in modulating relaxation induced by perivascular adipose tissue," *Life Sciences*, vol. 89, no. 13-14, pp. 467–472, 2011.
- [105] R. M. K. W. Lee, C. Lu, L. Y. Su, and Y. J. Gao, "Endothelium-dependent relaxation factor released by perivascular adipose tissue," *Journal of Hypertension*, vol. 27, no. 4, pp. 782–790, 2009.
- [106] W. O. Sampaio, C. H. De Castro, R. A. S. Santos, E. L. Schiffrin, and R. M. Touyz, "Angiotensin-(1-7) counterregulates angiotensin II signaling in human endothelial cells," *Hypertension*, vol. 50, no. 6, pp. 1093–1098, 2007.
- [107] L. A. Rabelo, N. Alenina, and M. Bader, "ACE2-angiotensin-(1-7)-Mas axis and oxidative stress in cardiovascular disease," *Hypertension Research*, vol. 34, no. 2, pp. 154–160, 2011.
- [108] V. Esteban, S. Heringer-Walther, A. Sterner-Kock et al., "Angiotensin-(1-7) and the G protein-coupled receptor Mas are key players in renal inflammation," *PLoS ONE*, vol. 4, no. 4, Article ID e5406, 2009.
- [109] E. Velkoska, R. G. Dean, K. Griggs, L. Burchill, and L. M. Burrell, "Angiotensin-(1-7) infusion is associated with increased blood pressure and adverse cardiac remodelling in rats with subtotal nephrectomy," *Clinical Science*, vol. 120, no. 8, pp. 335–345, 2011.
- [110] M. J. Davies, "A macro and micro view of coronary vascular insult in ischemic heart disease," *Circulation*, vol. 82, no. 3, pp. II38–II46, 1990.
- [111] L. H. Opie, P. J. Commerford, B. J. Gersh, and M. A. Pfeffer, "Controversies in ventricular remodelling," *The Lancet*, vol. 367, no. 9507, pp. 356–367, 2006.
- [112] J. N. Cohn, R. Ferrari, and N. Sharpe, "Cardiac remodeling-concepts and clinical implications: a consensus paper from an International Forum on Cardiac Remodeling," *Journal of the American College of Cardiology*, vol. 35, no. 3, pp. 569–582, 2000.


- [113] S. B. Gurley, A. Allred, T. H. Le et al., "Altered blood pressure responses and normal cardiac phenotype in ACE2-null mice," *Journal of Clinical Investigation*, vol. 116, no. 8, pp. 2218–2225, 2006.
- [114] M. J. Huentelman, J. L. Grobe, J. Vazquez et al., "Protection from angiotensin II-induced cardiac hypertrophy and fibrosis by systemic lentiviral delivery of ACE2 in rats," *Experimental Physiology*, vol. 90, no. 5, pp. 783–790, 2005.
- [115] C. Díez-Freire, J. Vázquez, M. F. Correa De Adjounian et al., "ACE2 gene transfer attenuates hypertension-linked pathophysiological changes in the SHR," *Physiological Genomics*, vol. 27, no. 1, pp. 12–19, 2006.
- [116] K. Yamamoto, M. Ohishi, T. Katsuya et al., "Deletion of angiotensin-converting enzyme 2 accelerates pressure overload-induced cardiac dysfunction by increasing local angiotensin II," *Hypertension*, vol. 47, no. 4, pp. 718–726, 2006.
- [117] J. L. Grobe, A. P. Mecca, M. Lingis et al., "Prevention of angiotensin II-induced cardiac remodeling by angiotensin-(1-7)," *American Journal of Physiology*, vol. 292, no. 2, pp. H736–H742, 2007.

















Submit your manuscripts at http://www.hindawi.com





