

上例: \mathbb{R}^n 中, $\text{cpt} \Leftrightarrow \text{seq. cpt.} \Leftrightarrow \text{closed \& bounded}$

But it is not true in general metric space (eg: l^∞)

Today: fix it for general metric space.

closed $\xrightarrow{\text{stronger}}$ complete (i.e. every Cauchy seq. conv.)

bdd $\xrightarrow{\text{stronger}}$ totally bdd.




Def Totally bounded.

X a metric space.

$E \subseteq X$ 称为 ttl. bdd. if:

$\forall \varepsilon > 0$, 都 \exists 一个 finite cover of E by balls of radius ε

ε :-



即: $\exists x_1, \dots, x_n \in E$ s.t

$$E \subseteq \bigcup_{i=1}^n B_\varepsilon(x_i)$$

In any MS, ttl. bdd. \Rightarrow bdd

In \mathbb{R}^d , ttl. bdd. \Leftrightarrow bdd A closed \Leftrightarrow complete

eg: the unit ball in l^∞ is not ttl. bdd.

Thm X be a M.S.

for $E \subseteq X$, TFAE:

- (1) E cpt.
- (2) E seq. cpt.
- (3) E complete & ttl. bdd.

(Rmk 1: \mathbb{R} generally, $\text{cpt} \Leftrightarrow$
 $\text{seq. cpt.} \Leftrightarrow \text{complete \& ttl. bdd.}$)

(Rmk 2 (exercise): if X complete, then $E \subseteq X$ is closed iff it is complete.)

(Rmk 3: 对于 general topological space, 并不 true)

Pf. $1 \Rightarrow 2$: Already done

$2 \Rightarrow 3$ seq. cpt. implies complete & ttl. bdd.

Pf of ttl. bdd.

Fix $\varepsilon > 0$

Pick $p_1 \in E$

Given p_1, \dots, p_{n-1} :

if $E \subseteq \bigcup_{i=1}^{n-1} B_\varepsilon(p_i)$, stop

otherwise pick $p_n \in E$ s.t. $p_n \notin \bigcup_{i=1}^{n-1} B_\varepsilon(p_i)$

Claim This process must stop.

if not, we will get seq $(p_n)_{n=1}^\infty$ with

$$d(p_i, p_j) > \varepsilon \quad \forall i \neq j$$

Then it is not Cauchy thus not conv.

So eventually will have $E \subseteq \bigcup_{i=1}^n B_\varepsilon(p_i)$ conflicting seq. cpt.

Pf of complete

Let (x_n) be a Cauchy seq. in E

Since E seq. cpt., \exists subseq. $(x_{n_k}) \rightarrow p$ for some $p \in E$

Claim $(x_n) \rightarrow p$ also.

Given $\varepsilon > 0$

Pick N s.t. $\forall n, m \geq N, d(x_n, x_m) < \frac{\varepsilon}{2}$

Pick $n_k \geq N$ s.t. $d(x_{n_k}, p) < \frac{\varepsilon}{2}$

$\Rightarrow \forall m \geq N, \underline{d(x_m, p) \leq d(x_m, x_{n_k}) + d(x_{n_k}, p) < \varepsilon}$

$3 \Rightarrow 2$ ttl. bdd. & complete implies seq. cpt.

Let (x_n) be a seq. in E

WTS: (x_n) has conv. subseq.

Fix a finite cover of E by balls of radius 1

One of them must have infly many terms of (x_n)

Call it B_1

Given B_1, \dots, B_{n-1} st. there are infly many m
with $x_m \in \bigcap_{i=1}^{n-1} B_i$

Let radius $(B_i) = \frac{1}{2^{i-1}}$

Find a finite cover of E by balls of radius $\frac{1}{2^{n-1}}$

By pigeonhole principle, one of them, call it B_n ,
 \exists infly many m s.t. $x_m \in \bigcap_{i=1}^n B_i$

Now pick n_1 with $x_{n_1} \in B_1$,
 n_2 with $x_{n_2} \in B_1 \cap B_2$ with $n_2 > n_1$,
 \vdots
 n_k with $x_{n_k} \in \bigcap_{i=1}^k B_k$ with $n_k > n_{k-1}$

For $\forall k_1, k_2 \geq k$,

$$x_{k_1}, x_{k_2} \in B_k \text{ so } d(x_{k_1}, x_{k_2}) \leq 2 \cdot \frac{1}{2^{k-1}} = \frac{1}{2^{k-2}}$$


So (x_{n_k}) is Cauchy, by completeness it conv.

2 \Rightarrow 1 Seq. cpt. implies cpt.

Intermediate Lemma (Leasbegue covering thm)

Let $\{U_\alpha\}$ be an open cover of any subset of a cpt set.

$\Rightarrow \exists \varepsilon > 0$ s.t. $(\forall p \in E, \exists \alpha \text{ s.t. } \underline{B}_\varepsilon(p) \subseteq U_\alpha$



Rmk: 表达了一件事, 即如果 E cpt, 那么 E 的 open cover 中存在 U_α 和 E 边缘及其他 U_β 边缘都有一定的 uniformly large distance

ex $[0, 1] \subseteq \mathbb{R}$ is cpt. $\{U_\alpha\} = \{(-1, \frac{2}{3}), (\frac{1}{3}, 2)\}$
 $(\overline{U_1} \cap \overline{U_2}) = \{(\frac{1}{3}, \frac{2}{3})\}$ Take $\varepsilon = \frac{1}{6}$, works.

Pf of Lebesgue covering thm:

suppose not then $\forall n \in \mathbb{N}$, $\exists p_n$ s.t. $\forall \alpha$, $B_{\frac{1}{n}}(p_n) \not\subseteq U_\alpha$

$\Rightarrow (p_n)$ has a conv. subseq. $(p_{n_k}) \rightarrow p \in E$ for some p


As $p \in U_{\alpha_0}$ for some α_0 , U_{α_0} open $\Rightarrow \exists \delta$ s.t. $B_\delta(p) \subseteq U_{\alpha_0}$

而 (p_{n_k}) conv. Take k_0 s.t. $d(p, p_{n_k}) < \frac{\delta}{2}$ whenever $k \geq k_0$

$\Rightarrow \delta - \frac{1}{n_k} < \frac{\delta}{2}$

$\Rightarrow B_{\frac{1}{n_k}}(p_{n_k}) \subseteq B_\delta(p) \subseteq U_{\alpha_0}$, contradicts

□



然否回到 2 \Rightarrow 1 的 pf:

Let $\{U_\alpha\}$ be an open cover of E

By lemma, $\exists \varepsilon > 0$ s.t. $\forall p, \exists \alpha$ with $B_\varepsilon(p) \subseteq U_\alpha$

Since seq. cpt \Rightarrow ttl. bdd

$\exists p_1, \dots, p_N$ s.t. $E \subseteq \bigcup_{i=1}^N B_\varepsilon(p_i)$

And for each p_i , $\exists \alpha_i$ s.t. $B_\varepsilon(p_i) \subseteq U_{\alpha_i}$

$\Rightarrow E \subseteq \bigcup_{i=1}^N U_{\alpha_i}$ □

