

Problem A: Let $F : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be satisfy

$$F(tx) = tF(x)$$

for all all positive real numbers t and all $x \in \mathbb{R}^n$. Assume F is differentiable at the origin. Show F is linear.

Pf Construct $r_0(h) = F(0+h) - F(0) - Df(0)h = \underline{F(h) - Df(0)h}$

Then that for $\underline{t \in \mathbb{R}_{\geq 0}}$

$$\underline{r_0(th) = F(th) - Df(0)(th) = tF(h) - tDf(0)h = t r_0(h)}$$

Claim: $\forall h \in \mathbb{R}^n, r_0(h) = 0$

(pf) Suppose for contradiction that for some $h_0 \in \mathbb{R}^n, r_0(h_0) \neq 0$

$$\text{Then } \frac{\|r_0(h_0)\|}{\|h_0\|} = c \text{ for some } c > 0$$

By homogeneity, for any $t > 0$, $\frac{\|r_0(th_0)\|}{\|th_0\|} = \frac{t\|r_0(h_0)\|}{t\|h_0\|} = \frac{\|r_0(h_0)\|}{\|h_0\|} = \underline{c}$,

then for $t \rightarrow 0, \|th_0\| \rightarrow 0$ while $\frac{\|r_0(h_0)\|}{\|h_0\|} = \underline{c}$,

contradicting that $\lim_{\|h\| \rightarrow 0} \left\| \frac{r_0(h)}{h} \right\| = 0$

This proves the claim.

So $\forall h \in \mathbb{R}^n, \underline{F(h) = Df(0)h}$ where $Df(0) \in \text{Hom}(\mathbb{R}^n, \mathbb{R}^m)$

Thus F is a linear transformation.

□

Problem B: Let $A \subset \mathbb{R}^n$ be open and $f: A \rightarrow \mathbb{R}^m$. Suppose that the partial derivatives $\frac{\partial f_i}{\partial x_j}$ ($1 \leq i \leq m, 1 \leq j \leq n$) exist and are bounded on A . Show that f is continuous on A .

Pf Claim $f: A \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^m = \begin{pmatrix} f_1: \mathbb{R}^n \rightarrow \mathbb{R} \\ \vdots \\ f_m: \mathbb{R}^n \rightarrow \mathbb{R} \end{pmatrix}$ is continuous

at $x_0 \in A$ iff $\forall i \in \{1, \dots, m\}$, f_i is continuous at x_0 .

This directly follows from $\|f(x) - f(x_0)\|_2 = \sqrt{\sum_i (f_i(x) - f_i(x_0))^2}$
 (if $\forall x \in B_\delta(x_0)$ we have $f(x) \in B_\varepsilon(f(x_0))$, then $\forall f_i, f_i(x) \in B_\varepsilon(f_i(x_0))$
 if for all $i, \forall x \in B_\delta(x_0)$ we have $f_i(x) \in B_{\frac{\varepsilon}{\sqrt{m}}}(f_i(x_0)) \Rightarrow f(x) \in B_\varepsilon(f(x_0))$)

There WLOG we can set $m=1$

Assume $\frac{\partial f}{\partial x_j}$ ($1 \leq j \leq n$) exist (for all $x \in A$) and bounded

WTS: f is continuous on A .

Let $\varepsilon > 0$.

let $x = x_0 + h$ where $h \in \mathbb{R}^n$


Then $h = \begin{pmatrix} h_1 \\ \vdots \\ h_n \end{pmatrix}$ for some $h_1, \dots, h_n \in \mathbb{R}$

let $p_0 = x_0$

$p_1 = p_0 + h_1 e_1$

\vdots

$p_n = p_{n-1} + h_n e_n = \underline{x_0 + h}$



For each $i = 1, \dots, n$, let $\varphi_i: [0, h_i] \rightarrow \mathbb{R}$

map $s \mapsto f(p_{i-1} + s e_i)$

Then $\forall s_0 \in (0, h_i), \frac{d}{ds} \Big|_{s=s_0} \varphi_i(s) = \frac{d}{ds} \Big|_{s=s_0} f(p_{i-1} + s e_i) = \frac{\partial}{\partial x_i} f(p_{i-1} + s e_i)$

Since all partials exist on A and bounded,
all φ_i are differentiable on $(0, h_i)$,

So by MVT, $\forall i, \varphi_i(h_i) - \varphi_i(0) = \left(\frac{\partial}{\partial x_i} f(p_{i-1} + s_i e_i) \right) \cdot h_i$

for some $s_i \in (0, h_i)$, we write $p_{i-1} + s_i e_i$ as q_i

$$\begin{aligned} \text{Then } |f(x+h) - f(x)| &= \left| \sum_{i=1}^n (f(p_i) - f(p_{i-1})) \right| = \left| \sum_{i=1}^n (\varphi_i(h_i) - \varphi_i(0)) \right| \\ &= \left| \sum_{i=1}^n \left(\frac{\partial}{\partial x_i} f(q_i) \right) h_i \right| \end{aligned}$$

Since all partials are bounded by some $M \in \mathbb{R}$ and
 $|h_i| \leq \|h\| = \|x - x_0\|$

Thus we have:

$$\underline{|f(x) - f(x_0)| \leq n M \|x - x_0\|}$$

This implies that f is Lipschitz on A , thus
 (uniformly) continuous (by hw 3). \square

Problem C: Let $f: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be defined by the equation:


$$f(r, \theta) = (r \cos \theta, r \sin \theta).$$

- (1) Calculate Df and $\det Df$.
- (2) Let $S = [1, 2] \times [0, \pi/2]$. Find $f(S)$ and sketch it.
- (3) Show that f is a homeomorphism from S on $f(S)$ and compute the inverse function f^{-1} .
- (4) Compute Df^{-1} and $\det Df^{-1}$.
- (5) What relation can you find between Df and Df^{-1} ?

$$(1) Df = \begin{pmatrix} \frac{\partial f_1}{\partial r} & \frac{\partial f_1}{\partial \theta} \\ \frac{\partial f_2}{\partial r} & \frac{\partial f_2}{\partial \theta} \end{pmatrix} = \begin{pmatrix} \cos \theta & -r \sin \theta \\ \sin \theta & r \cos \theta \end{pmatrix}$$

$$\det Df = r \cos^2 \theta + r \sin^2 \theta = r$$

$$(2) f(S) = \{(x, y) : 1 \leq x^2 + y^2 \leq 4 \text{ and } x, y \geq 0\}$$



(3) Claim 1. $f: S \rightarrow f(S)$ is bijective

Pf it is surjective since we take $f(S)$ to be in place of codomain

Now prove injectivity: suppose $f(r_1, \theta_1) = f(r_2, \theta_2)$

$$\text{then } r_1 \cos \theta_1 = r_2 \cos \theta_2$$

$$r_1 \sin \theta_1 = r_2 \sin \theta_2$$

$$\Rightarrow r_1^2 \cos^2 \theta_1 + r_1^2 \sin^2 \theta_1 = r_2^2 \cos^2 \theta_2 + r_2^2 \sin^2 \theta_2 \Rightarrow r_1^2 = r_2^2 \Rightarrow r_1 = r_2 \text{ since } r_1, r_2 > 0$$

$$\Rightarrow \cos \theta_1 = \cos \theta_2 \Rightarrow \theta_1 = \theta_2 \text{ since } \theta_1, \theta_2 \in (0, \frac{\pi}{2})$$

Claim ②. f is continuous.

Since $f(r, \theta) = \begin{pmatrix} f_1(r, \theta) = r \cos \theta \\ f_2(r, \theta) = r \sin \theta \end{pmatrix}$ where f_1, f_2 are all continuous functions, f is always continuous.

Claim ③ f^{-1} is continuous

$$\text{let } f(r, \theta) = \begin{pmatrix} x \\ y \end{pmatrix} \Rightarrow x = r \cos \theta, y = r \sin \theta \Rightarrow x^2 + y^2 = r^2$$

$$\Rightarrow r = \sqrt{x^2 + y^2} \text{ since } r > 0;$$

$$\text{and } \tan \theta = \frac{y}{x} \Rightarrow \theta = \tan^{-1}\left(\frac{y}{x}\right) \text{ since } x, y > 0, \theta \in [0, \frac{\pi}{2}]$$

So $f^{-1}: f(S) \rightarrow S$

sending $\begin{pmatrix} x \\ y \end{pmatrix} \mapsto \begin{pmatrix} \sqrt{x^2 + y^2} \\ \tan^{-1}(\frac{y}{x}) \end{pmatrix}$, which is continuous since f_1^{-1}, f_2^{-1} are continuous.

Claim ①②③ proves that f is a homeomorphism.

$$(4) \frac{\partial}{\partial x} r = \frac{x}{\sqrt{x^2 + y^2}}, \frac{\partial}{\partial y} r = \frac{y}{\sqrt{x^2 + y^2}}, \frac{\partial}{\partial x} \theta = \frac{\partial}{\partial x} \arctan\left(\frac{y}{x}\right) = \frac{-y}{x^2 + y^2},$$

$$\Rightarrow Df^{-1}(x, y) = \begin{pmatrix} \frac{x}{\sqrt{x^2 + y^2}} & \frac{y}{\sqrt{x^2 + y^2}} \\ \frac{-y}{x^2 + y^2} & \frac{x}{x^2 + y^2} \end{pmatrix} = \begin{pmatrix} \cos \theta & \sin \theta \\ -\frac{\sin \theta}{r} & \frac{\cos \theta}{r} \end{pmatrix} \quad \frac{\partial}{\partial y} \theta = \frac{x}{x^2 + y^2}$$

$$\det Df^{-1} = \frac{x^2}{r^3} + \frac{y^2}{r^3} = \frac{r^2}{r^3} = \frac{1}{r} = \frac{1}{\sqrt{x^2 + y^2}}$$

$$(5) Df \cdot Df^{-1} = \begin{pmatrix} \cos \theta & \sin \theta \\ \sin \theta & r \cos \theta \end{pmatrix} \begin{pmatrix} \cos \theta & \sin \theta \\ -\frac{\sin \theta}{r} & \frac{\cos \theta}{r} \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \Rightarrow (Df)(Df^{-1}) = I_2$$

Problem D: Give an example of a function $F: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ such that, at the origin, all directions derivatives exist and are zero, but F is not differentiable at the origin.

Consider $F\left(\begin{pmatrix} x \\ y \end{pmatrix}\right) = \begin{pmatrix} \frac{x^2 y}{x^2 + y^2} \\ \frac{x^2 y}{x^2 + y^2} \end{pmatrix}$ for $\begin{pmatrix} x \\ y \end{pmatrix} \neq \begin{pmatrix} 0 \\ 0 \end{pmatrix}$ and $F\left(\begin{pmatrix} 0 \\ 0 \end{pmatrix}\right) = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$

$$D_x F(0, 0) = \lim_{t \rightarrow 0} \frac{F\left(\begin{pmatrix} 0+t \\ 0 \end{pmatrix}\right) - F\left(\begin{pmatrix} 0 \\ 0 \end{pmatrix}\right)}{t} = \lim_{t \rightarrow 0} (0, 0) = (0, 0)$$

Similarly $D_{e_2} F(0,0) = \lim_{t \rightarrow 0} (0,0) = (0,0)$

Since $\forall u \in \mathbb{R}^2$, u is a linear comb of e_1, e_2 and $D_u F(0)$ is linear in u , all directional derivatives exist and are 0 at origin:

$$\underline{D_u f(0)} = D_{u_1 e_1 + u_2 e_2} f(0) = u_1 D_{e_1} f(0) + u_2 D_{e_2} f(0) = \underline{(0,0)}$$

The Jacobian matrix $J_f(0) = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$

$$\lim_{\|(x,y)\| \rightarrow 0} \frac{f(x,y) - f(0,0) - J_f(0) \begin{pmatrix} x \\ y \end{pmatrix}}{\sqrt{x^2 + y^2}} = \lim_{\|(x,y)\| \rightarrow 0} \underbrace{\left(\frac{\frac{\pi y}{(x^2+y^2)^{\frac{3}{2}}}}{\frac{x^2 y}{(x^2+y^2)^{\frac{3}{2}}}} \right)}$$

Consider the sequence $(x_n, y_n) = (\frac{1}{n}, \frac{1}{n})_{n \in \mathbb{N}}$
this sequence converge to 0 by norm

But $\lim_{n \rightarrow \infty} \frac{\pi n^2 y_n}{(x_n^2 + y_n^2)^{\frac{3}{2}}} = \lim_{n \rightarrow \infty} \frac{1}{n^3} \frac{2\sqrt{2}}{n^3} = \underline{\frac{\sqrt{2}}{4} \neq 0}$

Hence the Jacobian matrix is not the derivative of f at 0,
which suffices to indicate that f is not differentiable at 0

Problem E: Define $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ by setting $f(0) = 0$ and

$$f(x, y) = xy \frac{x^2 - y^2}{x^2 + y^2}.$$

- (1) Show that $\frac{\partial f}{\partial x}$ and $\frac{\partial f}{\partial y}$ exist at 0.
- (2) Compute $\frac{\partial f}{\partial x}$ and $\frac{\partial f}{\partial y}$ for $(x, y) \neq 0$.
- (3) Show that $f \in C^1(\mathbb{R}^2)$.
- (4) Show that

$$\frac{\partial}{\partial x} \frac{\partial f}{\partial y} \quad \text{and} \quad \frac{\partial}{\partial y} \frac{\partial f}{\partial x}$$

exist everywhere on \mathbb{R}^2 , but they are not equal at $(x, y) = 0$.

□

$$(1) \frac{\partial f}{\partial x}(0) = \lim_{t \rightarrow 0} \frac{f\left(\begin{smallmatrix} t \\ 0 \end{smallmatrix}\right) - f\left(\begin{smallmatrix} 0 \\ 0 \end{smallmatrix}\right)}{t} = \lim_{t \rightarrow 0} \frac{0}{t} = 0$$

$$\frac{\partial f}{\partial y}(0) = \lim_{t \rightarrow 0} \frac{f\left(\begin{smallmatrix} 0 \\ t \end{smallmatrix}\right) - f\left(\begin{smallmatrix} 0 \\ 0 \end{smallmatrix}\right)}{t} = \lim_{t \rightarrow 0} \frac{0}{t} = 0$$

(2) Chain product and quotient rule of differentiation holds for partial derivatives.

$$\begin{aligned} \text{PF } \frac{\partial}{\partial x}(u(x,y)v(x,y)) &= \lim_{t \rightarrow 0} \frac{u(x+t,y)v(x+t,y) - u(x,y)v(x,y)}{t} \\ &= \lim_{t \rightarrow 0} \frac{u(x+t,y)(v(x+t,y) - v(x,y))}{t} + \frac{v(x,y)(u(x+t,y) - u(x,y))}{t} \\ &= u(x,y) \frac{\partial v(x,y)}{\partial x} + v(x,y) \frac{\partial u(x,y)}{\partial x} \end{aligned}$$

Quotient rule follows from the product rule.

Now we use the rules for calculation:

$$\text{for } \begin{pmatrix} x \\ y \end{pmatrix} \neq 0, \frac{\partial}{\partial x} f(x,y) = \frac{x^2 - y^2}{x^2 + y^2} \frac{\partial}{\partial x} (xy) + xy \frac{\partial}{\partial x} \left(\frac{x^2 - y^2}{x^2 + y^2} \right)$$

$$\text{where } \frac{\partial}{\partial x} \left(\frac{x^2 - y^2}{x^2 + y^2} \right) = \frac{(x^2 + y^2)2x - (x^2 - y^2)2x}{(x^2 + y^2)^2} = \frac{4xy^2}{(x^2 + y^2)^2}$$

$$\Rightarrow \frac{\partial}{\partial x} f(x,y) = y \frac{x^2 - y^2}{x^2 + y^2} + \frac{4x^2 y^3}{(x^2 + y^2)^2}$$

$$\frac{\partial}{\partial y} f(x,y) = \frac{x^2 - y^2}{x^2 + y^2} \frac{\partial}{\partial y} (xy) + xy \frac{\partial}{\partial y} \left(\frac{x^2 - y^2}{x^2 + y^2} \right)$$

$$= x \frac{x^2 - y^2}{x^2 + y^2} + x \frac{-4x^3 y^2}{(x^2 + y^2)^2}$$

(3) Since we have shown that for all $(x,y) \in \mathbb{R}^2$, all partials at (x,y) exist, it suffices to show that $\forall (x,y) \in \mathbb{R}^2$, all partials are continuous, in order to show that $f \in C^1(\mathbb{R}^2)$. And since any directional derivative $D_u f(x_0)$ is linear in u , it suffices to show that $\forall (x,y) \in \mathbb{R}^2$, $\frac{\partial}{\partial x} f(x,y)$ and $\frac{\partial}{\partial y} f(x,y)$ are continuous. Let $(x,y) \in \mathbb{R}^2$

Since $\frac{\partial}{\partial x} f(x,y)$ and $\frac{\partial}{\partial y} f(x,y)$ are rational functions (thus ctn.) except at $x=0$, we only need to show that $\frac{\partial}{\partial x} f(x,y)$, $\frac{\partial}{\partial y} f(x,y)$ are ctn. at $x=0$.

$$\lim_{(x,y) \rightarrow (0,0)} \frac{\partial}{\partial x} f(x,y) = \lim_{(x,y) \rightarrow (0,0)} y \frac{x^2 - y^2}{x^2 + y^2} + y \frac{4xy^2}{x^4 + 2x^2y^2 + y^4}$$

the expression is bounded by $|3y|$, so its limit when $(x,y) \rightarrow 0$ is 0.

$$\text{Similarly, } \lim_{(x,y) \rightarrow (0,0)} \frac{\partial}{\partial y} f(x,y) = \lim_{(x,y) \rightarrow (0,0)} x \frac{x^2 - y^2}{x^2 + y^2} + x \frac{-4x^3y^2}{x^4 + 2x^2y^2 + y^4}$$

the expression is bounded by $|3x|$, so its limit when $(x,y) \rightarrow 0$ is 0.

Notice that \mathbb{R}^2 has no isolated pt., so f is ctn. at origin and thus ctn. since it is rational elsewhere

(4) Let $(x,y) \in \mathbb{R}^2$

$$\frac{\partial}{\partial x} \frac{\partial}{\partial y} f(x,y) = \frac{\partial}{\partial x} \left(x \frac{x^2 - y^2}{x^2 + y^2} + x \frac{-4x^3y^2}{(x^2 + y^2)^2} \right) = \frac{x^6 + 9x^4y^2 - 9x^2y^4 - y^6}{(x^2 + y^2)^3}$$

$$\frac{\partial}{\partial y} \frac{\partial}{\partial x} f(x,y) = \frac{\partial}{\partial y} \left(y \frac{x^2 - y^2}{x^2 + y^2} + \frac{4xy^3}{(x^2 + y^2)^2} \right) = \frac{x^6 + 9x^4y^2 - 9x^2y^4 - y^6}{(x^2 + y^2)^3}$$

So $\frac{\partial^2}{\partial x \partial y}$ and $\frac{\partial^2}{\partial y \partial x}$ exists everywhere and equal except on the origin

$$\text{On the origin: } \frac{\partial}{\partial x} \left(\frac{\partial}{\partial y} f(0,0) \right) = \lim_{t \rightarrow 0} \frac{\frac{\partial}{\partial y} f(t,0) - \frac{\partial}{\partial y} f(0,0)}{t} = \lim_{t \rightarrow 0} \frac{t-0}{t} = 1$$

$$\text{but } \frac{\partial}{\partial y} \left(\frac{\partial}{\partial x} f(0,0) \right) = \lim_{t \rightarrow 0} \frac{\frac{\partial}{\partial x} f(0,t) - \frac{\partial}{\partial x} f(0,0)}{t} = \lim_{t \rightarrow 0} \frac{-t-0}{t} = -1$$

Bonus: Recall that an ultrametric space is a metric space where one has the following stronger than usual form of the triangle inequality:

$$d(x, z) \leq \max(d(x, y), d(y, z)).$$

- (1) Show that, in an ultrametric space, open balls are closed.
- (2) Show that, in an ultrametric space, if two balls intersect, one of the two must be contained in the other.
- (3) Show that, in an ultrametric space, every point of a ball is the center of the ball. That is, if $y \in B_r(x)$, then $B_r(x) = B_r(y)$.
- (4) Let G be a connected weighted undirected graph. (The weighting is the assignment of a positive number to each edge). Let $V(G)$ be the set of vertices.

Given a path in the graph (a sequence of adjacent edges), define the length of the path to be the largest weight of an edge crossed by the path.

Given $v, w \in V(G)$, define $d(v, w)$ to be the smallest length of a path from v to w .

Show that d is an ultrametric on $V(G)$.

- (5) Show that any finite ultrametric arises as in the previous part.

Just for fun (don't hand in): Imagine you have an electric car, and you live in a country that provides free charging stations, and you're not in a hurry. Why might you end up thinking about an ultrametric?

(1) Let (X, d) be an ultra-metric space.

Suppose $B = \{x : d(x, c) < r\}$ is an open ball in X centered at $c \in X$

let $x \in X \setminus B \Rightarrow d(x, c) \geq r$

Consider $B_r(z)$: let $a \in B_r(z)$, then $d(a, z) < r$

By ultrametric, $d(z, c) \leq \max\{d(a, c), d(a, z)\}$

And since $d(z, c) \geq r \Rightarrow \max\{d(c, a), d(a, z)\} \geq r$

We already know that $d(a, z) < r$

Therefore $d(c, a) \geq r \Rightarrow a \in X \setminus B_r(c)$

$\Rightarrow B_r(z) \subseteq X \setminus B_r(c)$

Since z is arbitrary, this proves that $X \setminus B_r(c)$ is open

$\Rightarrow B_r(c)$ is closed

Then we can conclude that every open ball is also closed in X .

(2) let (X, d) be an ultrametric space

Let $B_r(x), B_s(y) \subseteq X$ be two open balls with $B_r(x) \cap B_s(y) \neq \emptyset$

We only need to consider the case when $x \neq y$ since if $x = y$

WLOG suppose $r \leq s$. then one ball must contain the other one.

let $a \in B_r(x)$, $z \in B_r(x) \cap B_s(y)$

$\Rightarrow d(z, x) < r, d(z, y) < r$

$\Rightarrow d(a, y) \leq \max\{d(a, z), d(z, y)\} < r \Rightarrow$ $a \in B_s(y)$

$\Rightarrow B_r(x) \subseteq B_s(y)$


□

(3) This directly follows from (2):

let $y \in B_r(x) \Rightarrow B_r(y) \cap B_r(x) \neq \emptyset$

$\Rightarrow B_r(y) \subseteq B_r(x)$ and $B_r(x) \subseteq B_r(y)$

$\Rightarrow B_r(y) = B_r(x)$



(4) Positivity follows from the definition of the graph and $\forall x, y \in V(G), d(x, y) = d(y, x)$

since the graph is undirected (every path commutes)

So it suffices to show d is an ultrametric by showing the ultra-triangular property

Let $x, y, z \in V(G)$ st. there is at least one path from x to z and z to y
 We write the weight of an edge e as $w(e)$ and the smallest weight of an edge cross a path p as $L(p)$

Case 1: the smallest-length path between x, y , say p_{xy} , goes through z .

Then $p_{xy} = p_{xz} \cup p_{zy}$ where p_{xz} is a path between x, z and p_{zy} is a path between z, y

$$\text{Then } d(x, y) = L(p_{xy}) = \max\{L(p_{xz}), L(p_{zy})\}$$

$$\text{and } d(x, z) = \min\{L(p): \text{path through } x, z\}$$

$$d(z, y) = \min\{L(p): \text{path through } z, y\}$$

$$\text{so } L(p_{xz}) \leq d(x, z), L(p_{zy}) \leq d(z, y)$$

$$\text{Thus } d(x, y) = L(p_{xy}) \leq \max\{d(x, z), d(z, y)\}$$

Case 2: the smallest-length path between x, y , say p_{xy} , does not go through z .

Take path p_{xz}, p_{zy} st. $L(p_{xz}) = d(x, z)$, $L(p_{zy}) = d(z, y)$

Then let $p_{xy}' = p_{xz} \cup p_{zy}$, we have $L(p_{xy}') = \max\{L(p_{xz}), L(p_{zy})\}$

$$\text{Since } d(x, y) = L(p_{xy}) \Rightarrow L(p_{xy}) \geq L(p_{xy}')$$

$$\Rightarrow \underline{d(x, y) = L(p_{xy})} > \max\{L(p_{xz}), L(p_{zy})\} \\ = \max\{d(x, z), d(z, y)\}$$

In both case the ultra-triangular ineq. holds true

This finishes the proof that d is an ultrametric on $V(G)$

(5) Let (X, d_X) be a finite ultrametric field with $\#|X| = C$

WTS: we can construct a graph $G = (X, E(G))$ endowed with

metric d_G in (4), st (X, d_X) is isometrically embedded into (G, d_G)

Construction: for each $v, w \in X$, add an edge $e(v, w)$ to $E(G)$
with $w(e(v, w)) = d_u(v, w)$

Then the graph will be a complete C -graph

By d_u , $\forall x \in X$, $d(v, w) \leq \max\{d(v, x), d(x, w)\}$

So every path P through v, w has $L(P) \geq w(e(v, w))$

So $d_g(v, w) = w(e(v, w)) = d_u(v, w)$