

hw3

Question 1. Use the complex form to find the Fourier series of $f(x) = e^x$, $-L < x < L$.

Solution. $e^x = \sum_{n=-\infty}^{\infty} (-1)^n \frac{L + in\pi}{L^2 + n^2\pi^2} (\sinh L) \exp\left(\frac{in\pi}{L}x\right)$

$$\text{Fourier series: } F(x) = \sum_{n \in \mathbb{Z}} a_n e^{\frac{in\pi}{L}x}, \quad -L < x < L$$

$$\text{where } a_n = \frac{1}{2L} \int_{-L}^L e^x e^{-\frac{in\pi}{L}x} dx$$

$$= \frac{1}{2L} \int_{-L}^L e^{(1 - \frac{in\pi}{L})x} dx$$

$$= \frac{1}{2L} \frac{1}{1 - \frac{in\pi}{L}} \int_{-L}^L e^{(1 - \frac{in\pi}{L})x} d\left(\left(1 - \frac{in\pi}{L}\right)x\right)$$

$$= \frac{1}{2L - 2in\pi} (e^{L - in\pi} - e^{-L + in\pi}) \quad (\text{note: } e^{in\pi} = (-1)^n)$$

$$= \frac{(-1)^n}{2L - 2in\pi} (e^L - e^{-L}) = \frac{(-1)^n}{2L - 2in\pi} 2 \sinh(L)$$

$$= \frac{(-1)^n}{L - in\pi} \sinh(L)$$

$$= (-1)^n \frac{L + in\pi}{L^2 + n^2\pi^2} \sinh(L)$$

$$\text{Therefore } f_L(x) = \sum_{n \in \mathbb{Z}} (-1)^n \frac{L + in\pi}{L^2 + n^2\pi^2} \sinh(L) \exp\left(\frac{in\pi}{L}x\right)$$

Question 2. Derive the following formula

- 1) Let $0 < r < 1$, $f(x) = 1/(1 - re^{ix})$, $-\pi < x < \pi$. Find the complex Fourier series of f
- 2) Let $0 \leq r < 1$. Use the Fourier series in 1) to derive

$$\frac{1 - r \cos x}{1 + r^2 - 2r \cos x} = 1 + \sum_{n=1}^{\infty} r^n \cos nx,$$

$$\frac{r \sin x}{1 + r^2 - 2r \cos x} = \sum_{n=1}^{\infty} r^n \sin nx.$$

Solution. 1) Expand f as a power series in r . The answer is $f(x) = \sum_{n=0}^{\infty} r^n e^{inx}$. 2) Use Euler's formula and consider the real part and imaginary part.

(1) note that $f(x) = \frac{1}{1 - re^{ix}} = \sum_{k=0}^{\infty} (re^{ix})^k = \sum_{k=0}^{\infty} r^k e^{ikx}$ by geometric series

(Here we can apply geometric series since $|re^{ix}| = r|\cos x + i \sin x| < 1$)

the Fourier series $F(x) = \sum_{n \in \mathbb{Z}} d_n e^{\frac{in\pi x}{\pi}} = \sum_{n \in \mathbb{Z}} d_n e^{inx}$

where $d_n = \frac{1}{2\pi} \int_{-\pi}^{\pi} \sum_{k=0}^{\infty} (re^{ix})^k e^{-inx} dx$

By the uniform convergence of integrand,

$$d_n = \frac{1}{2\pi} \sum_{k=0}^{\infty} \int_{-\pi}^{\pi} r^k e^{i(k-n)x} dx = 2\pi \delta_{kn} r^k \text{ by orthogonality}$$

Since $n \geq 0$, $d_n = \frac{2\pi}{2\pi} r^n = r^n$ for $n \geq 0$ and $d_n = 0$ for $n < 0$

$$\Rightarrow \underline{F(x) = \sum_{n \in \mathbb{Z}} d_n e^{inx} = \sum_{n=0}^{\infty} r^n e^{inx}}$$

(2) Notice that for f in (1):

$$f(x) = \frac{1}{1 - re^{ix}} = \frac{1}{(1 - r \cos x) + i r \sin x} = \frac{1 - r \cos x + i r \sin x}{(1 - r \cos x)^2 + r^2 \sin^2 x}$$

$$= \frac{1 - r \cos x}{1 - 2r \cos x + r^2} + i \frac{r \sin x}{1 - 2r \cos x + r^2}$$

$$\text{Also } f(x) = \sum_{n=0}^{\infty} r^n e^{inx} = \sum_{n=0}^{\infty} r^n \cos nx + i \sum_{n=0}^{\infty} r^n \sin nx$$

$$= \sum_{n=0}^{\infty} r^n \cos nx + i \sum_{n=0}^{\infty} r^n \sin nx$$

$$\Rightarrow \frac{1-r \cos x}{1+r^2-2r \cos x} = \sum_{n=0}^{\infty} r^n \cos nx = 1 + \sum_{n=1}^{\infty} r^n \cos nx$$

$$\frac{r \sin x}{1+r^2-2r \cos x} = \sum_{n=0}^{\infty} r^n \sin nx = \sum_{n=1}^{\infty} r^n \sin nx$$

Question 3. Find the mean square error for the Fourier series of the function $f(x) = 1$ for $0 < x < \pi$, $f(0) = 0$, and $f(x) = -1$ for $-\pi < x < 0$. Then, show that $\sigma_N^2 = O(N^{-1})$ as $N \rightarrow \infty$.

Solution. $\sigma_N^2 = \frac{2}{\pi^2} \sum_{n=N+1}^{\infty} \frac{[(-1)^n - 1]^2}{n^2}$. To show $O(N^{-1})$, define $n = 2m - 1$ and replace the summation with $\sum_{m=(N+2)/2}^{\infty}$ or $\sum_{m=(N+3)/2}^{\infty}$ depending on if N is even or odd. Then use integrals to estimate the sum.

Sol Note that this is an odd function on $(-\pi, \pi)$,

$$\text{so } A_0 = A_n = 0, \forall n$$

$$B_n = \frac{1}{\pi} \left(\int_{-\pi}^0 -\sin nx \, dx + \int_0^{\pi} \sin nx \, dx \right)$$

$$= \frac{1}{\pi} (\cos nx \Big|_{-\pi}^0 - \cos nx \Big|_0^{\pi})$$

$$= \frac{1}{\pi} (1 - (-1)^n - (-1)^n + 1) = \frac{2}{\pi} (1 - (-1)^n)$$

$$\text{So for } N \in \mathbb{N}, \sigma_N^2 = \frac{1}{2} \sum_{n=N+1}^{\infty} \left(\frac{2}{\pi n} (1 - (-1)^n) \right)^2$$

$$= \frac{2}{\pi^2} \sum_{n=N+1}^{\infty} \frac{(1 - (-1)^n)^2}{n^2}$$

$$= \frac{2}{\pi^2} \left(\sum_{m=\frac{N+2}{2}}^{\infty} \frac{(1 - (-1)^{2m-1})^2}{(2m-1)^2} + \sum_{m=\frac{N+3}{2}}^{\infty} \frac{(1 - (-1)^{2m})^2}{(2m)^2} \right)$$

always = $(-1)^{N+1}$ always = $(-1)^{N+2}$

$$= \frac{2}{\pi^2} \sum_{m=\frac{N+2}{2}}^{\infty} \frac{2}{(2m-1)^2} \text{ if } N \text{ is even}$$

$$\text{or } \frac{2}{\pi^2} \sum_{m=\frac{N+3}{2}}^{\infty} \frac{2}{(2m-1)^2} \text{ if } N \text{ is odd}$$

In whatever circumstance, we have

$$\begin{aligned}\sigma_N^2 &\leq \frac{2}{\pi^2} \int_{\frac{N}{2}}^{\infty} \frac{2}{(2m-1)^2} dm = \frac{2}{\pi^2} \int_{N-1}^{\infty} \frac{1}{(2m-1)^2} d(2m-1) \\ &= \frac{2}{\pi^2} \left[-\frac{1}{x} \right]_{N-1}^{\infty} = \frac{2}{\pi^2} \frac{1}{N-1} \in O(N^{-1})\end{aligned}$$

Question 4. Find the mean square error for the Fourier series of $f(x) = x^2$, $-\pi \leq x \leq \pi$. Then, show that $\sigma_N^2 = O(N^{-3})$ as $N \rightarrow \infty$.

Solution. $\sigma_N^2 = 8 \sum_{n=N+1}^{\infty} \frac{1}{n^4}$.

Sol Since f is an even function, $b_n = 0$

$$\begin{aligned}A_n &= \frac{2}{\pi} \int_0^{\pi} x^2 \cos nx \, dx \\ &= \frac{2}{n\pi} \int_0^{\pi} x^2 d(\sin nx) \\ &= \left[\frac{2}{n\pi} x^2 \sin(nx) \right]_0^{\pi} - \frac{4}{n\pi} \int_0^{\pi} \sin(nx) x \, dx \\ &= \frac{4}{n^2\pi} \int_0^{\pi} x d(\cos nx) \\ &= \left[\frac{4}{n^2\pi} x \cos nx \right]_0^{\pi} - \frac{4}{n^2\pi} \int_0^{\pi} \cos(nx) \, dx \\ &= \frac{4}{n^2} (-1)^n\end{aligned}$$

$$\begin{aligned}\text{So } \sigma_N^2 &= \frac{1}{2} \sum_{n=N+1}^{\infty} \frac{4}{n^4} = \frac{1}{8} \sum_{n=N+1}^{\infty} \frac{1}{n^4} \leq \frac{1}{8} \int_N^{\infty} \frac{1}{n^4} \, dn \text{ by integral test} \\ &= \frac{1}{8} \left[-\frac{1}{3} \right]_N^{\infty} \\ &= \frac{1}{24 N^3} \in O(N^{-3})\end{aligned}$$

Question 5. Write out Parseval's theorem for the Fourier series of

1) $f(x) = 1$ for $0 < x < \pi$, $f(0) = 0$, and $f(x) = -1$ for $-\pi < x < 0$,

2) $f(x) = x^2$, $-\pi \leq x \leq \pi$.

Solution. 1) $\pi^2/8 = 1 + \frac{1}{9} + \frac{1}{25} + \dots$, and 2) $\pi^4/90 = 1 + \frac{1}{16} + \frac{1}{81} + \dots$.

Sol. (1) Since f is odd $\Rightarrow A_0, A_n = 0 (\forall n)$

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} f(x)^2 dx = \frac{1}{2} \sum_{n=1}^{\infty} B_n^2$$

$$\Leftrightarrow 1 = \frac{1}{2} \sum_{n=1}^{\infty} \left(\frac{4}{n\pi}\right)^2$$

$$\Leftrightarrow \frac{\pi^2}{8} = \sum_{n=1}^{\infty} \frac{1}{n^2} = 1 + \frac{1}{9} + \frac{1}{25} + \dots$$

(2) f is even $\Rightarrow B_n = 0, \forall n$

$$A_0 = \frac{1}{2\pi} \int_{-\pi}^{\pi} x^2 dx = \frac{1}{2\pi} \cdot \frac{2}{3} \pi^3 = \frac{1}{3} \pi^2$$

$$A_n = \frac{1}{\pi} \int_{-\pi}^{\pi} x^2 \cos n\pi = \frac{4}{n^2} (-1)^n \text{ by the last problem}$$

$$\Rightarrow A_n^2 = \frac{16}{n^4}$$

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} |x^2|^2 dx = A_0^2 + \frac{1}{2} \sum_{n=1}^{\infty} A_n^2$$

$$\Leftrightarrow \frac{1}{2\pi} \frac{2}{5} \pi^5 = \frac{1}{9} \pi^4 + \frac{1}{2} \sum_{n=1}^{\infty} \frac{16}{n^4}$$

$$\Leftrightarrow \left(\frac{1}{5} - \frac{1}{9}\right) \pi^4 = \sum_{n=1}^{\infty} \frac{8}{n^4}$$

$$\Leftrightarrow \frac{4}{45} \pi^4 = \sum_{n=1}^{\infty} \frac{8}{n^4}$$

$$\Leftrightarrow \frac{1}{90} \pi^4 = \sum_{n=1}^{\infty} \frac{1}{n^4} = 1 + \frac{1}{16} + \frac{1}{81} + \dots$$

Question 6. Prove the following Parseval's theorem for complex, cosine and sine Fourier coefficients.

1) $f(x) = \sum_{n=-\infty}^{\infty} \alpha_n e^{in\pi x/L}$ implies that

$$\frac{1}{2L} \int_{-L}^L |f(x)|^2 dx = \sum_{n=-\infty}^{\infty} |\alpha_n|^2.$$

2) $f(x) = A_0 + \sum_{n=1}^{\infty} A_n \cos \frac{n\pi x}{L}$ implies that

$$\frac{1}{L} \int_0^L f(x)^2 dx = A_0^2 + \frac{1}{2} \sum_{n=1}^{\infty} A_n^2.$$

3) $f(x) = \sum_{n=1}^{\infty} B_n \sin \frac{n\pi x}{L}$ implies that

$$\frac{1}{L} \int_0^L f(x)^2 dx = \frac{1}{2} \sum_{n=1}^{\infty} B_n^2.$$

Solution. Try to mimic the proof of Parseval's theorem for Fourier series.

$$\begin{aligned} \text{w) pf } \frac{1}{L} \int_{-L}^L |f(x)|^2 dx &= \frac{1}{L} \int_{-L}^L f(x) \overline{f(x)} dx \\ &= \frac{1}{L} \int_{-L}^L \sum_{n=-\infty}^{\infty} \sum_{m=-\infty}^{\infty} \alpha_n \overline{\alpha_m} e^{\frac{i(n-m)\pi x}{L}} dx \end{aligned}$$

As we have proved: $\int_{-L}^L e^{\frac{i(n-m)\pi x}{L}} dx = 2L \delta_{nm}$, $\neq 0$ iff $n=m$

$$\Rightarrow \frac{1}{L} \int_{-L}^L |f(x)|^2 dx = \sum_{n=-\infty}^{\infty} |\alpha_n|^2$$

□

$$(2) f(x) = A_0 + \sum_{n=1}^{\infty} A_n \cos \frac{n\pi x}{L}$$

$$|f(x)|^2 = A_0^2 + 2A_0 \sum_{n=1}^{\infty} A_n \cos \frac{n\pi x}{L} + \left(\sum_{n=1}^{\infty} A_n \cos \frac{n\pi x}{L} \right)^2$$

$$\begin{aligned} \Rightarrow \frac{1}{L} \int_0^L |f(x)|^2 dx &= \underbrace{\frac{1}{L} \int_0^L A_0^2 dx}_{=A_0^2} + \underbrace{\left(\sum_{n=1}^{\infty} \frac{2}{L} A_0 \int_0^L \cos \frac{n\pi x}{L} dx \right)}_{=0} \\ &\quad + \underbrace{\frac{1}{L} \int_0^L \left(\sum_{n=1}^{\infty} A_n \cos \frac{n\pi x}{L} \right) \left(\sum_{m=1}^{\infty} A_m \cos \frac{m\pi x}{L} \right) dx}_{=\frac{1}{2} \delta_{nm}, \neq 0 \text{ iff } n=m} \end{aligned}$$

$$\Rightarrow \frac{1}{L} \int_0^L |f(x)|^2 dx = A_0^2 + \frac{1}{L} \sum_{n=1}^{\infty} A_n^2 \int_0^L \cos^2 dx = \frac{1}{2L}$$

$$\Rightarrow \frac{1}{L} \int_0^L |f(x)|^2 dx = A_0^2 + \sum_{n=1}^{\infty} A_n^2 \quad \square$$

$$\begin{aligned} (3) \quad \frac{1}{L} \int_0^L |f(x)|^2 dx &= \frac{1}{L} \int_0^L \left(\sum_{n=1}^{\infty} B_n \sin nx \right)^2 dx \\ &= \frac{1}{L} \int_0^L \left(\sum_{n=1}^{\infty} B_n \sin nx \right) \left(\sum_{m=1}^{\infty} B_m \sin mx \right) dx \\ &= \frac{1}{L} \sum_{n=1}^{\infty} B_n^2 \int_0^L \sin nx \sin nx dx = \frac{1}{2L} \\ &= \frac{1}{2} \sum_{n=1}^{\infty} B_n^2 \quad \square \end{aligned}$$

Question 7. Let us solve the heat equation in the slab $0 < z < L$:

$$\begin{cases} u_t = K u_{zz} & 0 < z < L, t > 0, \\ u(0, t) = u(L, t) = 0 & t > 0, \\ u(z, 0) = 1 & 0 < z < L, \end{cases}$$

where $K > 0$ is the thermal conductivity.

- 1) Find the separated solution depending on λ .
- 2) Find the general solution which satisfies the boundary conditions.
- 3) Find the particular solution which satisfies the initial and boundary conditions.

Solution. 1) For $\lambda > 0$, $u = (A \cos \sqrt{\lambda} z + B \sin \sqrt{\lambda} z) e^{-\lambda K t}$, for $\lambda = 0$, $u = (Az + B)$, for $\lambda < 0$, $u = (A e^{\sqrt{-\lambda} z} + B e^{-\sqrt{-\lambda} z}) e^{-\lambda K t}$. 2) $u = \sum_{n=1}^{\infty} A_n \sin(n\pi z/L) e^{-(n\pi/L)^2 K t}$. 3) $u = \frac{2}{\pi} \sum_{n=1}^{\infty} \frac{1 - (-1)^n}{n} \sin \frac{n\pi z}{L} e^{-(n\pi/L)^2 K t}$.

(1) Suppose $u(x, t) = Z(z) T(t)$ for the separated solutions

$$\Rightarrow Z T' = K T Z''$$

$$\frac{T'}{K T} = \frac{Z''}{Z} = -\lambda$$

$$\Rightarrow T' = -K \lambda T, \quad Z'' = -\lambda Z$$

$$\Rightarrow T(t) = C e^{-\lambda K t}$$

$$\Rightarrow Z = \begin{cases} A_1 \cos \sqrt{\lambda} z + A_2 \sin \sqrt{\lambda} z, & \lambda > 0 \\ A_1 z + A_2, & \lambda = 0 \\ A_1 e^{\sqrt{-\lambda} z} + A_2 e^{-\sqrt{-\lambda} z}, & \lambda < 0 \end{cases}$$

Thus the general separated solutions are

$$u(x,t) = \begin{cases} e^{-\lambda^2 t} (A_1 \cos(\sqrt{\lambda} x) + A_2 \sinh(\sqrt{\lambda} x)) & , \lambda > 0 \\ A_1 x + A_2 & , \lambda = 0 \\ e^{-\lambda^2 t} (A_1 e^{\sqrt{\lambda} x} + A_2 e^{-\sqrt{\lambda} x}) & , \lambda < 0 \end{cases}$$

(2) For $\lambda > 0$: $A_1 = 0$, $A_2 \sinh(\sqrt{\lambda} L) e^{-\lambda^2 t} = 0$

$$\Rightarrow \sinh(\sqrt{\lambda} L) = 0$$

$$\Rightarrow \sqrt{\lambda} L = n\pi \Rightarrow \lambda = \left(\frac{n\pi}{L}\right)^2$$

So $u = \sum_{n=1}^{\infty} A_n \sin_n e^{-(\frac{n\pi}{L})^2 kt}$ in this case

for $\lambda = 0 \Rightarrow A_1 = A_2 = 0 \Rightarrow u = 0 \forall x, t$.

for $\lambda < 0$, we have $\forall t$, $A_1 e^{-\lambda^2 t} = 0 \Rightarrow A_1 = 0$

$$\forall t, A_2 \sinh \sqrt{\lambda} L \cdot e^{-\lambda^2 t} = 0 \Rightarrow A_2 = 0$$

$\Rightarrow u = 0 \forall x, t$ in this case by boundary conditions

Therefore overall, $u = \sum_{n=1}^{\infty} A_n \sinh \frac{n\pi x}{L} e^{-\left(\frac{n\pi}{L}\right)^2 kt}$, $0 < x < L$, $t > 0$

is the general solution satisfying the boundary conditions

(3) $u(x,0) = \sum_{n=1}^{\infty} A_n \sinh_n = 1$

Since there is no A_0 , we calculate the Fourier sine coeffs to get the A_n , new s.t. $\sum_{n=1}^{\infty} A_n \sinh_n = 1$

$$\Rightarrow A_n = \frac{2}{L} \int_0^L \sinh \frac{n\pi x}{L} dx$$

$$= \left[\frac{2}{L} \cos \frac{n\pi x}{L} \right]_0^L \cdot \frac{L}{n\pi} = \frac{-2}{n\pi} (1 - (-1)^n) = \frac{2}{n\pi} (1 - (-1)^n)$$

Therefore $u(x,t) = \frac{2}{n\pi} \sum_{n=1}^{\infty} (1 - (-1)^n) \sinh \frac{n\pi x}{L} e^{-\left(\frac{n\pi}{L}\right)^2 kt}$, $0 < x < L$, $t > 0$

is the solution satisfying boundary and initial conditions

