


Finger-tapping home record

EXAMPLE DATA

2026-09-23 to 2026-10-07 · 56 tests · printed 2026-10-06

Hand	Medication	Tests	Taps/s	Size change	Rhythm unevenness*
Right	On	25	3.0	-9%	0.11
Right	On, with troublesome dyskinesia	3	3.2	-15%	0.10
Right	Off	21	2.5	-21%	0.18
Right	Not sure	7	2.7	-16%	0.11

Medians per group. Size change = mean size of the last 3 taps vs the first 3 taps in a 10-second test (one test can be off by about ± 6 points). *Rhythm unevenness = how much the gaps between taps vary (standard deviation $\div$ mean); 0 is perfectly even.



Date · time	Hand	Medication	Min. since dose	Taps/s	Size Δ
2026-10-04 13:06	R	Off · before 1st dose	—	2.3	-19%
2026-10-04 16:06	R	On	86	2.8	-13%

Date · time	Hand	Medication	Min. since dose	Taps/s	Size Δ
2026-10-04 20:06	R	Off	206	2.7	-25%
2026-10-05 00:06	R	On	45	3.0	-7%
2026-10-05 13:06	R	Off · before 1st dose	—	2.1	-21%
2026-10-05 16:06	R	On	94	3.0	-12%
2026-10-05 20:06	R	Not sure	211	2.5	-21%
2026-10-06 00:06	R	On, with troublesome dyskinesia	77	3.3	-8%
2026-10-06 13:06	R	Off · before 1st dose	—	2.9	-19%
2026-10-06 16:06	R	On	105	2.9	0%
2026-10-06 20:06	R	Off	218	2.6	-21%
2026-10-07 00:06	R	On	47	3.0	-20%

Last 12 tests shown.

Self-recorded home measurements from an ordinary webcam. Not a diagnosis and not an MDS-UPDRS score. Do not change medication based on these numbers — discuss them with your neurologist. Recorded with TapTen (open source). Method: MediaPipe hand landmarks, thumb–index 3D distance ÷ palm length.