

Finger-tapping home record

EXAMPLE DATA

2026-09-22 to 2026-10-05 · 56 tests · printed 2026-10-06

Hand	Medication	Tests	Taps/s	Size change	Rhythm unevenness*
Right	On	25	3.0	-9%	0.11
Right	On, with troublesome dyskinesia [†]	3 (few)	3.2	-15%	0.10
Right	Off	21	2.5	-21%	0.18
Right	Not sure	7	2.7	-16%	0.11


Medians per group. Size change = mean size of the last 3 taps vs the first 3 taps in a 10-second test (one test can be off by about ±6 points). *Rhythm unevenness = how much the gaps between taps vary (standard deviation ÷ mean); 0 is perfectly even. " (few)" = fewer than 5 tests: treat the median with caution. [†] Dyskinesia can disturb the measurement. On our synthetic-hand benchmark a single test's size change is typically within ±6 points, and 95% of tests are within ±17 points; accuracy on real hands is not yet known.

● On ◆ On, troublesome dyskinesia ▲ Off ■ Not sure

Taps per second, by date



Taps per second, by minutes since last dose



Date · time	Hand	Medication	Min. since dose	Taps/s	Size Δ
2026-10-04 07:00	R	Off · before 1st dose	—	2.1	-21%
2026-10-04 10:00	R	On	94	3.0	-12%
2026-10-04 14:00	R	Not sure	211	2.5	-21%
2026-10-04 18:00	R	On, with troublesome dyskinesia	77	3.3	-8%
2026-10-05 07:00	R	Off · before 1st dose	—	2.9	-19%
2026-10-05 10:00	R	On	105	2.9	0%
2026-10-05 14:00	R	Off	218	2.6	-21%
2026-10-05 18:00	R	On	47	3.0	-20%

Last 8 tests shown; times are local.

Self-recorded home measurements from an ordinary webcam. Not a diagnosis and not an MDS-UPDRS score. Do not change medication based on these numbers — discuss them with your neurologist. Recorded with TapTen (open source). Method: MediaPipe hand landmarks, thumb-index 3D distance ÷ palm length.