

Exercise sheet 1

Max Wisniewski, Alexander Steen

Problem 1 Computing the minimum

1. Let $X_i \in \{0, 1\}$ be the random variable that indicates whether line (*) is executed in the i -th iteration of the for-loop. Show that $E[X] = \sum_{i=2}^n E[X_i]$.

Since $X = \sum_{i=2}^n X_i$, it holds that

$$E[X] = E\left[\sum_{i=2}^n X_i\right] \stackrel{(*)}{=} \sum_{i=2}^n E[X_i] \quad (1)$$

where (*) holds by linearity of the expected value.

2. Find $E[X_i]$.

Since the numbers are pairwise distinct, the probability that at some fixed position k $A[k]$ is minimal, is given by $Pr[A[k] \text{ minimal}] = \frac{1}{i}$, where i is the number of distinct numbers. It follows that

$$\begin{aligned} E[X_i] &\stackrel{\text{Def.}}{=} \sum_{a \in \{0,1\}} a \cdot Pr[X_i = a] \\ &= Pr[X_i = 1] = Pr[A[i] \text{ minimal in } i \text{ elements}] = \frac{1}{i} \end{aligned} \quad (2)$$

3. Conclude that $E[X] = O(\log n)$.

$$\begin{aligned} E[X] &\stackrel{(1)}{=} \sum_{i=2}^n E[X_i] = \sum_{i=2}^n \frac{1}{i} \\ &\leq \sum_{i=1}^n \frac{1}{i} = H_n = O(\log n) \end{aligned} \quad (3)$$

Problem 2 Induction

Problem 3 O-Notation

1. $\log(n!) = \Theta(n \log(n))$
(1) $\log(n!) = O(n \log(n))$

$$\log(n!) = \sum_{i=1}^n \log i \leq n \cdot \log n \quad (4)$$

- (2) $\log(n!) = \Omega(n \log(n))$

$$TBD \quad (5)$$

2. $\log(mn) = O(\log(n + m))$ holds since

$$\begin{aligned}\log(mn) &= \log(m) + \log(n) \leq \log(m + n) + \log(n + m) \\ &= 2\log(n + m) = O(\log(n + m))\end{aligned}\tag{6}$$

3. Let $f, g \geq 2$ and $f(n) = O(g(n))$

(a) $\sqrt{f(n)} = O(\sqrt{g(n)})$

$$\begin{aligned}\sqrt{f(n)} &= (f(n))^{\frac{1}{2}} \leq (c \cdot g(n))^{\frac{1}{2}} \\ &= \sqrt{c} \cdot \sqrt{g(n)} = O(\sqrt{g(n)})\end{aligned}\tag{7}$$

(b) $2^f(n) = O(2^g(n))$

$$TBD\tag{8}$$