

# Modulation Instability in Semiconductor Quantum Dots

Y-type Excitation Scheme: Interpretation of Plots and Experimental Implications

Shaon Samanta


Department of Physics

April 19, 2025

# Overview

- Motivation: Understanding Modulation Instability (MI) in SQDs (Y-type excitation).
- Tools: Density matrix formalism, NLSE.
- Focus: Interpretation of numerical plots & physical insights.


# Light Matter Interaction




- Quantum dots confine charge carriers in all three dimensions.
- Enhanced interaction with electromagnetic fields due to confinement.
- Leads to strong nonlinear effects: Kerr nonlinearity, EIT, and MI.

# What are Semiconductor Quantum Dots

- Tiny semiconductor nanocrystals (2–10 nm in size).
- Exhibit size-dependent fluorescence, unique electronic properties, enhanced optical properties.
- Intermediate properties between bulk semiconductors and discrete molecules.




# Y-type Excitation Scheme in SQDs




- Probe field couples levels  $|1\rangle \rightarrow |2\rangle$ .
- Control fields modulate transitions  $|2\rangle \rightarrow |3\rangle$ ,  $|3\rangle \rightarrow |4\rangle$ .
- The interacting electric field inside QD is:  $\vec{E} = \sum_{j=p,c,b} \hat{e}_j E_j e^{i(k_j z - \omega_j t)}$

# Absorption & Dispersion Spectra




(a)



(b)


- $\Omega_d$  modifies absorption: transparency windows form.
- Side peaks grow  $\Rightarrow$  indicative of coherent interference.
- $\text{Re } \chi^{(1)}$  slope changes  $\Rightarrow$  group velocity control.

## Kerr Nonlinearity: $\text{Re } \chi^{(3)}$



- $\Omega_c, \Omega_d$  enhance Kerr nonlinearity.
- Enables efficient four-wave mixing and phase modulation.
- $\chi^{(3)}$  control  $\Rightarrow$  quantum gates, optical switches.


# Modulation Instability Gain



- MI gain shows symmetric sidebands  $\Rightarrow$  phase matching.
- Gain increases with laser power.
- Peaks shift outward with higher nonlinear refractive index.



# MI Gain



- Threshold at  $P_0 \approx 5$  mW.
- Fourth-order dispersion ( $\beta_4$ ) broadens bandwidth.
- Application: broadband supercontinuum generation.

# Conclusion & Applications

- SQDs exhibit tunable nonlinearity through coherent control.
- Applications:
  - Quantum memory and communication
  - Slow light and optical buffers
  - Frequency combs and all-optical logic