

Exercício 1

Regression

Descriptive Statistics

	Mean	Std. Deviation	N
I (intensidade de corrente)	1,5460	,72807	5
V (dif. Potencial)	1,3600	,61074	5

Correlations

		I (intensidade de corrente)	V (dif. Potencial)
Pearson Correlation	I (intensidade de corrente)	1,000	,994
	V (dif. Potencial)	,994	1,000
Sig. (1-tailed)	I (intensidade de corrente)	.	,000
	V (dif. Potencial)	,000	.
N	I (intensidade de corrente)	5	5
	V (dif. Potencial)	5	5

Variables Entered/Removed^b

Model	Variables Entered	Variables Removed	Method
1	V (dif. Potencial) ^a	.	Enter

a. All requested variables entered.

b. Dependent Variable: I (intensidade de corrente)

Model Summary^b

Model	R	R Square	Adjusted R Square	Std. Error of the Estimate
1	,994 ^a	,987	,983	,09515

a. Predictors: (Constant), V (dif. Potencial)

b. Dependent Variable: I (intensidade de corrente)

ANOVA^b

Model		Sum of Squares	df	Mean Square	F	Sig.
1	Regression	2,093	1	2,093	231,210	,001 ^a
	Residual	,027	3	,009		
	Total	2,120	4			

a. Predictors: (Constant), V (dif. Potencial)

b. Dependent Variable: I (intensidade de corrente)

Coefficients^a

Model		Unstandardized Coefficients		Standardized Coefficients		
		B	Std. Error	Beta	t	Sig.
1	(Constant)	-,065	,114		-,568	,610
	V (dif. Potencial)	1,184	,078	,994	15,206	,001


a. Dependent Variable: I (intensidade de corrente)

Coefficients^a


Model		95,0% Confidence Interval for B	
		Lower Bound	Upper Bound
1	(Constant)	-,428	,298
	V (dif. Potencial)	,937	1,432

a. Dependent Variable: I (intensidade de corrente)


Graph



Graph



Graph



Explore

Case Processing Summary						
	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
Studentized Residual	5	100,0%	0	,0%	5	100,0%

Descriptives

		Statistic	Std. Error
Studentized Residual	Mean	,0353615	,48184425
	95% Confidence Interval for Mean		
	Lower Bound	-1,3024526	
	Upper Bound	1,3731756	
	5% Trimmed Mean	,0224441	
	Median	-,1404695	
	Variance	1,161	
	Std. Deviation	1,07743649	
	Minimum	-1,08794	
	Maximum	1,39117	
	Range	2,47911	
	Interquartile Range	2,10873	
	Skewness	,313	,913
	Kurtosis	-2,322	2,000

NPar Tests

One-Sample Kolmogorov-Smirnov Test

		Studentized Residual
	N	5
Normal Parameters ^{a,b}	Mean	,0353615
	Std. Deviation	1,07743649
Most Extreme Differences	Absolute	,198
	Positive	,198
	Negative	-,182
	Kolmogorov-Smirnov Z	,442
	Asymp. Sig. (2-tailed)	,990

a. Test distribution is Normal.

b. Calculated from data.

Exercício 2

Variables Entered/Removed

Model	Variables Entered	Variables Removed	Method
1	X2, X1 ^a	.	Enter

a. All requested variables entered.

Model Summary^b

Model	R	R Square	Adjusted R Square	Std. Error of the Estimate
1	,990 ^a	,979	,973	2,433

a. Predictors: (Constant), X2, X1

b. Dependent Variable: Y

ANOVA^b

Model		Sum of Squares	df	Mean Square	F	Sig.
1	Regression	1953,076	2	976,538	165,021	,000 ^a
	Residual	41,424	7	5,918		
	Total	1994,500	9			

a. Predictors: (Constant), X2, X1

b. Dependent Variable: Y

Coefficients^a

Model		Unstandardized Coefficients		Standardized Coefficients		
		B	Std. Error	Beta	t	Sig.
1	(Constant)	51,720	2,624		19,707	,000
	X1	1,515	,130	,639	11,651	,000
	X2	,669	,053	,688	12,553	,000

a. Dependent Variable: Y

Coefficients^a

Model	95,0% Confidence Interval for B	
	Lower Bound	Upper Bound
1 (Constant)	45,514	57,926
X1	1,207	1,822
X2	,543	,794

a. Dependent Variable: Y

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
Studentized Residual	10	100,0%	0	,0%	10	100,0%

Descriptives

		Statistic	Std. Error
Studentized Residual	Mean	,0116925	,31869746
	95% Confidence Interval for Mean		
	Lower Bound	-,7092513	
	Upper Bound	,7326362	
	5% Trimmed Mean	,0149189	
	Median	-,3835987	
	Variance	1,016	
	Std. Deviation	1,00780986	
	Minimum	-1,77009	
	Maximum	1,73540	
	Range	3,50550	
	Interquartile Range	1,26311	
	Skewness	,078	,687
	Kurtosis	,013	1,334


Tests of Normality



	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
	Statistic	df	Sig.	Statistic	df	Sig.
Studentized Residual	,237	10	,117	,928	10	,430

a. Lilliefors Significance Correction

Studentized Residual

Normal Q-Q Plot of Studentized Residual





Exercício 3

Warnings

The Cubic model could not be fitted due to near-collinearity among model terms.

Model Description

	Model Name	MOD_3
Dependent Variable	1	Percentagem
Equation	1	Linear
	2	Quadratic
	3	Cubic
	Independent Variable	Temperatura
	Constant	Included
	Variable Whose Values	Unspecified
	Label Observations in Plots	
	Tolerance for Entering	,0001
	Terms in Equations	

Case Processing Summary

	N
Total Cases	10
Excluded Cases ^a	0
Forecasted Cases	0
Newly Created Cases	0

a. Cases with a missing value in any variable are excluded from the analysis.

Variable Processing Summary

		Variables	
		Dependent	Independent
		Percentagem	Temperatura
	Number of Positive Values	10	10
	Number of Zeros	0	0
	Number of Negative Values	0	0
Number of Missing Values	User-Missing	0	0
	System-Missing	0	0

Model Summary and Parameter Estimates

Dependent Variable:Percentagem

Equation	Model Summary				
	R Square	F	df1	df2	Sig.
Linear	,149	1,401	1	8	,271
Quadratic	,673	7,189	2	7	,020
Cubic	,742	10,065	2	7	,009


The independent variable is Temperatura.

Model Summary and Parameter Estimates

Dependent Variable:Percentagem

Equation	Parameter Estimates			
	Constant	b1	b2	b3
Linear	6,104	-,018		
Quadratic	-22,360	,410	-,002	
Cubic	-4,599	,000	,002	-7,619E-6

The independent variable is Temperatura.



Exercício 4

Variables Entered/Removed^b

Model	Variables Entered	Variables Removed	Method
1	log10d ^a	.	Enter

a. All requested variables entered.

b. Dependent Variable: log10h

Model Summary^b

Model	R	R Square	Adjusted R Square	Std. Error of the Estimate
1	,994 ^a	,988	,985	,01544

a. Predictors: (Constant), log10d

b. Dependent Variable: log10h

ANOVA^b

Model		Sum of Squares	df	Mean Square	F	Sig.
1	Regression	,079	1	,079	333,302	,000 ^a
	Residual	,001	4	,000		
	Total	,080	5			

a. Predictors: (Constant), log10d

b. Dependent Variable: log10h

Coefficients^a

Model		Unstandardized Coefficients		Standardized Coefficients		
		B	Std. Error	Beta	t	Sig.
1	(Constant)	4,968	,141		35,305	,000
	log10d	-1,865	,102	-,994	-18,257	,000

a. Dependent Variable: log10h

Coefficients^a

Model	95,0% Confidence Interval for B	
	Lower Bound	Upper Bound
1 (Constant)	4,577	5,358
log10d	-2,149	-1,582

a. Dependent Variable: log10h

Explore**Case Processing Summary**

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
Studentized Residual	6	100,0%	0	,0%	6	100,0%

Descriptives

		Statistic	Std. Error
Studentized Residual	Mean	-,0567132	,45801918
	95% Confidence Interval for Mean		
	Lower Bound	-1,2340890	
	Upper Bound	1,1206626	
	5% Trimmed Mean	-,0242651	
	Median	-,1038435	
	Variance	1,259	
	Std. Deviation	1,12191328	
	Minimum	-1,91378	
	Maximum	1,21629	
	Range	3,13007	
	Interquartile Range	1,79655	
	Skewness	-,710	,845
	Kurtosis	,642	1,741

Tests of Normality


	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
	Statistic	df	Sig.	Statistic	df	Sig.
Studentized Residual	,215	6	,200 [*]	,935	6	,623



a. Lilliefors Significance Correction

*. This is a lower bound of the true significance.

Studentized Residual

Normal Q-Q Plot of Studentized Residual





Exercício 5

Variables Entered/Removed^b

Model	Variables Entered	Variables Removed	Method
1	temperatura ^a	.	Enter

a. All requested variables entered.

b. Dependent Variable: oxigénio

Model Summary^b

Model	R	R Square	Adjusted R Square	Std. Error of the Estimate
1	,990 ^a	,981	,978	,16825

a. Predictors: (Constant), temperatura

b. Dependent Variable: oxigénio

ANOVA^b

Model		Sum of Squares	df	Mean Square	F	Sig.
1	Regression	8,745	1	8,745	308,933	,000 ^a
	Residual	,170	6	,028		
	Total	8,915	7			

a. Predictors: (Constant), temperatura

b. Dependent Variable: oxigénio

Coefficients^a

Model		Unstandardized Coefficients		Standardized Coefficients		
		B	Std. Error	Beta	t	Sig.
1	(Constant)	3,471	,060		57,738	,000
	temperatura	-,088	,005	-,990	-17,576	,000

a. Dependent Variable: oxigénio

Coefficients^a

Model		95,0% Confidence Interval for B	
		Lower Bound	Upper Bound
1	(Constant)	3,324	3,619
	temperatura	-,100	-,076

a. Dependent Variable: oxigénio

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
Studentized Residual	8	100,0%	0	,0%	8	100,0%

Descriptives

		Statistic	Std. Error
Studentized Residual	Mean	,0223358	,36441898
	95% Confidence Interval for Mean		
	Lower Bound	-,8393782	
	Upper Bound	,8840497	
	5% Trimmed Mean	,0735599	
	Median	,2022084	
	Variance	1,062	
	Std. Deviation	1,03073253	
	Minimum	-1,98166	
	Maximum	1,10430	
	Range	3,08595	
	Interquartile Range	1,50204	
	Skewness	-1,011	,752
	Kurtosis	,836	1,481

Tests of Normality



	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
	Statistic	df	Sig.	Statistic	df	Sig.
Studentized Residual	,157	8	,200*	,916	8	,396


a. Lilliefors Significance Correction

*. This is a lower bound of the true significance.

Studentized Residual

Normal Q-Q Plot of Studentized Residual





Exercício 6

Variables Entered/Removed^b

Model	Variables Entered	Variables Removed	Method
1	% S2 ^a	.	Enter

a. All requested variables entered.

b. Dependent Variable: Decrescimento

Model Summary^b

Model	R	R Square	Adjusted R Square	Std. Error of the Estimate
1	,925 ^a	,855	,819	66,4186

a. Predictors: (Constant), % S2

b. Dependent Variable: Decrescimento

ANOVA^b

Model		Sum of Squares	df	Mean Square	F	Sig.
1	Regression	104404,290	1	104404,290	23,667	,008 ^a
	Residual	17645,710	4	4411,428		
	Total	122050,000	5			

a. Predictors: (Constant), % S2

b. Dependent Variable: Decrescimento

Coefficients^a

Model		Unstandardized Coefficients		Standardized Coefficients		
		B	Std. Error	Beta	t	Sig.
1	(Constant)	374,534	45,784		8,181	,001
	% S2	474,988	97,637	,925	4,865	,008

a. Dependent Variable: Decrescimento

Coefficients^a

Model		95,0% Confidence Interval for B	
		Lower Bound	Upper Bound
1	(Constant)	247,418	501,649
	% S2	203,905	746,071

a. Dependent Variable: Decrescimento

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
Studentized Residual	6	100,0%	0	,0%	6	100,0%

Descriptives

		Statistic	Std. Error
Studentized Residual	Mean	-,0949053	,45070076
	95% Confidence Interval for Mean		
	Lower Bound	-1,2534685	
	Upper Bound	1,0636579	
	5% Trimmed Mean	-,0814108	
	Median	-,0811354	
	Variance	1,219	
	Std. Deviation	1,10398690	
	Minimum	-1,63763	
	Maximum	1,20492	
	Range	2,84255	
	Interquartile Range	2,17538	
	Skewness	-,190	,845
	Kurtosis	-1,314	1,741

Tests of Normality



	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
	Statistic	df	Sig.	Statistic	df	Sig.
Studentized Residual	,170	6	,200*	,956	6	,791


a. Lilliefors Significance Correction

*. This is a lower bound of the true significance.

Studentized Residual

Normal Q-Q Plot of Studentized Residual





Variables Entered/Removed^b

Model	Variables Entered	Variables Removed	Method
1	lnx ^a	.	Enter

a. All requested variables entered.

b. Dependent Variable: Decrescimento

Model Summary^b

Model	R	R Square	Adjusted R Square	Std. Error of the Estimate
1	,996 ^a	,992	,990	15,6077

a. Predictors: (Constant), lnx

b. Dependent Variable: Decrescimento

ANOVA^b

Model		Sum of Squares	df	Mean Square	F	Sig.
1	Regression	121075,594	1	121075,594	497,023	,000 ^a
	Residual	974,406	4	243,602		
	Total	122050,000	5			

a. Predictors: (Constant), lnx

b. Dependent Variable: Decrescimento

Coefficients^a

Model	Unstandardized Coefficients		Standardized Coefficients		
	B	Std. Error	Beta	t	Sig.
1 (Constant)	735,928	10,353		71,081	,000
lnx	127,589	5,723	,996	22,294	,000

a. Dependent Variable: Decrecimiento

Coefficients^a

Model	95,0% Confidence Interval for B	
	Lower Bound	Upper Bound
1 (Constant)	707,182	764,673
lnx	111,699	143,479

a. Dependent Variable: Decrecimiento

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
Studentized Residual	6	100,0%	0	,0%	6	100,0%

Descriptives

		Statistic	Std. Error
Studentized Residual	Mean	,0248247	,43991670
	95% Confidence Interval for Mean		
	Lower Bound	-1,1060172	
	Upper Bound	1,1556666	
	5% Trimmed Mean	,0618698	
	Median	,3885030	
	Variance	1,161	
	Std. Deviation	1,07757144	
	Minimum	-1,85487	
	Maximum	1,23770	

	Range	3,09257	
	Interquartile Range	1,54894	
	Skewness	-1,153	,845
	Kurtosis	1,449	1,741


Tests of Normality



	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
	Statistic	df	Sig.	Statistic	df	Sig.
Studentized Residual	,296	6	,110	,896	6	,349

a. Lilliefors Significance Correction

Studentized Residual

Normal Q-Q Plot of Studentized Residual





Exercício 7

Variables Entered/Removed

Model	Variables Entered	Variables Removed	Method
1	Horas, QI ^a	.	Enter

a. All requested variables entered.

Model Summary^b

Model	R	R Square	Adjusted R Square	Std. Error of the Estimate
1	,884 ^a	,782	,719	10,096

a. Predictors: (Constant), Horas, QI

b. Dependent Variable: Resultado

ANOVA^b

Model		Sum of Squares	df	Mean Square	F	Sig.
1	Regression	2554,156	2	1277,078	12,530	,005 ^a
	Residual	713,444	7	101,921		
	Total	3267,600	9			

a. Predictors: (Constant), Horas, QI

b. Dependent Variable: Resultado

Coefficients^a

Model		Unstandardized Coefficients		Standardized Coefficients		
		B	Std. Error	Beta	t	Sig.
1	(Constant)	-124,568	47,874		-2,602	,035
	QI	1,659	,446	,733	3,719	,007
	Horas	1,439	1,067	,266	1,349	,219

a. Dependent Variable: Resultado

Coefficients^a

Model		95,0% Confidence Interval for B	
		Lower Bound	Upper Bound
1	(Constant)	-237,773	-11,364
	QI	,604	2,714
	Horas	-1,083	3,962

a. Dependent Variable: Resultado

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
Studentized Residual	10	100,0%	0	,0%	10	100,0%

Descriptives

		Statistic	Std. Error
Studentized Residual	Mean	,0095740	,31978363
95% Confidence Interval for Mean	Lower Bound	-,7138268	
	Upper Bound	,7329748	
	5% Trimmed Mean	,0325599	
	Median	,4270071	
	Variance	1,023	
	Std. Deviation	1,01124464	
	Minimum	-1,63807	
	Maximum	1,24347	
	Range	2,88154	
	Interquartile Range	1,70487	
	Skewness	-,405	,687
	Kurtosis	-1,443	1,334



Tests of Normality


	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
	Statistic	df	Sig.	Statistic	df	Sig.
Studentized Residual	,260	10	,054	,893	10	,184

a. Lilliefors Significance Correction

Studentized Residual

Normal Q-Q Plot of Studentized Residual





Exercício 8

NPar Tests

One-Sample Kolmogorov-Smirnov Test

		motor	intelectual
Normal Parameters ^{a, b}	N	9	9
	Mean	87,0000	97,0000
	Std. Deviation	10,18577	12,39960
Most Extreme Differences	Absolute	,128	,151
	Positive	,128	,146
	Negative	-,121	-,151
	Kolmogorov-Smirnov Z	,383	,453
	Asymp. Sig. (2-tailed)	,999	,986

a. Test distribution is Normal.

b. Calculated from data.

Correlations

Correlations

		motor	intelectual
motor	Pearson Correlation	1	,743*
	Sig. (2-tailed)		,022
	N	9	9
intelectual	Pearson Correlation	,743*	1
	Sig. (2-tailed)	,022	
	N	9	9

*. Correlation is significant at the 0.05 level (2-tailed).

Exercício 9

Explore

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
QI	30	100,0%	0	,0%	30	100,0%
CM	30	100,0%	0	,0%	30	100,0%

Descriptives

		Statistic	Std. Error
QI	Mean	97,2667	2,42446
	95% Confidence Interval for Mean		
	Lower Bound	92,3081	
	Upper Bound	102,2252	
	5% Trimmed Mean	96,6111	
	Median	95,0000	
	Variance	176,340	
	Std. Deviation	13,27932	
	Minimum	79,00	
	Maximum	131,00	
	Range	52,00	
	Interquartile Range	23,25	
CM	Mean	2,5663	,15423
	95% Confidence Interval for Mean		
	Lower Bound	2,2509	
	Upper Bound	2,8818	
	5% Trimmed Mean	2,5859	
	Median	2,5850	
	Variance	,714	
	Std. Deviation	,84476	
	Minimum	,67	
	Maximum	4,00	
	Range	3,33	
	Interquartile Range	1,00	


Tests of Normality

	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
	Statistic	df	Sig.	Statistic	df	Sig.
QI	,134	30	,175	,942	30	,101
CM	,114	30	,200*	,967	30	,470


a. Lilliefors Significance Correction



*. This is a lower bound of the true significance.

QI



CM





Correlations

Correlations

		QI	CM
QI	Pearson Correlation	1	,708**
	Sig. (2-tailed)		,000
	N	30	30
CM	Pearson Correlation	,708**	1
	Sig. (2-tailed)	,000	
	N	30	30

**. Correlation is significant at the 0.01 level (2-tailed).

Exercício 10

Correlations

		X	Y
X	Pearson Correlation	1	-,334
	Sig. (2-tailed)		,071
	N	30	30
Y	Pearson Correlation	-,334	1
	Sig. (2-tailed)	,071	
	N	30	30

NPar Tests

One-Sample Kolmogorov-Smirnov Test

		X	Y
Normal Parameters ^{a,b}	N	30	30
	Mean	3,8977	8,6683
	Std. Deviation	2,83786	6,99143
Most Extreme Differences	Absolute	,140	,140
	Positive	,140	,140
	Negative	-,114	-,108
	Kolmogorov-Smirnov Z	,769	,769
	Asymp. Sig. (2-tailed)	,595	,595

a. Test distribution is Normal.

b. Calculated from data.

Exercício 11

Correlations		cálcio	colesterol
cálcio	Pearson Correlation	1	,857**
	Sig. (2-tailed)		,000
	N	12	12
colesterol	Pearson Correlation	,857**	1
	Sig. (2-tailed)	,000	
	N	12	12

** . Correlation is significant at the 0.01 level (2-tailed).

Correlations			cálcio	colesterol
Spearman's rho	cálcio	Correlation Coefficient	1,000	,689*
		Sig. (2-tailed)		,013
		N	12	12
	colesterol	Correlation Coefficient	,689*	1,000
		Sig. (2-tailed)	,013	
		N	12	12

*. Correlation is significant at the 0.05 level (2-tailed).

Exercício 12

Correlations			ele	ela
Spearman's rho	ele	Correlation Coefficient	1,000	-,613
		Sig. (2-tailed)		,060
		N	10	10
	ela	Correlation Coefficient	-,613	1,000
		Sig. (2-tailed)	,060	
		N	10	10

Exercício 13

Nonparametric Correlations

Correlations			tempo	velocidade
Spearman's rho	tempo	Correlation Coefficient	1,000	1,000**
		Sig. (1-tailed)	.	.
		N	5	5
	velocidade	Correlation Coefficient	1,000**	1,000
		Sig. (1-tailed)	.	.
		N	5	5

** . Correlation is significant at the 0.01 level (1-tailed).

NPar Tests

One-Sample Kolmogorov-Smirnov Test				tempo	velocidade
		N		5	5
Normal Parameters ^{a, b}	Mean			,4000	3,8540
	Std. Deviation			,31623	3,16245
Most Extreme Differences	Absolute			,136	,152
	Positive			,136	,142
	Negative			-,136	-,152
	Kolmogorov-Smirnov Z			,305	,341
	Asymp. Sig. (2-tailed)			1,000	1,000

a. Test distribution is Normal.

b. Calculated from data.

Correlations				tempo	velocidade
tempo	Pearson Correlation			1	,998**
	Sig. (1-tailed)				,000
	N			5	5
velocidade	Pearson Correlation			,998**	1
	Sig. (1-tailed)			,000	
	N			5	5

** . Correlation is significant at the 0.01 level (1-tailed).

Exercício 14

Correlations			DocenteA	DocenteB
Spearman's rho	DocenteA	Correlation Coefficient	1,000	,460
		Sig. (2-tailed)		,212
		N	9	9
	DocenteB	Correlation Coefficient	,460	1,000
		Sig. (2-tailed)	,212	
		N	9	9

Exercício 15

Correlations			idade	lugar
Spearman's rho	idade	Correlation Coefficient	1,000	,680*
		Sig. (1-tailed)		,046
		N	7	7
	lugar	Correlation Coefficient	,680*	1,000
		Sig. (1-tailed)	,046	
		N	7	7

*. Correlation is significant at the 0.05 level (1-tailed).