

Geant4: *L-qBar QUARTIC*


Bartłomiej Rzeszotarski

AGH - Wydział Fizyki i Informatyki Stosowanej


31 marca 2014

ATLAS Forward Protons

- QUARTIC - *QUARz Timing Cherenkov*




location: symmetrically, $\approx 214\text{m}$ from collision point



regular quartz L-qbar with aluminized elbow (90% reflectivity):

- a. vertical arm: length 80 mm, cross profile 3x6 mm,
- b. horizontal arm: length 80 mm, cross profile 3x6 mm,



regular quartz L-qbar with aluminized elbow (90% reflectivity):

- a. vertical arm: length 80 mm, cross profile 3x6 mm,
- b. horizontal arm: length 80 mm, cross profile 3x6 mm,

Configurations:


- a) four bars in one line without any isolation (acts like one bar)
- b) four bars in one line, separated using air isolation (acts like four independent bars)


Configurations:

- a) four bars in one line without any isolation (acts like one bar)
- b) four bars in one line, separated using air isolation (acts like four independent bars)

Configurations:

- a) four bars in one line without any isolation (acts like one bar)
- b) four bars in one line, separated using air isolation (acts like four independent bars)





Fused silica:

- Refractive index 1.45 - 1.51 gives Cherenkov radiation angle $\approx 48^\circ$ - total internal reflection
- Absorption length ($\approx 1m$) allows to reach the photodetector by almost all photons

Fused silica:

- Refractive index 1.45 - 1.51 gives Cherenkov radiation angle $\approx 48^\circ$ - total internal reflection
- Absorption length ($\approx 1m$) allows to reach the photodetector by almost all photons


Fused silica:

- Refractive index 1.45 - 1.51 gives Cherenkov radiation angle $\approx 48^\circ$ - total internal reflection
- Absorption length ($\approx 1m$) allows to reach the photodetector by almost all photons

L-qBar is designed to detect the protons scattered at small angles:

- high time resolution ($10 - 15ps$) allows us to specify the vertex point, and;
- to identify exactly which protons were involved in the interaction


With this information, we can more thoroughly study the *exclusive reactions*, in which is produced the Higgs boson or W^+W^- bosons



L-qBar is designed to detect the protons scattered at small angles:

- high time resolution ($10 - 15ps$) allows us to specify the vertex point, and;
- to identify exactly which protons were involved in the interaction


With this information, we can more thoroughly study the *exclusive reactions*, in which is produced the Higgs boson or W^+W^- bosons



L-qBar is designed to detect the protons scattered at small angles:

- high time resolution ($10 - 15\text{ps}$) allows us to specify the vertex point, and;
- to identify exactly which protons were involved in the interaction


With this information, we can more thoroughly study the *exclusive reactions*, in which is produced the Higgs boson or W^+W^- bosons



L-qBar is designed to detect the protons scattered at small angles:

- high time resolution ($10 - 15ps$) allows us to specify the vertex point, and;
- to identify exactly which protons were involved in the interaction

With this information, we can more thoroughly study the *exclusive reactions*, in which is produced the Higgs boson or W^+W^- bosons



L-qBar is designed to detect the protons scattered at small angles:

- high time resolution ($10 - 15ps$) allows us to specify the vertex point, and;
- to identify exactly which protons were involved in the interaction

With this information, we can more thoroughly study the *exclusive reactions*, in which is produced the Higgs boson or W^+W^- bosons

$$p + p \rightarrow p + H + p$$

L-qBar is designed to detect the protons scattered at small angles:

- high time resolution ($10 - 15ps$) allows us to specify the vertex point, and;
- to identify exactly which protons were involved in the interaction

With this information, we can more thoroughly study the *exclusive reactions*, in which is produced the Higgs boson or W^+W^- bosons

$$p + p \rightarrow p + H + p$$

$$p + p \rightarrow p + W^+W^- + p$$

