


EXPERIMENT- 10

1.

Dataset

```
survey <- data.frame(  
  SurveyID = c(1,2,3),  
  Question1 = c("A","B","C"),  
  Question2 = c("B","A","A"),  
  Question3 = c("C","D","B")  
)  
  
# Frequency of Question 1 responses  
q1 <- table(survey$Question1)  
  
# Grouped Bar Chart  
barplot(q1,  
  beside = TRUE,  
  col = c("red","green","blue"),  
  main = "Distribution of Answers for Question 1",  
  xlab = "Responses",  
  ylab = "Frequency")
```



2.

Dataset

```
survey <- data.frame(  
  SurveyID = c(1,2,3),  
  Question1 = c("A","B","C"),  
  Question2 = c("B","A","A"),  
  Question3 = c("C","D","B")  
)
```


Create response table

```
response_table <- rbind(  
  table(survey$Question1),  
  table(survey$Question2),  
  table(survey$Question3)  
)
```

```
rownames(response_table) <- c("Question1","Question2","Question3")
```

Stacked Bar Chart

```
barplot(response_table,  
  col = rainbow(ncol(response_table)),  
  main = "Responses for All Questions",  
  xlab = "Questions",  
  ylab = "Frequency",  
  legend = colnames(response_table))
```




3.

Dataset

```
survey <- data.frame(  
  SurveyID = c(1,2,3),  
  Question1 = c("A","B","C"),  
  Question2 = c("B","A","A"),  
  Question3 = c("C","D","B")  
)
```

Display Table

```
print(survey)
```



4.

