The Greatest Common Divisor and The Least Common Multiple

By Mathew Mouchamel

The Greatest Common Divisor

What is a divisor?

A divisor is a number that divides another number

However, in the context of integers, it is a number that divides another number completely. Thus, if a number A is the divisor of another number B, it means that when B is divided by A, there is no remainder.

What is the greatest common divisor?

The Greatest Common Divisor (GCD) of two numbers is the largest number that divides both of them without leaving a remainder. Thus, it is the largest divisor that both numbers have in common.

The three common methods of finding the GCD include:

- 1. The Brute Force Method or Listing Method
- 2. The Prime Factorization Method
- 3. The Euclidean Algorithm

Method #1

1. The Brute Force Method

This method involves listing out all the divisors of each number and then finding the largest number that appears on both lists. It's a straightforward approach but can be inefficient for large numbers.

12 :	1, 2,	3, 4,	6,	12
------	-------	-------	----	----

$$12 / 3 = 4$$

Example 1: Greatest Common Divisor of 12 and 18.

12: 1, 2, 3, 4, 6, 12

18: 1, 2, 3, 6, 9, 18

$$12 / 4 = 3$$

Example 1: Greatest Common Divisor of 12 and 18.

12: 1, 2, 3, 4, 6, 12

18: 1, 2, 3, 6, 9, 18

12 / 2 = 6

18 / 3 = 6

GCD = 6

Example 2: The GCD of 720 and 840

720: 1, 2, 3, 4, 5, 6, 8, 9, 10, 12, 15, 16, 18, 20, 24, 30, 36, 40, 45, 48, 60, 72, 80, 90, 120, 144, 180, 240, 360, 720

840 : 1, 2, 3, 4, 5, 6, 7, 8, 10, 12, 14, 15, 20, 21, 24, 28, 30, 35, 40, 42, 56, 60, 70, 84, 105, 120, 140, 168, 210, 280, 420, 840

Example 2: The GCD of 720 and 840

720: 1, 2, 3, 4, 5, 6, 8, 10, 12, 15, 20, 24, 30, 40, 60, 120

840: 1, 2, 3, 4, 5, 6, 8, 10, 12, 15, 20, 24, 30, 40, 60, 120

Example 2: The GCD of 720 and 840

720: 1, 2, 3, 4, 5, 6, 8, 10, 12, 15, 20, 24, 30, 40, 60, 120

120

840: 1, 2, 3, 4, 5, 6, 8, 10, 12, 15, 20, 24, 30, 40, 60,

Example 2: The GCD of 720 and 840

720: 1, 2, 3, 4, 5, 6, 8, 10, 12, 15, 20, 24, 30, 40, 60, 120

840: 1, 2, 3, 4, 5, 6, 8, 10, 12, 15, 20, 24, 30, 40, 60,

GCD = 120

Method #2

2. The Prime Factorization Method

Here, each number is broken down into its prime factors. The GCD is then found by multiplying the smallest power of all common prime factors present in both numbers. This method provides a clear illustration of the common factors but can be time-consuming if the prime factorization is complicated.

- $12: (2 \wedge 2) * (3 \wedge 1)$
- 18: (2 ^ 1) * (3 ^ 2)

Example 1: Greatest Common Divisor of 12 and 18.

12: (2 ^ 2) * (3 ^ 1)

18: (2 ^ 1) * (3 ^ 2)

Prime Factors: 2, 3

$$GCD = (2 \land min(2, 1)) * (3 \land min(1, 2))$$

$$GCD = (2 \land min(2, 1)) * (3 \land min(1, 2))$$

$$GCD = (2 ^1) * (3 ^1)$$

$$GCD = (2 \land min(2, 1)) * (3 \land min(1, 2))$$

$$GCD = (2 \land 1) * (3 \land 1) = 2 * 3$$

ı							
	Example 1:	Crostost	Common	Dirrigan	of 12	and	10
	Example 1:	Greatest	Common	DIVISOI	OI 1Z	allu	10.
	1						

$$GCD = (2 \land min(2, 1)) * (3 \land min(1, 2))$$

GCD =
$$(2 \land 1) * (3 \land 1) = 2 * 3$$

$$GCD = 6$$

```
720: (2 ^ 4) * (3 ^ 2) * (5 ^ 1)
```

```
840: (2 ^ 3) * (3 ^ 1) * (5 ^ 1) * (7 ^ 1)
```

Example 2: Greatest Common Divisor of 720 and 840

```
720: (2 ^ 4) * (3 ^ 2) * (5 ^ 1)
```

Prime Factors: 2, 3, 5, 7

Example 2: Greatest Common Divisor of 720 and 840

```
720: (2 ^ 4) * (3 ^ 2) * (5 ^ 1)
840: (2 ^ 3) * (3 ^ 1) * (5 ^ 1) * (7 ^ 1)
```

Prime Factors: 2, 3, 5, 7

GCD =
$$(2 \land \min(4, 3)) * (3 \land \min(2, 1)) * (5 \land \min(1, 1)) * (7 \land \min(0, 1))$$

720:
$$(2 ^4) * (3 ^2) * (5 ^1)$$

840: $(2 ^3) * (3 ^1) * (5 ^1) * (7 ^1)$
Prime Factors: 2, 3, 5, 7
GCD = $(2 ^min(4, 3)) * (3 ^min(2, 1)) * (5 ^min(1, 1)) * (7 ^min(0, 1))$
GCD = $(2 ^3) * (3 ^1) * (5 ^1) * (7 ^0) = 8 * 3 * 5 * 1$

Method #3

3. The Euclidean algorithm

A more efficient method named after the ancient Greek mathematician Euclid. The Euclidean algorithm is based on the principle that the GCD of two numbers also divides their difference. This method involves a series of divisions where the divisor at each step becomes the dividend for the next step, and the remainder becomes the new divisor. This process is repeated until the remainder is zero, at which point the last non-zero divisor is the GCD.

Example 1: Greatest Common Divisor of 12 and 18.

Divide the larger (18) by the smaller (12) to obtain:

18 = 12 * 1 + 6

Example 1: Greatest Common Divisor of 12 and 18.

Example 1: Greatest Common Divisor of 12 and 18.

$$12 = 6 * 2 + 0$$

Example 1: Greatest Common Divisor of 12 and 18.

$$18 = 12 * 1 + 6$$

$$12 = 6 * 2 + 0$$

$$12 = |6| * 2 + 0$$

Example 1: Greatest Common Divisor of 12 and 18.

$$18 = \boxed{12} * 1 + \boxed{6}$$

$$12 = \boxed{6} * 2 + 0$$

$$12 = |6| * 2 + 0$$

$$GCD = 6$$

Example 2: Greatest Common Divisor of 720 and 840

$$840 = 720 * 1 + 120$$

Example 2: Greatest Common Divisor of 720 and 840

Example 2: Greatest Common Divisor of 720 and 840

$$720 = 120 * 6 + 0$$

Example 2: Greatest Common Divisor of 720 and 840

Example 2: Greatest Common Divisor of 720 and 840

$$720 = 120 * 6 + 0$$

$$GCD = 120$$

The Least Common Multiple

What is a multiple?

A multiple is a number that is obtained by multiplying another number by an integer

Thus, if a number A is the multiple of another number B, it means that when A is divided by B, there is no remainder.

What is the Least Common Multiple?

The Least Common Multiple (LCM) is the lowest multiple shared by two or more numbers.

The two common methods of finding the LCM include:

- 1. The Brute Force Method or Listing Method
- 2. The Prime Factorization Method

Method #1

1. The Brute Force Method

This method involves listing out all the multiples of each number and then finding the smallest number that appears on both lists. It's a straightforward approach but can be inefficient for large numbers.

Example 1: The LCM of 6 and 15

6: 6, 12, 18, 24, 30, 36, 42, 48, 54, 60, 66, 72, 78, 84, 90

15: 15, 30, 45, 60, 75, 90, 105, 120, 135, 150, 165, 180

Example 1: The LCM of 6 and 15

6:

30,

60,

15: 30, 60, 90

90

Example 1: The LCM of 6 and 15

6: 30, 60, 90

15: 30, 60, 90

Example 1: The LCM of 6 and 15

6: 30,

90

15: | 30, | 60, 90

LCM = 30

Example 2: The LCM of 320 and 480

320: 320, 640, 960, 1280, ...

480: 480, 960, 1440, ...

Example 2: The LCM of 320 and 480

320: 960

480: 960

Example 2: The LCM of 320 and 480

320: 960

480: 960

Example 2: The LCM of 320 and 480

320: 960

480: 960

LCM = 960

Method #2

2. The Prime Factorization Method

Here, each number is broken down into its prime factors. The LCM is then found by multiplying the largest power of the aggregated prime factors present in both numbers. This method provides a clear illustration of the common factors but can be time-consuming if the prime factorization is complicated.

Example 1: The LCM of 6 and 15

- 6: (2 ^ 1) * (3 ^ 1)
- 15: (3 ^ 1) * (5 ^ 1)

Example 1: The LCM of 6 and 15

6: (2 ^ 1) * (3 ^ 1)

15: (3 ^ 1) * (5 ^ 1)

Example 1: The LCM of 6 and 15

```
6: (2^{1}) * (3^{1})
```

$$LCM = (2 \land max(1,0)) * (3 \land max(1,1)) * (5 \land max(0,1))$$

Example 1: The LCM of 6 and 15

```
6: (2 ^ 1) * (3 ^ 1)
```

$$LCM = (2 \land max(1,0)) * (3 \land max(1,1)) * (5 \land max(0,1))$$

$$LCM = (2 ^1) * (3 ^1) * (5 ^1)$$

Example 1: The LCM of 6 and 15

```
6: (2 ^ 1) * (3 ^ 1)
```

$$LCM = (2 \land max(1,0)) * (3 \land max(1,1)) * (5 \land max(0,1))$$

$$LCM = (2 ^1) * (3 ^1) * (5 ^1) = 2 * 3 * 5$$

Example 1: The LCM of 6 and 15

LCM =
$$(2 \land max(1,0)) * (3 \land max(1,1)) * (5 \land max(0,1))$$

LCM = $(2 \land 1) * (3 \land 1) * (5 \land 1) = 2 * 3 * 5$
LCM = 30

Example 2: The LCM of 320 and 480

```
320: (2 ^ 6) * (5 ^ 1)
```

480: (2 ^ 5) * (3 ^ 1) * (5 ^ 1)

Example 2: The LCM of 320 and 480

Example 2: The LCM of 320 and 480

320: (2 ^ 6) * (5 ^ 1)

Prime Factors: 2, 3, 5

 $LCM = (2 \land max(6, 5)) * (3 \land max(0, 1)) * (5 \land max(1,1))$

Example 2: The LCM of 320 and 480

320: (2 ^ 6) * (5 ^ 1)

Example 2: The LCM of 320 and 480

320: (2 ^ 6) * (5 ^ 1)

Example 2: The LCM of 320 and 480

480: (2 ^ 5) * (3 ^ 1) * (5 ^ 1)

320: (2 ^ 6) * (5 ^ 1)

LCM = 960

Prime Factors: 2, 3, 5
$$LCM = (2 \land max(6, 5)) * (3 \land max(0, 1)) * (5 \land max(1, 1))$$

 $LCM = (2 \land 6) * (3 \land 1) * (5 \land 1) = 64 * 3 * 5$

Key Takeaways

- The Euclidean algorithm is the most efficient method for finding the GCD of two or more numbers.
- The Prime Factorization is the most efficient method for finding the LCM of two or more numbers.

Thank you!