

Adrift in the California Current:

Clustered drifting recorders describe spatial variation in soundscapes and marine mammal presence within offshore wind energy areas along the US West Coast

 **Anne Simonis**, Cory Hom-Weaver, Kourtney Burger, Kaitlin Palmer, Taiki Sakai, Shannon Rankin
NOAA Southwest Fisheries Science Center
asimonis@sfsu.edu


BACKGROUND:

Baseline data on marine ecosystems is needed to inform the planning and management of proposed offshore wind energy areas along the US West Coast. However, there is scarce historical data on the seasonal presence of protected species and overall sound levels in these regions due to the expense and challenges of surveying offshore wind energy areas.

METHODS

All field and analysis methods freely available:  <https://github.com/SAEL-SWFSC>

1. Collect passive acoustic recordings using clusters of 4-8 drifting buoys in Oregon, Humboldt, and Morro Bay wind energy areas
2. Describe seasonal and spatial variation in soundscapes and presence of 17 cetacean species including:
 - Baleen whales
 - blue, fin, gray, humpback, minke, Sei, Bryde's
 - Dolphins
 - Risso's, Pacific white-sided, unidentified
 - Beaked whales
 - Cuvier's, Baird's, Hubbs', Stejneger's, BW43, BWC
 - NBHF
 - Kogia spp., Dall's & harbor porpoise




This study was funded in part by the U.S. Department of the Interior, Bureau of Ocean Energy Management through Interagency Agreement M20PG00013 with the U.S. Department of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service (NMFS), Southwest Fisheries Science Center (SWFSC)


Clustered drifting recorders

describe spatial variation in soundscapes & habitat use.



2021-2023: 90 drifting buoy deployments



Seasonal Acoustic Scenes



Beaked whale presence: Morro Bay wind energy area



Collaborative cruises leverage resources & build community




Clustered deployments complement long-term stationary moorings

- Beaked whale density estimation
- Species habitat models
- Validate sound propagation models



Blue whale presence



Blue whale peak calling period in WEAs during post-upwelling, consistent with seafloor studies in other regions

Baleen whale localization accuracy improves with more sensors in the cluster



Take a picture to contact me about the project

