

2021 NOAA HPP Projects

Maya Philipp, Shannon Rankin, Alex Curtis

With special thanks to Cory Hom-Weaver, Emmanis Dorval, and Jack Barkowski



Humpback Whale Acoustics

- Automatic detectors can process large amounts of data
- Humpback whales are difficult to classify via detectors
- Large vocal range
- GPL Detector shows potential
- Create dataset of manual detections to compare GPL performance

All Drifts





Figure 1. Locations of DASBR deployments (black squares), retrievals or last known location (black triangles), and buoy drifts (recovered = black lines, lost = gray lines).


Img: simonis et all CCES2018








- 10 days of audio files from Monterey, CA
- Manually go through and annotate files for low frequency calls
- 150Hz-1KHz
- Identify humpbacks or unique sounds when possible
- 7000+ sounds manually boxed







California Sea Lion & Market Squid


- Determine if California Sea Lion diet data is useful in determining Market Squid stock
- Squid fishery not active constantly
- Population affected by environmental factors
 - El Nino
 - La Nina









California Sea Lion and Market Squid



- Looked for patterns with SSB total and CSL diet data
- Frequency of occurrence, fraction of species, and mean number of squid per scat
 - Calculated correlation and created linear models

SSB Total and Mean Squid from CSL Scat Analysis Linear Model Values for diff. Mortality rates (M)				
	Coefficients	Intercepts	R Squared	P-Value
M=0.01	2.809312	-9.243479	0.1974	0.0009672
M=0.15	3.344764	-14.35458	0.285	4.589e-05
M=0.30	3.504013	-17.26447	0.4651	6.839e-08





Thank you!

OCEANIL AND ATMOSPARIO ROMAN STRATION C.O. OF COMMERCE

If you have any questions, or would like to know more about anything, feel free to contact me at mphilipp@ucsd.edu