



## 2019 National Terrestrial Carbon Sinks Assessment for South Africa District Municipality Profile | Joe Gqabi

### Organic carbon pools




Figure 1. Split of organic carbon by carbon pool.

Organic carbon pools within the district of Joe Gqabi are estimated to be 295 Tg and distributed per carbon pool as depicted in the Figure (1) above. Most carbon is to be found as soil organic carbon (SOC). The models that estimate the proposition of carbon as above ground woody carbon (AGW) are based on satellite imagery and do not differentiate between natural woody vegetation or planted trees. Below ground woody (BGW) carbon is based on AGW and varies across the country based on climate. Above ground herb (AGH) and below ground herb (BGH) is based on both natural vegetation and crop fields, on the proportional split per 1 km<sup>2</sup> land unit. Litter is based on biome level estimates.

### Land cover classes in 2018 (from national land cover map)

Land cover is used to determine changes in carbon stocks, and land cover change is seen as one of the key drivers for change in terrestrial carbon. Data from three national land cover products, NLC 1990, NLC 2014 and NLC 2018 are compared against 17 (or 18) land cover classes. Only the 2018 land cover includes fallow land, which was mapped as natural land or agricultural land in earlier land cover products. The bare ground land cover classes was found to show large changes in extent between land cover products, possibly as a consequence of that years rainfall. Figure 2 gives a summary of land cover classes as mapped in the 2018 NLC. Table 1 summarises important changes in land cover for the district.

Joe Gqabi is the 34 highest rate of land lost to urbanisation between 1990 and 2018 out of the 52 South African districts, with 62 km<sup>2</sup> of natural or agricultural land converted to urban expansion.

Joe Gqabi has the 33 highest rate of land lost to irrigation between 1990 and 2018 out of the 52 South African districts.

Joe Gqabi lost 129 ha of dryland agricultural land between 1990 and 2018.




Figure 2. Estimates of the proportion of the district in each land cover class based on 2018 NLC data.

Table 1. Drivers of change based on national land cover data from 1990, 2014 and 2018

	Area in km2			% of total land area			Area lost (-) or gained (+) 1990 to 2018	Area lost (-) or gained (+) 1990 to 2018
	1990	2014	2018	1990	2014	2018	Km2	% of total area
Water	244	202	288	1	1	1	43	0
Wetlands	616	360	292	2	1	1	-324	-1
Indigenous Forest	1	2	9	0	0	0	8	0
Natural Vegetation	22024	22211	20288	86	87	79	-1736	-7
Commercial Agriculture	1283	1212	997	5	5	4	-285	-1
Pivot Agriculture	18	73	301	0	0	1	283	1
Orchards	9	9	2	0	0	0	-7	0
Viticulture	0	0	0	0	0	0	0	0
Pineapple	0	0	0	0	0	0	0	0
Subsistence Agriculture	510	506	521	2	2	2	12	0
Sugarcane Irrigated	0	0	0	0	0	0	0	0
Sugarcane Dry	0	0	0	0	0	0	0	0
Plantation Forest	164	306	421	1	1	2	256	1
Mines	6	5	4	0	0	0	-2	0
Bare degraded	342	345	1624	1	1	6	1282	5
Built-up	383	368	445	1	1	2	62	0
Fallow	0	0	406	0	0	2	406	2

### Loss in soil organic carbon (SOC) over time

Soil organic carbon loss is based on estimates of loss due to land cover change. Estimates of the total loss of SOC in the district based on both historic and recent land cover is given in table 2. In addition crude estimates are given on total carbon gains that might be possible from changed farming practices. Actual gains that can be realised will require local data based on local research, local crop choices, local farming practices and soil and climatic conditions.

Table 2. Estimated Soil Organic Carbon (SOC) loss due to agricultural activities and estimates of potential gains if all land cover was converted to conservation agriculture. Note: the extent to which conservation agriculture can restore SOC has not been established for the municipality, but will depend of crop choice, managements methods, soil type and climate.

	Estimated loss (-) or gain (+) in soil organic carbon due to land use. Tg C			Estimated potential 20 year gains from conservation agriculture, assuming a 25%, 50% or 75% of lost SOC is regained Tg C		
	Loss before 1990	Total loss by 2014*	Total loss by 2018*	25%	50%	75%
Commercial Agriculture	0.244	0.230	0.187	0.047	0.093	0.140
Pivot Agriculture	0.000	0.003	0.018	0.005	0.009	0.014
Orchards	0.001	0.001	0.000	0.000	0.000	0.000
Viticulture	0.000	0.000	0.000	0.000	0.000	0.000
Pineapple	0.000	0.000	0.000	0.000	0.000	0.000
Subsistence Agriculture	0.105	0.106	0.108	0.027	0.054	0.081
Sugarcane Irrigated	0.000	0.000	0.000	0.000	0.000	0.000
Sugarcane Dry	0.000	0.000	0.000	0.000	0.000	0.000
Bare degraded	0.003	0.002	0.004	0.001	0.002	0.003
Fallow	Not estimated in 1990 or 2014		0.019	0.047	0.038	0.057
Built-up	Zero loss was assumed based on IPCC recommendations, but in reality this will depend on land use within the build-up area					
Total	0.399	0.388	0.656	0.164	0.328	0.492

- Note, total change might reduce within a class over time because land in that class reverts to a different land use class. Change between 2014 and 2018 is due to the inclusion of fallow land losses in the 2018 data, and should not be interpreted as a change between those two years. The 1990 to 2014 change data should be used to best understand long-term rates of loss, as the methodology for these two data sets was very similar.