






(a) Primary Error Taxonomy (N=186)
Error Count
113
(60.8% errors)
[21.5% cohort]
73
(39.2% errors)
[13.9% cohort]
False Positives
(Benign as Malignant)
False Negatives
(Malignant as Benign)



(b) Error Counts by Lesion Type
False Positive
False Negative
Error Count
64
35
49
38
Mass Lesions
(N=267)
Calcifications
(N=258)



(c) Error Counts by Projection View
False Positive
False Negative
Error Count
52
33
61
40
CC View
(N=250)
MLO View
(N=275)


(d) Error Rate by Proposal IoU
Error Rate (%)
33.6%
(42/125)
41.9%
(36/86)
42.2%
(35/83)
31.0%
(31/100)
32.1%
(42/131)
0.0
(Miss)
0.01-0.1
(Weak)
0.1-0.3
(Partial)
0.3-0.5
(Good)
>0.5
(Strong)


(e) Decision Confidence (Correct vs Error)
Correct (Mean=0.92)
Error (Mean=0.92)
Density
Max Posterior Likelihood max(P)


(f) Subgroup Accuracy Profile
MLO View
63.27%
CC View
66.00%
Calc
66.28%
Mass
62.92%
Overall
64.57%
Subgroup Accuracy (%)


(g) Historical Error Trade-off Evolution
False Positives (FP)
False Negatives (FN)
Absolute Error Count
116
68
116
71
113
73
144
51
V5
V6
V8 (Final)
V9 (Ext.)


(h) Scientific Governance Audit
SUBGROUP AUDIT & GOVERNANCE
SAMPLE SIZES (LOCKED TEST):
• Total Cohort: N = 525 (100%)
• Mass Abnormalities: N = 267 (50.9%)
• Calcifications: N = 258 (49.1%)
• CC Views: N = 250 (47.6%)
• MLO Views: N = 275 (52.4%)
• True Benign: N = 310 (59.0%)
• True Malignant: N = 215 (41.0%)
METHODOLOGICAL CAVEATS:
1. Subgroup analyses are purely descriptive and observational.
2. No causal inference should be deduced solely from stratified error variances.
3. Zero post-hoc threshold tuning or subgroup re-calibration was permitted.