



Scan QR code to check
report authenticity



MC-2202

sterling
ACCURIS
Pathology lab that cares

Passport No :

LABORATORY TEST REPORT

Patient Information	Sample Information	Client/Location Information
Name : Lyubochka Svetka	Lab Id : 02232160XXXX	Client Name : Sterling Accuris Buddy
Sex/Age : Male / 41 Y 01-Feb-1982	Registration on : 20-Feb-2023 09:10	Location :
Ref. Id :	Collected at : non SAWPL	Approved on : 20-Feb-2023 12:33 Status : Final
Ref. By :	Collected on : 20-Feb-2023 08:53	Printed On : 28-Feb-2023 10:26
	Sample Type : Serum	Process At : 1. NRL SAWPL Gujarat Ahmedabad Paldi

Immunoassay

Test	Result	Unit	Biological Ref. Interval
25(OH) Vitamin D <small>CLIA</small>	8.98	ng/mL	Deficiency : <10 Insufficiency : 10 - 30 Sufficiency : 30 - 100 Toxicity : >100

Vitamin D is a fat soluble vitamin and exists in two main forms as cholecalciferol(vitamin D3) which is synthesized in skin from 7-dehydrocholesterol in response to sunlight exposure & Ergocalciferol(vitamin D2) present mainly in dietary sources.Both cholecalciferol & Ergocalciferol are converted to 25(OH)vitamin D in liver.

Interpretation:

Increased In

- Vitamin D intoxication
- Excessive exposure to sunlight

Decreased In

- Malabsorption
- Steatorrhea
- Dietary osteomalacia, anticonvulsant osteomalacia
- Biliary and portal cirrhosis
- Thyrotoxicosis
- Pancreatic insufficiency
- Celiac disease
- Rickets
- Alzheimer disease

Limitations:

More recently, it has become clear that receptors for vitamin D are present in a wide variety of cells and that this hormone has biologic effects extending beyond the control of mineral metabolism. Vitamin D deficiency is not clear. Levels needed to prevent rickets and osteomalacia (15 ng/mL) are lower than those that dramatically suppress parathyroid hormone levels (20–30 ng/mL). In turn, those levels are lower than levels needed to optimize intestinal calcium absorption (34 ng/mL). Neuromuscular peak performance is associated with levels approximately 38 ng/mL. A recent study states that increasing mean baseline levels from 29 to 38 ng/mL was associated with a 50% lower risk for colon cancer and levels of 52 ng/mL with a 50% reduction in the incidence of breast cancer. It is recommended to have clinical correlation with serum 25(OH)vitamin D, serum calcium, serum PTH & serum alkaline phosphatase.

Dhote.

DR. TEJASWINI DHOTE

M.D. Pathology

Dr. Sanjeev Shah

MD Path

Dr. Yash Shah

MD Path

This is an Electronically Authenticated Report.

Referred Test

Page 12 of 19

Sterling Accuris Pathology Laboratory

National Reference Laboratory: 101 to 109, First Floor, Sankalp Square II, Nr. Old Sharda Mandir Cross Road, B/s. Jalaram Mandir Railway Crossing, Ellisbridge, Ahmedabad, Gujarat- 380006, **Ph.: 812 813 0000**

Email: customer.care@sterlingaccuris.com | **Website:** www.sterlingaccuris.com