

CHAPTER-9
AREAS OF PARALLELOGRAMS AND TRIANGLES

1 Exercise 9.2

Q1. In the figure given below, $ABCD$ is a parallelogram, $AE \perp DC$ and $CF \perp AD$. If $AB = 16\text{cm}$, $AE = 8\text{cm}$ and $CF = 10\text{cm}$, find AD .

Construction




Figure 1: Parallelogram ABCD

The following table consists of given input parameters of the above parallelo-

gram ABCD :

Symbol	Value	Description
x	16cm	$\ \mathbf{B} - \mathbf{A}\ $
a	10cm	$\ \mathbf{F} - \mathbf{C}\ $
b	8cm	$\ \mathbf{E} - \mathbf{A}\ $
$\angle CFD$	90°	$CF \perp AD$
$\angle AED$	90°	$AE \perp CD$

Table 1: Parameters

Table below has the given input co-ordinates of the parallelogram :

Point	Co-ordinates
\mathbf{D}	(0,0)

Table 2: Co-ordinates

Following table are the unknown lengths and angles and their symbols:

Symbol	Description
c	$\ \mathbf{D} - \mathbf{C}\ $
r	$\ \mathbf{A} - \mathbf{D}\ $
d	$\ \mathbf{D} - \mathbf{E}\ $
b	$\ \mathbf{A} - \mathbf{E}\ $
θ	$\angle \mathbf{D}$

Table 3: Symbols and Corresponding Vectors

The point co-ordinates are derived in the following way :

1. **To derive the co-ordinates of C :**

As mentioned in the 3, $\|\mathbf{D} - \mathbf{C}\| = c$. In the above parallelogram it is given that $\|\mathbf{B} - \mathbf{A}\| = 16cm$. According to the properties of a parallelogram the parallel sides are equal in length. So, it can be said that :

$$\|\mathbf{B} - \mathbf{A}\| = \|\mathbf{D} - \mathbf{C}\| = c \quad (1)$$

$$(2)$$

As point \mathbf{C} lies on x axis, it can be expressed in the following way :

$$\mathbf{C} = c\mathbf{e}_1 \quad (3)$$

$$\mathbf{C} = c \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} c \\ 0 \end{pmatrix} \quad (4)$$

$$(5)$$

2. **To derive the co-ordinates of \mathbf{A} :**

\mathbf{A} can be expressed in the form of $r \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix}$. In order to obtain r and θ , the following can be done :

(a) **To find out θ :**

To find out θ , let us assume that $\|\mathbf{C} - \mathbf{F}\| = a$

$$\text{from } \triangle CFD, \quad (6)$$

$$\sin \theta = \frac{a}{c} \quad (7)$$

$$\implies \theta = \sin^{-1} \frac{a}{c} \quad (8)$$

$$(9)$$

(b) **To find out r :**

As mentioned in 3, $\|\mathbf{D} - \mathbf{A}\| = r$ and $\|\mathbf{E} - \mathbf{A}\| = b$. In order to find out r ,

$$\text{from } \triangle ADE, \quad (10)$$

$$\sin \theta = \frac{b}{r} \quad (11)$$

$$r = \frac{b}{\sin \theta} \quad (12)$$

$$(13)$$

So, the co-ordinates of \mathbf{A} can be written as :

$$\mathbf{A} = \frac{b}{\sin \theta} \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} \quad (14)$$

$$\mathbf{A} = \begin{pmatrix} b \cot \theta \\ b \end{pmatrix} \quad (15)$$

$$(16)$$

3. **To derive the co-ordinates of \mathbf{B} :**

From parallelogram law of vectors, \mathbf{B} can be expressed as the sum of \mathbf{A}

and \mathbf{C} . So, it can be written as,

$$\mathbf{B} = \mathbf{A} + \mathbf{C} \quad (17)$$

$$(18)$$

4. **To derive the co-ordinates of \mathbf{E} :**

As mentioned in the table 3, $\|\mathbf{D} - \mathbf{E}\| = d$. As, \mathbf{E} lies on x-axis it can be written in the form of $d\mathbf{e}_1$. So, the co-ordinates can be found out in the following way :

$$\text{from } \triangle DAE, \quad (19)$$

$$\cos \theta = \frac{d}{r} \quad (20)$$

$$d = r \cos \theta \quad (21)$$

$$(22)$$

$$\mathbf{E} = d\mathbf{e}_1 = r \cos \theta \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} r \cos \theta \\ 0 \end{pmatrix}.$$

5. **To derive the co-ordinates of \mathbf{F} :**

As point \mathbf{F} divides \mathbf{AD} in the ratio $k : 1$. The co-ordinates of \mathbf{F} can be found out in the following way :

$$\mathbf{F} = \frac{k\mathbf{A} + \mathbf{D}}{k + 1} \quad (23)$$

$$(24)$$

The following table displays the unknown lengths and angles which were derived from the given quantities :

Symbol	value	Description
c	x	$\ \mathbf{D} - \mathbf{C}\ $
r	$\frac{b}{\sin \theta}$	$\ \mathbf{D} - \mathbf{A}\ $
d	$r \cos \theta$	$\ \mathbf{D} - \mathbf{E}\ $
θ	$\sin^{-1} \frac{a}{c}$	$\angle D$

Table 4: Co-ordinates in terms of given and derived lengths and angles

The following table displays the point co-ordinates in terms of known and de-

rived quantities:

Point	Co-ordinates
A	$\begin{pmatrix} b \cot \theta \\ b \end{pmatrix}$
B	$\mathbf{A} + \mathbf{C}$
C	$\begin{pmatrix} c \\ 0 \end{pmatrix}$
E	$\begin{pmatrix} r \cos \theta \\ 0 \end{pmatrix}$
F	$\frac{k\mathbf{A}+\mathbf{D}}{k+1}$

Table 5: Co-ordinates in terms of known and derived quantities

Finding the co-ordinates of the above parallelogram:

1. **Co-ordinates of C:**

As $c = \|\mathbf{D} - \mathbf{C}\| = 16$, the co-ordinates of **C** are:

$$\mathbf{C} = \begin{pmatrix} c \\ 0 \end{pmatrix} = \begin{pmatrix} 16 \\ 0 \end{pmatrix} \quad (25)$$

$$(26)$$

So, the co-ordinates of **C** are $\begin{pmatrix} 16 \\ 0 \end{pmatrix}$.

2. **Co-ordinates of A:**

(a) **Finding θ :**

$$\theta = \sin^{-1} \frac{a}{c} \quad (27)$$

$$\theta = \sin^{-1} \frac{10}{16} \quad (28)$$

$$\theta = 38.68^\circ \quad (29)$$

$$(30)$$

From the above derivations, we got that:

$$\mathbf{A} = \begin{pmatrix} b \cot \theta \\ b \end{pmatrix} \quad (31)$$

$$\Rightarrow \begin{pmatrix} 8 \cot 38.68 \\ 8 \end{pmatrix} \quad (32)$$

$$\mathbf{A} = \begin{pmatrix} 10 \\ 8 \end{pmatrix} \quad (33)$$

$$(34)$$

so, the co-ordinates of \mathbf{A} are $\begin{pmatrix} 10 \\ 8 \end{pmatrix}$.

3. Co-ordinates of \mathbf{B} :

From above derivation, we got that $\mathbf{B} = \mathbf{A} + \mathbf{C}$. Then, \mathbf{C} is:

$$\mathbf{B} = \begin{pmatrix} 10 \\ 8 \end{pmatrix} + \begin{pmatrix} 16 \\ 0 \end{pmatrix} \quad (35)$$

$$\mathbf{B} = \begin{pmatrix} 26 \\ 8 \end{pmatrix} \quad (36)$$

$$(37)$$

So, the co-ordinates of \mathbf{B} are $\begin{pmatrix} 26 \\ 8 \end{pmatrix}$.

4. Co-ordinates of \mathbf{E} :

(a) Finding r :

From above derivation, r can be found out in the following way:

$$r = \frac{b}{\sin \theta} \quad (38)$$

$$r = \frac{8}{\sin 38.68} \quad (39)$$

$$r = 12.8 \text{ cm} \quad (40)$$

$$(41)$$

The co-ordinates of \mathbf{E} are:

$$\mathbf{E} = \begin{pmatrix} r \cos \theta \\ 0 \end{pmatrix} \quad (42)$$

$$\Rightarrow \begin{pmatrix} (12.8) \cos 38.68 \\ 0 \end{pmatrix} \quad (43)$$

$$\mathbf{E} = \begin{pmatrix} 10 \\ 0 \end{pmatrix} \quad (44)$$

$$(45)$$

So, the co-ordinates for **E** are $\begin{pmatrix} 10 \\ 0 \end{pmatrix}$.

5. **Co-ordinates for F:**

F divides **AD** in the ratio 39 : 1. So, the co-ordinates of **F** are:

$$\mathbf{F} = \frac{k\mathbf{A} + \mathbf{D}}{k + 1} \quad (46)$$

$$\Rightarrow \frac{(39) \begin{pmatrix} 10 \\ 8 \end{pmatrix} + \begin{pmatrix} 0 \\ 0 \end{pmatrix}}{39 + 1} \quad (47)$$

$$\Rightarrow \frac{\begin{pmatrix} 390 \\ 312 \end{pmatrix}}{40} \quad (48)$$

$$\mathbf{F} = \begin{pmatrix} 9.75 \\ 7.8 \end{pmatrix} \quad (49)$$

So, the co-ordinates of **F** are $\begin{pmatrix} 9.75 \\ 7.8 \end{pmatrix}$.

The following table displays the final co-ordinates of the vertices of the parallelogram :

Point	Co-ordinates
A	$\begin{pmatrix} 10 \\ 8 \end{pmatrix}$
B	$\begin{pmatrix} 26 \\ 8 \end{pmatrix}$
C	$\begin{pmatrix} 16 \\ 0 \end{pmatrix}$
D	$\begin{pmatrix} 0 \\ 0 \end{pmatrix}$
E	$\begin{pmatrix} 10 \\ 0 \end{pmatrix}$
F	$\begin{pmatrix} 9.75 \\ 7.8 \end{pmatrix}$

Table 6: Final co-ordinates of the parallelogram

The length of $\|\mathbf{D} - \mathbf{A}\|$ was found out in the above process and it is $r = 12.8\text{cm}$.