

On topological solutions to a generalized Chern-Simons equation on lattice graphs

Songbo Hou*

Department of Applied Mathematics, College of Science, China Agricultural University, Beijing, 100083, P.R. China

Xiaoqing Kong

Department of Applied Mathematics, College of Science, China Agricultural University, Beijing, 100083, P.R. China

Abstract

In this paper, we study a generalized self-dual Chern-Simons equation on the lattice graph \mathbb{Z}^n for $n \geq 2$ as given by

$$\Delta u = \lambda e^u (e^u - 1)^{2p+1} + 4\pi \sum_{j=1}^M n_j \delta_{p_j},$$

where Δ denotes the Laplacian operator, λ is a positive constant, p is a non-negative integer, n_1, \dots, n_M are positive integers, and p_1, \dots, p_M are distinct points with δ_{p_j} representing the Dirac delta function at p_j . We establish the existence of a topological solution that is maximal among all possible solutions. Our findings extend those of of Hua et al. [arXiv:2310.13905], Chao and Hou [J. Math. Anal. Appl. **519**(1), 126787(2023)], and Hou and Qiao [J. Math. Phys. **65**(8), 081503(2024)].

Keywords: Chern-Simons equation, lattice graph, topological solution, maximal solution
2020 MSC: 35A01 35A16 35J91 35R02

1. Introduction

The Chern-Simons theory was indeed proposed by Shiing-Shen Chern and James Simons in 1974. This theory was initially developed within the field of mathematics to study the geometric structures on three-dimensional manifolds. Later, it found broad applications in physics, particularly in quantum physics and condensed matter physics. The theory quickly gained significance in understanding topological phase transitions, especially in the context of quantum Hall effects, topological insulators, and high-temperature superconductors [8, 44, 45].

Considerable research has been conducted in the area of field theory models influenced by Chern-Simons type dynamics, such as [34, 35, 41] and so on. From a mathematical perspective, the equations of motion for various Chern-Simons models pose significant analytical challenges, even in scenarios involving radial symmetry and static conditions [32]. The discovery of the

*Corresponding author: Songbo Hou

Email addresses: housb@cau.edu.cn (Songbo Hou), kxq@cau.edu.cn (Xiaoqing Kong)

Preprint submitted to ***

October 25, 2024

self-dual structure in the Abelian Chern-Simons model in [21, 31] has inspired a lot of subsequent research. Many problems of existence can be transformed into studies of elliptic partial differential equations or systems of equations, with a particular focus on exploring topological and non-topological solutions [3, 4, 18, 19, 29, 49].

Partial differential equations on discrete graphs have recently garnered significant interest and are now widely applied in diverse fields such as image processing, social network analysis, bioinformatics, and machine learning. Recent advancements have extended some traditional methods for solving partial differential equations in Euclidean spaces to the context of graphs.

In seminal works by Grigor'yan et al. [15–17], a variational approach was developed to study Yamabe type equations, the Kazdan-Warner equation, and some nonlinear equations. These studies primarily focused on establishing the existence of solutions. Subsequent research has extensively explored various types of partial differential equations on graphs. Notably, the existence results for Yamabe type equations have been detailed in references such as [11, 13, 52]. Studies addressing Kazdan-Warner equations are found in [10, 12, 37, 47, 51], and the results concerning Schrödinger equations appear in [5, 40, 43, 50]. Additionally, significant findings related to the heat equations are documented in [36, 38].

In this paper, we study a generalized self-dual Chern-Simons equation given by

$$\Delta u = \lambda e^u (e^u - 1)^{2p+1} + 4\pi \sum_{j=1}^M n_j \delta_{p_j}, \quad (1.1)$$

where Δ denotes the Laplacian operator, λ is a positive constant, p is a non-negative integer, n_1, \dots, n_M are positive integers, and p_1, \dots, p_M are distinct points with δ_{p_j} being the Dirac delta function at p_j . In the case of Euclidean spaces, a solution $u(x)$ to equation (1.1) is classified as topological if $u(x) \rightarrow 0$ as $|x| \rightarrow +\infty$, and non-topological if $u(x) \rightarrow -\infty$ as $|x| \rightarrow +\infty$.

When $p = 0$ and $n_i = 1$ (for $i = 1, 2, \dots, M$), Eq.(1.1) reduces to

$$\Delta u = \lambda e^u (e^u - 1) + 4\pi \sum_{j=1}^M \delta_{p_j}. \quad (1.2)$$

Caffarelli et al. [2] and Tarantello [48] investigated Eq.(1.2) in a doubly periodic region or the 2-torus in \mathbb{R}^2 and established the existence of solutions. Huang et al. [27] and Hou et al. [24] studied Eq.(1.2) on finite graphs and obtained the existence of solutions. When $p = 2$ and $n_i = 1$ (for $i = 1, 2, \dots, M$), Eq.(1.1) is reduced to

$$\Delta u = \lambda e^u (e^u - 1)^5 + 4\pi \sum_{j=1}^M \delta_{p_j}. \quad (1.3)$$

Han [20] established the existence of multi-vortices for Eq.(1.3) over a doubly periodic region in \mathbb{R}^2 . Chao et al. [6] and Hu [25] obtained multiple solutions to Eq.(1.3) on finite graphs. Further studies on the Chern-Simons models on graphs include [7, 9, 22, 23, 26, 30, 33, 39].

Next, we introduce the lattice \mathbb{Z}^n for $n \geq 2$. The lattice \mathbb{Z}^n is a discrete graph which consists of the vertex set V and the edge set E :

$$V = \mathbb{Z}^n = \{x = (x_1, \dots, x_n) \in \mathbb{R}^n : x_i \in \mathbb{Z} \text{ for all } i \in \{1, \dots, n\}\},$$

$$E = \left\{ \{x, y\} : x, y \in V \text{ with } \sum_{i=1}^n |x_i - y_i| = 1 \right\}.$$

Write $x \sim y$ if $\{x, y\}$ is an edge in E . The distance on \mathbb{Z}^n is defined by

$$d(x, y) = \sum_{i=1}^n |x_i - y_i|, \quad x, y \in \mathbb{Z}^n,$$

with $d(x) = d(x, 0)$ representing the distance from the origin.

For a finite subset $\Omega \subset \mathbb{Z}^n$, the boundary of Ω is delineated by

$$\delta\Omega := \{y \in \mathbb{Z}^n \setminus \Omega : \exists x \in \Omega \text{ such that } y \sim x\},$$

and the closure of Ω is designated as $\bar{\Omega} = \Omega \cup \delta\Omega$.

Now, we introduce function spaces and operators on graphs to prepare for the subsequent analysis. Denote $C(\mathbb{Z}^n)$ as the set of functions defined on \mathbb{Z}^n . For a finite subset $\Omega \subset \mathbb{Z}^n$, we define $C(\Omega)$ as the set of functions defined on Ω . The l^p -norm ($1 \leq p < \infty$) of $u \in C(\mathbb{Z}^n)$ is defined as

$$\|u\|_{l^p(\mathbb{Z}^n)} = \left(\sum_{x \in \mathbb{Z}^n} |u(x)|^p \right)^{\frac{1}{p}},$$

and the l^∞ -norm of $u \in C(\mathbb{Z}^n)$ is defined as

$$\|u\|_{l^\infty(\mathbb{Z}^n)} = \sup_{x \in \mathbb{Z}^n} |u(x)|.$$

We also define the following seminorm:

$$|u|_{1,p} := \left(\sum_{x \in \mathbb{Z}^n} \sum_{y \sim x} |u(y) - u(x)|^p \right)^{\frac{1}{p}}.$$

For any $u \in C(\mathbb{Z}^n)$, the Laplacian operator is expressed as

$$\Delta u(x) = \sum_{d(x,y)=1} (u(y) - u(x)).$$

Additionally, the difference operator is specified by

$$\nabla_{xy} u = u(y) - u(x), \quad \text{for any } u \in C(\mathbb{Z}^n) \text{ and } x, y \in \mathbb{Z}^n.$$

For functions $u, v \in C(\bar{\Omega})$, a bilinear form is introduced:

$$E_\Omega(u, v) := \frac{1}{2} \sum_{\substack{x, y \in \Omega \\ x \sim y}} \nabla_{xy} u \nabla_{xy} v + \sum_{\substack{x \in \Omega, y \in \delta\Omega \\ x \sim y}} \nabla_{xy} u \nabla_{xy} v, \quad (1.4)$$

and the Dirichlet energy of u on Ω is defined as $E_\Omega(u) = E_\Omega(u, u)$. The directional derivative operator $\frac{\partial u}{\partial \mathbf{n}}$ at $x \in \delta\Omega$ is

$$\frac{\partial u}{\partial \mathbf{n}}(x) := \sum_{\substack{y \in \Omega \\ x \sim y}} (u(x) - u(y)).$$

There also holds Green's identity [14]:

$$E_\Omega(u, v) = - \sum_{x \in \Omega} u(x) \Delta v(x) + \sum_{x \in \delta\Omega} u(x) \frac{\partial v}{\partial \mathbf{n}}(x). \quad (1.5)$$

In this study, we focus primarily on topological solutions to the generalized Chern-Simons equation defined on \mathbb{Z}^n :

$$\begin{cases} \Delta u = \lambda e^u (e^u - 1)^{2p+1} + 4\pi \sum_{j=1}^M n_j \delta_{p_j}, \\ \lim_{d(x) \rightarrow +\infty} u(x) = 0, \end{cases} \quad (1.6)$$

where we aim to construct a topological solution that is maximal among all possible solutions. Our main result is presented as follows.

Theorem 1.1. *The equation (1.6) admits a topological solution $u \in l^{2p+2}(\mathbb{Z}^n)$ on \mathbb{Z}^n for $n \geq 2$, which is maximal among all possible solutions.*

In our proof, we employ methods from [26, 46]. First, we establish an iterative scheme, which yields a monotone sequence $\{u_k\}$ addressing the Dirichlet problem. Subsequently, we introduce the functional $J_\Omega(u)$ and demonstrate that $J_\Omega(u_k)$ is uniformly bounded. By using the discrete Gagliardo-Nirenberg-Sobolev inequality, we ensure that the sequence $\{\|u_k\|_{l^{2p+2}(\Omega)}\}$ is uniformly bounded. As we pass to the limit, we ascertain a solution on Ω . Utilizing the exhaustion method, we extend this solution to \mathbb{Z}^n . Additionally, we establish the maximality of the solution.

2. Proof of Theorem 1.1

Initially, we recall a foundational result: the maximum principle.

Lemma 2.1. [26] *Let Ω be a finite subset of \mathbb{Z}^n . Consider any positive function $f \in C(\overline{\Omega})$. Suppose that a function $v \in C(\overline{\Omega})$ satisfies the following conditions:*

$$\begin{cases} (\Delta - f)v \geq 0 & \text{on } \Omega, \\ v \leq 0 & \text{on } \partial\Omega. \end{cases}$$

Then, it follows that $v \leq 0$ on $\overline{\Omega}$.

Next, we use the maximum principle to establish the iterative sequence. Let Ω_0 be a finite subset of \mathbb{Z}^n such that $\Omega_0 \supset \{p_j\}_{j=1}^M$. Additionally, let Ω be an arbitrary connected finite subset satisfying $\Omega_0 \subset \Omega \subset \mathbb{Z}^n$. Let $g = 4\pi \sum_{j=1}^M n_j \delta_{p_j}$ and $N = 4\pi \sum_{j=1}^M n_j$. Choose a constant $K > (2p+2)\lambda > 0$. Set $u_0 = 0$ and consider the following iterative equations:

$$\begin{cases} (\Delta - K)u_k = \lambda e^{u_{k-1}} (e^{u_{k-1}} - 1)^{2p+1} + g - Ku_{k-1} & \text{on } \Omega, \\ u_k = 0 & \text{on } \partial\Omega. \end{cases} \quad (2.1)$$

Lemma 2.2. *Let the sequence $\{u_k\}$ be defined as in (2.1). Then, for each k , u_k is uniquely defined and satisfies*

$$0 = u_0 \geq u_1 \geq u_2 \geq \dots$$

Proof. First, we establish the following equation for u_1 :

$$\begin{cases} (\Delta - K)u_1 = g & \text{on } \Omega, \\ u_1 = 0 & \text{on } \partial\Omega. \end{cases} \quad (2.2)$$

By the variation method in Lemma 2.2 in [22], we get that (2.2) admits a unique solution. Utilizing Lemma 2.1, it follows that $u_1 \leq 0$.

Assuming that

$$0 = u_0 \geq u_1 \geq u_2 \geq \dots \geq u_i,$$

and given that

$$\lambda e^{u_i} (e^{u_i} - 1)^{2p+1} + g - Ku_i \in l^2(\Omega),$$

we can also guarantee the existence and uniqueness of u_{i+1} by the variation method again.

Analyzing the iterative equation (2.1), we derive that

$$\begin{aligned} (\Delta - K)(u_{i+1} - u_i) &= \lambda e^{u_i} (e^{u_i} - 1)^{2p+1} - \lambda e^{u_{i-1}} (e^{u_{i-1}} - 1)^{2p+1} - K(u_i - u_{i-1}) \\ &\geq \lambda e^\omega (e^\omega - 1)^{2p} [(2p+2)e^\omega - 1] (u_i - u_{i-1}) - K(u_i - u_{i-1}) \\ &\geq K(e^{2\omega} - 1)(u_i - u_{i-1}) \geq 0, \end{aligned}$$

where ω is a function satisfying $u_i \leq \omega \leq u_{i-1}$. This indicates that $u_{i+1} \leq u_i$, thereby affirming the lemma using Lemma 2.1. \square

Next, we define the natural functional $J_\Omega(u)$ and demonstrate that $J_\Omega(u_k)$ decreases as k increases. Utilizing Green's identities, we introduce the following functional defined over Ω :

$$J_\Omega(u) = \frac{1}{2} E_\Omega(u) + \sum_{x \in \Omega} \left[\frac{\lambda}{2p+2} (e^{u(x)} - 1)^{2p+2} + g(x)u(x) \right]. \quad (2.3)$$

Lemma 2.3. *Let $\{u_k\}$ be the sequence defined in Eq.(2.1). Then, the following inequality holds:*

$$C \geq J_\Omega(u_1) \geq J_\Omega(u_2) \geq \dots \geq J_\Omega(u_k) \geq \dots,$$

where the constant C depends only on n , λ , p and N .

Proof. By multiplying Eq.(2.1) by the difference $u_k - u_{k-1}$ and performing a summation across the domain Ω , we derive:

$$\begin{aligned} &\sum_{x \in \Omega} (\Delta - K) u_k(x) [u_k(x) - u_{k-1}(x)] \\ &= \sum_{x \in \Omega} \left[\lambda e^{u_{k-1}} (e^{u_{k-1}} - 1)^{2p+1} (u_k - u_{k-1}) - K u_{k-1} (u_k - u_{k-1}) + g(u_k - u_{k-1}) \right] (x). \end{aligned} \quad (2.4)$$

Utilizing the Green's identity as specified in (1.5), we have

$$\sum_{x \in \Omega} \Delta u_k(x) (u_k(x) - u_{k-1}(x)) = -E_\Omega(u_k - u_{k-1}, u_k) = -E_\Omega(u_k) + E_\Omega(u_{k-1}, u_k). \quad (2.5)$$

Integrating this with equation (2.4), we conclude

$$\begin{aligned} &E_\Omega(u_k) - E_\Omega(u_{k-1}, u_k) + \sum_{x \in \Omega} K (u_k(x) - u_{k-1}(x))^2 \\ &= - \sum_{x \in \Omega} \left[\lambda e^{u_{k-1}} (e^{u_{k-1}} - 1)^{2p+1} (u_k - u_{k-1}) + g(x) (u_k - u_{k-1}) \right]. \end{aligned} \quad (2.6)$$

Now we introduce a concave function for $x \leq 0$:

$$h(x) = \frac{\lambda}{2p+2} (e^x - 1)^{2p+2} - \frac{K}{2} x^2.$$

It is easy to see that

$$h(u_{k-1}) - h(u_k) \geq h'(u_{k-1})(u_{k-1} - u_k) = \left[\lambda (e^{u_{k-1}} - 1)^{2p+1} e^{u_{k-1}} - K u_{k-1} \right] (u_{k-1} - u_k),$$

which yields that

$$\begin{aligned} \frac{\lambda}{2p+2} (e^{u_k} - 1)^{2p+2} &\leq \frac{\lambda}{2p+2} (e^{u_{k-1}} - 1)^{2p+2} + \frac{K}{2} (u_k - u_{k-1})^2 \\ &\quad + \lambda e^{u_{k-1}} (e^{u_{k-1}} - 1)^{2p+1} (u_k - u_{k-1}). \end{aligned} \quad (2.7)$$

It follows from (1.4) that

$$\begin{aligned} |E_\Omega(u_{k-1}, u_k)| &\leq \frac{1}{2} \sum_{\substack{x, y \in \Omega \\ x \sim y}} |\nabla_{xy} u_{k-1} \nabla_{xy} u_k| + \sum_{\substack{x \in \Omega, y \in \partial\Omega \\ x \sim y}} |\nabla_{xy} u_{k-1} \nabla_{xy} u_k| \\ &\leq \frac{1}{4} \sum_{\substack{x, y \in \Omega \\ x \sim y}} (|\nabla_{xy} u_{k-1}|^2 + |\nabla_{xy} u_k|^2) + \frac{1}{2} \sum_{\substack{x \in \Omega, y \in \partial\Omega \\ x \sim y}} (|\nabla_{xy} u_{k-1}|^2 + |\nabla_{xy} u_k|^2) \\ &= \frac{1}{2} E_\Omega(u_{k-1}) + \frac{1}{2} E_\Omega(u_k). \end{aligned} \quad (2.8)$$

Combining (2.6), (2.7) and (2.8), we conclude that

$$J_\Omega(u_k) \leq J_\Omega(u_k) + \frac{K}{2} \|u_{k-1} - u_k\|_{L^2(\Omega)}^2 \leq J_\Omega(u_{k-1}).$$

Next, we estimate the upper bound for $J_\Omega(u_1)$. Noting that

$$\begin{aligned} E_\Omega(u_1) &= \frac{1}{2} \sum_{\substack{x, y \in \Omega \\ x \sim y}} |\nabla_{xy} u_1|^2 + \sum_{\substack{x \in \Omega, y \in \partial\Omega \\ x \sim y}} |\nabla_{xy} u_1|^2 \\ &\leq \sum_{\substack{x, y \in \Omega \\ x \sim y}} (u_1(x)^2 + u_1(y)^2) + 2 \sum_{\substack{x \in \Omega, y \in \partial\Omega \\ x \sim y}} (u_1(x)^2 + u_1(y)^2) \\ &\leq 4n \|u_1\|_{L^2(\Omega)}^2, \end{aligned}$$

and $|e^{u_1} - 1| = 1 - e^{u_1} \leq -u_1$, we get

$$\begin{aligned} J_\Omega(u_1) &\leq \frac{1}{2} \cdot 4n \|u_1\|_{L^2(\Omega)}^2 + \frac{\lambda}{2p+2} \sum_{x \in \Omega} u_1(x)^{2p+2} + \frac{1}{2} \sum_{x \in \Omega} [g(x)^2 + u_1(x)^2] \\ &= c_1 + c_2 (\|u_1\|_{L^2(\Omega)}^2 + \|u_1\|_{L^{2p+2}(\Omega)}^{2p+2}), \end{aligned}$$

where c_1, c_2 are constants that only depend on n, p, λ and N .

Multiplying (2.2) by u_1 and summing over Ω , we have

$$E_\Omega(u_1) + K \sum_{x \in \Omega} u_1(x)^2 = - \sum_{x \in \Omega} g(x) u_1(x).$$

This yields

$$K \sum_{x \in \Omega} u_1(x)^2 \leq \frac{1}{2K} \sum_{x \in \Omega} g(x)^2 + \frac{K}{2} \sum_{x \in \Omega} u_1(x)^2.$$

Hence,

$$\sum_{x \in \Omega} u_1(x)^2 \leq \frac{\|g\|_{l^2(\mathbb{Z}^n)}^2}{K^2},$$

which yields

$$\sum_{x \in \Omega} u_1(x)^{2p+2} \leq \left(\sum_{x \in \Omega} u_1(x)^2 \right)^{p+1} \leq \left(\frac{\|g\|_{l^2(\mathbb{Z}^n)}^2}{K^2} \right)^{p+1}.$$

We derive that $J_\Omega(u_1) \leq C$, where C depends only on n, λ, p and N , and complete the proof. \square

Next, utilizing the discrete Gagliardo-Nirenberg-Sobolev inequality along with Lemma 2.3, we establish the bound of $\|u_k\|_{l^{2p+2}(\Omega)}$. The proof of Theorem 4.1 in [42] provides the following discrete Gagliardo-Nirenberg-Sobolev inequality:

Lemma 2.4. [42] Let $n \geq 2, q > 1, \gamma \geq q$ and $q' = \frac{q}{q-1}$. Then for any $u \in l^q(\mathbb{Z}^n)$, we have

$$\|u\|_{l^{\frac{\gamma n}{n-1}}(\mathbb{Z}^n)}^\gamma \leq C(q, n, \gamma) \|u\|_{1,q} \|u\|_{l^{(\gamma-1)q'}(\mathbb{Z}^n)}^{\gamma-1}.$$

Lemma 2.1 in [28] implies that

$$\|u\|_{l^{p''}(\mathbb{Z}^n)} \leq \|u\|_{l^{p'}(\mathbb{Z}^n)},$$

for any $p'' \geq p'$. Letting $\gamma = 2(p+1)$, $q = 2$ and $q' = 2$, we have for $u \in l^2(\mathbb{Z}^n)$,

$$\|u\|_{l^{4p+4}(\mathbb{Z}^n)} \leq \|u\|_{l^{\frac{2n(p+1)}{n-1}}(\mathbb{Z}^n)} \leq C(n, p) \|u\|_{1,2}^{\frac{1}{2p+2}} \|u\|_{l^{4p+2}(\mathbb{Z}^n)}^{\frac{2p+1}{2p+2}}. \quad (2.9)$$

Lemma 2.5. Let $\{u_k\}$ be the sequence defined by Eq.(2.1). For any $k \geq 1$, we have

$$\|u_k\|_{l^{2p+2}(\Omega)} \leq C_2 (J_\Omega(u_k) + 1) \leq C_1, \quad (2.10)$$

where C_2 and C_1 depend only on n, λ, p and N .

Proof. Let \tilde{u}_k denote the null extension of u_k to \mathbb{Z}^n as follows:

$$\tilde{u}_k(x) = \begin{cases} u_k(x) & \text{on } \Omega, \\ 0 & \text{on } \Omega^c. \end{cases} \quad (2.11)$$

It is evident that $\tilde{u}_k \in l^2(\mathbb{Z}^n)$. By (2.9), the following inequality holds:

$$\|\tilde{u}_k\|_{l^{4p+4}(\mathbb{Z}^n)}^{2p+2} \leq (C(n, p))^{2p+2} \|\tilde{u}_k\|_{1,2} \|\tilde{u}_k\|_{l^{4p+2}(\mathbb{Z}^n)}^{2p+1}.$$

Referring to (2.11), we obtain:

$$\begin{aligned} \|\tilde{u}_k\|_{l^{4p+4}(\mathbb{Z}^n)}^{4p+4} &= \sum_{x \in \Omega} u_k(x)^{4p+4}, \\ \|\tilde{u}_k\|_{l^{4p+2}(\mathbb{Z}^n)}^{4p+2} &= \sum_{x \in \Omega} u_k(x)^{4p+2}, \end{aligned}$$

and

$$|\tilde{u}_k|_{1,2} \leq (2E_\Omega(u_k))^{\frac{1}{2}}.$$

Thus, it follows that:

$$\sum_{x \in \Omega} u_k(x)^{4p+4} \leq C_3 E_\Omega(u_k) \sum_{x \in \Omega} u_k(x)^{4p+2}, \quad (2.12)$$

where $C_3 = 2(C(n, p))^{4p+4}$.

Using the identity $1 - e^{u_k} = 1 - e^{-|u_k|} = \frac{e^{|u_k|} - 1}{e^{|u_k|}} \geq \frac{|u_k|}{1 + |u_k|}$, and (2.3), we deduce:

$$\begin{aligned} J_\Omega(u_k) &= \frac{1}{2} E_\Omega(u_k) + \sum_{x \in \Omega} \left[\frac{\lambda}{2p+2} (e^{u_k(x)} - 1)^{2p+2} + g(x) u_k(x) \right] \\ &\geq \frac{1}{2} E_\Omega(u_k) + \frac{\lambda}{2p+2} \sum_{x \in \Omega} \left(\frac{|u_k(x)|}{1 + |u_k(x)|} \right)^{2p+2} - \|g\|_{\frac{4p+4}{4p+3}(\Omega)} \|u_k\|_{L^{4p+4}(\Omega)} \\ &\geq \frac{1}{2} E_\Omega(u_k) + \frac{\lambda}{2p+2} \sum_{x \in \Omega} \left(\frac{|u_k(x)|}{1 + |u_k(x)|} \right)^{2p+2} - C_4 (E_\Omega(u_k))^{\frac{1}{4p+4}} \left(\sum_{x \in \Omega} u_k(x)^{4p+2} \right)^{\frac{1}{4p+4}}, \end{aligned}$$

where C_4 is a uniform constant depending only on n, p and N .

Let $\epsilon > 0$ is a constant to be chosen later. By Yung's inequality, we have

$$\begin{aligned} &C_4 (E_\Omega(u_k))^{\frac{1}{4p+4}} \left(\sum_{x \in \Omega} u_k(x)^{4p+2} \right)^{\frac{1}{4p+4}} \\ &= \left[C_4 \epsilon^{-\frac{2p+1}{2p+2}} \left(\frac{2p+2}{2p+1} \right)^{-\frac{2p+1}{2p+2}} (E_\Omega(u_k))^{\frac{1}{4p+4}} \right] \left[\epsilon^{\frac{2p+1}{2p+2}} \left(\frac{2p+2}{2p+1} \right)^{\frac{2p+1}{2p+2}} \left(\sum_{x \in \Omega} u_k(x)^{4p+2} \right)^{\frac{1}{4p+4}} \right] \\ &\leq \epsilon \|\tilde{u}_k\|_{L^{4p+2}(\mathbb{Z}^n)} + C_5 E_\Omega(u_k)^{\frac{1}{2}} \\ &\leq \epsilon \|\tilde{u}_k\|_{L^{2p+2}(\mathbb{Z}^n)} + \frac{1}{4} E_\Omega(u_k) + C_6, \end{aligned}$$

where C_5 and C_6 depend only on ϵ, n, N and p .

Hence, we obtain

$$\begin{aligned} J_\Omega(u_k) &\geq \frac{1}{2} E_\Omega(u_k) + \frac{\lambda}{2p+2} \sum_{x \in \Omega} \left(\frac{|u_k(x)|}{1 + |u_k(x)|} \right)^{2p+2} - \epsilon \|\tilde{u}_k\|_{L^{2p+2}(\mathbb{Z}^n)} - \frac{1}{4} E_\Omega(u_k) - C_6 \\ &= \frac{1}{4} E_\Omega(u_k) + \frac{\lambda}{2p+2} \sum_{x \in \Omega} \left(\frac{|u_k(x)|}{1 + |u_k(x)|} \right)^{2p+2} - \epsilon \|u_k\|_{L^{2p+2}(\Omega)} - C_6. \end{aligned} \quad (2.13)$$

By the inequality (2.12), we have the following estimate:

$$\begin{aligned}
\left(\sum_{x \in \Omega} u_k(x)^{2p+2} \right)^2 &= \left[\sum_{x \in \Omega} \left(\frac{|u_k(x)|}{1 + |u_k(x)|} \right)^{p+1} (1 + |u_k(x)|)^{p+1} |u_k(x)|^{p+1} \right]^2 \\
&\leq \sum_{x \in \Omega} \left(\frac{|u_k(x)|}{1 + |u_k(x)|} \right)^{2p+2} \sum_{x \in \Omega} (1 + |u_k(x)|)^{2p+2} u_k(x)^{2p+2} \\
&\leq 2^{2p+2} \sum_{x \in \Omega} \left(\frac{|u_k(x)|}{1 + |u_k(x)|} \right)^{2p+2} \sum_{x \in \Omega} (u_k(x)^{2p+2} + u_k(x)^{4p+4}) \\
&\leq 2^{2p+2} \sum_{x \in \Omega} \left(\frac{|u_k(x)|}{1 + |u_k(x)|} \right)^{2p+2} \sum_{x \in \Omega} u_k(x)^{2p+2} + 2^{2p+2} C_3 \sum_{x \in \Omega} \left(\frac{|u_k(x)|}{1 + |u_k(x)|} \right)^{2p+2} E_\Omega(u_k) \sum_{x \in \Omega} u_k(x)^{4p+4} \\
&\leq 2^{2p+2} \sum_{x \in \Omega} \left(\frac{|u_k(x)|}{1 + |u_k(x)|} \right)^{2p+2} \sum_{x \in \Omega} u_k(x)^{2p+2} + 2^{2p+2} C_3 \sum_{x \in \Omega} \left(\frac{|u_k(x)|}{1 + |u_k(x)|} \right)^{2p+2} E_\Omega(u_k) \left(\sum_{x \in \Omega} u_k(x)^{2p+2} \right)^{\frac{2p+1}{p+1}} \\
&\leq \frac{1}{4} \left(\sum_{x \in \Omega} u_k(x)^{2p+2} \right)^2 + C_7 \left[\left(\sum_{x \in \Omega} \left(\frac{|u_k(x)|}{1 + |u_k(x)|} \right)^{2p+2} \right)^2 + \sum_{x \in \Omega} \left(\frac{|u_k(x)|}{1 + |u_k(x)|} \right)^{2p+2} E_\Omega(u_k) \left(\sum_{x \in \Omega} u_k(x)^{2p+2} \right)^{\frac{2p+1}{p+1}} \right] \\
&\leq \frac{1}{2} \left(\sum_{x \in \Omega} u_k(x)^{2p+2} \right)^2 + C_8 \left[\left(\sum_{x \in \Omega} \left(\frac{|u_k(x)|}{1 + |u_k(x)|} \right)^{2p+2} \right)^2 + \left(\sum_{x \in \Omega} \left(\frac{|u_k(x)|}{1 + |u_k(x)|} \right)^{2p+2} \right)^{2p+2} E_\Omega(u_k)^{2p+2} \right] \\
&\leq \frac{1}{2} \left(\sum_{x \in \Omega} u_k(x)^{2p+2} \right)^2 + C_9 \left[1 + \left(\sum_{x \in \Omega} \left(\frac{|u_k(x)|}{1 + |u_k(x)|} \right)^{2p+2} \right)^{4p+4} + E_\Omega(u_k)^{4p+4} \right],
\end{aligned}$$

which yields that

$$\|u_k\|_{L^{2p+2}(\Omega)} \leq C_{10} \left[1 + \sum_{x \in \Omega} \left(\frac{|u_k(x)|}{1 + |u_k(x)|} \right)^{2p+2} + E_\Omega(u_k) \right], \quad (2.14)$$

where C_7 - C_{10} are constants depending only on n and p .

Choosing $\epsilon = \frac{\min\{\frac{1}{8}, \frac{4}{4p+4}\}}{C_{10}}$ and combining (2.13) and (2.14), we get

$$\|u_k\|_{L^{2p+2}(\Omega)} \leq C_2(J_\Omega(u_k) + 1).$$

Furthermore, by Lemma 2.3, we conclude

$$\|u_k\|_{L^{2p+2}(\Omega)} \leq C_2(J_\Omega(u_k) + 1) \leq C_1,$$

where C_2 and C_1 depend only on n , λ , p and N . \square

The boundedness of $\|u_k\|_{L^{2p+2}(\Omega)}$ ensures the existence of a solution to the Chern-Simons equation on Ω .

Lemma 2.6. *Let Ω be a finite subset of \mathbb{Z}^n containing the distinct points $\{p_j\}_{j=1}^M$. Consider the boundary value problem*

$$\begin{cases} \Delta u = \lambda e^u (e^u - 1)^{2p+1} + g & \text{on } \Omega, \\ u(x) = 0 & \text{on } \delta\Omega. \end{cases}$$

A solution $u_\Omega : \bar{\Omega} \rightarrow \mathbb{R}$ exists for this problem. This solution is maximal among all possible solutions and satisfies the condition $\|u\|_{l^{2p+2}(\Omega)} \leq C_1$, where C_1 depends only on n, λ, p and N .

Proof. By Lemmas 2.5 and 2.2, we have

$$u_k \rightarrow u_\Omega \text{ in } l^{2p+2}(\Omega),$$

and

$$\|u_\Omega\|_{l^{2p+2}(\Omega)} \leq C_0.$$

Since Δ is a local operator, and given the pointwise convergence, the function u_Ω in $l^{2p+2}(\Omega)$ is the solution to the equation

$$\begin{cases} \Delta u = \lambda e^u (e^u - 1)^{2p+1} + g \text{ on } \Omega, \\ u(x) = 0 \text{ on } \partial\Omega. \end{cases}$$

This completes the main proof.

The remaining task is to demonstrate that this solution is maximal. We claim that for any function $V \in C(\bar{\Omega})$ satisfying

$$\begin{cases} \Delta V \geq \lambda e^V (e^V - 1)^{2p+1} + g \text{ on } \Omega, \\ V(x) \leq 0 \text{ on } \partial\Omega, \end{cases}$$

it holds that

$$u_0 \geq u_1 \geq \dots \geq u_k \geq \dots \geq u_\Omega \geq V. \quad (2.15)$$

Initially, it is noted that

$$\Delta V \geq \lambda e^V (e^V - 1)^{2p+1} + g \geq \lambda e^V (e^V - 1)^{2p+1}.$$

We assert that $\sup_{x \in \Omega} V(x) \leq 0$. Should this not hold, and $V(x_0) = \sup_{x \in \Omega} V(x) > 0$ for some $x_0 \in \Omega$, then

$$0 \geq \Delta V(x_0) \geq \lambda e^{V(x_0)} (e^{V(x_0)} - 1)^{2p+1} > 0,$$

resulting in a contradiction and thus establishing the claim.

Assuming $V \leq u_k$, then

$$\begin{aligned} (\Delta - K)(u_{k+1} - V) &\leq \lambda e^{u_k} (e^{u_k} - 1)^{2p+1} - \lambda e^V (e^V - 1)^{2p+1} - K(u_k - V) \\ &\leq \lambda e^\omega (e^\omega - 1)^{2p} [(2p+2)e^\omega - 1] (u_k - V) - K(u_k - V) \\ &\leq K(e^{2\omega} - 1)(u_k - V) \leq 0, \end{aligned}$$

where the function ω satisfies $V \leq \omega \leq u_k \leq 0$. This ensures that $V \leq u_{k+1}$ by Lemma 2.1. Hence $V \leq u_\Omega$. If u'_Ω is another solution, we conclude that $u'_\Omega \leq u_\Omega$, which means that u_Ω is maximal. \square

We are now in a position to prove Theorem 1.1. Let Ω_i be finite and connected subsets satisfying

$$\Omega_0 \subset \Omega_1 \subset \dots \subset \Omega_k \subset \dots, \quad \bigcup_{i=1}^{\infty} \Omega_i = \mathbb{Z}^n.$$

We will use these lemmas to establish Theorem 1.1. For any integers $1 \leq j \leq k$, since $\Omega_j \subset \Omega_k$, and noting that $u_{\Omega_k} \leq 0$ on $\overline{\Omega_j}$, by (2.15), we conclude

$$u_{\Omega_k} \leq u_{\Omega_j} \text{ on } \overline{\Omega_j}.$$

Let \tilde{u}_{Ω_k} be the null extension of u_{Ω_k} to \mathbb{Z}^n . Then,

$$0 \geq \tilde{u}_{\Omega_1} \geq \tilde{u}_{\Omega_2} \geq \dots \geq \tilde{u}_{\Omega_k} \geq \dots$$

on \mathbb{Z}^n . Noting that $\|\tilde{u}_{\Omega_k}\|_{l^{2p+2}(\mathbb{Z}^n)} \leq C_1$ for any $k \geq 1$, we have the pointwise convergence

$$\tilde{u}_{\Omega_k}(x) \rightarrow u(x), \quad \forall x \in \mathbb{Z}^n,$$

and $u \in l^{2p+2}(\mathbb{Z}^n)$. Hence, u satisfies the equations

$$\begin{cases} \Delta u = \lambda e^u (e^u - 1)^{2p+1} + 4\pi \sum_{j=1}^M \delta_{p_j} \text{ on } \mathbb{Z}^n, \\ \lim_{d(x) \rightarrow +\infty} u(x) = 0, \end{cases}$$

which constitutes a topological solution.

Assume there exists another topological solution \hat{u} to (1.6). We now prove that $\hat{u}(x) \leq 0$ for all $x \in \mathbb{Z}^n$. If not, there exists $x_0 \in \mathbb{Z}^n$ such that $\hat{u}(x_0) > 0$. Noting that $\lim_{d(x) \rightarrow +\infty} \hat{u}(x) = 0$, we conclude there exists a domain Ω_i such that $\hat{u}(x_1) = \sup_{x \in \Omega_i} \hat{u}(x) > 0$ for some $x_1 \in \Omega_i$. Then, we obtain

$$0 \geq \Delta \hat{u}(x_1) \geq \lambda e^{\hat{u}(x_1)} (e^{\hat{u}(x_1)} - 1)^{2p+1} > 0,$$

which yields a contradiction.

Applying (2.15) on Ω_j , we have

$$\hat{u} \leq \tilde{u}_{\Omega_j}.$$

For a fixed integer $k \geq 1$, and for $j \geq k$, we have

$$\hat{u}(x) \leq \lim_{j \rightarrow \infty} \tilde{u}_{\Omega_j}(x) = u(x) \text{ on } \Omega_k.$$

Thus, we obtain $\hat{u} \leq u$ on \mathbb{Z}^n , and the solution u is maximal among all possible solutions.

ACKNOWLEDGMENTS

This work is partially supported by the National Key Research and Development Program of China 2020YFA0713100 and by the National Natural Science Foundation of China (Grant No. 11721101).

References

- [1] Martin T. Barlow. *Random walks and heat kernels on graphs*, volume 438 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 2017.
- [2] Luis A. Caffarelli and Yi Song Yang. Vortex condensation in the Chern-Simons Higgs model: an existence theorem. *Comm. Math. Phys.*, 168(2):321–336, 1995. 2
- [3] Dongho Chae and Namkwon Kim. Topological multivortex solutions of the self-dual Maxwell-Chern-Simons-Higgs system. *J. Differential Equations*, 134(1):154–182, 1997. 2

- [4] Hsungrow Chan, Chun-Chieh Fu, and Chang-Shou Lin. Non-topological multi-vortex solutions to the self-dual Chern-Simons-Higgs equation. *Comm. Math. Phys.*, 231(2):189–221, 2002. [2](#)
- [5] Xiaojun Chang, Ru Wang, and Duokui Yan. Ground states for logarithmic Schrödinger equations on locally finite graphs. *J. Geom. Anal.*, 33(7):Paper No. 211, 26, 2023. [2](#)
- [6] Ruixue Chao and Songbo Hou. Multiple solutions for a generalized Chern-Simons equation on graphs. *J. Math. Anal. Appl.*, 519(1):Paper No. 126787, 16, 2023. [2](#)
- [7] Ruixue Chao, Songbo Hou, and Jiamin Sun. Existence of solutions to a generalized self-dual Chern-Simons system on finite graphs. *arXiv preprint arXiv:2206.12863*, 2022. [2](#)
- [8] Gil Young Cho and Joel E. Moore. Topological BF field theory description of topological insulators. *Ann. Physics*, 326(6):1515–1535, 2011. [1](#)
- [9] Jia Gao and Songbo Hou. Existence theorems for a generalized Chern-Simons equation on finite graphs. *J. Math. Phys.*, 64(9):Paper No. 091502, 12, 2023. [2](#)
- [10] Huabin Ge. Kazdan-Warner equation on graph in the negative case. *J. Math. Anal. Appl.*, 453(2):1022–1027, 2017. [2](#)
- [11] Huabin Ge. A p -th Yamabe equation on graph. *Proc. Amer. Math. Soc.*, 146(5):2219–2224, 2018. [2](#)
- [12] Huabin Ge. The p th Kazdan-Warner equation on graphs. *Commun. Contemp. Math.*, 22(6):Paper No. 1950052, 17, 2020. [2](#)
- [13] Huabin Ge and Wenfeng Jiang. Yamabe equations on infinite graphs. *J. Math. Anal. Appl.*, 460(2):885–890, 2018. [2](#)
- [14] Alexander Grigor'yan. *Introduction to analysis on graphs*, volume 71 of *University Lecture Series*. American Mathematical Society, Providence, RI, 2018. [3](#)
- [15] Alexander Grigor'yan, Yong Lin, and Yunyan Yang. Kazdan-Warner equation on graph. *Calc. Var. Partial Differential Equations*, 55(4):Art. 92, 13, 2016. [2](#)
- [16] Alexander Grigor'yan, Yong Lin, and Yunyan Yang. Yamabe type equations on graphs. *J. Differential Equations*, 261(9):4924–4943, 2016.
- [17] Alexander Grigor'yan, Yong Lin, and Yunyan Yang. Existence of positive solutions to some nonlinear equations on locally finite graphs. *Sci. China Math.*, 60(7):1311–1324, 2017. [2](#)
- [18] Boling Guo and Fangfang Li. Existence of topological vortices in an Abelian Chern-Simons model. *J. Math. Phys.*, 56(10):101505, 10, 2015. [2](#)
- [19] Jongmin Han and Hee-Seok Nam. On the topological multivortex solutions of the self-dual Maxwell-Chern-Simons gauged $O(3)$ sigma model. *Lett. Math. Phys.*, 73(1):17–31, 2005. [2](#)
- [20] Xiaosen Han. The existence of multi-vortices for a generalized self-dual Chern-Simons model. *Nonlinearity*, 26(3):805–835, 2013. [2](#)
- [21] Jooyoo Hong, Yoonbai Kim, and Pong Youl Pac. Multivortex solutions of the abelian Chern-Simons-Higgs theory. *Phys. Rev. Lett.*, 64(19):2230–2233, 1990. [2](#)
- [22] Songbo Hou and Xiaoqing Kong. Existence and asymptotic behaviors of solutions to Chern-Simons systems and equations on finite graphs. *arXiv preprint arXiv:2211.04237*, 2022. [2](#), [5](#)
- [23] Songbo Hou and Wenjie Qiao. Solutions to a generalized Chern-Simons Higgs model on finite graphs by topological degree. *J. Math. Phys.*, 65(8):Paper No. 081503, 2024. [2](#)
- [24] Songbo Hou and Jiamin Sun. Existence of solutions to Chern-Simons-Higgs equations on graphs. *Calc. Var. Partial Differential Equations*, 61(4):Paper No. 139, 13, 2022. [2](#)
- [25] Yuanyang Hu. Existence of solutions to a generalized self-dual Chern-Simons equation on finite graphs. *J. Korean Math. Soc.*, 61(1):133–147, 2024. [2](#)
- [26] Bobo Hua, Genggeng Huang, and Jiaxuan Wang. The existence of topological solutions to the chern-simons model on lattice graphs. *arXiv preprint arXiv:2310.13905*, 2023. [2](#), [4](#)
- [27] An Huang, Yong Lin, and Shing-Tung Yau. Existence of solutions to mean field equations on graphs. *Comm. Math. Phys.*, 377(1):613–621, 2020. [2](#)
- [28] Genggeng Huang, Congming Li, and Ximing Yin. Existence of the maximizing pair for the discrete Hardy-Littlewood-Sobolev inequality. *Discrete Contin. Dyn. Syst.*, 35(3):935–942, 2015. [7](#)
- [29] Hsin-Yuan Huang, Youngae Lee, and Chang-Shou Lin. Uniqueness of topological multi-vortex solutions for a skew-symmetric Chern-Simons system. *J. Math. Phys.*, 56(4):041501, 12, 2015. [2](#)
- [30] Hsin-Yuan Huang, Jun Wang, and Wen Yang. Mean field equation and relativistic Abelian Chern-Simons model on finite graphs. *J. Funct. Anal.*, 281(10):Paper No. 109218, 36, 2021. [2](#)
- [31] R. Jackiw and Erick J. Weinberg. Self-dual Chern-Simons vortices. *Phys. Rev. Lett.*, 64(19):2234–2237, 1990. [2](#)
- [32] C. Nagraj Kumar and Avinash Khare. Charged vortex of finite energy in nonabelian gauge theories with Chern-Simons term. *Phys. Lett. B*, 178(4):395–399, 1986. [1](#)
- [33] Jiayu Li, Linlin Sun, and Yunyan Yang. Topological degree for Chern-Simons Higgs models on finite graphs. *Calc. Var. Partial Differential Equations*, 63(4):Paper No. 81, 21, 2024. [2](#)
- [34] Chang-Shou Lin and Jyotshana V. Prajapat. Vortex condensates for relativistic abelian Chern-Simons model with

- two Higgs scalar fields and two gauge fields on a torus. *Comm. Math. Phys.*, 288(1):311–347, 2009. [1](#)
- [35] Chang-Shou Lin and Yisong Yang. Non-Abelian multiple vortices in supersymmetric field theory. *Comm. Math. Phys.*, 304(2):433–457, 2011. [1](#)
- [36] Yong Lin and Yiting Wu. The existence and nonexistence of global solutions for a semilinear heat equation on graphs. *Calc. Var. Partial Differential Equations*, 56(4):Paper No. 102, 22, 2017. [2](#)
- [37] Yang Liu and Yunyan Yang. Topological degree for Kazdan-Warner equation in the negative case on finite graph. *Ann. Global Anal. Geom.*, 65(4):Paper No. 29, 20, 2024. [2](#)
- [38] Yang Liu and Mengjie Zhang. A heat flow with sign-changing prescribed function on finite graphs. *J. Math. Anal. Appl.*, 528(2):Paper No. 127529, 17, 2023. [2](#)
- [39] Yingshu Lü and Peirong Zhong. Existence of solutions to a generalized self-dual Chern-Simons equation on graphs. *arXiv preprint arXiv:2107.12535*, 2021. [2](#)
- [40] Shoudong Man. On a class of nonlinear Schrödinger equations on finite graphs. *Bull. Aust. Math. Soc.*, 101(3):477–487, 2020. [2](#)
- [41] Kwan Hui Nam. Vortex condensation in $U(1) \times U(1)$ Chern-Simons model with a general Higgs potential on a torus. *J. Math. Anal. Appl.*, 407(2):305–315, 2013. [1](#)
- [42] Alessio Porretta. A note on the Sobolev and Gagliardo-Nirenberg inequality when $p > N$. *Adv. Nonlinear Stud.*, 20(2):361–371, 2020. [7](#)
- [43] Zidong Qiu and Yang Liu. Existence of solutions to the nonlinear Schrödinger equation on locally finite graphs. *Arch. Math. (Basel)*, 120(4):403–416, 2023. [2](#)
- [44] S. Randjbar-Daemi, A. Salam, and J. Strathdee. Chern-Simons superconductivity at finite temperature. *Nuclear Phys. B*, 340(2-3):403–447, 1990. [1](#)
- [45] Gordon W. Semenoff and Pasquale Sodano. Nonabelian adiabatic phases and the fractional quantum Hall effect. *Phys. Rev. Lett.*, 57(10):1195–1198, 1986. [1](#)
- [46] Joel Spruck and Yi Song Yang. Topological solutions in the self-dual Chern-Simons theory: existence and approximation. *Ann. Inst. H. Poincaré C Anal. Non Linéaire*, 12(1):75–97, 1995. [4](#)
- [47] Linlin Sun and Liuquan Wang. Brouwer degree for Kazdan-Warner equations on a connected finite graph. *Adv. Math.*, 404:Paper No. 108422, 29, 2022. [2](#)
- [48] Gabriella Tarantello. Multiple condensate solutions for the Chern-Simons-Higgs theory. *J. Math. Phys.*, 37(8):3769–3796, 1996. [2](#)
- [49] Guofang Wang and Liqun Zhang. Non-topological solutions of the relativistic $SU(3)$ Chern-Simons Higgs model. *Comm. Math. Phys.*, 202(3):501–515, 1999. [2](#)
- [50] Ning Zhang and Liang Zhao. Convergence of ground state solutions for nonlinear Schrödinger equations on graphs. *Sci. China Math.*, 61(8):1481–1494, 2018. [2](#)
- [51] Xiaoxiao Zhang and Yanxun Chang. p -th Kazdan-Warner equation on graph in the negative case. *J. Math. Anal. Appl.*, 466(1):400–407, 2018. [2](#)
- [52] Xiaoxiao Zhang and Aijin Lin. Positive solutions of p -th Yamabe type equations on infinite graphs. *Proc. Amer. Math. Soc.*, 147(4):1421–1427, 2019. [2](#)