

# Matrix Theory EE5609 - Assignment 5

Sandhya Addetla  
PhD Artificial Intelligence Department  
AI20RESCH14001

**Abstract**—Find the asymptotes of the given hyperbola and also the equation to its conjugate hyperbola

Download all python codes from

<https://github.com/SANDHYA-A/Assignment5/blob/master/Assignment5.py>

## 1 PROBLEM STATEMENT

Find the asymptotes of the given hyperbola and also the equation to its conjugate hyperbola

$$19x^2 + 24xy + y^2 - 22x - 6y = 0 \quad (1.0.1)$$

## 2 SOLUTION

The general equation of second degree is given by

$$ax^2 + 2bxy + cy^2 + 2dx + 2ey + f = 0 \quad (2.0.1)$$

and can be expressed as

$$\mathbf{x}^T \mathbf{V} \mathbf{x} + 2\mathbf{u}^T \mathbf{x} + f = 0 \quad (2.0.2)$$

where

$$\mathbf{V} = \mathbf{V}^T = \begin{pmatrix} a & b \\ b & c \end{pmatrix} \quad (2.0.3)$$

$$\mathbf{u} = \begin{pmatrix} d \\ e \end{pmatrix} \quad (2.0.4)$$

Comparing equations 1.0.1 and 2.0.2 we get

$$\mathbf{V} = \mathbf{V}^T = \begin{pmatrix} 19 & 12 \\ 12 & 1 \end{pmatrix} \quad (2.0.5)$$

$$\mathbf{u} = \begin{pmatrix} -11 \\ -3 \end{pmatrix} \quad (2.0.6)$$

$$f = 0 \quad (2.0.7)$$

Expanding the Determinant of  $\mathbf{V}$ .

$$\Delta_V = \begin{vmatrix} 19 & 12 \\ 12 & 1 \end{vmatrix} < 0 \quad (2.0.8)$$

Hence from 2.0.8 given equation represents the hyperbola.

The characteristic equation of  $\mathbf{V}$  is obtained by evaluating the determinant

$$|V - \lambda \mathbf{I}| = 0 \quad (2.0.9)$$

$$\begin{vmatrix} 19 - \lambda & 12 \\ 12 & 1 - \lambda \end{vmatrix} = 0 \quad (2.0.10)$$

$$(19 - \lambda)(1 - \lambda) - 144 = 0 \quad (2.0.11)$$

$$\lambda_1 = -5, \lambda_2 = 25 \quad (2.0.12)$$

The eigenvector  $\mathbf{p}$  is defined as

$$\mathbf{V}\mathbf{p} = \lambda\mathbf{p} \quad (2.0.13)$$

$$\Rightarrow (\mathbf{V} - \lambda\mathbf{I})\mathbf{p} = 0 \quad (2.0.14)$$

For  $\lambda_1 = -5$ ,

$$(\mathbf{V} - \lambda_1\mathbf{I}) = \begin{pmatrix} 19 + 5 & 12 \\ 12 & 1 + 5 \end{pmatrix} \quad (2.0.15)$$

By row reduction ,

$$\begin{pmatrix} 24 & 12 \\ 12 & 6 \end{pmatrix} \quad (2.0.16)$$

$$\xleftrightarrow{R_2 \leftarrow 2R_2 - R_1} \begin{pmatrix} 24 & 12 \\ 0 & 0 \end{pmatrix} \quad (2.0.17)$$

$$\xleftrightarrow{R_1 \leftarrow \frac{R_1}{12}} \begin{pmatrix} 2 & 1 \\ 0 & 0 \end{pmatrix} \quad (2.0.18)$$

Substituting equation 2.0.18 in equation 2.0.14 we get

$$\begin{pmatrix} 2 & 1 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} u_1 \\ u_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \quad (2.0.19)$$

Where,  $\mathbf{p} = \begin{pmatrix} u_1 \\ u_2 \end{pmatrix}$  Let  $u_1 = t$

$$u_2 = -2t \quad (2.0.20)$$

Eigen vector  $\mathbf{p}_1$  is given by

$$\mathbf{p}_1 = \begin{pmatrix} t \\ -2t \end{pmatrix} \quad (2.0.21)$$

Let  $t = 1$ , we get

$$\mathbf{p}_1 = \begin{pmatrix} 1 \\ -2 \end{pmatrix} \quad (2.0.22)$$

For  $\lambda_2 = 25$ ,

$$(\mathbf{V} - \lambda_2 \mathbf{I}) = \begin{pmatrix} 19 - 25 & 12 \\ 12 & 1 - 25 \end{pmatrix} \quad (2.0.23)$$

By row reduction ,

$$\begin{pmatrix} -6 & 12 \\ 12 & -24 \end{pmatrix} \quad (2.0.24)$$

$$\xrightarrow{R_2 \leftarrow R_2 + 2R_1} \begin{pmatrix} -6 & 12 \\ 0 & 0 \end{pmatrix} \quad (2.0.25)$$

$$\xrightarrow{R_1 \leftarrow \frac{R_1}{6}} \begin{pmatrix} -1 & 2 \\ 0 & 0 \end{pmatrix} \quad (2.0.26)$$

Substituting equation 2.0.26 in equation 2.0.14 we get

$$\begin{pmatrix} -1 & 2 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \quad (2.0.27)$$

Where,  $\mathbf{p} = \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}$  Let  $v_1 = t$

$$v_2 = \frac{t}{2} \quad (2.0.28)$$

Eigen vector  $\mathbf{p}_2$  is given by

$$\mathbf{p}_2 = \begin{pmatrix} t \\ \frac{t}{2} \end{pmatrix} \quad (2.0.29)$$

Let  $t = 1$ , we get

$$\mathbf{p}_2 = \begin{pmatrix} 1 \\ \frac{1}{2} \end{pmatrix} \quad (2.0.30)$$

By eigen decomposition  $\mathbf{V}$  can be represented by

$$\mathbf{V} = \mathbf{P} \mathbf{D} \mathbf{P}^T \quad (2.0.31)$$

where

$$\mathbf{P} = (\mathbf{p}_1 \quad \mathbf{p}_2) \quad (2.0.32)$$

$$\mathbf{D} = \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix} \quad (2.0.33)$$

Substituting equations 2.0.22, 2.0.30 in equation 2.0.32 we get

$$\mathbf{P} = \begin{pmatrix} 1 & 1 \\ -2 & \frac{1}{2} \end{pmatrix} \quad (2.0.34)$$

Substituting equation 2.0.12 in 2.0.33 we get

$$\mathbf{D} = \begin{pmatrix} -5 & 0 \\ 0 & 25 \end{pmatrix} \quad (2.0.35)$$

Equation of a hyperbola and the combined equation of the Asymptotes differ only in the constant term.

$$19x^2 + 24xy + y^2 - 22x - 6y + K = 0 \quad (2.0.36)$$

The above equation can be expressed in the form

$$\mathbf{x}^T \mathbf{V} \mathbf{x} + 2\mathbf{u}^T \mathbf{x} + f = 0 \quad (2.0.37)$$

Comparing equation we get

$$\mathbf{V} = \mathbf{V}^T = \begin{pmatrix} 19 & 12 \\ 12 & 1 \end{pmatrix} \quad (2.0.38)$$

$$\mathbf{u} = \begin{pmatrix} -11 \\ -3 \end{pmatrix} \quad (2.0.39)$$

$$f = K \quad (2.0.40)$$

$$\Delta = \begin{vmatrix} 19 & 12 & -11 \\ 12 & 1 & -3 \\ -11 & -3 & K \end{vmatrix} \quad (2.0.41)$$

Since the equations represent pair of straight lines, equating the determinant to zero, we can get the value of K

$$\implies K = 4 \quad (2.0.42)$$

Let  $(\alpha, \beta)$  be their point of intersection, then

$$\begin{pmatrix} a & b \\ b & c \end{pmatrix} \begin{pmatrix} \alpha \\ \beta \end{pmatrix} = \begin{pmatrix} -d \\ -e \end{pmatrix} \quad (2.0.43)$$

Substituting the values, we obtain,

$$\begin{pmatrix} 19 & 12 \\ 12 & 1 \end{pmatrix} \begin{pmatrix} \alpha \\ \beta \end{pmatrix} = \begin{pmatrix} 11 \\ 3 \end{pmatrix} \quad (2.0.44)$$

$$\text{We get, } \alpha = \frac{1}{5}, \beta = \frac{3}{5} \quad (2.0.45)$$

Using Affine transformation and Spectral decomposition, we get

$$X' = \pm \sqrt{-\frac{\lambda_2}{\lambda_1}} Y' \quad (2.0.46)$$

$$\text{where } X' = Xu_1 + Yu_2 \quad (2.0.47)$$

$$Y' = Xv_1 + Yv_2 \quad (2.0.48)$$

$$X = x - \alpha \text{ and } Y = y - \beta \quad (2.0.49)$$

Therefore,

$$u_1(x - \alpha) + u_2(y - \beta) = \pm \sqrt{-\frac{\lambda_2}{\lambda_1}}(v_1(x - \alpha) + v_2(y - \beta)) \quad (2.0.50)$$

Substituting values, we get

$$(x - \frac{1}{5}) - 2(y - \frac{3}{5}) = \pm \sqrt{\frac{25}{5}}(x - \frac{1}{5}) + \frac{1}{2}(y - \frac{3}{5}) \quad (2.0.51)$$

Simplifying above equation

$$8x + 9y - 7 = 0 \quad (2.0.52)$$

$$12x + y + 7 = 0 \quad (2.0.53)$$

$$\implies (8x + 9y - 7)(12x + y + 7) = 0 \quad (2.0.54)$$

Thus the equation of lines are

$$(8 \ 9) \mathbf{x} = 7 \quad (2.0.55)$$

$$(12 \ 1) \mathbf{x} = -7 \quad (2.0.56)$$

The Equation of Conjugate hyperbola is given by:

2(Equation of Asymptotes)- Equation of hyperbola.

From Eq 1.0.1 and 2.0.36, we obtain equation of Conjugate hyperbola as:-

$$19x^2 + 24xy + y^2 - 22x - 6y + 8 = 0 \quad (2.0.57)$$

The general equation of second degree is given by

$$ax^2 + 2bxy + cy^2 + 2dx + 2ey + f = 0 \quad (2.0.58)$$

comparing equation 2.0.57 with the general equation of second degree given at 2.0.58, it can be expressed as

$$\mathbf{x}^T \mathbf{V} \mathbf{x} + 2\mathbf{u}^T \mathbf{x} + f = 0 \quad (2.0.59)$$

where

$$\mathbf{V} = \mathbf{V}^T = \begin{pmatrix} a & b \\ b & c \end{pmatrix} \quad (2.0.60)$$

$$\mathbf{u} = \begin{pmatrix} d \\ e \end{pmatrix} \quad (2.0.61)$$

Comparing equations 2.0.57 and 2.0.59 we get

$$\mathbf{V} = \mathbf{V}^T = \begin{pmatrix} 19 & 12 \\ 12 & 1 \end{pmatrix} \quad (2.0.62)$$

$$\mathbf{u} = \begin{pmatrix} -11 \\ -3 \end{pmatrix} \quad (2.0.63)$$

$$f = 8 \quad (2.0.64)$$

Therefore, the equation of the conjugate hyperbola is as given below:-

$$\mathbf{x}^T \begin{pmatrix} 19 & 12 \\ 12 & 1 \end{pmatrix} \mathbf{x} + 2 \begin{pmatrix} -11 & -3 \end{pmatrix} \mathbf{x} + 8 = 0 \quad (2.0.65)$$




Fig. 1: Hyperbola, Conjugate Hyperbola and Asymptotes