

Government of India
India Meteorological Department
State Agro-Meteorological Centre
Visakhapatnam-530017




भारत सरकार
भारत मौसम विज्ञान विभाग
राज्यकृषि- मौसम विज्ञान केंद्र
विशाखापत्तनम-530017

Agromet Advisory Service Bulletin for the State of ANDHRA PRADESH
(Advisories mentioned in the bulletin are prepared by different AMFUs in ANDHRA PRADESH)

Bulletin No.: 55/2026

Issued on: 10.07.2026



From the period (11.07.2026 to 15.07.2026)
(5 days from Saturday to Wednesday)
State Agro-meteorological Services Center (SAMC), Amaravati

Agro-climatic Zone	Districts	In-charge stations
North coastal zone	Srikakulam, Vizianagaram, Visakhapatnam & Anakapalli	RARS, Anakapalle
Southern zone	SPSR Nellore, Tirupati, Chittoor, YSR-Kadapa & Annamayya	RARS, Tirupati
Godavari zone	East Godavari, Kakinada & Dr. B.R. Ambedkar Konaseema	RARS, Chintapalle
	West Godavari & Eluru	RARS, Lam, Guntur
Krishna zone	Guntur, Krishna, Bapatla, NTR, Palnadu & Prakasam	RARS, Lam, Guntur
Scarce rainfall zone	Ananthapuramu, Sri Sathyasai, Kurnool & Nandyal	RARS, Ananthapuramu
High Altitude zone	Parvathipuram Manyam & Alluri Sitharama Raju	RARS, Chintapalle

Weather Warnings (Met-Subdivision Level): Coastal Andhra Pradesh & Rayalaseema:

Day 1 (10.07.2026): Thunderstorm accompanied with lightening likely at isolated places over North Coastal Andhra Pradesh and Yanam and Rayalaseema. Strong surface winds speed reaching 40-50 Kmph likely at isolated places over Coastal Andhra Pradesh, Yanam and Rayalaseema.

Day 2 (11.07.2026): Strong surface winds speed reaching 40-50 Kmph likely at isolated places over Coastal Andhra Pradesh, Yanam and Rayalaseema.

Day 3 (12.07.2026): Strong surface winds speed reaching 30-40 Kmph likely at isolated places over Coastal Andhra Pradesh, Yanam and Rayalaseema.

Day 4 (13.07.2026): Strong surface winds speed reaching 30-40 Kmph likely at isolated places over Coastal Andhra Pradesh, Yanam and Rayalaseema.

Day 5 (14.07.2026): Strong surface winds speed reaching 30-40 Kmph likely at isolated places over Coastal Andhra Pradesh, Yanam and Rayalaseema.

NCAP: North Coastal Andhra Pradesh	SCAP: South Coastal Andhra Pradesh
---	---

Likely impacts of weather warnings (General)

- Partial reduction in visibility, slippery roads.
- Major disruption in traffic flow.
- Moderate water logging in parts of low-lying areas and adjacent to river banks.
- Possibility of falling small tree branches.
- Loosening of rocks due to persistent rains leading to possibility of minor land / mud slides in elevated hilly areas (Ghat areas).

Impact based advisories (General)

- Avoid travel during times of heavy rain.
- Do not take shelter under tree during heavy rain.
- Follow guidelines issued by the government.

Likely impacts of weather warnings on Agriculture due to heavy rain

- Heavy rainfall can cause soil erosion and wash away valuable topsoil, harming agriculture.
- Excessive rainfall can cause delays in planting and harvesting.
- heavy rains may restrict the growth of roots of plants.
- Excess rain can lead to oxygen deficiency and nutrient loss, resulting in reduced crop yields.
- Pests, diseases and fungus may develop on the crops due to high humidity.
- Weed growth increases.
- Long exposure to strong winds causes morphological changes.
- Exposure of growing plants to hot wind results in dwarfing due to desiccation of plant tissue and reduces growth.
- Strong wind may increase crop water requirements due to increased evapotranspiration
- Heavy wind during flowering reduces pollination, causes flower shed, increases sterility and reduces fruit set in all crops.
- Livestock may be infected by mites, ticks, flies & other seasonal diseases.
- The monsoon season is congenial for vector transmission of viral diseases.

Agromet advisories

- Maintain 5 to 10 cm water level in rice fields,
- Drain out excess water from rice, pigeon pea, black gram, green gram, maize, sunflower, groundnut, soybean, vegetables and orchards,
- Advisories issued by AMFUs may be followed in respect of various crops depending on the stage of the crop while applying fertilizers, pesticides, herbicides and other management practices.
- Livestock may be vaccinated in time.
- Keep ready insect repellents, insecticides
- provide staking in vegetables and newly planted orchards,
- keep farm animals indoor during thunderstorm & lightning periods.

Summary of four days' rainfall over the State (06th July 2026 – 09th July 2026):

Met Sub Division	06-07-2026	07-07-2026	08-07-2026	09-07-2026
Coastal Andhra Pradesh	ISOL	ISOL	ISOL	ISOL
Rayalaseema	ISOL	ISOL	ISOL	ISOL
ISOL: Isolated i.e. Rain at 1 or 2 places SCT: Scattered i.e. Rain at a few places DRY: No rain		FWS: Fairly Widespread i.e. Rain at many places WS: Widespread i.e. Rain at most places NSC: No Significant Change		

Rainfall forecast for next 5 days:

Met Sub Division	10.07.2026	11.07.2026	12.07.2026	13.07.2026	14.07.2026
COASTAL ANDHRA PRADESH	ISOL	ISOL	ISOL	ISOL	ISOL
RAYALASEEMA	ISOL	ISOL	DRY	DRY	DRY
MET SUBDIVISION OF COASTAL ANDHRA PRADESH	10.07.2026	11.07.2026	12.07.2026	13.07.2026	14.07.2026
NORTH COASTAL ANDHRA PRADESH & YANAM	SCT	ISOL	ISOL	ISOL	ISOL
SOUTH COASTAL ANDHRA PRADESH	ISOL	ISOL	DRY	DRY	DRY
ISOL: Isolated i.e. Rain at 1 or 2 places SCT: Scattered i.e. Rain at a few places DRY: No rain		FWS: Fairly Widespread i.e. Rain at many places WS: Widespread i.e. Rain at most places NSC: No Significant Change			

Agromet Advisories issued by different Agromet Field Units (AMFUs) of Andhra Pradesh

Meteorological Subdivision of Coastal Andhra Pradesh:

AMFU ANAKAPALLI

AGRO METEOROLOGICAL ADVISORY FOR SRIKAKULAM DISTRICT

S. No.	Crop	Stage and condition of the crop	Problems / Pest and disease	Agro Advisory
1	Paddy	Nursery management	-	• Select short and medium duration varieties • If there is sufficient soil moisture,

				prepare the land for sowing paddy nursery, prepare raised beds and apply Well-decomposed FYM @ 200 kg/5cents nursery apply 2 kg urea, 6 kg single super phosphate, 1 kg of murate of potash and sow the seed @ 20kg/acre.
2	Dry Direct sown paddy	Sowing		- Farmers can go for sowing of dry direct sown paddy if there is sufficient moisture with suitable varieties @ 30 kg/acre duly following the seed treatment with carbendazim @ 3g/kg seed. - Go for pre-emergence application of pendimethalin @ 700 ml/acre within 24-48 hours after sowing.
3	Mesta		-	Go sowing of mesta with varieties like Bhimili-1, AMV-9, AMV-10, AMV-7, AMV-5, AMV-11 etc., duly following the seed treatment with 5g of imidacloprid and 3g of mancozeb for one kg seed.
4	Maize	Sowing	-	If sufficient moisture is available, farmers can go for sowing of maize using the varieties like DHM-115, DHM-11, DHM-17, DHM-19 and DHM 121 with a seed rate of 8 kg/acre at a spacing of 60x20 cm.
		Seedling stage	Fall armyworm	Go for spraying with 1500 PPM neem oil @ 5 ml/L.
5	Groundnut	Sowing	-	Go for land preparation and sowing of <i>kharif</i> groundnut with high yielding groundnut varieties like visishta, Kadiri 9 with a seed rate of 56 kg/acre. Go for seed treatment with imidacloprid 600 FS @ 2ml along with 4ml water for kg seed and after drying treat the seed with tebuconazole 2DS @ 2g/kg seed and then after two days before sowing of the crop go for seed treatment with rhizobium culture @ 10g or 10 ml/kg seed.
6	Cotton			- If sufficient moisture is available Select cotton varieties like L.A.H.H-5, L.A.H.H-7, N.D.L.H.H-240 and go for sowing at a seed rate of 1 kg/acre adopting a spacing of 90x45 cm. - To avoid the incidence of sucking pests at early crop growth stages, go for seed treatment with imidachloprid 70% WS @ 5g/kg.

AGRO METEOROLOGICAL ADVISORY FOR VIZIANAGARAM DISTRICT

S. No.	Crop	Stage and condition of the crop	Problems / Pest and disease	Agro Advisory
1	Paddy	Nursery management	-	- Select short and medium duration varieties - If there is sufficient soil moisture, prepare the land for sowing paddy nursery, prepare raised beds and apply Well-decomposed FYM @ 200 kg/ 5cents nursery apply 2 kg urea, 6 kg single super phosphate, 1 kg of murate of potash and sow the seed @ 20kg/acre.
		Seedling	-	- Apply urea @ 2 kg as top dressing for a nursery bed of 5 cents at 12-14 days after sowing. - Apply carbofuran granules @ 160g/cent at 15-20 days after sowing nursery.
		Transplanting		- Go for clipping of leaf tips of paddy seedlings and transplant at a spacing of 20x15 cm @ 2-3 seedlings per hill and see that there are 33 hills per square metre. - Make alley ways of 20cm for every 2 metres after transplanting
2	Dry Direct sown paddy	Sowing		- Farmers can go for sowing of dry direct sown paddy if there is sufficient moisture with suitable varieties @ 30 kg/acre duly following the seed treatment with carbendazim @ 3g/kg seed. - Go for pre-emergence application of pendimethalin @ 700 ml/acre within 24-48 hours after sowing.
		Seedling stage		If the crop is at 20 days age, go for spraying of bispyribac sodium @ 100 ml/acre to control grasses and broad leaf weeds. If grasses are dominating, go for spraying with fenoxoprop P ethyl @ 350 ml/acre.
3	Maize	Sowing	-	Based on available soil moisture, Maize growing farmers can go for sowing of <i>kharif</i> maize using the varieties like DHM-115, DHM-11, DHM-17, DHM-19 and DHM 121 with a seed rate of 8 kg/acre at a spacing of 60x20 cm.
		Seedling stage	Fall armyworm	If insect scrappings are noticed, spray 1500 PPM neem oil @ 5 ml/L or metarhizium/MI 35 @ 5 g/L for twice at week interval.
4	Blackgram	Sowing		Select blackgram varieties like LBG 932, LBG 904,TBG 129, LBG 884, TRG 59,TBG 104 and go for sowing duly following the seed treatment with carbendazim 2.5 g and imidacloprid 600 FS

				@ 5 ml per kg seed to protect the crop from sucking pests and diseases.
		Seedling stage	Flea beetles	Spray chlorpyrifos 2.5 ml per litre.
5	Sugarcane	Formative stage		Go for earthing up for the crop of 120 days and also propping to avoid crop lodging due to winds.
6	<i>Rainfed</i> sugarcane	Planting	-	Planting of sugarcane may be done before 15th July duly following sett treatment with 10% lime water and using a seed rate of 20,000 three budded setts/acre at a spacing of 60 cm between rows.
7	Cotton			- If sufficient moisture is available Select cotton varieties like L.A.H.H-5, L.A.H.H-7, N.D.L.H.H-240 and go for sowing at a seed rate of 1 kg/acre adopting a spacing of 90x45 cm. - To avoid the incidence of sucking pests at early crop growth stages, go for seed treatment with imidachloprid 70% WS @ 5g/kg.
8	Mesta		-	Go sowing of mesta with varieties like Bhimili-1, AMV-9, AMV-10, AMV-7, AMV-5, AMV-11 etc., duly following the seed treatment with 5g of imidacloprid and 3g of mancozeb for one kg seed.
9	Cashew	-	-	Farmers are suggested to take up pruning in cashew orchards to ensure good air circulation and sunlight penetration.

AGRO METEOROLOGICAL ADVISORY FOR VISAKHAPATNAM DISTRICT

S. No.	Crop	Stage and condition of the crop	Problems / Pest and disease	Agro Advisory
1	Paddy	Nursery management	-	- Select short and medium duration varieties - If there is sufficient soil moisture, prepare the land for sowing paddy nursery, prepare raised beds and apply Well-decomposed FYM @ 200 kg/5cents nursery apply 2 kg urea, 6 kg single super phosphate, 1 kg of murate of potash and sow the seed @ 20kg/acre.
2	Bajra	Sowing	-	Select varieties like ABV 04 or APHB 126 and go for sowing with a seed rate of 1.6 kg/acre duly following the seed treatment
3	Sesamum	Vegetative Stage	Sucking pest	Sucking pest incidence may occur during prevailing weather conditions which acts as vector for phyllody

				disease. Spray dimethoate @ 2 ml/L for vector control.
4	Cashew	-	-	Farmers are suggested to take up pruning in cashew orchards to ensure good air circulation and sunlight penetration.
5	Chilli/Brinjal	Nursery	-	Nurseries can be raised in pro-trays to avoid damping off in nurseries
		Seedling stage	Damping off	Drench the soil with COC @ 3 g/L.S

AGRO METEOROLOGICAL ADVISORY FOR ANAKAPALLI DISTRICT

S. No.	Crop	Stage and condition of the crop	Problems / Pest and disease	Agro Advisory
1	Paddy	Nursery management	-	- Select short and medium duration varieties - If there is sufficient soil moisture, prepare the land for sowing paddy nursery, prepare raised beds and apply Well-decomposed FYM @ 200 kg/ 5cents nursery apply 2 kg urea, 6 kg single super phosphate, 2 kg of murate of potash and sow the seed @ 20kg/acre.
		Seedling	-	- Apply urea @ 2 kg as top dressing for a nursery bed of 5 cents at 12-14 days after sowing. - Apply carbofuran granules @ 160g/cent at 15-20 days after sowing nursery.
2	Maize	Sowing	-	- Based on available soil moisture, go for sowing of <i>kharif</i> maize using the varieties like DHM-115, DHM-11, DHM-17, DHM-19 and DHM 121 with a seed rate of 8 kg/acre at a spacing of 60x20 cm. - Install pheromone traps @ 10/acre before sowing to forecast the pest.
		Seedling stage	Fall armyworm	If insect scrappings are noticed, spray 1500 PPM neem oil @ 5 ml/L or metarhizium/MI 35 @ 5 g/L for twice at week interval.
3	Blackgram	Sowing		Select blackgram varieties like LBG 932, LBG 904,TBG 129, LBG 884, TRG 59,TBG 104 and go for sowing duly following the seed treatment with carbendazim 2.5 g and imidacloprid 600 FS @ 5 ml per kg seed to protect the crop from sucking pests and diseases.
		Seedling stage	Flea beetles	Spray chlorpyrifos 2.5 ml per litre.
4	Greengram	Sowing		Select greengram varieties like LGG 630, LGG 460 and IPM 2-14 and go for sowing duly following the seed treatment.
5	Groundnut	Sowing	-	Go for land preparation and sowing of groundnut with high yielding groundnut varieties like visishta, Kadiri 9 with a seed rate

				of 56 kg/acre. Go for seed treatment with imidacloprid 600 FS @ 2ml along with 4ml water for kg seed and after drying treat the seed with tebuconazole 2DS @ 2g/kg seed and then after two days before sowing of the crop go for seed treatment with rhizobium culture @ 10g or 10 ml/kg seed.
6	Sesamum	Vegetative Stage	Sucking pest	Sucking pest incidence may occur during prevailing weather conditions which acts as vector for phyllody disease. Spray dimethoate @ 2 ml/L for vector control.
7	Sugarcane	Formative stage	-	Go for earthing up for the crop of 120 days and also propping to avoid crop lodging due to winds.
8	<i>Rainfed</i> sugarcane	Planting	-	Planting of sugarcane may be done before 15th July duly following sett treatment with 10% lime water and using a seed rate of 20,000 three budded setts/acre at a spacing of 60 cm between rows.
9	Cashew	-	-	Farmers are suggested to take up pruning in cashew orchards to ensure good air circulation and sunlight penetration.

AMFU CHINTAPALLI

AGRO METEOROLOGICAL ADVISORY FOR PARVATHIPURAM MANYAM DISTRICT

Crop	Stage of the crop/ Management	Recommendation
Paddy	Nursery Stage	- Utilize the available soil moisture for nursery bed preparation and sowing of recommended varieties. Ensure proper drainage to prevent water stagnation in nursery beds. - Treat seed with Carbendazim @ 2 g/kg seed or Trichoderma viride @ 10 g/kg seed before sowing to protect against seed-borne diseases. - Apply DAP @ 20 kg/acre of nursery area and monitor for blast disease under humid conditions. If symptoms appear, spray Tricyclazole 75 WP @ 0.6 g/L of water.
Ragi	Nursery	- The expected rainfall provides favorable moisture for sowing; farmers are advised to begin sowing of ragi varieties such as Local Peddachodi and Indravathi using a seed rate of 4–5 kg/acre. - Treat seed with Trichoderma viride @ 10 g/kg seed or Pseudomonas fluorescens @ 10 g/kg seed before sowing to minimize seedling diseases and promote vigorous crop establishment.
Mango	Fruit development to harvest stage	- Maintain orchard sanitation by removing and destroying damaged and fallen fruits to reduce fruit fly and fruit borer incidence. If infestation is noticed, spray neem oil @ 2–3 ml or Spinosad 45 SC @ 0.3 ml per litre of water during clear weather conditions. - Due to warm and humid weather, monitor the orchard for mango hoppers and other sucking pests. If incidence is observed, spray Thiamethoxam @ 0.3 g or Imidacloprid @ 0.3 ml per litre of water during dry weather. - Harvest mature fruits during clear weather and avoid harvesting during rainfall to reduce fruit rotting and improve shelf life. Keep harvested fruits in a shaded and well-ventilated place.
Maize	Sowing Stage	- The forecasted rainfall provides favorable soil moisture for sowing; farmers are advised to take up sowing of suitable maize hybrids using a seed rate of 8–10 kg/acre and incorporate well-decomposed FYM or compost @ 2–4 t/acre during land preparation to improve soil health and moisture retention. - Treat seed with Trichoderma viride @ 10 g/kg seed or Metalaxyl + Mancozeb @ 2 g/kg seed before sowing to protect against seed- and soil-borne diseases and ensure

		uniform germination and healthy crop establishment.
Turmeric	Early Crop Establishment Stage	- Maintain proper field drainage to prevent water stagnation and rhizome rot. Carry out gap filling wherever germination is poor to maintain optimum plant population. - Apply well-decomposed FYM/compost @ 2–3 t/acre as mulch or top dressing and keep the field weed-free to promote uniform sprouting and early growth. - Monitor for rhizome rot incidence under humid conditions. If symptoms are observed, drench the soil around affected plants with Metalaxyl + Mancozeb @ 2 g/L of water and remove severely affected plants to prevent further spread during rain free period.
Farm Preparatory Advisories		- Incorporation of farmyard manure, compost and crop residues during field preparation is advised to improve soil organic carbon and moisture retention capacity. - Undertake nursery bed preparation for Kharif vegetables and other seasonal crops with adequate drainage provisions to prevent seedling mortality under intermittent rainfall conditions. - Farmers are advised to undertake soil sample collection and testing before kharif crop establishment to assess soil fertility status and enable site-specific nutrient management based on soil test recommendations. - Sow Dhaincha (Sesbania aculeata) @ 25–30 kg seed/ha or Sunhemp (Crotalaria juncea) @ 40–50 kg seed/ha as green manure crops and incorporate them into the soil at 45–50 days after sowing to improve soil organic carbon, soil structure, and nutrient availability for the succeeding crop.
Livestock		- Prevailing humid and cloudy weather conditions may favor the incidence of Hemorrhagic Septicemia (HS) and Black Quarter (BQ) in livestock; farmers are advised to vaccinate animals in consultation with veterinary officials. - Maintain proper sanitation, dry floor conditions and avoid water stagnation near cattle sheds to minimize the spread of bacterial and parasitic diseases during intermittent rainfall conditions.

AGRO METEOROLOGICAL ADVISORY FOR ALLURI SITARAMA RAJU DISTRICT

Crop	Stage of the crop/ Management	Recommendation
Rice	nursery	- The forecasted light rainfall and favorable temperature conditions are suitable for initiating nursery sowing; farmers are advised to start nursery raising with recommended rice varieties such as MTU 1121, RNR 15049, RGL 2537, suitable local land races, and private hybrids using a seed rate of 30–40 kg/acre of main field area. - Treat seed with Carbendazim @ 2 g/kg seed or Trichoderma viride @ 10 g/kg seed before sowing to protect against seed- and soil-borne diseases and ensure healthy seedling establishment. - Apply DAP @ 20 kg/acre of nursery area and monitor for blast disease under humid conditions. If symptoms appear, spray Tricyclazole 75 WP @ 0.6 g/L of water.
Ragi	nursery	- The expected rainfall provides favorable moisture for sowing; farmers are advised to begin sowing of ragi varieties such as Local Peddachodi and Indravathi using a seed rate of 4–5 kg/acre. - Treat seed with Trichoderma viride @ 10 g/kg seed or Pseudomonas fluorescens @ 10 g/kg seed before sowing to minimize seedling diseases and promote vigorous crop establishment.
Turmeric	Early Crop Establishment Stage	- Maintain proper field drainage to prevent water stagnation and rhizome rot. Carry out gap filling wherever germination is poor to maintain optimum plant population. - Apply well-decomposed FYM/compost @ 2–3 t/acre as mulch or top dressing and keep the field weed-free to promote uniform sprouting and early growth. - Monitor for rhizome rot incidence under humid conditions. If symptoms are observed, drench the soil around affected plants with Metalaxyl + Mancozeb @ 2 g/L of water and remove severely affected plants to prevent further spread during rain free period.
Coffee	init development stage	- Due to humid and rainy weather conditions, monitor plantations for black rot incidence and spray 10 g Bordeaux mixture per litre of water during dry weather for effective control. - Avoid spraying before rainfall; ensure at least 5–6 rain-free hours after application for

		better effectiveness. Do not carry out spraying during rainfall; perform spraying only when the weather is calm and dry.
Black pepper	fruit development stage	- For quick wilt / foot rot prevention, drench the vine base with Copper oxychloride @ 3 g/L and maintain proper drainage to avoid water stagnation around the root zone, especially during rainfall periods. - Do not carry out spraying during rainfall; perform spraying only when the weather is calm and dry.
Mango	Fruit development to harvest stage	- Maintain orchard sanitation by removing and destroying damaged and fallen fruits to reduce fruit fly and fruit borer incidence. If infestation is noticed, spray neem oil @ 2–3 ml or Spinosad 45 SC @ 0.3 ml per litre of water during clear weather conditions. - Due to warm and humid weather, monitor the orchard for mango hoppers and other sucking pests. If incidence is observed, spray Thiamethoxam @ 0.3 g or Imidacloprid @ 0.3 ml per litre of water during dry weather. - Harvest fully matured fruits before heavy rains to avoid fruit drop and spongy tissue. Protect harvested fruit using graded packaging and shade drying. Prune diseased/dry twigs post-harvest and apply Bordeaux paste.
Maize	Sowing Stage	- The forecasted rainfall provides favorable soil moisture for sowing; farmers are advised to take up sowing of suitable maize hybrids using a seed rate of 8–10 kg/acre and incorporate well-decomposed FYM or compost @ 2–4 t/acre during land preparation to improve soil health and moisture retention. - Treat seed with Trichoderma viride @ 10 g/kg seed or Metalaxyl + Mancozeb @ 2 g/kg seed before sowing to protect against seed- and soil-borne diseases and ensure uniform germination and healthy crop establishment.
Livestock		- Prevailing humid weather conditions may increase the risk of Hemorrhagic Septicemia (HS), Black Quarter (BQ), and parasitic infestations in livestock. Farmers are advised to vaccinate susceptible animals against HS and BQ in consultation with veterinary officials. - Maintain proper shed hygiene, drainage, and dry bedding to reduce pathogen buildup. Ensure adequate ventilation and provide clean drinking water to minimize disease incidence and environmental stress under humid conditions

AGRO METEOROLOGICAL ADVISORY FOR KAKINADA DISTRICT

Crop	Stage of the crop/ Management	Recommendation
Paddy	Nursery Stage	- Utilize the available soil moisture for nursery bed preparation and sowing of recommended varieties. Ensure proper drainage to prevent water stagnation in nursery beds. - Treat seed with Carbendazim @ 2 g/kg seed or Trichoderma viride @ 10 g/kg seed before sowing to protect against seed-borne diseases. - Apply DAP @ 20 kg/acre of nursery area and monitor for blast disease under humid conditions. If symptoms appear, spray Tricyclazole 75 WP @ 0.6 g/L of water.
Coconut	Nut development	- Apply FYM/compost @ 25–50 kg/palm and neem cake @ 5 kg/palm under adequate soil moisture conditions. - Monitor for Rhinoceros beetle and Red palm weevil; install Rhynchophorol traps @ 1 trap/ha and apply neem cake @ 1–2 kg/palm in basins to reduce infestation.
Mango	Fruit development to harvest stage	- Maintain proper orchard drainage and remove fallen fruits to minimize fruit rotting and disease buildup under intermittent rainfall conditions. - Harvest mature fruits during rain-free periods and keep harvested produce in a shaded, well-ventilated place to improve shelf life and reduce post-harvest losses. - Monitor for fruit fly, anthracnose, and mango hopper incidence. If infestation is observed, spray Spinosad 45 SC @ 0.3 ml/L for fruit fly or Carbendazim @ 1 g/L for anthracnose during clear weather conditions.
Banana	bunch development stage	- Maintain proper drainage to prevent waterlogging and root-zone hypoxia. Provide propping to bearing plants to prevent lodging and pseudostem breakage under strong winds. - Apply neem cake @ 1–2 kg/plant and recommended fertilizers in split doses under adequate soil moisture conditions to support vegetative growth and bunch development - Monitor for Sigatoka leaf spot, pseudostem weevil, aphids, and thrips under humid conditions. For Sigatoka management, spray Propiconazole @ 1 ml/L or Hexaconazole @ 2 ml/L of water. - For severe aphid or thrips infestation, spray Imidacloprid 17.8 SL @ 0.3 ml/L or Thiamethoxam 25 WG @ 0.25 g/L of water during rain-free periods.
Sugarcane	Management	- Maintain proper field drainage to prevent waterlogging and ensure adequate soil aeration. Carry out earthing-up to improve root anchorage and minimize lodging. - Apply urea @ 55 kg/acre (or as per recommended split dose) under adequate soil moisture conditions and incorporate it into the soil to improve nitrogen-use efficiency. - Monitor for Pyrilla incidence, which may increase under warm and humid conditions. If infestation is observed, spray Imidacloprid 17.8 SL @ 0.3 ml/L or Thiamethoxam 25 WG @ 0.25 g/L of water during clear weather conditions.
Maize	Sowing Stage	- The forecasted rainfall provides favorable soil moisture for sowing; farmers are advised to take up sowing of suitable maize hybrids using a seed rate of 8–10 kg/acre and incorporate well-decomposed FYM or compost @ 2–4 t/acre during land preparation to improve soil health and moisture retention. - Treat seed with Trichoderma viride @ 10 g/kg seed or Metalaxyl + Mancozeb @ 2 g/kg seed before sowing to protect against seed- and soil-borne diseases and ensure uniform germination and healthy crop establishment.
Farm Preparatory Advisories		- Incorporation of farmyard manure, compost and crop residues during field preparation is advised to improve soil organic carbon and moisture retention capacity. - Undertake nursery bed preparation for Kharif vegetables and other seasonal crops with adequate drainage provisions to prevent seedling mortality under intermittent rainfall conditions. - Farmers are advised to undertake soil sample collection and testing before

	kharif crop establishment to assess soil fertility status and enable site-specific nutrient management based on soil test recommendations. Sow Dhaincha (<i>Sesbania aculeata</i>) @ 25–30 kg seed/ha or Sunhemp (<i>Crotalaria juncea</i>) @ 40–50 kg seed/ha as green manure crops and incorporate them into the soil at 45–50 days after sowing to improve soil organic carbon, soil structure, and nutrient availability for the succeeding crop.
Livestock	- Prevailing humid weather conditions may increase the risk of Hemorrhagic Septicemia (HS), Black Quarter (BQ), and parasitic infestations in livestock. Farmers are advised to vaccinate susceptible animals against HS and BQ in consultation with veterinary officials. - Maintain proper shed hygiene, drainage, and dry bedding to reduce pathogen buildup. Ensure adequate ventilation and provide clean drinking water to minimize disease incidence and environmental stress under humid conditions

AGRO METEOROLOGICAL ADVISORY FOR EAST GODAVARI DISTRICT

Crop	Stage of the crop/ Management	Recommendation
Paddy	Nursery Stage	- Utilize the available soil moisture for nursery bed preparation and sowing of recommended varieties. Ensure proper drainage to prevent water stagnation in nursery beds. - Treat seed with Carbendazim @ 2 g/kg seed or Trichoderma viride @ 10 g/kg seed before sowing to protect against seed-borne diseases. - Apply DAP @ 20 kg/acre of nursery area and monitor for blast disease under humid conditions. If symptoms appear, spray Tricyclazole 75 WP @ 0.6 g/L of water.
Coconut	Nut development	- Apply FYM/compost @ 25–50 kg/palm and neem cake @ 5 kg/palm under adequate soil moisture conditions. - Monitor for Rhinoceros beetle and Red palm weevil; install Rhynchophorol traps @ 1 trap/ha and apply neem cake @ 1–2 kg/palm in basins to reduce infestation during rain free period
Mango	Fruit development to harvest stage	- Maintain proper orchard drainage and remove fallen fruits to minimize fruit rotting and disease buildup under intermittent rainfall conditions. - Harvest mature fruits during rain-free periods and keep harvested produce in a shaded, well-ventilated place to improve shelf life and reduce post-harvest losses. - Monitor for fruit fly, and mango hopper incidence. If infestation is observed, spray Spinosad 45 SC @ 0.3 ml/L for fruit fly
Banana	bunch development stage	- Maintain proper drainage to prevent waterlogging and root-zone hypoxia. Provide propping to bearing plants to prevent lodging and pseudostem breakage under strong winds. - Apply neem cake @ 1–2 kg/plant and recommended fertilizers in split doses under adequate soil moisture conditions to support vegetative growth and bunch development - Monitor for Sigatoka leaf spot, pseudostem weevil, aphids, and thrips under humid conditions. For Sigatoka management, spray Propiconazole @ 1 ml/L or Hexaconazole @ 2 ml/L of water. - For severe aphid or thrips infestation, spray Imidacloprid 17.8 SL @ 0.3 ml/L or Thiamethoxam 25 WG @ 0.25 g/L of water during rain-free periods.
Sugarcane	Management	- Maintain proper field drainage to prevent waterlogging and ensure adequate soil aeration. Carry out earthing-up to improve root anchorage and minimize lodging. - Apply urea @ 55 kg/acre (or as per recommended split dose) under adequate soil moisture conditions and incorporate it into the soil to improve nitrogen-use efficiency. - Monitor for Pyrilla incidence, which may increase under warm and humid

		conditions. If infestation is observed, spray Imidacloprid 17.8 SL @ 0.3 ml/L or Thiamethoxam 25 WG @ 0.25 g/L of water during clear weather conditions.
Maize	Sowing Stage	- The forecasted rainfall provides favorable soil moisture for sowing; farmers are advised to take up sowing of suitable maize hybrids using a seed rate of 8–10 kg/acre and incorporate well-decomposed FYM or compost @ 2–4 t/acre during land preparation to improve soil health and moisture retention. - Treat seed with Trichoderma viride @ 10 g/kg seed or Metalaxyl + Mancozeb @ 2 g/kg seed before sowing to protect against seed- and soil-borne diseases and ensure uniform germination and healthy crop establishment.
Farm Preparatory Advisories		- Incorporation of farmyard manure, compost and crop residues during field preparation is advised to improve soil organic carbon and moisture retention capacity. - Undertake nursery bed preparation for Kharif vegetables and other seasonal crops with adequate drainage provisions to prevent seedling mortality under intermittent rainfall conditions. - Farmers are advised to undertake soil sample collection and testing before kharif crop establishment to assess soil fertility status and enable site-specific nutrient management based on soil test recommendations. - Sow Dhaincha (Sesbania aculeata) @ 25–30 kg seed/ha or Sunhemp (Crotalaria juncea) @ 40–50 kg seed/ha as green manure crops and incorporate them into the soil at 45–50 days after sowing to improve soil organic carbon, soil structure, and nutrient availability for the succeeding crop.
Livestock		- Prevailing humid weather conditions may increase the risk of Hemorrhagic Septicemia (HS), Black Quarter (BQ), and parasitic infestations in livestock. Farmers are advised to vaccinate susceptible animals against HS and BQ in consultation with veterinary officials. - Maintain proper shed hygiene, drainage, and dry bedding to reduce pathogen buildup. Ensure adequate ventilation and provide clean drinking water to minimize disease incidence and environmental stress under humid conditions

AGRO METEOROLOGICAL ADVISORY FOR KONASEEMA DISTRICT

Crop	Stage of the crop/ Management	Recommendation
Paddy	Nursery Stage	- Utilize the available soil moisture for nursery bed preparation and sowing of recommended varieties. Ensure proper drainage to prevent water stagnation in nursery beds. - Treat seed with Carbendazim @ 2 g/kg seed or Trichoderma viride @ 10 g/kg seed before sowing to protect against seed-borne diseases. - Apply DAP @ 20 kg/acre of nursery area and monitor for blast disease under humid conditions. If symptoms appear, spray Tricyclazole 75 WP @ 0.6 g/L of water.
Coconut	Nut development	- Apply FYM/compost @ 25–50 kg/palm and neem cake @ 5 kg/palm under adequate soil moisture conditions. - Monitor for Rhinoceros beetle and Red palm weevil; install Rhynchophorol traps @ 1 trap/ha and apply neem cake @ 1–2 kg/palm in basins to reduce infestation.
Mango	Fruit development to harvest stage	- Maintain proper orchard drainage and remove fallen fruits to minimize fruit rotting and disease buildup under intermittent rainfall conditions. - Harvest mature fruits during rain-free periods and keep harvested produce in a shaded, well-ventilated place to improve shelf life and reduce post-harvest losses. - Monitor for fruit fly, anthracnose, and mango hopper incidence. If

		infestation is observed, spray Spinosad 45 SC @ 0.3 ml/L for fruit fly or Carbendazim @ 1 g/L for anthracnose during clear weather conditions.
Banana	bunch development stage	• Maintain proper drainage to prevent waterlogging and root-zone hypoxia. Provide propping to bearing plants to prevent lodging and pseudostem breakage under strong winds. • Apply neem cake @ 1–2 kg/plant and recommended fertilizers in split doses under adequate soil moisture conditions to support vegetative growth and bunch development • Monitor for Sigatoka leaf spot, pseudostem weevil, aphids, and thrips under humid conditions. For Sigatoka management, spray Propiconazole @ 1 ml/L or Hexaconazole @ 2 ml/L of water. • For severe aphid or thrips infestation, spray Imidacloprid 17.8 SL @ 0.3 ml/L or Thiamethoxam 25 WG @ 0.25 g/L of water during rain-free periods.
Sugarcane	Management	• Maintain proper field drainage to prevent waterlogging and ensure adequate soil aeration. Carry out earthing-up to improve root anchorage and minimize lodging. • Apply urea @ 55 kg/acre (or as per recommended split dose) under adequate soil moisture conditions and incorporate it into the soil to improve nitrogen-use efficiency. • Monitor for Pyrilla incidence, which may increase under warm and humid conditions. If infestation is observed, spray Imidacloprid 17.8 SL @ 0.3 ml/L or Thiamethoxam 25 WG @ 0.25 g/L of water during clear weather conditions.
Maize	Sowing Stage	• The forecasted rainfall provides favorable soil moisture for sowing; farmers are advised to take up sowing of suitable maize hybrids using a seed rate of 8–10 kg/acre and incorporate well-decomposed FYM or compost @ 2–4 t/acre during land preparation to improve soil health and moisture retention. • Treat seed with Trichoderma viride @ 10 g/kg seed or Metalaxyl + Mancozeb @ 2 g/kg seed before sowing to protect against seed- and soil-borne diseases and ensure uniform germination and healthy crop establishment.
Farm Preparatory Advisories		• Incorporation of farmyard manure, compost and crop residues during field preparation is advised to improve soil organic carbon and moisture retention capacity. • Undertake nursery bed preparation for Kharif vegetables and other seasonal crops with adequate drainage provisions to prevent seedling mortality under intermittent rainfall conditions. • Farmers are advised to undertake soil sample collection and testing before kharif crop establishment to assess soil fertility status and enable site-specific nutrient management based on soil test recommendations. • Sow Dhaincha (Sesbania aculeata) @ 25–30 kg seed/ha or Sunhemp (Crotalaria juncea) @ 40–50 kg seed/ha as green manure crops and incorporate them into the soil at 45–50 days after sowing to improve soil organic carbon, soil structure, and nutrient availability for the succeeding crop.
Livestock		• Prevailing humid weather conditions may increase the risk of Hemorrhagic Septicemia (HS), Black Quarter (BQ), and parasitic infestations in livestock. Farmers are advised to vaccinate susceptible animals against HS and BQ in consultation with veterinary officials. • Maintain proper shed hygiene, drainage, and dry bedding to reduce pathogen buildup. Ensure adequate ventilation and provide clean drinking water to minimize disease incidence and environmental stress under humid conditions

AMFU GUNTUR**AGRO METEOROLOGICAL ADVISORY FOR GUNTUR & BAPATLA DISTRICTS****CROP ADVISORY**

Crop	Stage	Crop Specific Advisory
General Advisory		• Apply well decomposed farm yard manure @ 4 to 5 tonnes per acre before taking up the ploughing operation. • Farmers are advised to follow the contingency crop planning and weather-based agro-advisories issued by ANGRAU and the Department of Agriculture from time to time. • In view of the likely El Niño conditions, adopt moisture conservation practices to reduce moisture loss. • Prefer short-duration and drought-tolerant varieties in rainfed areas to minimize the risk of moisture stress during crop growth. • Strengthen rainwater harvesting through farm ponds and field channels to provide supplemental irrigation during dry spells. • Use irrigation water judiciously and prioritize irrigation during critical crop growth stages. • Keep contingency crop seeds and alternate cropping options ready in case of delayed monsoon onset or prolonged dry spells. • Regularly monitor weather forecasts and undertake sowing, fertilizer application, and other crop management practices accordingly.
Rice	Nursey to Transplanting stage	• The forecasted light rainfall and favorable temperature conditions are suitable for initiating paddy nursery sowing. Prepare the nursery field with 2–3 ploughings followed by leveling and ensure adequate irrigation and drainage facilities. • Use 20 kg of certified seed acre⁻¹ for nursery raising. Select high-yielding, locally recommended varieties suitable for the region.. • Treat seed with carbendazim before sowing (dry treatment: 3 g kg⁻¹seed; wet treatment: 1 g kg⁻¹ seed, soak for 24 hours and incubate for another 24 hours). • Apply 1 kg urea, 2 kg SSP and 0.5 kg MOP per 100 m² nursery area as a basal dose. If seedlings become pale, apply 1 kg urea per 100 m² at 10–12 DAS. • Maintain optimum moisture with a 2–3 cm water depth and ensure proper drainage to prevent waterlogging and seedling diseases. • Transplant healthy 20–25-day-old seedlings during the morning or evening. • Spray 19:19:19 @ 1% or 13:00:45 (potassium nitrate) @ 1% after transplanting to improve crop establishment and reduce heat stress. • If nursery sowing is delayed, raise and incorporate green manure crops such as sunhemp or dhaincha before transplanting.
Maize	Sowing to seedling stage	• Farmers are advised to take up maize sowing after receipt of adequate rainfall and availability of sufficient soil moisture. Under prevailing El Niño conditions, prefer medium-duration, drought-tolerant popular hybrids.

		• Treat seed with <i>Trichoderma viride</i> @ 10 g kg⁻¹ seed before sowing to minimize seed and soil-borne diseases. • Apply fertilizers based on soil test recommendations. Where soil test results are not available, follow the recommended fertilizer dose and apply nutrients in split applications for efficient utilization. • Apply Atrazine @ 800 g acre⁻¹ (heavy soils) or 1200 g acre⁻¹ (light soils) within 1–2 days after sowing. • Ensure proper drainage during heavy rains and conserve soil moisture through timely intercultivation and earthing-up. • Monitor regularly for fall armyworm and manage promptly. • Provide protective irrigation during dry spells. • Spray 19:19:19 @ 1%, 20:20:20 @ 1%, or ZnSO₄ @ 0.5% + lime @ 0.25% as needed to improve crop growth and correct nutrient deficiency.
Cotton	Sowing to seedling stage	• Initiate cotton sowing after receiving adequate rainfall to ensure proper seed germination and crop establishment. • Under prevailing El Niño conditions, prefer short to medium-duration hybrids and complete sowing within a short and uniform period across the area to minimize pink bollworm incidence. • Treat seed with <i>Trichoderma viride</i> @ 10 g kg⁻¹ seed before sowing and maintain the recommended plant spacing. • Apply 4–5 tonnes FYM acre⁻¹ during land preparation and fertilizers as per soil test recommendations. • Provide light protective irrigation whenever required and carry out intercultivation to conserve soil moisture. • Foliar nutrient recommendation: Spray 19:19:19 @ 1% + MgSO₄ @ 1% or boron @ 0.2% (where deficiency is observed).
Redgram	Sowing	• Light, red and black soils with good drainage is good for redgram. Saline soils and water logged soils are not suitable. • Sow redgram after receipt of adequate rainfall to ensure sufficient soil moisture for successful crop establishment. Use certified, drought-tolerant varieties. • Treat seed with <i>Rhizobium</i> and PSB cultures along with <i>Trichoderma viride</i> @ 10 g kg⁻¹ seed before sowing to improve nodulation and reduce wilt incidence. • Apply well-decomposed FYM @ 4–5 tonnes acre⁻¹ during land preparation and follow soil test-based fertilizer recommendations.
Blackgram	Sowing to seedling stage	• Based on available soil moisture, farmers are advised to take up sowing of blackgram using suitable varieties such as LBG-884, LBG-904 & LBG-932 & TB-104 and sow at a seed rate of 8-10 kg acre⁻¹ duly following seed treatment with Thiram / Captan/ Mancozeb/ Carbendazim @ 2.5 g kg⁻¹ seed along with Imidacloprid @ 5 ml kg⁻¹ or Thiamethoxam @ 5 g kg⁻¹ seed to protect the crop from sucking pests and diseases. • At the time of sowing, apply the recommended basal fertilizer dose of 8 kg nitrogen and 20 kg phosphorus acre⁻¹. • Apply Pendimethalin 30% EC @ 1.0–1.5 litres acre⁻¹ immediately after sowing or on the following day as a pre-

		emergence herbicide. • Conserve soil moisture through timely intercultivation and other moisture conservation practices. • Foliar nutrient recommendation: Spray DAP @ 2%, 19:19:19 @ 1%, or a multi-micronutrient mixture @ 0.5% to promote crop growth and correct nutrient deficiencies.
Greengram	Sowing to seedling stage	• Based on available soil moisture, farmers are advised to take up sowing of greengram using suitable varieties such as LGG-630, LGG-607, WGG-42, and IPM-2-14 at a seed rate of 8 kg acre⁻¹ duly following seed treatment with Thiram / Captan/ Mancozeb/ Carbendazim @ 2.5 g kg⁻¹ seed along with Imidacloprid @ 5 ml kg⁻¹ or Thiamethoxam @ 5 g kg⁻¹ seed to protect the crop from sucking pests and diseases.. • At the time of sowing, apply the recommended basal fertilizer dose of 8 kg nitrogen and 20 kg phosphorus acre⁻¹. • Apply Pendimethalin 30% EC @ 1.3-1.6 litres acre⁻¹ immediately after sowing or on the following day as a pre-emergence herbicide. • Conserve soil moisture through timely intercultivation and other moisture conservation practices. • Foliar nutrient recommendation: Spray DAP @ 2%, 19:19:19 @ 1%, or a multi-micronutrient mixture @ 0.5% to promote crop growth and correct nutrient deficiencies.
Livestock		Livestock Specific Advisory
Buffaloes		• During thunderstorms and lightning, keep buffaloes indoors in a safe, dry, and well-ventilated shed. Do not allow grazing in open fields or near trees during thunderstorms. • Provide clean drinking water, quality green fodder, dry fodder, and mineral mixture daily. • Avoid waterlogged areas and maintain proper drainage around the animal shed. • Maintain udder hygiene and keep the shed clean and dry to prevent mastitis and other infections. • Monitor animals for signs of haemorrhagic septicaemia (HS), black quarter (BQ), foot-and-mouth disease (FMD), and parasitic infestations. Vaccinate as per the veterinary schedule and consult a veterinarian immediately if any symptoms of illness are observed.

AGRO METEOROLOGICAL ADVISORY FOR PALNADU & PRAKASAM DISTRICTS

CROP ADVISORY		
Crop	Stage	Crop Specific Advisory
General Advisory		• Apply well decomposed farm yard manure @ 4 to 5 tonnes per acre before taking up the ploughing operation. • Farmers are advised to follow the contingency crop planning and weather-based agro-advisories issued by ANGRAU and the Department of Agriculture from time to time. • In view of the likely El Niño conditions, adopt moisture conservation practices to reduce moisture loss. • Prefer short-duration and drought-tolerant varieties in rainfed areas to minimize the risk of moisture stress during crop

		growth. • Strengthen rainwater harvesting through farm ponds and field channels to provide supplemental irrigation during dry spells. • Use irrigation water judiciously and prioritize irrigation during critical crop growth stages. • Keep contingency crop seeds and alternate cropping options ready in case of delayed monsoon onset or prolonged dry spells. • Regularly monitor weather forecasts and undertake sowing, fertilizer application, and other crop management practices accordingly.
Rice	Nursey to Transplanting stage	• The forecasted light rainfall and favorable temperature conditions are suitable for initiating paddy nursery sowing. Prepare the nursery field with 2–3 ploughings followed by leveling and ensure adequate irrigation and drainage facilities. • Use 20 kg of certified seed acre⁻¹ for nursery raising. Select high-yielding, locally recommended varieties suitable for the region.. • Treat seed with carbendazim before sowing (dry treatment: 3 g kg⁻¹seed; wet treatment: 1 g kg⁻¹ seed, soak for 24 hours and incubate for another 24 hours). • Apply 1 kg urea, 2 kg SSP and 0.5 kg MOP per 100 m² nursery area as a basal dose. If seedlings become pale, apply 1 kg urea per 100 m² at 10–12 DAS. • Maintain optimum moisture with a 2–3 cm water depth and ensure proper drainage to prevent waterlogging and seedling diseases. • Transplant healthy 20–25-day-old seedlings during the morning or evening. • Spray 19:19:19 @ 1% or 13:00:45 (potassium nitrate) @ 1% after transplanting to improve crop establishment and reduce heat stress. • If nursery sowing is delayed, raise and incorporate green manure crops such as sunhemp or dhaincha before transplanting.
Maize	Sowing to seedling stage	• Farmers are advised to take up maize sowing after receipt of adequate rainfall and availability of sufficient soil moisture. Under prevailing El Niño conditions, prefer medium-duration, drought-tolerant popular hybrids. • Treat seed with <i>Trichoderma viride</i> @ 10 g kg⁻¹ seed before sowing to minimize seed and soil-borne diseases. • Apply fertilizers based on soil test recommendations. Where soil test results are not available, follow the recommended fertilizer dose and apply nutrients in split applications for efficient utilization. • Apply Atrazine @ 800 g acre⁻¹ (heavy soils) or 1200 g acre⁻¹ (light soils) within 1–2 days after sowing. • Ensure proper drainage during heavy rains and conserve soil moisture through timely intercultivation and earthing-up. • Monitor regularly for fall armyworm and manage promptly.

		• Provide protective irrigation during dry spells. • Spray 19:19:19 @ 1%, 20:20:20 @ 1%, or ZnSO₄ @ 0.5% + lime @ 0.25% as needed to improve crop growth and correct nutrient deficiency.
Cotton	Sowing to seedling stage	• Initiate cotton sowing after receiving adequate rainfall to ensure proper seed germination and crop establishment. • Under prevailing El Niño conditions, prefer short to medium-duration hybrids and complete sowing within a short and uniform period across the area to minimize pink bollworm incidence. • Treat seed with <i>Trichoderma viride</i> @ 10 g kg⁻¹ seed before sowing and maintain the recommended plant spacing. • Apply 4–5 tonnes FYM acre⁻¹ during land preparation and fertilizers as per soil test recommendations. • Provide light protective irrigation whenever required and carry out intercultivation to conserve soil moisture. • Foliar nutrient recommendation: Spray 19:19:19 @ 1% + MgSO₄ @ 1% or boron @ 0.2% (where deficiency is observed).
Redgram	Sowing	• Light, red and black soils with good drainage is good for redgram. Saline soils and water logged soils are not suitable. • Sow redgram after receipt of adequate rainfall to ensure sufficient soil moisture for successful crop establishment. Use certified, drought-tolerant varieties. • Treat seed with <i>Rhizobium</i> and PSB cultures along with <i>Trichoderma viride</i> @ 10 g kg⁻¹ seed before sowing to improve nodulation and reduce wilt incidence. • Apply well-decomposed FYM @ 4–5 tonnes acre⁻¹ during land preparation and follow soil test-based fertilizer recommendations.
Blackgram	Sowing to seedling stage	• Based on available soil moisture, farmers are advised to take up sowing of blackgram using suitable varieties such as LBG-884, LBG-904 & LBG-932 & TB-104 and sow at a seed rate of 8-10 kg acre⁻¹ duly following seed treatment with Thiram / Captan/ Mancozeb/ Carbendazim @ 2.5 g kg⁻¹ seed along with Imidacloprid @ 5 ml kg⁻¹ or Thiamethoxam @ 5 g kg⁻¹ seed to protect the crop from sucking pests and diseases. • At the time of sowing, apply the recommended basal fertilizer dose of 8 kg nitrogen and 20 kg phosphorus acre⁻¹. • Apply Pendimethalin 30% EC @ 1.0–1.5 litres acre⁻¹ immediately after sowing or on the following day as a pre-emergence herbicide. • Conserve soil moisture through timely intercultivation and other moisture conservation practices. • Foliar nutrient recommendation: Spray DAP @ 2%, 19:19:19 @ 1%, or a multi-micronutrient mixture @ 0.5% to promote crop growth and correct nutrient deficiencies.
Greengram	Sowing to seedling stage	• Based on available soil moisture, farmers are advised to take up sowing of greengram using suitable varieties such as LGG-630, LGG-607, WGG-42, and IPM-2-14 at a seed rate of 8 kg acre⁻¹ duly following seed treatment with Thiram / Captan/ Mancozeb/ Carbendazim @ 2.5 g kg⁻¹ seed along

		with Imidacloprid @ 5 ml kg⁻¹ or Thiamethoxam @ 5 g kg⁻¹ seed to protect the crop from sucking pests and diseases.. • At the time of sowing, apply the recommended basal fertilizer dose of 8 kg nitrogen and 20 kg phosphorus acre⁻¹. • Apply Pendimethalin 30% EC @ 1.3-1.6 litres acre⁻¹ immediately after sowing or on the following day as a pre-emergence herbicide. • Conserve soil moisture through timely intercultivation and other moisture conservation practices. • Foliar nutrient recommendation: Spray DAP @ 2%, 19:19:19 @ 1%, or a multi-micronutrient mixture @ 0.5% to promote crop growth and correct nutrient deficiencies.
Livestock	Livestock Specific Advisory	
Buffaloes		• During thunderstorms and lightning, keep buffaloes indoors in a safe, dry, and well-ventilated shed. Do not allow grazing in open fields or near trees during thunderstorms. • Provide clean drinking water, quality green fodder, dry fodder, and mineral mixture daily. • Avoid waterlogged areas and maintain proper drainage around the animal shed. • Maintain udder hygiene and keep the shed clean and dry to prevent mastitis and other infections. • Monitor animals for signs of haemorrhagic septicaemia (HS), black quarter (BQ), foot-and-mouth disease (FMD), and parasitic infestations. Vaccinate as per the veterinary schedule and consult a veterinarian immediately if any symptoms of illness are observed.

AGRO METEOROLOGICAL ADVISORY FOR KRISHNA & NTR DISTRICTS

CROP ADVISORY		
Crop	Stage	Crop Specific Advisory
General Advisory		• Apply well decomposed farm yard manure @ 4 to 5 tonnes per acre before taking up the ploughing operation. • Farmers are advised to follow the contingency crop planning and weather-based agro-advisories issued by ANGRAU and the Department of Agriculture from time to time. • In view of the likely El Niño conditions, adopt moisture conservation practices to reduce moisture loss. • Prefer short-duration and drought-tolerant varieties in rainfed areas to minimize the risk of moisture stress during crop growth. • Strengthen rainwater harvesting through farm ponds and field channels to provide supplemental irrigation during dry spells. • Use irrigation water judiciously and prioritize irrigation during critical crop growth stages. • Keep contingency crop seeds and alternate cropping options ready in case of delayed monsoon onset or prolonged dry spells. • Regularly monitor weather forecasts and undertake sowing, fertilizer application, and other crop management practices accordingly.

Rice	Nursey to Transplanting stage	• The forecasted light rainfall and favorable temperature conditions are suitable for initiating paddy nursery sowing. Prepare the nursery field with 2–3 ploughings followed by leveling and ensure adequate irrigation and drainage facilities. • Use 20 kg of certified seed acre⁻¹ for nursery raising. Select high-yielding, locally recommended varieties suitable for the region.. • Treat seed with carbendazim before sowing (dry treatment: 3 g kg⁻¹seed; wet treatment: 1 g kg⁻¹ seed, soak for 24 hours and incubate for another 24 hours). • Apply 1 kg urea, 2 kg SSP and 0.5 kg MOP per 100 m² nursery area as a basal dose. If seedlings become pale, apply 1 kg urea per 100 m² at 10–12 DAS. • Maintain optimum moisture with a 2–3 cm water depth and ensure proper drainage to prevent waterlogging and seedling diseases. • Transplant healthy 20–25-day-old seedlings during the morning or evening. • Spray 19:19:19 @ 1% or 13:00:45 (potassium nitrate) @ 1% after transplanting to improve crop establishment and reduce heat stress. • If nursery sowing is delayed, raise and incorporate green manure crops such as sunhemp or dhaincha before transplanting.
Maize	Sowing to seedling stage	• Farmers are advised to take up maize sowing after receipt of adequate rainfall and availability of sufficient soil moisture. Under prevailing El Niño conditions, prefer medium-duration, drought-tolerant popular hybrids. • Treat seed with <i>Trichoderma viride</i> @ 10 g kg⁻¹ seed before sowing to minimize seed and soil-borne diseases. • Apply fertilizers based on soil test recommendations. Where soil test results are not available, follow the recommended fertilizer dose and apply nutrients in split applications for efficient utilization. • Apply Atrazine @ 800 g acre⁻¹ (heavy soils) or 1200 g acre⁻¹ (light soils) within 1–2 days after sowing. • Ensure proper drainage during heavy rains and conserve soil moisture through timely intercultivation and earthing-up. • Monitor regularly for fall armyworm and manage promptly. • Provide protective irrigation during dry spells. • Spray 19:19:19 @ 1%, 20:20:20 @ 1%, or ZnSO₄ @ 0.5% + lime @ 0.25% as needed to improve crop growth and correct nutrient deficiency.
Cotton	Sowing to seedling stage	• Initiate cotton sowing after receiving adequate rainfall to ensure proper seed germination and crop establishment. • Under prevailing El Niño conditions, prefer short to medium-duration hybrids and complete sowing within a short and uniform period across the area to minimize pink bollworm incidence. • Treat seed with <i>Trichoderma viride</i> @ 10 g kg⁻¹ seed before sowing and maintain the recommended plant spacing. • Apply 4–5 tonnes FYM acre⁻¹ during land preparation and fertilizers as per soil test recommendations. • Provide light protective irrigation whenever required and carry

		out intercultivation to conserve soil moisture. Foliar nutrient recommendation: Spray 19:19:19 @ 1% + MgSO₄ @ 1% or boron @ 0.2% (where deficiency is observed).
Redgram	Sowing	- Light, red and black soils with good drainage is good for redgram. Saline soils and water logged soils are not suitable. - Sow redgram after receipt of adequate rainfall to ensure sufficient soil moisture for successful crop establishment. Use certified, drought-tolerant varieties. - Treat seed with Rhizobium and PSB cultures along with <i>Trichoderma viride</i> @ 10 g kg⁻¹ seed before sowing to improve nodulation and reduce wilt incidence. - Apply well-decomposed FYM @ 4–5 tonnes acre⁻¹ during land preparation and follow soil test-based fertilizer recommendations.
Blackgram	Sowing to seedling stage	- Based on available soil moisture, farmers are advised to take up sowing of blackgram using suitable varieties such as LBG-884, LBG-904 & LBG-932 & TB-104 and sow at a seed rate of 8-10 kg acre⁻¹ duly following seed treatment with Thiram / Captan/ Mancozeb/ Carbendazim @ 2.5 g kg⁻¹ seed along with Imidacloprid @ 5 ml kg⁻¹ or Thiamethoxam @ 5 g kg⁻¹ seed to protect the crop from sucking pests and diseases. - At the time of sowing, apply the recommended basal fertilizer dose of 8 kg nitrogen and 20 kg phosphorus acre⁻¹. - Apply Pendimethalin 30% EC @ 1.0–1.5 litres acre⁻¹ immediately after sowing or on the following day as a pre-emergence herbicide. - Conserve soil moisture through timely intercultivation and other moisture conservation practices. - Foliar nutrient recommendation: Spray DAP @ 2%, 19:19:19 @ 1%, or a multi-micronutrient mixture @ 0.5% to promote crop growth and correct nutrient deficiencies.
Greengram	Sowing to seedling stage	- Based on available soil moisture, farmers are advised to take up sowing of greengram using suitable varieties such as LGG-630, LGG-607, WGG-42, and IPM-2-14 at a seed rate of 8 kg acre⁻¹ duly following seed treatment with Thiram / Captan/ Mancozeb/ Carbendazim @ 2.5 g kg⁻¹ seed along with Imidacloprid @ 5 ml kg⁻¹ or Thiamethoxam @ 5 g kg⁻¹ seed to protect the crop from sucking pests and diseases.. - At the time of sowing, apply the recommended basal fertilizer dose of 8 kg nitrogen and 20 kg phosphorus acre⁻¹. - Apply Pendimethalin 30% EC @ 1.3-1.6 litres acre⁻¹ immediately after sowing or on the following day as a pre-emergence herbicide. - Conserve soil moisture through timely intercultivation and other moisture conservation practices. - Foliar nutrient recommendation: Spray DAP @ 2%, 19:19:19 @ 1%, or a multi-micronutrient mixture @ 0.5% to promote crop growth and correct nutrient deficiencies.
Livestock	Livestock Specific Advisory	
Buffaloes		- During thunderstorms and lightning, keep buffaloes indoors in a safe, dry, and well-ventilated shed. Do not allow grazing in open fields or near trees during thunderstorms. - Provide clean drinking water, quality green fodder, dry fodder,

	and mineral mixture daily. • Avoid waterlogged areas and maintain proper drainage around the animal shed. • Maintain udder hygiene and keep the shed clean and dry to prevent mastitis and other infections. • Monitor animals for signs of haemorrhagic septicaemia (HS), black quarter (BQ), foot-and-mouth disease (FMD), and parasitic infestations. Vaccinate as per the veterinary schedule and consult a veterinarian immediately if any symptoms of illness are observed.
--	--

AGRO METEOROLOGICAL ADVISORY FOR WEST GODAVARI & ELURU DISTRICTS

Crop	Stage of the crop/ Management	Recommendation
Paddy	Nursery Stage	• Prepare nursery beds and sow recommended varieties under adequate soil moisture conditions. Ensure proper drainage for healthy seedling growth. • Treat seed with Carbendazim @ 2 g kg⁻¹ seed or Trichoderma viride @ 10 g kg⁻¹ seed before sowing to prevent seed-borne diseases. • Apply DAP @ 20 kg acre⁻¹ of nursery area. Monitor for blast disease under humid conditions and spray Tricyclazole 75 WP @ 0.6 g L⁻¹ water if symptoms appear.
Coconut	But development	• Apply well-decomposed farmyard manure (FYM) or compost @ 25–50 kg palm⁻¹ along with neem cake @ 5 kg palm⁻¹ when sufficient soil moisture is available. • Regularly inspect palms for Rhinoceros beetle and Red palm weevil infestation. Install Rhynchophorol pheromone traps @ 1 trap ha⁻¹ and incorporate neem cake @ 1–2 kg palm⁻¹ in the basin area to minimize pest incidence.
Mango	Harvesting and Post-harvest Operations	• Harvest mature mango fruits during dry weather conditions and avoid harvesting during rainfall, thunderstorms and strong winds to minimize fruit damage. • Collect fallen fruits separately and remove damaged or diseased fruits to prevent post-harvest losses. • Grade, pack and store harvested fruits in well-ventilated and dry places protected from rainfall, direct sunlight and high temperatures. • Secure harvested fruits during transportation and marketing to avoid mechanical injury caused by squally winds and adverse weather conditions.
Banana	Bunch development stage	• Ensure efficient field drainage to avoid water stagnation and oxygen deficiency in the root zone. Support fruit-bearing plants with sturdy props to protect them from lodging and pseudostem damage during periods of strong winds. • Apply neem cake @ 1–2 kg plant⁻¹ and supply the recommended fertilizer dose in splits under adequate soil moisture conditions to promote healthy plant growth and improved bunch formation. • Regularly inspect the crop for Sigatoka leaf spot, pseudostem weevil, aphids, and thrips, especially during

		humid weather. To manage Sigatoka disease, spray Propiconazole @ 1 ml L⁻¹ or Hexaconazole @ 2 ml L⁻¹ of water. • In case of heavy infestation of aphids or thrips, apply Imidacloprid 17.8 SL @ 0.3 ml L⁻¹ or Thiamethoxam 25 WG @ 0.25 g L⁻¹ of water during dry weather conditions free from rainfall.
Sugarcane	Management	• Maintain proper field drainage to prevent waterlogging and ensure adequate soil aeration. Carry out earthing-up to improve root anchorage and minimize lodging. • Apply urea @ 55 kg acre⁻¹ (or as per recommended split dose) under adequate soil moisture conditions and incorporate it into the soil to improve nitrogen-use efficiency. • Monitor for <i>Pyrilla</i> incidence, which may increase under warm and humid conditions. If infestation is observed, spray Imidacloprid 17.8 SL @ 0.3 ml L⁻¹ or Thiamethoxam 25 WG @ 0.25 g L⁻¹ of water during clear weather conditions.
Maize	Sowing to seedling stage	• Farmers are advised to take up maize sowing after receipt of adequate rainfall and availability of sufficient soil moisture. Under prevailing El Niño conditions, prefer medium-duration, drought-tolerant popular hybrids. • Treat seed with <i>Trichoderma viride</i> @ 10 g kg⁻¹ seed before sowing to minimize seed and soil-borne diseases. • Apply fertilizers based on soil test recommendations. Where soil test results are not available, follow the recommended fertilizer dose and apply nutrients in split applications for efficient utilization. • Apply Atrazine @ 800 g acre⁻¹ (heavy soils) or 1200 g acre⁻¹ (light soils) within 1–2 days after sowing. • Ensure proper drainage during heavy rains and conserve soil moisture through timely intercultivation and earthing-up. • Monitor regularly for fall armyworm and manage promptly. • Provide protective irrigation during dry spells. • Spray 19:19:19 @ 1%, 20:20:20 @ 1%, or ZnSO₄ @ 0.5% + lime @ 0.25% as needed to improve crop growth and correct nutrient deficiency.
General Advisory		• Apply well decomposed farm yard manure @ 4 to 5 tonnes per acre before taking up the ploughing operation. • Farmers are advised to follow the contingency crop planning and weather-based agro-advisories issued by ANGRAU and the Department of Agriculture from time to time. • In view of the likely El Niño conditions, adopt moisture conservation practices to reduce moisture loss. • Prefer short-duration and drought-tolerant varieties in rainfed areas to minimize the risk of moisture stress during crop growth. • Strengthen rainwater harvesting through farm ponds and field channels to provide supplemental irrigation during dry spells. • Use irrigation water judiciously and prioritize irrigation during critical crop growth stages. • Keep contingency crop seeds and alternate cropping options ready in case of delayed monsoon onset or prolonged dry spells.

	Regularly monitor weather forecasts and undertake sowing, fertilizer application, and other crop management practices accordingly.
Livestock	- During thunderstorms and lightning, keep buffaloes indoors in a safe, dry, and well-ventilated shed. Do not allow grazing in open fields or near trees during thunderstorms. - Provide clean drinking water, quality green fodder, dry fodder, and mineral mixture daily. - Avoid waterlogged areas and maintain proper drainage around the animal shed. - Maintain udder hygiene and keep the shed clean and dry to prevent mastitis and other infections. - Monitor animals for signs of haemorrhagic septicaemia (HS), black quarter (BQ), foot-and-mouth disease (FMD), and parasitic infestations. Vaccinate as per the veterinary schedule and consult a veterinarian immediately if any symptoms of illness are observed.

Meteorological Subdivision of Rayalaseema:

AMFU TIRUPATI

AGRO METEOROLOGICAL ADVISORY FOR TIRUPATI DISTRICT

Likely impacts of weather warnings on Agriculture and associated Agromet advisories

In view of thunderstorms, lightening and squalls farmers are advised to harvest the matured fruits and vegetables and take care of strong staking in banana gardens.

GENERAL ADVISORY

- Farmers who have not yet sown groundnut crop, still have time to sow until the end of this month. Sowing should be taken up only after receipt of sufficient rainfall and enough moisture is available in the soil.

SMS ADVISORY

- In view of scattered and very light rainfall and high temperature, farmers are advised to monitor and irrigate the crops if necessary

CROP SPECIFIC ADVISORY

Select category: crop

Select Type: G.Nut

- Farmers who have not yet sown groundnut crop, still have time to sow until the end of this month. Sowing should be taken up only after receipt of sufficient rainfall and enough moisture is available in the soil. Before sowing, treat the seed with Imidachloprid 600 F.S @ 2 ml. + 4 ml water per kg of seed followed by Mancozeb 3 g/kg or Tebuconazole @ 1 g/kg of seed to control sucking pest and other seed borne diseases.
- Farmers who have sown groundnut during June month and is at flowering stage are advised to provide one irrigation to prevent moisture stress conditions.

CROP SPECIFIC ADVISORY

Select category: crop

Select Type: Paddy

- The farmers who have taken up paddy nurseries are advised to apply carbofuran 3g granules @ 160 g. per cent of nursery 1 week prior to pulling of the nursery.

Farmers who are taking up transplanting of paddy nurseries are advised to

- To avoid attack of stem borer and leaf folder clip the tips of the leaves before transplanting
- To avoid BPH attack leave 20cm. spacing between the rows
- Apply entire 'P2O5' & 'K2O' as basal and 1/3 'N' as basal during last puddling .
- Transplant 30-35 days old rice seedlings.
- Leave 20 cm pathways for every 2 mt. of transplanted crop.

Select category:

Select Type: Maize

Select variety:

- The present weather conditions are favourable for fall army worm in Maize crop under irrigated conditions at vegetative stage. To control spray Neem oil @ 5ml/L of water in early vegetative stage crop.
- If the maize crop is at 30 DAS, spray Larvin @ 1 g/L of water.

Select category:

Select Type: Minor millets

Select variety:

- Farmers who are taking up ragi, korra (minor millets) sowings are advised to treat the seed compulsorily with carbendazim @ 2 g. or mancozeb @ 3 g./kg of seed.

Select category:

Select Type: CITRUS

Select variety:

- The present weather conditions are favourable for citrus canker in citrus. To control spray copper oxychloride @ 3 g./L of water + Streptocyclin @ 0.1 g./L of water.

Select category:

Select Type: Cotton

Stage:

- The present weather conditions are favourable for sucking pests in cotton. To control spray fipronil @ 2 ml./L (or) profenophos @ 2 ml./L of water.

Likely impacts of weather warnings (General)

- Due to the possibility of heavy winds, livestock and goats should be kept in safe areas and provided with fodder.

Impact based advisories (General)

- In view of heavy winds stake the plants in banana gardens.

AGRO METEOROLOGICAL ADVISORY FOR CHITTOOR DISTRICT

Likely impacts of weather warnings on Agriculture and associated Agromet advisories

In view of thunderstorms, lightening and squalls farmers are advised to harvest the matured fruits and vegetables and take care of strong staking in banana gardens.

GENERAL ADVISORY

- Farmers who have not yet sown groundnut crop, still have time to sow until the end of this month. Sowing should be taken up only after receipt of sufficient rainfall and enough moisture is available in the soil.

SMS ADVISORY

- In view of scattered and very light rainfall and high temperature, farmers are advised to monitor and irrigate the crops if necessary

CROP SPECIFIC ADVISORY

Select category: crop

Select Type: Paddy

- The farmers who have taken up paddy nurseries are advised to apply carbofuran 3g granules @ 160 g. per cent of nursery 1 week prior to pulling of the nursery.

Farmers who are taking up transplanting of paddy nurseries are advised to

- To avoid attack of stem borer and leaf folder clip the tips of the leaves before transplanting
- To avoid BPH attack leave 20cm. spacing between the rows
- Apply entire 'P2O5' & 'K2O' as basal and 1/3 'N' as basal during last puddling.
- Transplant 30-35 days old rice seedlings.
- Leave 20 cm pathways for every 2 mt. of transplanted crop.

CROP SPECIFIC ADVISORY

Select category: crop

Select Type: G.Nut

- Farmers who have not yet sown groundnut crop, still have time to sow until the end of this month. Sowing should be taken up only after receipt of sufficient rainfall and enough moisture is available in the soil. Before sowing, treat the seed with Imidachloprid 600 F.S @ 2 ml. + 4 ml water per kg of seed followed by Mancozeb 3 g/kg or Tebuconazole @ 1 g/kg of seed to control sucking pest and other seed borne diseases.

- Farmers who have sown groundnut during June month and is at flowering stage are advised to provide one irrigation to prevent moisture stress conditions.

Select category: Crop

Select Type: Black gram

Select variety:

- The present weather conditions are favourable for sucking pest in black gram. To control spray Imidachloprid @ 0.4ml/L + 19-19-19 @ 10g L of water
- Farmers who are taking up sowing of blackgram and green gram under irrigations are advised to follow seed treatment with 2.5 g thiram / kaptan / mancozeb / carbendazem and imidachloprid @ 5 ml./kg of seed.

Select category:

Select Type: Maize

Select variety:

- The present weather conditions are favourable for fall army worm in Maize crop under irrigated conditions at vegetative stage. To control spray Neem oil @ 5ml/L of water in early vegetative stage crop.
- If the maize crop is at 30 DAS, spray Larvin @ 1 g/L of water.

Select category:

Select Type: Minor millets

Select variety:

- Farmers who are taking up ragi, korra (minor millets) sowings are advised to treat the seed compulsorily with carbendazim @ 2 g. or mancozeb @ 3 g./kg of seed.

Likely impacts of weather warnings (General)

- Due to the possibility of surface winds, livestock and goats should be kept in safe areas and provided with fodder.

Impact based advisories (General)

- In view of heavy winds stake the plants in banana gardens.

AGRO METEOROLOGICAL ADVISORY FOR SPSR-NELLORE DISTRICT

Likely impacts of weather warnings on Agriculture and associated Agromet advisories

In view of strong surface winds farmers are advised to harvest the matured fruits and vegetables and take care of strong staking in banana gardens.

GENERAL ADVISORY

- Farmers who have not yet sown groundnut crop, still have time to sow until the end of this month. Sowing should be taken up only after receipt of sufficient rainfall and enough moisture is available in the soil.

SMS ADVISORY

- In view of scattered and very light rainfall and high temperature, farmers are advised to monitor and irrigate the crops if necessary

CROP SPECIFIC ADVISORY

Select category: crop

Select Type: Paddy

- The farmers who have taken up paddy nurseries are advised to apply carbofuran 3g granules @ 160 g. per cent of nursery 1 week prior to pulling of the nursery.

Farmers who are taking up transplanting of paddy nurseries are advised to

- To avoid attack of stem borer and leaf folder clip the tips of the leaves before transplanting
- To avoid BPH attack leave 20cm. spacing between the rows
- Apply entire 'P2O5' & 'K2O' as basal and 1/3 'N' as basal during last puddling.
- Transplant 30-35 days old rice seedlings.
- Leave 20 cm pathways for every 2 mt. of transplanted crop.

CROP SPECIFIC ADVISORY

Select category: crop

Select Type: G.Nut

- Farmers who have not sown groundnut crop, still have time to sow until the end of this month. Sowing should be taken up only after receipt of sufficient rainfall and enough moisture is available in the soil. Before sowing, treat the seed with Imidachloprid 600 F.S @ 2 ml. + 4 ml water per kg of

seed followed by Mancozeb 3 g/kg or Tebuconazole @ 1 g/kg of seed to control sucking pest and other seed borne diseases.

- Farmers who have sown groundnut during June month and is at flowering stage are advised to provide one irrigation to prevent moisture stress conditions.

Select category: **Select Type: Maize** **Select variety:**

- The present weather conditions are favourable for fall army worm in Maize crop under irrigated conditions at vegetative stage. To control spray Neem oil @ 5ml/L of water in early vegetative stage crop.
- If the maize crop is at 30 DAS, spray Larvin @ 1 g/L of water.

Select category: **Select Type: Minor millets** **Select variety:**

- Farmers who are taking up ragi, korra (minor millets) sowings are advised to treat the seed compulsorily with carbendazim @ 2 g. or mancozeb @ 3 g./kg of seed.

Select category: Crop **Select Type: Black gram** **Select variety:**

- The present weather conditions are favourable for sucking pest in black gram. To control spray Imidachloprid @ 0.4ml/L + 19-19-19 @ 10g L of water
- Farmers who are taking up sowing of blackgram and green gram under irrigations are advised to follow seed treatment with 2.5 g thiram / kaptan / mancozeb / carbendzem and imidachloprid @ 5 ml./kg of seed.

Select category: **Select Type: Cotton** **Stage:**

- The present weather conditions are favourable for sucking pests in cotton. To control spray fipronil @ 2 ml./L (or) profenophos @ 2 ml./L of water.

Likely impacts of weather warnings (General)

- Due to the possibility of heavy winds, livestock and goats should be kept in safe areas and provided with fodder.

Impact based advisories (General)

- In view of heavy winds stake the plants in banana gardens.

AGRO METEOROLOGICAL ADVISORY FOR YSR KADAPA DISTRICT

Likely impacts of weather warnings on Agriculture and associated Agromet advisories

In view of thunderstorms, lightening and squalls farmers are advised to harvest the matured fruits and vegetables and take care of strong staking in banana gardens.

GENERAL ADVISORY

- Farmers who have not yet sown groundnut crop, still have time to sow until the end of this month. Sowing should be taken up only after receipt of sufficient rainfall and enough moisture is available in the soil.

SMS ADVISORY

- In view of scattered and very light rainfall and high temperature, farmers are advised to monitor and irrigate the crops if necessary

CROP SPECIFIC ADVISORY **Select category: crop** **Select Type: G.Nut**

- Farmers who have not yet sown groundnut crop, still have time to sow until the end of this month. Sowing should be taken up only after receipt of sufficient rainfall and enough moisture is available in the soil. Before sowing, treat the seed with Imidachloprid 600 F.S @ 2 ml. + 4 ml water per kg of seed followed by Mancozeb 3 g/kg or Tebuconazole @ 1 g/kg of seed to control sucking pest and other seed borne diseases.

- Farmers who have sown groundnut during June month and is at flowering stage are advised to provide one irrigation to prevent moisture stress conditions.

Select category: **Select Type: Maize** **Select variety:**

- The present weather conditions are favourable for fall army worm in Maize crop under irrigated conditions at vegetative stage. To control spray Neem oil @ 5ml/L of water in early vegetative stage crop.
- If the maize crop is at 30 DAS, spray Larvin @ 1 g/L of water.

CROP SPECIFIC ADVISORY **Select category: crop** **Select Type: Paddy**

- The farmers who have taken up paddy nurseries are advised to apply carbofuran 3g granules @ 160 g. per cent of nursery 1 week prior to pulling of the nursery.

Farmers who are taking up transplanting of paddy nurseries are advised to

- To avoid attack of stem borer and leaf folder clip the tips of the leaves before transplanting
- To avoid BPH attack leave 20cm. spacing between the rows
- Apply entire 'P2O5' & 'K2O' as basal and 1/3 'N' as basal during last puddling .
- Transplant 30-35 days old rice seedlings.
- Leave 20 cm pathways for every 2 mt. of transplanted crop.

Select category: **Select Type: Minor millets** **Select variety:**

- Farmers who are taking up ragi, korra (minor millets) sowings are advised to treat the seed compulsorily with carbendazim @ 2 g. or mancozeb @ 3 g./kg of seed.

Select category: **Select Type: Cotton** **Stage:**

- The present weather conditions are favourable for sucking pests in cotton. To control spray fipronil @ 2 ml./L (or) profenophos @ 2 ml./L of water.

Likely impacts of weather warnings (General)

- Due to the possibility of heavy winds, livestock and goats should be kept in safe areas and provided with fodder.

Impact based advisories (General)

- In view of heavy winds stake the plants in banana gardens.

AGRO METEOROLOGICAL ADVISORY FOR ANNAMAYYA DISTRICT

Likely impacts of weather warnings on Agriculture and associated Agromet advisories

In view of thunderstorms, lightening and squalls farmers are advised to harvest the matured fruits and vegetables and take care of strong staking in banana gardens.

GENERAL ADVISORY

- Farmers who have not sown groundnut crop, still have time to sow until the end of this month. Sowing should be taken up only after receipt of sufficient rainfall and enough moisture is available in the soil.

SMS ADVISORY

- In view of scattered and very light rainfall and high temperature, farmers are advised to monitor and irrigate the crops if necessary

CROP SPECIFIC ADVISORY **Select category: crop** **Select Type: Paddy**

- The farmers who have taken up paddy nurseries are advised to apply carbofuran 3g granules @ 160 g. per cent of nursery 1 week prior to pulling of the nursery.

Farmers who are taking up transplanting of paddy nurseries are advised to

- To avoid attack of stem borer and leaf folder clip the tips of the leaves before transplanting
- To avoid BPH attack leave 20cm. spacing between the rows
- Apply entire 'P2O5' & 'K2O' as basal and 1/3 'N' as basal during last puddling.
- Transplant 30-35 days old rice seedlings.
- Leave 20 cm pathways for every 2 mt. of transplanted crop.

CROP SPECIFIC ADVISORY**Select category: crop****Select Type: G.Nut**

- Farmers who have not yet sown groundnut crop, still have time to sow until the end of this month. Sowing should be taken up only after receipt of sufficient rainfall and enough moisture is available in the soil. Before sowing, treat the seed with Imidachloprid 600 F.S @ 2 ml. + 4 ml water per kg of seed followed by Mancozeb 3 g/kg or Tebuconazole @ 1 g/kg of seed to control sucking pest and other seed borne diseases.
- Farmers who have sown groundnut during June month and is at flowering stage are advised to provide one irrigation to prevent moisture stress conditions.

Select category:**Select Type: Maize****Select variety:**

- The present weather conditions are favourable for fall army worm in Maize crop under irrigated conditions at vegetative stage. To control spray Neem oil @ 5ml/L of water in early vegetative stage crop.
- If the maize crop is at 30 DAS, spray Larvin @ 1 g/L of water.

Select category: Crop**Select Type: Black gram****Select variety:**

- The present weather conditions are favourable for sucking pest in black gram. To control spray Imidachloprid @ 0.4ml/L + 19-19-19 @ 10g L of water
- Farmers who are taking up sowing of blackgram and green gram under irrigations are advised to follow seed treatment with 2.5 g thiram / kaptan / mancozeb / carbendazim and imidachloprid @ 5 ml./kg of seed.

Select category:**Select Type: Minor millets****Select variety:**

- Farmers who are taking up ragi, korra (minor millets) sowings are advised to treat the seed compulsorily with carbendazim @ 2 g. or mancozeb @ 3 g./kg of seed.

Select category:**Select Type: Tomato****Select variety:**

- The present weather conditions are favourable for pin worm in tomato. Erect pheromone traps @ 8/ac. at 3-6 feet from ground level. If the intensity of the pest is more, spray flubendamide @ 0.25 ml/L of water.

Likely impacts of weather warnings (General)

- Due to the possibility of heavy winds, livestock and goats should be kept in safe areas and provided with fodder.

Impact based advisories (General)

- In view of heavy winds stake the plants in banana gardens.

AMFU ANANTHAPURAMU**AGRO METEOROLOGICAL ADVISORY FOR ANANTHAPURAMU & SRI SATHYASAI DISTRICTS****Likely impacts of weather warnings on Agriculture and associated Agromet advisories**

- Provide mechanical support and propping to horticultural crops and staking to vegetables to avoid lodging due to strong winds under your direct control.

General Advisory:

- Farmers are advised to take up sowings of rainfed crops viz., Groundnut, Redgram, Castor, Cotton and Bajra under sufficient moisture conditions.

SMS Advisory:

- Farmers are advised to take up seed treatment for rainfed crops to avoid the initial stages of pest and diseases.

Crop Specific Advisory

Crop (Varieties)	Crop Specific Advisory
Groundnut	• Farmers are advised to select high yielding varieties viz., Kadiri-6, Kadiri Harithandhra, Kadiri lepakshi and TCGS-1694 and also advised to follow seed treatment with imidacloprid 600 FS @ 2 ml + 4 ml of water/kg seed followed by tebuconazole 2 DS @ 1g or mancozeb @ 3 g/kg and

	Trichoderma viride @ 8 g/kg seed. - Farmers are advised to sow 4-6 rows of bajra/jowar as border crops all around the field to avoid transmission of vectors like thrips which spread PSND/PBND diseases. - Farmers are advised to adopt intercropping with redgram at 8:1/11:1/15:1 ratio as per their convenience and availability of sowing equipment. - Formation of conservation furrows every 3.6 m interval at sowing/20-30 days after sowing will help in soil moisture conservation. - Early-sown groundnut is at the vegetative stage. As dry weather conditions are prevailing, farmers are advised to spray 19:19:19 @ 5 g per litre of water as a foliar spray to help the crop withstand moisture stress and minimize the adverse effects of drought.
Redgram	- Farmers are advised to take up sowing only after receiving adequate rainfall. Farmers are advised to select medium duration varieties viz., PRG-176, LRG 52 and take up sowing windows up to 31st July. Farmers are also suggested to treat the seeds with rhizobium @10 g /kg seed. - For early sown redgram, Farmers are advised to take up intercultivation at 20 to 30 days after sowing to maintain weed free condition and also take up thinning. - Early sown redgram is at vegetative stage. As dry weather conditions are prevailing, farmers are advised to spray 19:19:19 @ 5 g per litre of water as a foliar spray to help the crop withstand moisture stress and minimize the adverse effects of drought.
Castor	- Under sufficient moisture condition, farmers are advised to take up sowing up to July 2nd fortnight. - Farmers are advised to select high yielding varieties such as Haritha, Kiran, Jwala, Pragathi and hybrids i.e DCH. 117, PCH-111, GCH.8, ICH.66 and ICH 5. At the time of sowing apply recommended dose of fertilizers (12 kg of nitrogen, 16 kg of phosphorus, and 12 kg of potassium per acre). - Early sown Castor crop is at vegetative stage. whenever there is soil moisture advised to take up tinning to maintain optimum plant population. As dry weather conditions are prevailing, farmers are advised to spray 19:19:19 @ 5 g per litre of water as a foliar spray to help the crop withstand moisture stress and minimize the adverse effects of drought.
Cotton	The cotton crop is at the vegetative to flowering stage. As dry weather conditions are prevailing, farmers are advised to spray NPK 19:19:19 @ 5 g per litre of water as a foliar spray during the vegetative stage. At the flowering stage, spray Potassium Nitrate (13:0:45) @ 5 g per litre of water as a foliar spray to help the crop withstand moisture stress and minimize the adverse effects of drought.
Bajra	Farmers are advised to take up sowing of bajra crop up to July. Take up seed treatment with Thiram or Captan @ 3 g/kg of seed, seed rate is 1.6 kg/acre and to follow spacing 45 cm x 15 cm to maintain optimum plant population. Farmers are advised to select the varieties viz., ABV 04, Dhanalakshmi, HHB -67, I.C.M.H -356, Raj 171 and Pusa 61 and PSV 56. Before sowing soak the seed in salt water for 10 minutes to remove ergot if present and then treat the seed with metalaxil @ 6 g/kg seed.
Mango	Farmers are advised to take up pruning after harvesting in mango orchards, dry twigs and irregular branches should be pruned to ensure good air circulation and sunlight penetration. After receiving good amount of rainfall, take up sowing of green manure crops in between fruit trees.
Banana	Farmers are advised to utilize the available soil moisture and apply recommended dose of fertilizers at 45, 90, 135 and 180 days for long duration varieties and 40, 80, 120 and 160 for short duration varieties after planting in banana.
Livestock	- Feed a balanced diet consisting of green fodder, dry fodder, and concentrates to maintain good health and optimum productivity. - Keep animal sheds clean, dry, and well-drained, avoiding water stagnation to minimize the risk of disease outbreaks during the monsoon season.
General advice	Farmers using smartphones can download the Vyavasaya Vathavaranam and Meghdoot mobile app for free from Google Play Store to get timely weather forecast and district wise agro advisories.

AGRO METEOROLOGICAL ADVISORY FOR KURNOOL & NANDYAL DISTRICTS

Likely impacts of weather warnings on Agriculture and associated Agromet advisories

- Provide mechanical support and propping to horticultural crops and staking to vegetables to avoid lodging due to strong winds.

General Advisory:

- Farmers are advised to take up sowings of rainfed crops viz., Groundnut, Redgram, Castor, Cotton and Bajra under sufficient moisture conditions.

SMS Advisory:

- Farmers are advised to take up seed treatment for rainfed crops to avoid the initial stages of pest and diseases.

Crop Specific Advisory


Crop (Varieties)	Crop Specific Advisory
Groundnut	• Farmers are advised to select high yielding varieties viz., Kadiri-6, Kadiri-9, Kadiri Harithandhra, Anantha, Kadiri Iepakshi and TCGS-1694 and also advised to follow seed treatment with imidacloprid 600 FS @ 2 ml + 4 ml of water/kg seed followed by tebuconazole 2 DS @ 1g or mancozeb @ 3 g/kg and Trichoderma viride @ 8 g/kg seed. • Farmers are advised to sow 4-6 rows of bajra/jowar as border crops all around the field to avoid transmission of vectors like thrips which spread PSND/PBND diseases. • Farmers are advised to adopt intercropping with redgram at 8:1/11:1/15:1 ratio as per their convenience and availability of sowing equipment. • Formation of conservation furrows every 3.6 m interval at sowing/20-30 days after sowing will help in soil moisture conservation. • Early-sown groundnut is at the vegetative stage. As dry weather conditions are prevailing, farmers are advised to spray 19:19:19 @ 5 g per litre of water as a foliar spray to help the crop to minimize the adverse effects of drought.
Redgram	• Farmers are advised to take up sowing only after receiving adequate rainfall. Farmers are advised to select medium duration varieties viz., PRG-176, LRG 52 and take up sowing windows up to 31st July. Farmers are also suggested to treat the seeds with rhizobium @10 g /kg seed. • For early sown redgram, Farmers are advised to take up intercultivation at 20 to 30 days after sowing to maintain weed free condition and also take up thinning. • Early sown redgram is at vegetative stage. As dry weather conditions are prevailing, farmers are advised to spray 19:19:19 @ 5 g per litre of water as a foliar spray to help the crop withstand moisture stress and minimize the adverse effects of drought.
Castor	• Under sufficient moisture condition, farmers are advised to take up sowing up to July 2nd fortnight. • Farmers are advised to select high yielding varieties such as Haritha, Kiran, Jwala, Pragathi and hybrids i.e DCH. 117, PCH-111, GCH.8, ICH.66 and ICH 5. At the time of sowing apply recommended dose of fertilizers (12 kg of nitrogen, 16 kg of phosphorus, and 12 kg of potassium per acre). • Early sown Castor crop is at vegetative stage. whenever there is soil moisture advised to take up tinning to maintain optimum plant population. As dry weather conditions are prevailing, farmers are advised to spray 19:19:19 @ 5 g per litre of water as a foliar spray to help the crop to minimize the adverse effects of drought.
Cotton	• The cotton crop is at the vegetative to flowering stage. As dry weather conditions are prevailing, farmers are advised to spray NPK 19:19:19 @ 5 g per litre of water as a foliar spray during the vegetative stage. At the flowering stage, spray Potassium Nitrate (13:0:45) @ 5 g per litre of water as a foliar spray to help the crop withstand moisture stress and minimize the adverse effects of drought.
Bajra	• Farmers are advised to take up sowing of bajra crop up to July. Take up seed treatment with Thiram or Captan @ 3 g/kg of seed, seed rate is 1.6 kg/acre and to follow spacing 45 cm x 15 cm to maintain optimum plant population. Farmers are advised to select the varieties

	viz., ABV 04, Dhanalkshmi, HHB -67, I.C.M.H -356, Raj 171 and Pusa 61 and PSV 56. Before sowing soak the seed in salt water for 10 minutes to remove ergot if present and then treat the seed with metalaxil @ 6 g/kg seed.
Mango	Farmers are advised to takeup pruning after harvesting in mango orchards, dry twigs and irregular branches should be pruned to ensure good air circulation and sunlight penetration. After receiving good amount of rainfall, take up sowing of green manure crops in between fruit trees.
Banana	Farmers are advised to utilize the available soil moisture and apply recommended dose of fertilizers at 45, 90, 135 and 180 days for long duration varieties and 40, 80, 120 and 160 for short duration varieties after planting in banana
Livestock	- Feed a balanced diet consisting of green fodder, dry fodder, and concentrates to maintain good health and optimum productivity. - Keep animal sheds clean, dry, and well-drained, avoiding water stagnation to minimize the risk of disease outbreaks during the monsoon season.
General advice	Farmers using smartphones can download the Vyavasaya Vathavaranam and Meghdoot mobile app for free from Google Play Store to get timely weather forecast and district wise agro advisories.

Annexure - I

District-level Weather warnings issued by MC Amaravati

Weather warning maps for next 5 days:




Weather warnings for Andhra Pradesh and Yanam.

Valid from 0830 hrs IST of 11.07.2026 to 0830 hrs IST of 12.07.2026



Date of Issue:- 10 July 2026

Day 2- 11 July 2026




Weather warnings for Andhra Pradesh and Yanam.

Valid from 0830 hrs IST of 12.07.2026 to 0830 hrs IST of 13.07.2026



Date of Issue:- 10 July 2026

Day 3- 12 July 2026




Weather warnings for Andhra Pradesh and Yanam.

Valid from 0830 hrs IST of 13.07.2026 to 0830 hrs IST of 14.07.2026



Date of Issue:- 10 July 2026

Day 4 - 13 July 2026




Weather warnings for Andhra Pradesh and Yanam.

Valid from 0830 hrs IST of 14.07.2026 to 0830 hrs IST of 15.07.2026



Date of Issue:- 10 July 2026

Day 5 - 14 July 2026



Annexure - II

Meteorological Subdivision Level Rainfall Weather Forecast for next 5 days (maps):

-No Warnings-

District Forecast & Warnings for next 5 days:

DISTRICT LEVEL FORECAST FOR ANDHRA PRADESH & YANAM						
METEOROLOGICAL SUB-DIVISION	NAME OF THE DISTRICTS / जिलो का नाम	WEATHER FORECAST FOR NEXT 5 DAYS				
		DAY -1 VALID FROM 1300 HRS OF 10-Jul-2026 TO 0830 HOURS IST OF 11-Jul-2026	DAY -2 VALID FROM 0830 HRS OF 11-Jul-2026 TO 0830 HOURS IST OF 12-Jul-2026	DAY -3 VALID FROM 0830 HRS OF 12-Jul-2026 TO 0830 HOURS IST OF 13-Jul-2026	DAY -4 VALID FROM 0830 HRS OF 13-Jul-2026 TO 0830 HOURS IST OF 14-Jul-2026	DAY -5 VALID FROM 0830 HRS OF 14-Jul-2026 TO 0830 HOURS IST OF 15-Jul-2026
NORTH COASTAL ANDHRA PRADESH & YANAM	Srikakulam	SCATTERED	ISOLATED	ISOLATED	ISOLATED	ISOLATED
		Light to Moderate Rain/TS	Light to Moderate Rain	Light to Moderate Rain	Light to Moderate Rain	Light to Moderate Rain
		likely	likely	likely	likely	likely
	Parvathipuram Manyam	SCATTERED	ISOLATED	ISOLATED	ISOLATED	ISOLATED
		Light to Moderate Rain/TS	Light to Moderate Rain	Light to Moderate Rain	Light to Moderate Rain	Light to Moderate Rain
		likely	likely	likely	likely	likely
	Vizianagaram	SCATTERED	ISOLATED	ISOLATED	ISOLATED	ISOLATED
		Light to Moderate Rain/TS	Light to Moderate Rain	Light to Moderate Rain	Light to Moderate Rain	Light to Moderate Rain
		likely	likely	likely	likely	likely
	Visakhapatnam	ISOLATED	ISOLATED	ISOLATED	ISOLATED	ISOLATED
		Light to Moderate Rain/TS	Light Rain	Light Rain	Light to Moderate Rain	Light to Moderate Rain
		likely	likely	likely	likely	likely
	Sri Alluri Sitharamaraju District	SCATTERED	ISOLATED	ISOLATED	ISOLATED	ISOLATED
		Light to Moderate Rain/TS	Light to Moderate Rain	Light to Moderate Rain	Light to Moderate Rain	Light to Moderate Rain
		likely	likely	likely	likely	likely
	Anakapalli	SCATTERED	ISOLATED	ISOLATED	ISOLATED	ISOLATED
		Light to Moderate Rain/TS	Light Rain	Light to Moderate Rain	Light to Moderate Rain	Light to Moderate Rain
		likely	likely	likely	likely	likely
	Kakinada	ISOLATED	ISOLATED	ISOLATED	ISOLATED	ISOLATED
		Light to Moderate Rain/TS	Light Rain	Light to Moderate Rain	Light to Moderate Rain	Light to Moderate Rain
		likely	likely	likely	likely	likely
	East Godavari	SCATTERED	ISOLATED	ISOLATED	ISOLATED	ISOLATED
		Light to Moderate Rain/TS	Light Rain	Light to Moderate Rain	Light to Moderate Rain	Light to Moderate Rain
		likely	likely	likely	likely	likely
	Yanam	ISOLATED	ISOLATED	ISOLATED	ISOLATED	ISOLATED
		Light to Moderate Rain/TS	Light Rain	Light to Moderate Rain	Light to Moderate Rain	Light to Moderate Rain
		likely	likely	likely	likely	likely
	Dr B.R.Ambedkar Konaseema	ISOLATED	ISOLATED	ISOLATED	ISOLATED	ISOLATED
		Light to Moderate Rain/TS	Light Rain	Light to Moderate Rain	Light to Moderate Rain	Light to Moderate Rain
		likely	likely	likely	likely	likely
	Eluru	SCATTERED	ISOLATED	ISOLATED	ISOLATED	ISOLATED
		Light to Moderate Rain/TS	Light to Moderate Rain	Light to Moderate Rain	Light to Moderate Rain	Light to Moderate Rain
		likely	likely	likely	likely	likely
	West Godavari	ISOLATED	ISOLATED	ISOLATED	ISOLATED	ISOLATED
		Light to Moderate Rain/TS	Light to Moderate Rain	Light to Moderate Rain	Light to Moderate Rain	Light to Moderate Rain
		likely	likely	likely	likely	likely

SOUTH COASTAL ANDHRA PRADESH	NTR District	ISOLATED	ISOLATED	DRY	DRY	DRY
		Light to Moderate Rain	Light to Moderate Rain			
		likely	likely			
	Krishna	ISOLATED	ISOLATED	DRY	DRY	DRY
		Light to Moderate Rain	Light to Moderate Rain			
		likely	likely			
	Palnadu	ISOLATED	ISOLATED	DRY	DRY	DRY
		Light to Moderate Rain	Light Rain			
		likely	likely			
	Guntur	ISOLATED	ISOLATED	DRY	DRY	DRY
		Light to Moderate Rain	Light Rain			
		likely	likely			
	Bapatla	ISOLATED	ISOLATED	DRY	DRY	DRY
		Light Rain	Light to Moderate Rain			
		likely	likely			
	Prakasam	ISOLATED	ISOLATED	DRY	DRY	DRY
		Light Rain	Light Rain			
		likely	likely			
	SPSR Nellore	ISOLATED	ISOLATED	DRY	DRY	DRY
		Light to Moderate Rain	Light to Moderate Rain			
		likely	likely			
RAYALASEEMA	Kurnool	ISOLATED	ISOLATED	DRY	DRY	DRY
		Light Rain/TS	Light to Moderate Rain			
		likely	likely			
	Nandyal	ISOLATED	ISOLATED	DRY	DRY	DRY
		Light Rain/TS	Light Rain			
		likely	likely			
	Anantapuramu	ISOLATED	ISOLATED	DRY	DRY	DRY
		Light Rain/TS	Light to Moderate Rain			
		likely	likely			
	Sri Satyasai District	ISOLATED	ISOLATED	DRY	DRY	DRY
		Light to Moderate Rain/TS	Light to Moderate Rain			
		likely	likely			
	YSR District	ISOLATED	ISOLATED	DRY	DRY	DRY
		Light to Moderate Rain/TS	Light Rain			
		likely	likely			
	Annamayya	ISOLATED	ISOLATED	DRY	DRY	DRY
		Light to Moderate Rain/TS	Light Rain			
		likely	likely			
	Tirupathi	ISOLATED	ISOLATED	DRY	DRY	DRY
		Light to Moderate Rain/TS	Light Rain			
		likely	likely			
	Chittoor	ISOLATED	ISOLATED	DRY	DRY	DRY
		Light to Moderate Rain/TS	Light to Moderate Rain			
		likely	likely			
REMARKS		Maximum temperatures are No Large Change over Andhrapradesh&Yanam for D1 to D5				

ISOL: Isolated i.e. Rain at 1 or 2 places	FWS: Fairly Widespread i.e. Rain at many places
SCT: Scattered i.e. Rain at a few places	WS: Widespread i.e. Rain at most places
DRY: No rain	NSC: No Significant Change
NCAP: North Coastal Andhra Pradesh	SCAP: South Coastal Andhra Pradesh

FWS: Fairly Widespread i.e. Rain at many places
WS: Widespread i.e. Rain at most places
NSC: No Significant Change

NCAP: North Coastal Andhra Pradesh

SCAP: South Coastal Andhra Pradesh

LEGEND:					
Spatial Distribution of Rainfall	Descriptive term used	Criteria for observed/ forecast weather	Probability	Intensity of Rainfall	
Dry	Dry	No Station reported/ expecting rainfall		Very Light rainfall	Trace to 2.4 mm
Isolated	One or two places	≤25% of Stations get/ expected to get rainfall	Unlikely	Light rainfall	2.5 to 15.5 mm
Scattered	At a Few places	(26% to 50%) of Stations get/ expected to get rainfall	Likely	Moderate rainfall	15.6 to 64.4 mm
Fairly Widespread	At Many places	(51% to 75%) of Stations get/ expected to get rainfall	Very likely	Heavy Rainfall	64.5 to 115.5 mm
Wide spread	At Most places	76% to 100%) of Stations get/ expected to get rainfall	Most likely	Very Heavy Rainfall	115.6 to 204.4 mm
				Extremely Heavy Rainfall	204.5 mm or more
		Weather		<u>WARNINGS</u>	
		RA	Rain	Warning - Take action	
		TS	Thunder shower	Alert - Be Prepared	
		NLC	No Large Change	Watch - Be Updated	
				No Warning - No Action	



For Weather Forecast, Warnings & Inference Kindly refer the link below:

<https://mausam.imd.gov.in/amaravati/mcdata/apinf.pdf>

https://mausam.imd.gov.in/visakhapatnam/mcdata/heavyrain_warning.pdf

<https://mausam.imd.gov.in/amaravati/mcdata/APvaf.pdf>

<https://mausam.imd.gov.in/amaravati/mcdata/APAgromet.pdf>

For IMD AMARAVATI WEBSITE, INSTAGRAM, TWITTER, FACEBOOK, YOU TUBE links

	Instagram	https://www.instagram.com/mc_amaravati/
	Twitter	https://twitter.com/AmaravatiMc
	Facebook	https://www.facebook.com/MC-Amaravati-IMD-100787794998444/
	YouTube	https://www.youtube.com/channel/UCMhKw9DfBAIcfpacLYVppCw

Android apps: Mausam- weather forecast; Damini- lightning; Meghdoot- agriculture forecast.