REPUBLIC OF KENYA

KNBS
KENYA NATIONAL BUREAU OF STATISTICS
Keeping you informed

2025-26 Kenya Mini DEMOGRAPHIC and Health Survey



October 2026

REPUBLIC OF KENYA

KNBS
KENYA NATIONAL BUREAU OF STATISTICS
Keeping you informed

Kenya
Mini Demographic and Health Survey 2025–26

Kenya National Bureau of Statistics Nairobi, Kenya
Ministry of Health Nairobi, Kenya

The DHS Program
ICF
Rockville, Maryland, USA

October 2026

Gates Foundation

CONRAD N. Hilton FOUNDATION

ICF

THE AGA KHAN UNIVERSITY
INSTITUTE FOR HUMAN DEVELOPMENT

unicef
for every child

World Health Organization

UNFPA

UN WOMEN

FOR ALL WOMEN AND GIRLS

NATIONAL COUNCIL FOR POPULATION AND DEVELOPMENT
NCPD
MANAGING POPULATION FOR PROSPERITY

The 2025–26 Kenya Mini Demographic and Health Survey (KMDHS) was implemented by the Kenya National Bureau of Statistics (KNBS) in collaboration with the Ministry of Health (MoH) and other stakeholders. Funding for the survey was provided by the Government of Kenya, the Gates Foundation, the Conrad N. Hilton Foundation, the Aga Khan University Institute for Human Development, the United Nations Children's Fund (UNICEF), the United Nations Population Fund (UNFPA), and the World Health Organization (WHO). ICF provided technical assistance through the DHS Program.

Additional information about the 2025–26 KMDHS may be obtained from Kenya National Bureau of Statistics (KNBS), P.O. Box 30266-00100, GPO Nairobi, Kenya; +254-20-3317583, +254-20-2911000/1, +254-20-3317612/22/23/51; directorgeneral@knbs. or.ke, info@knbs.or.ke; www.knbs.or.ke. Information about The DHS Program may be obtained from ICF, 530 Gaither Road, Suite 500, Rockville, MD 20850, USA; +1-301-407-6500; fax: +1-301-407-6501; info@DHSprogram.com; www.DHSprogram.com. The contents of this report are the sole responsibility of KNBS and ICF and do not necessarily reflect the views of the donors.

ISBN: 978-9914-9428-7-3
DOI: https://doi.org/10.63814/k7m3xq21

Published 2026



Some rights reserved.

This work is made available under the Attribution-Noncommercial 4.0 International (CC BY-NC 4.0) available at https:/creativecommons. org/licenses/by-nc/4.0/ This license allows reusers to distribute, remix, adapt, and build upon the material in any medium or format for noncommercial purposes only so long as the work is appropriately cited and attribution is given to the creator.

The contents of this report are the sole responsibility of KNBS and ICF and do not necessarily reflect the views of the United States Government, or other donor agencies.

Image source: All images and visual elements in this publication were generated with the assistance of AI image generation tools under the author's creative direction. The images are original outputs generated for this work and any familiar resemblance is completely coincidental.

Recommended citation:

KNBS and ICF. 2026. *2025–26 Kenya Mini Demographic and Health Survey*. Nairobi, Kenya, and Rockville, Maryland, USA: KNBS and ICF.

CONTENTS

TABLES AND FIGURES ..... **ix**
PREFACE ..... **xxi**
ACRONYMS AND ABBREVIATIONS ..... **xxiii**
READING AND UNDERSTANDING TABLES FROM THE 2025–26 KENYA MINI DEMOGRAPHIC AND HEALTH SURVEY (KMDHS) ..... **xxvii**
SUSTAINABLE DEVELOPMENT GOAL INDICATORS ..... **xxxv**
MAP OF KENYA ..... **xxxviii**

1 INTRODUCTION AND SURVEY METHODOLOGY **1**
1.1 Survey Objectives 1
1.2 Sample Design 1
1.3 Questionnaires 2
1.4 Training of Trainers and Pretest 3
1.4.1 Training of Trainers 3
1.4.2 Pretest 3
1.5 Training of Field Staff 4
1.6 Fieldwork 4
1.7 Data Processing 5
1.8 Response Rates 5

2 HOUSING CHARACTERISTICS AND HOUSEHOLD POPULATION **7**
2.1 Housing Characteristics 7
2.1.1 Use of Clean Fuels and Technologies 8
2.1.2 Cooking 8
2.1.3 Heating and Lighting 8
2.1.4 Primary Reliance on Clean Fuels and Technologies 8
2.2 Household Wealth 10
2.2.1 Household Durable Goods 10
2.2.2 Wealth Index 10
2.3 Household Characteristics and Composition 11
2.4 Birth Registration 11
2.5 Educational Attainment 14
2.6 Health Insurance 14
2.6.1 Coverage 14
2.6.2 Status of Premium Payments under the Social Health Insurance Fund 15
2.7 Household Enterprises 16
2.7.1 Registration Status 16
2.7.2 Type of Economic Activity and Average Monthly Revenue 16
2.8 Household Member Deaths 16
2.8.1 Place of Death 16
2.8.2 Issuance of Death Certificates and Burial Permits 17
2.8.3 Reasons for Non-issuance of Burial Permits 19

3 CHARACTERISTICS OF RESPONDENTS 51
3.1 Basic Characteristics of Survey Respondents 52
3.2 Education and Literacy 52
3.3 Mass Media Exposure and Internet Usage 54
3.3.1 Mass Media Exposure 54
3.3.2 Internet Usage 55
3.4 Place of Birth and Recent Migration 56
3.4.1 Type of Migration 56
3.4.2 Reason for Migration 56
3.5 Social Health Authority (SHA) Coverage and Utilisation 57
3.5.1 Coverage: Social Health Authority (SHA) Registration 57
3.5.2 Reasons for Not Registering 57
3.5.3 Utilisation of Services 58
3.5.4 Services Accessed 58
3.5.5 Challenges Accessing Services 58
3.5.6 Benefits Received 59
3.5.7 Satisfaction with Services 59

4 MARRIAGE AND SEXUAL ACTIVITY 89
4.1 Marital Status 89
4.2 Marriage Registration 90
4.3 Age at First Sexual Intercourse 90
4.4 Recent Sexual Activity 91
4.5 Multiple Sexual Partners 92

5 FERTILITY 101
5.1 Current Fertility 101
5.2 Children Ever Born and Living 103
5.3 Birth Intervals 103
5.4 Insusceptibility to Pregnancy 104
5.5 Age at First Menstruation 105
5.6 Menopause 106
5.7 Age at First Birth 106
5.8 Teenage Pregnancy 106
5.9 Pregnancy Outcomes and Induced Abortion 108

6 FERTILITY PREFERENCES 125
6.1 Desire for Another Child 125
6.2 Ideal Family Size 126
6.3 Fertility Planning Status 127
6.4 Wanted Fertility Rates 128

7 FAMILY PLANNING 135
7.1 Contraceptive Knowledge and Use 135
7.1.1 Use of Emergency Contraception 138
7.1.2 Knowledge of the Fertile Period 139
7.2 Source of Modern Contraceptive Methods 139
7.3 Informed Choice 140
7.4 Discontinuation of Contraceptives 140

7.5 Demand for Family Planning 141
7.6 Future Use of Contraception 144
7.7 Exposure to Family Planning Messages 144
7.8 Contact of Non-users with Family Planning Providers 144

8 INFANT AND CHILD MORTALITY 169
8.1 Infant and Child Mortality 170
8.2 Perinatal Mortality 171
8.3 High-risk Fertility Behaviour 172

9 MATERNAL AND NEWBORN HEALTH CARE 177
9.1 Antenatal Care Coverage 178
9.1.1 Skilled Providers 178
9.1.2 Timing and Number of Antenatal Care Visits 178
9.2 Components of Antenatal Care 179
9.2.1 Deworming and Iron-containing Supplementation During Pregnancy 181
9.2.2 Source of Iron-containing Supplements 182
9.3 Protection against Neonatal Tetanus 182
9.4 Delivery Services 183
9.4.1 Institutional Deliveries 183
9.4.2 Delivery by Caesarean Section 184
9.4.3 Skilled Assistance during Delivery 186
9.5 Postnatal Care 188
9.5.1 Postnatal Health Check for Mothers 188
9.5.2 Postnatal Health Check for Newborns 189
9.5.3 Postnatal Health Checks for Mothers and Newborns 190
9.6 Breast and Cervical Cancer Examinations 191
9.7 Problems in Accessing Health Care 193
9.8 Distance and Means of Transport to the Nearest Health Facility 194
9.9 Vaccination against Human Papillomavirus 194

10 CHILD HEALTH 243
10.1 Child's Size and Birth Weight 243
10.2 Vaccination of Children 244
10.2.1 Vaccination Card Ownership and Availability 244
10.2.2 Basic Antigen Coverage 245
10.2.3 National Schedule Coverage 246
10.3 Symptoms of Acute Respiratory Infection and Care-seeking Behaviour 248
10.3.1 Treatment for Symptoms 249
10.3.2 Source of Advice or Treatment 249
10.4 Fever and Care-seeking Behaviour 249
10.5 Diarrhoeal Disease 250
10.5.1 Care-seeking Behaviour 250
10.5.2 Feeding Practices 250
10.5.3 Oral Rehydration Therapy, Zinc, Continued Feeding, and Other Treatments 251
10.6 Childhood Illness and Care-seeking Behaviour 252
10.7 Early Childhood Development 252

11 NUTRITION OF CHILDREN 279
11.1 Growth Monitoring and Promotion 279
11.2 Infant and Young Child Feeding Practices 280
11.2.1 Ever Breastfed, Early Initiation of Breastfeeding, and Exclusive Breastfeeding for the First 2 Days After Birth 280
11.2.2 Exclusive Breastfeeding and Mixed Milk Feeding 281
11.2.3 Continued Breastfeeding and Bottle Feeding 283
11.2.4 Introduction of Complementary Foods 283
11.2.5 Minimum Dietary Diversity, Minimum Meal Frequency, Minimum Milk Feeding Frequency, Minimum Acceptable Diet, and Egg and/or Flesh Food Consumption 284
11.2.6 Sweet Beverage Consumption, Unhealthy Food Consumption, and Zero Vegetable or Fruit Consumption 286
11.2.7 Infant and Young Child Feeding Indicators 288
11.3 Infant and Young Child Feeding Counselling 288
11.4 Micronutrient Supplementation and Deworming 289

12 MALARIA 305
12.1 Ownership of Insecticide-treated Nets 307
12.2 Household Access to and Use of ITNs 310
12.3 Use of ITNs by Children and Pregnant Women 312
12.4 Reasons Mosquito Nets Were Not Used 313
12.5 Malaria in Pregnancy 313
12.6 Case Management of Malaria in Children 314

13 HOUSEHOLD WATER AND SANITATION 337
13.1 Drinking Water Sources, Availability, and Treatment 337
13.2 Sanitation 340
13.2.1 Sanitation Service Ladder 341
13.2.2 Removal and Disposal of Excreta 343
13.3 Disposal of Children's Stools 344
13.4 Handwashing 344
13.5 Menstrual Hygiene 345

14 GENDER-BASED VIOLENCE 361
14.1 Measurement of Technology-facilitated Gender-based Violence 363
14.2 Experience of Technology-facilitated Gender-based Violence by Any Perpetrator 364
14.2.1 Experience of Various Forms of Technology-facilitated Gender-based Violence by Any Perpetrator 366
14.2.2 Experience of Technology-facilitated Gender-based Violence by Any Intimate Partner 366
14.2.3 Perpetrators of Technology-facilitated Gender-Based Violence 367
14.3 Forms of Controlling Behaviours 367
14.4 Experience of Technology-facilitated Gender-based Violence by Non- intimate Partners According to Platform 369
14.5 Community Perceptions on Violence against Women and Girls 369
14.5.1 Perceived Risks of Violence and Safety for Women and Girls 370
14.5.2 Perceived Community Response to Killing of Women and Girls 371
14.5.3 Justification for Violence against Women or Girls 372
14.5.4 Perceptions of Victim Blaming and Privacy of Domestic Violence 372

15 ADULT AND MATERNAL MORTALITY 395
15.1 Data 395
15.2 Direct Estimates of Adult Mortality 396
15.3 Trends in Adult Mortality 397
15.4 Direct Estimates of Maternal Mortality 397
15.5 Trends in Pregnancy-related Mortality 399

REFERENCES 403

Appendix A SAMPLE DESIGN 407
A.1 Introduction 407
A.2 Sampling Frame 407
A.3 Sample Design and Implementation 410
A.4 Sample Probabilities and Sampling Weights 412
A.4 Survey Results 414

Appendix B ESTIMATES OF SAMPLING ERRORS 415

Appendix C DATA QUALITY TABLES 475

Appendix D PERSONS INVOLVED IN THE 2025–26 KENYA MINI DEMOGRAPHIC AND HEALTH SURVEY 485

Appendix E QUESTIONNAIRES 489

TABLES, FIGURES, AND MAPS

1 INTRODUCTION AND SURVEY METHODOLOGY 1

Table 1.1 Results of the household and individual interviews 6

2 HOUSING CHARACTERISTICS AND HOUSEHOLD POPULATION 7

Table 2.1 Household characteristics: Housing 21
Table 2.2 Household characteristics: Cooking 22
Table 2.3 Household characteristics: Heating and lighting 23
Table 2.4 Primary reliance on clean fuels and technologies 24
Table 2.4C Primary reliance on clean fuels and technologies: County level 25
Table 2.5 Household possessions 26
Table 2.6 Wealth quintiles 27
Table 2.7 Household population by age, sex, and residence 28
Table 2.8 Household composition 29
Table 2.9 Birth registration of children under age 5 29
Table 2.9C Birth registration of children under age 5: County level 30
Table 2.10.1 Educational attainment of the female household population (standard DHS education categories) 31
Table 2.10.1C Educational attainment of the female household population (standard DHS education categories): County level 32
Table 2.10.2 Educational attainment of the female household population (CBE) 33
Table 2.10.2C Educational attainment of the female household population (CBE): County level 34
Table 2.10.3 Educational attainment of the male household population (standard DHS education categories) 35
Table 2.10.3C Educational attainment of the male household population (standard DHS education categories): County level 36
Table 2.10.4 Educational attainment of the male household population (CBE) 37
Table 2.10.4C Educational attainment of the male household population (CBE): County level 38
Table 2.11 Health insurance coverage 39
Table 2.11C Health insurance coverage: County level 40
Table 2.12 Status of premium payments under the Social Health Insurance Fund 41
Table 2.12C Status of premium payments under the Social Health Insurance Fund: County level 42
Table 2.13 Households operating an income-generating business/enterprise 43
Table 2.14 Household enterprises by sex of operator 44
Table 2.15 Registration status of household enterprises 45
Table 2.16 Type of economic activity of household enterprises 46
Table 2.17 Average monthly revenue of household enterprises 47
Table 2.18 Household member deaths 48
Table 2.19 Death certificate and burial permit issuance 48
Table 2.19C Death certificate and burial permit issuance: County level 49
Table 2.20 Reasons for non-issuance of burial permits 50

Figure 2.1 Primary reliance on clean fuels and technologies 9
Figure 2.2 Household wealth by residence 10
Figure 2.3 Population pyramid 11
Figure 2.4 Birth registration by household wealth 12
Figure 2.5 Birth registration 13
Figure 2.6 Educational attainment of the household population 14

Map 2.1 Primary reliance on clean fuels and technologies by county 9
Map 2.2 Birth registration by county 13
Map 2.3 Health insurance coverage by county 15

3 CHARACTERISTICS OF RESPONDENTS 51
Table 3.1 Background characteristics of respondents 61
Table 3.1C Background characteristics of respondents: County level 62
Table 3.2 Education 63
Table 3.2C Education: County level 64
Table 3.3 Literacy 65
Table 3.3C Literacy: County level 66
Table 3.4 Exposure to mass media 67
Table 3.4C Exposure to mass media: County level 68
Table 3.5 Internet usage 69
Table 3.5C Internet usage: County level 70
Table 3.6 Place of birth and recent migration 71
Table 3.6C Place of birth and recent migration: County level 72
Table 3.7 Type of migration 73
Table 3.8 Reasons for migration 73
Table 3.8C Reasons for migration: County level 74
Table 3.9 Social Health Authority (SHA) registration 75
Table 3.9C Social Health Authority (SHA) registration: County level 76
Table 3.10 Reasons for not registering with the Social Health Authority (SHA) scheme 77
Table 3.10C Reasons for not registering with the Social Health Authority (SHA): County level 78
Table 3.11 Utilisation of Social Health Authority (SHA) services 79
Table 3.11C Utilisation of Social Health Authority (SHA) services: County level 80
Table 3.12 Services received through the Social Health Authority (SHA) 81
Table 3.12C Services received through the Social Health Authority (SHA): County level 82
Table 3.13 Challenges accessing Social Health Authority (SHA) services 83
Table 3.13C Challenges accessing Social Health Authority (SHA) Services: County level 84
Table 3.14 Benefits received through Social Health Authority (SHA) coverage 85
Table 3.14C Benefits received through Social Health Authority (SHA) coverage: County level 86
Table 3.15 Satisfaction with services covered under the Social Health Authority (SHA) 87
Table 3.15C Satisfaction with services covered under the Social Health Authority (SHA): County level 88

Figure 3.1 Education 52
Figure 3.2 Secondary education by household wealth 53
Figure 3.3 Exposure to mass media 54
Figure 3.4 Trends in exposure to mass media 54
Figure 3.5 Internet usage by residence 55

Map 3.1 Secondary education by county 53

4 MARRIAGE AND SEXUAL ACTIVITY 89
Table 4.1 Current marital status 93
Table 4.2 Marriage registration 93
Table 4.2C Marriage registration by county 94
Table 4.3 Age at first sexual intercourse 95
Table 4.4 Median age at first sexual intercourse according to background characteristics 95
Table 4.4C Median age at first sexual intercourse by county 96
Table 4.5 Recent sexual activity 97
Table 4.5C Recent sexual activity by county 98
Table 4.6 Multiple sexual partners and higher-risk sexual intercourse in the past 12 months among young women 99

Figure 4.1 Marital status 89
Figure 4.2 Trends in marital status 90
Figure 4.3 Trends in first sexual intercourse 91

5 FERTILITY 101
Table 5.1 Current fertility 110
Table 5.2 Fertility by background characteristics 110
Table 5.2C Fertility by background characteristics: County level 111
Table 5.3.1 Trends in age-specific fertility rates 112
Table 5.3.2 Trends in age-specific and total fertility rates 112
Table 5.4 Children ever born and living 113
Table 5.5 Birth intervals 114
Table 5.5C Birth intervals: County level 115
Table 5.6 Postpartum amenorrhoea, abstinence, and insusceptibility 116
Table 5.7 Median durations of postpartum amenorrhoea, abstinence, and insusceptibility 116
Table 5.8 Age at first menstruation 117
Table 5.9 Menopause 117
Table 5.10 Age at first birth 117
Table 5.11 Median age at first birth 118
Table 5.11C Median age at first birth: County level 119
Table 5.12 Teenage pregnancy 120
Table 5.12C Teenage pregnancy: County level 121
Table 5.13 Sexual and reproductive health behaviours before age 15 122
Table 5.14 Pregnancy outcomes by background characteristics 122
Table 5.14C Pregnancy outcomes by background characteristics: County level 123
Table 5.15 Induced abortion rates 124

Figure 5.1 Trends in age-specific fertility 102
Figure 5.2 Trends in fertility by residence 102
Figure 5.3 Fertility by education 102
Figure 5.4 Birth intervals 104
Figure 5.5 Pregnancy outcomes 108

Map 5.1 Fertility by county 103
Map 5.2 Teenage pregnancy by county 107

6 FERTILITY PREFERENCES 125
Table 6.1 Fertility preferences according to number of living children 130
Table 6.2 Desire to limit childbearing 130
Table 6.3 Ideal number of children according to number of living children 131
Table 6.4 Mean ideal number of children according to background characteristics 131
Table 6.4C Mean ideal number of children according to background characteristics: County level 132
Table 6.5 Fertility planning status 133
Table 6.6 Wanted fertility rates 133
Table 6.6C Wanted fertility rates: County level 134

Figure 6.1 Trends in desire to limit childbearing by number of living children 126
Figure 6.2 Ideal family size by number of living children 126
Figure 6.3 Fertility planning status 127
Figure 6.4 Trends in fertility planning status 127
Figure 6.5 Trends in wanted and actual fertility 128

Map 6.1 Wanted fertility rates by county 129

7 FAMILY PLANNING 135
Table 7.1 Knowledge of contraceptive methods 146
Table 7.2 Knowledge of contraceptive methods according to background characteristics 147
Table 7.2C Knowledge of contraceptive methods: County level 148
Table 7.3 Current use of contraception by age 149
Table 7.4.1 Trends in current use of contraception 150
Table 7.4.2 Current use of contraception according to background characteristics 151
Table 7.4.2C Current use of contraception: County level 152
Table 7.5 Timing of sterilisation 153
Table 7.6 Use of subcutaneous depot-medroxyprogesterone acetate/Sayana Press 153
Table 7.7 Use of emergency contraception 154
Table 7.7C Use of emergency contraception: County level 155
Table 7.8 Knowledge of fertile period 156
Table 7.9 Knowledge of fertile period by age 156
Table 7.10 Source of modern contraceptive methods 157
Table 7.11 Informed choice 158
Table 7.12 Twelve-month contraceptive discontinuation rates 159
Table 7.13 Reasons for discontinuation 159
Table 7.14.1 Need and demand for family planning among currently married women 160
Table 7.14.1C Need and demand for family planning among currently married women: County level 161

Table 7.14.2 Need and demand for family planning among all women and sexually active unmarried women ... 162
Table 7.14.2C Need and demand for family planning among all women: County level ... 163
Table 7.15 Future use of contraception ... 164
Table 7.16 Exposure to family planning messages ... 164
Table 7.16C Exposure to family planning messages: County level ... 165
Table 7.17 Contact of non-users with family planning providers ... 166
Table 7.17C Contact of non-users with family planning providers: County level ... 167

Figure 7.1 Contraceptive use ... 136
Figure 7.2 Trends in contraceptive use ... 137
Figure 7.3 Use of modern contraceptive methods by education ... 137
Figure 7.4 Source of modern contraceptive methods ... 139
Figure 7.5 Source of specific modern contraceptive methods ... 140
Figure 7.6 Demand for family planning ... 142
Figure 7.7 Trends in demand for family planning ... 142
Figure 7.8 Unmet need for family planning by education ... 143

Map 7.1 Modern contraceptive use by county ... 138
Map 7.2 Unmet need for family planning by county ... 143

8 INFANT AND CHILD MORTALITY ... 169
Table 8.1 Early childhood mortality rates ... 173
Table 8.2 Five-year early childhood mortality rates according to background characteristics 173
Table 8.3 Ten-year early childhood mortality rates according to additional characteristics ... 174
Table 8.4 Perinatal mortality ... 175
Table 8.5 High-risk fertility behaviour ... 176

Figure 8.1 Trends in early childhood mortality rates ... 170

9 MATERNAL AND NEWBORN HEALTH CARE ... 177
Table 9.1 Antenatal care ... 197
Table 9.1C Antenatal care: County level ... 198
Table 9.2 Number of antenatal care visits and timing of first visit ... 199
Table 9.2C Number of antenatal care visits and timing of first visit: County level ... 200
Table 9.3.1 Components of antenatal care among women receiving antenatal care ... 201
Table 9.3.1C Components of antenatal care among women receiving antenatal care: County level ... 202
Table 9.3.2 Components of antenatal care among all women ... 203
Table 9.3.2C Components of antenatal care among all women: County level ... 204
Table 9.4 Deworming and iron-containing supplementation during pregnancy ... 205
Table 9.4C Deworming and iron-containing supplementation during pregnancy: County level ... 206
Table 9.5 Source of iron-containing supplements ... 207
Table 9.6 Tetanus toxoid injections ... 208
Table 9.6C Tetanus toxoid injections: County level ... 209
Table 9.7 Place of delivery ... 210
Table 9.7C Place of delivery: County level ... 211
Table 9.8 Caesarean section ... 212

Table 9.8C Caesarean section: County level 213
Table 9.9 Assistance during delivery 214
Table 9.9C Assistance during delivery: County level 215
Table 9.10 Duration of stay in a health facility after birth 216
Table 9.11 Timing of first postnatal check for the mother 217
Table 9.11C Timing of first postnatal check for the mother: County level 218
Table 9.12 Type of provider of first postnatal check for the mother 219
Table 9.12C Type of provider of first postnatal check for the mother: County level 220
Table 9.13 Content of postnatal care for the mother 221
Table 9.13C Content of postnatal care for the mother: County level 222
Table 9.14 Timing of first postnatal check for the newborn 223
Table 9.14C Timing of first postnatal check for the newborn: County level 224
Table 9.15 Type of provider of first postnatal check for the newborn 225
Table 9.15C Type of provider of first postnatal check for the newborn: County level 226
Table 9.16 Content of postnatal care for the newborn 227
Table 9.16C Content of postnatal care for the newborn: County level 228
Table 9.17 Postnatal health checks for mothers and newborns 229
Table 9.17C Postnatal health checks for mothers and newborns: County level 230
Table 9.18 Examinations for breast and cervical cancers 231
Table 9.18C Examinations for breast and cervical cancers: County level 232
Table 9.19 Problems in accessing health care 233
Table 9.19C Problems in accessing health care: County level 234
Table 9.20 Distance from a health care facility 235
Table 9.20C Distance from a health care facility: County level 236
Table 9.21.1 Human papillomavirus vaccination coverage among women age 15–19 237
Table 9.21.1C Human papillomavirus vaccination coverage among women age 15–19: County level 238
Table 9.21.2 Human papillomavirus vaccination coverage among women age 20–49 239
Table 9.21.2C Human papillomavirus vaccination coverage among women age 20–49: County level 240
Table 9.22 Source of human papillomavirus vaccinations 241
Table 9.22C Source of human papillomavirus vaccinations: County level 242

Figure 9.1 Trends in antenatal care coverage 178
Figure 9.2 Components of antenatal care 180
Figure 9.3 Trends in health facility births 183
Figure 9.4 Assistance during delivery 186
Figure 9.5 Trends in delivery assistance 186
Figure 9.6 Postnatal health checks for mothers and newborns 191
Figure 9.7 Breast and cervical cancer exams by residence 192

Map 9.1 Health facility births by county 184
Map 9.2 Caesarean section rates by county 185
Map 9.3 Skilled delivery assistance by county 187
Map 9.4 Cervical cancer screening by county 193

10 CHILD HEALTH 243
Table 10.1 Child's size and weight at birth 257
Table 10.1C Child's size and weight at birth: County level 258

Table 10.2 Possession and observation of vaccination cards according to background characteristics ... 259
Table 10.3 Vaccinations by source of documentation ... 260
Table 10.4 Vaccinations by background characteristics ... 261
Table 10.5 Source of vaccinations ... 263
Table 10.6.1 Children with symptoms of acute respiratory infection and care seeking for symptoms ... 264
Table 10.6.2 Treatment for children with symptoms of acute respiratory infection ... 265
Table 10.7 Source of advice or treatment for children with symptoms of acute respiratory infection ... 266
Table 10.8 Children with fever and care seeking for fever ... 267
Table 10.9 Children with diarrhoea and care seeking for diarrhoea ... 268
Table 10.10 Feeding practices during diarrhoea ... 269
Table 10.11 Oral rehydration salts, zinc, continued feeding, and other treatments for diarrhoea ... 270
Table 10.12 Source of advice or treatment for children with diarrhoea ... 271
Table 10.13 Global Scales of Early Development ... 272
Table 10.13C Global Scales of Early Development: County level ... 273
Table 10.14 Early Childhood Development Index 2030 ... 274
Table 10.14C Early Childhood Development Index 2030: County level ... 275
Table 10.15 Combined Early Childhood Development Index 2030 and Global Scales of Early Development ... 276
Table 10.15C Combined Early Childhood Development Index 2030 and Global Scales of Early Development: County level ... 277

Figure 10.1 Trends in basic antigen coverage in children ... 246
Figure 10.2 Childhood vaccinations ... 248
Figure 10.3 Diarrhoea prevalence by age ... 250
Figure 10.4 Feeding practices during diarrhoea ... 251
Figure 10.5 Treatment of diarrhoea ... 252
Figure 10.6 Symptoms of childhood illness and care seeking ... 252
Figure 10.7 Early childhood development ... 254

Map 10.1 Early childhood development ... 255

11 NUTRITION OF CHILDREN ... 279
Table 11.1 Child growth monitoring ... 291
Table 11.1C Child growth monitoring: County level ... 292
Table 11.2 Early breastfeeding ... 293
Table 11.3 Breastfeeding status according to age ... 294
Table 11.4 Infant feeding practices by age ... 295
Table 11.5 Liquids consumed by children in the day or night preceding the interview ... 296
Table 11.6 Foods consumed by children in the day or night preceding the interview ... 297
Table 11.7 Minimum dietary diversity, minimum meal frequency, and minimum acceptable diet among children ... 298
Table 11.8 Egg and/or flesh food consumption and unhealthy feeding practices among children ... 299
Table 11.9 Infant and young child feeding indicators ... 300
Table 11.10 Infant and young child feeding counselling ... 301
Table 11.11 Micronutrient supplementation and deworming among children ... 302
Table 11.11C Micronutrient supplementation and deworming among children: County level ... 303

Figure 11.1 Infant feeding practices by age 282
Figure 11.2 Trends of exclusive breastfeeding 282
Figure 11.3 IYCF indicators on Minimum Acceptable Diet (MAD), Minimum Meal Frequency (MMF) and Minimum Dietary Diversity (MDD) 285
Figure 11.4 IYCF indicators on Minimum Acceptable Diet (MAD) by mother's education 286
Figure 11.5 Unhealthy feeding practices among children age 6–23 months by mother's education 287
Figure 11.6 Unhealthy feeding practices among children age 6–23 months by residence 288

12 MALARIA 305
Table 12.1 Household possession of mosquito nets 317
Table 12.1C Household possession of mosquito nets by county 318
Table 12.2 Source of mosquito nets 319
Table 12.2C Source of mosquito nets by county 320
Table 12.3 Access to an insecticide-treated net (ITN) 321
Table 12.3C Access to an insecticide-treated net (ITN) by county 322
Table 12.4 Use of mosquito nets by persons in the household 323
Table 12.4C Use of mosquito nets by persons in the household by county 324
Table 12.5 Use of existing ITNs 325
Table 12.5C Use of existing ITNs by county 326
Table 12.6 Use of mosquito nets by children 327
Table 12.6C Use of mosquito nets by children by county 328
Table 12.7 Use of mosquito nets by pregnant women 329
Table 12.8 Main reason mosquito net was not used the night before the survey 330
Table 12.8C Main reason mosquito net was not used the night before the survey by county 331
Table 12.9 Use of Intermittent Preventive Treatment (IPTp) by women during pregnancy 332
Table 12.9C Use of Intermittent Preventive Treatment (IPTp) by women during pregnancy by county 333
Table 12.10 Children with fever and careseeking, prompt treatment, and diagnosis 334
Table 12.11 Source of advice or treatment for children with fever 335
Table 12.12 Type of antimalarial drugs used 336

Figure 12.1 Household ownership of ITNs 307
Figure 12.2 Trends in household ownership of ITNs 308
Figure 12.3 Source of ITNs 309
Figure 12.4 Trends in ITN access and use 310
Figure 12.5 ITN use 312
Figure 12.6 Trends in ITN use among pregnant women and children under age 5 312
Figure 12.7 Reason ITN was not used 313
Figure 12.8 Trends in IPTp use by pregnant women 314
Figure 12.9 Trend in ACT use by children with fever 315

Map 12.1 ITN ownership by county 309
Map 12.2 ITN use by county 311

13 HOUSEHOLD WATER AND SANITATION 337
Table 13.1 Household drinking water 347
Table 13.2 Drinking water service ladder 347
Table 13.2C Drinking water service ladder: County level 348

Table 13.3 Household sanitation facilities 349
Table 13.4 Sanitation service ladder 349
Table 13.4C Sanitation service ladder: County level 350
Table 13.5 Emptying and removal of wastes from on-site sanitation facilities 351
Table 13.5C Emptying and removal of waste from on-site sanitation facilities: County level 352
Table 13.6 Management of household excreta 353
Table 13.6C Management of household excreta: County level 354
Table 13.7 Disposal of children's stools 355
Table 13.7C Disposal of children's stools: County level 356
Table 13.8 Handwashing 357
Table 13.8C Handwashing: County level 358
Table 13.9 Menstrual hygiene 359
Table 13.9C Menstrual hygiene: County level 360

Figure 13.1 Household with improved source of drinking water service by residence 338
Figure 13.2 Household population Drinking water service by residence 339
Figure 13.3 Household with improved source of drinking water service 341
Figure 13.4 Household population sanitation service by residence 342
Figure 13.5 Appropriate management of household excreta 343
Figure 13.6 Appropriate management of household excreta by wealth 344
Figure 13.7 Menstrual hygiene by education 346

Map 13.1 At least basic service for drinking water by county 340

14 GENDER-BASED VIOLENCE 361
Table 14.1 Experience of technology-facilitated gender-based violence by any perpetrator 374
Table 14.1C Experience of technology-facilitated gender-based violence by any perpetrator: County level 375
Table 14.2 Experience of various forms of technology-facilitated gender-based violence by any perpetrator 376
Table 14.3 Experience of various forms of technology-facilitated gender-based violence by any perpetrator according to background characteristics 377
Table 14.3C Experience of various forms technology-facilitated gender-based violence by any perpetrator: County level 378
Table 14.4 Experience of technology-facilitated gender-based violence by any intimate partner 379
Table 14.4C Experience of technology-facilitated gender-based violence by any intimate partner: County level 380
Table 14.5 Persons committing technology-facilitated gender-based violence 381
Table 14.6 Forms of controlling behaviours 381
Table 14.7 Controlling behaviours of husband/intimate partner by background characteristics 382
Table 14.7C Controlling behaviours of husband/intimate partner: County level 383
Table 14.8 Experience of technology-facilitated gender-based violence by platform 384
Table 14.9 Experience of technology-facilitated gender-based violence by platform and background characteristics 385
Table 14.10 Perceived risk of violence and safety for women and girls 386
Table 14.10C Perceived risk of violence and safety for women and girls: County level 387
Table 14.11 Perceived community response to femicide 388
Table 14.11C Perceived community response to femicide: County level 389

Table 14.12 Perceived socio-cultural and religious beliefs on justification of violence against women and girls 390
Table 14.12C Perceived socio-cultural and religious beliefs on justification of violence against women and girls: County level 391
Table 14.13 Perceptions of victim blaming and privacy of domestic violence 392
Table 14.13C Perceptions of victim blaming and privacy of domestic violence: County level 393

Figure 14.1 Experience of various forms of technology-facilitated gender-based violence by any perpetrator 366
Figure 14.2 Forms of controlling behaviours 368
Figure 14.3 Experience of technology-facilitated gender-based violence by platform 369
Figure 14.4 Perceived community response to femicide by household wealth 371

Map 14.1 Experience of various forms of TFGBV by any perpetrator by county 365

15 ADULT AND MATERNAL MORTALITY 395
Table 15.1 Adult mortality rates 401
Table 15.2 Adult mortality probabilities 401
Table 15.3 Maternal mortality rates 402
Table 15.4 Maternal mortality ratio 402

Figure 15.1 Adult mortality rates by age 397
Figure 15.2 Trends in the pregnancy-related mortality ratio (PRMR) with confidence intervals 400

Appendix A SAMPLE DESIGN 407
Table A.1 Population and household distribution 408
Table A.2 Distribution of enumeration areas 409
Table A.3 Sample allocation of clusters and households by county 410
Table A.4 Expected number of women 15–49 interviews 411
Table A.5 Sample implementation 414

Appendix B ESTIMATES OF SAMPLING ERRORS 415
Table B.1 List of selected variables for sampling errors, Kenya Mini DHS 2025–26 417
Table B.2 Sampling errors: Total sample, Kenya Mini DHS 2025–26 419
Table B.3 Sampling errors: Urban sample, Kenya Mini DHS 2025–26 421
Table B.4 Sampling errors: Rural sample, Kenya Mini DHS 2025–26 423
Table B.5 Sampling errors: Mombasa sample, Kenya Mini DHS 2025–26 425
Table B.6 Sampling errors: Kwale sample, Kenya Mini DHS 2025–26 426
Table B.7 Sampling errors: Kilifi sample, Kenya Mini DHS 2025–26 427
Table B.8 Sampling errors: Tana River sample, Kenya Mini DHS 2025–26 428
Table B.9 Sampling errors: Lamu sample, Kenya Mini DHS 2025–26 429
Table B.10 Sampling errors: Taita Taveta sample, Kenya Mini DHS 2025–26 430
Table B.11 Sampling errors: Garissa sample, Kenya Mini DHS 2025–26 431
Table B.12 Sampling errors: Wajir sample, Kenya Mini DHS 2025–26 432
Table B.13 Sampling errors: Mandera sample, Kenya Mini DHS 2025–26 433
Table B.14 Sampling errors: Marsabit sample, Kenya Mini DHS 2025–26 434
Table B.15 Sampling errors: Isiolo sample, Kenya Mini DHS 2025–26 435
Table B.16 Sampling errors: Meru sample, Kenya Mini DHS 2025–26 436

Table B.17 Sampling errors: Tharaka-Nithi sample, Kenya Mini DHS 2025–26 437
Table B.18 Sampling errors: Embu sample, Kenya Mini DHS 2025–26 438
Table B.19 Sampling errors: Kitui sample, Kenya Mini DHS 2025–26 439
Table B.20 Sampling errors: Machakos sample, Kenya Mini DHS 2025–26 440
Table B.21 Sampling errors: Makueni sample, Kenya Mini DHS 2025–26 441
Table B.22 Sampling errors: Nyandarua sample, Kenya Mini DHS 2025–26 442
Table B.23 Sampling errors: Nyeri sample, Kenya Mini DHS 2025–26 443
Table B.24 Sampling errors: Kirinyaga sample, Kenya Mini DHS 2025–26 444
Table B.25 Sampling errors: Murang'a sample, Kenya Mini DHS 2025–26 445
Table B.26 Sampling errors: Kiambu sample, Kenya Mini DHS 2025–26 446
Table B.27 Sampling errors: Turkana sample, Kenya Mini DHS 2025–26 447
Table B.28 Sampling errors: West Pokot sample, Kenya Mini DHS 2025–26 448
Table B.29 Sampling errors: Samburu sample, Kenya Mini DHS 2025–26 449
Table B.30 Sampling errors: Trans Nzoia sample, Kenya Mini DHS 2025–26 450
Table B.31 Sampling errors: Uasin Gishu sample, Kenya Mini DHS 2025–26 451
Table B.32 Sampling errors: Elgeyo-Marakwet sample, Kenya Mini DHS 2025–26 452
Table B.33 Sampling errors: Nandi sample, Kenya Mini DHS 2025–26 453
Table B.34 Sampling errors: Baringo sample, Kenya Mini DHS 2025–26 454
Table B.35 Sampling errors: Laikipia sample, Kenya Mini DHS 2025–26 455
Table B.36 Sampling errors: Nakuru sample, Kenya Mini DHS 2025–26 456
Table B.37 Sampling errors: Narok sample, Kenya Mini DHS 2025–26 457
Table B.38 Sampling errors: Kajiado sample, Kenya Mini DHS 2025–26 458
Table B.39 Sampling errors: Kericho sample, Kenya Mini DHS 2025–26 459
Table B.40 Sampling errors: Bomet sample, Kenya Mini DHS 2025–26 460
Table B.41 Sampling errors: Kakamega sample, Kenya Mini DHS 2025–26 461
Table B.42 Sampling errors: Vihiga sample, Kenya Mini DHS 2025–26 462
Table B.43 Sampling errors: Bungoma sample, Kenya Mini DHS 2025–26 463
Table B.44 Sampling errors: Busia sample, Kenya Mini DHS 2025–26 464
Table B.45 Sampling errors: Siaya sample, Kenya Mini DHS 2025–26 465
Table B.46 Sampling errors: Kisumu sample, Kenya Mini DHS 2025–26 466
Table B.47 Sampling errors: Homa Bay sample, Kenya Mini DHS 2025–26 467
Table B.48 Sampling errors: Migori sample, Kenya Mini DHS 2025–26 468
Table B.49 Sampling errors: Kisii sample, Kenya Mini DHS 2025–26 469
Table B.50 Sampling errors: Nyamira sample, Kenya Mini DHS 2025–26 470
Table B.51 Sampling errors: Nairobi City sample, Kenya Mini DHS 2025–26 471
Table B.52 Sampling errors for adult and maternal mortality rates, Kenya Mini DHS 2025–26 472
Table B.53 Sampling errors for Early Childhood Development Index 2030, Kenya Mini DHS 2025–26 473

Appendix C DATA QUALITY TABLES 475
Table C.1 Household age distribution 475
Table C.2 Age distribution of eligible and interviewed women 477
Table C.3 Age displacement at ages 14/15 478
Table C.4 Age displacement at ages 49/50 479
Table C.5 Pregnancy outcomes by years preceding the survey 480
Table C.6 Completeness of reporting 480
Table C.7 Observation of mosquito nets 481
Table C.8 Observation of handwashing facility 482

Table C.9 Vaccination cards photographed 483
Table C.10 Completeness of information on siblings 484
Table C.11 Sibship size and sex ratio of siblings 484
Table C.12 Pregnancy-related mortality trends 484

Figure C1 Population pyramid 476

PREFACE

The 2025–26 Kenya Mini Demographic and Health Survey (KMDHS) is the latest in the series of demographic and health surveys conducted in Kenya since 1989. It follows seven full-scale surveys implemented in 1989, 1993, 1998, 2003, 2008–09, 2014, and 2022. The KMDHS was designed to provide a timely interim update of key demographic, nutrition, and health indicators between full survey rounds. The findings are intended to inform the design, implementation, monitoring, and evaluation of population, health, and development programmes at both the national and county levels. Data were collected from 5 November 2025 to 10 February 2026, using a nationally representative sample of 29,935 households drawn from the Kenya Household Master Sample Frame. The survey provides estimates at the national level, for rural and urban areas separately, and for selected indicators at the county level.

The 2025–26 KMDHS gathered a broad range of information on demographic, health, and nutrition indicators, including fertility levels, fertility preferences, and contraceptive use; maternal and newborn health, including antenatal and delivery care; child mortality and child health, including childhood illnesses and vaccination coverage; the nutritional status of children under age 5; ownership and use of insecticide-treated nets; health insurance coverage and access; and adult and maternal mortality. This information is intended to track Kenya's progress towards national health targets, including the Sustainable Development Goals and Vision 2030. The findings are also meant to support the evaluation of ongoing health initiatives, particularly the Universal Health Coverage programme, while providing regionally and county-disaggregated evidence to guide decentralised health planning, resource allocation, and the alignment of county strategies with national priorities.

Overall, the 2025–26 KMDHS results show continued improvement in many of Kenya's health indicators since the 2022 Kenya Demographic and Health Survey (KDHS). The total fertility rate declined from 3.4 children per woman in 2022 to 3.1 children per woman in 2025–26. The percentage of women age 15–19 who have ever been pregnant fell from 14.8% to 11.9% over the same period, while the median birth interval widened from 42.1 to 46.3 months. Total demand for family planning among currently married women stood at 76%, of which 61% was satisfied through the use of modern or other contraceptive methods. Maternal and newborn health services also strengthened: 72% of women had at least four antenatal care visits for their most recent pregnancy, up from 66% in 2022; 91% of births were delivered in a health facility in 2025 compared to 88% in 2022 and 92% were assisted by a skilled provider as compared to 89% in 2022; and 84% of mothers and 86% of newborns received a postnatal check within the first 2 days after birth.

Childhood survival and health continued to improve, with the under-5 mortality rate falling to 37 deaths per 1,000 live births –from 41 in 2022 and 115 in 2003 – while the infant and neonatal mortality rates stood at 30 and 19 deaths per 1,000 live births, respectively. Eight in ten children aged 12–23 months were fully vaccinated with all basic antigens. Nutrition indicators showed that 72% of children under age 6 months were exclusively breastfed and 68% of children age 6–59 months received vitamin A supplementation in the preceding 6 months. Access to essential services also expanded: 70% of households had electricity, up from 58% in the 2022 KDHS; 85% had access to an improved source of drinking water, up from 80% in 2022; 76% had access to an improved sanitation facility compared with 72% in 2022; and 40% of the household population had some form of health insurance, with most insured persons covered under the Social Health Authority.

The 2025–26 KMDHS was implemented by the Kenya National Bureau of Statistics (KNBS) in collaboration with the Ministry of Health and other stakeholders. Aga Khan University, Kenya, was the principal financing partner for the survey, and its support was central to the successful implementation of this round of data collection. ICF provided technical assistance through The DHS Program. Other agencies and organisations that facilitated the successful implementation of the survey through technical and/or financial support include the Gates Foundation, the World Bank, the United Nations Children's Fund (UNICEF), the United Nations Population Fund (UNFPA), and the World Health Organization. KNBS provided leadership in the overall survey planning, development of survey tools, training of personnel, data collection, and analysis.

KNBS appreciates the support provided by the various stakeholders, the steering committee, the technical committee, the secretariat, and all the survey personnel who worked tirelessly to collect the data and develop this report. We are equally grateful to Aga Khan University, Kenya, and to all our other partners for their continued commitment to strengthening Kenya's health information system. Finally, we thank the respondents across the country for taking the time to provide the valuable information that has made this report possible.



Macdonald G. Obudho, PhD, EBS, MBS
Director General
Kenya National Bureau of Statistics


ACRONYMS AND ABBREVIATIONS

ACT	artemisinin-based combination therapy
AD	age at death
AL	artemether-lumefantrine
ANC	antenatal care
ARI	acute respiratory infection
ASFR	age-specific fertility rate
BCG	Bacille Calmette-Guérin
CAPI	computer-assisted personal interviewing
CBE	Competency-based Education
CBR	crude birth rate
CHW	community health worker
CI	confidence interval
CRVS	Civil Registration and Vital Statistics
CSR	crude birth rate
CSR	caesarean section rate
CSPro	Census and Survey Processing System
DEFT	design effect
DHAP	dihydroartemisinin-piperaquine
DHS	Demographic and Health Survey
DMPA-SC	subcutaneous depot-medroxyprogesterone acetate
DPT	diphtheria, pertussis, and tetanus
EA	enumeration area
ECDI	Early Childhood Development Index
EWRR	eligible women's response rate
FBO	faith-based organization
FP	family planning
GAR	gross attendance ratio
GAR	general induced abortion rate
GBV	gender-based violence
GFR	general fertility rate
GoK	Government of Kenya
GPI	gender parity index
GPS	Global Positioning System
GSED	Global Scales for Early Development

HepB	hepatitis B
Hib	*Haemophilus influenzae* type b
HPV	human papillomavirus
HRR	household response rate
IFAS	iron and folic acid supplements
IPTp	intermittent preventive treatment (of malaria) in pregnancy
IPV	inactivated polio vaccine
IPV	intimate partner violence
ITN	insecticide-treated net
IUD	intrauterine device
IYCF	infant and young child feeding
JMP	Joint Monitoring Programme for Water Supply, Sanitation and Hygiene
K-HMSF	Kenya Household Master Sample Frame
KDHS	Kenya Demographic and Health Survey
KICA	Kenya Information and Communications Act
KMDHS	Kenya Mini Demographic and Health Survey
KMIS	Kenya Malaria Indicator Survey
KNBS	Kenya National Bureau of Statistics
KPHC	Kenya Population and Housing Census
KSh	Kenyan Shilling
LAM	lactational amenorrhoea method
LLIN	long-lasting insecticidal net
LPG	liquefied petroleum gas
MAD	minimum acceptable diet
MCPR	Modern Contraceptive Prevalence Rate
MDD	Minimum Dietary Diversity
MIS	Malaria Indicator Survey
MMF	minimum meal frequency
MMR	maternal mortality ratio
MoH	Ministry of Health
MR	measles-rubella
MUAC	mid-upper arm circumference
NAR	net attendance ratio
NGAO	National Government Administrative Officers
NGO	nongovernmental organisation
NHIF	National Hospital Insurance Fund
NMCP	National Malaria Control Program
NN	neonatal mortality
NSC	National Steering Committee

OPV	oral polio vaccine
ORS	oral rehydration salts
ORT	oral rehydration therapy
OWRR	overall women's response rate
PCV	pneumococcal conjugate vaccine
PNC	postnatal care
PNN	postneonatal mortality
PRMR	pregnancy-related mortality ratio
RHF	recommended homemade fluids
RV	rotavirus vaccine
SDG	Sustainable Development Goal
SDM	Standard Days Method
SGBV	Sexual and Gender Based Violence
SHA	Social Health Authority
SHIF	Social Health Insurance Fund
SP	sulfadoxine-pyrimethamine
TAR	total induced abortion rate
TFGBV	technology-facilitated gender-based violence
TFR	total fertility rate
TWC	Technical Working Committee
UNFPA	United Nations Population Fund
UNICEF	United Nations Children's Fund
VIP	ventilated improved pit
WHO	World Health Organization
YSD	years since death

READING AND UNDERSTANDING TABLES FROM THE 2025–26 KENYA MINI DEMOGRAPHIC AND HEALTH SURVEY (KMDHS)



- The percentage of women who have completed a secondary or higher level of education rises steadily with wealth, from 13% among women in the lowest wealth quintile to 72% among those in the highest quintile (**Figure 3.2**).
- By county, the percentage of women who have completed a secondary or higher level of education is highest in Kiambu (70%) and lowest in Mandera (14%) (**Table 3.2C** and **Map 3.1**).

Figure 3.2 **Secondary education by household wealth**

Percentage of women age 15–49 who have completed at least a secondary education by wealth quintile

Map 3.1 **Secondary education by county**

Percentage of women age 15–49 who have completed at least a secondary education

The 2025–26 Kenya Mini Demographic and Health Survey (KMDHS) final report is based on approximately 200 tables of data. For quick reference, they are located at the end of each chapter and can be accessed through links in the pertinent text (electronic version). Additionally, this more reader-friendly version features about 90 figures that clearly highlight trends, subnational patterns, and background characteristics. Large colourful maps display breakdowns for counties in Kenya. The text has been simplified to highlight key points in bullets and to clearly identify indicator definitions in boxes.

While the text and figures featured in each chapter highlight some of the most important findings from the tables, not every finding can be discussed or displayed graphically. For this reason, KMDHS data users should be comfortable reading and interpreting tables.

The following pages provide an introduction to the organisation of KMDHS tables and the presentation of background characteristics, along with a brief summary of sampling and understanding denominators. In addition, this section provides some exercises for users as they practice their new skills in interpreting KMDHS tables.

Example 1: Exposure to Mass Media

A Question Asked of All Survey Respondents

Table 3.4 Exposure to mass media 1

Percentage of women age 15–49 who are exposed to specific media on a weekly basis, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic 3	Reads a newspaper at least once a week	Watches television at least once a week	Listens to the radio at least once a week	Accesses all three media at least once a week	Accesses none of the three media at least once a week	Number of women 2
Age						
15–19	4.5	48.0	48.5	2.1	28.4	3,916
20–24	6.0	47.8	47.3	2.9	29.7	3,653
25–29	4.6	52.7	51.9	2.5	25.9	3,732
30–34	5.7	53.6	52.7	3.5	25.0	3,108
35–39	3.7	49.5	54.0	2.0	26.6	2,989
40–44	5.6	49.1	56.2	3.2	25.1	2,048
45–49	4.6	49.8	59.5	3.8	25.6	1,784
Residence						
Urban	6.4	63.7	44.1	3.6	24.4	9,829
Rural	3.8	38.4	58.7	2.0	29.1	11,402
Education[1]						
No education	0.1	17.7	22.1	0.0	65.4	1,161
Primary	2.2	38.0	54.7	1.3	31.7	6,053
Secondary	5.0	52.7	54.1	2.5	23.4	9,849
More than secondary	10.4	70.5	51.1	6.0	17.5	4,167
Wealth quintile						
Lowest	1.9	7.9	43.9	0.3	53.0	3,312
Second	2.5	26.8	61.2	1.0	30.7	3,852
Middle	4.6	53.0	59.5	2.4	23.2	4,133
Fourth	5.9	65.0	52.8	3.7	21.1	4,800
Highest	8.3	78.4	43.3	4.9	15.6	5,134
Total	5.0 4	50.1	51. 9	2.7	26.9	21,231

5

[1] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Step 1: Read the title and subtitle, highlighted in orange in the table above. They tell you the topic and the specific population group being described. In this case, the table is about women age 15–49 and their exposure to different types of media. All eligible female respondents age 15–49 were asked these questions.

Step 2: Scan the column headings—highlighted in green in Example 1. They describe how the information is categorised. In this table, the first three columns of data show different types of media that women access at least once a week. The fourth column shows women who access all three types of media, while the fifth column shows women who do not access any of the three types of media on a weekly basis. The last column lists the number of women age 15–49 interviewed in the survey.

Step 3: Scan the row headings—the first vertical column highlighted in blue in Example 1. These show the different ways the data are divided into categories based on population characteristics. In this case, the table presents women's exposure to media by age, urban-rural residence, level of education, and wealth quintile. Most of the tables in the KMDHS report will be divided into these same categories; selected additional tables are divided by county.

Step 4: Look at the row at the bottom of the table highlighted in pink. These percentages represent the totals of all women age 15–49 and their weekly access to different types of media. In this case, 5.0% of women age 15–

49 read a newspaper at least once a week, 50.1% watch television at least weekly, and 51.9% listen to the radio on a weekly basis.*

Step 5: Draw two imaginary lines, as shown on the table, to find out what percentage of women in rural areas watch television. This shows that 38.4% of women age 15–49 in rural areas watch television at least once a week.

By looking at patterns by background characteristics, we can see how exposure to mass media varies across Kenya. Mass media are often used to communicate health messages. Knowing how mass media exposure varies among different groups can help programme planners and policymakers determine how to most effectively reach their target populations.

*For the purpose of this document data are presented exactly as they appear in the table including decimal places. However, the text in the remainder of this report rounds data to the nearest whole percentage point.

Practice: Use the table in Example 1 to answer the following questions:

a) What percentage of women in Kenya do not access any of the three media at least once a week?
b) What age group of women is most likely to listen to the radio at least once a week?
c) Compare women by urban-rural residence—which group is more likely to read a newspaper at least once a week?
d) What are the lowest and the highest percentages (range) of women who access none of the three media at least once a week by education level?
e) Is there a clear pattern in weekly exposure to television by wealth quintile?

Answers:

a) 26.9%
b) Women ages 45–49: 59.5% of women in this age group listen to the radio at least once a week.
c) 6.4% of women in urban areas read a newspaper at least once a week, compared with 3.8% of women in rural areas.
d) The percentage of women who access none of the three media ranges from a low of 17.5% among women with more than secondary education to a high of 65.4% among women with no education.
e) Yes. The percentage of women who watch television at least once a week increases with increasing wealth, from 7.9% among those in the lowest wealth quintile to 78.4% among those in the highest wealth quintile.

Example 2: Use of Mosquito Nets by Children

A Question Asked of a Subgroup of Survey Respondents

Table 12.7 Use of mosquito nets by pregnant women 1

Percentage of pregnant women age 15–49 who slept under a mosquito net (treated or untreated) and under an insecticide-treated net (ITN) the night before the survey; and among pregnant women age 15–49 in households with at least one ITN, percentage who slept under an ITN the night before the survey, according to background characteristics, Kenya Mini DHS 2025–26

	Among pregnant women age 15–49 in all households:			2 Among pregnant women age 15–49 in households with at least one ITN[1]:	
Background characteristic	Percentage who slept under any mosquito net last night	Percentage who slept under an ITN[1] last night	Number of pregnant women	Percentage who slept under an ITN[1] last night	Number of pregnant women
Residence					
Urban	44.6	28.3	443	80.1	157
Rural	55.1	48.1	544	82.2	318
Malaria burden					
No transmission	(28.1)	(15.3)	127	* 4	30
Very low burden	39.6	24.3	147	74.6	48
Low burden	40.6	28.4	229	87.2	75
Moderate burden	55.6	48.3	143	84.5	82
Moderate to high burden	69.7	59.7	86	84.3	61
High burden	71.5	64.5	125	85.2	94
Very high burden	62.9	50.5	131	77.6	85
Education[2]					
No education	47.3	26.1	65	69.2	25
Primary	51.0	44.3	247	81.3	135
Secondary	49.5	39.7	492	84.1	233
More than secondary	53.2	35.7	183	78.3	84
Wealth quintile					
Lowest	48.0	42.2	205	78.6	110
Second	55.6	50.6	173	81.0	108
Middle	58.7	45.6	192	85.1	103
Fourth	50.4	35.9	224	90.5	89
Highest	40.0	23.4	194	(69.4)	66
Total	50.4	39.2	3 988	81.5	475

a b

Note: Figures in parentheses are based on 25–49 unweighted cases. An asterisk indicates that a figure is based on fewer than 25 unweighted cases and has been suppressed.
Table is based on women who stayed in the household the night before the interview.
[1] An insecticide-treated net (ITN) is a factory-treated net that does not require any further treatment. In KDHS and KMIS surveys conducted prior to 2020, this was known as a long-lasting insecticidal net (LLIN).
[2] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level

Step 1: Read the title and subtitle. In this case, the table is about two separate groups of women: pregnant women ages 15–49 in all households (a) and (b) pregnant women ages 15–49 in households with at least one ITN (b).

Step 2: Identify the two panels. First, identify the columns that refer to all pregnant women ages 15–49 (a), and then isolate the columns that refer only to pregnant women ages 15–49 in households with at least one ITN (b).

Step 3: Look at the first panel. How many pregnant women ages 15–49 in all households were interviewed? It is 988. Now look at the second panel. How many pregnant women in households with at least one ITN were interviewed? It's 475 women, or 48% of the 988. (with rounding). The second panel is a subset of the first panel.

Step 4: Approximately half (48%) of pregnant women ages 15–49 live in households with at least one ITN. Once these women are further divided into the background characteristic categories, there may be too few cases for the percentages to be reliable.

- What percentage of pregnant women ages 15–49 living in households with at least one ITN, and in the highest wealth quintile, slept under an ITN last night? 69.4%. This percentage is in parentheses because there are between 25 and 49 pregnant women (unweighted) in this category. Readers should use this number with caution—it may not be reliable. (For more information on weighted and unweighted numbers, see Example 3.)

- What percentage of pregnant women ages 15–49 living in households with at least one ITN in areas with a low malaria transmission burden slept under an ITN last night? There is no number in this cell—only an asterisk. This is because there are fewer than 25 unweighted cases, or that fewer than 25 pregnant women living in households with at least one ITN in areas with a low malaria transmission burden were interviewed. Results for this group are not reported. The subgroup is too small, and therefore the data are not reliable.

Note: When parentheses or asterisks are used in a table, the explanation will be noted under the table. If there are no parentheses or asterisks in a table, you can proceed with confidence that enough cases were included in all categories that the data are reliable.

Example 3: Understanding Sampling Weights in KMDHS Tables

A sample is a group of people who have been selected for a survey. In the KMDHS the sample is designed to represent the national population age 15–49. In addition to national data, most countries want to collect and report data on smaller geographical or administrative areas. However, doing so requires a large enough sample size in each area. For the 2025–26 KMDHS, the survey sample is representative at the national and county levels, and for urban and rural areas.

To generate statistics that are representative of the country as a whole and the 47 counties, the number of women surveyed in each county should contribute to the size of the total (national) sample in proportion to size of the county. However, if some counties have small populations, then a sample allocated in proportion to each county's population may not include sufficient women from each county for analysis. To solve this problem, counties with small populations are oversampled. For example, let's say that you have enough money to interview 21,231 women and want to produce results that are representative of Kenya as a whole and its counties (as in Table 3.1C). However, the total population of Kenya is not evenly distributed among the counties: some counties, such as Nairobi, are heavily populated while others, such as Lamu are not. Thus, Lamu must be oversampled.

A sampling statistician determines how many women should be interviewed in each county in order to get reliable statistics. The **blue column (1)** in the table to the rightshows the actual number of women interviewed in each county. Within the counties, the number of women interviewed ranges from 260 in Nyandarua to 632 in Kiambu. The number of interviews is sufficient to get reliable results in each county.

Table 3.1C Background characteristics of respondents by county

Percent distribution of women age 15–49 according to county, Kenya Mini DHS 2025–26

	Number of women		
County	3 Weighted percent	2 Weighted number	1 Unweighted number
Mombasa	3.8	802.6	478
Kwale	1.8	374.2	563
Kilifi	2.9	608.5	516
Tana River	0.6	137.1	400
Lamu	0.3	53.5	417
Taita Taveta	0.7	153.2	297
Garissa	0.8	168.3	516
Wajir	0.6	130.2	518
Mandera	0.6	123.2	484
Marsabit	0.7	141.8	416
Isiolo	0.4	75.8	431
Meru	2.5	530.3	454
Tharaka-Nithi	0.7	146.8	294
Embu	1.1	239.0	361
Kitui	1.9	412.0	468
Machakos	3.2	677.2	447
Makueni	1.6	349.9	389
Nyandarua	1.1	223.2	260
Nyeri	1.0	222.8	273
Kirinyaga	1.2	258.1	319
Murang'a	1.8	377.0	434
Kiambu	8.0	1,689.1	632
Turkana	0.8	178.0	344
West Pokot	1.1	225.4	444
Samburu	0.5	97.9	395
Trans Nzoia	1.7	370.4	500
Uasin Gishu	3.1	667.1	525
Elgeyo-Marakwet	0.8	177.5	439
Nandi	1.7	360.9	480
Baringo	0.9	187.2	364
Laikipia	0.9	195.2	321
Nakuru	5.1	1,078.4	505
Narok	2.0	427.6	470
Kajiado	2.2	467.4	369
Kericho	1.8	386.9	463
Bomet	2.3	480.4	594
Kakamega	3.1	665.1	526
Vihiga	1.1	233.0	443
Bungoma	2.9	624.9	574
Busia	1.8	372.7	441
Siaya	1.7	369.8	475
Kisumu	2.6	550.6	538
Homa Bay	2.3	485.4	570
Migori	2.4	509.1	586
Kisii	2.1	438.6	442
Nyamira	1.1	237.9	454
Nairobi	16.7	3,549.9	602
Total 15–49	100.0	21,231.0	21,231

With this distribution of interviews, some counties are overrepresented and some counties are underrepresented. For example, the population in Nairobi is about 16.7% of the population in Kenya, while Lamu's population contributes only 2.5% of the country's population. But as the blue column shows, the number of women interviewed in Nairobi accounts for only about 2.8% of the total sample of women interviewed (602/21,231) and the number of women interviewed in Lamu accounts for 2.0% (417/21,231). This unweighted distribution of women does not accurately represent the population.

In order to get statistics that are representative of Kenya, the distribution of the women in the sample needs to be weighted (or mathematically adjusted) such that it resembles the true distribution in the country. Women

from a small county, like Lamu, should contribute only a small amount to the national total. Women from a large county, like Nairobi, should contribute much more. Therefore, KMDHS statisticians mathematically calculate a "weight" that is used to adjust the number of women from each county so that each SUBUNIT's contribution to the total is proportional to the actual population of the county. The numbers in the **purple column (2)** represent the "weighted" values. The weighted values can be smaller or larger than the unweighted values at the county level. The total national sample size of 21,231 women has not changed after weighting, but the distribution of the women in the counties has been changed to represent their contribution to the total population size.

How do statisticians weight each category? They take into account the probability that a woman was selected in the sample. If you were to compare the **green column (3)** to the actual population distribution of Kenya, you would see that women in each county are contributing to the total sample with the same weight that they contribute to the population of the country. The weighted number of women in the survey now accurately represents the proportion of women who live in Nairobi and the proportion of women who live in Lamu.

With sampling and weighting, it is possible to interview enough women to provide reliable statistics at national and county levels. In general, only the weighted numbers are shown in each of the KMDHS tables, so don't be surprised if these numbers seem low: they may actually represent a larger number of women interviewed.

SUSTAINABLE DEVELOPMENT GOAL INDICATORS

Kenya Mini DHS 2025–26

Indicator	Residence		Total	DHS table number
	Urban	Rural		
1. No poverty				
1.4.1 Proportion of population living in households with access to basic services				
a) Access to basic drinking water services	95.2	59.3	73.4	13.2
b) Access to basic sanitation services	50.8	40.9	44.8	13.4
c) Access to basic hygiene services	59.5	49.2	53.2	13.8
d) Access to electricity[1]	93.5	47.5	65.5	2.3
e) Access to clean fuels and technologies[2]	61.1	5.7	27.3	2.4
	Sex			
	Male	Female	Total	
2. Zero hunger				
2.2.4 Prevalence of minimum dietary diversity, by population group				
a) Prevalence of minimum dietary diversity in children age 6–23 months	41.6	43.8	42.7	11.7
3. Good health and well-being				
3.1.1 Maternal mortality ratio[3]	na	na	263	15.4
3.1.2 Proportion of births attended by skilled health personnel	na	na	91.9	9.9
3.2.1 Under-five mortality rate[4]	45.0	30.0	37.0	8.2
3.2.2 Neonatal mortality rate[4]	24.0	14.0	19.0	8.2
3.7.1 Proportion of women of reproductive age (aged 15–49 years) who have their need for family planning satisfied with modern methods	na	73.2	na	7.14.2
3.7.2 Adolescent birth rates per 1,000 women				
a) Girls aged 10–14 years[5]	na	1.0	na	5.1
b) Women aged 15–19 years[6]	na	64.0	na	5.1
3.b.1 Proportion of the target population covered by all vaccines included in their national programme				
a) Coverage of DPT containing vaccine (3rd dose)[7]	88.9	90.9	89.9	10.4
b) Coverage of measles containing vaccine (2nd dose)[8]	70.4	73.6	72.0	10.4
c) Coverage of pneumococcal conjugate vaccine (last dose in schedule)[9]	92.3	92.4	92.4	10.4
d) Coverage of HPV vaccine (last dose in schedule)[10]	na	49.4	na	9.21.1
4. Quality education				
4.2.1 Proportion of children aged 24–59 months who are developmentally on track in health, learning and psychosocial well-being	80.3	85.2	82.8	10.14
5. Gender equality				
5.b.1 Proportion of individuals who own a mobile telephone[11]	na	81.5	na	B2
	Residence			
	Urban	Rural	Total	
6. Clean water and sanitation				
6.1.1 Proportion of population using safely managed drinking water services				
a) Proportion with basic drinking water services	95.2	59.3	73.4	13.2
6.2.1 Proportion of population using (a) safely managed sanitation services and (b) hand-washing facility with soap and water				
a) Proportion using basic sanitation service	50.8	40.9	44.8	13.4
b) Proportion in which excreta are safely disposed of in situ or treated off site	88.4	58.9	70.5	13.6
c) Proportion using a hand-washing facility with soap and water	59.5	49.2	53.2	13.8
d) Proportion using open defecation	0.4	11.1	6.9	13.4
7. Affordable clean energy				
7.1.1 Proportion of population with access to electricity[1]	93.5	47.5	65.5	2.3
7.1.2 Proportion of population with primary reliance on clean fuels and technology[2]	61.1	5.7	27.3	2.4
	Sex			
	Male	Female	Total	
16. Peace, justice, and strong institutions				
16.9.1 Proportion of children under 5 years of age whose births have been registered with a civil authority	76.8	77.9	77.4	2.9

Continued…

Continued

Indicator	Sex		Total	DHS table number
	Male	Female		
17. Partnerships for the goals				
17.8.1 Proportion of individuals using the Internet[12]	na	60.6	na	3.5

na = not applicable
[1] Persons living in households that report the primary source of lighting is electricity.
[2] Persons living in households that report no cooking, no space heating, or no lighting are not excluded from the numerator.
[3] Expressed in terms of maternal deaths per 100,000 live births in the 7-year period preceding the survey
[4] Expressed in terms of deaths per 1,000 live births for the 5-year period preceding the survey
[5] Equivalent to the age-specific fertility rate for girls age 10–14 for the 3-year period preceding the survey, expressed in terms of births per 1,000 girls age 10–14
[6] Equivalent to the age-specific fertility rate for women age 15–19 for the 3-year period preceding the survey, expressed in terms of births per 1,000 women age 15–19
[7] The percentage of children age 12–23 months who received three doses of DPT-HepB-Hib
[8] The percentage of children age 24–35 months who received two doses of MR
[9] The percentage of children age 12–23 months who received three doses of pneumococcal vaccine
[10] The percentage of women age 15–17 who received two doses of human papillomavirus (HPV) vaccine
[11] Data are available for women age 15–49 only.
[12] Data are available for women age 15–49 who have used the Internet in the last 12 months.

KENYA


SOUTH SUDAN
Lake Turkana
ETHIOPIA
UGANDA
SOMALIA
TANZANIA
Lake Victoria
Indian Ocean
Turkana
Marsabit
Mandera
Wajir
West Pokot
Samburu
Isiolo
Elgeyo-Marakwet
Bungoma
Trans Nzoia
Uasin Gishu
Baringo
Kakamega
Busia
Vihiga
Laikipia
Meru
Siaya
Nandi
Kisumu
Kericho
Nakuru
Nyandarua
Nyeri
Embu
Tharaka-Nithi
Garissa
Homa Bay
Bomet
Murang'a
Kirinyaga
Kisii
Migori
Kiambu
Narok
Machakos
Kitui
Tana River
Nairobi
Nyamira
Makueni
Kajiado
Lamu
Kilifi
Taita/Taveta
Kwale
Mombasa
N
0
125
250
500
Kilometers

INTRODUCTION AND SURVEY METHODOLOGY 1

The 2025–26 Kenya Mini Demographic and Health Survey (KMDHS) was implemented by the Kenya National Bureau of Statistics (KNBS) in collaboration with the Ministry of Health (MoH) and other stakeholders. Data collection took place from 5 November 2025 to 10 February 2026. ICF provided technical assistance through The Demographic and Health Surveys (DHS) Program; and the Gates Foundation, which offers financial support and technical assistance for population and health surveys in countries worldwide, provided financial support. Other agencies and organisations that facilitated the successful implementation of the survey through technical or financial support were the Conrad N. Hilton Foundation, the Aga Khan University Institute for Human Development, the United Nations Children's Fund (UNICEF), the United Nations Population Fund (UNFPA), and the World Health Organization (WHO).

1.1 SURVEY OBJECTIVES

The 2025–26 KMDHS was designed to provide timely and accurate data on key demographic and health indicators in Kenya to inform the design, implementation, monitoring, and evaluation of population, health, and development programmes at both national and county levels.

Specifically, the 2025–26 survey collected information on:

- Fertility levels, fertility preferences, and contraceptive use
- Maternal health, including antenatal and delivery care and maternal mortality
- Child mortality and child heath, including childhood diseases and vaccination coverage
- Nutritional status of children under age 5
- Availability of, access to, and use of insecticide-treated nets (ITNs)
- Perceptions of harm against women, gender-based violence, and technology-facilitated violence
- Health insurance coverage and access
- Adult and maternal mortality

Information was collected through the 2025–26 KMDHS to track Kenya's progress towards national health targets, including Sustainable Development Goals and Kenya Vision 2030, by providing reliable evidence on key health indicators, achievements, gaps, and areas requiring accelerated action. The survey aimed to support the evaluation of ongoing health initiatives, particularly the universal health coverage programme, by assessing their effectiveness and informing necessary refinements. The survey was also expected to produce evidence-based data to guide policy formulation, programme adjustments, prioritisation of interventions, and resource allocation while generating data segregated at the regional and county levels to support decentralised health planning, equitable resource distribution, and alignment of county strategies with national priorities.

1.2 SAMPLE DESIGN

The sample for the 2025–26 KMDHS was drawn from the Kenya Household Master Sample Frame (K-HMSF), which was developed and maintained by the KNBS for household-based sample surveys in Kenya. The K-HMSF, based on the enumeration areas (EAs) used in the 2019 Kenya Population and Housing Census, comprises 10,000 clusters selected using probability proportional to size. The 10,000 clusters were randomised into four equal subsamples, allowing surveys to utilise one or more subsamples depending on their sample size requirements. The 2025–26 KMDHS sample was drawn from subsample four of the frame. To design the

frame, each of the 47 counties in Kenya was stratified into rural and urban, resulting in 92 strata, as Nairobi City and Mombasa Counties are classified as entirely urban.

The sample was designed to provide estimates at the national level; for rural and urban areas separately; and, for selected indicators, at the county level. The computed sample size was 30,000 households, with a fixed allocation of 30 households per cluster, resulting in 1,000 clusters spread across the country: 646 in rural areas and 354 in urban areas. The sample was allocated to the different sampling strata using power allocation to ensure that the precision of the survey would be comparable across counties.

A two-stage stratified sample design was employed. In the first stage, 1,000 clusters were selected from the K-HMSF using the equal probability of selection method, independently within each sampling stratum. Households were relisted in 616 of the 1,000 selected clusters to update household information. Thereafter, the full list of households served as the sampling frame for the second stage of selection, in which 30 households were selected from each cluster. However, in clusters with fewer than 30 households (mostly in arid and semi-arid areas), all households were included in the sample. This resulted in a final sample of 29,935 households. Interviews were conducted only in the pre-selected households and clusters; no pre-selected units were allowed to be replaced during data collection. The survey was successfully implemented in 998 of the 1,000 selected clusters; one cluster in Garissa County was not visited during data collection due to insecurity, and one in Nairobi City County was excluded following refusal by residents.

As a result of the nonproportional allocation of the sample to the sampling strata and to account for nonresponse, the survey was not self-weighting. Sampling weights were therefore computed based on the selection probabilities at each stage of sampling and applied to ensure that the data were representative nationally and across all survey domains.

1.3 QUESTIONNAIRES

The Household and Woman's Questionnaires were used in the 2025–26 KMDHS. Both questionnaires, based on The DHS Program's model questionnaires, were adapted to reflect population and health issues relevant to Kenya.

The Household Questionnaire listed all members of and visitors to the selected households. Basic demographic information was collected on each person listed, including age, sex, marital status, education, and relationship to the head of the household. The data on age and sex of household members were used to identify women who were eligible for individual interviews. Information on the characteristics of the household's dwelling unit was also collected, including source of water; type of toilet facilities; materials used for flooring, external walls, and roofing; ownership of various household goods; and ownership and use of mosquito nets. In addition, respondents were asked about the death registration status of household members who had died, about any household enterprises they owned or operated, and about their perceptions of harm against women.

The Woman's Questionnaire collected information from women age 15–49 on:

- Background characteristics, including age, education, and media exposure
- Reproduction and child mortality
- Contraception
- Antenatal, delivery, and postnatal care
- Vaccinations and childhood illnesses
- Maternal and child health and nutrition
- Marriage and sexual activity
- Fertility preferences

- Other health issues and chronic diseases
- Adult mortality, including maternal mortality
- Technology-facilitated gender-based violence

The Fieldworker Questionnaire collects data on the basic characteristics of fieldworkers that can be used in analysing data quality. Fieldworkers filled out a two-page self-administered questionnaire on their general background characteristics. The KNBS distributed and collected this questionnaire before the fieldworkers entered the field. No personal identifiers were attached to the KMDHS fieldworkers' data files.

All questionnaires were programmed into tablet computers to allow computer-assisted personal interviewing (CAPI) to be used for data collection.

The CAPI survey instrument was designed to allow interviewers to collect GPS coordinates. GPS capture was integrated into the identification and interviewing process. GPS coordinates were captured to precisely georeference each household. This ensured that each household's position was logged for spatial analysis, coverage verification, and compliance checks.

The protocol for the 2025–26 KMDHS was reviewed by ICF's institutional review board. Country-specific reviews were not performed because the survey was conducted in accordance with Kenya's Statistics Act (Cap. 112).

1.4 Training of Trainers and Pretest

1.4.1 Training of Trainers

Thirty-eight trainers from the KNBS, the MoH, other government departments and agencies, universities, and development partners participated in the training of trainers. The training was supported by ICF and was held 17–23 August 2025.

The objectives of the training were to:

- Equip trainers with adult learning principles and effective facilitation methods
- Review and finalise the 2025–26 KMDHS questionnaires
- Familiarise trainers with the 2025–26 KMDHS CAPI programme
- Prepare and finalise materials for training survey personnel (interviewers and supervisors)

1.4.2 Pretest

Pretest training, which included classroom training and field practice for interviewers and supervisors, took place from 31 August to 20 September 2025. Five supervisors (four female and one male), five back-up supervisors (one female and four male), and 15 research assistants (13 female and 2 male) took part in the pilot training. Participants were grouped into five teams, each comprising one supervisor, one back-up supervisor, and three interviewers. The interviewer training focused on questionnaire content, general interviewing techniques, CAPI techniques, and fieldwork procedures. The training utilised a variety of learning tools including formal lectures, informal discussions on practice scenarios, videos, and hands-on demonstrations.

The objectives of the pretest were to:

- Test the adequacy of the training agenda for the main survey
- Test the data collection instruments (questionnaires, manuals, and forms)
- Test the accuracy and flow of the CAPI data collection programme

- Evaluate the competence of personnel
- Assess the workload of field interviewers and supervisors
- Test the adequacy of training procedures for the field personnel
- Test the adequacy of the planned duration of data collection
- Evaluate the overall administrative and financial structure as well as other general logistical issues
- Test the reliability of the central server's data transmission mechanisms and the robustness of the established system for monitoring the quality of data from the field
- Test the effectiveness of the publicity and advocacy strategy and of data processing strategies

Field practice was conducted 17–19 September 2025. Each team was assigned one cluster in Nakuru County consisting of 30 households for the team to interview over 3 days. Convenience sampling was used, and the teams selected households based on their composition, to gain experience interviewing individuals of different ages. The selected pretest clusters were not part of the 2025–26 KMDHS sample. Once the pretest fieldwork concluded, a debriefing session was held to review and discuss the survey aspects that were being tested. The outcomes of the debriefing session informed final adjustments to the questionnaires, the CAPI programme, and field logistics before the main training and data collection.

1.5 Training of Field Staff

One hundred ninety-five field personnel (47 supervisors and 148 research assistants) were engaged in the 2025–26 KMDHS main training, which took place at the Waterbuck Hotel in Nakuru, Kenya, on 5–30 October 2025. The training was led by staff from the KNBS, the MoH, and the Aga Khan University Institute for Human Development and backstopped by staff from ICF.

During the training, facilitators discussed concepts, procedures, and methodologies related to survey implementation and conducted a comprehensive step-by-step review of each questionnaire item, supported by guidance from the interviewer's manual. Participants were guided through the questionnaires both on paper and in the CAPI programme in both English and Kiswahili. Time was also allocated for local language practice. Training activities included lectures, role-play demonstrations, group discussions, paired practice interviews, and assessments to reinforce skills and ensure readiness for fieldwork.

At the end of the training, participants were grouped into 47 field teams, each comprising one supervisor and three to four interviewers, with team composition based on the languages that participants could speak and read. These teams were used for a mini-cluster simulation exercise as well as for field practice, further reinforcing the classroom learning.

A 4-day field practice took place in a mix of urban and rural areas in 19 EAs that had not been selected for the main survey. Each team was assigned 30 households to interview within an assigned EA. Teams used convenience sampling to select households for interviews, prioritising households with women with various numbers and ages of children. All teams successfully closed their clusters. Throughout the training, each individual's performance was evaluated. Those who performed well were placed on teams as supervisors or interviewers. The supervisors received additional training covering their roles and responsibilities, including how they should organise fieldwork, monitor interviews, and conduct quality control checks on both paper and CAPI questionnaires.

1.6 Fieldwork

Forty-seven field teams collected data for the 2025–26 KMDHS between 5 November and 10 February 2026. Each team consisted of a supervisor, two to three interviewers, and a driver. KNBS statistical officers supported the teams by linking them with National Government Administration Officers (NGAOs) at the

county level. Before fieldwork commenced, county mobilisation teams conducted targeted publicity within the selected clusters to prepare communities. KNBS staff and village elders assisted the teams in locating the sampled clusters and households. Throughout data collection, members of the Technical Working Committee (TWC) and the National Steering Committee (NSC) monitored progress to ensure adherence to survey protocols and to provide timely solutions to any challenges encountered.

1.7 Data Processing

The survey data were collected using tablet computers running the Android operating system and Census and Survey Processing System (CSPro) software jointly developed by the United States Census Bureau, ICF, and Serpro S.A. The DHS Program, with ICF and the KNBS, was responsible for programming the questionnaires into the Android application. Each of the field personnel was given a unique username, and devices were password protected to safeguard data integrity.

The CAPI programme was used for data collection. The programme accepted only valid responses, automatically performed checks on ranges of values, skipped to appropriate questions based on responses given, and checked the consistency of the data collected. Supervisors assigned work and transferred it to interviewers' tablets via Bluetooth. Answers to the survey questions were entered into the tablets by each interviewer. Supervisors downloaded interview data from interviewers' tablets to their tablets via Bluetooth, checked the data for completeness, and monitored fieldwork progress.

Each day, after completion of interviews, field supervisors submitted data to the central server. Data were sent to the central office via secure internet data transfer. The data processing monitors monitored the quality of the data received and downloaded data files for completed clusters into the system. ICF provided the CSPro software for data processing and offered technical assistance in the preparation of data capture, in data management, and in data editing programmes. Field check tables were produced weekly by the central office team and shared with data monitors and ICF staff. Secondary editing was conducted during and following data collection. ICF provided technical support for data processing, and the ICF team at the central office spearheaded further editing by KNBS staff to address unresolved issues in the field. A CSPro batch editing tool was applied to clean and tabulate data during analysis.

1.8 Response Rates

Table 1.1 shows the survey response rates. A total of 29,875 households were selected for the survey, of which 27,612 were found to be occupied. Overall, 26,380 households were successfully interviewed, resulting in a household response rate of 96%. In the interviewed households, 22,507 women age 15–49 were identified as eligible for individual interviews. Of these, 21,231 were successfully interviewed, yielding an eligible women's response rate of 94%. Overall, response rates varied little by residence, although they were slightly higher in rural areas than in urban areas.

Table 1.1 Results of the household and individual interviews

Number of households, number of interviews, and response rates, according to residence (unweighted), Kenya Mini DHS 2025–26

Result	Residence		Total
	Urban	Rural	
Household interviews			
Households selected	10,567	19,308	29,875
Households occupied	9,669	17,943	27,612
Households interviewed	8,950	17,430	26,380
Household response rate[1]	93	97	96
Interviews with women age 15–49			
Number of eligible women	7,771	14,736	22,507
Number of eligible women interviewed	7,241	13,990	21,231
Eligible women's response rate[2]	93	95	94

[1] Households interviewed/households occupied
[2] Women interviewed/eligible women

HOUSING CHARACTERISTICS AND HOUSEHOLD POPULATION 2

Key Findings

- ***Electricity:*** Nationally, 70% of households have electricity.
- ***Primary reliance on clean fuels and technologies:*** 27% of the population relies on clean fuels and technologies for cooking, space heating, and lighting.
- ***Birth registration:*** Three in four children (77%) under age 5 are registered with the civil registration authority.
- ***Education:*** The percentage of the household population with no education has dropped by almost half since 2003, from 23% to 12% among women and from 16% to 8% among men.
- ***Health insurance coverage:*** Nationally, 40% of the household population has some form of health insurance, with most insured households covered under the Social Health Authority.
- ***Household enterprises:*** One in five households (20%) in Kenya operates an enterprise (income-generating activity).
- ***Death certificates and permits:*** Nationally, a death certificate was issued for 55% of reported deaths among household members, and either a death certificate or a burial permit was issued for 90% of the reported deaths.

Information on the socioeconomic characteristics of the household population in the 2025–26 Kenya Mini Demographic and Health Survey (KMDHS) provides context for interpreting demographic and health indicators and for approximating the representativeness of the survey. The information also sheds light on the living conditions of the population.

This chapter presents housing characteristics and information on the household population including use of clean fuels and technologies (related to cooking, heating, and lighting), wealth, household composition, birth registration, educational attainment, health insurance, household enterprises, and deaths in the households.

2.1 HOUSING CHARACTERISTICS

Nationally, 70% of households have electricity. Most households in urban areas (94%) have electricity compared with 50% in rural areas (**Table 2.1**). Additionally, about half of the households (48%) use cement as their main flooring material.

2.1.1 Use of Clean Fuels and Technologies

Primary reliance on clean fuels and technologies

The percentage of the population using clean fuels and technologies for cooking, heating, and lighting, with each application defined as follows:

- **Clean cooking fuels and technologies**

  Includes stoves/cookers using electricity, liquefied petroleum gas (LPG)/natural gas/biogas, solar, and alcohol/ethanol.
- **Clean heating fuels and technologies**

  Includes central heating, electricity, LPG/natural gas/biogas, solar air heaters, and alcohol/ethanol.
- **Clean lighting fuels and technologies**

  Includes electricity, solar lanterns, battery-powered or rechargeable flashlights/torches/lanterns, and biogas lamps.

Sample: Households and de jure population

2.1.2 Cooking

More than half of the households in Kenya (52%) cook indoors, with 32% having no separate room or kitchen for cooking. A much higher percentage of households in rural areas (64%) than in urban areas (6%) have a separate building for cooking (**Table 2.2**).

Nationally, 37% of households use clean fuels and technologies for cooking. A higher percentage of households in urban areas (70%) than in rural areas (9%) use clean fuels and technologies for cooking. Among the 59% of households that use solid fuels for cooking, wood is the most common source of fuel (46%); however, although wood is the most common source in rural areas (79%), charcoal is the most common source in urban areas (17%).

2.1.3 Heating and Lighting

Most households (83%) do not heat their homes. Among those that do, 12% use traditional cook stoves, and only 2% use clean fuels and technologies as their heating fuel (**Table 2.3**).

Overall, 96% of households use clean fuels and technologies for lighting. The most common sources are electricity (71%) and solar lanterns (20%).

2.1.4 Primary Reliance on Clean Fuels and Technologies


Nationally, 27% of the population relies on clean fuels and technologies for cooking, space heating, and lighting. The percentage of the population relying primarily on clean fuels and technologies for cooking, space heating, and lighting increases with wealth, from 1% in the lowest wealth quintile to 83% in the highest wealth quintile (**Table 2.4**).

More than half (61%) of the urban population relies on clean fuels and technologies for cooking, space heating, and lighting compared with 6% of the rural population (**Figure 2.1**).

The percentage of the population relying primarily on clean fuels and technologies for cooking, space heating, and lighting differs across counties. The percentages of the populations in Nairobi City (81%), Kiambu (75%), Kajiado (52%), Mombasa (41%), Machakos (37%), and Kirinyaga (28%) that rely primarily on clean fuels and technologies are higher than the national average (27%). Wajir County has the lowest percentage (1%) of the population relying primarily on clean fuels and technologies, followed by West Pokot, Turkana, Mandera, and Samburu Counties (2% each) (**Table 2.4C** and **Map 2.1**).


Figure 2.1 Primary reliance on clean fuels and technologies

Percentage of de jure population relying on clean fuels and technologies for:



Map 2.1 Primary reliance on clean fuels and technologies by county

Percentage of the de jure population relying on clean fuels and technologies for cooking, space heating, and lighting



2.2 HOUSEHOLD WEALTH

2.2.1 Household Durable Goods

The socioeconomic status of households is reflected through household possessions. Most households in Kenya (95%) own a mobile phone. Almost all households in urban areas (98%) compared with 92% in rural areas own a mobile phone (**Table 2.5**).

Nationally, 63% of households have a radio (64% in urban areas and 62% in rural areas), 48% have a television (61% in urban areas and 37% in rural areas), and 9% own a computer (17% in urban areas and 3% in rural areas). Eleven percent of households own a motorcycle or scooter (9% in urban areas and 13% in rural areas).

Forty-one percent of households own agricultural land, with a higher percentage of rural households (60%) than of urban households (19%) owning the land. Ownership of farm animals is more common in rural households (67%) than in urban households (23%).

2.2.2 Wealth Index

Wealth index


Households are given scores based on the number and kinds of consumer goods they own, ranging from televisions to bicycles or cars, and on housing characteristics such as source of drinking water, toilet facilities, and flooring materials. These scores are derived using principal component analysis. National wealth quintiles are compiled by assigning the household score to each usual (de jure) household member and ranking each person in the household population by her or his score; the distribution is then divided into five equal categories, each comprising 20% of the population.

Sample: Households

Wealth index is a composite measure of a household's cumulative standard of living. In this report, it is used as a background characteristic to illustrate the effect of wealth on various population, health, and nutrition indicators.

Figure 2.2 **Household wealth by residence**

Percent distribution of de jure population by wealth quintiles



Almost half of the urban population (47%) compared with only 3% of the rural population are in the highest wealth quintile (**Table 2.6** and **Figure 2.2**). Conversely, 32% of the rural population compared with only 2% of the urban population are in the lowest wealth quintile. More than three-quarters (77%) of the population in Turkana County falls within the lowest wealth quintile, while none of the population in Nairobi City County falls within the same quintile. Nairobi City County has the highest percentage of the population in the highest wealth quintile (64%), followed by Kiambu County (51%). Mandera County (1%), Bomet County (1%), Nyamira County (2%), and West Pokot County (2%) have the lowest percentages of the population in the highest wealth quintile.

2.3 Household Characteristics and Composition

Household

A person or group of related or unrelated persons who live in the same dwelling unit(s), acknowledge one adult male or female as the head of the household, share the same housekeeping arrangements, and are considered a single unit.

De facto population

All persons who stayed in the selected households the night before the interview (whether usual residents or visitors).

De jure population

All persons who are usual residents of the selected households, whether or not they stayed in the household the night before the interview.

How data are calculated


All tables are based on the de facto population unless otherwise specified.

The age-sex structure of the Kenyan population is illustrated by the population pyramid in **Figure 2.3**. The pyramid has a broad base, indicating that young people make up the largest share of the Kenyan population. Nationally, 58% of the population falls within working ages (15–64 years), including 64% in urban areas and 54% in rural areas. The dependent population age 0–14 accounts for 37% of the national population, while those age 65 and older account for 5% (**Table 2.7**).

Female-headed households account for a third of households in Kenya. They constitute 31% of households in urban areas and 32% of households in rural areas (**Table 2.8**).

***Figure 2.3* Population pyramid**

Percent distribution of the household population



The mean household size in Kenya is 3.5 persons. Rural areas have a higher mean household size (3.9 persons) than urban areas (3.0 persons).

2.4 Birth Registration

Registered birth

A birth that is registered with the civil registration authority, whether or not the child has a birth certificate.

Sample: De jure children under age 5

All births in Kenya are required to be officially registered by the Directorate of Civil Registration Services (Republic of Kenya 1928). The registration of births establishes legal identity of children, thereby facilitating the protection of their rights and access to government services.


Nationally, 77% of children age 0–5 are registered with the civil registration authority, but only 38% have a birth certificate (**Table 2.9**).

Patterns by background characteristics

- The percentage of children with registered births is higher in urban areas (83%) than in rural areas (74%). Among children with registered births, the percentage with a birth certificate is also higher in urban areas (50%) than in rural areas (30%).
- The likelihood of birth registration increases with household wealth, from 62% in the lowest wealth quintile to 89% in the highest wealth quintile (**Figure 2.4**).
- Embu County has the highest percentage of children whose births are registered (96%), while Mandera County has the lowest (44%) (**Map 2.2**). West Pokot (12%) and Samburu (13%) Counties have the lowest percentages of children whose births are registered with a birth certificate, whereas Lamu County has the highest percentage (60%) (**Table 2.9C**).


Figure 2.4 **Birth registration by household wealth**

Percentage of de jure children under age 5 whose births are registered with the civil registration authority



Map 2.2 Birth registration by county


Percentage of de jure children under age 5 whose births are registered with the civil registration authority



Trends: The percentage of children under age 5 whose births are registered with the civil registration authority has increased steadily from 60% in the 2008–09 Kenya Demographic and Health Survey (KDHS) to 77% in the 2025–26 KMDHS. The percentage has increased more rapidly in rural areas, from 57% in 2008–09 to 74% in 2025–26 (**Figure 2.5**).

Figure 2.5 Birth registration

Percentage of de jure children under age 5 whose births are registered with the civil registration authority



2.5 EDUCATIONAL ATTAINMENT

Median educational attainment

When half of the population has completed less than the median number of years of schooling, and half of the population has completed more than the median number of years of schooling.

Sample: De facto household population age 6 and older


In 2017, Kenya introduced the Competency-based Education (CBE), to replace the 8-4-4 system that comprised 8 years of primary education, 4 years of secondary education, and a minimum of 4 years of tertiary education. The CBE was implemented gradually in phases across education levels and followed a 2-6-3-3-3 structure: 2 years of pre-primary education, 6 years of primary education, 3 years of junior secondary education, 3 years of senior secondary education, and a minimum of 3 years of tertiary education.

Patterns by background characteristics

- The median number of years of education completed is higher in urban areas (7.7 years for women and 8.1 years for men) than in rural areas (7.0 years for women and 7.1 years for men).
- The percentage of the population with more than a secondary education increases with wealth, from less than 1% of both women and men in the lowest wealth quintile to 15% of women and 23% of men in the highest wealth quintile.
- The median number of years of schooling is highest in Kiambu County at 8 years for women and 8.9 years for men. In Mandera, Wajir, Garissa, Turkana, West Pokot, and Samburu Counties, the median number of years of schooling is 5.5 or less for women, and between 5.2 and 6.2 for men (**Tables 2.10.1C**, **2.10.2C**, **2.10.3C**, and **2.10.4C**).

Figure 2.6 **Educational attainment of the household population**

Percentage of the household population age 6 and above with no education



2.6 HEALTH INSURANCE

2.6.1 Coverage


Patterns by background characteristics

- Forty-four percent of the population in urban areas compared with 38% of the population in rural areas has some form of health insurance.
- Health insurance coverage is highest among those in the highest wealth quintile (52%) and lowest among those in the lowest wealth quintile (24%).
- Health insurance coverage is highest in Nyeri County (67%) and lowest in Tana River County (14%). Other counties with low health insurance coverage include Isiolo (19%), Garissa (23%), and Mandera (24%) (**Map 2.3**).

- Murang'a County has the highest percentage of households covered under a county-level insurance scheme (3%), while all other counties record percentages below 1%.

Map 2.3 Health insurance coverage by county

Percentage of the de jure household population with any form of health insurance



2.6.2 Status of Premium Payments under the Social Health Insurance Fund

Patterns by background characteristics

- Fifty-six percent of the population in urban areas and 34% of the population in rural areas had paid their SHIF premiums for the current month.
- The percentage of the population that had paid their SHIF premiums for the current month increases with household wealth, from 18% in the lowest wealth quintile to 66% in the highest wealth quintile.
- Seventy-four percent of people age 0–14 and age 65 and older were up to date on their SHIF premium payments.
- Kiambu County had the highest percentage of the population that was up to date on SHIF premium payments (69%), while Kwale County had the lowest percentage (10%) (**Table 2.12C**).

2.7 HOUSEHOLD ENTERPRISES

Patterns by background characteristics

- The percentage of households operating an enterprise increases with household wealth, from 10% in the lowest wealth quintile to 27% in the highest wealth quintile.
- Female-operated enterprises are more common (59%) than male-operated enterprises (42%) (**Table 2.14**).
- The percentage of women operating enterprises declines with household wealth, from 67% in the lowest wealth quintile to 55% in the highest wealth quintile.

2.7.1 Registration Status

2.7.2 Type of Economic Activity and Average Monthly Revenue

2.8 HOUSEHOLD MEMBER DEATHS

Household member death
Death of a de jure household member that occurred in the 5 years preceding the survey (2021–2025).

Health facility
A hospital, health centre, dispensary, maternity home, nursing home, or any other recognised health facility.

Community
At home, in transit, at work, on the road, or in any other place that is not a health facility.

2.8.1 Place of Death

Patterns by background characteristics

- Deaths among children under age 5 occurred much more frequently in health facilities (72%) than in the community (28%). Among persons who died at age 50–64, 67% of the deaths occurred in health facilities, while 32% occurred in the community.
- The percentage of deaths occurring in health facilities was identical among male and female household members (59% for both sexes).
- Urban residents were slightly more likely to die in health facilities (61%) than were rural residents (58%).

2.8.2 Issuance of Death Certificates and Burial Permits

Death certificate

A document issued by the Registrar of Births and Deaths in Kenya, certifying the death of an individual. It contains vital information such as the deceased person's name, age, cause of death, date and place of death, and other relevant details.

Burial permit

An official document, usually issued only for a legally registered death, authorising the removal of the dead body to the cemetery or other final disposal place.

Sample: Usual household members who died in the 5 years preceding the survey (2021–2025)

Overall, about 4% of women and 7% of men have attained more than secondary education, while 2% of women and 3% of men have completed junior school. A higher percentage of women (12%) than of men (8%) have no education, and about 18% of women compared with 17% of men have completed only primary education. The median number of years of schooling completed is 7.2 for women and 7.4 for men (**Tables 2.10.1**, **2.10.2**, **2.10.3**, and **2.10.4**).

Trends: Between the 2003 KDHS and the 2025–26 KMDHS, the percentage of the household population age 6 and older with no education decreased from 23% to 12% among women and from 16% to 8% among men (**Figure 2.6**).

Health insurance provides households with protection against unforeseen medical expenses. It is a key component of financial protection, as out-of-pocket health expenses during illness can push households into poverty.

Nationally, 40% of the de jure household population has some form of health insurance. The largest percentage of insured households (39%) are covered under the Social Health Authority (SHA) (**Tables 2.11** and **2.11C**).

Up-to-date payment of premiums under the Social Health Insurance Fund (SHIF) is essential for accessing health services. Being registered is not enough, as premiums must also be up to date for coverage to be valid. Nationally, 44% of the household population that were registered with the SHA as contributing members (not dependents) had paid their SHIF premiums for the current month (**Table 2.12**).

Household enterprises are activities operated by household members to generate income. They refer to economic units engaged in the production of goods or the provision of services. Household income is an important determinant of access to health care and related services.

Nationally, one in five households (20%) operates an enterprise. The percentage of households operating an enterprise is higher in urban areas (25%) than in rural areas (15%) (**Table 2.13**).

Registration of an enterprise as a limited liability company is important because it gives the business a legal identity separate from that of its owner or owners. This means that the owners are not personally liable for the business debts beyond their investment in the company.

Thirty percent of household enterprises were registered or licensed by the County Government, while 7% were registered or licensed by the Business Registration Service, formerly known as the Registrar of Companies (**Table 2.15**).

The economic activity of an enterprise is defined by the type of goods produced or services offered by the economic unit or establishment. Most household enterprises operate in the services sector (89%), while 6% operate in the manufacturing sector and 5% in the agri-business sector. All household income-generating activities in Garissa and Wajir Counties are in the services sector (**Table 2.16**).

Nationally, the largest percentage of household enterprises (34%) reported an average monthly revenue of KSh 5,000 or less. This was followed by enterprises with average monthly revenues of KSh 10,000–19,999 (22%) and KSh 20,000–49,999 (21%) (**Table 2.17**).

Enterprises with an average monthly revenue of KSh 50,000 or more account for 15% of households in urban areas compared with 6% of households in rural areas.

Data on household deaths are useful for generating reliable mortality statistics. They help in understanding levels and patterns of mortality within the population, which are critical for planning public health services, reducing preventable deaths, monitoring development goals, and informing policy decisions.

Nationally, most household member deaths reported in the 5 years preceding the 2025–26 KMDHS occurred in health facilities (59%), while 41% occurred in the community (**Table 2.18**).

The issuance of death certificates and burial permits is important for strengthening Kenya's civil registration and vital statistics system and for improving the availability of reliable mortality data for planning, formulating policies, and implementing programmes (Ministry of Health 2022a). Death registration supports the production of official mortality statistics, including information on age, sex, and causes of death. These data are essential for monitoring population health trends, guiding health interventions, and allocating resources across sectors (United Nations Statistics Division 2013; World Health Organization 2024).

In Kenya, death certificates also serve important roles in legal and administrative functions, including settling inheritance claims, accessing insurance and pension benefits, and updating population registers. Burial permits facilitate the notification and recording of deaths occurring both in health facilities and in the community.

Nationally, a death certificate was issued for 55% of household member deaths reported in the 5 years preceding the survey, and a burial permit was issued for 35%. Overall, either a death certificate or a burial permit was issued for 90% of the deaths, while neither document was issued for 10% of the deaths (**Tables 2.19** and **2.19C**).

Table 2.20 presents the percentage distribution of household member deaths in the 5 years preceding the survey for which no burial permit was issued, according to the main reason for non-issuance. Nationally, the main reason for non-issuance was perceived lack of benefit (48%), followed by cultural and social reasons (19%) and cost (13%). Smaller percentages of respondents (<10%) cited other reasons including lack of access and geographical barriers.

For more information on housing characteristics and the household population, see the following tables:

Patterns by background characteristics

- Death certification increases with age, rising from 36% among deaths of children under age 5 to 67% among deaths of persons age 50–64.

- Death certification or issuance of a burial permit is least likely for deaths among children under age 5 (74%), compared with more than 90% among deaths of persons age 15 and older.
- Female deaths are more likely than male deaths to have no death certificate and no burial permit issued for them (12% versus 9%).
- The percentage of deaths for which either a death certificate or a burial permit was issued was similar in urban (89%) and rural (90%) areas.

2.8.3 Reasons for Non-issuance of Burial Permits

Patterns by background characteristics

- Perceived lack of benefit was the leading reason for non-issuance of burial permits across age groups, cited for 57% of deaths among children under age 5 and 54% of deaths among persons age 65 and older.
- Cost was cited as a reason for non-issuance for a higher percentage of reported female deaths (19%) than of reported male deaths (7%).
- Cultural and social reasons for non-issuance were reported at similar levels for male (18%) and female (20%) deaths.
- Rural residents were more likely to cite cost (15%) and access or geographical barriers (6%) as reasons for non-issuance than were urban residents (5% and 2%, respectively).

LIST OF TABLES

- **Table 2.1** Household characteristics: Housing
- **Table 2.2** Household characteristics: Cooking
- **Table 2.3** Household characteristics: Heating and lighting
- **Table 2.4** Primary reliance on clean fuels and technologies
- **Table 2.4C** Primary reliance on clean fuels and technologies: County level
- **Table 2.5** Household possessions
- **Table 2.6** Wealth quintile
- **Table 2.7** Household population by age, sex, and residence
- **Table 2.8** Household composition
- **Table 2.9** Birth registration of children under age 5
- **Table 2.9C** Birth registration of children under age 5: County level
- **Table 2.10.1** Educational attainment of the female household population (standard DHS education categories)
- **Table 2.10.1C** Educational attainment of the female household population (standard DHS education categories): County level
- **Table 2.10.2** Educational attainment of the female household population (new education system)
- **Table 2.10.2C** Educational attainment of the female household population (new education system): County level
- **Table 2.10.3** Educational attainment of the male household population (standard DHS education categories)
- **Table 2.10.3C** Educational attainment of the male household population (standard DHS education categories): County level
- **Table 2.10.4** Educational attainment of the male household population (new education system)
- **Table 2.10.4C** Educational attainment of the male household population (new education system): County level
- **Table 2.11** Health insurance coverage
- **Table 2.11C** Health insurance coverage: County level
- **Table 2.12** Status of premium payments under the Social Health Insurance Fund
- **Table 2.12C** Status of premium payments under the Social Health Insurance Fund: County level
- **Table 2.13** Households operating an income-generating business/enterprise
- **Table 2.14** Household enterprises by sex of operator
- **Table 2.15** Registration status of household enterprises
- **Table 2.16** Type of economic activity of household enterprises
- **Table 2.17** Average monthly revenue of household enterprises
- **Table 2.18** Household member deaths
- **Table 2.19** Death certificate and burial permit issuance
- **Table 2.19C** Death certificate and burial permit issuance: County level
- **Table 2.20** Reasons for non-issuance of burial permits

Table 2.1 Household characteristics: Housing

Percent distribution of households and de jure population by housing characteristics and percent distribution by frequency of smoking in the home, according to residence, Kenya Mini DHS 2025–26

	Households			Population		
	Residence			Residence		
Characteristic	Urban	Rural	Total	Urban	Rural	Total
Electricity						
Yes	94.2	49.8	69.9	93.7	46.0	64.7
No	5.8	50.2	30.1	6.3	54.0	35.3
Total	100.0	100.0	100.0	100.0	100.0	100.0
Flooring material						
Earth, sand	4.2	28.1	17.2	4.8	29.5	19.9
Dung	0.7	20.7	11.6	0.8	22.8	14.2
Wood/planks	0.0	0.1	0.1	0.0	0.1	0.1
Palm/bamboo	0.0	0.0	0.0	0.0	0.0	0.0
Parquet or polished wood	0.1	0.0	0.0	0.1	0.0	0.0
Vinyl or asphalt strips	0.1	0.1	0.1	0.2	0.0	0.1
Ceramic tiles	34.9	9.0	20.8	36.9	8.7	19.7
Cement	56.9	40.9	48.2	54.2	37.8	44.2
Carpet	3.1	1.1	2.0	3.1	1.0	1.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Rooms used for sleeping						
One	66.3	34.6	49.0	51.1	24.9	35.2
Two	24.2	40.4	33.0	32.1	43.9	39.3
Three or more	9.5	24.9	17.9	16.8	31.1	25.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Frequency of smoking in the home						
Daily	4.0	7.1	5.7	3.2	6.2	5.0
Weekly	1.3	2.1	1.7	1.0	2.0	1.6
Monthly	0.3	0.5	0.4	0.2	0.5	0.4
Less than once a month	0.5	0.6	0.6	0.4	0.6	0.5
Never	93.9	89.7	91.6	95.2	90.7	92.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Number of households/ population	11,975	14,405	26,380	36,354	56,592	92,946

Table 2.2 Household characteristics: Cooking

Percent distribution of households and de jure population by place for cooking, cooking technology, and cooking fuel, according to residence, Kenya Mini DHS 2025–26

	Households			Population		
	Residence			Residence		
Characteristic	Urban	Rural	Total	Urban	Rural	Total
Place for cooking						
In the house	85.8	24.0	52.1	82.6	19.2	44.0
Separate room/kitchen	33.3	9.9	20.5	38.6	8.7	20.4
No separate room/kitchen	52.5	14.1	31.5	44.0	10.5	23.6
In a separate building	6.4	64.0	37.9	9.7	70.4	46.7
Outdoors	5.3	10.1	8.0	6.7	9.8	8.6
No food cooked in household	2.4	1.9	2.1	0.9	0.6	0.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Main cooking technology						
Clean fuels and technologies	70.3	9.3	37.0	65.8	6.2	29.5
Electric stove	0.2	0.1	0.1	0.2	0.1	0.1
Solar cooker	0.0	0.0	0.0	0.0	0.0	0.0
LPG/natural gas stove	69.2	9.0	36.3	65.0	5.9	29.0
Piped natural gas stove	0.4	0.1	0.2	0.3	0.1	0.1
Biogas stove	0.0	0.1	0.1	0.0	0.1	0.1
Liquid fuel stove using alcohol/ethanol	0.4	0.0	0.2	0.3	0.0	0.1
Other fuels and technologies	27.3	88.8	60.9	33.3	93.2	69.8
Liquid fuel stove not using alcohol/ethanol	3.3	0.1	1.6	3.0	0.0	1.2
Manufactured solid fuel stove (Jiko)	16.0	9.1	12.2	19.3	7.3	12.0
With a chimney	2.8	1.9	2.3	3.6	1.7	2.5
Without a chimney	13.2	7.2	9.9	15.7	5.6	9.6
Traditional solid fuel stove	0.9	6.3	3.9	0.9	7.6	5.0
With a chimney	0.2	1.0	0.6	0.2	1.0	0.7
Without a chimney	0.7	5.3	3.3	0.7	6.5	4.3
Three stone stove/open fire	7.1	73.4	43.3	10.0	78.3	51.6
No food cooked in household	2.4	1.9	2.1	0.9	0.6	0.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Cooking fuel						
Clean fuels and technologies[1]	70.3	9.3	37.0	65.8	6.2	29.5
Solid fuels for cooking	23.6	88.7	59.2	29.9	93.2	68.4
Coal/lignite	0.0	0.0	0.0	0.0	0.0	0.0
Charcoal	16.6	9.0	12.4	19.9	7.3	12.2
Wood	6.9	79.3	46.4	9.9	85.3	55.8
Straw/shrubs/grass	0.0	0.2	0.1	0.0	0.3	0.2
Agricultural crop	0.0	0.1	0.1	0.0	0.1	0.1
Animal dung/waste	0.0	0.0	0.0	0.0	0.0	0.0
Processed biomass (pellets) or woodchips	0.0	0.1	0.1	0.0	0.1	0.1
Sawdust	0.0	0.0	0.0	0.0	0.0	0.0
Other fuels	3.7	0.1	1.7	3.4	0.0	1.3
Gasoline/diesel	0.5	0.0	0.2	0.4	0.0	0.1
Kerosene/paraffin	3.2	0.1	1.5	2.9	0.0	1.1
Other	0.1	0.0	0.0	0.2	0.0	0.1
No food cooked in household	2.4	1.9	2.1	0.9	0.6	0.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Number of households/ population	11,975	14,405	26,380	36,354	56,592	92,946

LPG = Liquefied petroleum gas

[1] Includes stoves/cookers using electricity, LPG/natural gas/biogas, solar, and alcohol/ethanol

Table 2.3 Household characteristics: Heating and lighting

Percent distribution of households and de jure population by heating technology, heating fuel, and main lighting fuel or technology, according to residence, Kenya Mini DHS 2025–26

	Households			Population		
	Residence			Residence		
Characteristic	Urban	Rural	Total	Urban	Rural	Total
Heating technology						
Central heating	0.7	0.2	0.4	0.8	0.1	0.4
Manufactured space heater	1.6	0.2	0.8	1.7	0.1	0.7
With a chimney	0.1	0.0	0.0	0.1	0.0	0.0
Without a chimney	1.6	0.2	0.8	1.6	0.1	0.7
Traditional space heater	0.0	0.0	0.0	0.0	0.0	0.0
With a chimney	0.0	0.0	0.0	0.0	0.0	0.0
Without a chimney	0.0	0.0	0.0	0.0	0.0	0.0
Manufactured cookstove	0.1	0.2	0.1	0.1	0.2	0.2
With a chimney	0.0	0.0	0.0	0.0	0.0	0.0
Without a chimney	0.1	0.2	0.1	0.1	0.2	0.2
Traditional cookstove	9.9	14.4	12.3	11.3	14.2	13.0
With a chimney	0.9	2.5	1.8	0.9	2.4	1.8
Without a chimney	9.0	11.9	10.6	10.4	11.7	11.2
Three stone stove/open fire	0.3	4.2	2.5	0.4	4.6	3.0
Other	0.2	1.5	0.9	0.2	1.6	1.1
No heating in household	87.2	79.4	82.9	85.5	79.1	81.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Heating fuel						
Clean fuels and technologies[1]	2.7	0.6	1.5	2.9	0.4	1.4
Central heating	0.7	0.2	0.4	0.8	0.1	0.4
Electricity	1.9	0.4	1.1	2.0	0.3	1.0
Other clean fuels and technologies[2]	0.1	0.0	0.0	0.1	0.0	0.0
Coal/lignite	0.0	0.3	0.2	0.0	0.3	0.2
Charcoal	9.6	13.6	11.8	11.0	13.4	12.5
Wood	0.5	6.1	3.5	0.6	6.6	4.3
Other fuel[3]	0.0	0.1	0.0	0.0	0.1	0.0
No heating in household	87.2	79.4	82.9	85.5	79.1	81.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Main lighting fuel or technology						
Clean fuels and technologies	98.4	94.6	96.3	98.8	95.4	96.7
Electricity	94.0	51.5	70.8	93.5	47.5	65.5
Solar lantern	3.3	33.4	19.7	4.2	37.8	24.7
Rechargeable flashlight/ torch/lantern	0.8	5.7	3.5	0.7	5.8	3.8
Battery powered flashlight/ torch/lantern	0.4	4.0	2.3	0.4	4.2	2.7
Biogas lamp	0.0	0.0	0.0	0.0	0.0	0.0
Kerosene/paraffin lamp	0.3	2.2	1.4	0.4	1.8	1.2
Wood	0.0	0.5	0.3	0.0	0.5	0.3
Straw/shrubs/grass	0.0	0.2	0.1	0.0	0.3	0.2
Oil lamp	0.1	0.4	0.2	0.0	0.3	0.2
Candle	0.9	1.0	1.0	0.6	0.7	0.6
Other fuel[4]	0.1	0.2	0.2	0.1	0.1	0.1
No lighting in household	0.1	0.9	0.5	0.1	1.0	0.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Number of households/ population	11,975	14,405	26,380	36,354	56,592	92,946

[1] Includes electricity, liquefied petroleum gas (LPG)/natural gas/biogas, solar air heater, and alcohol/ethanol
[2] Includes piped natural gas, solar air heater, LPG, biogas, and alcohol and ethanol
[3] Includes straw/shrubs/grass, agricultural crops, processed biomass, saw dust, and other heating sources
[4] Includes gasoline lamp, charcoal, and other fuels mentioned by the respondent

Table 2.4 Primary reliance on clean fuels and technologies

Percentage of de jure population relying on clean fuels and technologies for cooking; percentage relying on solid fuels for cooking; percentage relying on clean fuel and technologies for space heating; percentage relying on clean fuel and technologies for lighting; and percentage relying on clean fuels and technologies for cooking, space heating, and lighting, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Primary reliance on clean fuels and technologies for cooking[1]	Primary reliance on solid fuels for cooking[2]	Number of persons in households that reported cooking in the house	Primary reliance on clean fuels and technologies for space heating[3]	Number of persons in households that reported use of space heating	Primary reliance on clean fuels and technologies for lighting[4]	Number of persons in households that reported use of lighting	Primary reliance on clean fuels and technologies for cooking, space heating, and lighting[5]	Number of persons
Residence									
Urban	66.4	30.1	36,010	19.9	5,289	98.9	36,332	61.1	36,354
Rural	6.2	93.8	56,242	2.1	11,814	96.4	56,045	5.7	56,592
Wealth quintile									
Lowest	0.0	100.0	18,434	0.5	2,805	91.7	18,047	0.7	18,594
Second	1.0	98.5	18,440	0.8	3,487	95.9	18,573	1.6	18,589
Middle	8.0	89.6	18,355	2.1	4,597	99.0	18,579	7.4	18,584
Fourth	49.1	47.6	18,462	4.5	3,680	100.0	18,589	43.6	18,589
Highest	89.9	9.6	18,561	39.5	2,535	100.0	18,589	83.4	18,589
Total	29.7	68.9	92,253	7.6	17,103	97.3	92,377	27.3	92,946

[1] Includes stoves/cookers using electricity, liquefied petroleum gas (LPG)/natural gas/biogas, solar, and alcohol/ethanol
[2] Includes coal/lignite, charcoal, wood, straw/shrubs/grass, agricultural crops, and animal dung/waste, processed biomass (pellets) or woodchips, garbage/plastic, and sawdust
[3] Includes central heating, electricity, LPG/natural gas/biogas, solar air heater, and alcohol/ethanol
[4] Includes electricity, solar lantern, rechargeable flashlight/torch/lantern, battery powered flashlight/torch/lantern, and biogas lamp
[5] To calculate Sustainable Development Goal Indicator 7.1.2, persons living in households that report no cooking, no space heating, or no lighting are not excluded from the numerator.

Table 2.4C Primary reliance on clean fuels and technologies: County level

Percentage of de jure population relying on clean fuels and technologies for cooking; percentage relying on solid fuels for cooking; percentage relying on clean fuel and technologies for space heating; percentage relying on clean fuel and technologies for lighting; and percentage relying on clean fuels and technologies for cooking, space heating, and lighting, according to county, Kenya Mini DHS 2025–26

County	Primary reliance on clean fuels and technologies for cooking[1]	Primary reliance on solid fuels for cooking[2]	Number of persons in households that reported cooking in the house	Primary reliance on clean fuels and technologies for space heating[3]	Number of persons in households that reported use of space heating	Primary reliance on clean fuels and technologies for lighting[4]	Number of persons in households that reported use of lighting	Primary reliance on clean fuels and technologies for cooking, space heating, and lighting[5]	Number of persons
Mombasa	47.3	48.8	2,922	1.8	342	96.3	2,969	41.1	2,969
Kwale	5.9	93.7	1,577	0.0	2	95.5	1,570	6.4	1,586
Kilifi	12.2	87.8	2,650	0.0	2	97.8	2,661	12.6	2,666
Tana River	1.7	98.2	716	-	0	99.3	721	2.5	721
Lamu	7.7	92.3	256	20.9	3	98.3	253	9.5	261
Taita Taveta	16.4	83.6	687	2.7	153	98.2	691	15.5	691
Garissa	2.8	97.2	841	56.5	19	99.5	665	3.7	852
Wajir	0.3	99.7	841	0.0	15	98.2	693	1.4	850
Mandera	0.1	99.9	797	81.7	15	99.8	810	1.9	812
Marsabit	6.2	93.8	746	4.1	80	99.7	748	6.4	748
Isiolo	8.0	92.0	374	0.0	26	99.8	376	8.7	377
Meru	12.0	88.0	2,657	9.2	186	94.3	2,684	12.0	2,693
Tharaka-Nithi	7.0	93.0	701	4.7	16	95.6	703	6.9	704
Embu	26.0	73.9	1,021	7.6	48	96.7	1,035	26.0	1,035
Kitui	6.4	93.5	2,104	6.7	111	97.6	2,092	6.7	2,122
Machakos	38.1	60.9	2,950	0.7	578	98.3	2,991	36.7	2,993
Makueni	9.5	90.1	1,639	1.5	130	98.5	1,665	10.0	1,665
Nyandarua	13.7	86.3	1,055	3.9	835	96.4	1,060	9.0	1,060
Nyeri	19.4	80.3	1,138	6.7	274	96.9	1,145	18.8	1,148
Kirinyaga	28.0	71.8	1,197	23.3	32	94.0	1,209	27.7	1,210
Murang'a	18.4	81.5	1,704	13.8	153	95.3	1,707	17.5	1,708
Kiambu	77.1	22.0	6,044	45.7	573	99.4	6,075	75.2	6,076
Turkana	1.6	98.4	1,181	0.0	86	81.3	1,120	1.8	1,183
West Pokot	2.0	98.0	1,255	1.4	531	96.0	1,252	1.7	1,259
Samburu	3.1	96.9	554	2.5	394	94.3	547	2.2	554
Trans Nzoia	7.5	92.5	1,564	0.5	291	98.0	1,565	6.3	1,568
Uasin Gishu	33.6	66.4	2,647	3.1	1,155	98.4	2,675	23.0	2,677
Elgeyo-Marakwet	6.7	93.3	843	0.1	319	98.1	845	6.0	848
Nandi	6.6	93.4	1,539	0.8	643	98.4	1,549	5.2	1,550
Baringo	4.2	95.8	1,011	4.0	331	97.5	1,002	3.5	1,012
Laikipia	18.5	81.5	918	3.8	231	97.5	930	17.5	931
Nakuru	31.0	68.9	4,613	3.9	2,374	98.7	4,655	23.1	4,659
Narok	8.4	91.6	2,260	2.1	537	98.9	2,258	7.9	2,265
Kajiado	54.0	43.7	1,961	8.5	351	99.4	1,990	51.7	1,991
Kericho	9.2	90.8	1,767	0.9	881	99.3	1,777	5.4	1,777
Bomet	5.0	95.0	1,950	2.6	581	99.4	1,959	4.8	1,963
Kakamega	8.2	91.8	3,153	1.1	669	94.5	3,155	6.8	3,162
Vihiga	5.6	94.4	1,055	0.6	249	93.3	1,055	3.8	1,055
Bungoma	5.7	94.3	3,036	3.5	188	97.0	3,042	5.4	3,052
Busia	7.6	92.4	1,748	10.2	72	95.1	1,753	8.1	1,758
Siaya	6.0	94.0	1,784	0.4	339	96.2	1,802	6.0	1,804
Kisumu	21.5	78.1	2,281	4.0	587	96.1	2,288	18.5	2,293
Homa Bay	6.1	93.8	2,232	1.6	451	97.4	2,245	6.1	2,246
Migori	14.7	85.3	2,231	6.0	353	98.6	2,240	12.9	2,240
Kisii	10.9	89.1	2,196	2.3	220	91.1	2,208	10.9	2,210
Nyamira	7.6	92.4	1,044	0.0	119	93.9	1,050	7.5	1,050
Nairobi City	87.9	4.5	12,814	36.8	1,556	99.5	12,891	81.3	12,891
Total	29.7	68.9	92,253	7.6	17,103	97.3	92,377	27.3	92,946

[1] Includes stoves/cookers using electricity, liquefied petroleum gas (LPG)/natural gas/biogas, solar, and alcohol/ethanol
[2] Includes coal/lignite, charcoal, wood, straw/shrubs/grass, agricultural crops, and animal dung/waste, processed biomass (pellets) or woodchips, garbage/plastic, and sawdust
[3] Includes central heating, electricity, LPG/natural gas/biogas, solar air heater, and alcohol/ethanol
[4] Includes electricity, solar lantern, rechargeable flashlight/torch/lantern, battery powered flashlight/torch/lantern, and biogas lamp
[5] To calculate Sustainable Development Goal Indicator 7.1.2, persons living in households that report no cooking, no space heating, or no lighting are not excluded from the numerator.

Table 2.5 Household possessions

Percentage of households possessing various household effects, means of transportation, agricultural land, and livestock/farm animals, according to residence, Kenya Mini DHS 2025–26

Possession	Residence		Total
	Urban	Rural	
Household effects			
Radio	63.5	61.9	62.7
Television	60.8	37.1	47.9
Mobile phone	98.2	91.9	94.8
Computer	16.9	3.2	9.4
Non-mobile telephone	2.2	1.1	1.6
Refrigerator	19.1	3.4	10.5
Watch	39.7	17.4	27.5
Solar panel	6.8	37.7	23.7
Table	84.1	86.0	85.1
Chair	79.0	88.2	84.1
Sofa	67.8	54.9	60.8
Bed	93.4	91.9	92.5
Cupboard	45.1	41.6	43.2
Clock	15.7	7.0	11.0
Microwave oven	11.7	2.1	6.5
DVD player	6.8	3.4	4.9
Cassette or CD player	3.3	1.7	2.4
Means of transportation			
Bicycle	10.7	10.5	10.6
Animal-drawn cart	0.9	2.5	1.8
Motorcycle/scooter	8.8	13.1	11.1
Car/truck	7.9	4.2	5.9
Boat with a motor	0.7	0.4	0.5
Ownership of agricultural land	19.1	59.5	41.2
Ownership of non-agricultural land	10.9	14.9	13.1
Ownership of farm animals[1]	22.6	67.3	47.0
Number of households	11,975	14,405	26,380

[1] Local cattle (indigenous), exotic/grade cattle, horses, donkeys, camels, goats, sheep, chickens/other poultry, or pigs

Table 2.6 Wealth quintiles

Percent distribution of de jure population by wealth quintile, and the Gini coefficient, according to residence and county, Kenya Mini DHS 2025–26

Residence/County	Wealth quintile					Total	Number of persons	Gini coefficient[1]
	Lowest	Second	Middle	Fourth	Highest			
Residence								
Urban	2.0	4.7	12.2	34.1	47.0	100.0	36,354	0.13
Rural	31.6	29.8	25.0	11.0	2.6	100.0	56,592	0.21
County								
Mombasa	6.3	11.6	21.9	29.1	31.1	100.0	2,969	0.25
Kwale	58.7	17.6	9.6	11.6	2.5	100.0	1,586	0.42
Kilifi	40.0	21.2	14.6	16.2	8.0	100.0	2,666	0.34
Tana River	60.3	16.2	13.3	7.1	3.1	100.0	721	0.34
Lamu	31.1	21.9	17.7	21.0	8.4	100.0	261	0.24
Taita Taveta	12.8	25.4	24.5	25.8	11.4	100.0	691	0.25
Garissa	53.2	2.2	8.7	25.8	10.1	100.0	852	0.38
Wajir	67.3	5.3	7.7	14.7	5.0	100.0	850	0.40
Mandera	62.4	12.9	13.5	10.2	1.1	100.0	812	0.31
Marsabit	51.0	16.1	14.3	11.0	7.7	100.0	748	0.35
Isiolo	41.7	12.4	22.6	17.7	5.6	100.0	377	0.33
Meru	27.4	22.5	23.9	19.0	7.2	100.0	2,693	0.24
Tharaka-Nithi	34.7	26.5	19.3	12.6	7.0	100.0	704	0.32
Embu	15.0	22.2	26.4	23.1	13.2	100.0	1,035	0.28
Kitui	50.8	26.5	12.5	5.5	4.8	100.0	2,122	0.33
Machakos	5.9	20.1	28.7	28.3	17.1	100.0	2,993	0.30
Makueni	8.0	33.6	34.2	18.2	5.9	100.0	1,665	0.25
Nyandarua	5.7	19.6	43.0	27.6	4.1	100.0	1,060	0.24
Nyeri	1.9	10.4	43.0	35.1	9.6	100.0	1,148	0.17
Kirinyaga	3.5	14.5	39.9	30.7	11.3	100.0	1,210	0.20
Murang'a	5.6	20.7	39.3	23.4	10.9	100.0	1,708	0.21
Kiambu	0.4	3.0	11.0	34.2	51.4	100.0	6,076	0.19
Turkana	76.5	4.7	7.6	7.0	4.2	100.0	1,183	0.46
West Pokot	61.3	18.1	9.9	8.5	2.2	100.0	1,259	0.34
Samburu	68.8	9.1	7.6	8.4	6.1	100.0	554	0.49
Trans Nzoia	14.4	27.2	36.8	15.6	6.0	100.0	1,568	0.30
Uasin Gishu	5.6	17.2	24.5	24.8	27.9	100.0	2,677	0.26
Elgeyo-Marakwet	28.8	32.8	23.7	11.4	3.3	100.0	848	0.28
Nandi	20.0	30.3	29.3	16.3	4.0	100.0	1,550	0.20
Baringo	46.0	20.8	18.9	9.0	5.4	100.0	1,012	0.34
Laikipia	21.7	20.9	27.6	17.8	12.0	100.0	931	0.29
Nakuru	14.5	14.9	19.3	32.9	18.5	100.0	4,659	0.25
Narok	41.3	30.1	15.7	10.0	2.9	100.0	2,265	0.30
Kajiado	21.6	9.9	10.6	19.8	38.0	100.0	1,991	0.32
Kericho	13.4	28.0	36.5	18.9	3.2	100.0	1,777	0.22
Bomet	27.5	38.6	25.2	7.3	1.4	100.0	1,963	0.25
Kakamega	22.1	39.9	21.9	10.8	5.2	100.0	3,162	0.29
Vihiga	11.9	40.3	29.8	14.5	3.5	100.0	1,055	0.21
Bungoma	31.8	37.2	18.0	9.6	3.4	100.0	3,052	0.30
Busia	26.2	31.4	23.4	13.7	5.3	100.0	1,758	0.26
Siaya	16.4	41.8	26.2	11.2	4.4	100.0	1,804	0.21
Kisumu	12.1	26.2	27.3	21.1	13.3	100.0	2,293	0.31
Homa Bay	26.8	38.5	22.6	8.4	3.7	100.0	2,246	0.30
Migori	13.4	31.7	26.6	17.7	10.6	100.0	2,240	0.30
Kisii	21.0	42.2	22.7	10.9	3.2	100.0	2,210	0.34
Nyamira	28.1	37.4	24.1	8.6	1.8	100.0	1,050	0.28
Nairobi City	0.0	0.8	7.7	27.8	63.7	100.0	12,891	0.12
Total	20.0	20.0	20.0	20.0	20.0	100.0	92,946	0.25

[1] The Gini coefficient indicates the level of concentration of wealth, with 0 representing an equal wealth distribution and 1 representing a totally unequal distribution.

Table 2.7 Household population by age, sex, and residence

Percent distribution of the de facto household population by various age groups and percentage of the de facto household population age 10–19, according to sex and residence, Kenya Mini DHS 2025–26

	Urban			Rural			Total		
	Male	Female	Total	Male	Female	Total	Male	Female	Total
Age									
<5	12.7	12.2	12.4	11.8	11.7	11.8	12.1	11.9	12.0
5–9	12.1	11.1	11.6	14.2	12.9	13.5	13.4	12.2	12.8
10–14	9.8	10.0	9.9	14.2	13.7	14.0	12.4	12.3	12.4
15–19	8.8	8.3	8.5	12.4	9.9	11.1	11.0	9.3	10.1
20–24	9.9	11.5	10.7	7.1	6.3	6.7	8.2	8.4	8.3
25–29	11.6	12.4	12.0	5.9	6.4	6.1	8.1	8.7	8.4
30–34	9.2	9.5	9.4	5.3	5.5	5.4	6.8	7.1	7.0
35–39	7.6	7.6	7.6	5.2	6.4	5.8	6.1	6.9	6.5
40–44	6.0	5.0	5.5	4.8	4.6	4.7	5.3	4.8	5.0
45–49	4.1	3.7	3.9	4.2	4.1	4.1	4.1	3.9	4.0
50–54	3.1	3.8	3.5	3.6	4.8	4.2	3.4	4.4	3.9
55–59	2.2	1.5	1.9	2.7	3.1	2.9	2.5	2.5	2.5
60–64	1.3	1.3	1.3	2.7	3.0	2.8	2.1	2.3	2.2
65–69	0.7	0.8	0.8	2.1	2.5	2.3	1.6	1.8	1.7
70–74	0.5	0.5	0.5	1.8	2.1	1.9	1.3	1.5	1.4
75–79	0.2	0.3	0.3	1.0	1.2	1.1	0.7	0.8	0.8
80 +	0.2	0.5	0.3	1.1	1.7	1.4	0.7	1.2	1.0
Don't know/missing	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Dependency age groups									
0–14	34.6	33.3	33.9	40.1	38.4	39.3	38.0	36.4	37.2
15–64	63.7	64.5	64.1	53.7	54.1	53.9	57.6	58.2	57.9
65+	1.6	2.1	1.9	6.1	7.4	6.8	4.3	5.4	4.8
Don't know/missing	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Child and adult population age groups									
0–17	39.9	37.9	38.9	48.0	45.1	46.5	44.8	42.3	43.5
18+	60.0	62.0	61.0	51.9	54.9	53.4	55.1	57.7	56.4
Don't know/missing	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Adolescents 10–19	18.6	18.3	18.4	26.5	23.7	25.1	23.4	21.6	22.5
Number of persons	17,490	17,754	35,243	27,241	27,627	54,867	44,730	45,380	90,110

Table 2.8 Household composition

Percent distribution of households by sex of head of household, household size, and mean size of households, according to residence, Kenya Mini DHS 2025–26

Characteristic	Residence		Total
	Urban	Rural	
Household headship			
Male	69.4	68.5	68.9
Female	30.6	31.5	31.1
Total	100.0	100.0	100.0
Number of usual members			
1	28.9	18.2	23.1
2	17.7	12.7	15.0
3	17.4	15.2	16.2
4	15.7	16.7	16.3
5	10.6	14.1	12.5
6	4.1	9.9	7.3
7	2.6	5.8	4.4
8	1.1	3.5	2.4
9+	1.8	3.8	2.9
Total	100.0	100.0	100.0
Mean size of households	3.0	3.9	3.5
Number of households	11,975	14,405	26,380

Note: Table is based on de jure household members (i.e., usual residents).

Table 2.9 Birth registration of children under age 5

Percentage of de jure children under age 5 whose births are registered with the civil registration authority, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Percentage of children whose births are registered and who:		Total percentage of children whose births are registered	Number of children
	Had a birth certificate	Did not have birth certificate		
Age				
<1	13.8	60.0	73.7	1,948
1–4	43.5	34.6	78.2	8,882
Sex				
Male	39.3	37.5	76.8	5,431
Female	37.0	40.9	77.9	5,399
Residence				
Urban	49.8	32.8	82.6	4,431
Rural	30.2	43.6	73.8	6,400
Wealth quintile				
Lowest	19.4	42.7	62.1	2,517
Second	28.8	46.5	75.3	2,081
Middle	39.0	40.3	79.3	1,948
Fourth	43.3	40.5	83.8	2,209
Highest	64.1	25.2	89.4	2,075
Total	38.2	39.2	77.4	10,830

Table 2.9C Birth registration of children under age 5: County level

Percentage of de jure children under age 5 whose births are registered with the civil registration authority, according to county, Kenya Mini DHS 2025–26

County	Percentage of children whose births are registered and who:		Total percentage of children whose births are registered	Number of children
	Had a birth certificate	Did not have birth certificate		
Mombasa	42.7	11.9	54.6	345
Kwale	32.0	55.4	87.5	202
Kilifi	40.9	25.4	66.3	302
Tana River	31.4	33.6	65.0	106
Lamu	60.2	26.1	86.3	31
Taita Taveta	57.7	16.9	74.6	81
Garissa	40.2	8.0	48.3	123
Wajir	41.5	11.3	52.9	118
Mandera	38.5	5.7	44.2	135
Marsabit	46.3	25.1	71.3	105
Isiolo	40.2	42.4	82.6	46
Meru	34.0	26.6	60.7	270
Tharaka-Nithi	45.4	41.9	87.3	69
Embu	47.9	47.9	95.8	81
Kitui	23.0	56.1	79.1	208
Machakos	32.6	61.0	93.6	354
Makueni	40.4	25.6	66.0	167
Nyandarua	43.8	48.1	91.9	99
Nyeri	51.7	42.5	94.2	82
Kirinyaga	56.9	30.4	87.2	100
Murang'a	46.6	47.5	94.1	150
Kiambu	48.3	43.9	92.1	652
Turkana	16.0	53.1	69.1	187
West Pokot	12.3	34.7	46.9	201
Samburu	12.9	49.1	62.0	85
Trans Nzoia	31.8	42.2	74.0	173
Uasin Gishu	34.5	57.4	91.9	307
Elgeyo-Marakwet	42.9	44.7	87.7	102
Nandi	33.1	38.3	71.4	176
Baringo	26.8	51.6	78.3	128
Laikipia	39.3	26.3	65.5	105
Nakuru	39.3	35.1	74.4	572
Narok	25.3	24.9	50.1	254
Kajiado	37.2	56.4	93.6	289
Kericho	30.5	26.0	56.5	173
Bomet	30.0	47.2	77.2	224
Kakamega	40.4	50.1	90.5	360
Vihiga	44.0	44.0	88.0	120
Bungoma	21.2	54.7	75.9	368
Busia	37.7	29.9	67.6	204
Siaya	27.0	22.6	49.6	195
Kisumu	40.0	42.8	82.8	258
Homa Bay	28.9	61.1	90.0	265
Migori	23.1	66.9	90.0	335
Kisii	31.4	55.9	87.3	237
Nyamira	32.6	59.5	92.1	106
Nairobi City	55.1	26.5	81.7	1,582
Total	38.2	39.2	77.4	10,830

Table 2.10.1 Educational attainment of the female household population (standard DHS education categories)

Percent distribution of the de facto female household population age 6 and older by highest level of schooling attended or completed and median years completed, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	No education[1]	Some primary	Completed primary[2]	Some secondary[3]	Completed secondary[4]	More than secondary[5]	Total	Number	Median years completed
Age									
6–9	23.7	76.1	0.0	0.0	0.0	0.0	100.0	4,414	1.2
10–14	3.7	39.5	20.2	36.2	0.4	0.0	100.0	5,577	5.4
15–19	2.2	3.1	4.6	66.6	21.5	1.9	100.0	4,119	8.2
20–24	5.4	5.9	11.0	17.9	48.8	10.8	100.0	3,131	9.2
25–29	4.9	10.5	19.4	15.5	37.5	11.5	100.0	3,174	9.0
30–34	7.7	16.0	23.5	13.8	28.8	9.8	100.0	2,637	7.9
35–39	8.7	19.0	33.6	11.3	21.9	4.8	100.0	2,767	7.6
40–44	10.9	24.3	31.6	10.6	17.9	3.8	100.0	1,858	7.5
45–49	9.5	26.6	30.4	10.7	19.3	2.9	100.0	1,589	7.4
50–54	13.4	24.6	31.9	8.7	17.5	3.6	100.0	1,850	7.3
55–59	16.8	20.2	38.3	10.1	11.4	2.7	100.0	1,048	7.0
60–64	22.0	21.6	29.0	11.4	13.6	2.3	100.0	998	6.6
65+	47.9	24.9	15.6	3.7	6.4	0.7	100.0	2,372	5.9
Residence									
Urban	8.2	21.9	16.4	18.8	25.8	8.2	100.0	13,110	7.7
Rural	14.1	29.5	19.6	21.6	13.3	1.6	100.0	22,455	7.0
Wealth quintile									
Lowest	28.1	33.5	14.8	18.1	5.3	0.1	100.0	7,364	6.1
Second	10.2	31.3	21.8	23.9	11.9	0.6	100.0	7,528	6.9
Middle	7.6	26.3	23.0	23.4	17.9	1.5	100.0	7,337	7.3
Fourth	5.5	22.6	20.0	20.2	27.3	4.0	100.0	6,871	7.6
Highest	7.5	18.6	11.7	16.9	29.3	15.3	100.0	6,464	8.5
Total	12.0	26.7	18.4	20.6	17.9	4.0	100.0	35,565	7.2

Notes: Respondents whose current or previous level of education was vocational training have been excluded from this table. Total includes 31 women who were missing information on age who are not shown separately.

[1] No education includes informal education (e.g., madrassa/duksi/adult education).

[2] Completed grade 8 at the primary level for those under age 56; because of the change in the school system in the 1980s, those age 56 and older are considered to have completed primary if they completed grade 7.

[3] Those who have completed junior school are considered to have completed some secondary education if it's the highest level attained at the time of interview.

[4] Four years of secondary is considered completed secondary for purposes of the Demographic and Health Surveys indicator.

[5] More than secondary includes middle-level colleges and universities.

Table 2.10.1C Educational attainment of the female household population (standard DHS education categories): County level

Percent distribution of the de facto female household population age 6 and older by highest level of schooling attended or completed and median years completed, according to county, Kenya Mini DHS 2025–26

County	No education[1]	Some primary	Completed primary[2]	Some secondary[3]	Completed secondary[4]	More than secondary[5]	Total	Number	Median years completed
Mombasa	9.0	27.8	21.2	17.3	20.3	3.9	100.0	1,148	7.2
Kwale	22.8	31.8	18.3	17.2	8.1	1.4	100.0	653	6.4
Kilifi	22.7	26.8	19.1	18.0	11.8	1.5	100.0	1,086	6.9
Tana River	41.2	23.9	10.2	15.3	8.3	1.0	100.0	267	6.1
Lamu	23.3	25.0	19.5	17.6	14.2	0.4	100.0	103	7.0
Taita Taveta	8.7	25.8	28.5	17.2	16.2	3.5	100.0	269	7.3
Garissa	61.6	18.9	5.3	8.2	4.4	1.6	100.0	341	5.4
Wajir	59.3	19.0	4.0	10.9	5.2	1.6	100.0	289	5.1
Mandera	59.8	23.3	5.3	7.1	4.0	0.4	100.0	282	4.1
Marsabit	37.5	25.0	9.5	17.8	8.4	1.7	100.0	273	6.1
Isiolo	25.9	31.9	14.5	14.4	11.3	1.8	100.0	149	6.4
Meru	13.1	30.9	20.3	18.6	15.4	1.8	100.0	1,008	7.1
Tharaka-Nithi	10.2	31.3	18.5	21.7	15.3	2.9	100.0	291	7.1
Embu	6.8	28.2	20.0	20.5	19.9	4.0	100.0	406	7.3
Kitui	12.6	26.4	23.4	25.5	10.4	1.7	100.0	846	7.1
Machakos	5.3	20.2	23.9	23.5	23.9	3.0	100.0	1,095	7.5
Makueni	7.5	25.4	22.0	22.6	18.7	3.7	100.0	681	7.3
Nyandarua	7.1	26.0	24.7	21.5	17.5	2.4	100.0	430	7.2
Nyeri	6.4	21.4	24.1	16.6	28.7	2.3	100.0	454	7.6
Kirinyaga	9.4	21.7	26.6	16.3	23.5	2.2	100.0	486	7.4
Murang'a	5.6	29.5	24.7	20.8	17.7	1.8	100.0	739	7.1
Kiambu	4.4	18.5	15.7	20.2	27.4	13.2	100.0	2,246	8.0
Turkana	59.6	22.6	5.6	7.9	3.5	0.8	100.0	470	4.0
West Pokot	30.0	36.7	6.6	16.3	7.7	2.6	100.0	457	5.5
Samburu	47.1	24.5	7.3	14.2	6.0	1.0	100.0	206	5.3
Trans Nzoia	8.4	30.2	20.5	22.7	15.0	3.0	100.0	640	7.2
Uasin Gishu	4.9	24.3	17.7	22.7	23.4	6.6	100.0	916	7.6
Elgeyo-Marakwet	7.6	28.0	19.2	25.8	16.0	3.4	100.0	321	7.1
Nandi	4.9	32.4	19.6	23.3	16.7	2.4	100.0	601	7.2
Baringo	17.7	25.4	18.0	22.1	14.1	2.3	100.0	377	7.1
Laikipia	14.7	20.9	21.7	19.6	19.7	2.4	100.0	362	7.4
Nakuru	6.5	27.4	18.5	23.8	20.0	3.3	100.0	1,766	7.3
Narok	17.5	33.8	17.5	17.0	12.4	1.7	100.0	847	6.6
Kajiado	15.3	23.2	13.8	15.0	22.8	8.3	100.0	707	7.6
Kericho	6.6	28.3	19.3	25.4	16.6	3.7	100.0	638	7.3
Bomet	6.1	31.8	17.7	25.3	17.0	2.1	100.0	775	7.2
Kakamega	8.6	30.6	19.7	23.8	14.3	2.3	100.0	1,236	7.0
Vihiga	5.3	26.0	22.3	28.1	15.4	1.2	100.0	436	7.2
Bungoma	7.5	35.4	18.5	24.8	11.8	2.1	100.0	1,202	6.8
Busia	9.4	31.2	16.9	23.7	16.6	1.3	100.0	699	7.0
Siaya	10.3	31.5	24.5	21.6	11.2	0.8	100.0	791	6.8
Kisumu	6.9	29.3	16.9	26.4	16.4	3.3	100.0	875	7.1
Homa Bay	6.4	30.4	21.4	28.9	10.3	2.5	100.0	882	6.9
Migori	7.5	32.6	22.5	19.5	15.1	2.6	100.0	844	7.0
Kisii	7.6	31.2	17.3	25.9	16.1	1.7	100.0	901	7.0
Nyamira	5.3	26.5	18.7	29.0	17.4	2.5	100.0	459	7.1
Nairobi City	8.0	21.7	16.6	17.0	27.8	8.2	100.0	4,616	7.7
Total	12.0	26.7	18.4	20.6	17.9	4.0	100.0	35,565	7.2

Note: Respondents whose current or previous level of education was vocational training have been excluded from this table.
[1] No education includes informal education (e.g., madrassa/duksi/adult education).
[2] Completed grade 8 at the primary level for those under age 56; because of the change in the school system in the 1980s, those age 56 and older are considered to have completed primary if they completed grade 7.
[3] Those who have completed junior school are considered to have completed some secondary education if it's the highest level attained at the time of interview.
[4] Four years of secondary is considered completed secondary for purposes of the Demographic and Health Surveys indicator.
[5] More than secondary includes middle-level colleges and universities.

Table 2.10.2 Educational attainment of the female household population (CBE)

Percent distribution of the de facto female household population age 6 and older by highest level of schooling attended or completed and median years completed, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	No education [1]	Some primary	Completed primary[2]	Some junior	Completed junior	Some secondary	Completed secondary[3]	More than secondary[4]	Total	Number	Median years completed
Age											
6–9	23.7	76.1	0.0	0.0	0.0	0.0	0.0	0.0	100.0	4,414	1.2
10–14	3.7	39.5	20.2	29.7	4.9	1.5	0.4	0.0	100.0	5,577	5.4
15–19	2.2	3.1	4.6	11.1	13.7	41.8	21.5	1.9	100.0	4,119	8.2
20–24	5.4	5.9	11.0	0.0	0.0	17.9	48.8	10.8	100.0	3,131	9.2
25–29	4.9	10.5	19.4	0.0	0.0	15.5	37.5	11.5	100.0	3,174	9.0
30–34	7.7	16.0	23.5	0.0	0.0	13.8	28.8	9.8	100.0	2,637	7.9
35–39	8.7	19.0	33.6	0.0	0.0	11.3	21.9	4.8	100.0	2,767	7.6
40–44	10.9	24.3	31.6	0.0	0.0	10.6	17.9	3.8	100.0	1,858	7.5
45–49	9.5	26.6	30.4	0.0	0.0	10.7	19.3	2.9	100.0	1,589	7.4
50–54	13.4	24.6	31.9	0.0	0.0	8.7	17.5	3.6	100.0	1,850	7.3
55–59	16.8	20.2	38.3	0.0	0.0	10.1	11.4	2.7	100.0	1,048	7.0
60–64	22.0	21.6	29.0	0.0	0.0	11.4	13.6	2.3	100.0	998	6.6
65+	47.9	24.9	15.6	0.0	0.0	3.7	6.4	0.7	100.0	2,372	5.9
Residence											
Urban	8.2	21.9	16.4	4.7	2.0	12.1	25.8	8.2	100.0	13,110	7.7
Rural	14.1	29.5	19.6	6.6	2.6	12.4	13.3	1.6	100.0	22,455	7.0
Wealth quintile											
Lowest	28.1	33.5	14.8	6.8	2.4	8.9	5.3	0.1	100.0	7,364	6.1
Second	10.2	31.3	21.8	7.4	2.6	13.8	11.9	0.6	100.0	7,528	6.9
Middle	7.6	26.3	23.0	5.6	2.7	15.1	17.9	1.5	100.0	7,337	7.3
Fourth	5.5	22.6	20.0	4.7	2.2	13.4	27.3	4.0	100.0	6,871	7.6
Highest	7.5	18.6	11.7	5.0	1.8	10.1	29.3	15.3	100.0	6,464	8.5
Total	12.0	26.7	18.4	5.9	2.4	12.3	17.9	4.0	100.0	35,565	7.2

Notes: Respondents whose current or previous level of education was vocational training have been excluded from this table. Total includes 31 women who were missing information on age who are not shown separately.

[1] No education includes informal education (e.g., madrassa/duksi/adult education).

[2] Completed grade 8 at the primary level for those under age 56; because of the change in the school system in the 1980s, those age 56 and older are considered to have completed primary if they completed grade 7.

[3] Four years of secondary is considered completed secondary for purposes of the Demographic and Health Surveys indicator.

[4] More than secondary includes middle-level colleges and universities.

Table 2.10.2C Educational attainment of the female household population (CBE): County level

Percent distribution of the de facto female household population age 6 and older by highest level of schooling attended or completed and median years completed, according to county, Kenya Mini DHS 2025–26

County	No education [1]	Some primary	Completed primary[2]	Some junior	Completed junior	Some secondary	Completed secondary[3]	More than secondary[4]	Total	Number	Median years completed
Mombasa	9.0	27.8	21.2	6.4	1.6	9.3	20.3	3.9	100.0	1,148	7.2
Kwale	22.8	31.8	18.3	7.1	2.4	7.8	8.1	1.4	100.0	653	6.4
Kilifi	22.7	26.8	19.1	8.9	3.4	5.7	11.8	1.5	100.0	1,086	6.9
Tana River	41.2	23.9	10.2	6.9	2.4	6.0	8.3	1.0	100.0	267	6.1
Lamu	23.3	25.0	19.5	5.6	1.7	10.3	14.2	0.4	100.0	103	7.0
Taita Taveta	8.7	25.8	28.5	3.9	2.3	11.1	16.2	3.5	100.0	269	7.3
Garissa	61.6	18.9	5.3	3.7	1.4	3.0	4.4	1.6	100.0	341	5.4
Wajir	59.3	19.0	4.0	4.2	2.2	4.5	5.2	1.6	100.0	289	5.1
Mandera	59.8	23.3	5.3	3.6	1.4	2.1	4.0	0.4	100.0	282	4.1
Marsabit	37.5	25.0	9.5	7.0	2.4	8.4	8.4	1.7	100.0	273	6.1
Isiolo	25.9	31.9	14.5	5.2	2.3	6.9	11.3	1.8	100.0	149	6.4
Meru	13.1	30.9	20.3	6.1	1.4	11.1	15.4	1.8	100.0	1,008	7.1
Tharaka-Nithi	10.2	31.3	18.5	6.1	2.8	12.9	15.3	2.9	100.0	291	7.1
Embu	6.8	28.2	20.0	4.5	2.2	13.9	19.9	4.0	100.0	406	7.3
Kitui	12.6	26.4	23.4	10.3	2.6	12.6	10.4	1.7	100.0	846	7.1
Machakos	5.3	20.2	23.9	4.6	1.9	16.9	23.9	3.0	100.0	1,095	7.5
Makueni	7.5	25.4	22.0	5.0	2.1	15.5	18.7	3.7	100.0	681	7.3
Nyandarua	7.1	26.0	24.7	5.7	3.0	12.8	17.5	2.4	100.0	430	7.2
Nyeri	6.4	21.4	24.1	3.0	1.1	12.5	28.7	2.3	100.0	454	7.6
Kirinyaga	9.4	21.7	26.6	2.5	2.0	11.8	23.5	2.2	100.0	486	7.4
Murang'a	5.6	29.5	24.7	4.8	2.4	13.6	17.7	1.8	100.0	739	7.1
Kiambu	4.4	18.5	15.7	6.0	2.4	11.8	27.4	13.2	100.0	2,246	8.0
Turkana	59.6	22.6	5.6	3.3	0.8	3.7	3.5	0.8	100.0	470	4.0
West Pokot	30.0	36.7	6.6	6.1	1.9	8.3	7.7	2.6	100.0	457	5.5
Samburu	47.1	24.5	7.3	5.0	2.2	7.0	6.0	1.0	100.0	206	5.3
Trans Nzoia	8.4	30.2	20.5	4.9	2.6	15.2	15.0	3.0	100.0	640	7.2
Uasin Gishu	4.9	24.3	17.7	6.3	3.0	13.3	23.4	6.6	100.0	916	7.6
Elgeyo-Marakwet	7.6	28.0	19.2	7.6	3.3	14.9	16.0	3.4	100.0	321	7.1
Nandi	4.9	32.4	19.6	6.4	4.0	12.9	16.7	2.4	100.0	601	7.2
Baringo	17.7	25.4	18.0	7.6	4.1	10.5	14.1	2.3	100.0	377	7.1
Laikipia	14.7	20.9	21.7	6.9	2.0	10.6	19.7	2.4	100.0	362	7.4
Nakuru	6.5	27.4	18.5	5.1	2.2	16.5	20.0	3.3	100.0	1,766	7.3
Narok	17.5	33.8	17.5	5.6	1.9	9.5	12.4	1.7	100.0	847	6.6
Kajiado	15.3	23.2	13.8	4.3	2.2	8.6	22.8	8.3	100.0	707	7.6
Kericho	6.6	28.3	19.3	5.0	3.0	17.4	16.6	3.7	100.0	638	7.3
Bomet	6.1	31.8	17.7	6.9	3.8	14.7	17.0	2.1	100.0	775	7.2
Kakamega	8.6	30.6	19.7	7.2	2.2	14.4	14.3	2.3	100.0	1,236	7.0
Vihiga	5.3	26.0	22.3	7.7	3.5	16.9	15.4	1.2	100.0	436	7.2
Bungoma	7.5	35.4	18.5	6.0	3.8	15.0	11.8	2.1	100.0	1,202	6.8
Busia	9.4	31.2	16.9	8.7	2.5	12.6	16.6	1.3	100.0	699	7.0
Siaya	10.3	31.5	24.5	6.3	3.0	12.3	11.2	0.8	100.0	791	6.8
Kisumu	6.9	29.3	16.9	9.9	2.7	13.8	16.4	3.3	100.0	875	7.1
Homa Bay	6.4	30.4	21.4	12.1	3.1	13.7	10.3	2.5	100.0	882	6.9
Migori	7.5	32.6	22.5	5.1	1.6	12.8	15.1	2.6	100.0	844	7.0
Kisii	7.6	31.2	17.3	7.8	2.7	15.4	16.1	1.7	100.0	901	7.0
Nyamira	5.3	26.5	18.7	6.4	3.5	19.2	17.4	2.5	100.0	459	7.1
Nairobi City	8.0	21.7	16.6	3.4	1.5	12.1	27.8	8.2	100.0	4,616	7.7
Total	12.0	26.7	18.4	5.9	2.4	12.3	17.9	4.0	100.0	35,565	7.2

Note: Respondents whose current or previous level of education was vocational training have been excluded from this table.
[1] No education includes informal education (e.g., madrassa/duksi/adult education).
[2] Completed grade 8 at the primary level for those under age 56; because of the change in the school system in the 1980s, those age 56 and older are considered to have completed primary if they completed grade 7.
[3] Four years of secondary is considered completed secondary for purposes of the Demographic and Health Surveys indicator.
[4] More than secondary includes middle-level colleges and universities.

Table 2.10.3 Educational attainment of the male household population (standard DHS education categories)

Percent distribution of the de facto male household population age 6 and older by highest level of schooling attended or completed and median years completed, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	No education[1]	Some primary	Completed primary[2]	Some secondary[3]	Completed secondary[4]	More than secondary[5]	Don't know	Total	Number	Median years completed
Age										
6–9	25.8	74.1	0.0	0.0	0.0	0.0	0.1	100.0	4,772	1.0
10–14	3.7	43.5	20.7	31.8	0.3	0.0	0.0	100.0	5,566	5.2
15–19	2.6	6.2	6.8	65.2	17.7	1.4	0.0	100.0	4,834	8.0
20–24	3.1	6.7	10.9	15.7	47.5	16.0	0.2	100.0	3,130	9.3
25–29	3.2	9.6	15.1	10.4	43.0	18.1	0.6	100.0	2,947	9.3
30–34	3.9	12.0	20.5	9.5	37.5	15.5	1.1	100.0	2,517	9.1
35–39	4.1	16.8	27.1	10.2	28.9	12.4	0.6	100.0	2,357	7.9
40–44	6.8	19.5	29.7	8.0	26.2	8.6	1.2	100.0	2,079	7.7
45–49	6.3	22.5	31.5	8.7	23.6	6.3	1.2	100.0	1,649	7.6
50–54	6.6	18.3	28.5	8.6	29.9	6.4	1.7	100.0	1,385	7.8
55–59	9.6	13.9	31.8	6.5	29.3	8.1	0.9	100.0	1,008	7.7
60–64	9.2	15.4	31.1	8.6	29.4	5.6	0.8	100.0	870	7.3
65+	21.5	22.2	28.4	7.5	16.4	3.1	0.9	100.0	1,850	6.7
Residence										
Urban	5.7	20.4	14.7	15.8	29.7	13.1	0.6	100.0	13,028	8.1
Rural	9.7	29.6	18.9	22.1	16.3	2.9	0.4	100.0	21,976	7.1
Wealth quintile										
Lowest	20.3	35.3	15.9	18.5	9.1	0.6	0.3	100.0	7,234	6.1
Second	6.8	30.3	20.7	24.5	16.0	1.3	0.4	100.0	7,471	7.1
Middle	4.2	25.4	21.7	22.9	22.5	2.7	0.5	100.0	7,199	7.4
Fourth	3.4	19.6	17.4	18.5	32.7	7.8	0.7	100.0	6,742	7.9
Highest	5.6	18.9	10.3	13.2	27.8	23.4	0.7	100.0	6,359	9.1
Total	8.2	26.2	17.4	19.7	21.3	6.7	0.5	100.0	35,004	7.4

Notes: Respondents whose current or previous level of education was vocational training have been excluded from this table. Total includes 40 men who were missing information on age who are not shown separately.

[1] No education includes informal education (e.g., madrassa/duksi/adult education).

[2] Completed grade 8 at the primary level for those under age 56; because of the change in the school system in the 1980s, those age 56 and older are considered to have completed primary if they completed grade 7.

[3] Those who have completed junior school are considered to have completed some secondary education if it's the highest level attained at the time of interview.

[4] Four years of secondary is considered completed secondary for purposes of the Demographic and Health Surveys indicator.

[5] More than secondary includes middle-level colleges and universities.

Table 2.10.3C Educational attainment of the male household population (standard DHS education categories): County level

Percent distribution of the de facto male household population age 6 and older by highest level of schooling attended or completed and median years completed, according to county, Kenya Mini DHS 2025–26

County	No education[1]	Some primary	Completed primary[2]	Some secondary[3]	Completed secondary[4]	More than secondary[5]	Total	Number	Median years completed
Mombasa	5.3	24.5	19.2	19.1	23.8	7.5	100.0	1,064	7.5
Kwale	12.4	34.0	22.3	18.5	11.2	0.9	100.0	600	6.5
Kilifi	9.3	31.9	22.2	18.1	15.7	2.9	100.0	1,021	7.0
Tana River	28.9	26.4	11.9	17.7	12.1	3.0	100.0	261	6.5
Lamu	13.5	25.1	21.2	20.1	18.3	1.8	100.0	103	7.2
Taita Taveta	6.2	23.0	25.1	21.6	19.8	4.3	100.0	259	7.4
Garissa	52.4	21.1	5.5	9.5	8.8	2.6	100.0	316	6.2
Wajir	43.8	22.7	6.8	13.5	9.3	3.9	100.0	324	5.9
Mandera	42.9	26.1	7.2	15.1	6.4	2.3	100.0	324	5.3
Marsabit	27.4	29.6	10.0	19.2	9.7	3.9	100.0	233	6.4
Isiolo	17.7	28.4	13.6	18.8	18.5	2.8	100.0	144	6.9
Meru	8.0	32.2	18.8	19.4	18.7	2.4	100.0	1,105	7.1
Tharaka-Nithi	6.5	35.8	18.3	18.7	14.6	5.2	100.0	251	6.8
Embu	4.1	29.3	19.8	21.8	18.2	5.7	100.0	416	7.3
Kitui	6.8	28.5	20.7	24.3	16.0	3.5	100.0	849	7.2
Machakos	2.1	19.1	18.9	18.8	35.2	5.6	100.0	1,189	7.9
Makueni	2.1	27.6	20.0	24.9	19.8	5.3	100.0	637	7.3
Nyandarua	4.0	25.0	24.7	20.8	20.8	3.1	100.0	434	7.3
Nyeri	4.0	18.9	21.5	19.5	29.6	6.3	100.0	434	7.7
Kirinyaga	5.2	23.3	24.7	17.1	24.2	4.6	100.0	460	7.5
Murang'a	3.2	27.0	24.8	20.4	21.0	3.6	100.0	719	7.3
Kiambu	3.0	15.5	16.5	14.8	28.0	20.8	100.0	2,288	9.0
Turkana	50.4	23.5	7.3	9.8	6.6	2.4	100.0	397	5.2
West Pokot	23.8	37.4	7.0	18.5	10.6	2.6	100.0	454	5.6
Samburu	41.2	24.0	8.2	15.4	7.5	3.6	100.0	202	5.6
Trans Nzoia	7.3	27.9	19.1	21.5	18.3	4.1	100.0	618	7.1
Uasin Gishu	3.2	24.1	14.6	22.0	26.4	9.2	100.0	955	7.7
Elgeyo-Marakwet	6.4	27.4	17.4	23.1	20.7	5.0	100.0	325	7.3
Nandi	4.3	32.6	18.1	24.7	16.2	3.6	100.0	594	7.2
Baringo	12.0	27.4	14.9	18.2	21.5	4.8	100.0	394	7.4
Laikipia	12.4	18.8	19.8	20.0	22.6	5.2	100.0	350	7.5
Nakuru	3.7	23.3	19.2	22.5	25.4	5.2	100.0	1,725	7.6
Narok	12.9	33.5	17.6	16.1	15.9	3.6	100.0	946	6.8
Kajiado	11.4	24.4	11.4	15.4	22.0	14.6	100.0	747	7.7
Kericho	4.3	28.7	15.6	24.6	21.1	5.6	100.0	675	7.4
Bomet	4.1	31.4	15.4	26.7	17.8	4.4	100.0	738	7.2
Kakamega	6.4	31.1	18.8	23.4	16.1	3.8	100.0	1,250	6.9
Vihiga	5.0	32.9	21.6	21.7	15.7	2.3	100.0	376	6.9
Bungoma	5.3	35.5	16.5	23.5	15.6	3.4	100.0	1,208	6.8
Busia	6.0	33.7	17.7	25.6	12.5	4.4	100.0	669	7.0
Siaya	4.9	30.2	23.1	22.9	16.5	2.2	100.0	684	7.1
Kisumu	6.0	26.6	16.4	27.0	17.1	6.2	100.0	853	7.3
Homa Bay	5.6	27.3	20.1	29.1	13.7	4.2	100.0	858	7.1
Migori	5.3	31.0	21.5	22.0	14.1	4.8	100.0	799	7.0
Kisii	4.9	26.4	18.4	26.0	20.2	3.9	100.0	820	7.1
Nyamira	3.3	26.5	16.6	28.4	21.6	3.2	100.0	390	7.3
Nairobi City	5.8	20.5	14.1	12.4	33.9	12.8	100.0	4,548	8.6
Total	8.2	26.2	17.4	19.7	21.3	6.7	100.0	35,004	7.4

Note: Respondents whose current or previous level of education was vocational training have been excluded from this table.
[1] No education includes informal education (e.g., madrassa/duksi/adult education).
[2] Completed grade 8 at the primary level for those under age 56; because of the change in the school system in the 1980s, those age 56 and older are considered to have completed primary if they completed grade 7.
[3] Those who have completed junior school are considered to have completed some secondary education if it's the highest level attained at the time of interview.
[4] Four years of secondary is considered completed secondary for purposes of the Demographic and Health Surveys indicator.
[5] More than secondary includes middle-level colleges and universities.

Table 2.10.4 Educational attainment of the male household population (CBE)

Percent distribution of the de facto male household population age 6 and older by highest level of schooling attended or completed and median years completed, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	No education [1]	Some primary	Completed primary[2]	Some junior	Completed junior	Some secondary	Completed secondary[3]	More than secondary[4]	Total	Number	Median years completed
Age											
6–9	25.8	74.1	0.0	0.0	0.0	0.0	0.0	0.0	100.0	4,772	1.0
10–14	3.7	43.5	20.7	26.5	4.3	0.9	0.3	0.0	100.0	5,566	5.2
15–19	2.6	6.2	6.8	14.6	14.4	36.2	17.7	1.4	100.0	4,834	8.0
20–24	3.1	6.7	10.9	0.0	0.0	15.7	47.5	16.0	100.0	3,130	9.3
25–29	3.2	9.6	15.1	0.0	0.0	10.4	43.0	18.1	100.0	2,947	9.3
30–34	3.9	12.0	20.5	0.0	0.0	9.5	37.5	15.5	100.0	2,517	9.1
35–39	4.1	16.8	27.1	0.0	0.0	10.2	28.9	12.4	100.0	2,357	7.9
40–44	6.8	19.5	29.7	0.0	0.0	8.0	26.2	8.6	100.0	2,079	7.7
45–49	6.3	22.5	31.5	0.0	0.0	8.7	23.6	6.3	100.0	1,649	7.6
50–54	6.6	18.3	28.5	0.0	0.0	8.6	29.9	6.4	100.0	1,385	7.8
55–59	9.6	13.9	31.8	0.0	0.0	6.5	29.3	8.1	100.0	1,008	7.7
60–64	9.2	15.4	31.1	0.0	0.0	8.6	29.4	5.6	100.0	870	7.3
65+	21.5	22.2	28.4	0.0	0.0	7.5	16.4	3.1	100.0	1,850	6.7
	(12.4)	(19.9)	(26.1)	(1.4)	(0.0)	(0.0)	(16.2)	(10.3)	100.0	40	(7.5)
Residence											
Urban											
Rural	5.7	20.4	14.7	4.6	1.7	9.5	29.7	13.1	100.0	13,028	8.1
	9.7	29.6	18.9	7.2	3.3	11.6	16.3	2.9	100.0	21,976	7.1
Wealth quintile											
Lowest											
Second	20.3	35.3	15.9	7.1	3.0	8.4	9.1	0.6	100.0	7,234	6.1
Middle	6.8	30.3	20.7	7.8	3.4	13.2	16.0	1.3	100.0	7,471	7.1
Fourth	4.2	25.4	21.7	6.6	3.0	13.3	22.5	2.7	100.0	7,199	7.4
Highest	3.4	19.6	17.4	5.0	2.5	11.0	32.7	7.8	100.0	6,742	7.9
	5.6	18.9	10.3	4.2	1.3	7.7	27.8	23.4	100.0	6,359	9.1
Total											

Notes: Respondents whose current or previous level of education was vocational training have been excluded from this table. Total includes 40 men who were missing information on age who are not shown separately.

[1] No education includes informal education (e.g., madrassa/duksi/adult education).

[2] Completed grade 8 at the primary level for those under age 56; because of the change in the school system in the 1980s, those age 56 and older are considered to have completed primary if they completed grade 7.

[3] Four years of secondary is considered completed secondary for purposes of the Demographic and Health Surveys indicator.

[4] More than secondary includes middle-level colleges and universities.

Table 2.10.4C Educational attainment of the male household population (CBE): County level

Percent distribution of the de facto male household population age 6 and older by highest level of schooling attended or completed and median years completed, according to county, Kenya Mini DHS 2025–26

County	No education [1]	Some primary	Completed primary[2]	Some junior	Completed junior	Some secondary	Completed secondary[3]	More than secondary[4]	Total	Number	Median years completed
Mombasa	5.3	24.5	19.2	6.7	1.4	11.0	23.8	7.5	0.7	100.0	1,064
Kwale	12.4	34.0	22.3	7.4	3.1	8.0	11.2	0.9	0.7	100.0	600
Kilifi	9.3	31.9	22.2	7.6	2.8	7.7	15.7	2.9	0.0	100.0	1,021
Tana River	28.9	26.4	11.9	6.4	2.2	9.1	12.1	3.0	0.0	100.0	261
Lamu	13.5	25.1	21.2	5.3	2.6	12.2	18.3	1.8	0.0	100.0	103
Taita Taveta	6.2	23.0	25.1	6.0	3.3	12.3	19.8	4.3	0.0	100.0	259
Garissa	52.4	21.1	5.5	2.8	1.6	5.1	8.8	2.6	0.1	100.0	316
Wajir	43.8	22.7	6.8	3.5	1.8	8.1	9.3	3.9	0.0	100.0	324
Mandera	42.9	26.1	7.2	5.6	1.9	7.6	6.4	2.3	0.0	100.0	324
Marsabit	27.4	29.6	10.0	8.0	4.1	7.1	9.7	3.9	0.2	100.0	233
Isiolo	17.7	28.4	13.6	7.7	2.6	8.6	18.5	2.8	0.2	100.0	144
Meru	8.0	32.2	18.8	5.5	2.3	11.6	18.7	2.4	0.4	100.0	1,105
Tharaka-Nithi	6.5	35.8	18.3	3.7	1.5	13.5	14.6	5.2	1.0	100.0	251
Embu	4.1	29.3	19.8	5.6	3.5	12.7	18.2	5.7	1.0	100.0	416
Kitui	6.8	28.5	20.7	8.9	3.8	11.6	16.0	3.5	0.2	100.0	849
Machakos	2.1	19.1	18.9	4.6	2.2	12.0	35.2	5.6	0.3	100.0	1,189
Makueni	2.1	27.6	20.0	6.5	2.6	15.8	19.8	5.3	0.3	100.0	637
Nyandarua	4.0	25.0	24.7	6.3	3.1	11.3	20.8	3.1	1.7	100.0	434
Nyeri	4.0	18.9	21.5	6.1	2.4	11.0	29.6	6.3	0.2	100.0	434
Kirinyaga	5.2	23.3	24.7	3.8	2.3	11.0	24.2	4.6	1.0	100.0	460
Murang'a	3.2	27.0	24.8	5.1	2.9	12.4	21.0	3.6	0.0	100.0	719
Kiambu	3.0	15.5	16.5	4.3	1.4	9.0	28.0	20.8	1.4	100.0	2,288
Turkana	50.4	23.5	7.3	4.5	1.5	3.7	6.6	2.4	0.0	100.0	397
West Pokot	23.8	37.4	7.0	6.1	3.0	9.3	10.6	2.6	0.0	100.0	454
Samburu	41.2	24.0	8.2	4.6	2.2	8.7	7.5	3.6	0.1	100.0	202
Trans Nzoia	7.3	27.9	19.1	5.7	2.9	12.9	18.3	4.1	1.8	100.0	618
Uasin Gishu	3.2	24.1	14.6	5.5	2.9	13.6	26.4	9.2	0.6	100.0	955
Elgeyo-Marakwet	6.4	27.4	17.4	6.7	3.9	12.5	20.7	5.0	0.0	100.0	325
Nandi	4.3	32.6	18.1	7.7	4.1	12.8	16.2	3.6	0.5	100.0	594
Baringo	12.0	27.4	14.9	5.7	3.2	9.3	21.5	4.8	1.1	100.0	394
Laikipia	12.4	18.8	19.8	9.0	1.8	9.3	22.6	5.2	1.2	100.0	350
Nakuru	3.7	23.3	19.2	6.9	3.3	12.3	25.4	5.2	0.7	100.0	1,725
Narok	12.9	33.5	17.6	4.7	3.3	8.1	15.9	3.6	0.3	100.0	946
Kajiado	11.4	24.4	11.4	5.1	1.8	8.5	22.0	14.6	0.8	100.0	747
Kericho	4.3	28.7	15.6	6.7	4.0	13.9	21.1	5.6	0.2	100.0	675
Bomet	4.1	31.4	15.4	7.1	4.3	15.3	17.8	4.4	0.1	100.0	738
Kakamega	6.4	31.1	18.8	6.8	3.5	13.1	16.1	3.8	0.5	100.0	1,250
Vihiga	5.0	32.9	21.6	6.2	2.9	12.6	15.7	2.3	0.9	100.0	376
Bungoma	5.3	35.5	16.5	7.1	2.8	13.7	15.6	3.4	0.2	100.0	1,208
Busia	6.0	33.7	17.7	9.8	3.3	12.5	12.5	4.4	0.1	100.0	669
Siaya	4.9	30.2	23.1	9.1	4.7	9.1	16.5	2.2	0.2	100.0	684
Kisumu	6.0	26.6	16.4	9.4	3.9	13.7	17.1	6.2	0.8	100.0	853
Homa Bay	5.6	27.3	20.1	13.6	4.7	10.7	13.7	4.2	0.0	100.0	858
Migori	5.3	31.0	21.5	6.1	3.7	12.2	14.1	4.8	1.4	100.0	799
Kisii	4.9	26.4	18.4	9.7	4.2	12.1	20.2	3.9	0.1	100.0	820
Nyamira	3.3	26.5	16.6	9.7	2.7	16.1	21.6	3.2	0.4	100.0	390
Nairobi City	5.8	20.5	14.1	3.6	1.0	7.8	33.9	12.8	0.4	100.0	4,548
Total	8.2	26.2	17.4	6.2	2.7	10.8	21.3	6.7	0.5	100.0	35,004

Note: Respondents whose current or previous level of education was vocational training have been excluded from this table.
[1] No education includes informal education (e.g., madrassa/duksi/adult education).
[2] Completed grade 8 at the primary level for those under age 56; because of the change in the school system in the 1980s, those age 56 and older are considered to have completed primary if they completed grade 7.
[3] Four years of secondary is considered completed secondary for purposes of the Demographic and Health Surveys indicator.
[4] More than secondary includes middle-level colleges and universities.

Table 2.11 Health insurance coverage

Percentage of de jure household population with specific types of health insurance coverage, and percentage with any health insurance, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Social Health Authority (SHA)	Private medical insurance	Community-based health insurance	County insurance scheme	Employer-managed insurance scheme	None/ don't know	Any health insurance	Number of persons
Age								
0–14	35.3	0.8	0.1	0.1	1.4	64.1	35.9	33,886
15–64	41.4	1.4	0.1	0.2	2.0	57.5	42.5	54,520
65+	46.6	1.2	0.2	0.3	0.3	52.3	47.7	4,460
Don't know age	30.3	0.2	0.0	0.0	1.7	69.7	30.3	80
Sex								
Male	37.4	1.2	0.1	0.1	1.8	61.8	38.2	46,764
Female	41.4	1.1	0.2	0.2	1.6	57.6	42.4	46,182
Residence								
Urban	42.4	2.0	0.1	0.1	2.7	56.1	43.9	36,354
Rural	37.5	0.6	0.1	0.2	1.0	62.0	38.0	56,592
Wealth quintile								
Lowest	24.0	0.0	0.0	0.0	0.1	76.0	24.0	18,594
Second	35.3	0.2	0.1	0.1	0.2	64.5	35.5	18,589
Middle	41.6	0.6	0.1	0.2	0.7	57.9	42.1	18,584
Fourth	47.0	1.3	0.3	0.2	2.0	52.1	47.9	18,589
Highest	49.2	3.7	0.1	0.2	5.5	48.0	52.0	18,589
Total	39.4	1.1	0.1	0.2	1.7	59.7	40.3	92,946

Table 2.11C Health insurance coverage: County level

Percentage of de jure household population with specific types of health insurance coverage, and percentage with any health insurance, according to county, Kenya Mini DHS 2025–26

County	Social Health Authority (SHA)	Private medical insurance	Community-based health insurance	County insurance scheme	Employer managed insurance scheme	None/ don't know	Any health insurance	Number of persons
Mombasa	43.4	1.1	0.1	0.0	3.5	55.5	44.5	2,969
Kwale	37.1	0.0	0.0	0.0	1.0	62.6	37.4	1,586
Kilifi	33.1	0.4	0.1	0.1	0.4	66.7	33.3	2,666
Tana River	13.7	0.0	0.0	0.2	0.3	86.3	13.7	721
Lamu	43.8	0.3	0.0	0.0	0.3	56.2	43.8	261
Taita Taveta	55.9	1.3	0.0	0.4	3.2	43.7	56.3	691
Garissa	23.1	0.1	0.1	0.1	0.1	76.8	23.2	852
Wajir	33.4	0.0	0.0	0.0	0.3	66.5	33.5	850
Mandera	24.0	0.1	0.0	0.1	0.1	76.0	24.0	812
Marsabit	28.6	1.5	0.0	0.1	0.1	71.2	28.8	748
Isiolo	18.0	0.3	0.0	0.3	0.6	81.5	18.5	377
Meru	26.8	1.5	0.0	0.2	0.7	72.5	27.5	2,693
Tharaka-Nithi	45.1	1.3	0.3	0.0	2.3	54.2	45.8	704
Embu	51.5	1.1	0.1	0.4	4.9	47.4	52.6	1,035
Kitui	31.6	0.1	0.0	0.2	4.1	67.7	32.3	2,122
Machakos	40.2	0.3	0.0	0.1	1.7	59.0	41.0	2,993
Makueni	39.1	0.6	0.2	0.6	2.4	60.6	39.4	1,665
Nyandarua	48.8	0.4	0.0	0.0	0.2	50.9	49.1	1,060
Nyeri	65.7	2.2	0.5	0.0	0.8	33.1	66.9	1,148
Kirinyaga	59.6	1.9	0.4	0.4	1.7	39.3	60.7	1,210
Murang'a	29.6	1.2	1.4	2.9	1.3	66.6	33.4	1,708
Kiambu	38.0	3.8	0.0	0.0	1.1	59.4	40.6	6,076
Turkana	30.6	0.7	0.0	0.6	2.4	68.3	31.7	1,183
West Pokot	26.0	0.2	0.0	0.1	2.9	73.9	26.1	1,259
Samburu	39.7	0.7	0.0	0.3	1.8	60.2	39.8	554
Trans Nzoia	27.4	0.3	0.1	0.2	2.1	72.0	28.0	1,568
Uasin Gishu	46.3	1.6	0.0	0.0	3.7	52.3	47.7	2,677
Elgeyo-Marakwet	65.2	0.2	0.0	0.1	2.2	34.8	65.2	848
Nandi	46.4	0.6	0.0	0.0	2.0	53.0	47.0	1,550
Baringo	40.0	0.6	0.0	0.1	1.5	59.6	40.4	1,012
Laikipia	44.7	0.4	0.0	0.1	1.1	54.8	45.2	931
Nakuru	44.1	0.6	0.0	0.0	0.7	55.7	44.3	4,659
Narok	22.7	0.6	0.0	0.0	0.0	77.1	22.9	2,265
Kajiado	48.5	1.4	0.0	0.3	2.3	50.3	49.7	1,991
Kericho	49.1	0.8	0.3	0.0	2.1	50.8	49.2	1,777
Bomet	54.1	0.8	0.0	0.1	0.2	45.7	54.3	1,963
Kakamega	41.9	0.6	0.0	0.0	0.4	57.9	42.1	3,162
Vihiga	40.4	0.8	0.2	0.0	0.8	59.1	40.9	1,055
Bungoma	33.1	1.3	0.0	0.1	0.5	66.8	33.2	3,052
Busia	36.5	0.3	0.1	0.3	1.4	63.3	36.7	1,758
Siaya	32.4	0.8	0.2	0.0	0.3	67.1	32.9	1,804
Kisumu	48.1	1.1	0.1	0.1	2.7	51.3	48.7	2,293
Homa Bay	42.4	1.7	0.2	0.0	0.7	56.6	43.4	2,246
Migori	36.3	0.8	0.4	0.0	1.5	63.2	36.8	2,240
Kisii	37.8	0.2	0.0	0.3	0.8	62.0	38.0	2,210
Nyamira	37.8	0.8	0.1	0.0	0.8	61.3	38.7	1,050
Nairobi City	40.2	2.0	0.2	0.0	3.3	58.1	41.9	12,891
Total	39.4	1.1	0.1	0.2	1.7	59.7	40.3	92,946

Table 2.12 Status of premium payments under the Social Health Insurance Fund

Among de jure household members who are registered with the Social Health Authority (SHA) as contributing members, percentage whose premiums have been paid for the current month for the Social Health Insurance Fund (SHIF), according to background characteristics, Kenya Mini DHS, 2025–26

Background characteristic	Percentage of de jure population whose: Premiums have been paid for the current month for SHIF	Premiums have not been paid for the current month for SHIF	Don't know	Total	Number of persons
Age					
0–14	36.9	59.8	3.3	100.0	172
15–64	44.7	51.2	4.0	100.0	16,512
65+	37.4	58.5	4.1	100.0	1,838
Sex					
Male	49.5	44.6	5.9	100.0	9,258
Female	38.4	59.4	2.3	100.0	9,285
Residence					
Urban	55.8	40.2	3.9	100.0	8,285
Rural	34.3	61.5	4.2	100.0	10,259
Wealth quintile					
Lowest	17.9	78.1	4.0	100.0	2,057
Second	26.4	70.3	3.4	100.0	3,106
Middle	35.8	59.2	5.0	100.0	3,914
Fourth	51.1	45.2	3.7	100.0	4,616
Highest	65.9	30.0	4.1	100.0	4,851
Total	43.9	52.0	4.1	100.0	18,544

Note: Total includes 22 persons who were missing information on age who are not shown separately.

Table 2.12C Status of premium payments under the Social Health Insurance Fund: County level

Among de jure household members who are registered with the Social Health Authority (SHA) as contributing members, percentage whose premiums have been paid for the current month for the Social Health Insurance Fund (SHIF), according to county, Kenya Mini DHS, 2025–26

	Percentage of de jure population whose:				
County	Premiums have been paid for the current month for SHIF	Premiums have not been paid for the current month for SHIF	Don't know	Total	Number of persons
Mombasa	44.8	48.6	6.7	100.0	726
Kwale	10.2	82.6	7.3	100.0	230
Kilifi	40.6	57.7	1.7	100.0	476
Tana River	46.8	52.9	0.3	100.0	44
Lamu	33.3	66.5	0.2	100.0	58
Taita Taveta	28.6	66.5	4.9	100.0	258
Garissa	35.7	63.7	0.6	100.0	78
Wajir	24.7	69.0	6.2	100.0	91
Mandera	25.4	74.6	0.0	100.0	84
Marsabit	46.5	47.8	5.7	100.0	85
Isiolo	54.7	44.7	0.6	100.0	29
Meru	37.6	60.5	1.9	100.0	391
Tharaka-Nithi	36.2	62.5	1.2	100.0	172
Embu	37.7	53.5	8.8	100.0	310
Kitui	39.7	58.6	1.8	100.0	331
Machakos	40.0	55.3	4.7	100.0	691
Makueni	34.0	62.8	3.2	100.0	393
Nyandarua	41.6	50.8	7.6	100.0	234
Nyeri	33.4	62.5	4.1	100.0	417
Kirinyaga	29.3	64.0	6.7	100.0	377
Murang'a	55.6	42.5	1.9	100.0	283
Kiambu	68.6	27.1	4.4	100.0	1,342
Turkana	34.7	61.7	3.6	100.0	148
West Pokot	45.4	54.2	0.4	100.0	134
Samburu	20.2	71.9	7.9	100.0	112
Trans Nzoia	42.3	50.9	6.8	100.0	242
Uasin Gishu	46.2	49.8	3.9	100.0	554
Elgeyo-Marakwet	38.4	59.8	1.8	100.0	213
Nandi	45.4	53.4	1.2	100.0	437
Baringo	53.1	35.9	11.0	100.0	177
Laikipia	55.6	38.9	5.5	100.0	189
Nakuru	50.0	46.8	3.2	100.0	1,022
Narok	42.8	51.6	5.6	100.0	245
Kajiado	61.6	34.7	3.8	100.0	493
Kericho	53.7	44.9	1.4	100.0	441
Bomet	28.4	69.7	2.0	100.0	406
Kakamega	30.4	67.5	2.1	100.0	700
Vihiga	25.1	63.0	11.9	100.0	227
Bungoma	25.1	70.6	4.2	100.0	508
Busia	25.1	63.6	11.3	100.0	308
Siaya	30.6	67.1	2.2	100.0	285
Kisumu	38.4	55.9	5.7	100.0	477
Homa Bay	34.8	64.5	0.8	100.0	397
Migori	42.0	55.2	2.9	100.0	355
Kisii	35.5	60.1	4.4	100.0	360
Nyamira	26.4	67.7	5.9	100.0	179
Nairobi City	57.2	39.2	3.6	100.0	2,834
Total	43.9	52.0	4.1	100.0	18,544

Table 2.13 Households operating an income-generating business/enterprise

Percentage of households in which any member operated an income-generating business or enterprise in the past month, by background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Household operated an enterprise (%)	Number of households
Residence		
Urban	24.5	11,975
Rural	15.4	14,405
County		
Mombasa	24.6	866
Kwale	17.3	318
Kilifi	27.3	699
Tana River	13.4	143
Lamu	9.1	62
Taita Taveta	19.7	206
Garissa	12.0	145
Wajir	13.3	126
Mandera	6.8	137
Marsabit	15.4	138
Isiolo	17.8	87
Meru	12.7	840
Tharaka-Nithi	13.7	209
Embu	18.0	385
Kitui	19.4	526
Machakos	17.1	1,021
Makueni	16.6	506
Nyandarua	9.6	341
Nyeri	17.0	441
Kirinyaga	16.6	482
Murang'a	11.7	571
Kiambu	14.8	2,295
Turkana	42.7	232
West Pokot	13.4	251
Samburu	19.4	109
Trans Nzoia	16.5	393
Uasin Gishu	20.2	794
Elgeyo-Marakwet	9.5	209
Nandi	10.1	414
Baringo	11.2	242
Laikipia	10.7	293
Nakuru	18.1	1,442
Narok	11.0	576
Kajiado	14.2	653
Kericho	15.4	464
Bomet	12.1	437
Kakamega	24.2	850
Vihiga	18.0	267
Bungoma	15.0	746
Busia	19.6	429
Siaya	18.1	482
Kisumu	33.0	615
Homa Bay	32.4	568
Migori	31.8	499
Kisii	8.0	597
Nyamira	11.3	284
Nairobi City	29.7	3,992
Wealth quintile		
Lowest	10.4	4,299
Second	15.4	4,719
Middle	18.1	5,346
Fourth	23.3	6,192
Highest	26.9	5,825
Total	19.5	26,380

Table 2.14 Household enterprises by sex of operator

Percent distribution of household enterprises by sex of operator, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Male enterprise operators (%)	Female enterprise operators (%)	Total	Number of enterprises
Residence				
Urban	42.3	57.7	100.0	3,221
Rural	40.4	59.6	100.0	2,354
County				
Mombasa	31.0	69.0	100.0	237
Kwale	41.7	58.3	100.0	56
Kilifi	40.3	59.7	100.0	201
Tana River	40.1	59.9	100.0	20
Lamu	(47.8)	(52.2)	100.0	6
Taita Taveta	42.3	57.7	100.0	45
Garissa	47.7	52.3	100.0	17
Wajir	11.4	88.6	100.0	17
Mandera	(44.2)	(55.8)	100.0	9
Marsabit	30.3	69.7	100.0	21
Isiolo	39.7	60.3	100.0	16
Meru	42.1	57.9	100.0	114
Tharaka-Nithi	34.8	65.2	100.0	30
Embu	54.1	45.9	100.0	79
Kitui	42.1	57.9	100.0	114
Machakos	32.0	68.0	100.0	179
Makueni	33.1	66.9	100.0	87
Nyandarua	(41.6)	(58.4)	100.0	33
Nyeri	51.1	48.9	100.0	80
Kirinyaga	47.6	52.4	100.0	84
Murang'a	52.4	47.6	100.0	70
Kiambu	44.2	55.8	100.0	383
Turkana	22.8	77.2	100.0	102
West Pokot	43.2	56.8	100.0	37
Samburu	29.7	70.3	100.0	22
Trans Nzoia	33.5	66.5	100.0	66
Uasin Gishu	46.5	53.5	100.0	169
Elgeyo-Marakwet	36.5	63.5	100.0	22
Nandi	39.0	61.0	100.0	43
Baringo	34.0	66.0	100.0	27
Laikipia	48.5	51.5	100.0	34
Nakuru	44.5	55.5	100.0	280
Narok	46.5	53.5	100.0	68
Kajiado	52.1	47.9	100.0	97
Kericho	43.6	56.4	100.0	74
Bomet	59.1	40.9	100.0	57
Kakamega	34.9	65.1	100.0	210
Vihiga	46.2	53.8	100.0	55
Bungoma	45.1	54.9	100.0	119
Busia	33.0	67.0	100.0	89
Siaya	23.6	76.4	100.0	92
Kisumu	27.9	72.1	100.0	217
Homa Bay	40.1	59.9	100.0	196
Migori	38.7	61.3	100.0	176
Kisii	51.1	48.9	100.0	50
Nyamira	49.3	50.7	100.0	35
Nairobi City	46.3	53.7	100.0	1,338
Wealth quintile				
Lowest	32.6	67.4	100.0	455
Second	37.7	62.3	100.0	757
Middle	43.7	56.3	100.0	1,025
Fourth	40.4	59.6	100.0	1,532
Highest	45.1	54.9	100.0	1,806
Total	41.5	58.5	100.0	5,576

Note: Figures in parentheses are based on 25–49 unweighted cases.

Table 2.15 Registration status of household enterprises

Percentage of household enterprises registered or licensed with selected authorities, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Business Registration Service (formerly Registrar of Companies)	County Government	Micro and Small Enterprise Authority	Professional bodies	Social Development Department	Other	Not registered	Number of enterprises
Residence								
Urban	9.3	31.8	4.4	2.7	1.7	0.4	63.3	3,221
Rural	3.7	28.2	3.0	2.6	2.1	0.4	68.9	2,354
County								
Mombasa	15.6	28.5	14.9	8.2	7.0	0.0	66.8	237
Kwale	0.0	31.6	0.0	0.0	0.0	0.0	68.4	56
Kilifi	3.2	8.4	4.0	0.9	3.3	0.0	90.9	201
Tana River	3.9	38.5	2.1	2.1	0.0	0.0	61.5	20
Lamu	(0.9)	(34.9)	(0.9)	(2.6)	(2.6)	(0.0)	(65.1)	6
Taita Taveta	1.5	50.5	2.6	0.8	0.0	0.0	49.5	45
Garissa	1.3	22.1	0.0	0.0	1.3	0.0	77.9	17
Wajir	1.2	40.4	0.0	0.0	0.0	0.0	58.4	17
Mandera	(0.0)	(76.0)	(0.0)	(0.0)	(0.0)	(0.0)	(24.0)	9
Marsabit	0.0	22.2	1.0	0.0	0.0	0.0	76.8	21
Isiolo	1.4	15.3	0.0	1.5	0.0	0.0	83.3	16
Meru	4.1	34.3	2.0	3.9	2.4	4.2	57.1	114
Tharaka-Nithi	4.1	37.2	4.1	2.0	0.9	2.0	61.9	30
Embu	7.7	29.9	1.6	1.9	0.0	1.0	69.1	79
Kitui	4.6	40.3	3.2	0.7	0.0	0.0	59.5	114
Machakos	1.9	43.7	3.1	1.9	2.1	0.0	53.7	179
Makueni	6.7	55.3	6.7	1.6	0.0	3.1	40.2	87
Nyandarua	(32.7)	(67.6)	(7.4)	(2.2)	(0.0)	(2.2)	(25.2)	33
Nyeri	1.0	45.6	0.0	1.0	0.0	0.5	54.4	80
Kirinyaga	4.8	35.3	1.4	3.4	0.0	4.0	59.8	84
Murang'a	3.4	58.0	3.1	4.3	1.4	0.0	41.8	70
Kiambu	4.3	28.6	0.6	4.4	1.7	0.0	67.0	383
Turkana	0.6	7.3	1.5	0.7	0.0	0.0	91.1	102
West Pokot	0.0	34.2	0.8	0.5	2.1	0.0	63.7	37
Samburu	1.4	20.2	0.0	1.4	0.0	0.0	78.4	22
Trans Nzoia	2.8	9.7	2.1	0.8	0.8	0.0	90.3	66
Uasin Gishu	5.1	32.0	1.5	1.3	2.3	0.6	65.9	169
Elgeyo-Marakwet	0.0	30.4	2.9	0.0	0.0	0.0	66.7	22
Nandi	2.2	21.8	2.8	3.4	0.0	2.2	73.2	43
Baringo	4.2	24.7	8.5	6.6	0.6	0.0	66.3	27
Laikipia	14.9	58.6	3.0	11.9	3.0	1.6	41.4	34
Nakuru	2.9	38.6	1.4	0.0	0.0	0.0	60.1	280
Narok	1.3	19.8	2.9	1.3	0.0	0.0	77.3	68
Kajiado	19.6	40.4	17.4	12.4	11.5	0.0	53.4	97
Kericho	8.1	30.6	7.2	1.4	2.4	0.0	67.3	74
Bomet	0.2	31.1	0.0	6.5	0.2	0.0	65.9	57
Kakamega	5.6	31.9	2.3	0.0	0.0	0.0	66.3	210
Vihiga	7.1	33.0	2.6	2.9	5.8	0.6	63.7	55
Bungoma	2.8	37.5	0.0	2.2	1.9	0.3	59.4	119
Busia	3.2	12.4	1.8	0.9	0.9	0.0	85.3	89
Siaya	5.4	17.1	8.1	3.4	3.7	0.0	74.7	92
Kisumu	5.4	23.0	1.8	2.1	0.5	0.9	74.5	217
Homa Bay	5.4	42.4	4.0	12.8	15.7	0.0	51.5	196
Migori	11.6	30.5	8.7	3.6	2.6	0.5	64.5	176
Kisii	9.4	46.3	19.4	5.7	1.1	0.0	45.5	50
Nyamira	7.2	20.6	3.9	0.0	1.4	0.0	79.4	35
Nairobi City	11.3	25.5	3.4	1.0	0.0	0.3	67.4	1,338
Wealth quintile								
Lowest	0.7	11.5	1.3	0.7	0.8	0.0	87.4	455
Second	1.4	19.4	2.3	2.2	2.6	0.6	77.2	757
Middle	3.1	26.6	2.0	1.8	1.9	0.2	71.4	1,025
Fourth	5.4	31.2	3.5	2.3	1.4	0.5	64.8	1,532
Highest	14.4	40.9	6.3	4.1	2.2	0.4	52.8	1,806
Total	7.0	30.3	3.8	2.7	1.9	0.4	65.6	5,576

Notes: Enterprises may be registered with more than one authority. Figures in parentheses are based on 25–49 unweighted cases.

Table 2.16 Type of economic activity of household enterprises

Percent distribution of household enterprises by main economic activity, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Manufacturing	Services	Agri-business	Total	Number of enterprises
Residence					
Urban	3.5	92.6	3.8	100.0	3,221
Rural	8.7	83.9	7.4	100.0	2,354
County					
Mombasa	3.2	96.8	0.0	100.0	237
Kwale	29.3	66.3	4.4	100.0	56
Kilifi	5.8	88.8	5.5	100.0	201
Tana River	0.6	99.4	0.0	100.0	20
Lamu	(0.0)	(100.0)	(0.0)	100.0	6
Taita Taveta	0.7	95.6	3.6	100.0	45
Garissa	0.0	100.0	0.0	100.0	17
Wajir	0.0	100.0	0.0	100.0	17
Mandera	(0.0)	(95.5)	(4.5)	100.0	9
Marsabit	2.2	85.7	12.1	100.0	21
Isiolo	4.6	85.0	10.4	100.0	16
Meru	3.0	91.3	5.6	100.0	114
Tharaka-Nithi	20.4	75.9	3.7	100.0	30
Embu	6.0	93.1	0.9	100.0	79
Kitui	3.8	91.2	5.0	100.0	114
Machakos	2.9	91.4	5.7	100.0	179
Makueni	2.2	95.4	2.4	100.0	87
Nyandarua	(1.5)	(94.6)	(4.0)	100.0	33
Nyeri	3.3	96.7	0.0	100.0	80
Kirinyaga	1.9	91.3	6.8	100.0	84
Murang'a	14.9	80.9	4.3	100.0	70
Kiambu	2.2	89.2	8.6	100.0	383
Turkana	66.0	25.9	8.1	100.0	102
West Pokot	8.6	82.0	9.4	100.0	37
Samburu	8.1	83.6	8.3	100.0	22
Trans Nzoia	1.0	94.3	4.7	100.0	66
Uasin Gishu	8.1	86.6	5.3	100.0	169
Elgeyo-Marakwet	5.2	89.0	5.8	100.0	22
Nandi	6.3	89.7	4.0	100.0	43
Baringo	3.8	80.0	16.1	100.0	27
Laikipia	2.1	97.9	0.0	100.0	34
Nakuru	2.0	96.7	1.3	100.0	280
Narok	2.8	85.0	12.3	100.0	68
Kajiado	4.8	91.2	4.0	100.0	97
Kericho	0.0	98.9	1.1	100.0	74
Bomet	4.2	95.8	0.0	100.0	57
Kakamega	10.6	81.8	7.6	100.0	210
Vihiga	6.5	92.3	1.2	100.0	55
Bungoma	3.4	83.6	13.0	100.0	119
Busia	1.5	98.5	0.0	100.0	89
Siaya	4.4	75.1	20.5	100.0	92
Kisumu	4.0	89.3	6.7	100.0	217
Homa Bay	4.4	85.1	10.5	100.0	196
Migori	14.1	76.0	10.0	100.0	176
Kisii	16.2	81.8	2.0	100.0	50
Nyamira	0.0	83.2	16.8	100.0	35
Nairobi City	2.9	93.5	3.6	100.0	1,338
Wealth quintile					
Lowest	20.1	72.4	7.5	100.0	455
Second	6.7	85.1	8.2	100.0	757
Middle	5.0	89.3	5.7	100.0	1,025
Fourth	5.0	90.8	4.2	100.0	1,532
Highest	2.6	93.0	4.3	100.0	1,806
Total	5.7	89.0	5.3	100.0	5,576

Note: Figures in parentheses are based on 25–49 unweighted cases.

Table 2.17 Average monthly revenue of household enterprises

Percent distribution of household enterprises according to monthly revenue and background characteristics, Kenya Mini DHS 2025–26

Background characteristic	KSh <5,000 (%)	KSh 5,000–9,999 (%)	KSh 10,000–19,999 (%)	KSh 20,000–49,999 (%)	KSh 50,000+ (%)	Total	Number of enterprises
Residence							
Urban	27.5	9.7	22.5	25.8	14.6	100.0	3,221
Rural	43.1	14.7	21.4	14.8	6.0	100.0	2,354
County							
Mombasa	32.1	15.7	27.2	13.2	11.8	100.0	237
Kwale	64.6	11.0	15.9	6.5	2.0	100.0	56
Kilifi	56.2	13.1	15.4	11.8	3.5	100.0	201
Tana River	6.1	1.2	37.6	22.8	32.3	100.0	20
Lamu	(15.4)	(20.9)	(31.5)	(23.1)	(9.1)	100.0	6
Taita Taveta	29.2	13.7	32.8	18.5	5.8	100.0	45
Garissa	15.0	8.6	37.5	26.7	12.2	100.0	17
Wajir	54.1	26.9	13.0	6.0	0.0	100.0	17
Mandera	(34.8)	(22.1)	(18.7)	(4.1)	(20.3)	100.0	9
Marsabit	26.4	15.0	25.5	19.5	13.7	100.0	21
Isiolo	30.0	9.2	32.3	20.6	7.9	100.0	16
Meru	42.8	17.2	24.5	10.5	5.0	100.0	114
Tharaka-Nithi	56.0	13.2	12.9	7.6	10.4	100.0	30
Embu	37.8	13.3	21.9	15.5	11.6	100.0	79
Kitui	36.3	12.7	18.3	24.0	8.7	100.0	114
Machakos	34.1	7.6	23.0	20.4	14.9	100.0	179
Makueni	20.1	12.5	33.3	22.3	11.8	100.0	87
Nyandarua	(18.9)	(7.3)	(20.9)	(35.1)	(17.8)	100.0	33
Nyeri	35.6	17.7	23.6	18.0	5.1	100.0	80
Kirinyaga	37.3	11.6	25.3	18.9	6.9	100.0	84
Murang'a	32.7	22.0	20.6	18.2	6.6	100.0	70
Kiambu	22.7	8.3	24.6	29.4	15.1	100.0	383
Turkana	61.3	16.3	13.1	6.6	2.7	100.0	102
West Pokot	21.0	16.5	22.4	14.7	25.4	100.0	37
Samburu	51.4	10.4	16.5	19.4	2.4	100.0	22
Trans Nzoia	26.2	21.8	29.5	18.6	3.9	100.0	66
Uasin Gishu	18.9	9.6	24.1	37.1	10.3	100.0	169
Elgeyo-Marakwet	39.4	18.1	24.1	11.4	7.1	100.0	22
Nandi	44.8	6.6	19.1	25.9	3.5	100.0	43
Baringo	42.6	15.9	31.6	8.5	1.4	100.0	27
Laikipia	38.2	6.7	24.0	19.9	11.3	100.0	34
Nakuru	22.4	18.9	23.2	25.6	9.9	100.0	280
Narok	20.9	13.3	37.8	14.7	13.3	100.0	68
Kajiado	10.2	12.0	35.3	20.3	22.3	100.0	97
Kericho	7.2	10.8	37.2	26.5	18.2	100.0	74
Bomet	34.6	20.0	22.0	20.3	3.2	100.0	57
Kakamega	50.2	19.5	15.0	11.5	3.8	100.0	210
Vihiga	41.7	12.9	24.9	18.7	1.8	100.0	55
Bungoma	52.2	19.4	15.2	12.7	0.4	100.0	119
Busia	60.4	9.6	15.4	7.6	7.0	100.0	89
Siaya	53.0	5.9	15.0	19.4	6.7	100.0	92
Kisumu	38.8	12.1	23.9	16.2	8.9	100.0	217
Homa Bay	47.8	15.5	18.9	13.5	4.2	100.0	196
Migori	31.4	13.5	21.2	23.6	10.3	100.0	176
Kisii	31.8	10.0	25.1	29.1	4.0	100.0	50
Nyamira	22.4	26.5	31.8	11.3	8.0	100.0	35
Nairobi City	29.8	5.8	19.5	27.9	16.9	100.0	1,338
Wealth quintile							
Lowest	63.1	14.7	14.5	6.2	1.5	100.0	455
Second	49.8	18.0	19.4	9.9	2.8	100.0	757
Middle	40.3	15.6	22.1	16.3	5.8	100.0	1,025
Fourth	25.8	12.6	29.3	24.1	8.1	100.0	1,532
Highest	23.8	5.5	18.8	29.9	22.0	100.0	1,806
Total	34.1	11.8	22.0	21.2	10.9	100.0	5,576

Note: Figures in parentheses are based on 25–49 unweighted cases.

Table 2.18 Household member deaths

Percent distribution of household member deaths (2021–2025) by place of occurrence and background characteristics, Kenya Mini DHS, 2025–26

Background characteristic	Place of death			Total	Number of deaths (2021–2025)
	Health facility	Community	Don't know		
Age at death					
<5	71.9	28.1	0.0	100.0	201
5–14	55.9	44.1	0.0	100.0	69
15–49	56.6	42.4	0.9	100.0	509
50–64	67.4	32.0	0.6	100.0	288
65+	52.6	47.3	0.1	100.0	639
Sex					
Male	58.7	40.7	0.6	100.0	1,054
Female	58.7	41.2	0.2	100.0	652
Residence					
Urban	60.7	39.0	0.3	100.0	394
Rural	58.1	41.4	0.5	100.0	1,311
Total	58.7	40.9	0.4	100.0	1,705

Table 2.19 Death certificate and burial permit issuance

Percentage of household member deaths for which a death certificate and/or burial permit was issued (2021–2025), by background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Death certificate issued (Death Certification)[1]	Burial permit issued only	Either death certificate or burial permit issued (Death Notification)	Neither death certificate nor burial permit issued	Number of household deaths (2021–2025)
Age at death					
<5	36.2	37.4	73.5	26.5	201
5–14	42.4	44.2	86.6	13.4	69
15–49	53.3	39.1	92.4	7.6	509
50–64	67.2	26.8	94.0	6.0	288
65+	59.1	32.6	91.7	8.3	639
Male	56.7	34.2	90.9	9.1	1,054
Female	53.2	35.1	88.3	11.7	652
Residence					
Urban	55.6	33.6	89.3	10.7	394
Rural	55.3	34.9	90.1	9.9	1,311
Total	55.4	34.6	89.9	10.1	1,705

[1] A burial permit is required for issuance of a death certificate.

Table 2.19C Death certificate and burial permit issuance: County level

Percentage of household member deaths for which a death certificate and/or burial permit was issued (2021–2025), by county, Kenya Mini DHS 2025–26

County	Death certificate issued (Death Certification)[1]	Burial permit issued only	Either death certificate or burial permit issued (Death Notification)	Neither death certificate nor burial permit issued	Number of household deaths (2021–2025)
Mombasa	*	*	*	*	41
Kwale	39.0	20.9	59.9	40.1	32
Kilifi	(67.2)	(30.0)	(97.2)	(2.8)	48
Tana River	(29.8)	(20.5)	(50.3)	(49.7)	10
Lamu	68.0	30.1	98.1	1.9	7
Taita Taveta	57.7	37.2	94.9	5.1	24
Garissa	(25.1)	(0.0)	(25.1)	(74.9)	12
Wajir	33.0	0.0	33.0	67.0	13
Mandera	34.1	8.9	43.0	57.0	17
Marsabit	(24.9)	(0.0)	(24.9)	(75.1)	12
Isiolo	(60.7)	(18.4)	(79.2)	(20.8)	9
Meru	47.4	45.0	92.4	7.6	85
Tharaka-Nithi	(62.1)	(35.0)	(97.1)	(2.9)	21
Embu	(64.9)	(30.7)	(95.6)	(4.4)	28
Kitui	79.9	18.9	98.8	1.2	82
Machakos	58.7	39.9	98.7	1.3	79
Makueni	55.5	42.6	98.1	1.9	45
Nyandarua	(35.0)	(65.0)	(100.0)	(0.0)	35
Nyeri	(57.3)	(37.9)	(95.2)	(4.8)	33
Kirinyaga	(76.4)	(20.7)	(97.2)	(2.8)	34
Murang'a	54.8	43.9	98.7	1.3	55
Kiambu	(53.2)	(41.4)	(94.6)	(5.4)	62
Turkana	18.1	15.0	33.1	66.9	40
West Pokot	(42.7)	(14.7)	(57.4)	(42.6)	17
Samburu	(31.4)	(32.8)	(64.3)	(35.7)	8
Trans Nzoia	(70.0)	(27.0)	(96.9)	(3.1)	26
Uasin Gishu	(61.3)	(36.1)	(97.5)	(2.5)	37
Elgeyo-Marakwet	(65.2)	(28.2)	(93.5)	(6.5)	17
Nandi	(53.3)	(46.7)	(100.0)	(0.0)	21
Baringo	*	*	*	*	11
Laikipia	(79.5)	(12.0)	(91.6)	(8.4)	15
Nakuru	(52.5)	(37.9)	(90.4)	(9.6)	63
Narok	(53.4)	(32.7)	(86.0)	(14.0)	29
Kajiado	*	*	*	*	18
Kericho	(43.3)	(47.7)	(91.0)	(9.0)	31
Bomet	*	*	*	*	16
Kakamega	70.1	29.9	100.0	0.0	81
Vihiga	41.3	53.3	94.6	5.4	31
Bungoma	60.5	32.2	92.7	7.3	64
Busia	(60.0)	(32.8)	(92.8)	(7.2)	39
Siaya	50.7	39.1	89.8	10.2	56
Kisumu	(49.7)	(48.1)	(97.8)	(2.2)	46
Homa Bay	(57.0)	(37.8)	(94.8)	(5.2)	39
Migori	(57.8)	(33.3)	(91.0)	(9.0)	38
Kisii	63.8	31.8	95.6	4.4	68
Nyamira	(57.0)	(33.0)	(90.0)	(10.0)	24
Nairobi City	*	*	*	*	86
Total	55.4	34.6	89.9	10.1	1,705

Note: Figures in parentheses are based on 25–49 unweighted cases. An asterisk indicates that a figure is based on fewer than 25 unweighted cases and has been suppressed.

[1] A burial permit is required for issuance of a death certificate.

Table 2.20 Reasons for non-issuance of burial permits

Percent distribution of deaths in the 5 years preceding the survey (2021–2025) without a burial permit by main reason for non-issuance, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Main reason for non-issuance of burial permit						Total	Number of deaths (2021–2025)
	Cultural and social reasons	Access and geographical issues	Cost	Perceived lack of benefit	Other	Don't know		
Age at death								
<5	20.3	7.4	7.9	57.0	2.2	5.2	100.0	49
5–14	*	*	*	*	*	*	100.0	9
15–49	17.6	7.2	12.5	31.7	18.8	12.2	100.0	31
50–64	(29.4)	(1.3)	(35.3)	(28.2)	(0.0)	(5.8)	100.0	15
65+	13.5	0.7	7.7	54.4	12.3	11.4	100.0	36
Sex								
Male	17.9	4.8	7.1	47.1	12.5	10.6	100.0	76
Female	19.8	4.2	19.4	47.9	2.9	5.8	100.0	63
Residence								
Urban	19.9	1.5	4.7	46.6	18.5	8.8	100.0	34
Rural	18.4	5.5	15.2	47.7	4.9	8.3	100.0	105
Total	18.8	4.5	12.7	47.5	8.2	8.4	100.0	139

Note: Figures in parentheses are based on 25–49 unweighted cases. An asterisk indicates that a figure is based on fewer than 25 unweighted cases and has been suppressed.

CHARACTERISTICS OF RESPONDENTS 3

Key Findings

- ***Education:*** 6% of women have no education, 12% have some primary education, and 17% have completed primary school but pursued no further education. Overall, 45% of women have completed at least a secondary education.
- ***Exposure to mass media:*** The level of exposure to mass media is high among women in Kenya. Only 27% report that they do not access any of three media sources (newspapers, television, or radio).
- ***Internet usage:*** 61% of women age 15–49 used the internet in the past 12 months. Among those who used the internet in the past 12 months, 65% use the internet daily.
- ***Migration:*** 46% of women were not born in their current place of residence; among these women, 34% moved to their current place of residence in the 5 years prior to the survey. Economic or business reasons were the leading cause of migration, accounting for 37% of cases. Family reunification and other family-related reasons followed at 28%, while marriage formation contributed to 22% of migrations.
- ***Health insurance coverage:*** Overall, 42% of women age 15–49 are registered with the Social Health Authority (SHA). Among those registered, 77% found the registration process easy or very easy.
- ***Reasons for not registering with the Social Health Authority:*** Among women not registered with the SHA, the main reasons were cost (37%), documentation problems or inadequate coverage (24%), and either lack of interest or satisfaction with other coverage (17%).
- ***Utilisation of Social Health Authority services:*** 53% of women registered with the SHA used SHA services in the 12 months preceding the survey.

This chapter presents information on the demographic and socioeconomic characteristics of survey respondents, including age, education and literacy, marital status, religion, self-reported health status, health insurance coverage and utilisation, wealth, mass media exposure, internet use, type of current residence, place of birth and recent migration, and reasons for migration. These factors help explain access to and use of reproductive health services, use of contraceptives, and other health behaviours.

3.1 Basic Characteristics of Survey Respondents

The 2025–26 Kenya Mini Demographic and Health Survey (KMDHS) included 21,231 women age 15–49. The largest proportion of respondents were teenagers age 15–19, accounting for 18% of all women interviewed.

Fifty-seven percent of women rate their health as good, 23% rate it as very good, and a very small percentage rate it as bad (3%) or very bad (1%).

Fifty-five percent of women are currently married or living with a partner as though married; 33% have never been married; and 12% are divorced, separated, or widowed (**Table 3.1**). At the county level, the largest proportion of respondents (17%) are from Nairobi City (**Table 3.1C**).

3.2 Education and Literacy

Literacy


Respondents who had attended more than secondary school were assumed to be literate. All other respondents were considered literate if they could read aloud all or part of a sentence shown to them.

Sample: Women age 15–49

Six percent of women have no education, 17% have completed a primary education, 25% have completed a secondary education, and 20% have more than a secondary education (**Figure 3.1**). The median number of years of education completed is 8.7 years among the women (**Tables 3.2** and **3.2C**). Eight percent of women are illiterate (**Tables 3.3** and **3.3C**).

Figure 3.1 **Education**

Percent distribution of women age 15–49 by highest level of schooling



Trends: The percentage of women with no education has declined sharply from 13% in the 2003 Kenya Demographic and Health Survey (KDHS) to 6% in the 2025–26 KMDHS. Over the same period, the percentage of women with more than a secondary education has risen from 6% to 20%.


Patterns by background characteristics

- The percentage of women with no education increases with age. Only 3% of young women age 15–24 have no education, compared with 19% of women age 40–49 (**Table 3.2**).
- Women in rural areas are twice as likely as women in urban areas to have no education or to have not completed primary school (22% versus 11%).

- The percentage of women who have completed a secondary or higher level of education rises steadily with wealth, from 13% among women in the lowest wealth quintile to 72% among those in the highest quintile (**Figure 3.2**).
- By county, the percentage of women who have completed a secondary or higher level of education is highest in Kiambu (70%) and lowest in Mander (14%) (**Table 3.2C** and **Map 3.1**).


Figure 3.2 Secondary education by household wealth

Percentage of women age 15–49 who have completed at least a secondary education by wealth quintile



Map 3.1 Secondary education by county

Percentage of women age 15–49 who have completed at least a secondary education



3.3 Mass Media Exposure and Internet Usage

Exposure to mass media

Respondents were asked how often they read a newspaper, listen to the radio, and watch television. Those who responded *at least once a week* are considered regularly exposed to that form of media.

Sample: Women age 15–49

Use of the internet

Respondents were asked if they have ever used the internet from any device; if they had used the internet in the past 12 months; and, if they have used the internet in the past 12 months, how often they used it during the past month.

Sample: Women age 15–49


3.3.1 Mass Media Exposure

Exposure to mass media improves access to information, raises awareness about health and social issues, supports education, and promotes civic participation. **Tables 3.4** and **3.4C** show the percentages of women in Kenya who are exposed to different types of media by background characteristics. Radio and television are the most commonly used forms of mass media, used by 52% and 50% of women, respectively (**Figure 3.3**).

More than one-quarter of women (27%) reported that they do not access any form of mass media (radio, television, or newspapers) at least once a week, while only 3% reported accessing all three at least once a week.

Figure 3.3 **Exposure to mass media**


Percentage of women age 15–49 who are exposed to specific media on a weekly basis



Trends: Exposure to radio and television has declined steadily since 2003, from 75% in the 2003 KDHS to 52% in the 2025–26 KMDHS survey. Similarly, exposure to newspapers decreased from 23% to just 5% over the same period. However, exposure to television increased from 29% in 2003 to 50% in 2025–26 (**Figure 3.4**).

Figure 3.4 **Trends in exposure to mass media**

Percentage of women age 15–49 who are exposed to specific media on a weekly basis



Patterns by background characteristics

- Among the three forms of mass media, newspapers were the least accessed, with only 5% of women reporting exposure (**Table 3.4**).

- Access to all three forms of media (radio, television, and newspapers) increases with level of education. Sixty-five percent of women with no education versus 18% of women with more than a secondary education reported not accessing any of the three forms of media at least once a week.
- Exposure to newspapers and television increases with wealth. Among women in the lowest wealth quintile, only 2% reported exposure to newspapers and 8% to television. In contrast, 8% of women in the highest quintile reported exposure to newspapers and 78% to television.
- By county, the percentage of women who do not access any of the three forms of media at least once a week is highest in Wajir (91%) and lowest in Nyandarua (11%) (**Table 3.2C**).

3.3.2 Internet Usage

The internet provides substantial economic benefits because it enables new forms of employment, business, communication, entertainment, expression, and collaboration. It also improves access to a vast depth of knowledge and learning resources, as well as access to services where traditional ways of service provision are lacking.

Sixty-one percent of women age 15–49 used the internet in the 12 months preceding the survey. Among those who used the internet in the past 12 months, 65% used it daily during the past month (**Table 3.5**).


Trends: The percentage of women age 15–49 who used the internet in the past 12 months increased from 44% in the 2022 KDHS to 61% in the 2025–26 KMDHS.

Patterns by background characteristics

- Internet usage is more common in urban areas than in rural areas. In the past 12 months, 81% of women in urban areas compared with 43% of women in rural areas used the internet (**Figure 3.5**).
- The percentage of women who used the internet in the past 12 months increases with level of education. Twenty-five percent of women with no education compared with 95% of those with more than a secondary education used the internet in the past 12 months.
- The percentage of women who used the internet in the past 12 months rises steadily with wealth, with 37% of women in the lowest wealth quintile versus 84% of women in the highest wealth quintile using it almost every day.

Figure 3.5 **Internet usage by residence**

Percentage of women age 15–49 who used the internet in the last 12 months



- By county, internet usage in the past 12 months was highest in Nairobi City (87%) and lowest in Turkana (18%) (**Table 3.5C**).

3.4 Place of Birth and Recent Migration

Recent migration

Percentage of respondents who were born outside of their current place of residence and moved to their current place of residence in the 5 years preceding the survey.

Lifetime migrants

Percentage of respondents who were born outside their current place of residence.

Lifetime internal migrants

Percentage of respondents who were born in Kenya but outside their current place of residence.

Sample: Women age 15–49 who were born outside their current place of residence

Fifty-one percent of women age 15–49 have always lived in their current place of residence. Forty-six percent were born in Kenya but outside their current place of residence, while 3% were born outside of Kenya (**Tables 3.6** and **3.6C**).

Patterns by background characteristics

- The percentage of women age 15–49 who moved to their current place of residence in the past 5 years steadily declines with increasing age. Women age 20–24 have the highest rate of recent migration (67%), while women age 45–49 have the lowest (10%) (**Table 3.6**).
- Lifetime internal migration varies by residence. Women age 15–49 are more likely to be lifetime internal migrants if they live in urban areas (63%) than if they live in rural areas (32%).
- The likelihood of lifetime internal migration increases with level of education. Fifty-nine percent of women who have more than a secondary education were born outside their current place of residence, compared with only 23% of women who have no education.
- Lifetime internal migration varies by county. More than two-thirds of women in Kisii (78%), Laikipia (77%), Nairobi City (72%), and Narok (67%) are lifetime internal migrants. In contrast, Wajir and Mandera Counties report very low levels of lifetime internal migration (<1% each) (**Table 3.6C**).

3.4.1 Type of Migration

Recent migration patterns show that rural-to-urban movements account for the largest share of migration (37%), followed by urban-to-urban migration (34%). Rural-to-urban migration was most common among women age 20–24 (43%), while urban-to-rural migration was most common among women age 40–44 (29%) (**Table 3.7**).

3.4.2 Reason for Migration

Economic or business reasons were the leading cause of migration, accounting for 37% of cases. Family reunification and other family-related reasons followed at 28%, while marriage formation contributed to 22% of migrations. One in ten women (10%) migrated for education or training purposes.

Among women in urban areas, nearly half (49%) reported migrating for economic or business reasons. In contrast, most women in rural areas migrated due to marriage formation (42%), followed closely by family reunification or other family-related reasons (36%) (**Tables 3.8** and **3.8C**).

3.5 Social Health Authority (SHA) Coverage and Utilisation

The Social Health Authority (SHA) was established under the Social Health Insurance Act, 2023 (Republic of Kenya 2023) to replace the former National Hospital Insurance Fund. The SHA is central to Kenya's efforts to achieve universal health coverage. It manages three funds—the Primary Health Care Fund; the Social Health Insurance Fund (SHIF); and the Emergency, Chronic and Critical Illness Fund—with the aim of providing all households with affordable, equitable, and quality health care. Registration with the SHA is mandatory for all residents, and contributions are structured according to household income.

3.5.1 Coverage: Social Health Authority (SHA) Registration

In Kenya, 42% of women age 15–49 are registered with the SHA. Among those registered, 16% report that the process was very easy and 60% report it was easy. In contrast, 11% found registration difficult, while 3% described it as very difficult (**Table 3.9**).

Patterns by background characteristics

- SHA coverage increases with age, from 22% among women age 15–19 to 52% among women age 30–34.
- SHA coverage increases with level of education. Twenty-four percent of women with no education compared with 60% of women with more than a secondary education report being registered.
- Similarly, coverage increases with household wealth, from 30% among women in the lowest wealth quintile to 46% among those in the highest quintile.
- SHA coverage varies across counties, ranging from a low of 14% in Isiolo to a high of 68% in Taita Taveta. Counties with the lowest coverage are concentrated in the arid and semi-arid north, including Isiolo, Garissa, Tana River, and West Pokot (**Table 3.9C**).

3.5.2 Reasons for Not Registering

Fifty-eight percent of women age 15–49 are not registered with the SHA. Among those not registered, the leading reason was financial barriers, with 37% reporting that registration was too expensive. Other commonly cited reasons were problems with required documentation or inadequate coverage of needed services (24%) and lack of interest or satisfaction with other coverage (17%) (**Tables 3.10** and **3.10C**).

Patterns by background characteristics

- Reasons for SHA non-registration vary by age. Younger women age 15–19 cite problems with required documentation or inadequate coverage of needed services (40%), while women age 30–49 most often report financial barriers (42% to 50%).
- The percentage of women who are not registered with the SHA is similar in urban and rural areas (58% and 59%, respectively). However, the reasons for non-registration differ by residence. For example, problems with required documentation or inadequate coverage of needed services were cited by 15% of women in urban areas compared with 31% of women in rural areas.

- Non-registration with the SHA decreases with level of education, from 76% among women with no education to 40% among those with more than a secondary education. Similarly, lack of awareness as a reason for non-registration declines with level of education, from 22% among women with no education to only 3% among those with more than a secondary education.

3.5.3 Utilisation of Services

Among women age 15–49 registered with the SHA, 53% reported having used SHA services in the 12 months preceding the survey (**Table 3.11**).

Patterns by background characteristics

- Utilisation of SHA services is highest among women age 20–24 years (56%) and lowest among those age 15–19 years (44%).
- Among women registered with the SHA, service utilisation is nearly the same in urban and rural areas (51% and 54%, respectively).
- Use of SHA services in the past 12 months varied by the type of health facility where women most recently sought care. The percentage of women reporting SHA service utilisation was 58% among those who sought care in the public sector, compared with 44% among those who sought care in the private sector and 42% among those who sought care in medical-sector nongovernmental organisations or faith-based organisations.
- Utilisation of SHA services ranges from 35% in Bungoma County to 77% in Elgeyo-Marakwet County (**Table 3.11C**).

3.5.4 Services Accessed

Among women age 15–49 who used SHA services in the 12 months before the survey, the most commonly accessed services were outpatient consultations (62%), followed by antenatal care (9%). Delivery or maternity services and inpatient care/hospitalisation services were each accessed by 8% of women (**Table 3.12**).

Patterns by background characteristics

- Usage of SHA outpatient services/consultations increases slightly with wealth, from 59% among women in the lowest wealth quintile to 69% among women in the highest wealth quintile.
- Laikipia (49%), Lamu (24%), and Bomet (20%) Counties have the highest rates of SHA usage for delivery and maternity services. Wajir (98%), Kwale (83%), and Samburu (80%) Counties have the highest rates of SHA usage for outpatient/chronic care services (**Table 3.12C**).

3.5.5 Challenges Accessing Services

Among women age 15–49 who were registered with the SHA and received care from a health care provider in the 12 months before the survey, 73% reported facing no challenges accessing SHA-covered services. However, 27% reported difficulties accessing those services. The most commonly reported challenges were lack of medicine or supplies (9%), long waiting times (7%), the SHA system being down (7%), and inadequacy of benefits covered (6%) (**Table 3.13**).

Patterns by background characteristics

- Women in urban areas were more likely than those in rural areas (4% versus 1%) to report that the SHA was not accepted at their preferred health facility.
- Challenges accessing SHA services varied by wealth quintile. Affordability-related challenges, specifically unpaid premiums, were more commonly reported by women in the lowest wealth quintile (9%) than by those in the highest quintile (1%). In contrast, women in the highest wealth quintile were more likely than women in the lowest wealth quintile (6% versus 1%) to report that the SHA was not accepted at their preferred health facility.
- The leading challenges in accessing SHA services varied across counties. In Wajir, the main issue was the SHA system being down (42%). In Kitui, top concerns were premiums not being paid (35%) and lack of medicine or supplies (30%). In Machakos, 22% of women cited that some services were not available at their preferred health facility (**Table 3.13C**).

3.5.6 Benefits Received

Among women who used SHA services in the 12 months before the survey, 91% reported reduced out-of-pocket spending as a benefit, while 22% mentioned reduced stress over health expenses (**Table 3.14**).

Patterns by background characteristics

- Among women registered with the SHA, 16% in rural areas compared with 10% in rural areas reported increased access to quality health services as a benefit. Similarly, reduced stress over health expenses was more commonly cited as a benefit among women in rural areas (25%) than among women in urban areas (19%).
- Reduced out-of-pocket spending on health care was the most pronounced benefit in Lamu, Kericho, and Nyamira Counties (100% each) (**Table 3.14C**).

3.5.7 Satisfaction with Services

Among women who received SHA-covered services in the 12 months before the survey, overall satisfaction was high: 92% reported being satisfied with the services they received, and 91% said they would recommend SHA services. Most women also reported positive service experiences, including being treated with respect by health care workers (95%), receiving quality services (94%), and finding SHA services available when needed (92%) (**Table 3.15**).

Patterns by background characteristics

- Ninety-three percent of women in rural areas compared with 87% of women in urban areas are satisfied with the timeliness of SHA services.
- Satisfaction with SHA-covered services declines modestly with level of education. Overall satisfaction decreases from 98% among women with no education to 90% among those with more than a secondary education. Willingness to recommend SHA services also declines, from 95% among women with no education to 88% among those with more than a secondary education.
- Overall satisfaction with SHA services also declines modestly with wealth, from 93% among women in the lowest wealth quintile to 89% among those in the highest wealth quintile.

- Satisfaction with SHA services is generally high in most counties, ranging from 80% to 100% across service measures. Nyandarua County recorded the lowest levels of satisfaction, with 51% of women recommending SHA services and 84% reporting being treated with respect by health care workers (**Table 3.15C**).

List of Tables

For more information on the characteristics of survey respondents, see the following tables:

- **Table 3.1 Background characteristics of respondents**
- **Table 3.1C Background characteristics of respondents: County level**
- **Table 3.2 Education**
- **Table 3.2C Education: County level**
- **Table 3.3 Literacy**
- **Table 3.3C Literacy: County level**
- **Table 3.4 Exposure to mass media**
- **Table 3.4C Exposure to mass media: County level**
- **Table 3.5 Internet usage**
- **Table 3.5C Internet usage: County level**
- **Table 3.6 Place of birth and recent migration**
- **Table 3.6C Place of birth and recent migration: County level**
- **Table 3.7 Type of migration**
- **Table 3.8 Reasons for migration**
- **Table 3.8C Reasons for migration: County level**
- **Table 3.9 Social Health Authority (SHA) registration**
- **Table 3.9C Social Health Authority (SHA) registration: County level**
- **Table 3.10 Reasons for not registering with the Social Health Authority (SHA)**
- **Table 3.10C Reasons for not registering with the Social Health Authority (SHA): County level**
- **Table 3.11 Utilisation of Social Health Authority (SHA) services**
- **Table 3.11C Utilisation of Social Health Authority (SHA) services: County level**
- **Table 3.12 Services received through the Social Health Authority (SHA)**
- **Table 3.12C Services received through the Social Health Authority (SHA): County level**
- **Table 3.13 Challenges accessing Social Health Authority (SHA) services**
- **Table 3.13C Challenges accessing Social Health Authority (SHA) services: County level**
- **Table 3.14 Benefits received through Social Health Authority (SHA) coverage**
- **Table 3.14C Benefits received through Social Health Authority (SHA) coverage: County level**
- **Table 3.15 Satisfaction with services covered under the Social Health Authority (SHA)**
- **Table 3.15C Satisfaction with services covered under the Social Health Authority (SHA): County level**

Table 3.1 Background characteristics of respondents

Percent distribution of women age 15–49 by selected background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Number of women		
	Weighted percentage	Weighted number	Unweighted number
Age			
15–19	18.4	3,916	4,400
20–24	17.2	3,653	3,472
25–29	17.6	3,732	3,482
30–34	14.6	3,108	2,932
35–39	14.1	2,989	2,989
40–44	9.6	2,048	2,146
45–49	8.4	1,784	1,810
Self-reported health status			
Very good	23.4	4,963	5,085
Good	56.5	11,998	12,088
Moderate	16.9	3,583	3,408
Bad	2.7	569	561
Very bad	0.6	118	89
Religion			
Catholic	17.6	3,731	3,476
Protestant	40.8	8,657	8,203
Evangelical church	18.5	3,929	3,663
African-instituted church	9.0	1,906	1,714
Orthodox	0.3	71	77
Other Christian	3.1	661	361
Islam	9.0	1,903	3,357
Hindu	0.0	3	6
Traditionist	0.2	48	117
No religion/atheist	1.5	319	253
Other	0.0	2	4
Marital status			
Never married	32.9	6,983	7,108
Married	47.7	10,124	10,338
Living together	7.3	1,548	1,348
Divorced/separated	9.6	2,038	1,861
Widowed	2.5	538	576
Residence			
Urban	46.3	9,829	7,241
Rural	53.7	11,402	13,990
Education[1]			
No education	5.5	1,161	2,326
Primary	28.5	6,053	6,105
Secondary	46.4	9,849	9,451
More than secondary	19.6	4,167	3,349
Wealth quintile			
Lowest	15.6	3,312	5,181
Second	18.1	3,852	4,446
Middle	19.5	4,133	4,603
Fourth	22.6	4,800	4,209
Highest	24.2	5,134	2,792
Total 15–49	100.0	21,231	21,231

Note: Education categories refer to the highest level of education attended, whether or not that level was completed.

[1] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 3.1C Background characteristics of respondents: County level

Percent distribution of women age 15–49 according to county, Kenya Mini DHS 2025–26

County	Number of women		
	Weighted percentage	Weighted number	Unweighted number
Mombasa	3.8	802.6	478
Kwale	1.8	374.2	563
Kilifi	2.9	608.5	516
Tana River	0.6	137.1	400
Lamu	0.3	53.5	417
Taita Taveta	0.7	153.2	297
Garissa	0.8	168.3	516
Wajir	0.6	130.2	518
Mandera	0.6	123.2	484
Marsabit	0.7	141.8	416
Isiolo	0.4	75.8	431
Meru	2.5	530.3	454
Tharaka-Nithi	0.7	146.8	294
Embu	1.1	239.0	361
Kitui	1.9	412.0	468
Machakos	3.2	677.2	447
Makueni	1.6	349.9	389
Nyandarua	1.1	223.2	260
Nyeri	1.0	222.8	273
Kirinyaga	1.2	258.1	319
Murang'a	1.8	377.0	434
Kiambu	8.0	1,689.1	632
Turkana	0.8	178.0	344
West Pokot	1.1	225.4	444
Samburu	0.5	97.9	395
Trans Nzoia	1.7	370.4	500
Uasin Gishu	3.1	667.1	525
Elgeyo-Marakwet	0.8	177.5	439
Nandi	1.7	360.9	480
Baringo	0.9	187.2	364
Laikipia	0.9	195.2	321
Nakuru	5.1	1,078.4	505
Narok	2.0	427.6	470
Kajiado	2.2	467.4	369
Kericho	1.8	386.9	463
Bomet	2.3	480.4	594
Kakamega	3.1	665.1	526
Vihiga	1.1	233.0	443
Bungoma	2.9	624.9	574
Busia	1.8	372.7	441
Siaya	1.7	369.8	475
Kisumu	2.6	550.6	538
Homa Bay	2.3	485.4	570
Migori	2.4	509.1	586
Kisii	2.1	438.6	442
Nyamira	1.1	237.9	454
Nairobi City	16.7	3,549.9	602
Total 15–49	100.0	21,231.0	21,231

Table 3.2 Education

Percent distribution of women age 15–49 by highest level of schooling attended or completed and median years completed, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Highest level of schooling							Total	Median years completed	Number of women
	No education[1]	Some primary	Completed primary[2]	Some secondary[3]	Completed secondary[4]	More than secondary[5]	Missing			
Age										
15–24	3.0	4.0	6.3	40.7	30.1	15.7	0.1	100.0	8.8	7,516
15–19	2.0	2.9	4.3	65.1	21.3	4.5	0.0	100.0	8.3	3,891
20–24	4.2	5.3	8.5	14.4	39.6	27.8	0.2	100.0	9.5	3,624
25–29	3.7	7.8	16.3	12.3	30.5	29.3	0.0	100.0	9.4	3,700
30–34	6.4	14.1	18.5	10.8	23.0	27.2	0.0	100.0	9.1	3,094
35–39	7.9	18.1	30.8	10.1	16.8	16.3	0.0	100.0	7.7	2,949
40–44	9.9	21.8	27.5	9.0	14.6	17.3	0.0	100.0	7.7	2,034
45–49	9.3	24.2	28.4	8.6	16.7	12.8	0.0	100.0	7.6	1,769
Residence										
Urban	3.9	7.5	13.9	16.4	29.1	29.2	0.0	100.0	9.4	9,779
Rural	6.9	15.1	20.1	25.5	20.8	11.6	0.1	100.0	7.9	11,283
Wealth quintile										
Lowest	20.5	22.7	18.9	25.1	10.5	2.1	0.1	100.0	7.4	3,277
Second	3.1	17.2	23.7	29.0	20.5	6.5	0.1	100.0	7.7	3,812
Middle	2.2	11.5	21.5	25.0	25.8	13.9	0.1	100.0	8.1	4,096
Fourth	1.8	7.1	17.7	18.5	31.9	23.0	0.0	100.0	9.2	4,766
Highest	3.8	4.4	7.3	12.6	29.2	42.7	0.0	100.0	9.8	5,109
Total	5.5	11.6	17.2	21.3	24.7	19.8	0.0	100.0	8.7	21,062

Note: Respondents who reported vocational training as their highest current or previous level of education have been excluded from this table.
[1] No education includes informal education (e.g., madrassa/duksi/adult education).
[2] Completed grade 8 at the primary level or completed grade 8 at the junior school level for those under age 56; because of the change in the school system in the 1980s, those age 56 and older are considered to have completed primary if they completed grade 7.
[3] Those who completed grade 9 at the junior school level are considered to have completed some secondary education.
[4] Completed grade 4 at the secondary level
[5] More than secondary includes middle-level colleges and universities.

Table 3.2C Education: County level

Percent distribution of women age 15–49 by highest level of schooling attended or completed and median years completed, according to county, Kenya Mini DHS 2025–26

County	Highest level of schooling							Total	Median years completed	Number of women
	No education[1]	Some primary	Completed primary[2]	Some secondary[3]	Completed secondary[4]	More than secondary[5]	Missing			
Mombasa	5.5	12.3	22.1	12.9	25.8	21.4	0.0	100.0	9.0	798
Kwale	15.9	19.6	22.2	22.5	10.8	8.7	0.4	100.0	7.5	368
Kilifi	9.6	17.1	21.6	22.3	19.3	10.2	0.0	100.0	7.8	608
Tana River	37.1	11.3	13.3	17.0	14.5	6.1	0.7	100.0	7.8	137
Lamu	13.1	16.2	19.4	24.5	22.4	4.4	0.0	100.0	7.8	53
Taita Taveta	1.3	8.4	29.0	19.9	20.8	20.6	0.0	100.0	8.0	145
Garissa	55.7	11.9	6.2	11.2	7.1	7.7	0.1	100.0	7.8	168
Wajir	61.0	2.9	2.5	16.4	11.6	5.5	0.0	100.0	8.7	130
Mandera	62.4	7.8	6.0	10.1	8.9	4.8	0.1	100.0	8.1	123
Marsabit	38.0	10.7	8.1	19.7	14.9	8.6	0.0	100.0	8.1	141
Isiolo	18.3	16.3	22.2	18.1	17.4	7.4	0.2	100.0	7.8	75
Meru	4.4	22.1	20.0	20.5	20.3	12.7	0.0	100.0	7.8	526
Tharaka-Nithi	1.2	15.4	15.9	26.0	24.0	17.5	0.0	100.0	8.4	140
Embu	0.2	14.2	15.2	21.4	26.9	21.8	0.3	100.0	8.9	236
Kitui	2.2	18.3	31.2	17.9	19.7	10.7	0.0	100.0	7.7	380
Machakos	0.2	3.1	19.4	20.7	31.2	25.4	0.0	100.0	9.2	652
Makueni	0.3	5.2	22.7	22.0	29.6	20.2	0.0	100.0	8.9	336
Nyandarua	0.0	8.3	22.5	23.9	26.5	18.9	0.0	100.0	8.6	222
Nyeri	1.0	3.2	17.9	18.3	38.5	21.0	0.0	100.0	9.3	220
Kirinyaga	0.9	8.1	23.5	16.7	31.6	19.2	0.0	100.0	9.0	258
Murang'a	0.3	11.6	22.1	26.6	23.7	15.7	0.0	100.0	8.2	377
Kiambu	0.9	4.8	10.4	14.0	30.3	39.6	0.0	100.0	9.7	1,689
Turkana	51.8	12.5	4.5	13.5	8.7	8.9	0.1	100.0	7.9	178
West Pokot	18.2	24.7	8.1	19.9	14.4	14.8	0.0	100.0	7.8	223
Samburu	41.1	12.7	5.5	19.2	11.4	10.2	0.0	100.0	8.0	98
Trans Nzoia	1.3	17.7	19.2	26.0	19.5	16.3	0.0	100.0	7.9	369
Uasin Gishu	0.5	8.0	11.8	22.2	27.1	30.2	0.1	100.0	9.3	667
Elgeyo-Marakwet	0.2	8.4	18.7	25.2	24.4	23.1	0.0	100.0	8.7	176
Nandi	0.3	17.8	17.4	26.3	22.9	15.3	0.0	100.0	8.1	361
Baringo	8.5	7.5	17.7	24.9	22.3	19.2	0.0	100.0	8.6	185
Laikipia	8.4	7.9	17.2	16.6	29.6	20.2	0.0	100.0	9.1	194
Nakuru	2.0	10.2	15.4	28.6	25.9	17.8	0.0	100.0	8.6	1,077
Narok	13.1	19.5	17.1	19.9	21.6	8.8	0.0	100.0	7.7	428
Kajiado	9.1	8.5	13.9	15.1	28.2	25.1	0.0	100.0	9.3	464
Kericho	0.2	13.1	18.3	24.5	24.4	19.5	0.0	100.0	8.4	384
Bomet	0.2	15.6	19.4	23.9	24.4	16.6	0.0	100.0	8.2	478
Kakamega	2.1	17.0	19.3	25.9	22.4	13.2	0.2	100.0	7.9	660
Vihiga	0.3	10.3	19.6	33.8	21.1	14.8	0.0	100.0	8.1	233
Bungoma	0.9	14.8	19.4	33.7	16.1	15.2	0.0	100.0	8.0	624
Busia	1.9	17.9	14.4	27.8	25.8	12.3	0.0	100.0	8.0	373
Siaya	2.0	15.3	27.4	28.3	19.2	7.8	0.0	100.0	7.8	370
Kisumu	1.1	11.9	18.5	24.2	22.9	21.4	0.0	100.0	8.5	547
Homa Bay	0.3	16.7	21.2	29.8	17.3	14.2	0.5	100.0	7.8	484
Migori	1.1	16.5	21.7	23.3	22.4	14.9	0.0	100.0	7.9	509
Kisii	0.8	12.7	15.9	31.5	27.6	11.6	0.0	100.0	8.2	439
Nyamira	0.2	8.3	14.3	39.0	24.7	13.2	0.2	100.0	8.2	237
Nairobi City	5.1	8.0	15.3	15.3	30.4	25.8	0.0	100.0	9.3	3,521
Total	5.5	11.6	17.2	21.3	24.7	19.8	0.0	100.0	8.7	21,062

Note: Respondents who reported vocational training as their highest current or previous level of education have been excluded from this table.
[1] No education includes informal education (e.g., madrassa/duksi/adult education).
[2] Completed grade 8 at the primary level or completed grade 8 at the junior school level for those under age 56; because of the change in the school system in the 1980s, those age 56 and older are considered to have completed primary if they completed grade 7.
[3] Those who completed grade 9 at the junior school level are considered to have completed some secondary education.
[4] Completed grade 4 at the secondary level
[5] More than secondary includes middle-level colleges and universities.

Table 3.3 Literacy

Percent distribution of women age 15–49 by level of schooling attended and level of literacy, and percentage literate, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Among women with no, primary, or secondary schooling:						Total	Percent-age literate[1]	Number of women
	Higher than secondary schooling	Can read a whole sentence	Can read part of a sentence	Cannot read at all	No card with required language	Blind/ visually impaired			
Age									
15–24	15.6	75.1	4.7	4.5	0.0	0.0	100.0	95.4	7,569
15–19	4.4	87.6	4.3	3.7	0.0	0.0	100.0	96.3	3,916
20–24	27.6	61.7	5.1	5.5	0.0	0.0	100.0	94.5	3,653
25–29	29.1	58.2	6.6	5.8	0.0	0.2	100.0	94.0	3,732
30–34	27.1	56.2	7.9	8.8	0.0	0.1	100.0	91.1	3,108
35–39	16.1	60.7	11.4	11.5	0.0	0.3	100.0	88.2	2,989
40–44	17.2	56.5	10.7	15.1	0.0	0.6	100.0	84.3	2,048
45–49	12.7	60.9	12.5	13.0	0.0	0.9	100.0	86.1	1,784
Residence									
Urban	29.1	60.4	4.8	5.5	0.0	0.2	100.0	94.2	9,829
Rural	11.5	67.7	10.2	10.3	0.0	0.2	100.0	89.4	11,402
Wealth quintile									
Lowest	2.1	58.0	13.6	26.0	0.1	0.2	100.0	73.8	3,312
Second	6.4	74.6	11.9	6.8	0.0	0.3	100.0	92.9	3,852
Middle	13.8	72.9	8.3	4.5	0.0	0.5	100.0	95.0	4,133
Fourth	22.9	68.1	5.3	3.6	0.0	0.0	100.0	96.3	4,800
Highest	42.5	50.4	2.3	4.6	0.0	0.2	100.0	95.2	5,134
Total	19.6	64.4	7.7	8.1	0.0	0.2	100.0	91.7	21,231

[1] Refers to women who attended schooling higher than the secondary level and women with less schooling who can read a whole sentence or part of a sentence

Table 3.3C Literacy: County level

Percent distribution of women age 15–49 by level of schooling attended and level of literacy, and percentage literate, according to county, Kenya Mini DHS 2025–26

		Among women with no, primary, or secondary schooling:							
County	Higher than secondary schooling	Can read a whole sentence	Can read part of a sentence	Cannot read at all	No card with required language	Blind/ visually impaired	Total	Percent-age literate[1]	Number of women
Mombasa	21.3	63.6	7.3	7.8	0.0	0.0	100.0	92.2	803
Kwale	8.6	67.5	5.1	18.9	0.0	0.0	100.0	81.1	374
Kilifi	10.2	70.2	6.2	13.4	0.0	0.0	100.0	86.6	608
Tana River	6.1	45.9	8.4	39.6	0.0	0.0	100.0	60.4	137
Lamu	4.4	63.6	15.0	16.6	0.0	0.3	100.0	83.0	54
Taita Taveta	19.5	74.2	3.1	2.9	0.0	0.4	100.0	96.8	153
Garissa	7.7	25.9	9.8	56.6	0.0	0.0	100.0	43.4	168
Wajir	5.5	27.5	4.6	62.4	0.0	0.0	100.0	37.6	130
Mandera	4.8	17.0	12.0	66.0	0.0	0.1	100.0	33.8	123
Marsabit	8.6	39.7	10.0	41.8	0.0	0.0	100.0	58.2	142
Isiolo	7.3	60.8	11.7	19.9	0.0	0.3	100.0	79.8	76
Meru	12.6	67.3	10.5	9.3	0.0	0.2	100.0	90.4	530
Tharaka-Nithi	16.6	72.5	2.9	7.6	0.0	0.4	100.0	92.0	147
Embu	21.5	62.0	12.0	3.9	0.0	0.6	100.0	95.5	239
Kitui	9.9	74.0	9.2	7.0	0.0	0.0	100.0	93.0	412
Machakos	24.4	70.4	5.2	0.0	0.0	0.0	100.0	100.0	677
Makueni	19.4	64.7	15.0	0.6	0.0	0.3	100.0	99.1	350
Nyandarua	18.8	75.4	4.5	1.3	0.0	0.0	100.0	98.7	223
Nyeri	20.8	76.1	1.5	1.6	0.0	0.0	100.0	98.4	223
Kirinyaga	19.2	71.3	4.9	4.4	0.0	0.2	100.0	95.3	258
Murang'a	15.7	77.3	3.5	3.3	0.0	0.3	100.0	96.4	377
Kiambu	39.6	55.2	3.0	2.2	0.0	0.0	100.0	97.8	1,689
Turkana	8.9	34.3	3.0	53.8	0.0	0.0	100.0	46.2	178
West Pokot	14.6	40.5	22.5	22.4	0.0	0.0	100.0	77.6	225
Samburu	10.1	42.6	4.2	43.1	0.0	0.0	100.0	56.9	98
Trans Nzoia	16.2	65.5	11.6	6.1	0.1	0.5	100.0	93.3	370
Uasin Gishu	30.2	60.3	5.1	4.3	0.0	0.0	100.0	95.7	667
Elgeyo-Marakwet	23.0	60.1	14.0	3.0	0.0	0.0	100.0	97.0	178
Nandi	15.3	68.8	11.8	3.8	0.0	0.3	100.0	95.9	361
Baringo	19.0	62.1	9.6	8.9	0.0	0.4	100.0	90.7	187
Laikipia	20.1	70.2	3.3	6.0	0.4	0.0	100.0	93.6	195
Nakuru	17.8	68.4	8.4	5.4	0.0	0.0	100.0	94.6	1,078
Narok	8.8	58.0	13.6	19.6	0.0	0.0	100.0	80.4	428
Kajiado	25.0	57.8	4.4	12.8	0.0	0.0	100.0	87.2	467
Kericho	19.3	66.0	10.0	4.6	0.0	0.0	100.0	95.4	387
Bomet	16.5	69.3	13.2	1.1	0.0	0.0	100.0	98.9	480
Kakamega	13.1	67.3	14.1	5.3	0.0	0.2	100.0	94.5	665
Vihiga	14.8	77.6	4.8	2.7	0.0	0.0	100.0	97.3	233
Bungoma	15.1	67.2	13.7	3.5	0.0	0.6	100.0	95.9	625
Busia	12.3	75.5	6.4	5.6	0.3	0.0	100.0	94.1	373
Siaya	7.8	76.2	10.8	4.6	0.0	0.5	100.0	94.8	370
Kisumu	21.3	69.8	5.7	3.2	0.0	0.0	100.0	96.8	551
Homa Bay	14.2	63.9	15.7	6.3	0.0	0.0	100.0	93.7	485
Migori	14.9	71.5	7.4	6.3	0.0	0.0	100.0	93.7	509
Kisii	11.6	77.0	7.9	2.2	0.0	1.3	100.0	96.5	439
Nyamira	13.2	72.4	9.3	3.0	0.0	2.1	100.0	94.9	238
Nairobi City	25.6	62.6	4.8	6.4	0.0	0.6	100.0	93.0	3,550
Total	19.6	64.4	7.7	8.1	0.0	0.2	100.0	91.7	21,231

[1] Refers to women who attended schooling higher than the secondary level and women with less schooling who can read a whole sentence or part of a sentence

Table 3.4 Exposure to mass media

Percentage of women age 15–49 who are exposed to specific media on a weekly basis, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Reads a newspaper at least once a week	Watches television at least once a week	Listens to the radio at least once a week	Accesses all three media at least once a week	Accesses none of the three media at least once a week	Number of women
Age						
15–19	4.5	48.0	48.5	2.1	28.4	3,916
20–24	6.0	47.8	47.3	2.9	29.7	3,653
25–29	4.6	52.7	51.9	2.5	25.9	3,732
30–34	5.7	53.6	52.7	3.5	25.0	3,108
35–39	3.7	49.5	54.0	2.0	26.6	2,989
40–44	5.6	49.1	56.2	3.2	25.1	2,048
45–49	4.6	49.8	59.5	3.8	25.6	1,784
Residence						
Urban	6.4	63.7	44.1	3.6	24.4	9,829
Rural	3.8	38.4	58.7	2.0	29.1	11,402
Education[1]						
No education	0.1	17.7	22.1	0.0	65.4	1,161
Primary	2.2	38.0	54.7	1.3	31.7	6,053
Secondary	5.0	52.7	54.1	2.5	23.4	9,849
More than secondary	10.4	70.5	51.1	6.0	17.5	4,167
Wealth quintile						
Lowest	1.9	7.9	43.9	0.3	53.0	3,312
Second	2.5	26.8	61.2	1.0	30.7	3,852
Middle	4.6	53.0	59.5	2.4	23.2	4,133
Fourth	5.9	65.0	52.8	3.7	21.1	4,800
Highest	8.3	78.4	43.3	4.9	15.6	5,134
Total	5.0	50.1	51.9	2.7	26.9	21,231

[1] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 3.4C Exposure to mass media: County level

Percentage of women age 15–49 who are exposed to specific media on a weekly basis, according to county, Kenya Mini DHS 2025–26

County	Reads a newspaper at least once a week	Watches television at least once a week	Listens to the radio at least once a week	Accesses all three media at least once a week	Accesses none of the three media at least once a week	Number of women
Mombasa	3.7	42.9	25.2	0.9	45.4	803
Kwale	4.0	37.4	52.3	2.2	29.8	374
Kilifi	0.7	21.0	15.6	0.2	69.0	608
Tana River	0.7	28.7	23.9	0.6	60.2	137
Lamu	2.6	38.2	27.8	0.4	44.5	54
Taita Taveta	7.8	59.9	68.9	4.2	12.0	153
Garissa	4.4	14.2	9.0	0.7	78.5	168
Wajir	1.1	7.9	2.1	0.5	90.8	130
Mandera	0.3	11.7	1.4	0.0	87.4	123
Marsabit	1.0	24.2	19.5	0.0	59.2	142
Isiolo	1.7	41.2	34.3	0.6	47.0	76
Meru	1.8	35.3	42.1	1.6	42.2	530
Tharaka-Nithi	9.1	36.9	55.9	4.0	26.3	147
Embu	2.1	53.4	42.2	1.4	26.5	239
Kitui	1.5	25.9	60.1	0.9	29.0	412
Machakos	13.0	62.3	62.3	8.4	17.7	677
Makueni	5.9	40.9	77.5	3.0	13.3	350
Nyandarua	4.7	75.1	69.5	2.6	10.5	223
Nyeri	3.5	66.4	55.6	2.9	21.3	223
Kirinyaga	3.6	65.8	51.9	1.8	19.2	258
Murang'a	3.2	62.1	57.7	1.0	20.1	377
Kiambu	7.6	78.4	57.3	3.8	12.5	1,689
Turkana	1.7	17.0	29.5	0.7	62.1	178
West Pokot	11.4	20.4	54.2	4.4	38.7	225
Samburu	3.1	25.2	54.9	1.0	32.3	98
Trans Nzoia	5.1	58.7	54.1	3.4	18.9	370
Uasin Gishu	6.5	61.5	58.7	3.6	18.1	667
Elgeyo-Marakwet	6.3	32.0	63.9	3.9	27.8	178
Nandi	6.5	49.1	71.0	4.0	16.2	361
Baringo	3.0	29.5	61.0	2.0	31.6	187
Laikipia	3.3	61.7	55.8	2.0	22.9	195
Nakuru	5.1	54.3	57.6	2.5	21.6	1,078
Narok	1.7	22.2	44.9	0.0	46.0	428
Kajiado	2.9	46.0	31.4	1.3	37.7	467
Kericho	9.4	46.5	72.5	6.1	16.7	387
Bomet	0.8	26.1	67.1	0.2	27.1	480
Kakamega	3.0	38.4	61.1	1.5	25.8	665
Vihiga	5.9	44.2	70.8	3.3	15.7	233
Bungoma	3.9	35.3	72.6	1.5	18.9	625
Busia	3.5	43.5	60.9	1.5	24.1	373
Siaya	1.2	39.4	66.0	0.6	21.3	370
Kisumu	4.9	53.5	55.5	2.9	21.9	551
Homa Bay	4.4	35.7	63.2	2.8	23.8	485
Migori	4.8	62.6	72.4	4.1	16.3	509
Kisii	8.2	44.9	65.6	5.1	21.4	439
Nyamira	2.8	39.4	62.8	1.9	26.7	238
Nairobi City	6.3	66.7	41.4	3.7	22.2	3,550
Total	5.0	50.1	51.9	2.7	26.9	21,231

Table 3.5 Internet usage

Percentage of women age 15–49 who have ever used the internet; percentage who have used the internet in the past 12 months; and among women who have used the internet in the past 12 months, percent distribution by frequency of internet use in the past month, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Ever used the internet	Used the internet in the past 12 months	Number	Among respondents who have used the internet in the past 12 months, percentage who in the past month used the internet: Almost every day	At least once a week	Less than once a week	Not at all	Total	Number
Age									
15–19	50.3	46.9	3,916	51.1	26.7	17.0	5.1	100.0	1,837
20–24	78.3	75.8	3,653	72.5	18.4	7.4	1.8	100.0	2,769
25–29	77.9	75.6	3,732	69.4	21.3	6.4	2.9	100.0	2,820
30–34	70.2	66.5	3,108	68.9	22.8	6.3	2.1	100.0	2,065
35–39	57.8	53.9	2,989	60.3	28.3	9.1	2.3	100.0	1,612
40–44	54.0	49.9	2,048	58.1	28.2	8.4	5.3	100.0	1,022
45–49	46.3	41.8	1,784	54.9	31.6	12.0	1.5	100.0	746
Residence									
Urban	83.7	80.8	9,829	72.5	19.2	6.1	2.3	100.0	7,944
Rural	47.0	43.2	11,402	51.5	31.0	13.6	3.9	100.0	4,928
Education[1]									
No education	26.1	25.0	1,161	62.2	22.1	11.8	3.9	100.0	291
Primary	44.5	39.7	6,053	45.9	34.2	14.9	5.0	100.0	2,404
Secondary	66.8	63.2	9,849	59.5	26.7	10.7	3.1	100.0	6,229
More than secondary	96.2	94.8	4,167	83.8	12.8	2.3	1.1	100.0	3,949
Wealth quintile									
Lowest	22.9	20.1	3,312	36.8	34.6	23.3	5.2	100.0	666
Second	42.7	38.4	3,852	40.5	36.6	17.0	6.0	100.0	1,478
Middle	62.4	57.6	4,133	52.6	31.5	12.7	3.2	100.0	2,379
Fourth	80.6	76.8	4,800	62.0	26.7	8.2	3.1	100.0	3,687
Highest	92.1	90.8	5,134	84.0	11.7	3.0	1.3	100.0	4,664
Total	64.0	60.6	21,231	64.5	23.7	8.9	2.9	100.0	12,873

[1] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 3.5C Internet usage: County level

Percentage of women age 15–49 who have ever used the internet; percentage who have used the internet in the past 12 months; and among women who have used the internet in the past 12 months, percent distribution by frequency of internet use in the past month, according to county, Kenya Mini DHS 2025–26

County	Ever used the internet	Used the internet in the past 12 months	Number	Among respondents who have used the internet in the past 12 months, percentage who in the past month used the internet: Almost every day	At least once a week	Less than once a week	Not at all	Total	Number
Mombasa	78.8	76.2	803	64.9	23.3	11.4	0.4	100.0	611
Kwale	48.1	45.9	374	63.2	29.4	6.9	0.5	100.0	172
Kilifi	57.0	56.3	608	51.8	28.8	17.9	1.6	100.0	343
Tana River	50.9	46.2	137	35.6	28.6	35.5	0.4	100.0	63
Lamu	59.0	52.5	54	64.7	24.9	9.3	1.1	100.0	28
Taita Taveta	76.3	70.1	153	58.7	31.9	3.6	5.8	100.0	107
Garissa	53.9	51.0	168	63.6	21.6	11.7	3.1	100.0	86
Wajir	44.1	40.3	130	73.6	12.4	7.9	6.1	100.0	52
Mandera	32.3	30.6	123	68.8	16.4	6.5	8.3	100.0	38
Marsabit	37.9	37.7	142	60.7	31.1	8.2	0.0	100.0	53
Isiolo	47.9	44.4	76	50.0	33.9	14.0	2.1	100.0	34
Meru	53.0	47.3	530	60.2	21.5	10.8	7.5	100.0	251
Tharaka-Nithi	63.7	58.8	147	53.7	31.4	8.0	7.0	100.0	86
Embu	61.3	58.6	239	61.8	25.7	10.0	2.6	100.0	140
Kitui	44.6	40.0	412	53.6	27.3	15.5	3.6	100.0	165
Machakos	71.6	69.2	677	70.5	21.5	6.1	1.9	100.0	469
Makueni	73.4	61.8	350	60.0	27.4	4.1	8.4	100.0	216
Nyandarua	65.7	59.0	223	54.5	31.8	8.8	4.9	100.0	132
Nyeri	65.8	63.9	223	65.8	25.1	8.6	0.4	100.0	142
Kirinyaga	71.8	67.4	258	60.9	24.7	10.7	3.7	100.0	174
Murang'a	69.2	59.7	377	53.4	26.3	17.8	2.5	100.0	225
Kiambu	82.7	79.5	1,689	85.2	10.6	3.3	0.9	100.0	1,344
Turkana	17.8	17.8	178	62.3	26.8	8.2	2.7	100.0	32
West Pokot	22.1	21.1	225	65.8	30.8	3.4	0.0	100.0	48
Samburu	28.0	26.6	98	48.7	38.7	5.0	7.6	100.0	26
Trans Nzoia	54.4	54.1	370	62.2	24.9	12.8	0.0	100.0	200
Uasin Gishu	74.1	71.7	667	67.3	22.5	8.3	1.9	100.0	478
Elgeyo-Marakwet	44.3	43.0	178	70.7	21.4	7.3	0.6	100.0	76
Nandi	44.0	41.6	361	69.3	22.1	6.5	2.1	100.0	150
Baringo	47.3	42.8	187	44.2	32.1	18.5	5.2	100.0	80
Laikipia	55.6	53.4	195	71.4	21.9	5.5	1.2	100.0	104
Nakuru	63.2	60.4	1,078	59.5	27.2	9.4	3.9	100.0	652
Narok	39.3	34.9	428	64.8	30.0	4.3	0.9	100.0	149
Kajiado	69.0	64.1	467	70.2	14.6	13.7	1.6	100.0	299
Kericho	50.6	48.6	387	56.4	31.0	11.4	1.1	100.0	188
Bomet	35.8	35.0	480	60.0	29.2	10.3	0.5	100.0	168
Kakamega	62.1	56.9	665	47.2	31.4	15.2	6.2	100.0	378
Vihiga	60.7	52.5	233	49.4	31.5	5.9	13.2	100.0	122
Bungoma	36.8	34.2	625	58.7	30.4	7.1	3.8	100.0	214
Busia	49.6	48.1	373	64.2	22.4	9.7	3.7	100.0	179
Siaya	47.1	45.1	370	37.2	40.7	21.7	0.4	100.0	167
Kisumu	61.2	56.9	551	59.6	24.4	9.1	6.9	100.0	313
Homa Bay	51.9	50.3	485	38.7	41.4	17.4	2.5	100.0	244
Migori	64.1	57.5	509	46.4	28.6	18.6	6.4	100.0	293
Kisii	41.6	40.3	439	59.6	33.5	5.9	1.1	100.0	177
Nyamira	45.0	42.1	238	49.1	39.7	7.2	4.0	100.0	100
Nairobi City	90.5	87.4	3,550	70.3	20.8	6.0	2.9	100.0	3,103
Total	64.0	60.6	21,231	64.5	23.7	8.9	2.9	100.0	12,873

Table 3.6 Place of birth and recent migration

Percent distribution of women age 15–49 who have always lived in their current place of residence, who were born in Kenya but outside of their current place of residence, and who were born in another country; and among women who were born outside of their current place of residence, percentage who moved to their current place of residence in the past 5 years, according to background characteristics, Kenya Mini DHS 2025–26

	Percent distribution by residence and place of birth					Among women who were born outside of current place of residence:	
Background characteristic	Always lived in current place of residence[1]	Born in Kenya but outside of current place of residence	Born outside of Kenya	Total	Number of women	Percentage who moved to current place of residence in the past 5 years	Number of women[2]
Age							
15–19	70.5	26.5	3.0	100.0	3,851	44.4	1,136
20–24	46.9	50.1	3.0	100.0	3,600	67.1	1,912
25–29	43.1	53.9	3.0	100.0	3,683	41.5	2,096
30–34	44.4	52.5	3.2	100.0	3,082	23.1	1,715
35–39	49.3	49.5	1.2	100.0	2,978	13.5	1,511
40–44	50.8	47.1	2.0	100.0	2,043	12.3	1,005
45–49	50.2	47.7	2.1	100.0	1,779	10.0	887
Residence							
Urban	32.1	63.0	4.9	100.0	9,729	37.6	6,609
Rural	67.6	31.8	0.6	100.0	11,287	26.9	3,653
Education[3]							
No education	69.3	22.8	7.9	100.0	1,149	30.0	353
Primary	53.4	44.1	2.5	100.0	6,036	21.3	2,815
Secondary	52.9	44.8	2.3	100.0	9,721	37.9	4,578
More than secondary	38.8	59.1	2.1	100.0	4,112	40.8	2,516
Wealth quintile							
Lowest	76.1	23.2	0.7	100.0	3,290	23.9	786
Second	65.8	33.6	0.6	100.0	3,829	24.6	1,310
Middle	58.3	40.8	0.8	100.0	4,080	32.9	1,700
Fourth	40.3	58.3	1.3	100.0	4,749	37.8	2,833
Highest	28.3	63.7	8.0	100.0	5,069	36.6	3,632
Total	51.2	46.2	2.6	100.0	21,017	33.8	10,261

Note: Respondents who are visitors in the household are excluded from this table.
[1] May include respondents who were born elsewhere in Kenya but moved to their current place of residence when very young
[2] Includes respondents who reported both that they were born outside of Kenya and that they have always lived in their current place of residence; such respondents assumed not to have moved in the past 5 years
[3] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level

Table 3.6C Place of birth and recent migration: County level

Percent distribution of women age 15–49 who have always lived in their current place of residence, who were born in Kenya but outside of their current place of residence, and who were born in another country; and among women who were born outside of their current place of residence, percentage who moved to their current place of residence in the past 5 years, according to county, Kenya Mini DHS 2025–26

	Percent distribution by residence and place of birth					Among women who were born outside of current place of residence:	
County	Always lived in current place of residence[1]	Born in Kenya but outside of current place of residence	Born outside of Kenya	Total	Number of women	Percentage who moved to current place of residence in the past 5 years	Number of women[2]
Mombasa	33.0	65.7	1.3	100.0	789	37.7	529
Kwale	85.0	14.7	0.3	100.0	369	37.9	55
Kilifi	84.0	16.0	0.0	100.0	607	41.3	97
Tana River	84.0	16.0	0.0	100.0	137	21.9	22
Lamu	69.0	30.8	0.2	100.0	53	24.3	16
Taita Taveta	44.1	53.8	2.1	100.0	146	29.6	82
Garissa	85.2	13.8	1.0	100.0	168	43.3	25
Wajir	100.0	0.0	0.0	100.0	130	*	0
Mandera	98.3	0.3	1.4	100.0	123	*	2
Marsabit	87.4	7.2	5.4	100.0	140	16.1	18
Isiolo	77.1	22.7	0.2	100.0	75	40.9	17
Meru	85.6	14.4	0.0	100.0	530	25.3	76
Tharaka-Nithi	78.2	21.8	0.0	100.0	146	28.4	32
Embu	58.4	41.5	0.1	100.0	238	25.3	99
Kitui	70.8	29.2	0.0	100.0	409	31.4	119
Machakos	47.9	52.1	0.0	100.0	673	52.8	351
Makueni	72.4	27.6	0.0	100.0	346	32.5	96
Nyandarua	51.8	48.2	0.0	100.0	218	34.3	105
Nyeri	65.7	34.3	0.0	100.0	221	37.3	76
Kirinyaga	64.5	35.5	0.0	100.0	257	38.0	91
Murang'a	71.8	28.0	0.3	100.0	368	42.6	104
Kiambu	35.1	63.5	1.4	100.0	1,686	48.7	1,094
Turkana	90.6	9.2	0.1	100.0	178	(15.7)	17
West Pokot	88.6	10.5	0.8	100.0	225	28.1	26
Samburu	88.4	11.6	0.0	100.0	97	28.3	11
Trans Nzoia	68.1	30.9	1.0	100.0	370	15.4	118
Uasin Gishu	44.3	54.9	0.8	100.0	660	35.5	368
Elgeyo-Marakwet	82.4	17.1	0.5	100.0	177	42.9	31
Nandi	74.4	24.8	0.8	100.0	361	20.0	93
Baringo	86.3	13.7	0.0	100.0	187	45.5	26
Laikipia	22.6	77.4	0.0	100.0	192	24.3	149
Nakuru	50.9	49.1	0.0	100.0	1,073	32.1	526
Narok	32.8	67.0	0.3	100.0	428	19.2	288
Kajiado	37.7	58.2	4.1	100.0	460	46.8	287
Kericho	73.4	26.6	0.0	100.0	384	34.0	102
Bomet	78.6	21.4	0.0	100.0	469	40.0	100
Kakamega	44.2	55.1	0.7	100.0	659	37.4	368
Vihiga	51.8	47.7	0.5	100.0	230	35.8	111
Bungoma	73.7	25.9	0.4	100.0	624	27.6	164
Busia	45.7	49.6	4.6	100.0	367	23.9	199
Siaya	69.0	30.3	0.7	100.0	366	24.6	113
Kisumu	49.5	49.6	0.9	100.0	543	34.3	274
Homa Bay	60.1	38.7	1.1	100.0	478	27.1	191
Migori	41.2	55.6	3.1	100.0	497	43.7	292
Kisii	22.1	77.8	0.1	100.0	434	13.7	338
Nyamira	82.2	17.5	0.3	100.0	235	23.7	42
Nairobi City	16.3	72.0	11.6	100.0	3,493	29.7	2,923
Total	51.2	46.2	2.6	100.0	21,017	33.8	10,261

Notes: Figures in parentheses are based on 25–49 unweighted cases. An asterisk indicates that a figure is based on fewer than 25 unweighted cases and has been suppressed. Respondents who are visitors in the household are excluded from this table.

[1] May include respondents who were born elsewhere in Kenya but moved to their current place of residence when very young

[2] Includes respondents who reported both that they were born outside of Kenya and that they have always lived in their current place of residence; such respondents assumed not to have moved in the past 5 years.

Table 3.7 Type of migration

Percent distribution of women age 15–49 who have moved to their current place of residence in the past 5 years by type of migration, according to age, Kenya Mini DHS 2025–26

Age	Type of migration				Total	Number
	Urban to urban	Urban to rural	Rural to urban	Rural to rural		
15–19	28.0	12.3	39.4	20.3	100.0	504
20–24	33.1	10.0	43.1	13.9	100.0	1,284
25–29	39.9	12.4	34.0	13.7	100.0	869
30–34	34.3	14.9	36.3	14.5	100.0	396
35–39	28.9	22.1	27.2	21.9	100.0	204
40–44	38.7	28.6	19.7	13.0	100.0	123
45–49	43.2	15.6	25.9	15.4	100.0	89
Total 15–49	34.4	13.0	37.3	15.3	100.0	3,468

Note: Type of migration is based on categorising the previous place of residence and the current place of residence as urban or rural. The previous place of residence is the place the person moved from just before moving to the current place of residence.

Table 3.8 Reasons for migration

Percent distribution of women age 15–49 who have moved to their current place of residence by reason for migration, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Economic reasons/ business	Education/ training	Marriage formation	Family reunification/ other family-related reason	Forced displacement	Other	Total	Number of women
Age								
15–19	12.0	18.5	4.5	59.2	4.9	0.9	100.0	1,133
20–24	35.9	14.9	16.6	30.4	1.4	0.9	100.0	1,911
25–29	43.6	7.9	22.4	23.2	2.0	0.9	100.0	2,096
30–34	41.5	7.3	25.2	23.0	2.4	0.6	100.0	1,715
35–39	43.7	5.0	28.1	21.3	1.5	0.3	100.0	1,511
40–44	39.0	7.7	27.0	22.6	3.1	0.6	100.0	1,004
45–49	36.8	3.8	29.6	24.3	4.8	0.6	100.0	886
Timing of move to current place of residence								
0–4 years	43.9	10.0	16.4	27.6	1.3	0.8	100.0	3,468
5–9 years	43.4	10.1	19.6	23.2	3.4	0.3	100.0	2,325
10 years or more	28.9	8.8	27.0	31.4	3.0	0.8	100.0	4,462
Type of migration[1]								
Urban to urban	45.1	12.2	9.7	29.1	1.8	1.9	100.0	1,193
Urban to rural	22.9	5.8	20.3	48.9	1.5	0.6	100.0	451
Rural to urban	56.6	11.2	12.2	18.7	1.2	0.1	100.0	1,294
Rural to rural	28.5	5.7	38.0	27.5	0.3	0.0	100.0	531
Residence								
Urban	48.7	12.6	10.8	24.1	3.0	0.7	100.0	6,605
Rural	16.6	3.8	41.5	35.7	1.7	0.7	100.0	3,651
Wealth quintile								
Lowest	10.5	3.5	46.9	35.3	3.2	0.6	100.0	785
Second	14.9	3.6	44.1	35.0	1.8	0.6	100.0	1,309
Middle	29.7	6.9	30.9	30.7	1.2	0.5	100.0	1,700
Fourth	48.9	8.3	16.2	25.4	0.7	0.4	100.0	2,833
Highest	45.6	15.0	8.2	25.4	4.7	1.1	100.0	3,629
Total	37.3	9.5	21.7	28.3	2.5	0.7	100.0	10,256

Notes: Respondents who are visitors in the household are excluded from this table. Respondents who stated both that they were born outside of Kenya and that they have always lived in their current place of residence were not asked about the reason for migration and are excluded from this table.
[1]Restricted to respondents who migrated within the past 5 years

Table 3.8C Reasons for migration: County level

Percent distribution of women age 15–49 who have moved to their current place of residence by reason for migration, according to county, Kenya Mini DHS 2025–26

County	Economic reasons/ business	Education/ training	Marriage formation	Family reunification/ other family-related reason	Forced displacement	Other	Total	Number of women
Mombasa	49.1	14.5	5.4	31.0	0.0	0.0	100.0	526
Kwale	36.4	3.3	27.0	27.5	3.2	2.6	100.0	55
Kilifi	34.4	6.1	31.0	28.5	0.0	0.0	100.0	97
Tana River	53.3	0.0	14.3	32.5	0.0	0.0	100.0	22
Lamu	26.3	1.4	20.1	39.7	11.1	1.4	100.0	16
Taita Taveta	26.9	6.8	19.1	46.6	0.0	0.5	100.0	82
Garissa	40.8	11.2	21.6	25.3	1.1	0.0	100.0	25
Mandera	*	*	*	*	*	*	100.0	2
Marsabit	36.9	0.0	28.5	34.7	0.0	0.0	100.0	16
Isiolo	49.9	3.2	20.5	26.5	0.0	0.0	100.0	17
Meru	24.2	7.5	30.5	25.7	12.2	0.0	100.0	76
Tharaka-Nithi	34.3	10.6	23.3	23.2	3.0	5.5	100.0	32
Embu	31.5	11.9	13.8	31.4	0.0	11.4	100.0	99
Kitui	13.9	3.3	40.6	35.8	0.0	6.4	100.0	119
Machakos	50.0	2.8	27.9	18.9	0.4	0.0	100.0	351
Makueni	32.0	8.8	31.0	28.3	0.0	0.0	100.0	96
Nyandarua	41.7	3.7	25.3	21.7	7.3	0.4	100.0	105
Nyeri	33.6	7.9	29.5	29.0	0.0	0.0	100.0	76
Kirinyaga	35.0	7.9	27.6	28.1	0.6	0.9	100.0	91
Murang'a	28.0	5.0	19.6	47.0	0.4	0.0	100.0	104
Kiambu	52.6	13.5	12.2	18.3	2.2	1.1	100.0	1,094
Turkana	(26.3)	(4.6)	(36.3)	(31.2)	(1.6)	(0.0)	100.0	17
West Pokot	30.8	1.7	34.3	32.5	0.0	0.7	100.0	26
Samburu	38.3	5.4	16.7	37.2	2.4	0.0	100.0	11
Trans Nzoia	18.3	5.4	41.4	32.7	0.3	1.9	100.0	118
Uasin Gishu	29.2	13.6	23.3	32.1	1.8	0.0	100.0	368
Elgeyo-Marakwet	27.0	2.0	44.4	23.1	3.6	0.0	100.0	31
Nandi	26.5	4.1	35.5	33.9	0.0	0.0	100.0	93
Baringo	27.7	8.3	31.6	32.3	0.0	0.0	100.0	26
Laikipia	21.6	1.2	20.1	54.2	2.9	0.0	100.0	149
Nakuru	39.0	4.8	27.0	26.0	3.2	0.0	100.0	526
Narok	21.4	4.2	37.8	32.8	2.1	1.7	100.0	288
Kajiado	52.2	7.0	15.5	21.6	1.0	2.8	100.0	287
Kericho	31.8	5.8	27.2	35.2	0.0	0.0	100.0	102
Bomet	18.5	10.4	45.6	25.4	0.0	0.0	100.0	100
Kakamega	14.8	1.8	46.6	35.0	0.9	0.8	100.0	368
Vihiga	5.8	10.5	23.8	59.5	0.0	0.5	100.0	111
Bungoma	13.4	5.3	39.6	40.2	0.8	0.8	100.0	164
Busia	15.9	1.9	25.6	55.1	1.2	0.3	100.0	199
Siaya	11.2	4.4	40.9	42.6	0.9	0.0	100.0	113
Kisumu	28.2	10.6	34.1	26.6	0.5	0.0	100.0	274
Homa Bay	8.7	8.2	55.9	27.1	0.0	0.0	100.0	191
Migori	18.7	8.5	35.4	35.9	1.1	0.3	100.0	290
Kisii	6.9	2.6	59.2	30.7	0.6	0.0	100.0	338
Nyamira	13.4	6.9	61.4	16.6	1.6	0.0	100.0	42
Nairobi City	50.3	14.0	6.0	23.7	5.4	0.5	100.0	2,923
Total	37.3	9.5	21.7	28.3	2.5	0.7	100.0	10,256

Notes: Respondents who are visitors in the household are excluded from this table. Respondents who stated both that they were born outside of Kenya and that they have always lived in their current place of residence were not asked about the reason for migration and are excluded from this table. Figures in parentheses are based on 25–49 unweighted cases. An asterisk indicates that a figure is based on fewer than 25 unweighted cases and has been suppressed.

Table 3.9 Social Health Authority (SHA) registration

Percentage of women age 15–49 years registered with the Social Health Authority (SHA) scheme; and among those registered, percent distribution by how easy it was to register/enroll in the SHA scheme, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Percentage of women registered with SHA	Number of women	Ease of registration among those registered in the SHA scheme					Total	Number of women registered with SHA
			Very easy	Easy	Difficult	Very difficult	Don't know		
Age									
15–19	21.6	3,916	7.0	39.9	4.8	0.5	47.9	100.0	847
20–24	32.9	3,653	17.4	64.9	9.7	1.3	6.8	100.0	1,202
25–29	51.0	3,732	19.0	63.1	10.6	2.4	5.0	100.0	1,903
30–34	51.6	3,108	19.0	60.8	11.8	3.2	5.1	100.0	1,604
35–39	50.0	2,989	15.4	58.9	15.4	4.6	5.6	100.0	1,495
40–44	48.2	2,048	16.4	61.6	12.8	3.6	5.5	100.0	988
45–49	47.3	1,784	14.4	67.2	9.2	3.7	5.5	100.0	844
Residence									
Urban	42.4	9,829	19.0	62.4	9.1	2.6	6.8	100.0	4,166
Rural	41.4	11,402	14.0	58.3	12.8	3.0	11.9	100.0	4,718
Education[1]									
No education	24.2	1,161	6.4	69.3	13.8	2.2	8.3	100.0	282
Primary	39.1	6,053	12.4	63.2	14.2	4.0	6.2	100.0	2,367
Secondary	38.0	9,849	15.2	57.3	9.8	2.8	14.9	100.0	3,743
More than secondary	59.8	4,167	22.9	60.8	9.6	1.9	4.8	100.0	2,492
Wealth quintile									
Lowest	29.7	3,312	10.6	59.9	14.8	4.9	9.7	100.0	983
Second	39.4	3,852	12.9	57.9	14.3	3.1	11.8	100.0	1,516
Middle	44.4	4,133	14.6	60.0	11.9	2.7	10.8	100.0	1,837
Fourth	46.0	4,800	16.2	61.5	10.3	3.1	9.0	100.0	2,209
Highest	45.5	5,134	22.4	60.9	7.4	1.7	7.5	100.0	2,338
Total 15–49	41.8	21,231	16.3	60.2	11.1	2.8	9.5	100.0	8,883

[1] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 3.9C Social Health Authority (SHA) registration: County level

Percentage of women age 15–49 years registered with the Social Health Authority (SHA) scheme; and among those registered, percent distribution by how easy it was to register/enroll in the SHA scheme, according to county, Kenya Mini DHS 2025–26

County	Percentage of women registered with SHA	Number of women	Ease of registration among those registered in the SHA scheme					Total	Number of women registered with SHA
			Very easy	Easy	Difficult	Very difficult	Don't know		
Mombasa	48.3	803	8.1	80.5	6.5	0.9	3.9	100.0	388
Kwale	38.8	374	0.0	90.1	6.6	0.0	3.3	100.0	145
Kilifi	41.8	608	15.1	80.1	3.1	0.5	1.2	100.0	254
Tana River	22.8	137	2.5	75.9	18.4	0.0	3.3	100.0	31
Lamu	33.5	54	4.6	80.4	12.9	2.1	0.0	100.0	18
Taita Taveta	68.0	153	9.8	60.2	19.6	5.3	5.1	100.0	104
Garissa	22.5	168	3.0	90.7	1.0	3.5	1.8	100.0	38
Wajir	28.0	130	0.5	52.2	9.4	0.0	37.9	100.0	36
Mandera	24.7	123	14.8	80.2	3.1	0.9	1.1	100.0	30
Marsabit	27.5	142	0.0	74.2	6.6	0.3	18.8	100.0	39
Isiolo	14.4	76	9.6	74.0	13.8	0.0	2.6	100.0	11
Meru	31.1	530	11.7	66.8	15.9	2.1	3.4	100.0	165
Tharaka-Nithi	49.5	147	7.0	55.4	21.3	11.5	4.7	100.0	73
Embu	61.5	239	6.0	64.3	11.5	1.0	17.3	100.0	147
Kitui	37.0	412	4.1	76.9	11.4	2.0	5.5	100.0	153
Machakos	41.5	677	9.7	73.9	11.8	3.9	0.8	100.0	281
Makueni	48.7	350	17.7	72.8	7.6	1.3	0.6	100.0	171
Nyandarua	52.3	223	6.2	57.1	25.9	6.4	4.4	100.0	117
Nyeri	65.3	223	14.0	45.5	12.3	10.3	17.9	100.0	146
Kirinyaga	66.0	258	4.7	56.9	20.2	2.6	15.7	100.0	170
Murang'a	25.2	377	21.6	63.1	9.2	2.9	3.2	100.0	95
Kiambu	39.9	1,689	51.1	31.6	7.0	4.3	6.0	100.0	675
Turkana	44.9	178	9.4	51.2	23.7	5.0	10.6	100.0	80
West Pokot	24.0	225	9.1	85.1	5.9	0.0	0.0	100.0	54
Samburu	55.1	98	13.4	70.0	7.3	1.3	8.0	100.0	54
Trans Nzoia	30.6	370	14.1	57.8	6.1	6.9	15.1	100.0	113
Uasin Gishu	41.2	667	5.8	71.7	8.2	2.1	12.3	100.0	275
Elgeyo-Marakwet	64.2	178	16.1	54.4	5.5	0.8	23.3	100.0	114
Nandi	51.1	361	32.8	40.7	10.6	1.9	14.0	100.0	185
Baringo	52.8	187	25.6	36.6	16.5	1.0	20.3	100.0	99
Laikipia	36.6	195	12.0	71.4	9.2	2.3	5.2	100.0	72
Nakuru	45.0	1,078	19.7	49.1	12.6	5.1	13.4	100.0	486
Narok	24.1	428	12.5	51.8	12.5	5.0	18.2	100.0	103
Kajiado	48.6	467	6.9	72.6	7.7	4.7	8.0	100.0	227
Kericho	46.9	387	6.2	65.3	5.7	2.5	20.3	100.0	182
Bomet	55.4	480	51.0	24.5	6.7	0.0	17.9	100.0	266
Kakamega	46.9	665	6.7	64.6	16.7	5.2	6.8	100.0	312
Vihiga	51.6	233	1.7	66.4	12.8	2.2	16.9	100.0	120
Bungoma	40.4	625	4.9	64.6	10.7	2.2	17.5	100.0	253
Busia	42.6	373	22.2	32.4	14.3	4.6	26.5	100.0	159
Siaya	34.9	370	5.1	70.5	12.4	0.5	11.5	100.0	129
Kisumu	52.0	551	21.7	42.1	20.8	8.1	7.4	100.0	286
Homa Bay	41.2	485	42.0	43.7	9.8	3.3	1.3	100.0	200
Migori	39.4	509	4.7	72.7	8.0	2.2	12.4	100.0	201
Kisii	35.1	439	6.4	62.1	12.9	3.6	15.0	100.0	154
Nyamira	41.8	238	0.9	53.3	10.4	9.4	26.1	100.0	100
Nairobi City	38.7	3,550	13.4	69.1	11.5	0.0	5.9	100.0	1,374
Total 15–49	41.8	21,231	16.3	60.2	11.1	2.8	9.5	100.0	8,883

Table 3.10 Reasons for not registering with the Social Health Authority (SHA) scheme

Percentage of women age 15–49 years not registered with the Social Health Authority (SHA) scheme, by reason for not registering, according to background characteristics, Kenya Mini DHS 2025–26

			Among women not registered, reason for not registering:								
Background characteristic	Percentage of women not registered with SHA	Number of women	Inconvenient/ burdensome	Have not heard of SHA/not aware	Aware but don't know how to register	Distrust in system/ government	Financial barriers/ too expensive	Problems with required documentation/ inadequate coverage of needed services	Not interested/ satisfied with other coverage	Other	Number of women not registered with SHA
Age											
15–19	78.4	3,916	3.9	7.0	23.2	2.1	24.9	39.7	10.9	4.7	3,069
20–24	67.1	3,653	7.6	4.7	17.4	4.3	33.0	25.9	19.8	2.4	2,450
25–29	49.0	3,732	11.9	3.8	12.9	6.9	37.4	17.8	21.9	2.3	1,829
30–34	48.4	3,108	13.2	3.6	14.5	7.9	41.6	15.5	18.6	4.2	1,505
35–39	50.0	2,989	10.7	3.9	14.8	8.6	44.7	14.2	18.2	2.7	1,494
40–44	51.8	2,048	9.3	4.1	13.4	9.8	47.3	15.1	17.2	1.6	1,060
45–49	52.7	1,784	10.3	4.4	14.1	6.9	49.6	13.5	16.8	3.4	940
Residence											
Urban	57.6	9,829	10.2	4.8	17.1	7.1	27.3	15.1	27.0	3.0	5,664
Rural	58.6	11,402	7.5	4.8	16.8	4.6	44.5	30.8	8.7	3.5	6,684
Education[1]											
No education	75.8	1,161	5.9	22.4	26.8	3.3	32.3	16.0	11.8	2.4	880
Primary	60.9	6,053	8.5	3.9	14.1	6.2	49.5	21.7	13.0	2.0	3,686
Secondary	62.0	9,849	8.6	3.4	18.2	4.8	32.6	29.3	15.7	3.7	6,107
More than secondary	40.2	4,167	11.2	2.6	13.3	9.3	25.0	11.0	34.0	4.6	1,675
Wealth quintile											
Lowest	70.3	3,312	6.7	9.2	19.9	3.6	48.3	32.1	4.8	2.4	2,329
Second	60.6	3,852	7.0	2.3	17.3	4.5	48.5	31.5	7.3	3.3	2,336
Middle	55.6	4,133	8.4	2.6	13.4	5.8	41.1	25.3	14.1	3.9	2,296
Fourth	54.0	4,800	10.0	1.9	14.7	8.1	33.4	21.1	20.9	2.9	2,591
Highest	54.5	5,134	10.9	7.7	19.0	6.4	16.2	10.8	34.5	3.6	2,796
Total 15–49	58.2	21,231	8.7	4.8	16.9	5.8	36.6	23.6	17.1	3.2	12,348

[1] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 3.10C Reasons for not registering with the Social Health Authority (SHA): County level

Percentage of women age 15–49 years not registered with the Social Health Authority (SHA) by reason for not registering, according to county, Kenya Mini DHS 2025–26

			Among women not registered, reason for not registering								
County	Percentage of women not registered with SHA	Number of women	Inconvenient/ burdensome	Have not heard of SHA/not aware	Aware but don't know how to register	Distrust in system/ government	Financial barriers/ too expensive	Problems with required documentation/ inadequate coverage of needed services	Not interested/ satisfied with other coverage	Other	Number of women not registered with SHA
Mombasa	51.7	803	8.5	0.4	2.6	12.2	12.0	21.2	54.0	2.2	415
Kwale	61.2	374	1.2	1.3	19.7	7.2	46.7	40.6	6.1	11.6	229
Kilifi	58.2	608	2.5	7.5	57.2	2.0	19.5	12.0	8.0	0.8	354
Tana River	77.2	137	14.4	35.6	24.7	9.7	17.1	2.8	0.8	0.0	106
Lamu	66.5	54	0.3	10.6	19.9	3.4	36.8	24.4	4.5	2.7	36
Taita Taveta	32.0	153	6.9	3.7	18.1	17.5	27.2	21.9	7.7	1.3	49
Garissa	77.5	168	12.4	28.2	28.1	0.0	11.4	11.0	14.7	1.1	130
Wajir	72.0	130	2.0	6.3	39.4	0.9	47.7	0.2	22.4	0.0	94
Mandera	75.3	123	0.7	25.3	41.6	6.8	32.7	13.3	3.3	0.9	93
Marsabit	72.5	142	31.2	22.0	21.5	4.4	56.3	7.2	7.2	0.0	103
Isiolo	85.6	76	13.0	14.9	2.0	0.0	43.6	4.2	26.6	0.0	65
Meru	68.9	530	10.6	3.0	12.1	6.5	49.3	25.7	8.6	2.9	365
Tharaka-Nithi	50.5	147	8.1	1.1	5.5	0.9	44.8	23.4	12.6	17.2	74
Embu	38.5	239	7.4	3.7	23.4	7.2	53.1	16.5	18.3	1.8	92
Kitui	63.0	412	30.5	1.8	16.6	4.8	65.2	16.5	15.9	3.2	259
Machakos	58.5	677	15.9	0.0	10.3	2.8	48.9	22.1	24.8	2.7	396
Makueni	51.3	350	8.6	8.9	7.5	6.2	32.0	43.0	4.3	6.0	179
Nyandarua	47.7	223	10.5	3.2	5.9	12.9	37.9	18.7	14.9	16.7	106
Nyeri	34.7	223	9.3	0.9	9.2	14.4	17.6	21.1	35.2	1.2	77
Kirinyaga	34.0	258	0.6	0.0	17.9	10.8	51.3	5.5	15.8	9.2	88
Murang'a	74.8	377	4.3	4.1	18.2	6.4	48.3	14.3	20.8	1.1	282
Kiambu	60.1	1,689	7.9	2.9	11.4	6.0	12.8	7.9	52.5	3.0	1,014
Turkana	55.1	178	7.8	11.2	33.0	5.2	49.5	22.3	9.6	0.7	98
West Pokot	76.0	225	5.3	1.1	5.3	12.6	60.3	56.3	3.7	0.3	171
Samburu	44.9	98	4.1	19.3	17.3	1.3	7.2	43.6	3.1	9.4	44
Trans Nzoia	69.4	370	7.6	2.3	10.8	4.2	49.8	36.0	3.4	5.0	257
Uasin Gishu	58.8	667	7.5	0.9	8.4	7.4	41.5	27.2	19.1	2.9	392
Elgeyo-Marakwet	35.8	178	29.9	3.9	15.5	3.9	27.0	64.4	3.5	0.2	64
Nandi	48.9	361	4.6	1.6	6.7	4.1	27.7	49.8	6.6	0.5	176
Baringo	47.2	187	6.3	13.0	14.3	0.0	46.8	59.1	4.8	6.1	88
Laikipia	63.4	195	1.8	12.0	38.8	7.0	13.3	7.4	4.8	19.5	124
Nakuru	55.0	1,078	9.9	1.4	12.8	7.9	24.4	38.3	13.7	2.4	593
Narok	75.9	428	13.9	10.0	25.2	6.6	20.1	17.8	9.0	9.5	325
Kajiado	51.4	467	4.9	7.9	13.9	6.3	22.1	28.9	23.7	3.1	240
Kericho	53.1	387	5.4	0.5	7.3	1.5	40.2	32.7	15.9	5.1	205
Bomet	44.6	480	0.1	0.9	32.3	0.0	56.6	22.7	2.4	1.3	214
Kakamega	53.1	665	7.0	2.7	18.1	5.1	49.8	32.0	10.7	3.9	353
Vihiga	48.4	233	3.4	2.3	9.8	2.6	53.3	25.8	2.6	12.8	113
Bungoma	59.6	625	3.4	2.3	5.6	4.5	53.8	50.7	1.0	1.3	372
Busia	57.4	373	5.9	4.3	10.5	1.3	42.4	37.6	1.6	4.3	214
Siaya	65.1	370	3.4	1.3	11.7	2.1	55.0	27.4	7.5	0.0	241
Kisumu	48.0	551	20.3	0.0	18.6	5.6	43.1	31.6	8.7	0.7	264
Homa Bay	58.8	485	4.3	0.3	18.7	9.1	73.2	28.5	5.2	0.0	285
Migori	60.6	509	6.3	2.4	12.3	2.9	48.7	43.3	2.5	1.2	308
Kisii	64.9	439	3.7	0.8	5.5	14.6	64.4	30.7	4.4	0.0	284
Nyamira	58.2	238	1.2	0.0	2.5	8.3	53.7	40.6	1.3	7.6	138
Nairobi City	61.3	3,550	11.2	7.9	23.7	4.8	27.5	10.0	21.2	2.6	2,176
Total 15–49	58.2	21,231	8.7	4.8	16.9	5.8	36.6	23.6	17.1	3.2	12,348

Note: Multiple responses were allowed, so total percentages may sum to more than 100%.

Table 3.11 Utilisation of Social Health Authority (SHA) services

Percentage of women age 15–49 registered with the Social Health Authority (SHA) who received care from a health provider or pharmacy in the past 12 months whose most recent health care services were covered under the SHA scheme, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Percentage whose services were covered under SHA	Number of women registered with SHA who received care from a health provider in the past 12 months
Age		
15–19	43.8	409
20–24	56.3	831
25–29	53.7	1,409
30–34	52.8	1,114
35–39	54.3	911
40–44	51.2	601
45–49	49.2	503
Residence		
Urban	51.1	2,870
Rural	54.1	2,909
Education[1]		
No education	50.0	165
Primary	51.2	1,483
Secondary	53.1	2,304
More than secondary	53.4	1,827
Wealth quintile		
Lowest	52.0	599
Second	52.2	920
Middle	53.1	1,154
Fourth	57.8	1,470
Highest	48.1	1,636
Where care was sought most recently		
Public sector	57.8	4,111
Government hospital	59.1	2,150
Government health centre	60.5	824
Government dispensary	53.6	1,132
Other public sector	*	6
Private medical sector	43.8	1,945
Private hospital	55.3	1,273
Private clinic	32.4	314
Pharmacy	12.6	342
Other private sector	*	16
NGO/FBO medical sector	42.2	107
NGO hospital	*	16
FBO mission hospital	43.7	84
Other FBO medical sector	*	6
Other	*	10
Total 15–49	52.6	5,779

Note: An asterisk indicates that a figure is based on fewer than 25 unweighted cases and has been suppressed.
FBO = faith-based organisation
NGO = nongovernmental organisation
[1] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 3.11C Utilisation of Social Health Authority (SHA) services: County level

Percentage of women age 15–49 registered with the Social Health Authority (SHA) who received care from a health provider or pharmacy in the past 12 months whose most recent health care services were covered under the SHA scheme, according to county, Kenya Mini DHS 2025–26

County	Percentage whose services were covered under SHA	Number of women registered with SHA who received care from a health provider in the past 12 months
Mombasa	43.8	231
Kwale	53.4	101
Kilifi	57.6	183
Tana River	37.0	26
Lamu	74.2	11
Taita Taveta	57.6	82
Garissa	(45.4)	17
Wajir	52.8	22
Mandera	*	6
Marsabit	40.7	27
Isiolo	(38.8)	6
Meru	39.3	126
Tharaka-Nithi	60.4	65
Embu	46.5	89
Kitui	62.4	119
Machakos	47.8	204
Makueni	50.1	128
Nyandarua	63.3	84
Nyeri	65.7	107
Kirinyaga	40.5	118
Murang'a	61.5	53
Kiambu	67.3	409
Turkana	55.3	40
West Pokot	66.7	33
Samburu	67.3	38
Trans Nzoia	50.4	54
Uasin Gishu	66.6	187
Elgeyo-Marakwet	77.4	74
Nandi	76.1	79
Baringo	65.6	73
Laikipia	53.1	33
Nakuru	52.3	278
Narok	58.3	50
Kajiado	50.6	174
Kericho	58.2	136
Bomet	63.7	85
Kakamega	45.1	173
Vihiga	65.3	70
Bungoma	34.9	183
Busia	42.8	132
Siaya	(55.3)	37
Kisumu	56.8	185
Homa Bay	42.4	124
Migori	44.3	158
Kisii	49.7	84
Nyamira	48.3	46
Nairobi City	45.9	1,042
Total 15–49	52.6	5,779

Note: Figures in parentheses are based on 25–49 unweighted cases. An asterisk indicates that a figure is based on fewer than 25 unweighted cases and has been suppressed.

Table 3.12 Services received through the Social Health Authority (SHA)

Percent distribution of women age 15–49 who are registered with the Social Health Authority (SHA) and received services under the SHA scheme in the past 12 months by type of service received the last time they accessed care covered by SHA, according to background characteristics, Kenya Mini DHS 2025–26

	Percentage of women receiving a specific service the last time they accessed care:											
Background characteristic	Outpatient services/ consul-tations	Inpatient care/ hospitali-sation	Accident/ injury	Antenatal care services	Delivery and maternity services	Postnatal care services	Chronic illness manage-ment	Routine medical checkup/ examina-tion	Labora-tory/ diagnostic services	Medi-cines/ pharmacy	Total	Number of women
Age												
15–19	66.4	4.4	3.5	8.0	7.9	3.9	1.6	2.2	1.5	0.6	100.0	179
20–24	46.1	4.5	0.3	19.4	13.7	12.6	0.7	1.2	1.1	0.4	100.0	468
25–29	53.9	9.4	0.5	10.2	11.7	10.5	0.6	1.2	0.9	1.1	100.0	757
30–34	64.4	7.0	0.1	8.4	7.0	9.1	0.5	0.9	1.9	0.8	100.0	588
35–39	66.8	9.6	0.3	6.7	7.1	6.7	1.5	0.9	0.2	0.3	100.0	495
40–44	76.4	9.7	0.8	3.9	3.0	2.4	1.2	2.4	0.2	0.1	100.0	308
45–49	76.1	9.9	0.3	0.0	1.5	0.0	4.8	5.6	0.7	1.1	100.0	247
Residence												
Urban	63.2	8.1	0.4	9.5	8.3	7.3	1.0	0.9	0.7	0.6	100.0	1,467
Rural	60.2	7.9	0.7	8.7	8.5	8.4	1.4	2.3	1.2	0.7	100.0	1,574
Education[1]												
No education	71.7	4.8	0.0	7.4	9.7	3.9	2.2	0.1	0.0	0.0	100.0	82
Primary	60.3	11.9	0.6	7.7	9.3	4.8	2.0	1.4	1.6	0.5	100.0	760
Secondary	57.8	7.0	0.8	11.0	9.3	9.1	1.1	1.9	0.8	1.0	100.0	1,224
More than secondary	66.6	6.4	0.2	8.0	6.5	9.0	0.5	1.7	0.7	0.4	100.0	975
Wealth quintile												
Lowest	59.3	5.7	0.9	11.5	10.8	7.5	1.1	1.5	0.7	0.9	100.0	311
Second	54.1	11.0	0.4	9.0	9.6	9.8	1.4	1.8	2.2	0.7	100.0	480
Middle	59.5	11.4	0.4	8.3	6.8	7.8	1.9	1.8	1.2	1.0	100.0	613
Fourth	61.1	7.6	0.5	8.5	10.9	6.9	0.7	2.3	0.8	0.8	100.0	850
Highest	69.3	4.8	0.7	9.6	5.3	8.0	1.0	0.7	0.3	0.2	100.0	787
Total 15–49	61.6	8.0	0.6	9.1	8.4	7.9	1.2	1.6	1.0	0.7	100.0	3,041

[1] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 3.12C Services received through the Social Health Authority (SHA): County level

Percent distribution of women 15–49 who are registered with the Social Health Authority (SHA) and received services under the SHA scheme in the past 12 months by type of service received the last time they accessed care covered by SHA, Kenya Mini DHS 2025–26

	Percentage of women receiving a specific service the last time they accessed care:											
County	Outpatient services/ consul-tation	Inpatient care/ hospitali-sation	Accident/ injury	Antenatal care services	Delivery and maternity services	Postnatal care services	Chronic illness manage-ment	Routine medical checkup/ examina-tion	Labora-tory/ diagnostic services	Medi-cines/ pharmacy	Total	Number of women
Mombasa	78.7	6.9	0.0	9.0	3.7	1.7	0.0	0.0	0.0	0.0	100.0	101
Kwale	83.2	3.2	0.0	13.6	0.0	0.0	0.0	0.0	0.0	0.0	100.0	54
Kilifi	50.6	5.9	0.7	14.1	7.7	4.3	10.7	5.0	1.1	0.0	100.0	105
Tana River	(56.7)	(1.3)	(6.0)	(11.1)	(11.1)	(3.9)	(0.0)	(9.9)	(0.0)	(0.0)	100.0	10
Lamu	69.5	2.1	0.0	2.0	23.5	0.0	2.8	0.0	0.0	0.0	100.0	8
Taita Taveta	61.5	2.3	1.2	5.5	5.8	12.8	3.7	3.2	2.0	2.1	100.0	47
Garissa	*	*	*	*	*	*	*	*	*	*	100.0	8
Wajir	(98.3)	(0.0)	(0.0)	(0.0)	(1.7)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	100.0	12
Mandera	*	*	*	*	*	*	*	*	*	*	100.0	4
Marsabit	(64.1)	(3.4)	(0.0)	(2.0)	(4.8)	(25.6)	(0.0)	(0.0)	(0.0)	(0.0)	100.0	11
Isiolo	*	*	*	*	*	*	*	*	*	*	100.0	2
Meru	(68.5)	(6.8)	(0.0)	(6.6)	(6.6)	(6.8)	(0.0)	(2.5)	(2.2)	(0.0)	100.0	49
Tharaka-Nithi	67.0	7.2	1.4	4.4	7.5	11.0	0.0	0.8	0.0	0.8	100.0	39
Embu	68.8	10.3	0.0	7.4	2.7	9.2	0.0	1.5	0.0	0.0	100.0	41
Kitui	62.5	1.4	0.0	15.8	1.2	18.6	0.0	0.0	0.5	0.0	100.0	74
Machakos	49.6	7.2	0.0	17.8	8.5	12.5	0.0	3.3	0.0	1.1	100.0	97
Makueni	61.3	2.7	1.5	1.8	9.6	17.7	1.4	2.8	1.2	0.0	100.0	64
Nyandarua	56.0	1.7	0.0	17.5	4.6	13.6	5.2	1.5	0.0	0.0	100.0	53
Nyeri	70.0	3.1	0.0	3.2	10.5	8.8	0.0	1.4	3.1	0.0	100.0	70
Kirinyaga	64.6	9.8	0.0	15.6	2.3	7.6	0.0	0.0	0.0	0.0	100.0	48
Murang'a	(66.4)	(9.7)	(2.9)	(9.4)	(8.4)	(3.1)	(0.0)	(0.0)	(0.0)	(0.0)	100.0	33
Kiambu	67.6	1.4	1.7	7.1	11.9	8.9	0.4	0.4	0.5	0.0	100.0	275
Turkana	(76.0)	(8.2)	(2.1)	(5.5)	(0.6)	(4.3)	(0.0)	(0.0)	(3.2)	(0.0)	100.0	22
West Pokot	(49.3)	(12.2)	(0.0)	(8.3)	(7.8)	(2.1)	(2.1)	(6.4)	(4.9)	(6.9)	100.0	22
Samburu	79.9	2.1	0.0	3.3	7.6	4.5	0.0	0.7	2.0	0.0	100.0	25
Trans Nzoia	(76.8)	(7.9)	(0.0)	(5.8)	(3.2)	(6.2)	(0.0)	(0.0)	(0.0)	(0.0)	100.0	27
Uasin Gishu	47.8	6.4	1.3	17.6	11.5	9.5	0.8	2.9	1.7	0.8	100.0	124
Elgeyo-Marakwet	66.5	3.4	2.3	12.3	8.8	0.0	0.0	4.4	0.0	2.3	100.0	57
Nandi	66.6	14.7	0.0	2.2	12.4	1.4	1.6	0.0	1.2	0.0	100.0	60
Baringo	74.8	5.1	3.1	5.1	5.1	4.3	0.0	2.5	0.0	0.0	100.0	48
Laikipia	(26.8)	(9.7)	(0.0)	(14.6)	(48.9)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	100.0	18
Nakuru	65.1	7.8	0.0	13.7	2.5	9.6	0.0	1.3	0.0	0.0	100.0	146
Narok	(65.7)	(4.8)	(3.2)	(11.3)	(9.8)	(5.2)	(0.0)	(0.0)	(0.0)	(0.0)	100.0	29
Kajiado	54.6	7.8	0.0	4.5	16.9	14.0	0.0	0.0	0.0	2.2	100.0	88
Kericho	68.6	7.9	0.0	12.4	6.3	3.5	0.0	1.3	0.0	0.0	100.0	79
Bomet	49.0	9.7	1.7	11.7	19.5	8.4	0.0	0.0	0.0	0.0	100.0	54
Kakamega	36.0	15.8	0.0	7.4	18.1	12.8	0.0	8.2	0.0	1.7	100.0	78
Vihiga	58.0	9.7	1.7	8.9	6.9	13.5	1.2	0.0	0.0	0.0	100.0	45
Bungoma	40.2	14.5	0.0	18.9	8.1	17.0	0.0	0.0	1.2	0.0	100.0	64
Busia	56.3	8.9	0.0	8.6	9.0	13.6	3.5	0.0	0.0	0.0	100.0	56
Siaya	(56.8)	(14.0)	(0.0)	(10.9)	(11.8)	(4.9)	(0.0)	(1.7)	(0.0)	(0.0)	100.0	21
Kisumu	67.0	6.2	0.0	1.8	11.2	2.1	1.9	4.9	3.0	1.9	100.0	105
Homa Bay	48.4	18.9	0.0	4.5	1.9	0.0	7.7	12.2	4.0	2.4	100.0	53
Migori	57.0	16.9	0.6	5.5	5.6	6.3	0.0	1.3	5.2	1.5	100.0	70
Kisii	(37.1)	(50.6)	(0.0)	(0.0)	(12.3)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	100.0	42
Nyamira	(67.2)	(10.0)	(0.0)	(7.9)	(9.9)	(0.0)	(2.3)	(2.7)	(0.0)	(0.0)	100.0	22
Nairobi City	64.3	9.2	0.0	8.1	6.8	7.6	1.3	0.0	1.3	1.4	100.0	478
Total 15–49	61.6	8.0	0.6	9.1	8.4	7.9	1.2	1.6	1.0	0.7	100.0	3,041

Note: Figures in parentheses are based on 25–49 unweighted cases. An asterisk indicates that a figure is based on fewer than 25 unweighted cases and has been suppressed.

Table 3.13 Challenges accessing Social Health Authority (SHA) services

Percentage of women age 15–49 registered with the Social Health Authority (SHA) who received care from a health provider in the past 12 months and reported difficulties accessing SHA-covered health services, by type of difficulty encountered the most recent time they accessed SHA-covered care, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	No challenges faced	Long waiting times	Some services not available at preferred health facility	Lack of medicines or supplies	Poor provider attitude	Confusion about benefits	Inadequacy of benefits covered	Not accepted at preferred health facility	Premiums not paid	SHA system down	Number of women
Age											
15–19	85.7	5.9	0.9	1.8	0.6	1.5	2.1	0.5	2.1	3.0	179
20–24	81.9	2.9	2.6	6.8	1.4	1.1	3.5	0.9	3.0	5.0	468
25–29	69.3	8.9	4.7	9.2	0.7	1.4	5.8	2.9	3.3	8.6	757
30–34	71.7	5.9	6.1	7.9	0.5	2.6	5.4	3.5	3.1	6.8	588
35–39	68.8	10.7	7.9	10.4	0.4	1.2	6.9	4.4	3.3	7.7	495
40–44	71.1	9.0	5.8	9.2	0.2	2.1	10.3	1.6	6.6	7.2	308
45–49	71.1	4.7	4.4	14.7	1.0	2.3	6.3	0.9	5.5	4.9	247
Residence											
Urban	69.5	8.3	5.0	9.6	0.7	1.1	6.5	4.1	2.1	7.6	1,467
Rural	76.2	6.1	5.0	8.0	0.6	2.2	5.2	1.0	5.0	6.1	1,574
Education[1]											
No education	83.1	5.0	3.3	7.4	0.9	0.8	1.8	0.9	4.3	4.4	82
Primary	76.2	6.6	5.0	9.4	0.5	1.5	5.0	1.4	5.7	3.6	760
Secondary	73.3	7.5	5.9	8.6	0.5	1.9	6.6	1.5	3.7	6.7	1,224
More than secondary	69.1	7.4	4.1	8.7	1.0	1.6	5.9	4.8	1.9	9.5	975
Wealth quintile											
Lowest	75.5	6.3	4.2	11.1	0.7	3.3	4.5	0.7	8.8	3.9	311
Second	80.2	6.3	4.1	5.4	0.5	2.5	4.2	0.2	5.1	5.0	480
Middle	73.7	6.6	4.2	10.7	0.2	0.9	3.6	1.8	4.4	7.7	613
Fourth	73.1	8.7	5.8	8.6	1.1	2.0	7.6	1.6	2.4	7.6	850
Highest	66.7	6.9	5.7	8.6	0.6	0.8	7.2	6.2	1.4	7.5	787
Total 15–49	72.9	7.2	5.0	8.8	0.7	1.7	5.8	2.5	3.6	6.8	3,041

[1] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level

Table 3.13C Challenges accessing Social Health Authority (SHA) Services: County level

Percentage of women age 15–49 registered with the Social Health Authority (SHA) who received care from a health provider in the past 12 months and reported difficulties accessing SHA-covered health services, by type of difficulty encountered the most recent time they accessed SHA-covered care, according to county, Kenya Mini DHS 2025–26

County	No challenges faced	Long waiting times	Some services not available at preferred health facility	Lack of medicines or supplies	Poor provider attitude	Confusion about benefits	Inadequacy of benefits covered	Not accepted at preferred health facility	Premiums not paid	SHA system down	Number of women
Mombasa	86.1	4.0	4.0	1.5	0.0	2.7	0.0	0.0	4.5	1.7	101
Kwale	80.1	0.0	0.0	2.5	0.0	1.2	6.8	0.0	9.0	11.5	54
Kilifi	85.3	5.0	5.9	2.0	0.0	0.0	4.8	1.1	3.2	1.5	105
Tana River	(68.5)	(6.0)	(6.5)	(10.0)	(1.9)	(9.2)	(0.0)	(0.0)	(0.0)	(1.6)	10
Lamu	93.3	1.6	0.0	3.4	0.0	0.0	0.0	0.0	1.6	1.6	8
Taita Taveta	73.6	7.0	1.8	12.0	2.7	3.8	6.0	1.3	4.8	4.5	47
Garissa	*	*	*	*	*	*	*	*	*	*	8
Wajir	(55.0)	(4.5)	(0.0)	(0.0)	(0.0)	(0.0)	(1.5)	(0.0)	(0.0)	(42.0)	12
Mandera	*	*	*	*	*	*	*	*	*	*	4
Marsabit	(78.2)	(9.4)	(0.0)	(6.6)	(0.0)	(0.0)	(0.0)	(0.0)	(7.8)	(7.4)	11
Isiolo	*	*	*	*	*	*	*	*	*	*	2
Meru	(64.4)	(1.3)	(7.5)	(17.7)	(0.0)	(5.5)	(15.8)	(0.0)	(3.1)	(3.5)	49
Tharaka-Nithi	50.3	12.7	8.2	23.7	1.2	2.5	9.6	2.0	7.7	17.7	39
Embu	80.2	7.2	11.1	5.3	2.3	3.6	8.1	5.8	1.3	0.0	41
Kitui	51.3	14.8	5.3	30.2	0.0	4.9	0.0	0.5	35.1	2.9	74
Machakos	71.0	9.9	22.0	15.1	0.0	1.8	2.4	0.0	5.7	10.2	97
Makueni	83.0	2.0	3.7	1.4	1.4	2.9	7.8	1.7	4.1	3.7	64
Nyandarua	48.0	21.9	11.9	21.7	5.8	0.0	13.0	1.7	8.0	12.9	53
Nyeri	79.4	2.8	2.5	0.0	0.8	3.4	8.7	3.4	6.8	5.4	70
Kirinyaga	77.3	1.4	5.3	7.8	0.0	0.0	0.0	3.6	2.7	6.4	48
Murang'a	(70.0)	(8.4)	(1.2)	(14.5)	(0.0)	(4.2)	(11.4)	(1.2)	(0.0)	(9.6)	33
Kiambu	67.0	7.2	0.3	13.8	1.1	0.0	5.6	5.2	1.4	2.1	275
Turkana	(83.4)	(2.1)	(11.4)	(4.3)	(0.0)	(3.1)	(0.0)	(0.0)	(0.0)	(2.1)	22
West Pokot	(83.6)	(9.1)	(7.8)	(3.2)	(0.0)	(0.0)	(0.0)	(0.0)	(3.2)	(7.4)	22
Samburu	82.3	4.3	3.2	5.1	0.7	0.7	1.9	3.3	3.9	5.2	25
Trans Nzoia	(67.8)	(13.4)	(4.8)	(4.9)	(6.7)	(8.0)	(9.7)	(4.8)	(2.9)	(13.2)	27
Uasin Gishu	70.7	17.0	6.1	14.0	1.5	0.0	6.0	0.8	1.2	15.4	124
Elgeyo-Marakwet	95.1	1.5	1.0	0.0	0.0	0.0	3.2	0.0	2.2	1.4	57
Nandi	73.4	3.0	6.4	6.5	0.0	0.0	2.2	1.4	3.9	10.6	60
Baringo	78.3	8.6	6.4	2.9	1.4	2.7	1.8	0.0	4.1	9.1	48
Laikipia	(82.3)	(6.1)	(3.4)	(0.0)	(0.0)	(0.0)	(3.1)	(0.0)	(2.0)	(12.4)	18
Nakuru	62.4	8.4	12.9	13.2	0.0	3.3	15.8	7.8	6.2	6.3	146
Narok	(63.0)	(8.8)	(13.1)	(8.7)	(0.0)	(0.0)	(2.9)	(4.0)	(5.9)	(0.0)	29
Kajiado	84.1	9.0	4.8	2.4	1.9	1.9	4.9	1.9	1.1	3.0	88
Kericho	85.3	1.8	0.0	2.3	0.0	0.0	3.4	0.0	4.0	9.2	79
Bomet	83.1	8.9	3.1	5.0	0.0	0.0	0.2	0.0	3.3	1.5	54
Kakamega	72.1	10.3	4.7	7.4	1.7	3.2	3.8	0.6	2.9	16.6	78
Vihiga	88.3	1.5	2.3	2.5	0.0	0.0	0.6	0.0	2.2	4.4	45
Bungoma	82.3	3.3	0.0	4.4	0.0	2.7	3.0	0.0	2.0	9.1	64
Busia	75.6	6.3	2.2	10.6	0.0	3.9	4.3	1.0	1.0	6.1	56
Siaya	(65.9)	(15.9)	(4.2)	(0.0)	(4.2)	(4.2)	(8.6)	(0.0)	(1.7)	(9.5)	21
Kisumu	71.6	13.4	3.3	9.4	0.5	5.5	8.0	0.0	5.8	10.9	105
Homa Bay	80.2	6.0	4.4	4.6	0.0	3.8	8.6	0.0	0.0	8.1	53
Migori	68.7	12.4	7.1	3.3	0.0	0.0	6.5	1.3	2.6	16.0	70
Kisii	(95.7)	(0.0)	(0.0)	(3.0)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(1.2)	42
Nyamira	(64.5)	(5.8)	(4.9)	(8.8)	(0.0)	(14.0)	(13.5)	(2.6)	(3.4)	(6.8)	22
Nairobi City	65.9	5.2	4.0	10.0	0.0	0.0	7.1	6.0	0.0	5.9	478
Total 15–49	72.9	7.2	5.0	8.8	0.7	1.7	5.8	2.5	3.6	6.8	3,041

Note: Figures in parentheses are based on 25–49 unweighted cases. An asterisk indicates that a figure is based on fewer than 25 unweighted cases and has been suppressed.

Table 3.14 Benefits received through Social Health Authority (SHA) coverage

Percentage of women age 15–49 registered with the Social Health Authority (SHA) who received care from a health provider in the past 12 months and received benefits through the SHA scheme for their most recent health care services, by type of benefit received, according to background characteristics, Kenya Mini DHS 2025–26

	Percentage of women receiving specific benefits through SHA:							
Background characteristic	Reduced out-of-pocket spending on health care	Increased access to quality health services	Timely medical attention	Improved personal health	Reduced stress over health expenses	Inclusion of dependents	Other	Number of women
Age								
15–19	89.5	7.4	12.0	5.6	17.1	4.4	1.0	179
20–24	92.7	11.4	7.1	5.0	21.1	7.1	0.4	468
25–29	91.0	12.4	11.6	6.7	20.5	10.2	0.5	757
30–34	88.6	14.7	12.4	7.0	25.1	9.4	3.2	588
35–39	91.2	11.7	9.2	8.0	22.0	9.4	1.5	495
40–44	90.9	17.2	16.0	10.8	24.7	10.4	0.7	308
45–49	91.1	13.8	17.5	9.3	24.7	8.0	0.0	247
Residence								
Urban	92.1	10.1	10.2	5.6	19.2	8.5	2.0	1,467
Rural	89.5	15.5	12.9	8.9	25.2	9.3	0.4	1,574
Education[1]								
No education	89.6	11.4	9.7	2.5	23.5	2.5	0.3	82
Primary	92.1	9.8	12.9	5.5	23.7	8.5	0.2	760
Secondary	89.1	13.2	11.7	8.2	24.3	10.1	1.0	1,224
More than secondary	91.8	15.0	10.6	8.0	18.6	8.3	2.2	975
Wealth quintile								
Lowest	90.5	13.9	10.6	8.3	29.0	9.8	0.2	311
Second	90.3	13.8	12.9	8.9	28.6	7.5	0.5	480
Middle	89.4	15.4	13.6	7.2	23.5	10.4	0.3	613
Fourth	92.6	9.2	11.8	6.2	18.5	10.0	0.7	850
Highest	90.1	14.0	9.5	7.2	19.0	7.2	3.2	787
Total 15–49	90.7	12.9	11.6	7.3	22.3	8.9	1.2	3,041

[1] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 3.14C Benefits received through Social Health Authority (SHA) coverage: County level

Percentage of women age 15–49 registered with the Social Health Authority (SHA) who received care from a health provider in the past 12 months and received benefits through the SHA scheme for their most recent health care services, by type of benefit received, according to county, Kenya Mini DHS 2025–26

	Percentage of women receiving specific benefits through SHA:							
County	Reduced out-of-pocket spending on health care	Increased access to quality health services	Timely medical attention	Improved personal health	Reduced stress over health expenses	Inclusion of dependents	Other	Number of women
Mombasa	79.1	11.2	13.4	2.7	1.8	6.7	0.0	101
Kwale	96.1	1.0	3.3	0.0	1.6	1.2	0.0	54
Kilifi	97.0	13.5	6.5	14.6	42.9	4.3	0.0	105
Tana River	(94.1)	(27.7)	(23.0)	(13.1)	(32.0)	(7.0)	(0.0)	10
Lamu	100.0	0.0	0.0	0.0	1.6	1.6	0.0	8
Taita Taveta	93.2	5.5	4.2	1.8	15.3	6.3	0.7	47
Garissa	*	*	*	*	*	*	*	8
Wajir	(89.2)	(0.0)	(0.0)	(0.0)	(12.5)	(0.0)	(0.0)	12
Mandera	*	*	*	*	*	*	*	4
Marsabit	(96.6)	(14.3)	(6.2)	(0.0)	(17.3)	(22.3)	(0.0)	11
Isiolo	*	*	*	*	*	*	*	2
Meru	(85.9)	(5.8)	(16.2)	(3.6)	(18.1)	(2.5)	(0.0)	49
Tharaka-Nithi	99.2	8.9	5.2	6.2	16.0	4.7	0.0	39
Embu	67.4	21.1	12.7	2.3	40.3	0.0	3.5	41
Kitui	68.0	16.0	5.3	4.2	9.3	16.4	0.0	74
Machakos	93.0	25.4	3.7	2.6	20.7	10.0	0.0	97
Makueni	98.7	8.0	3.0	3.3	43.3	0.0	0.0	64
Nyandarua	84.2	10.0	6.9	2.2	13.9	3.1	0.0	53
Nyeri	97.7	1.2	0.0	1.3	0.0	0.0	1.1	70
Kirinyaga	68.8	11.7	12.6	3.1	17.0	34.7	2.8	48
Murang'a	(92.0)	(17.6)	(4.1)	(2.9)	(14.2)	(13.4)	(0.0)	33
Kiambu	98.4	0.6	1.0	3.7	11.2	0.0	1.1	275
Turkana	(72.8)	(20.4)	(36.4)	(3.8)	(78.0)	(1.8)	(0.0)	22
West Pokot	(96.8)	(10.5)	(9.8)	(1.4)	(4.4)	(3.4)	(0.0)	22
Samburu	87.9	20.5	3.9	8.5	25.2	6.0	2.7	25
Trans Nzoia	(75.8)	(18.0)	(4.5)	(2.9)	(25.0)	(0.0)	(1.7)	27
Uasin Gishu	89.5	27.4	23.4	18.4	23.9	23.5	0.0	124
Elgeyo-Marakwet	97.0	67.6	64.6	60.6	74.2	72.2	0.0	57
Nandi	58.7	40.8	26.7	22.4	20.1	2.2	0.0	60
Baringo	96.2	27.7	49.1	25.1	34.0	9.9	1.5	48
Laikipia	(91.9)	(18.4)	(0.0)	(20.1)	(61.4)	(0.0)	(3.1)	18
Nakuru	89.2	1.5	4.9	1.3	13.6	4.3	4.1	146
Narok	(96.8)	(3.2)	(6.8)	(0.0)	(18.1)	(2.9)	(0.0)	29
Kajiado	97.5	1.6	0.0	9.1	51.5	0.0	0.0	88
Kericho	100.0	7.3	3.3	2.4	22.5	3.1	0.0	79
Bomet	96.9	9.2	8.6	1.9	3.4	1.6	0.0	54
Kakamega	80.9	19.2	4.6	11.1	26.3	3.2	3.4	78
Vihiga	95.8	17.0	30.7	4.9	31.8	0.0	1.3	45
Bungoma	86.8	9.8	10.4	14.3	36.0	15.5	0.0	64
Busia	96.7	15.0	13.0	14.4	8.0	2.8	0.0	56
Siaya	(82.9)	(18.8)	(4.4)	(15.8)	(9.9)	(3.0)	(0.0)	21
Kisumu	91.5	18.3	32.8	11.0	42.3	8.2	0.0	105
Homa Bay	88.5	36.4	24.7	18.4	55.9	8.6	0.0	53
Migori	98.8	5.1	5.3	0.0	6.0	17.3	0.0	70
Kisii	(97.0)	(0.0)	(2.1)	(0.0)	(11.8)	(0.0)	(0.0)	42
Nyamira	(100.0)	(0.0)	(2.3)	(0.0)	(2.2)	(0.0)	(0.0)	22
Nairobi City	92.2	11.2	13.9	3.6	20.1	15.7	3.6	478
Total 15–49	90.7	12.9	11.6	7.3	22.3	8.9	1.2	3,041

Note: Figures in parentheses are based on 25–49 unweighted cases. An asterisk indicates that a figure is based on fewer than 25 unweighted cases and has been suppressed.

Table 3.15 Satisfaction with services covered under the Social Health Authority (SHA)

Among women age 15–49 registered with the Social Health Authority (SHA) who received care in the past 12 months whose most recent services were covered under the SHA scheme, percentage reporting satisfaction with key aspects of SHA services, by background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Percentage of women satisfied with:							Percentage of women who would recommend SHA services	Number of women
	Availability of SHA services	Quality of services	Respect by health care workers	Timeliness of service	Coverage of essential medicines and diagnostics	Communi-cation and public information provided	Overall satisfaction		
Age									
15–19	93.7	96.3	97.4	92.3	90.9	92.7	95.6	92.9	179
20–24	95.1	97.2	95.9	93.1	89.8	92.3	97.7	95.6	468
25–29	94.0	96.1	96.0	90.3	82.7	93.1	91.9	92.5	757
30–34	87.9	90.1	93.0	87.4	80.8	87.5	87.6	86.2	588
35–39	89.1	91.1	95.7	87.7	82.3	89.2	89.3	86.8	495
40–44	91.3	94.2	95.0	89.8	80.8	91.5	91.3	92.3	308
45–49	94.8	95.6	96.6	93.7	81.3	94.6	92.5	92.6	247
Residence									
Urban	90.9	92.9	94.3	86.9	83.0	90.1	90.9	90.6	1,467
Rural	93.0	95.2	96.4	93.0	84.1	92.2	92.5	91.1	1,574
Education[1]									
No education	97.1	97.4	97.5	95.9	91.3	97.3	97.7	95.2	82
Primary	93.5	92.7	96.0	92.1	84.5	93.1	91.2	90.2	760
Secondary	92.2	95.2	96.1	91.1	84.0	91.8	93.0	93.5	1,224
More than secondary	90.0	93.5	93.9	86.8	81.5	88.5	89.9	87.6	975
Wealth quintile									
Lowest	93.3	93.4	96.1	94.0	87.0	95.0	93.0	91.1	311
Second	96.8	96.4	97.3	93.1	86.6	94.6	96.7	92.7	480
Middle	92.5	95.4	96.3	91.5	81.0	91.3	91.3	91.8	613
Fourth	90.6	93.5	94.6	88.7	82.0	90.6	91.8	92.6	850
Highest	89.5	92.6	94.1	87.0	84.0	88.2	88.5	86.9	787
Total 15–49	92.0	94.1	95.4	90.1	83.5	91.2	91.7	90.8	3,041

[1] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 3.15C Satisfaction with services covered under the Social Health Authority (SHA): County level

Among women age 15–49 registered with the Social Health Authority (SHA) who received care in the past 12 months whose most recent services were covered by the SHA scheme, percentage reporting satisfaction with key aspects of SHA services, by county, Kenya Mini DHS 2025–26

	Percentage of women satisfied with:							Percentage of women who would recommend SHA services	Number of women
County	Availability of SHA services	Quality of services	Respect by health care workers	Timeliness of service	Coverage of essential medicines and diagnostics	Communi-cation and public information provided	Overall satisfaction		
Mombasa	93.7	95.4	95.9	90.1	92.3	95.9	95.5	98.0	101
Kwale	89.4	93.9	94.8	94.8	85.9	96.6	91.0	90.1	54
Kilifi	99.0	99.0	96.3	94.9	90.8	92.5	99.0	95.8	105
Tana River	(100.0)	(100.0)	(98.1)	(98.1)	(91.1)	(96.7)	(100.0)	(95.4)	10
Lamu	96.6	95.2	100.0	100.0	86.2	96.6	96.6	90.2	8
Taita Taveta	89.3	92.6	97.0	90.8	75.5	90.8	93.4	85.4	47
Garissa	*	*	*	*	*	*	*	*	8
Wajir	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	12
Mandera	*	*	*	*	*	*	*	*	4
Marsabit	(100.0)	(100.0)	(100.0)	(100.0)	(95.8)	(93.8)	(100.0)	(100.0)	11
Isiolo	*	*	*	*	*	*	*	*	2
Meru	(87.8)	(94.0)	(97.7)	(88.5)	(71.3)	(91.5)	(89.4)	(91.9)	49
Tharaka-Nithi	84.7	98.8	95.3	82.1	71.7	98.3	89.3	97.6	39
Embu	81.1	85.1	84.4	79.2	74.2	70.1	80.9	77.8	41
Kitui	96.1	94.9	96.1	93.5	87.7	92.9	92.4	82.8	74
Machakos	89.6	92.1	100.0	88.3	81.0	79.9	90.1	87.4	97
Makueni	98.3	98.3	96.6	94.9	89.7	94.4	95.7	93.5	64
Nyandarua	69.5	72.6	84.3	72.8	54.0	68.5	57.0	50.7	53
Nyeri	83.3	93.7	95.0	85.8	68.0	76.2	77.1	81.0	70
Kirinyaga	75.0	88.8	96.1	91.1	54.7	83.4	83.8	78.6	48
Murang'a	(92.4)	(90.9)	(94.0)	(83.8)	(66.9)	(89.7)	(92.1)	(88.0)	33
Kiambu	95.9	98.9	98.9	92.7	94.8	95.2	95.2	97.8	275
Turkana	(97.9)	(97.9)	(97.9)	(97.9)	(90.3)	(97.9)	(97.9)	(96.8)	22
West Pokot	(96.1)	(96.8)	(100.0)	(100.0)	(93.6)	(96.8)	(100.0)	(96.1)	22
Samburu	96.9	96.9	95.2	82.6	87.7	99.5	98.1	98.4	25
Trans Nzoia	(95.2)	(98.3)	(90.2)	(93.4)	(96.5)	(93.4)	(95.2)	(88.7)	27
Uasin Gishu	95.4	96.1	99.0	92.2	83.7	97.6	98.1	96.1	124
Elgeyo-Marakwet	98.8	98.8	100.0	98.8	98.1	100.0	98.8	98.8	57
Nandi	95.7	97.1	97.1	90.8	93.9	94.4	98.5	100.0	60
Baringo	97.9	98.5	98.5	97.5	85.7	97.1	98.5	94.8	48
Laikipia	(81.8)	(93.2)	(94.7)	(84.3)	(87.6)	(85.3)	(84.5)	(74.6)	18
Nakuru	82.6	87.5	85.3	78.9	79.5	86.2	86.4	92.0	146
Narok	(93.1)	(93.1)	(90.6)	(89.6)	(85.5)	(93.1)	(93.1)	(97.1)	29
Kajiado	95.9	92.9	98.1	83.6	82.2	92.6	96.0	95.0	88
Kericho	97.4	98.1	100.0	95.2	87.2	93.3	95.2	95.5	79
Bomet	100.0	100.0	98.6	97.2	80.8	96.1	98.4	98.4	54
Kakamega	88.8	93.4	95.1	88.3	83.1	94.6	85.2	81.6	78
Vihiga	96.9	98.7	98.7	98.1	94.5	98.7	98.7	96.7	45
Bungoma	97.8	94.2	97.8	94.4	81.8	92.4	96.1	92.7	64
Busia	92.9	95.3	97.2	89.3	81.2	87.6	88.5	81.4	56
Siaya	(94.8)	(100.0)	(100.0)	(100.0)	(94.8)	(95.8)	(100.0)	(100.0)	21
Kisumu	91.6	96.4	93.5	94.6	87.8	93.6	91.0	91.0	105
Homa Bay	95.4	95.4	98.1	91.7	78.1	96.5	95.4	95.4	53
Migori	94.6	96.8	97.0	92.3	94.7	94.2	96.2	94.6	70
Kisii	(97.5)	(94.7)	(100.0)	(88.0)	(71.8)	(82.6)	(94.7)	(89.5)	42
Nyamira	(89.7)	(88.1)	(97.7)	(90.9)	(86.2)	(86.2)	(88.2)	(79.0)	22
Nairobi City	89.1	90.1	91.7	87.4	78.7	89.0	87.7	87.5	478
Total 15–49	92.0	94.1	95.4	90.1	83.5	91.2	91.7	90.8	3,041

Note: Figures in parentheses are based on 25–49 unweighted cases. An asterisk indicates that a figure is based on fewer than 25 unweighted cases and has been suppressed.

MARRIAGE AND SEXUAL ACTIVITY 4

Key Findings

- ***Current marital status:*** 55% of women age 15–49 are currently in union.
- ***Marriage registration:*** 18% of currently married women age 15–49 are in a union registered with a civil authority.
- ***Age at first sexual intercourse:*** 15% of women age 25–49 had their first sexual intercourse before age 15.
- ***Recent sexual activity:*** 47% of women age 15–49 had sexual intercourse within the 4 weeks before the survey.
- ***Multiple sexual partners:*** 22% of women age 15–24 had intercourse with a person who was neither their husband nor someone who lived with them in the past 12 months; among those women, 36% used a condom during their last sexual intercourse.

Marriage and sexual activity help determine the extent to which women are exposed to the risk of pregnancy. Thus, they are important determinants of fertility levels (Shallo 2020; Ayele and Malesse 2017). However, the timing and circumstances of marriage and sexual activity also have profound consequences for the lives of both women and men.

4.1 MARITAL STATUS

Currently in union


Women and men who report being married or living together with a partner as though married at the time of the survey. In this report, the terms 'currently in union' and 'currently married' are used interchangeably except where noted.

Sample: Women and men age 15–49

Overall, 55% of women age 15–49 are currently in union (48% are married and 7% are living together), 33% have never been married, 10% are divorced or separated, and 3% are widowed (**Table 4.1** and **Figure 4.1**).

Figure 4.1 **Marital status**

Percent distribution of women age 15–49 by marital status




Note: Sum adds to more than 100% due to rounding.

Trends: The percentage of women in union decreased from 60% in the 2003 Kenya Demographic and Health Survey (KDHS) to 55% in the 2025–26 Kenya Mini Demographic and Health Survey (KMDHS), whereas the percentage who have never married increased from 30% to 33% over the same period (**Figure 4.2**).

The share of women who are divorced or separated increased from 6% in the 2003 KDHS to 10% in the 2025–26 KMDHS, while the percentage widowed remained relatively stable.

Figure 4.2 **Trends in marital status**



4.2 MARRIAGE REGISTRATION

Registered marriage

A marriage that is registered with civil authorities regardless of whether a woman has a marriage certificate.

Sample: Currently married women age 15–49 (does not include women who are living with a man as if married but do not report themselves to be married)

Eighteen percent of women age 15–49 who are in union have registered their union with a civil authority. Sixteen percent have documentation recognising their marriage or union, and 14% have a marriage certificate (**Table 4.2**).

Patterns by background characteristics

- The percentage of in-union women whose marriages are registered and who have a marriage certificate is higher in urban areas (18%) than in rural areas (11%) (**Table 4.2**).
- The percentage of in-union women whose marriages are registered and who have a marriage certificate increases with wealth, from 5% among women in the lowest wealth quintile to 27% among women in the highest wealth quintile.
- Across counties, the percentage of in-union women whose marriages are registered and who have a marriage certificate varies considerably. It is highest in Garissa (52%), Isiolo (44%), and Lamu (31%) and lowest in Mandera (2%), Wajir (3%), and Vihiga (3%) (**Table 4.2C**).

4.3 AGE AT FIRST SEXUAL INTERCOURSE

Median age at first sexual intercourse

Age by which half of respondents have had sexual intercourse.

Sample: Women age 20–49 and women age 25–49


Among women age 25–49, 15% had their first sexual intercourse before age 15, and 51% had it by age 18. Additionally, 1% of women in the same age group have never had sexual intercourse (**Table 4.3**).

The median age at first sexual intercourse for women age 25–49 is 18 years (**Table 4.4**).

Trends: The percentage of women age 25–49 who had their first sexual intercourse by age 15 decreased from 18% in the 2003 KDHS to 15% in the 2025–26 KMDHS. Similarly, the percentage who had their first sexual intercourse by age 18 decreased from 54% to 51% during the same period. The percentage of women age 25–49 who have never had sexual intercourse has remained stable at 1% (**Figure 4.3**).

***Figure 4.3* Trends in first sexual intercourse**

Percentage of women age 25–49 by timing of first sexual intercourse



Patterns by background characteristics

- The median age at first sexual intercourse among women age 25–49 is higher in urban areas (18.5 years) than in rural areas (17.3 years) (**Table 4.4**).
- The median age at first sexual intercourse among women age 25–49 increases with household wealth, from 16.7 years among women in the lowest wealth quintile to 19.1 years among women in the highest wealth quintile.
- Counties with the highest median age at first sexual intercourse among women age 25–49 are Wajir (19.8 years), Nyeri (19.1 years), and Embu (19.1 years). In contrast, counties with the lowest median age at first sexual intercourse are Samburu (15.4 years), Migori (15.9 years), and Kisii (16.0 years) (**Table 4.4C**).

4.4 Recent Sexual Activity

Forty-seven percent of women age 15–49 had sexual intercourse within the 4 weeks preceding the 2025–26 KMDHS, while 17% had never had sexual intercourse by the time of the survey (**Table 4.5**).

Trends: The percentage of women age 15–49 who had sexual intercourse within the 4 weeks preceding the survey has remained relatively stable, declining slightly from 48% in 2003 to 47% in 2025–26, despite peaking at 51% in 2014.

Patterns by background characteristics

- The percentage of women age 15–49 who had never had sexual intercourse by the time of the survey decreases as age increases, from 75% among those age 15–19 to 16% among those age 20–24 to less than 1% among those age 45–49 (**Table 4.5**).
- West Pokot (61%), Machakos (60%), Tana River (58%), Narok (57%), and Migori (56%) Counties have the highest percentages of women age 15–49 who had sexual intercourse within the 4 weeks before the survey. Conversely, counties with the lowest percentages are Vihiga (33%), Wajir (35%), Garissa (38%), Marsabit (39%), and Nyeri (40%) (**Table 4.5C**).

- Counties with the highest percentages of women age 15–49 who had never had sexual intercourse by the time of the survey are Wajir (38%), Garissa (37%), and Mandera (31%). Counties with the lowest percentages are Kirinyaga (10%), Machakos (11%), and Kajiado (11%).

4.5 Multiple Sexual Partners

Two percent of women age 15–24 had more than two sexual partners in the past 12 months, of whom 30% reported using a condom during their last sexual intercourse. Additionally, 22% of women had sexual intercourse in the past 12 months with a person who was neither their husband nor someone who lived with them, of whom 36% reported using a condom during their last sexual intercourse with such a partner (**Table 4.6**).

Patterns by background characteristics

- The percentage of women age 15–24 who had intercourse in the past 12 months with a person who was neither their husband nor someone who lived with them is highest among those with more than secondary education (41%) and lowest among those with no education (2%) (**Table 4.6**).
- Condom use during last sexual intercourse among women with more than two sexual partners in the past 12 months is higher among never-married women (39%) than among ever-married women (19%).
- The percentage of women age 15–24 who had more than two sexual partners in the past 12 months and reported using a condom during their last sexual intercourse is higher in rural areas (35%) than in urban areas (27%).
- Similarly, the percentage of women age 15–24 who reported using a condom during last sexual intercourse within the past 12 months with a person who was neither their husband nor someone who lived with them is higher in rural areas (38%) than in urban areas (33%).

List of Tables

For more information on marriage and sexual activity, see the following tables:

- **Table 4.1 Current marital status**
- **Table 4.2 Marriage registration**
- **Table 4.2C Marriage registration by county**
- **Table 4.3 Age at first sexual intercourse**
- **Table 4.4 Median age at first sexual intercourse according to background characteristics**
- **Table 4.4C Median age at first sexual intercourse by county**
- **Table 4.5 Recent sexual activity**
- **Table 4.5C Recent sexual activity by county**
- **Table 4.6 Multiple sexual partners and higher-risk sexual intercourse in the past 12 months among young women**

Table 4.1 Current marital status

Percent distribution of women age 15–49 by current marital status, according to age, Kenya Mini DHS 2025–26

Age	Marital status						Total	Percentage of respondents currently in union	Number of respondents
	Never married	Married	Living together	Divorced	Separated	Widowed			
15–19	93.1	4.3	1.6	0.0	0.9	0.0	100.0	5.9	3,916
20–24	51.5	34.0	9.4	0.3	4.6	0.2	100.0	43.4	3,653
25–29	20.4	57.6	11.2	1.4	9.1	0.3	100.0	68.9	3,732
30–34	9.0	67.2	7.8	1.9	12.5	1.6	100.0	75.1	3,108
35–39	6.9	65.6	8.5	1.9	13.0	4.0	100.0	74.1	2,989
40–44	6.0	65.1	7.0	2.9	11.7	7.3	100.0	72.1	2,048
45–49	4.9	66.1	4.5	2.9	10.5	11.2	100.0	70.6	1,784
Total 15–49	32.9	47.7	7.3	1.4	8.2	2.5	100.0	55.0	21,231

Table 4.2 Marriage registration

Percentage of in-union women age 15–49 whose current marriage or union is registered, percentage whose current marriage or union is registered and who have any documentation recognising the marriage/union, and percentage whose current marriage or union is registered and who have a marriage certificate, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Percentage whose current marriage or union is registered[1]	Percentage whose current marriage or union is registered and who have any documentation recognising the marriage/union	Percentage whose current marriage is registered and who have a marriage certificate	Number of women in union[1]
Age				
15–19	7.3	2.4	2.4	231
20–24	7.1	6.1	5.4	1,584
25–29	12.7	10.9	9.2	2,570
30–34	20.5	18.9	16.2	2,334
35–39	19.6	18.4	16.7	2,216
40–44	23.3	21.1	19.2	1,478
45–49	27.2	24.4	22.3	1,259
Marital status				
Currently married	20.1	18.2	16.1	10,124
Living together	1.5	0.8	0.6	1,548
Residence				
Urban	22.6	20.4	18.2	5,174
Rural	13.6	12.2	10.7	6,498
Education[2]				
No education	15.7	13.3	11.6	896
Primary	12.4	11.2	10.4	4,288
Secondary	13.7	12.0	10.0	4,207
More than secondary	35.3	32.6	29.3	2,280
Wealth quintile				
Lowest	7.0	6.4	5.4	1,993
Second	8.8	7.6	6.3	2,107
Middle	14.7	13.3	11.9	2,230
Fourth	19.2	17.0	15.2	2,626
Highest	33.0	30.3	27.0	2,716
Total	17.6	15.9	14.0	11,672

[1] Includes women who report that they are currently married or living with a man as if married
[2] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 4.2C Marriage registration by county

Percentage of in-union women age 15–49 whose current marriage or union is registered, percentage whose current marriage or union is registered and who have any documentation recognising the marriage/union, and percentage whose current marriage or union is registered and who have a marriage certificate, according to county, Kenya Mini DHS 2025–26

County	Percentage whose current marriage or union is registered[1]	Percentage whose current marriage or union is registered and who have any documentation recognising the marriage/union	Percentage whose current marriage is registered and who have a marriage certificate	Number of women in union[1]
Mombasa	31.3	28.5	25.3	444
Kwale	28.6	24.8	24.1	205
Kilifi	13.6	13.3	11.9	302
Tana River	19.5	19.0	15.3	88
Lamu	32.2	31.9	30.6	27
Taita Taveta	25.3	22.4	17.9	85
Garissa	53.8	52.0	52.0	90
Wajir	2.8	2.8	2.8	72
Mandera	8.9	8.0	1.8	77
Marsabit	30.2	27.1	27.1	77
Isiolo	52.8	43.8	43.8	36
Meru	19.1	18.0	17.6	308
Tharaka-Nithi	16.2	11.6	11.6	80
Embu	16.6	11.0	7.5	129
Kitui	10.7	8.4	8.0	251
Machakos	9.9	9.4	8.0	420
Makueni	19.2	18.2	15.1	192
Nyandarua	16.4	13.7	10.4	122
Nyeri	12.5	10.2	10.2	104
Kirinyaga	13.3	12.2	8.8	141
Murang'a	14.5	14.2	13.3	182
Kiambu	17.8	16.7	15.8	819
Turkana	13.4	13.0	4.3	116
West Pokot	5.5	4.8	4.3	145
Samburu	11.0	10.7	8.9	58
Trans Nzoia	13.2	9.8	9.8	197
Uasin Gishu	22.8	22.5	20.6	335
Elgeyo-Marakwet	21.7	20.5	19.0	92
Nandi	32.8	30.9	21.8	188
Baringo	11.0	8.6	4.4	109
Laikipia	15.7	15.7	15.7	105
Nakuru	11.4	10.4	9.0	614
Narok	11.1	10.0	10.0	283
Kajiado	18.3	17.9	17.7	261
Kericho	19.1	18.3	15.9	198
Bomet	15.7	13.1	13.1	242
Kakamega	10.9	9.7	8.2	406
Vihiga	5.0	3.4	3.4	118
Bungoma	9.8	8.3	8.0	349
Busia	15.9	11.7	11.7	212
Siaya	10.1	8.7	7.7	206
Kisumu	13.3	11.1	10.4	296
Homa Bay	12.7	11.5	9.1	279
Migori	11.6	10.6	8.1	314
Kisii	7.0	6.6	5.0	253
Nyamira	13.1	12.7	9.3	123
Nairobi City	25.7	22.8	19.9	1,920
Total	17.6	15.9	14.0	11,672

[1] Includes women who report that they are currently married or living with a man as if married

Table 4.3 Age at first sexual intercourse

Percentage of women age 15–49 who had first sexual intercourse by specific ages, percentage who never had sexual intercourse, and median age at first sexual intercourse, according to current age, Kenya Mini DHS 2025–26

Current age	Percentage who had first sexual intercourse by exact age					Percentage who never had intercourse	Number	Median age at first intercourse
	15	18	20	22	25			
15–19	5.1	na	na	na	na	75.2	3,916	a
20–24	7.8	36.4	65.5	na	na	16.0	3,653	18.8
25–29	12.1	46.1	70.3	86.2	94.1	3.1	3,732	18.3
30–34	14.8	50.8	73.2	88.0	93.3	0.4	3,108	17.9
35–39	16.1	53.4	73.6	85.3	92.9	0.2	2,989	17.7
40–44	16.6	53.1	73.7	86.9	91.9	0.1	2,048	17.7
45–49	14.6	55.2	76.0	89.0	93.6	0.2	1,784	17.6
20–49	13.2	47.9	71.4	na	na	4.2	17,315	18.1
25–49	14.6	51.0	72.9	86.9	93.3	1.0	13,662	17.9
15–24	6.4	na	na	na	na	46.6	7,569	a

na = Not applicable (due to censoring)
a = Omitted because less than 50% of respondents had sexual intercourse for the first time before reaching the beginning of the age group

Table 4.4 Median age at first sexual intercourse according to background characteristics

Median age at first sexual intercourse among women age 20–49 and women age 25–49, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Age	
	20–49	25–49
Residence		
Urban	18.7	18.5
Rural	17.5	17.3
Education[1]		
No education	16.9	16.8
Primary	16.7	16.7
Secondary	18.3	18.1
More than secondary	20.0	20.0
Wealth quintile		
Lowest	16.8	16.7
Second	17.1	16.9
Middle	17.9	17.6
Fourth	18.5	18.3
Highest	19.3	19.1
Total	18.1	17.9

[1] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 4.4C Median age at first sexual intercourse by county

Median age at first sexual intercourse among women age 20–49 and women age 25–49, according to county, Kenya Mini DHS 2025–26

	Age	
County	20–49	25–49
Mombasa	19.0	18.8
Kwale	18.6	18.4
Kilifi	18.2	18.0
Tana River	18.6	18.5
Lamu	17.9	17.6
Taita Taveta	18.8	18.8
Garissa	18.2	17.8
Wajir	a	19.8
Mandera	18.4	18.2
Marsabit	18.7	18.6
Isiolo	17.5	17.3
Meru	17.9	17.9
Tharaka-Nithi	18.8	18.6
Embu	19.3	19.1
Kitui	17.3	17.1
Machakos	18.1	18.2
Makueni	18.5	18.4
Nyandarua	18.5	18.1
Nyeri	19.2	19.1
Kirinyaga	18.8	18.9
Murang'a	18.7	18.4
Kiambu	19.0	18.7
Turkana	17.7	17.5
West Pokot	16.1	16.2
Samburu	15.7	15.4
Trans Nzoia	17.8	17.8
Uasin Gishu	17.9	17.5
Elgeyo-Marakwet	17.0	16.7
Nandi	16.9	16.5
Baringo	17.8	18.0
Laikipia	17.8	17.3
Nakuru	18.5	18.3
Narok	17.2	16.9
Kajiado	18.2	17.9
Kericho	17.8	17.5
Bomet	17.9	17.8
Kakamega	17.7	17.3
Vihiga	17.9	17.6
Bungoma	17.7	17.4
Busia	18.3	18.1
Siaya	16.9	16.6
Kisumu	17.3	16.9
Homa Bay	16.3	16.1
Migori	16.1	15.9
Kisii	16.2	16.0
Nyamira	16.4	16.1
Nairobi City	18.8	18.5
Total	18.1	17.9

a = Omitted because less than 50% of the respondents had intercourse for the first time before reaching the beginning of the age group

Table 4.5 Recent sexual activity

Percent distribution of women age 15–49 by timing of last sexual intercourse, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Timing of last sexual intercourse				Never had sexual intercourse	Total	Number of women
	Within the past 4 weeks	Within the past 1 year[1]	1 or more years ago	Missing			
Age							
15–19	7.7	11.7	5.4	0.0	75.2	100.0	3,916
20–24	40.5	31.4	12.2	0.0	16.0	100.0	3,653
25–29	59.7	27.6	9.4	0.2	3.1	100.0	3,732
30–34	64.7	23.7	10.8	0.4	0.4	100.0	3,108
35–39	61.1	24.1	14.4	0.2	0.2	100.0	2,989
40–44	59.0	22.3	18.6	0.0	0.1	100.0	2,048
45–49	52.0	23.9	23.9	0.2	0.2	100.0	1,784
Marital status							
Never married	8.3	23.9	15.0	0.3	52.5	100.0	6,983
Married or living together	76.8	19.8	3.3	0.1	0.0	100.0	11,672
Divorced/separated/ widowed	17.0	38.5	44.5	0.0	0.0	100.0	2,576
Residence							
Urban	46.5	23.8	13.6	0.2	15.9	100.0	9,829
Rural	47.5	23.1	10.9	0.0	18.4	100.0	11,402
Education[2]							
No education	51.3	25.1	16.6	0.0	7.0	100.0	1,161
Primary	58.8	23.3	14.8	0.2	2.9	100.0	6,053
Secondary	38.0	20.8	10.0	0.2	31.1	100.0	9,849
More than secondary	50.0	29.3	12.2	0.1	8.4	100.0	4,167
Wealth quintile							
Lowest	47.8	22.6	11.0	0.0	18.7	100.0	3,312
Second	45.1	24.8	11.2	0.1	18.9	100.0	3,852
Middle	47.4	22.2	12.9	0.1	17.5	100.0	4,133
Fourth	48.5	25.1	13.1	0.0	13.4	100.0	4,800
Highest	46.3	22.4	12.2	0.4	18.7	100.0	5,134
Total	47.0	23.4	12.1	0.1	17.3	100.0	21,231

[1] Excludes women who had sexual intercourse within the past 4 weeks

[2] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 4.5C Recent sexual activity by county

Percent distribution of women age 15–49 by timing of last sexual intercourse, according to county, Kenya Mini DHS 2025–26

County	Timing of last sexual intercourse				Never had sexual intercourse	Total	Number of women
	Within the past 4 weeks	Within the past 1 year[1]	1 or more years ago	Missing			
Mombasa	47.2	23.3	13.8	0.0	15.7	100.0	803
Kwale	42.3	23.6	10.2	0.0	23.9	100.0	374
Kilifi	44.3	22.5	8.8	0.0	24.3	100.0	608
Tana River	58.2	11.4	5.4	0.0	24.9	100.0	137
Lamu	43.0	18.9	11.0	0.0	27.1	100.0	54
Taita Taveta	50.5	21.4	13.0	0.0	15.1	100.0	153
Garissa	38.3	16.5	7.9	0.0	37.3	100.0	168
Wajir	35.0	13.9	13.0	0.0	38.1	100.0	130
Mandera	42.6	21.7	5.1	0.0	30.6	100.0	123
Marsabit	39.4	19.1	12.3	0.0	29.2	100.0	142
Isiolo	41.1	19.9	15.5	0.0	23.5	100.0	76
Meru	51.2	17.8	12.2	0.0	18.8	100.0	530
Tharaka-Nithi	47.6	23.0	11.5	0.0	17.9	100.0	147
Embu	50.6	20.7	14.1	0.0	14.6	100.0	239
Kitui	53.2	24.7	8.3	0.0	13.7	100.0	412
Machakos	60.0	22.7	6.6	0.0	10.7	100.0	677
Makueni	45.2	28.1	9.3	0.0	17.4	100.0	350
Nyandarua	50.1	20.9	13.8	0.0	15.2	100.0	223
Nyeri	39.7	28.9	16.0	0.0	15.5	100.0	223
Kirinyaga	52.1	20.4	15.2	2.3	10.0	100.0	258
Murang'a	41.5	21.9	18.1	0.3	18.2	100.0	377
Kiambu	45.0	24.3	14.0	0.0	16.7	100.0	1,689
Turkana	45.3	25.3	12.3	0.0	17.1	100.0	178
West Pokot	61.4	21.2	4.7	0.0	12.8	100.0	225
Samburu	44.3	24.5	17.5	0.0	13.7	100.0	98
Trans Nzoia	43.8	22.2	13.6	0.0	20.5	100.0	370
Uasin Gishu	45.5	25.6	14.1	0.0	14.8	100.0	667
Elgeyo-Marakwet	51.6	21.2	10.2	0.0	17.0	100.0	178
Nandi	44.7	23.6	13.8	0.0	17.8	100.0	361
Baringo	48.8	25.6	10.2	0.0	15.4	100.0	187
Laikipia	46.2	19.7	16.1	0.0	18.1	100.0	195
Nakuru	49.5	23.8	11.6	0.0	15.2	100.0	1,078
Narok	56.5	20.8	9.1	0.0	13.7	100.0	428
Kajiado	44.1	27.2	17.5	0.0	11.2	100.0	467
Kericho	45.6	28.2	9.3	0.0	16.9	100.0	387
Bomet	43.1	24.5	13.1	0.0	19.3	100.0	480
Kakamega	46.7	26.9	8.4	0.0	18.1	100.0	665
Vihiga	33.4	28.4	11.3	0.0	26.9	100.0	233
Bungoma	45.1	25.8	9.2	0.0	19.8	100.0	625
Busia	41.4	24.7	12.5	0.3	21.1	100.0	373
Siaya	44.5	25.6	10.9	0.0	19.0	100.0	370
Kisumu	45.4	29.1	9.6	0.0	15.9	100.0	551
Homa Bay	49.7	24.3	9.0	0.0	17.0	100.0	485
Migori	56.4	21.5	10.1	0.0	12.0	100.0	509
Kisii	45.2	27.3	15.1	0.0	12.4	100.0	439
Nyamira	43.6	25.9	12.8	0.0	17.6	100.0	238
Nairobi City	46.5	21.3	14.4	0.6	17.1	100.0	3,550
Total	47.0	23.4	12.1	0.1	17.3	100.0	21,231

[1] Excludes women who had sexual intercourse within the past 4 weeks

Table 4.6 Multiple sexual partners and higher-risk sexual intercourse in the past 12 months among young women

Among all young women age 15–24, percentage who had sexual intercourse with more than one sexual partner in the past 12 months and percentage who had intercourse in the past 12 months with a person who was neither their husband nor someone who lived with them; among those with more than one partner in the past 12 months, percentage who used a condom during last intercourse; and among those who had sexual intercourse in the past 12 months with a person who was neither their husband nor someone who lived with them, percentage who used a condom during last intercourse with such a partner, according to background characteristics, Kenya Mini DHS 2025–26

	Women age 15–24			Women age 15–24 who had 2+ partners in the past 12 months		Women age 15–24 who had intercourse in the past 12 months with a person who was neither their husband nor someone who lived with them	
Background characteristic	Percentage who had 2+ partners in the past 12 months	Percentage who had intercourse in the past 12 months with a person who was neither their husband nor someone who lived with them	Number of women	Percentage who reported using a condom during last sexual intercourse	Number of women	Percentage who reported using a condom during last sexual intercourse with such a partner	Number of women
Age							
15–19	0.9	13.5	3,916	(24.6)	36	39.4	527
15–17	0.4	8.2	2,433	*	10	37.6	200
18–19	1.8	22.1	1,483	(25.1)	26	40.5	328
20–24	3.5	30.4	3,653	32.0	128	33.6	1,111
20–22	3.3	32.4	2,230	26.1	73	33.6	722
23–24	3.9	27.3	1,423	39.8	55	33.6	389
Marital status							
Never married	1.7	26.2	5,528	38.9	93	36.7	1,450
Ever married	3.5	9.2	2,041	19.4	71	25.9	188
Residence							
Urban	2.6	23.2	3,360	26.8	89	33.1	778
Rural	1.8	20.4	4,209	34.6	75	37.6	861
Education[1]							
No education	0.1	1.6	228	*	0	*	4
Primary	2.5	18.9	777	(10.0)	19	20.6	147
Secondary	2.0	18.8	5,383	28.9	108	36.8	1,009
More than secondary	3.1	40.5	1,181	(45.8)	37	37.4	478
Total 15–24	2.2	21.6	7,569	30.4	164	35.5	1,639

Note: Figures in parentheses are based on 25–49 unweighted cases. An asterisk indicates that a figure is based on fewer than 25 unweighted cases and has been suppressed.

[1] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

FERTILITY 5

Key Findings

- ***Total fertility rate:*** The total fertility rate declined from 3.4 children per woman in the 2022 Kenya Demographic and Health Survey (KDHS) to 3.1 children per woman in the 2025–26 Kenya Mini Demographic and Health Survey (KMDHS).
- ***Birth intervals:*** The median birth interval increased from 42.1 months in the 2022 KDHS to 46.3 months in the 2025–26 KMDHS.
- ***Age-specific fertility rate:*** Fertility rises from 63 births per 1,000 women age 15–19 to a peak of 161 births per 1,000 women age 25–29 and then declines among older age groups.
- ***Teenage pregnancy:*** The percentage of women age 15–19 years who have ever been pregnant declined from 14.8% in the 2022 KDHS to 11.9% in the 2025–26 KMDHS.

The number of children that a woman bears depends on many factors, including the age she begins childbearing, how long she waits between births, and her fecundity. Postponing first births and extending the interval between births have played a role in reducing fertility levels in many countries. Delaying first births and spacing births also have positive health consequences. In contrast, short birth intervals (<24 months) can lead to harmful outcomes such as preterm birth, low birth weight, and death for both newborns and their mothers. Childbearing at a very young age is associated with an increased risk of complications during pregnancy and childbirth and higher rates of neonatal mortality.

This chapter describes the current level of fertility in Kenya and some known proximate determinants of fertility. It also presents information on the total fertility rate, birth intervals, insusceptibility to pregnancy (because of postpartum amenorrhoea, postpartum abstinence, or menopause), age at first birth, teenage pregnancy, and rates of induced abortion.

5.1 CURRENT FERTILITY

Total fertility rate

The average number of children a woman would have by the end of her childbearing years if she bore children at the current age-specific fertility rates. Age-specific fertility rates are calculated for the 3 years before the survey, based on detailed pregnancy histories provided by women.

Sample: Women age 15–49

The total fertility rate (TFR) is 3.1 children per woman, meaning that if fertility were to remain at current levels, a woman in Kenya would bear an average of 3.1 children in her lifetime (**Table 5.1**, **Table 5.2**, and **Table 5.2C**).

Table 5.3.1 shows the age-specific fertility rates for the 5 years preceding the survey.

Fertility rises from 63 births per 1,000 women age 15–19 to a peak of 161 births per 1,000 women age 25–29 and then declines among older age groups. In previous surveys, fertility peaked among women age 20–24 and declined thereafter (**Table 5.3.2** and **Figure 5.1**).

Trends: The overall TFR has declined by 1.8 children since the 2003 KDHS (from 4.9 to 3.1 children per woman). The decline has been just as steep in rural areas, where the fertility rate decreased from 5.4 in the 3 years preceding the 2003 KDHS to 3.6 in the 3 years preceding the 2025–26 KMDHS (a decline of 1.8 children). In urban areas, the fertility rate declined more gradually, from 3.3 in the 3 years preceding the 2003 survey to 2.6 in the 3 years preceding the 2025–26 survey (a decline of 0.7 children) (**Table 5.2**, **Table 5.3.2**, and **Figure 5.2**).

Patterns by background characteristics

- The TFR declines with increasing level of education, from 5.4 children per woman among women with no education to 2.7 children per woman among those with more than a secondary education (**Table 5.2** and **Figure 5.3**).
- Similarly, the TFR declines with increasing wealth, from 4.5 children per woman in the lowest wealth quintile to 2.3 children per woman in the highest quintile.
- Fertility levels vary substantially across counties, reflecting marked differentials in reproductive behaviour. Mandera County has the highest TFR (6.8 children per woman), while Embu County has the lowest (2.2 children per woman). Thirty-one of the 47 counties have a TFR above the national estimate of 3.1 children per woman (**Table 5.2C** and **Map 5.1**).

Figure 5.1 **Trends in age-specific fertility**

Births per 1,000 women




Figure 5.2 **Trends in fertility by residence**

Total fertility rates for the 3 years before each survey






Figure 5.3 **Fertility by education**

Total fertility rate for the 3 years before the survey



Map 5.1 Fertility by county

Total fertility rate for the 3 years before the survey



5.2 Children Ever Born and Living

Among all women age 15–49, the mean number of children ever born is 2.1, with a mean of 2.0 surviving children (**Table 5.4**). Among the subset of currently married women, the mean number of children ever born is higher at 3.0, with a mean of 2.9 surviving children.

The number of children ever born increases with age; women age 45–49 have given birth to a mean of 4.5 children, among whom a mean of 4.2 were still living at the time of the survey. Among currently married women of the same age, the mean number of children ever born is 4.8, with a mean of 4.5 surviving children. About 1% of women age 45–49 have never given birth (**Table 5.4**).

5.3 Birth Intervals

Median birth interval

Number of months since the preceding birth by which half of children are born.


Sample: Non-first births among women age 15–49 in the 5 years before the survey

The World Health Organization (WHO) and other international organisations advise individuals and couples to wait at least 24–36 months between births to lower the likelihood of adverse maternal and child health outcomes (WHO 2006). The median birth interval in Kenya is 46.3 months (**Table 5.5**).

Six percent of non-first births occur less than 18 months after the preceding birth, while 22% occur within 2 years after the preceding birth (**Table 5.5** and **Figure 5.4**).

***Figure 5.4* Birth intervals**

Percent distribution of non-first births by number of months since the preceding birth



Trends: The median birth interval in Kenya has consistently exceeded the WHO-recommended 24 months in all DHS surveys, surpassing 36 months by 2014. Between the 2003 and 2025–26 surveys, the median birth interval increased by 13.7 months—from 32.6 to 46.3 months. Only 14% of births occur within 24 months of the preceding birth

Patterns by background characteristics

- The median birth interval increases with maternal age, from 38.7 months for births among women age 20–29 to 61.1 months for births among women age 40–49.
- The median birth interval is substantially shorter if the mother's preceding child has died (27.7 months) than if the preceding child is still alive (47.1 months).
- The median birth interval is shorter if mothers have no education (30.5 months) than if they have a primary, secondary, or more than secondary education (48.5–49.0 months).
- The median birth interval is longer in urban areas (49.8 months) than in rural areas (43.4 months).
- The shortest median birth intervals are in Mandera (24.8 months), Garissa (25.3 months), and Wajir (26.7 months) Counties, while the longest ones are in Embu (68.5 months), Kirinyaga (64.4 months), and Machakos (57.6 months) Counties (**Table 5.5C**).

5.4 INSUSCEPTIBILITY TO PREGNANCY

Postpartum amenorrhoea

The period between the end of a pregnancy and the resumption of menstruation.

Postpartum abstinence

The period between the end of a pregnancy and the resumption of sexual intercourse.

Postpartum insusceptibility

The period during which a woman is considered not at risk of pregnancy because she is postpartum amenorrhoeic and/or abstaining from sexual intercourse postpartum.

Median duration of postpartum amenorrhoea

Number of months after the end of a pregnancy by which half of women have begun menstruating.

Sample: Women age 15–49 who had a live birth or stillbirth in the 3 years before the survey

Median duration of postpartum insusceptibility

Number of months after the end of a pregnancy by which half of women are no longer protected against pregnancy by either postpartum amenorrhoea or abstinence from sexual intercourse.

Sample: Women age 15–49 who had a live birth or stillbirth in the 3 years before the survey

Overall, women are currently insusceptible to pregnancy for 34% of the births that occurred in the 3 years preceding the survey, postpartum amenorrhoeic for 25% of the births, and abstaining from sexual intercourse for 19% of the births (**Table 5.6**).

Trends: Overall, the median duration of postpartum amenorrhoea has declined, from 9.7 months in the 2003 KDHS to 5.8 months in the 2025–26 KMDHS, despite a recent increase from 4.5 months in the 2022 KDHS. In contrast, the median duration of postpartum abstinence has remained relatively stable over time, fluctuating narrowly from 2.9 to 3.7 months and standing at 3.5 months in the 2025–26 KMDHS. The median duration of postpartum insusceptibility has also declined overall, from 11.8 months in the 2003 KDHS to 9.4 months in the 2025–26 KMDHS, although it increased substantially from 7.0 to 9.4 months between 2022 and 2025–26.

Patterns by background characteristics

- The median duration of postpartum amenorrhoea is longer among rural women than among urban women (6.8 versus 4.9 months). The same is true for the median duration of postpartum abstinence (3.9 versus 3.1 months) and the median duration of postpartum insusceptibility (11.2 versus 6.6 months) (**Table 5.7**).
- The median duration of postpartum insusceptibility generally declines with increasing level of education, from 10.6 months among women with no education to 7.2 months for women with more than a secondary education.
- The median duration of postpartum amenorrhoea also decreases with increasing level of education, from 9.7 months for women with no education to 4.0 months among women with more than a secondary education. No clear relationship is found between education and postpartum abstinence.

5.5 Age at First Menstruation

The age at which a woman experiences her first menstrual period marks the biological beginning of her reproductive life, during which she is potentially susceptible to conception. The most common mean age at which women experienced menarche was age 14 (24%), followed by age 15 (22%) (**Table 5.8**).

5.6 MENOPAUSE

Menopause

Women are considered to have reached menopause if they are neither pregnant nor postpartum amenorrhoeic and (1) have not had a menstrual period in the 6 months before the survey, (2) report being menopausal or having had a hysterectomy, or (3) have never menstruated.

Sample: Women age 30–49

Approximately 11% of women age 30–49 are menopausal. The percentage of women who are menopausal increases with age, from 4% among women age 30–34 to 44% among women age 48–49 (**Table 5.9**).

5.7 AGE AT FIRST BIRTH

Median age at first birth

Age by which half of women have had their first child.

Sample: Women age 25–49

The age at which childbearing starts has important consequences for overall fertility as well as the health and welfare of both the mother and her child. Early initiation of childbearing lengthens the reproductive period. The median age at first birth for women age 25–49 is 20.9 years.

The percentage of women who have begun childbearing increases steadily with age. Among women age 25–49, about 6% had a live birth by age 15, 23% by age 18, 42% by age 20, 60% by age 22, and 79% by age 25 (**Table 5.10**).

Women age 35–39 reported the lowest median age at first birth (20.4 years), while women age 25–29 had the highest median age at first birth (21.4 years) (**Table 5.10**).

Patterns by background characteristics

- The median age at first birth increases with level of education, rising from 19.8 years among women with no education to 24.3 years among women with more than a secondary education.
- Similarly, the median age at first birth increases with wealth, from 19.4 years among women in the lowest wealth quintile to 23.0 years among women in the highest wealth quintile (**Table 5.11**).
- The highest median ages at first birth are in Kiambu (22.5 years), Mombasa (22.3 years), Nairobi City (22.0 years), and Embu (21.9 years) Counties, while the lowest are in Homa Bay (18.6 years), Kisumu (19.0 years), Migori (19.0 years), and Samburu (19.1 years) (**Table 5.11C**).

5.8 TEENAGE PREGNANCY

Teenage pregnancy

Percentage of women age 15–19 who have ever been pregnant.

Sample: Women age 15–19

Teenage pregnancy and motherhood remain a major public health and social concern given its association with increased risks of death for both the mother and her child. Twelve percent of women age 15–19 years have

ever been pregnant, 9% have ever had a live birth, and 3% are currently pregnant with their first child (**Table 5.12**).


Among women age 15–19, 1% were pregnant before age 15, less than 1% gave birth before age 15, and 5% had sexual intercourse before age 15 (**Table 5.13**).

Patterns by background characteristics

- The percentage of women age 15–19 who have ever been pregnant increases with age, from 2% among women age 15 to 27% among women age 19 (**Table 5.12**).
- The percentage of women age 15–19 who have ever been pregnant is 13% in rural areas compared with 11% in urban areas.
- The rate of teenage pregnancy is roughly three times higher among women in the lowest wealth quintile than among those in the highest wealth quintile (15% versus 5%).
- Rates of teenage pregnancy vary widely across counties. The highest percentages of women age 15–19 who have ever been pregnant are in West Pokot (25%), Migori (23%), and Kajiado (20%). The lowest percentages are in Isiolo (3%), Makueni (4%), Garissa (5%), and Murang'a (5%) (**Map 5.2**).

Map 5.2 **Teenage pregnancy by county**

Percentage of women age 15–19 who have ever been pregnant



5.9 PREGNANCY OUTCOMES AND INDUCED ABORTION

Live birth

A child who was born alive, even if for a very short time.

Stillbirth

A child who was born dead (no signs of life) following a pregnancy that lasted 7 months (28 weeks) or longer.

Miscarriage

A pregnancy that ended involuntarily before completing 7 months (28 weeks).

Induced abortion

A pregnancy that was voluntarily ended.

Sample: Pregnancies among women age 15–49 ending in the 3 years preceding the survey


Of all pregnancy outcomes in the 3 years before the survey, 89% resulted in live births, 9% in miscarriages, 1% in stillbirths, and less than 1% in induced abortions (**Table 5.14** and **Figure 5.5**). **Table 5.15** also shows age-specific and total induced abortion rates, as well as general abortion rates, for the 3 years preceding the survey. Induced abortions are likely underreported due to stigma and legal restrictions in Kenya.

Patterns by background characteristics

- The percentage of pregnancies that result in a miscarriage increases with age, from 7% among women younger than age 20 to 13% among women age 35–44.

Figure 5.5 **Pregnancy outcomes**

Percent distribution of pregnancies ending in the 3 years before the survey



LIST OF TABLES

For more information on fertility and its determinants, see the following tables:

- **Table 5.1 Current fertility**
- **Table 5.2 Fertility by background characteristics**
- **Table 5.2C Fertility by background characteristics: County level**
- **Table 5.3.1 Trends in age-specific fertility rates**
- **Table 5.3.2 Trends in age-specific and total fertility rates**
- **Table 5.4 Children ever born and living**
- **Table 5.5 Birth intervals**
- **Table 5.5C Birth intervals: County level**
- **Table 5.6 Postpartum amenorrhoea, abstinence, and insusceptibility**
- **Table 5.7 Median durations of postpartum amenorrhoea, abstinence, and insusceptibility**
- **Table 5.8 Age at first menstruation**
- **Table 5.9 Menopause**
- **Table 5.10 Age at first birth**
- **Table 5.11 Median age at first birth**
- **Table 5.11C Median age at first birth: County level**
- **Table 5.12 Teenage pregnancy**
- **Table 5.12C Teenage pregnancy: County level**
- **Table 5.13 Sexual and reproductive health behaviours before age 15**
- **Table 5.14 Pregnancy outcomes by background characteristics**
- **Table 5.14C Pregnancy outcomes by background characteristics: County level**
- **Table 5.15 Induced abortion**

Table 5.1 Current fertility

Age-specific and total fertility rates, the general fertility rate, and the crude birth rate among women for the 3 years preceding the survey, by residence, Kenya Mini DHS 2025–26

	Residence		
Age group	Urban	Rural	Total
10–14	[1]	[1]	[1]
15–19	54	70	63
20–24	127	195	157
25–29	145	181	161
30–34	108	148	128
35–39	62	83	74
40–44	29	34	32
45–49	[3]	[7]	[6]
TFR (15–49)	2.6	3.6	3.1
GFR	98	120	110
CBR	27.0	23.3	24.7

Notes: Age-specific fertility rates are per 1,000 women. Estimates in brackets are truncated. Rates are for the period 1–36 months preceding the interview. Rates for the 10–14 age group are based on retrospective data from women age 15–17.
CBR = crude birth rate (per 1,000 population)
GFR = general fertility rate (per 1,000 women age 15–44)
TFR = total fertility rate (per woman)

Table 5.2 Fertility by background characteristics

Total fertility rate for the 3 years preceding the survey, percentage of women age 15–49 currently pregnant, and mean number of children ever born to women age 40–49 years, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Total fertility rate	Percentage of women age 15–49 currently pregnant	Mean number of children ever born to women age 40–49
Residence			
Urban	2.6	4.7	3.5
Rural	3.6	4.8	4.7
Education[1]			
No education	5.4	7.1	6.2
Primary	3.6	4.5	4.6
Secondary	3.0	4.6	3.6
More than secondary	2.7	4.9	2.8
Wealth quintile			
Lowest	4.5	6.2	5.9
Second	3.5	4.5	4.9
Middle	3.0	4.8	4.2
Fourth	2.9	4.8	3.4
Highest	2.3	4.0	3.2
Total	3.1	4.7	4.2

Note: Total fertility rates are for the period 1–36 months prior to interview.
[1] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 5.2C Fertility by background characteristics: County level

Total fertility rate for the 3 years preceding the survey, percentage of women age 15–49 currently pregnant, and mean number of children ever born to women age 40–49 years, according to county Kenya Mini DHS 2025–26

County	Total fertility rate	Percentage of women age 15–49 currently pregnant	Mean number of children ever born to women age 40–49
Mombasa	2.3	6.1	3.4
Kwale	3.4	6.6	5.6
Kilifi	3.0	4.4	5.3
Tana River	5.7	9.8	6.2
Lamu	3.6	6.6	4.6
Taita Taveta	3.2	4.6	4.0
Garissa	5.3	8.6	6.8
Wajir	5.8	6.5	7.5
Mandera	6.8	8.9	7.7
Marsabit	4.9	6.6	5.6
Isiolo	3.6	5.0	4.9
Meru	2.8	4.1	3.6
Tharaka-Nithi	2.7	3.9	3.8
Embu	2.2	4.8	3.1
Kitui	2.8	4.9	4.8
Machakos	3.2	5.6	3.7
Makueni	3.3	3.8	3.7
Nyandarua	3.8	2.7	4.0
Nyeri	2.8	2.9	3.0
Kirinyaga	2.7	4.5	2.7
Murang'a	3.4	3.1	3.4
Kiambu	2.3	4.0	3.4
Turkana	5.6	6.1	5.9
West Pokot	5.1	7.5	6.1
Samburu	5.0	5.0	5.8
Trans Nzoia	3.0	4.1	4.3
Uasin Gishu	3.0	4.4	3.9
Elgeyo-Marakwet	2.9	3.5	5.0
Nandi	3.4	2.4	4.1
Baringo	4.3	6.2	5.7
Laikipia	3.7	2.8	3.8
Nakuru	3.5	4.7	4.0
Narok	3.2	7.7	4.8
Kajiado	3.6	7.0	3.6
Kericho	2.6	4.0	4.3
Bomet	3.4	4.0	5.2
Kakamega	3.5	5.2	4.8
Vihiga	3.5	5.1	4.6
Bungoma	3.6	7.5	5.5
Busia	3.6	5.1	4.9
Siaya	2.9	2.7	4.8
Kisumu	3.2	3.6	4.4
Homa Bay	3.3	5.1	5.0
Migori	3.8	7.2	5.7
Kisii	2.9	1.9	4.4
Nyamira	3.2	4.2	4.3
Nairobi City	2.6	4.0	3.4
Total	3.1	4.7	4.2

Note: Total fertility rates are for the period 1–36 months preceding the interview.

Table 5.3.1 Trends in age-specific fertility rates

Age-specific fertility rates for 5-year periods preceding the survey, according to age group, Kenya Mini DHS 2025–26

	Number of years preceding survey			
Age group	0–4	5–9	10–14	15–19
10–14	[1]	[5]	[7]	[12]
15–19	66	81	91	120
20–24	162	178	191	216
25–29	165	175	187	208
30–34	133	147	151	[206]
35–39	81	97	[115]	
40–44	35	[50]		
45–49	[6]			

Notes: Age-specific fertility rates are per 1,000 women. Estimates in brackets are truncated. Rates exclude the month of the interview. For the 0–4 year period, rates for the 10–14 age group are based on retrospective data from women age 15–19.

Table 5.3.2 Trends in age-specific and total fertility rates

Age-specific and total fertility rates for the 3-year period preceding several surveys, according to mother's age at the time of the birth, Kenya Mini DHS 2025–26

Mother's age at birth	KDHS 1988–89	KDHS 1993	KDHS 1998	KDHS 2003	KDHS 2008–09	KDHS 2014	KDHS 2022	KMDHS 2025–26
10–14	[2]	[3]	[3]	[3]	[2]	[2]	[2]	[1]
15–19	153	110	111	114	103	96	73	63
20–24	324	257	248	243	238	206	179	157
25–29	301	241	218	231	216	183	172	161
30–34	243	197	188	196	175	148	137	128
35–39	184	154	109	123	118	100	87	74
40–44	99	70	51	55	50	28	35	32
45–49	[29]	[50]	[16]	[15]	[12]	[9]	[5]	[6]
TFR (15–49)	6.7	5.4	4.7	4.9	4.6	3.9	3.4	3.1

Notes: Age-specific fertility rates are per 1,000 women. Rates for the 45–49 age group may be slightly biased due to truncation and are therefore displayed in brackets.
KDHS = Kenya Demographic and Health Survey
KMDHS = Kenya Mini Demographic and Health Survey
TFR = total fertility rate

Table 5.4 Children ever born and living

Percent distribution of all women and currently married women age 15–49 by number of children ever born, mean number of children ever born, and mean number of living children, according to age group, Kenya Mini DHS 2025–26

Age group	Number of children ever born											Total	Number of women	Mean number of children ever born	Mean number of living children
	0	1	2	3	4	5	6	7	8	9	10+				
ALL WOMEN															
15–19	90.6	8.5	0.9	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	100.0	3,916	0.10	0.10
20–24	46.3	37.0	12.7	3.2	0.6	0.1	0.0	0.0	0.0	0.0	0.0	100.0	3,653	0.75	0.73
25–29	14.8	30.5	31.7	14.9	5.6	1.9	0.5	0.1	0.0	0.0	0.0	100.0	3,732	1.74	1.68
30–34	6.1	15.8	26.5	25.8	14.5	5.5	3.9	1.1	0.5	0.2	0.0	100.0	3,108	2.70	2.60
35–39	2.5	7.5	20.3	24.0	19.5	12.7	6.5	3.8	1.9	0.8	0.6	100.0	2,989	3.56	3.41
40–44	1.9	7.4	14.5	21.0	20.8	13.4	8.4	5.6	3.7	1.9	1.4	100.0	2,048	4.00	3.78
45–49	1.3	4.3	13.1	20.2	18.9	14.7	9.4	6.8	5.2	3.1	3.0	100.0	1,784	4.45	4.16
Total	28.8	17.7	17.2	14.1	9.6	5.5	3.2	1.8	1.1	0.6	0.5	100.0	21,231	2.11	2.01
CURRENTLY MARRIED WOMEN															
15–19	34.6	52.2	12.7	0.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	100.0	231	0.79	0.76
20–24	18.9	49.9	23.9	5.6	1.4	0.2	0.1	0.0	0.0	0.0	0.0	100.0	1,584	1.22	1.18
25–29	5.2	27.8	37.1	19.2	7.1	2.6	0.8	0.1	0.0	0.0	0.0	100.0	2,570	2.07	1.99
30–34	3.6	10.0	26.8	29.6	16.3	6.5	4.8	1.4	0.7	0.3	0.1	100.0	2,334	2.97	2.87
35–39	1.3	4.6	17.1	26.3	21.1	13.9	7.3	4.6	2.2	1.0	0.8	100.0	2,216	3.82	3.67
40–44	0.5	4.9	11.1	22.9	24.3	13.4	9.0	5.3	4.6	2.3	1.9	100.0	1,478	4.27	4.03
45–49	1.0	2.0	11.2	17.0	19.1	17.0	11.0	8.2	6.4	3.2	3.9	100.0	1,259	4.82	4.51
Total	5.5	17.6	22.9	20.6	14.1	8.1	4.8	2.7	1.8	0.9	0.8	100.0	11,672	3.02	2.88

Table 5.5 Birth intervals

Percent distribution of non-first live births among women age 15–49 in the 5 years preceding the survey by number of months since preceding live birth, and median number of months since preceding live birth, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Months since preceding birth						Total	Number of non-first births	Median number of months since preceding birth
	7–17	18–23	24–35	36–47	48–59	60+			
Mother's age									
15–19	(22.8)	(18.4)	(48.2)	(9.6)	(0.9)	(0.0)	100.0	36	(26.5)
20–29	7.5	11.6	26.8	18.1	13.3	22.8	100.0	2,871	38.7
30–39	4.2	6.8	19.8	14.8	13.5	40.9	100.0	3,701	51.4
40–49	4.3	5.5	16.3	12.8	9.3	51.8	100.0	741	61.1
Sex of preceding birth									
Male	6.1	8.5	21.6	16.3	13.8	33.7	100.0	3,729	46.3
Female	5.0	8.7	23.0	15.5	12.0	35.8	100.0	3,621	46.3
Survival status of preceding birth									
Living	4.7	8.1	22.5	16.0	13.1	35.6	100.0	7,040	47.1
Dead	25.8	19.4	17.2	13.8	7.5	16.3	100.0	309	27.7
Birth order									
2–3	5.3	8.0	19.8	15.9	13.6	37.2	100.0	4,622	48.5
4–6	5.1	8.5	24.6	15.7	12.3	33.7	100.0	2,239	44.0
7+	10.1	13.9	35.0	15.8	9.1	16.0	100.0	489	32.6
Residence									
Urban	4.8	8.0	19.8	14.6	12.1	40.8	100.0	3,011	49.8
Rural	6.2	9.0	24.0	16.8	13.5	30.6	100.0	4,338	43.4
Mother's education[1]									
No education	10.3	15.8	39.8	13.7	7.8	12.6	100.0	861	30.5
Primary	4.8	7.1	20.7	17.1	12.1	38.2	100.0	2,764	48.5
Secondary	5.8	8.6	19.7	14.3	14.4	37.3	100.0	2,519	48.8
More than secondary	3.6	6.9	19.0	18.1	15.3	37.2	100.0	1,206	49.0
Wealth quintile									
Lowest	8.5	12.3	32.2	17.1	10.7	19.4	100.0	1,798	35.0
Second	4.6	8.0	21.8	16.9	14.8	34.0	100.0	1,418	47.2
Middle	6.1	6.2	17.8	15.3	13.3	41.3	100.0	1,236	52.4
Fourth	3.7	7.5	17.0	15.6	12.6	43.5	100.0	1,579	53.6
Highest	4.4	7.7	19.9	14.0	14.0	39.9	100.0	1,320	50.5
Total	5.6	8.6	22.3	15.9	12.9	34.7	100.0	7,349	46.3

Notes: First-order births are excluded. The interval for multiple births is the number of months since the preceding pregnancy that ended in a live birth. Figures in parentheses are based on 25–49 unweighted cases.

[1] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 5.5C Birth intervals: County level

Percent distribution of non-first live births among women age 15–49 in the 5 years preceding the survey by number of months since preceding live birth, and median number of months since preceding live birth, according to county, Kenya Mini DHS 2025–26

County	Months since preceding birth						Total	Number of non-first births	Median number of months since preceding birth
	7–17	18–23	24–35	36–47	48–59	60+			
Mombasa	1.2	7.9	20.2	16.5	18.9	35.2	100.0	234	50.6
Kwale	4.7	10.9	26.1	19.4	17.7	21.3	100.0	157	40.1
Kilifi	5.6	15.5	28.3	13.4	10.1	27.0	100.0	200	36.4
Tana River	8.4	12.0	33.4	15.1	12.2	18.9	100.0	81	34.0
Lamu	5.7	8.7	26.6	16.5	11.3	31.2	100.0	22	42.3
Taita Taveta	7.1	8.5	20.6	14.9	12.8	36.2	100.0	67	47.2
Garissa	19.6	23.8	33.7	11.2	5.4	6.3	100.0	107	25.3
Wajir	14.1	23.1	36.4	16.4	5.6	4.4	100.0	85	26.7
Mandera	24.0	22.4	29.5	14.7	4.9	4.5	100.0	109	24.8
Marsabit	4.2	10.3	31.6	25.0	15.3	13.7	100.0	83	37.9
Isiolo	4.3	12.1	23.4	24.2	13.3	22.8	100.0	36	41.3
Meru	3.5	5.8	15.2	14.9	16.6	44.0	100.0	163	56.6
Tharaka-Nithi	6.4	6.9	16.4	18.2	11.8	40.3	100.0	41	48.7
Embu	2.6	4.1	9.7	9.0	10.1	64.4	100.0	52	68.5
Kitui	6.5	5.3	20.4	16.9	17.5	33.3	100.0	121	48.4
Machakos	2.9	5.3	17.4	16.6	11.3	46.4	100.0	212	57.6
Makueni	4.0	12.3	6.4	20.4	14.7	42.2	100.0	104	53.3
Nyandarua	2.6	10.0	14.7	12.2	16.2	44.4	100.0	71	57.3
Nyeri	3.1	1.4	14.1	22.3	12.8	46.2	100.0	53	55.6
Kirinyaga	4.3	4.2	12.4	10.8	12.2	56.1	100.0	60	64.4
Murang'a	4.1	1.5	20.1	10.9	16.5	46.8	100.0	102	57.1
Kiambu	4.2	7.8	15.7	15.2	13.2	44.0	100.0	494	55.4
Turkana	3.4	13.2	50.0	16.3	9.3	7.8	100.0	136	32.3
West Pokot	10.4	19.8	31.2	13.1	12.8	12.7	100.0	140	30.5
Samburu	5.4	15.4	39.0	16.7	12.2	11.4	100.0	60	32.4
Trans Nzoia	0.7	6.5	13.9	21.6	14.3	43.0	100.0	110	55.4
Uasin Gishu	5.6	6.1	12.9	19.0	12.4	43.9	100.0	199	53.7
Elgeyo-Marakwet	3.7	12.3	33.4	15.7	13.1	22.0	100.0	63	36.5
Nandi	2.1	10.8	17.5	19.0	16.1	34.5	100.0	114	49.6
Baringo	8.3	9.6	27.5	18.1	11.6	24.8	100.0	89	38.3
Laikipia	7.6	12.9	18.4	15.4	11.5	34.2	100.0	76	46.9
Nakuru	4.3	4.1	22.5	18.1	13.3	37.6	100.0	386	50.2
Narok	8.5	12.0	29.7	15.4	9.1	25.3	100.0	181	35.9
Kajiado	4.3	12.7	30.7	11.8	13.5	26.9	100.0	188	38.4
Kericho	4.6	8.0	18.8	16.4	14.0	38.1	100.0	106	50.3
Bomet	3.4	7.8	31.0	20.6	12.0	25.3	100.0	156	41.7
Kakamega	4.9	6.4	21.4	18.6	13.8	34.8	100.0	234	45.9
Vihiga	3.7	7.3	22.2	14.4	15.2	37.2	100.0	69	49.5
Bungoma	5.3	7.2	16.8	12.9	18.5	39.2	100.0	217	51.0
Busia	6.6	8.8	18.9	15.4	17.1	33.2	100.0	111	48.2
Siaya	5.6	8.8	22.2	15.2	18.4	29.7	100.0	129	45.6
Kisumu	2.6	5.5	18.7	17.3	14.2	41.8	100.0	186	51.4
Homa Bay	4.9	6.8	24.9	17.3	11.9	34.2	100.0	180	45.3
Migori	6.6	5.6	19.2	17.0	17.4	34.3	100.0	225	50.0
Kisii	8.6	5.1	24.0	16.7	8.6	36.9	100.0	157	43.6
Nyamira	4.2	5.0	15.0	23.0	14.5	38.3	100.0	71	50.2
Nairobi City	5.9	6.2	21.6	13.0	9.9	43.3	100.0	1,109	49.2
Total	5.6	8.6	22.3	15.9	12.9	34.7	100.0	7,349	46.3

Notes: First-order births are excluded. The interval for multiple births is the number of months since the preceding pregnancy that ended in a live birth.

Table 5.6 Postpartum amenorrhoea, abstinence, and insusceptibility

Percentage of live births and stillbirths among women 15–49 in the 3 years preceding the survey for which mothers are postpartum amenorrhoeic, abstaining, or insusceptible, by number of months since birth; and median and mean durations, Kenya Mini DHS 2025–26

Months since birth	Percentage of births for which the mother is:			Number of births[2]
	Amenorrhoeic	Abstaining	Insusceptible[1]	
< 2	85.1	89.7	95.4	351
2–3	61.5	55.8	78.8	354
4–5	52.3	22.8	59.6	369
6–7	47.3	22.5	56.3	348
8–9	35.0	24.9	52.7	363
10–11	30.7	18.2	43.8	318
12–13	28.3	15.3	37.8	315
14–15	20.1	14.8	30.3	355
16–17	12.4	12.0	22.6	345
18–19	15.5	11.1	22.5	364
20–21	13.2	8.6	20.9	300
22–23	8.4	7.5	15.0	343
24–25	7.2	6.3	12.8	334
26–27	4.4	6.4	10.6	364
28–29	9.6	7.5	17.1	354
30–31	4.3	8.2	12.1	318
32–33	4.0	6.1	10.0	379
34–35	4.3	4.6	8.6	318
Total	24.9	19.3	34.0	6,191
Median	5.8	3.5	9.4	na
Mean	9.9	7.8	13.1	na

Note: Estimates are based on status at the time of the survey.
na = not applicable
[1] Includes live births and stillbirths for which mothers are either still amenorrhoeic or still abstaining (or both) following birth
[2] Includes live birth and stillbirths

Table 5.7 Median durations of postpartum amenorrhoea, abstinence, and insusceptibility

Median number of months of postpartum amenorrhoea, postpartum abstinence, and postpartum insusceptibility following live births and stillbirths among women 15–49 in the 3 years preceding the survey, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Postpartum amenorrhoea	Postpartum abstinence	Postpartum insusceptibility[1]
Mother's age			
15–29	5.1	3.8	10.3
30–49	6.7	2.9	8.5
Residence			
Urban	4.9	3.1	6.6
Rural	6.8	3.9	11.2
Mother's education[2]			
No education	9.7	3.2	10.6
Primary	6.8	3.4	11.7
Secondary	5.4	4.0	8.3
More than secondary	4.0	2.9	7.2
Wealth quintile			
Lowest	9.3	3.5	11.6
Second	7.0	4.2	10.3
Middle	5.8	4.1	11.2
Fourth	5.6	3.2	10.1
Highest	4.1	(2.9)	4.7
Total	5.8	3.5	9.4

Notes: Medians are based on the status at the time of the survey (current status). Figures in parentheses are based on 25–49 unweighted cases.
[1] Includes births for which mothers are either still amenorrhoeic or still abstaining (or both) following birth
[2] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 5.8 Age at first menstruation

Percent distribution of women age 15–49 by age at menarche, and mean age at menarche, according to current age, Kenya Mini DHS 2025–26

	Age at menarche								Percentage who have never menstruated	Total	Number of women	Mean age at menarche	Number of women who have ever menstruated[1]
Current age	≤10	11	12	13	14	15	≥16	Don't know					
15–19	1.9	2.4	11.9	22.1	29.5	20.8	8.9	0.1	2.4	100.0	3,916	13.8	3,817
20–24	0.6	1.4	9.1	18.5	23.7	22.2	23.8	0.7	0.1	100.0	3,653	14.4	3,626
25–29	1.1	2.3	10.2	18.9	24.8	20.9	20.9	0.9	0.1	100.0	3,732	14.3	3,692
30–34	2.0	1.5	10.0	20.2	22.6	22.8	19.6	1.0	0.3	100.0	3,108	14.2	3,068
35–39	0.9	1.8	9.6	19.4	23.1	21.4	22.0	1.6	0.1	100.0	2,989	14.4	2,938
40–44	0.7	0.4	9.6	16.9	22.3	22.7	24.9	2.4	0.2	100.0	2,048	14.5	1,997
45–49	1.0	0.9	10.6	17.2	21.2	24.8	22.3	1.7	0.4	100.0	1,784	14.5	1,747
Total	1.2	1.7	10.2	19.3	24.4	22.0	19.7	1.0	0.6	100.0	21,231	14.3	20,883

[1] Number of women who gave a numeric response

Table 5.9 Menopause

Percentage of women age 30–49 who are menopausal, according to age, Kenya Mini DHS 2025–26

Age	Percentage menopausal[1]	Number of women
30–34	4.3	3,108
35–39	6.9	2,989
40–41	7.9	987
42–43	11.0	799
44–45	14.7	703
46–47	26.4	784
48–49	43.8	561
Total	10.7	9,930

[1] Percentage of women 1) who are not pregnant, 2) who have had a birth in the past 5 years and are not postpartum amenorrhoeic, and 3) for whom one of the following conditions applies: their last menstrual period occurred 6 or more months preceding the survey, they declared they are in menopause or have had a hysterectomy, or they have never menstruated

Table 5.10 Age at first birth

Percentage of women age 15–49 who had a live birth by exact age, percentage who have never had a live birth, and median age at first live birth, according to current age, Kenya Mini DHS 2025–26

	Percentage who had a live birth by age:					Percentage who have never had a live birth	Number of women	Median age at first live birth
Current age	15	18	20	22	25			
15–19	0.7	na	na	na	na	90.6	3,916	a
20–24	2.4	13.8	31.7	na	na	46.3	3,653	a
25–29	3.8	19.2	37.2	55.6	77.2	14.8	3,732	21.4
30–34	5.4	22.9	40.4	60.4	80.0	6.1	3,108	21.0
35–39	7.1	26.3	46.1	63.0	79.0	2.5	2,989	20.4
40–44	6.9	26.1	42.5	62.7	79.9	1.9	2,048	20.7
45–49	5.6	23.0	44.1	62.5	81.4	1.3	1,784	20.6
20–49	4.9	21.2	39.5	na	na	14.8	17,315	a
25–49	5.6	23.1	41.6	60.3	79.2	6.4	13,662	20.9

na = not applicable (due to censoring)
a = omitted because less than 50% of women had a birth before reaching the beginning of the age group

Table 5.11 Median age at first birth

Median age at first live birth among women age 25–49, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Women age 25–49
Residence	
Urban	21.9
Rural	20.1
Education[1]	
No education	19.8
Primary	19.0
Secondary	21.3
More than secondary	24.3
Wealth quintile	
Lowest	19.4
Second	19.6
Middle	20.3
Fourth	21.5
Highest	23.0
Total	20.9

[1] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 5.11C Median age at first birth: County level

Median age at first live birth among women age 25–49, by county, Kenya Mini DHS 2025–26

County	Women age 25–49
Mombasa	22.3
Kwale	19.8
Kilifi	20.7
Tana River	20.9
Lamu	20.5
Taita Taveta	21.6
Garissa	21.7
Wajir	21.2
Mandera	20.8
Marsabit	21.1
Isiolo	20.5
Meru	20.6
Tharaka-Nithi	21.3
Embu	21.9
Kitui	20.3
Machakos	21.2
Makueni	21.6
Nyandarua	20.8
Nyeri	21.2
Kirinyaga	21.5
Murang'a	21.2
Kiambu	22.5
Turkana	21.8
West Pokot	19.6
Samburu	19.1
Trans Nzoia	20.0
Uasin Gishu	21.6
Elgeyo-Marakwet	20.7
Nandi	19.8
Baringo	20.5
Laikipia	21.0
Nakuru	20.9
Narok	20.1
Kajiado	21.2
Kericho	19.9
Bomet	20.3
Kakamega	20.3
Vihiga	20.1
Bungoma	19.6
Busia	19.8
Siaya	19.3
Kisumu	19.0
Homa Bay	18.6
Migori	19.0
Kisii	19.4
Nyamira	19.6
Nairobi City	22.0
Total	20.9

Table 5.12 Teenage pregnancy

Percentage of women age 15–19 who have ever had a live birth, percentage who have ever had a pregnancy loss, percentage who are currently pregnant, and percentage who have ever been pregnant, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Percentage of women age 15–19 who:				Number of women
	Have ever had a live birth	Have ever had a pregnancy loss[1]	Are currently pregnant	Have ever been pregnant	
Age					
15	0.8	0.0	0.9	1.7	771
16	3.3	0.1	1.3	4.8	809
17	8.2	0.8	1.6	10.3	853
18	13.1	1.0	5.3	17.0	776
19	23.2	3.4	4.6	27.3	707
Residence					
Urban	8.3	1.2	2.5	10.6	1,398
Rural	10.0	0.9	2.8	12.6	2,519
Education[2]					
No education	19.7	1.4	6.0	24.3	77
Primary	28.3	4.3	7.7	34.4	278
Secondary	7.5	0.5	2.3	9.7	3,388
More than secondary	12.0	5.2	1.3	13.3	174
Wealth quintile					
Lowest	12.2	1.1	3.2	15.1	771
Second	11.8	0.9	2.9	14.3	905
Middle	8.3	0.9	3.5	11.8	827
Fourth	9.6	0.7	3.4	12.4	651
Highest	4.7	1.4	0.6	5.4	762
Total	9.4	1.0	2.7	11.9	3,916

[1] Stillbirth, miscarriage, or abortion
[2] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 5.12C Teenage pregnancy: County level

Percentage of women age 15–19 who have ever had a live birth, percentage who have ever had a pregnancy loss, percentage who are currently pregnant, and percentage who have ever been pregnant, according to county, Kenya Mini DHS 2025–26

	Percentage of women age 15–19 who:				
County	Have ever had a live birth	Have ever had a pregnancy loss[1]	Are currently pregnant	Have ever been pregnant	Number of women
Mombasa	4.4	0.0	1.9	6.3	96
Kwale	9.0	0.6	5.1	13.4	94
Kilifi	4.4	0.0	1.0	5.4	137
Tana River	7.3	0.7	5.1	13.0	35
Lamu	8.7	1.0	0.9	8.7	14
Taita Taveta	4.4	0.0	4.5	8.9	25
Garissa	3.1	0.7	1.0	4.8	45
Wajir	5.8	0.0	1.3	7.1	38
Mandera	6.9	0.0	1.5	7.5	31
Marsabit	4.2	1.1	1.8	6.0	40
Isiolo	3.0	0.0	0.4	3.4	18
Meru	8.6	0.0	0.6	9.2	103
Tharaka-Nithi	7.1	0.0	0.0	7.1	27
Embu	6.8	5.4	8.2	18.6	42
Kitui	12.7	1.2	3.3	14.7	82
Machakos	8.2	0.7	4.1	12.3	98
Makueni	1.8	1.8	1.8	3.7	64
Nyandarua	(9.3)	(0.0)	(0.9)	(10.2)	36
Nyeri	(2.0)	(0.0)	(2.0)	(2.0)	33
Kirinyaga	(3.7)	(0.0)	(6.3)	(10.1)	30
Murang'a	5.0	1.1	0.0	5.0	77
Kiambu	4.6	0.0	1.5	5.6	245
Turkana	18.0	1.7	3.2	19.0	37
West Pokot	23.1	2.0	2.0	24.5	52
Samburu	11.7	0.6	2.7	15.0	22
Trans Nzoia	11.7	0.0	1.0	12.7	84
Uasin Gishu	13.1	2.4	2.2	14.3	112
Elgeyo-Marakwet	5.6	0.0	0.0	5.6	35
Nandi	11.6	0.0	2.1	13.7	86
Baringo	9.2	0.0	3.7	12.9	39
Laikipia	4.9	0.0	0.0	4.9	32
Nakuru	12.8	0.0	4.7	15.8	188
Narok	13.6	3.3	2.5	16.2	78
Kajiado	16.2	4.4	3.4	19.6	65
Kericho	10.4	0.0	0.0	10.4	70
Bomet	9.0	0.0	3.4	12.4	106
Kakamega	11.0	4.1	3.2	15.5	139
Vihiga	3.8	0.6	3.9	7.7	66
Bungoma	9.5	0.8	2.0	11.3	158
Busia	9.1	1.0	2.3	11.4	87
Siaya	5.4	0.0	1.7	5.4	76
Kisumu	10.8	2.4	2.3	14.2	106
Homa Bay	13.6	0.0	2.8	15.8	131
Migori	18.3	0.9	6.5	23.3	96
Kisii	16.2	0.0	2.8	17.6	85
Nyamira	8.9	1.7	3.8	13.3	61
Nairobi City	9.6	1.8	3.6	13.2	498
Total	9.4	1.0	2.7	11.9	3,916

Note: Figures in parentheses are based on 25–49 unweighted cases.
[1] Stillbirth, miscarriage, or abortion

Table 5.13 Sexual and reproductive health behaviours before age 15

Among women age 15–19, percentage who initiated sexual intercourse before age 15, percentage who had a live birth before age 15, and percentage who were ever pregnant before age 15, Kenya Mini DHS 2025–26

Age	Had sexual intercourse before age 15	Had a live birth before age 15	Were ever pregnant before age 15	Number of women
15–19	5.1	0.7	1.4	3,916

Table 5.14 Pregnancy outcomes by background characteristics

Percent distribution of pregnancies among women age 15–49 ending in the 3 years preceding the survey by type of outcome, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Pregnancy outcome				Total	Number of pregnancies
	Live birth	Stillbirth[1]	Miscarriage[2]	Induced abortion		
Mother's age at pregnancy outcome						
<20	90.8	0.6	7.1	1.5	100.0	811
20–24	90.4	1.1	8.1	0.4	100.0	1,980
25–34	88.7	1.5	9.1	0.8	100.0	3,200
35–44	86.2	1.1	12.6	0.2	100.0	952
45–49	(79.6)	(1.2)	(19.2)	(0.0)	100.0	25
Pregnancy order						
First	91.7	0.7	7.0	0.6	100.0	1,933
Second	89.7	0.8	8.6	0.8	100.0	1,646
Third	88.0	1.2	10.0	0.8	100.0	1,334
Fourth	85.3	2.1	11.9	0.8	100.0	892
Fifth or higher	87.6	2.1	10.0	0.3	100.0	1,161
Residence						
Urban	85.7	1.6	11.7	1.0	100.0	3,144
Rural	91.7	0.9	7.0	0.4	100.0	3,822
Mother's education[3]						
No education	91.9	3.2	4.9	0.0	100.0	593
Primary	88.3	1.6	9.4	0.7	100.0	1,944
Secondary	90.3	0.5	8.4	0.9	100.0	2,960
More than secondary	86.2	1.5	11.9	0.4	100.0	1,469
Wealth quintile						
Lowest	94.1	1.1	4.4	0.4	100.0	1,362
Second	90.5	0.9	7.5	1.0	100.0	1,281
Middle	90.4	1.3	7.7	0.6	100.0	1,239
Fourth	87.6	0.7	11.0	0.7	100.0	1,617
Highest	83.4	2.1	14.0	0.6	100.0	1,468
Total	89.0	1.2	9.1	0.7	100.0	6,967

Note: Figures in parentheses are based on 25–49 unweighted cases.
[1] Stillbirths are foetal deaths in pregnancies lasting 28 or more weeks. When pregnancy duration is reported in months, stillbirths are foetal deaths in pregnancies lasting 7 or more months.
[2] Miscarriages are foetal deaths in pregnancies lasting less than 28 weeks. When pregnancy duration is reported in months, miscarriages are foetal deaths in pregnancies lasting less than 7 months.
[3] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 5.14C Pregnancy outcomes by background characteristics: County level

Percent distribution of pregnancies among women age 15–49 ending in the 3 years preceding the survey by type of outcome, according to county, Kenya Mini DHS 2025–26

County	Pregnancy outcome				Total	Number of pregnancies
	Live birth	Stillbirth[1]	Miscarriage[2]	Induced abortion		
Mombasa	83.7	3.0	7.6	5.7	100.0	226
Kwale	92.1	0.8	7.1	0.0	100.0	115
Kilifi	87.4	1.1	9.8	1.7	100.0	192
Tana River	89.2	1.9	8.9	0.0	100.0	68
Lamu	89.2	3.0	7.9	0.0	100.0	18
Taita Taveta	91.3	1.1	7.6	0.0	100.0	50
Garissa	89.7	2.3	7.5	0.5	100.0	76
Wajir	91.2	3.8	5.1	0.0	100.0	62
Mandera	94.2	1.0	4.8	0.0	100.0	72
Marsabit	96.2	0.9	3.0	0.0	100.0	58
Isiolo	90.3	1.8	7.9	0.0	100.0	26
Meru	93.9	1.4	4.6	0.0	100.0	129
Tharaka-Nithi	90.3	0.0	9.7	0.0	100.0	40
Embu	87.4	0.0	6.9	5.7	100.0	52
Kitui	95.7	0.0	4.3	0.0	100.0	106
Machakos	95.4	0.0	4.6	0.0	100.0	230
Makueni	90.6	0.0	9.4	0.0	100.0	110
Nyandarua	91.4	0.0	8.6	0.0	100.0	76
Nyeri	89.8	1.6	8.6	0.0	100.0	61
Kirinyaga	89.8	0.0	10.2	0.0	100.0	65
Murang'a	92.3	0.7	6.3	0.7	100.0	106
Kiambu	90.5	0.3	9.1	0.0	100.0	447
Turkana	91.0	1.1	7.8	0.0	100.0	103
West Pokot	97.0	0.4	2.6	0.0	100.0	113
Samburu	95.0	0.3	3.9	0.7	100.0	51
Trans Nzoia	91.7	0.3	7.2	0.9	100.0	106
Uasin Gishu	91.4	2.4	6.3	0.0	100.0	204
Elgeyo-Marakwet	92.7	0.9	6.4	0.0	100.0	50
Nandi	93.4	2.7	3.9	0.0	100.0	114
Baringo	92.9	0.7	6.5	0.0	100.0	83
Laikipia	92.6	0.6	6.8	0.0	100.0	71
Nakuru	89.2	1.2	9.6	0.0	100.0	379
Narok	91.9	0.4	7.7	0.0	100.0	146
Kajiado	85.2	0.5	12.2	2.0	100.0	205
Kericho	93.9	1.6	4.5	0.0	100.0	105
Bomet	94.1	0.5	5.4	0.0	100.0	148
Kakamega	85.1	0.0	13.6	1.3	100.0	234
Vihiga	82.8	3.0	12.7	1.5	100.0	74
Bungoma	93.4	0.0	6.6	0.0	100.0	202
Busia	88.9	0.7	10.5	0.0	100.0	133
Siaya	88.6	0.0	10.1	1.3	100.0	108
Kisumu	88.6	0.6	8.0	2.8	100.0	188
Homa Bay	93.5	0.8	5.1	0.6	100.0	150
Migori	86.3	1.1	12.1	0.4	100.0	220
Kisii	95.9	2.0	1.1	1.0	100.0	125
Nyamira	88.7	2.5	7.4	1.4	100.0	70
Nairobi City	81.6	2.5	15.5	0.5	100.0	1,202
Total	89.0	1.2	9.1	0.7	100.0	6,967

[1] Stillbirths are foetal deaths in pregnancies lasting 28 or more weeks. When pregnancy duration is reported in months, stillbirths are foetal deaths in pregnancies lasting 7 or more months.
[2] Miscarriages are foetal deaths in pregnancies lasting less than 28 weeks. When pregnancy duration is reported in months, miscarriages are foetal deaths in pregnancies lasting less than 7 months.

Table 5.15 Induced abortion rates

Age-specific and total induced abortion rates, and general abortion rates, among women age 15–49 for the 3 years preceding the survey, according to residence, Kenya Mini DHS 2025–26

	Residence		
Age group	Urban	Rural	Total
10–14	[0]	[0]	[0]
15–19	1	1	1
20–24	0	1	1
25–29	3	0	2
30–34	1	0	1
35–39	0	0	0
40–44	0	0	0
45–49	[0]	[0]	[0]
TAR (15–49)	0.0	0.0	0.0
GAR	1	0	1

Notes: Age-specific induced abortion rates are per 1,000 women. Estimates in brackets are truncated. Rates are for the period 1–36 months preceding the interview. Rates for the 10–14 age group are based on retrospective data from women age 15–17.
GAR = general induced abortion rate (per 1,000 women age 15–44)
TAR = total induced abortion rate (per woman)

FERTILITY PREFERENCES 6

Key Findings

- ***Desire for another child:*** About half (49%) of currently married women age 15–49 want to have another child in the future, while 43% want no more children.
- ***Desire to limit childbearing:*** The desire to limit childbearing increases with the number of living children, rising from 1% among women with no children to 75% among women with six or more children.
- ***Ideal family size:*** The mean ideal number of children among all women age 15–49 is 3.7.
- ***Fertility planning status:*** 66% of live births and pregnancies among women age 15–49 that occurred in the 3 years before the survey were wanted at the time; 27% were wanted later, and 8% were not wanted at all.
- ***Wanted fertility:*** The wanted fertility rate is 2.7 children per woman, compared with a total fertility rate of 3.1 children per woman.

Information on fertility preferences can help planners of family planning programme assess the desire for children, the extent of mistimed and unwanted pregnancies, and the demand for contraception to space or limit births. This information suggests the direction that fertility patterns could take in the future.

This chapter presents information on whether and when married women want more children, their ideal family size, whether their last birth was wanted, and the theoretical fertility rate if all unwanted births were prevented.

6.1 DESIRE FOR ANOTHER CHILD

Desire for another child

Women were asked whether they wanted more children and, if so, how long they would prefer to wait before the birth of the next child. Women who are sterilised are assumed not to want any more children.

Sample: Currently married women age 15–49

Among currently married women age 15–49, 49% want to have another child; 18% want to have another child soon (within the next 2 years), while 31% want to wait at least 2 years before having another child (**Table 6.1**). In contrast, 43% of currently married women want no more children.

The percentage of women wanting another child declines steadily as the number of living children increases. The percentage of currently married women who desire another child is highest among women with no children (91%) and those with one child (88%), and lowest (17%) among those with six or more children.


The desire to limit childbearing increases with the number of living children. Among women with no children, 1% want no more children; among women with six or more children, 75% want no more children (**Table 6.2**).

Trends: Across all survey years from 2003 to 2025–26, the desire to stop childbearing increased with the number of living children a woman had (**Figure 6.1**). Figure 6.1 focuses on women with two to four children, although this pattern holds for all women, from those with no children to those with six or more children.

Between the 2003 KDHS and the 2014 KDHS, the percentage of women who wanted to stop childbearing rose for women with three and four children, peaking at 60% among those with three children and 74% among those with four children. Since 2014, the percentage has declined modestly in all groups. By the 2025–26 KMDHS, the percentage stood at 31% for women with two children, 57% for those with three children, and 69% for those with four children (**Figure 6.1**).

Figure 6.1 **Trends in desire to limit childbearing by number of living children**

Percentage of currently married women age 15–49 who want no more children



6.2 IDEAL FAMILY SIZE

Ideal family size

Respondents with no children were asked "If you could choose exactly the number of children to have in your whole life, how many would that be?" Respondents who had children were asked "If you could go back to the time when you did not have any children and could choose exactly the number of children to have in your whole life, how many would that be?"


Sample: Women age 15–49

The mean ideal number of children among all women age 15–49 is 3.7. Among currently married women, the mean ideal number is slightly higher, at 4.1 (**Table 6.3**).

The mean ideal number of children is 3.2 among all women with no living children and 6.5 among those with six or more children (**Figure 6.2**).

Figure 6.2 **Ideal family size by number of living children**

Mean ideal number of children



Patterns by background characteristics

- The mean ideal number of children among women age 15–49 increases steadily with age, from 3.3 among women age 15–19 to 4.6 among women age 45–49 (**Table 6.4**).
- The mean ideal number of children is higher among women in rural areas (3.9) than among women in urban areas (3.5).
- The mean ideal number of children is 4.9 in the lowest wealth quintile compared with 3.4 in the highest wealth quintile.
- The mean ideal number of children is highest in Mandera (8.3), Wajir (8.1), and Garissa (7.7) Counties. It is lowest in Nyeri County (2.8); Kirinyaga County (2.8); and Embu, Makueni, Nandi, Kisii, and Nyamira Counties (2.9 each) (**Table 6.4C**).

6.3 FERTILITY PLANNING STATUS

Status of planning for births/pregnancies

Women reported whether their births/pregnancies were wanted at the time (planned birth), at a later time (mistimed birth), or not at all (unwanted birth).

Sample: Current pregnancies and live births in the 3 years before the survey among women age 15–49 and all pregnancy outcomes in the 3 years before the survey among women age 15–49

Sixty-six percent of live births and current pregnancies among women age 15–49 that occurred in the 3 years before the survey were wanted at the time, while 27% were wanted later and 8% were not wanted at all (**Table 6.5**). Eighty-nine percent of stillbirths in the 3 years before the survey were wanted at the time, compared with 65% of live births and 78% of miscarriages (**Figure 6.3**).

Trends: The percentage of current pregnancies and live births in the 3 years before the survey that were wanted at the time of conception increased from 53% in the 2003 KDHS to 63% in the 2022 KDHS and further to 66% in the 2025–26 KMDHS. During the same period, the percentage of pregnancies that were unwanted declined from 20% in 2003 to 8% in both 2022 and 2025–26, indicating a gradual improvement in fertility planning over time (**Figure 6.4**).

Patterns by background characteristics

- Higher-order births were more likely to be unwanted than lower-order births. Four percent of first births were unwanted, compared with 18% of fourth- and higher-order births (**Table 6.5**).
- Mistimed births and pregnancies decrease with age, from 58% among women younger than age 20 to 9% among women age 40–44.

Figure 6.3 **Fertility planning status**

Percent distribution of pregnancy outcomes to women age 15–49 in the 3 years before the survey by planning status of pregnancy





Figure 6.4 **Trends in fertility planning status**

Percentage of live births and current pregnancies among women age 15–49 in the 3 years before the survey



6.4 WANTED FERTILITY RATES

Unwanted birth

Any birth exceeding the number of children a woman reported as her ideal number.

Wanted birth

Any birth less than or equal to the number of children a woman reported as her ideal number.

Wanted fertility rate

The average number of children a woman would have by the end of her childbearing years if she bore children at the current age-specific fertility rate, excluding unwanted births.


Sample: Women age 15–49

The wanted fertility rate is a hypothetical estimate of the number of children a woman would give birth to during her childbearing years if all unwanted births were avoided. The total wanted fertility rate and the total fertility rate are calculated in the same manner. A birth is considered wanted if the number of living children at the time of conception is lower than the ideal number of children reported by the woman. The total wanted fertility rate for the 3 years preceding the survey was 2.7 children per woman, compared with a total fertility rate of 3.1 children per woman (**Table 6.6**).

Trends: The wanted fertility rate declined from 3.6 children per woman in the 2003 KDHS to 2.9 in the 2022 KDHS and further to 2.7 in the 2025–26 KMDHS. The total fertility rate also declined from 4.9 to 3.1 children per woman during the same period. Consequently, the gap between total and wanted fertility rates narrowed from 1.3 children in 2003 to 0.5 children in 2022 and further to 0.4 children in 2025–26 (**Figure 6.5**).

Figure 6.5 **Trends in wanted and actual fertility**

Wanted and actual number of children per woman




Patterns by background characteristics

- The wanted fertility rate is higher in rural areas than in urban areas, at 3.0 and 2.4 children per woman, respectively.
- The wanted fertility rate decreases with increasing level of education and wealth. It declines from 5.1 children per women with no education to 2.5 children per woman with more than secondary education. Similarly, it declines from 3.7 children per woman in the lowest wealth quintile to 2.1 children per woman in the highest wealth quintile (**Table 6.6**).
- Mandera, Wajir, Tana River, Garissa, and Turkana Counties have wanted fertility rates of more than 5.0 children per woman, while Embu, Kisii, Mombasa, Kitui, Kiambu, Kericho, and Nyamira have wanted fertility rates of less than 2.2 children per woman (**Table 6.6C** and **Map 6.1**).

Map 6.1 Wanted fertility rates by county

Total wanted fertility rates for the 3 years preceding the survey, according to county



LIST OF TABLES

For more information on fertility preferences, see the following tables:

- **Table 6.1 Fertility preferences according to number of living children**
- **Table 6.2 Desire to limit childbearing**
- **Table 6.3 Ideal number of children according to number of living children**
- **Table 6.4 Mean ideal number of children according to background characteristics**
- **Table 6.4C Mean ideal number of children according to background characteristics: County level**
- **Table 6.5 Fertility planning status**
- **Table 6.6 Wanted fertility rates**
- **Table 6.6C Wanted fertility rates: County level**

Table 6.1 Fertility preferences according to number of living children

Percent distribution of currently married women age 15–49 by desire for children, according to number of living children, Kenya Mini DHS 2025–26

	Number of living children[1]							
Desire for children	0	1	2	3	4	5	6+	Total
Have another soon[2]	73.4	27.8	20.2	10.6	6.9	7.1	8.3	17.7
Have another later[3]	16.7	59.7	40.5	24.2	16.8	13.0	8.2	31.0
Have another, undecided when	1.3	0.5	1.3	0.6	0.1	0.2	0.1	0.6
Undecided	2.1	3.7	5.6	6.1	5.0	4.1	3.8	4.8
Want no more	0.9	7.4	30.8	55.4	66.1	70.1	70.6	42.5
Sterilised[4]	0.5	0.0	0.3	1.8	3.3	3.2	4.1	1.6
Declared infecund	5.2	0.8	1.2	1.2	1.8	2.2	4.8	1.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Number	471	2,123	2,883	2,563	1,584	900	1,148	11,672

[1] The number of living children includes a woman's current pregnancy.
[2] Wants next birth within 2 years
[3] Wants to delay next birth for 2 or more years
[4] Includes both female and male sterilisation

Table 6.2 Desire to limit childbearing

Percentage of currently married women age 15–49 who want no more children by number of living children, according to background characteristics, Kenya Mini DHS 2025–26

Background	Number of living children[1]							
characteristic	0	1	2	3	4	5	6+	Total
Residence								
Urban	0.6	7.8	34.4	58.1	67.5	67.5	65.0	36.1
Rural	3.1	6.9	27.4	56.6	70.1	75.3	76.6	50.4
Education[2]								
No education	(0.6)	5.0	16.7	23.0	35.1	40.8	49.2	35.0
Primary	2.7	9.9	26.3	57.4	72.4	79.6	86.0	56.8
Secondary	0.5	6.5	32.7	58.4	72.2	75.6	89.8	38.2
More than secondary	2.0	7.8	35.1	63.1	68.6	68.8	*	34.6
Wealth quintile								
Lowest	0.4	3.9	16.1	38.0	58.6	61.5	63.5	43.2
Second	6.3	5.8	23.8	53.2	71.3	83.0	85.7	52.5
Middle	0.0	11.9	30.7	64.7	76.1	81.8	88.7	51.2
Fourth	0.7	8.0	35.1	63.7	71.6	76.3	69.1	40.9
Highest	1.2	6.2	37.1	58.2	66.7	(58.5)	(75.7)	35.6
Total	1.4	7.5	31.2	57.3	69.3	73.4	74.7	44.1

Notes: Women who have been sterilised or whose husbands have been sterilised are considered to want no more children. Figures in parentheses are based on 25–49 unweighted cases. An asterisk indicates that a figure is based on fewer than 25 unweighted cases and has been suppressed.
[1] The number of living children includes a woman's current pregnancy.
[2] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 6.3 Ideal number of children according to number of living children

Percent distribution of women age 15–49 by ideal number of children, and mean ideal numbers of children for all women and currently married women, according to number of living children, Kenya Mini DHS 2025–26

	Number of living children[1]							
Ideal number of children	0	1	2	3	4	5	6+	Total
0	4.3	3.5	1.7	1.3	1.5	2.6	2.0	2.7
1	3.4	5.1	2.4	2.0	0.9	0.8	0.2	2.7
2	27.0	24.0	22.7	10.8	12.3	8.0	3.7	19.5
3	29.8	37.7	26.0	28.4	9.9	11.7	6.4	26.1
4	19.5	18.7	32.9	31.7	37.1	19.6	17.2	25.1
5	7.0	5.6	8.3	14.1	14.1	23.9	11.3	9.8
6+	8.3	5.0	6.0	10.9	24.1	32.7	57.2	13.4
Non-numeric responses	0.6	0.4	0.1	0.8	0.1	0.8	2.0	0.6
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Number	5,885	3,931	3,886	3,152	1,938	1,105	1,333	21,231
Mean ideal number of children for women 15–49[2]								
All	3.2	3.1	3.4	3.8	4.5	5.0	6.5	3.7
Number	5,847	3,914	3,883	3,126	1,935	1,096	1,307	21,109
Currently married	3.8	3.2	3.5	3.9	4.5	5.1	6.6	4.1
Number of currently married	468	2,114	2,882	2,549	1,581	891	1,122	11,607

[1] The number of living children includes current pregnancy for women.
[2] Means are calculated excluding respondents who gave non-numeric responses.

Table 6.4 Mean ideal number of children according to background characteristics

Mean ideal number of children for all women age 15–49, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Mean	Number of women[1]
Age		
15–19	3.3	3,901
20–24	3.3	3,640
25–29	3.6	3,710
30–34	3.9	3,089
35–39	4.1	2,970
40–44	4.2	2,039
45–49	4.6	1,759
Residence		
Urban	3.5	9,747
Rural	3.9	11,362
Education[2]		
No education	6.9	1,135
Primary	4.2	6,025
Secondary	3.3	9,805
More than secondary	3.1	4,143
Wealth quintile		
Lowest	4.9	3,291
Second	3.8	3,842
Middle	3.6	4,124
Fourth	3.3	4,781
Highest	3.4	5,070
Total	3.7	21,109

[1] Number of women who gave a numeric response
[2] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 6.4C Mean ideal number of children according to background characteristics: County level

Mean ideal number of children for all women age 15–49, according to county, Kenya Mini DHS 2025–26

County	Mean	Number of women[1]
Mombasa	3.6	795
Kwale	5.1	371
Kilifi	4.7	608
Tana River	5.8	137
Lamu	4.3	53
Taita Taveta	3.8	153
Garissa	7.7	168
Wajir	8.1	130
Mandera	8.3	122
Marsabit	6.0	142
Isiolo	5.0	76
Meru	3.1	529
Tharaka-Nithi	3.1	147
Embu	2.9	239
Kitui	3.3	412
Machakos	3.0	670
Makueni	2.9	350
Nyandarua	3.5	221
Nyeri	2.8	223
Kirinyaga	2.8	255
Murang'a	3.3	377
Kiambu	3.1	1,677
Turkana	5.6	171
West Pokot	6.3	225
Samburu	5.4	96
Trans Nzoia	3.6	368
Uasin Gishu	3.1	666
Elgeyo-Marakwet	3.8	178
Nandi	2.9	359
Baringo	4.6	186
Laikipia	3.8	195
Nakuru	3.8	1,074
Narok	4.4	425
Kajiado	3.6	466
Kericho	3.2	384
Bomet	3.6	480
Kakamega	3.6	665
Vihiga	3.6	230
Bungoma	3.9	623
Busia	3.7	367
Siaya	3.6	370
Kisumu	3.6	551
Homa Bay	4.1	485
Migori	4.0	508
Kisii	2.9	439
Nyamira	2.9	238
Nairobi City	3.6	3,505
Total	3.7	21,109

[1] Number of women who gave a numeric response

Table 6.5 Fertility planning status

Percent distribution of live births and current pregnancies among women age 15–49 in the 3 years preceding the survey by planning status of the pregnancy, according to birth order and mother's age at birth; and percent distribution of all pregnancy outcomes among women age 15–49 in the 3 years preceding the survey by planning status of the pregnancy, according to type of pregnancy outcome, Kenya Mini DHS 2025–26

Characteristic	Planning status of pregnancy outcome			Total	Number of pregnancy outcomes[1]
	Wanted then	Wanted later	Wanted no more		
LIVE BIRTHS AND CURRENT PREGNANCIES					
Birth order					
1	55.3	41.0	3.7	100.0	2,225
2	75.0	21.9	3.1	100.0	1,822
3	71.8	21.4	6.8	100.0	1,352
4+	64.5	17.6	17.9	100.0	1,809
Mother's age at birth[2]					
<20	36.9	57.9	5.2	100.0	839
20–24	64.1	32.1	3.8	100.0	2,065
25–29	72.4	22.6	5.0	100.0	1,970
30–34	74.7	16.0	9.3	100.0	1,343
35–39	69.1	12.1	18.7	100.0	746
40–44	63.7	9.4	27.0	100.0	223
45–49	(60.2)	(0.0)	(39.8)	100.0	23
Total	65.7	26.6	7.7	100.0	7,209
ALL PREGNANCY OUTCOMES					
Pregnancy outcome					
Current pregnancy	69.2	26.0	4.7	100.0	1,007
Live birth	65.1	26.7	8.2	100.0	6,202
Stillbirth	88.8	7.8	3.4	100.0	85
Miscarriage	77.5	16.8	5.7	100.0	634
Abortion	(42.1)	(42.2)	(15.7)	100.0	46
Total	66.7	25.7	7.5	100.0	7,974

Notes: A pregnancy outcome refers to a miscarriage, abortion, live birth or stillbirth. Some pregnancies produce multiple outcomes, for example in the case of twins. In this table, each pregnancy outcome is counted individually. Therefore, a pregnancy will be counted more than once if it produces multiple births (live births or stillbirths). Current pregnancies, miscarriages, and abortions are always counted as one pregnancy outcome. Figures in parentheses are based on 25–49 unweighted cases.
[1] For pregnancies that resulted in multiple outcomes (for example, twins), each outcome is counted individually.
[2] For current pregnancies, the maternal age at birth is estimated as the mother's expected age at the time of the birth.

Table 6.6 Wanted fertility rates

Total wanted fertility rates and total fertility rates for the 3 years preceding the survey, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Total wanted fertility rate	Total fertility rate
Residence		
Urban	2.4	2.6
Rural	3.0	3.6
Education[1]		
No education	5.1	5.4
Primary	3.0	3.6
Secondary	2.6	3.0
More than secondary	2.5	2.7
Wealth quintile		
Lowest	3.7	4.5
Second	2.9	3.5
Middle	2.6	3.0
Fourth	2.6	2.9
Highest	2.1	2.3
Total	2.7	3.1

Notes: Rates are calculated based on births among women age 15–49 in the 1–36 months preceding the survey. The total fertility rates are the same as those presented in Table 5.2.
[1] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 6.6C Wanted fertility rates: County level

Total wanted fertility rates and total fertility rates for the 3 years preceding the survey, according to county, Kenya Mini DHS 2025–26

County	Total wanted fertility rate	Total fertility rate
Mombasa	2.1	2.3
Kwale	3.1	3.4
Kilifi	2.6	3.0
Tana River	5.2	5.7
Lamu	3.2	3.6
Taita Taveta	2.7	3.2
Garissa	5.2	5.3
Wajir	5.6	5.8
Mandera	6.7	6.8
Marsabit	4.7	4.9
Isiolo	3.2	3.6
Meru	2.6	2.8
Tharaka-Nithi	2.2	2.7
Embu	1.9	2.2
Kitui	2.1	2.8
Machakos	2.8	3.2
Makueni	2.9	3.3
Nyandarua	3.4	3.8
Nyeri	2.2	2.8
Kirinyaga	2.5	2.7
Murang'a	3.1	3.4
Kiambu	2.1	2.3
Turkana	5.1	5.6
West Pokot	4.8	5.1
Samburu	4.3	5.0
Trans Nzoia	2.4	3.0
Uasin Gishu	2.7	3.0
Elgeyo-Marakwet	2.6	2.9
Nandi	2.3	3.4
Baringo	3.7	4.3
Laikipia	3.0	3.7
Nakuru	3.0	3.5
Narok	2.7	3.2
Kajiado	3.2	3.6
Kericho	2.1	2.6
Bomet	2.3	3.4
Kakamega	3.0	3.5
Vihiga	2.8	3.5
Bungoma	3.0	3.6
Busia	2.9	3.6
Siaya	2.2	2.9
Kisumu	2.4	3.2
Homa Bay	2.6	3.3
Migori	3.2	3.8
Kisii	1.9	2.9
Nyamira	2.1	3.2
Nairobi City	2.5	2.6
Total	2.7	3.1

Notes: Rates are calculated based on births to women age 15–49 in the 1–36 months preceding the survey. The total fertility rates are the same as those presented in Table 5.2C.

FAMILY PLANNING 7

Key Findings

- ***Contraceptive knowledge:*** Awareness of family planning methods is almost universal in Kenya, with 99% of currently married women age 15–49 having heard of at least one modern method.
- ***Contraceptive use:*** 61% of currently married women use a contraceptive method, with 56% using modern methods and 5% using traditional methods.
- ***Contraceptive discontinuation:*** 28% of contraceptive use episodes in the 5 years preceding the survey were discontinued within 12 months. The desire to become pregnant (25%) and side effects or health concerns (24%) were the main reasons for discontinuing a method.
- ***Unmet need for family planning:*** Among currently married women age 15–49, total demand for family planning is 76%. Sixteen percent of these women have an unmet need for family planning, and 61% have a met need.
- ***Future use of family planning:*** 44% of currently married women who are not using a contraceptive method intend to use one in the future.

Couples can use contraceptive methods to limit or space the number of children they would like to have. This chapter presents information on knowledge, use, and sources of various family planning methods as well as rates and reasons for discontinuing them. It also discusses informed choice, unmet need, demand, non-use, and plans to use contraceptive methods in the future. These topics are of practical use to policymakers in forming efficient and effective family planning strategies and policies. Comparisons are made with findings from previous surveys to evaluate trends and progress made in family planning in Kenya over time.

At the 2017 Family Planning Summit, Kenya outlined its commitments for advancing various aspects of family planning by 2030 (Government of Kenya 2017). The country aims to increase modern contraceptive prevalence from 58% in 2017 to 64% by 2030, while reducing unmet need for family planning among all women from 14% to 10% by 2030.

7.1 CONTRACEPTIVE KNOWLEDGE AND USE

According to results of the 2025–26 Kenya Mini Demographic and Health Survey (KMDHS), knowledge of contraceptive methods is nearly universal in Kenya. Almost all currently married women age 15–49 (99%) know at least one modern method of contraception, and 90% are aware of at least one traditional contraceptive method.

The most well-known methods among currently married women are injectables (98%), implants (97%), male condoms (96%), and pills (95%), while the Standard Days Method is the least-known modern contraceptive method (46%). On average, currently married women know of 10 contraceptive methods (**Table 7.1**).

Knowledge of at least one contraceptive method is nearly universal across all background characteristics among currently married women age 15–49. However, slightly lower levels of knowledge are observed among women with no education (92%) and those residing in Mandera (85%) and West Pokot (82%) Counties (**Table 7.2** and **Table 7.2C**).

Contraceptive prevalence

Percentage of women who use any contraceptive method.

Sample: All women age 15–49, currently married women age 15–49, and sexually active unmarried women age 15–49

Modern methods

Include male and female sterilisation, intrauterine devices (IUDs), injectables, implants, contraceptive pills, male and female condoms, emergency contraception, the Standard Days Method, and the lactational amenorrhoea method.


As shown in **Table 7.3**, 44% of women age 15–49 are using a contraceptive method, including 40% who are using a modern method and 4% who are using a traditional method.

Among currently married women, 61% use a contraceptive method, including 56% who use a modern method and 5% who use a traditional method. The most commonly used methods among currently married women are implants (23%), injectables (19%), and contraceptive pills (7%). Among sexually active unmarried women age 15–49, 61% use a contraceptive method, including 53% who use a modern method. For sexually active unmarried women, injectables (16%) and implants (15%) are the most commonly used modern contraceptive methods, followed by male condoms (12%) (**Figure 7.1**).

The use of traditional methods is more common among sexually active unmarried women (8%) than among currently married women (5%).

Figure 7.1 **Contraceptive use**

Percentage of women age 15–49 currently using a contraceptive method



Trends: The use of modern contraceptive methods by currently married women age 15–49 has increased over time, from 32% in the 2003 Kenya Demographic and Health Survey (KDHS) to 53% in the 2014 KDHS and further to 56% in the 2025–26 KMDHS. The greatest gains were in the use of implants, which increased from less than 1% among currently married women before the 2003 survey to 23% in the 2025–26 survey. Use of injectables increased from 14% in 2003 to 26% in 2014 before declining to 20% in 2022 and then to 19% in 2025–26 (**Table 7.4.1** and **Figure 7.2**).

Patterns by background characteristics

- The use of modern contraceptive methods among currently married women increases with number of living children, from 12% among women with no living children to 65% among women with three to four living children (**Table 7.4.2**).
- The use of modern contraceptive methods among currently married women with no education (21%) is lower than that among currently married women with at least a primary education (58%) (**Figure 7.3**).
- The percentage of sexually active unmarried women using traditional methods is two times higher in urban areas (11%) than in rural areas (6%).
- The use of modern contraceptive methods among currently married women is highest in Murang'a (76%), Makueni (72%), Meru (72%), and Kirinyaga (71%) Counties and lowest in Mandera (<1%), Wajir (1%), and Garissa (5%) Counties (**Table 7.4.2C** and **Map 7.1**).

Figure 7.2 Trends in contraceptive use

Percentage of currently married women age 15–49 currently using a contraceptive method






Figure 7.3 Use of modern contraceptive methods by education

Percentage of currently married women age 15–49 using a modern contraceptive method



Map 7.1 Modern contraceptive use by county

Percentage of currently married women age 15–49 using a modern contraceptive method



Timing of Sterilisation

The median age at sterilisation among women age 15–49 is 33.7 years (**Table 7.5**). Most women (62%) reported having undergone sterilisation between age 30 and age 39.

Use of Subcutaneous Depot-medroxyprogesterone Acetate/Sayana Press

Sixteen percent of current injectable users age 15–49 are using subcutaneous depot-medroxyprogesterone acetate (DMPA-SC)/Sayana Press (**Table 7.6**).

Ninety-seven percent of DMPA-SC/Sayana Press users reported to have been injected by a health care provider for their last injection.

7.1.1 Use of Emergency Contraception

Five percent of women age 15–49 reported using emergency contraception in the past 12 months (**Table 7.7**).

Patterns by background characteristics

- The uptake of emergency contraception is highest among women age 20–24 (8%) and women age 25–29 (7%).

- Use of emergency contraception among women is two times higher in urban areas (6%) than in rural areas (3%).
- Less than one percent of women with no education use emergency contraception compared with 9% of women with more than a secondary education.
- Machakos (10%), Kericho (9%), Kajiado (8%), and Kilifi (8%) Counties have the highest percentages of women who used emergency contraception in the 12 months before the survey. Less than 1% of women in Kwale, Lamu, Mandera, Garissa, Wajir, Siaya, and Turkana Counties used emergency contraception in the past 12 months (**Table 7.7C**).

7.1.2 Knowledge of the Fertile Period

Thirty-five percent of women age 15–49 have correct knowledge of the fertile period during the ovulatory cycle, which is 'halfway between two menstrual periods' (**Table 7.8**).

Nearly half (47%) of women using the rhythm method cited 'halfway between two menstrual periods' as the fertile period.

Women age 15–19 are less likely to have correct knowledge of the fertile period (22%) than are women in other age groups (35%–41%) (**Table 7.9**).

7.2 Source of Modern Contraceptive Methods

Source of modern contraceptive methods


The place where the modern method currently being used was obtained the last time it was acquired.

Sample: Women age 15–49 currently using a modern contraceptive method

Sixty-eight percent of all modern contraceptive users obtain their methods from the public sector, including government hospitals (31%), government health centres (15%), and dispensaries (22%). The second most common source of modern methods is the non-nongovernmental organisation private medical sector (30%), including pharmacies (15%), private hospitals (10%), and private clinics (5%) (**Table 7.10** and **Figure 7.4**).

***Figure 7.4* Source of modern contraceptive methods**


Percent distribution of current users of modern methods age 15–49 by most recent source of method



The public sector is the most popular source for implants (88%), female sterilisation (78%), and IUDs (76%). The private sector is the most popular source for emergency contraception (91%), contraceptive pills (66%), and male condoms (64%) (**Figure 7.5**).

***Figure 7.5* Source of specific modern contraceptive methods**

Percent distribution of current users of modern methods age 15–49 by most recent source of method



7.3 Informed Choice

Informed choice

Informed choice indicates that women were informed about their method's side effects, about what to do if they experience side effects, and about other methods they could use.

Sample: Women age 15–49 who are currently using selected modern contraceptive methods and who started the last episode of use within the 5 years before the survey

Sixty-two percent of women using modern contraceptive methods reported being informed about the side effects of their chosen method, 64% were told what to do if they experienced side effects, and 70% were informed about alternative contraceptive methods. Overall, 55% of women received all three types of information. Women who obtained their contraceptives from the public sector were more likely to receive all three types of information (60%) than were women who obtained their contraceptives from the private sector (42%). Additionally, 74% of current users were informed that they could switch to another method if they wanted or needed to (**Table 7.11**).

7.4 Discontinuation of Contraceptives

Contraceptive discontinuation rate

Percentage of contraceptive use episodes discontinued within 12 months.

Sample: Episodes of contraceptive use in the 5 years before the survey experienced by women who are currently age 15–49 (one woman may contribute more than one episode)

Twenty-eight percent of contraceptive use episodes in the 5 years preceding the survey were discontinued within 12 months. The discontinuation rates for pills (38%), injectables (37%), and male condoms (36%) are higher than the overall contraceptive discontinuation rate. Six percent of contraceptive use episodes were discontinued because women switched to another method (**Table 7.12**).

Women cited desire to become pregnant (25%) and other side effects or health concerns (24%) as the primary reasons for discontinuing a method. Among the other cited reasons were infrequent sex/husband away (10%), became pregnant while using (9%), and wanted a more effective method (9%) (**Table 7.13**).

7.5 Demand for Family Planning

Unmet need for family planning

Percentage of women who:

(1) are not pregnant and not postpartum amenorrhoeic, are considered fecund and want to postpone their next birth for 2 or more years or stop childbearing altogether, and are not using a contraceptive method; or

(2) have a mistimed or unwanted current pregnancy; or

(3) are postpartum amenorrhoeic and their most recent birth in the past 2 years was mistimed or unwanted.

Met need for family planning

Current contraceptive use (any method).

Sample: All women age 15–49, currently married women age 15–49, and sexually active unmarried women age 15–49


Demand for family planning	Unmet need for family planning + met need (current contraceptive use [any method])
Proportion of demand satisfied	Current contraceptive use (any method) / Unmet need + current contraceptive use (any method)
Proportion of demand satisfied by modern methods	Current contraceptive use (any modern method) / Unmet need + current contraceptive use (any method)

Seventy-six percent of currently married women age 15–49 have a demand for family planning; 38% want to space births, and another 39% want to limit births. Among those with a demand for family planning, 61% are already using a contraceptive method either to space births (30%) or to limit births (30%), and therefore have their need met. However, 16% of currently married women have an unmet need for family planning; 7% want to space births and 8% want to limit births but are not currently using contraception (**Table 7.14.1** and **Figure 7.6**).

If all currently married women who want to space or limit their children were to use a family planning method, the contraceptive prevalence among currently married women would increase from 61% (currently using) to 76% (total demand).

Figure 7.6 Demand for family planning

Percent distribution of currently married women age 15–49 by need for family planning




Note: Total does not add to 100% due to rounding.

Table 7.14.2 provides insight into the family planning needs of all women and sexually active unmarried women. The total demand for family planning among all women is 55%, with 44% having their need met and 11% having an unmet need. The demand for family planning is higher among sexually active unmarried women (91%) than among currently married women (76%). Among sexually active unmarried women, 61% have their family planning needs met, while 30% have an unmet need.

Trends: Total demand for family planning among currently married women age 15–49 increased from 67% in the 2003 KDHS to 76% in the 2014 KDHS and then remained unchanged at 76% in both the 2022 KDHS and the 2025–26 KMDHS. Unmet need for family planning among currently married women declined from 27% in 2003 to 18% in 2014 and then further to 14% in 2022, before increasing slightly to 16% in 2025–26 (**Figure 7.7**).

Figure 7.7 Trends in demand for family planning

Percentage of currently married women age 15–49 by need for family planning




Patterns by background characteristics

- Unmet need for family planning decreases with increasing level of education, from 29% among currently married women with no education to 13% among those with more than a secondary education (**Table 7.14.1** and **Figure 7.8**).
- The counties with the highest unmet need for family planning among currently married women are West Pokot (42%), Turkana (34%), Tana River (29%), Wajir (28%), Lamu (25%), and Mandera (25%). The county with the lowest total unmet need is Makueni (4%), followed by Embu, Murang'a, Kajiado, and Kirinyanga (7% each) (**Table 7.14.1C** and **Map 7.2**).

Figure 7.8 Unmet need for family planning by education


Percentage of currently married women age 15–49 with unmet need for family planning



- Among all women, the counties with the highest unmet need for family planning are West Pokot (33%), Turkana (24%), and Tana River (20%), while the counties with the lowest total unmet need are Makueni (4%), Murang'a (5%), Kirinyaga (5%), Meru (5%), and Machakos (6%) (**Table 7.14.2C**).

Map 7.2 Unmet need for family planning by county

Percentage of currently married women age 15–49 with unmet need for family planning



7.6 Future Use of Contraception

Forty-four percent of currently married women age 15–49 who are not using a contraceptive method intend to use one in the future, 53% do not intend to use one, and 3% are unsure (**Table 7.15**).

Among currently married women age 15–49 who are not using a contraceptive method, 66% of those with four or more living children do not intend to use one in the future, which is a higher percentage than among women with fewer children.

7.7 Exposure to Family Planning Messages

Peers or friends, radio, social media, and television were the most common sources of family planning messages among women age 15–49 during the 12 months preceding the survey. Eighty percent of the women received family planning messages from peers or friends, 55% from the radio, 50% during community meetings and events, 49% through social media, and 48% from television. Other popular sources of family planning messages were posters/leaflets/brochures (47%), the internet (46%), and outdoor signs or billboards (36%) (**Table 7.16**).

Patterns by background characteristics

- Women age 15–19 were the most likely to report no exposure to family planning messages in the past 12 months. Specifically, 19% of women age 15–19 reported no exposure, compared with 5%–7% of women in other age groups.
- Exposure to family planning messages through radio varies by residence. Fifty-nine percent of women living in rural areas reported seeing family planning messages on the radio, compared with 61% of women living in urban areas.
- Exposure to family planning messages through social media increases with level of education, from only 14% among women with no education to 84% among women with more than a secondary education.
- The counties with the highest percentages of women reporting social media as a source of family planning messages are Kiambu (75%), Nairobi City (62%), Mombasa (62%), and Machakos (60%). The counties with the lowest percentages are Mandera (12%), Tana River (20%), West Pokot (23%), and Marsabit (24%) (**Table 7.16C**).

7.8 Contact of Non-users with Family Planning Providers

Contact of non-users with family planning providers

Percentage of women who discussed family planning in the 12 months before the survey with a fieldworker or during a visit to a health facility.

Sample: Women age 15–49 who are not currently using any contraceptive method

Among women age 15–49 who are not using a contraceptive method, 82% said they had not discussed family planning with a fieldworker or with a provider at a health facility in the 12 months before the survey. Overall, only 5% of non-users reported discussing family planning with a fieldworker and 16% with a provider at a health facility in the past 12 months. However, 39% of non-users reported visiting a health facility and not discussing family planning during that visit (**Table 7.17**).

The percentage of non-users of contraception who did not discuss family planning in the 12 months before the survey ranges from 60% in Turkana County to 94% in Bomet and Wajir Counties (**Table 7.17C**).

List of Tables

For more information on family planning, see the following tables:

- **Table 7.1 Knowledge of contraceptive methods**
- **Table 7.2 Knowledge of contraceptive methods according to background characteristics**
- **Table 7.2C Knowledge of contraceptive methods: County level**
- **Table 7.3 Current use of contraception by age**
- **Table 7.4.1 Trends in current use of contraception**
- **Table 7.4.2 Current use of contraception according to background characteristics**
- **Table 7.4.2C Current use of contraception: County level**
- **Table 7.5 Timing of sterilisation**
- **Table 7.6 Use of subcutaneous depot-medroxyprogesterone acetate/Sayana Press**
- **Table 7.7 Use of emergency contraception**
- **Table 7.7C Use of emergency contraception: County level**
- **Table 7.8 Knowledge of fertile period**
- **Table 7.9 Knowledge of fertile period by age**
- **Table 7.10 Source of modern contraceptive methods**
- **Table 7.11 Informed choice**
- **Table 7.12 Twelve-month contraceptive discontinuation rates**
- **Table 7.13 Reasons for discontinuation**
- **Table 7.14.1 Need and demand for family planning among currently married women**
- **Table 7.14.1C Need and demand for family planning among currently married women: County level**
- **Table 7.14.2 Need and demand for family planning among all women and sexually active unmarried women**
- **Table 7.14.2C Need and demand for family planning among all women: County level**
- **Table 7.15 Future use of contraception**
- **Table 7.16 Exposure to family planning messages**
- **Table 7.16C Exposure to family planning messages: County level**
- **Table 7.17 Contact of non-users with family planning providers**
- **Table 7.17C Contact of non-users with family planning providers: County level**

Table 7.1 Knowledge of contraceptive methods

Percentages of all women, currently married women, and sexually active unmarried women age 15–49 who know any contraceptive methods, according to specific method, Kenya Mini DHS 2025–26

Method	All women	Currently married women	Sexually active unmarried women[1]
Any method	97.2	99.3	99.7
Any modern method	97.1	99.2	99.7
Female sterilisation	65.5	73.4	70.4
Male sterilisation	47.2	52.6	51.4
Intrauterine device (IUD)	75.8	87.5	82.4
Injectables	91.7	97.7	97.1
Implants	89.1	96.8	95.6
Pills	88.6	95.3	95.2
Male condoms	94.3	96.3	98.4
Female condoms	67.1	72.9	75.1
Emergency contraception	76.0	78.4	87.7
Standard Days Method	40.0	45.8	42.0
Lactational amenorrhoea method	56.3	69.1	54.3
Other modern method	0.4	0.5	0.7
Any traditional method	83.3	89.7	91.6
Rhythm	76.1	82.2	83.9
Withdrawal	68.4	76.1	79.4
Other traditional method	2.6	3.6	2.7
Mean number of methods known by women 15–49	9.4	10.3	10.2
Number of women	21,231	11,672	1,018

[1] Had last sexual intercourse within 30 days preceding the survey

Table 7.2 Knowledge of contraceptive methods according to background characteristics

Percentages of currently married women age 15–49 who have heard of at least one contraceptive method and who have heard of at least one modern method, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Heard of any method	Heard of any modern method[1]	Number
Age			
15–19	95.6	95.6	231
20–24	98.6	98.6	1,584
25–29	99.6	99.6	2,570
30–34	99.5	99.5	2,334
35–39	99.5	99.5	2,216
40–44	99.3	99.3	1,478
45–49	99.1	98.8	1,259
Residence			
Urban	99.7	99.7	5,174
Rural	98.9	98.9	6,498
Education[2]			
No education	92.0	91.5	896
Primary	99.7	99.7	4,288
Secondary	100.0	100.0	4,207
More than secondary	99.9	99.9	2,280
Wealth quintile			
Lowest	96.5	96.4	1,993
Second	99.9	99.9	2,107
Middle	99.9	99.9	2,230
Fourth	99.9	99.9	2,626
Highest	99.6	99.5	2,716
Total 15–49	99.3	99.2	11,672

[1] Female sterilisation, male sterilisation, intrauterine devices, injectables, implants, pills, male condoms, female condoms, emergency contraception, Standard Days Method, lactational amenorrhoea method, and other modern methods

[2] No education includes informal education (e.g., madrassa/ duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 7.2C Knowledge of contraceptive methods: County level

Percentages of currently married women age 15–49 who have heard of at least one contraceptive method and who have heard of at least one modern method, according to county, Kenya Mini DHS 2025–26

County	Heard of any method	Heard of any modern method[1]	Number
Mombasa	100.0	100.0	444
Kwale	99.7	99.7	205
Kilifi	99.7	99.7	302
Tana River	99.7	99.7	88
Lamu	92.9	92.9	27
Taita Taveta	100.0	100.0	85
Garissa	91.1	90.2	90
Wajir	94.5	94.5	72
Mandera	85.4	85.2	77
Marsabit	98.6	98.6	77
Isiolo	99.4	99.4	36
Meru	100.0	100.0	308
Tharaka-Nithi	100.0	100.0	80
Embu	100.0	100.0	129
Kitui	100.0	100.0	251
Machakos	100.0	100.0	420
Makueni	100.0	100.0	192
Nyandarua	100.0	100.0	122
Nyeri	99.4	99.4	104
Kirinyaga	100.0	100.0	141
Murang'a	100.0	100.0	182
Kiambu	100.0	100.0	819
Turkana	94.6	94.6	116
West Pokot	82.2	82.2	145
Samburu	99.6	99.6	58
Trans Nzoia	100.0	100.0	197
Uasin Gishu	99.7	99.7	335
Elgeyo-Marakwet	100.0	100.0	92
Nandi	99.6	99.6	188
Baringo	92.1	92.1	109
Laikipia	99.3	99.3	105
Nakuru	99.7	99.7	614
Narok	99.8	99.8	283
Kajiado	99.6	99.6	261
Kericho	100.0	100.0	198
Bomet	100.0	100.0	242
Kakamega	100.0	100.0	406
Vihiga	100.0	100.0	118
Bungoma	99.9	99.9	349
Busia	100.0	99.5	212
Siaya	99.8	99.8	206
Kisumu	100.0	100.0	296
Homa Bay	100.0	100.0	279
Migori	100.0	100.0	314
Kisii	100.0	100.0	253
Nyamira	100.0	100.0	123
Nairobi City	99.4	99.3	1,920
Total 15–49	99.3	99.2	11,672

[1] Female sterilisation, male sterilisation, intrauterine devices, injectables, implants, pills, male condoms, female condoms, emergency contraception, Standard Days Method, lactational amenorrhoea method, and other modern methods

Table 7.3 Current use of contraception by age

Percent distribution of all women, currently married women, and sexually active unmarried women age 15–49 by contraceptive method currently used, according to age, Kenya Mini DHS 2025–26

				Modern method											Traditional method					
Age	Any method	Any modern method	Female sterilisation	IUD	Injectables	Implants	Pills	Male condoms	Female condoms	Emergency contraception	SDM	LAM	Other	Any traditional method	Rhythm	Withdrawal	Other	Not currently using	Total	Number of women
										ALL WOMEN										
15–19	8.0	7.1	0.0	0.0	1.6	2.8	0.4	2.0	0.0	0.1	0.0	0.1	0.0	0.9	0.7	0.1	0.0	92.0	100.0	3,916
20–24	40.8	35.7	0.0	0.8	11.0	15.4	3.0	4.5	0.0	0.5	0.1	0.3	0.0	5.1	2.8	2.3	0.0	59.2	100.0	3,653
25–29	56.3	52.5	0.0	1.5	17.5	22.9	5.3	3.3	0.0	0.6	0.1	1.3	0.0	3.8	2.4	1.2	0.2	43.7	100.0	3,732
30–34	57.3	53.5	0.4	2.8	18.1	23.9	6.4	1.2	0.0	0.1	0.1	0.3	0.1	3.8	2.9	0.8	0.1	42.7	100.0	3,108
35–39	58.7	54.5	1.8	3.8	18.4	21.4	6.8	1.6	0.0	0.2	0.2	0.5	0.0	4.2	3.4	0.5	0.3	41.3	100.0	2,989
40–44	54.7	49.1	4.1	5.5	13.8	15.8	7.5	1.3	0.0	0.1	0.2	0.7	0.0	5.6	4.5	0.4	0.6	45.3	100.0	2,048
45–49	39.8	35.8	5.1	5.3	9.6	9.5	4.3	1.7	0.0	0.1	0.2	0.1	0.0	4.0	3.2	0.3	0.5	60.2	100.0	1,784
Total	43.7	39.9	1.1	2.3	12.6	16.0	4.5	2.4	0.0	0.3	0.1	0.5	0.0	3.7	2.7	0.9	0.2	56.3	100.0	21,231
										CURRENTLY MARRIED WOMEN										
15–19	41.9	39.9	0.0	0.0	13.2	20.6	4.1	0.6	0.0	0.6	0.0	0.8	0.0	2.0	1.7	0.3	0.0	58.1	100.0	231
20–24	57.2	51.9	0.0	0.8	18.4	24.4	5.8	1.8	0.0	0.3	0.0	0.3	0.1	5.3	2.5	2.8	0.0	42.8	100.0	1,584
25–29	61.7	58.7	0.0	1.8	20.0	26.1	7.1	1.8	0.0	0.2	0.1	1.6	0.0	3.1	1.9	0.8	0.3	38.3	100.0	2,570
30–34	63.3	59.3	0.5	2.9	20.1	27.0	7.3	0.8	0.0	0.1	0.1	0.4	0.0	4.0	2.9	1.0	0.1	36.7	100.0	2,334
35–39	66.1	61.6	2.0	4.6	21.8	23.3	7.8	1.0	0.0	0.2	0.2	0.6	0.0	4.5	3.5	0.6	0.3	33.9	100.0	2,216
40–44	63.8	56.9	3.7	7.1	16.3	17.7	9.3	1.4	0.0	0.0	0.3	1.0	0.1	6.9	5.5	0.5	0.9	36.2	100.0	1,478
45–49	47.9	42.7	5.8	5.7	12.6	11.7	4.9	1.7	0.0	0.1	0.3	0.1	0.0	5.1	4.2	0.4	0.6	52.1	100.0	1,259
Total	60.7	56.1	1.6	3.5	18.7	22.8	7.1	1.4	0.0	0.1	0.2	0.7	0.0	4.5	3.2	1.0	0.3	39.3	100.0	11,672
										SEXUALLY ACTIVE UNMARRIED WOMEN[1]										
15–19	47.3	34.3	0.0	1.1	2.9	8.5	1.3	16.9	0.8	2.2	0.0	0.0	0.6	12.9	9.8	3.1	0.0	52.7	100.0	120
20–24	59.8	49.1	0.0	0.5	11.6	9.1	4.0	21.1	0.0	2.8	0.0	0.0	0.0	10.6	7.6	3.1	0.0	40.2	100.0	279
25+	63.7	57.6	0.9	1.9	20.0	19.6	8.2	6.4	0.1	0.3	0.0	0.0	0.1	6.2	3.2	2.8	0.1	36.3	100.0	620
Total	60.7	52.5	0.6	1.4	15.7	15.4	6.2	11.7	0.2	1.2	0.0	0.0	0.2	8.2	5.2	2.9	0.1	39.3	100.0	1,018

Notes: If more than one method is used, only the most effective method is considered in this tabulation. Male sterilisation is not shown due to too few cases.
IUD = intrauterine device
LAM = lactational amenorrhoea method
SDM = Standard Days Method
[1] Had last sexual intercourse within 30 days preceding the survey

Table 7.4.1 Trends in current use of contraception

Percent distribution of currently married women age 15–49 by contraceptive method currently used, according to several surveys, Kenya Mini DHS 2025–26

Method	KDHS 1988–89	KDHS 1993	KDHS 1998	KDHS 2003	KDHS 2008–09	KDHS 2014	KDHS 2022	KMDHS 2025–26
Any method	**26.9**	**32.7**	**39.0**	**39.3**	**45.5**	**58.0**	**62.5**	**60.7**
Any modern method	**17.9**	**27.3**	**31.5**	**31.5**	**39.4**	**53.2**	**56.9**	**56.1**
Female sterilisation	4.7	5.5	6.2	4.3	4.8	3.2	2.3	1.6
Intrauterine device (IUD)	3.7	4.2	2.7	2.4	1.6	3.4	4.4	3.5
Injectables	3.3	7.2	11.8	14.3	21.6	26.4	19.9	18.7
Implants	0.0	0.0	0.8	1.7	1.9	9.9	18.5	22.8
Pills	5.2	9.5	8.5	7.5	7.2	8.0	8.1	7.1
Male condoms	0.5	0.8	1.3	1.2	1.8	2.2	1.8	1.4
Standard Days Method	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.2
Lactational amenorrhoea method	0.0	0.0	0.0	0.0	0.5	0.1	0.9	0.7
Other modern method	0.5	0.1	0.2	0.1	0.0	0.0	0.5	0.1
Any traditional method	**9.0**	**5.5**	**7.5**	**7.8**	**6.0**	**4.8**	**5.6**	**4.5**
Rhythm	7.5	4.4	6.1	6.3	4.7	3.8	4.0	3.2
Withdrawal	0.2	0.4	0.6	0.6	0.7	0.7	1.2	1.0
Other traditional method	1.3	0.7	0.8	0.9	0.6	0.3	0.4	0.3
Not currently using	**73.1**	**67.3**	**61.0**	**60.7**	**54.5**	**42.0**	**37.5**	**39.3**
Total	100	100	100	100	100	100	100	100
Number of women	4,765	4,629	4,834	4,919	4,928	18,549	17, 822	11,672

KDHS = Kenya Demographic and Health Survey
KMDHS = Kenya Mini Demographic and Health Survey

Table 7.4.2 Current use of contraception according to background characteristics

Percent distribution of currently married and sexually active unmarried women age 15–49 by contraceptive method currently used, according to background characteristics, Kenya Mini DHS 2025–26

			Modern method												Traditional method					
Background characteristic	Any method	Any modern method	Female sterilisation	Male sterilisation	IUD	Injectables	Implants	Pills	Male condoms	Female condoms	Emergency contraception	SDM	LAM	Other	Any traditional method	Rhythm	Withdrawal	Other	Not currently using	Total	Number of women
CURRENTLY MARRIED WOMEN																					
Number of living children																					
0	19.3	11.7	0.3	0.0	0.2	3.0	0.9	1.9	4.5	0.0	0.9	0.0	0.0	0.0	7.6	2.4	5.1	0.0	80.7	100.0	678
1–2	62.1	58.0	0.2	0.0	3.2	19.1	24.3	9.3	1.0	0.0	0.1	0.1	0.5	0.0	4.1	3.0	0.9	0.2	37.9	100.0	4,959
3–4	69.9	65.2	2.4	0.0	4.8	21.8	25.7	7.6	1.5	0.0	0.1	0.2	1.1	0.0	4.7	3.6	0.5	0.5	30.1	100.0	4,045
5+	52.3	48.1	3.8	0.1	2.4	16.9	20.5	2.2	1.1	0.0	0.1	0.2	0.8	0.0	4.2	3.1	0.7	0.4	47.7	100.0	1,989
Residence																					
Urban	60.4	54.7	0.8	0.0	4.6	15.6	20.6	10.0	1.8	0.0	0.2	0.2	0.9	0.0	5.7	4.0	1.3	0.4	39.6	100.0	5,174
Rural	60.9	57.3	2.3	0.0	2.6	21.3	24.5	4.7	1.0	0.0	0.1	0.2	0.6	0.0	3.6	2.6	0.7	0.3	39.1	100.0	6,498
Education[1]																					
No education	24.1	21.3	0.9	0.0	0.4	8.4	8.6	0.9	0.2	0.0	0.1	0.0	1.9	0.0	2.8	1.3	1.3	0.2	75.9	100.0	896
Primary	62.2	58.4	2.7	0.0	2.5	21.7	24.0	5.4	1.0	0.0	0.1	0.2	0.9	0.0	3.8	3.0	0.3	0.5	37.8	100.0	4,288
Secondary	64.7	60.2	0.9	0.0	2.5	21.3	25.1	7.9	1.6	0.0	0.2	0.2	0.5	0.0	4.5	3.3	0.9	0.3	35.3	100.0	4,207
More than secondary	64.7	58.0	1.1	0.0	8.3	12.6	21.8	11.2	2.0	0.0	0.2	0.3	0.4	0.1	6.7	4.2	2.3	0.2	35.3	100.0	2,280
Wealth quintile																					
Lowest	48.4	45.2	1.3	0.0	0.8	18.5	21.2	1.7	0.7	0.0	0.0	0.1	0.8	0.0	3.2	2.2	0.9	0.1	51.6	100.0	1,993
Second	65.2	62.5	2.4	0.1	1.8	23.1	28.4	4.4	1.2	0.0	0.1	0.2	0.7	0.0	2.7	2.2	0.4	0.2	34.8	100.0	2,107
Middle	63.8	60.1	2.4	0.0	2.7	22.1	25.3	5.2	1.4	0.0	0.2	0.2	0.5	0.1	3.7	2.9	0.5	0.4	36.2	100.0	2,230
Fourth	62.7	57.6	1.2	0.0	4.1	20.7	20.2	8.2	2.1	0.0	0.2	0.2	0.6	0.0	5.0	3.7	0.7	0.6	37.3	100.0	2,626
Highest	61.6	54.5	0.8	0.0	6.7	10.9	20.1	13.4	1.3	0.0	0.1	0.1	1.0	0.0	7.1	4.5	2.2	0.3	38.4	100.0	2,716
Total	60.7	56.1	1.6	0.0	3.5	18.7	22.8	7.1	1.4	0.0	0.1	0.2	0.7	0.0	4.5	3.2	1.0	0.3	39.3	100.0	11,672
SEXUALLY ACTIVE UNMARRIED WOMEN[2]																					
Residence																					
Urban	64.6	54.1	0.0	0.0	1.8	14.9	13.0	9.3	14.0	0.0	1.0	0.0	0.0	0.0	10.5	6.3	4.2	0.0	35.4	100.0	544
Rural	56.3	50.8	1.2	0.0	1.0	16.6	18.2	2.7	9.0	0.3	1.5	0.0	0.0	0.3	5.5	3.9	1.5	0.2	43.7	100.0	475
Total	60.7	52.5	0.6	0.0	1.4	15.7	15.4	6.2	11.7	0.2	1.2	0.0	0.0	0.2	8.2	5.2	2.9	0.1	39.3	100.0	1,018

Note: If more than one method is used, only the most effective method is considered in this tabulation.
IUD = intrauterine device
LAM = lactational amenorrhoea method
SDM = Standard Days Method
[1] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.
[2] Had sexual intercourse within 30 days preceding the survey

Table 7.4.2C Current use of contraception: County level

Percent distribution of currently married women age 15–49 by contraceptive method currently used, according to county, Kenya Mini DHS 2025–26

			Modern method												Traditional method					
County	Any method	Any modern method	Female sterilisation	Male sterilisation	IUD	Injectables	Implants	Pills	Male condoms	Emergency contraception	SDM	LAM	Other	Any traditional method	Rhythm	Withdrawal	Other	Not currently using	Total	Number of women
Mombasa	47.5	44.3	0.7	0.0	2.7	17.4	15.3	4.0	1.4	0.9	0.4	1.5	0.0	3.2	3.2	0.0	0.0	52.5	100.0	444
Kwale	45.8	45.5	0.0	0.0	0.8	20.8	20.2	3.1	0.0	0.4	0.0	0.0	0.0	0.3	0.3	0.0	0.0	54.2	100.0	205
Kilifi	63.0	57.2	2.1	0.0	1.0	21.2	24.8	5.2	1.3	0.0	0.0	1.6	0.0	5.8	5.4	0.4	0.0	37.0	100.0	302
Tana River	21.8	21.8	0.2	0.0	0.5	5.6	14.4	1.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	78.2	100.0	88
Lamu	39.9	39.6	2.3	0.0	3.1	17.7	13.1	2.9	0.5	0.0	0.0	0.0	0.0	0.4	0.0	0.4	0.0	60.1	100.0	27
Taita Taveta	72.2	65.0	1.5	0.0	4.1	20.5	26.1	8.6	2.9	0.8	0.0	0.6	0.0	7.1	7.1	0.0	0.0	27.8	100.0	85
Garissa	5.4	5.0	0.0	0.0	0.3	2.3	2.0	0.4	0.0	0.0	0.0	0.0	0.0	0.4	0.1	0.3	0.0	94.6	100.0	90
Wajir	0.8	0.8	0.0	0.0	0.2	0.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	99.2	100.0	72
Mandera	0.3	0.3	0.0	0.0	0.0	0.2	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	99.7	100.0	77
Marsabit	35.0	30.4	0.0	0.0	0.0	10.3	8.1	2.0	0.4	0.5	0.0	9.1	0.0	4.6	1.6	2.5	0.5	65.0	100.0	77
Isiolo	34.9	32.3	0.5	0.0	0.0	12.6	14.9	2.9	1.4	0.0	0.0	0.0	0.0	2.6	0.8	1.1	0.7	65.1	100.0	36
Meru	76.4	71.5	3.1	0.0	3.8	29.2	24.6	9.5	0.0	0.0	0.0	1.1	0.0	5.0	3.3	1.3	0.4	23.6	100.0	308
Tharaka-Nithi	69.1	64.9	2.9	0.8	9.0	28.2	18.0	4.0	1.9	0.0	0.0	0.0	0.0	4.3	2.2	2.0	0.0	30.9	100.0	80
Embu	73.4	69.7	1.0	0.0	5.1	26.4	20.9	15.3	1.0	0.0	0.0	0.0	0.0	3.7	0.6	3.1	0.0	26.6	100.0	129
Kitui	64.3	62.6	3.2	0.0	3.2	24.2	28.9	2.3	0.0	0.2	0.0	0.2	0.4	1.8	1.3	0.4	0.0	35.7	100.0	251
Machakos	75.1	68.1	0.7	0.0	1.6	21.6	32.2	8.3	1.1	0.0	0.0	2.4	0.3	7.0	4.7	0.5	1.7	24.9	100.0	420
Makueni	79.6	71.7	3.4	0.0	6.7	27.2	27.4	5.5	1.0	0.0	0.0	0.0	0.4	7.9	5.0	1.5	1.4	20.4	100.0	192
Nyandarua	68.6	65.6	1.5	0.0	8.2	19.0	22.2	11.3	3.3	0.0	0.0	0.0	0.0	2.9	2.4	0.5	0.0	31.4	100.0	122
Nyeri	61.5	60.7	1.8	0.0	8.5	11.2	21.0	18.2	0.0	0.0	0.0	0.0	0.0	0.7	0.7	0.0	0.0	38.5	100.0	104
Kirinyaga	76.7	71.2	1.3	0.0	8.9	19.6	19.6	20.9	0.5	0.0	0.5	0.0	0.0	5.5	4.3	0.4	0.8	23.3	100.0	141
Murang'a	78.4	75.6	4.3	0.0	8.5	26.8	20.4	13.8	1.2	0.0	0.5	0.0	0.0	2.8	2.6	0.2	0.0	21.6	100.0	182
Kiambu	61.6	57.3	0.8	0.0	7.1	13.0	19.7	15.3	0.5	0.5	0.0	0.4	0.0	4.3	1.5	2.3	0.5	38.4	100.0	819
Turkana	21.4	17.0	0.0	0.0	0.6	6.8	8.1	0.8	0.0	0.0	0.0	0.7	0.0	4.3	3.7	0.6	0.0	78.6	100.0	116
West Pokot	18.3	17.6	0.0	0.0	0.5	5.7	8.8	1.2	0.9	0.0	0.5	0.0	0.0	0.7	0.7	0.0	0.0	81.7	100.0	145
Samburu	44.6	37.4	1.0	0.0	0.5	21.0	11.3	0.8	2.0	0.0	0.0	0.9	0.0	7.2	4.2	3.0	0.0	55.4	100.0	58
Trans Nzoia	56.7	54.3	2.5	0.0	4.5	25.1	18.5	2.2	0.6	0.0	0.8	0.0	0.0	2.4	1.4	1.0	0.0	43.3	100.0	197
Uasin Gishu	63.1	56.5	0.8	0.0	4.7	16.6	27.6	4.9	1.9	0.0	0.0	0.0	0.0	6.6	3.7	2.9	0.0	36.9	100.0	335
Elgeyo-Marakwet	69.3	63.3	2.2	0.0	1.4	19.4	30.6	3.5	1.8	0.0	0.5	3.9	0.0	6.0	4.5	1.5	0.0	30.7	100.0	92
Nandi	55.9	55.5	1.8	0.4	3.9	19.0	25.5	3.6	0.9	0.0	0.4	0.0	0.0	0.5	0.5	0.0	0.0	44.1	100.0	188
Baringo	66.6	47.7	1.1	0.0	1.4	14.3	24.4	2.9	2.7	0.0	0.4	0.6	0.0	18.8	17.1	1.1	0.6	33.4	100.0	109
Laikipia	62.5	55.8	2.9	0.0	5.6	26.6	4.8	13.7	1.0	0.0	0.0	1.2	0.0	6.7	4.9	1.8	0.0	37.5	100.0	105
Nakuru	65.4	62.2	1.3	0.0	3.5	25.6	23.0	7.4	1.1	0.0	0.0	0.2	0.0	3.2	3.2	0.0	0.0	34.6	100.0	614
Narok	59.8	55.3	1.3	0.0	4.7	17.0	30.3	1.1	0.0	0.0	0.9	0.0	0.0	4.5	0.6	3.9	0.0	40.2	100.0	283
Kajiado	63.3	53.8	1.8	0.0	3.8	17.9	17.7	10.0	1.1	0.3	0.4	0.7	0.0	9.5	6.2	3.3	0.0	36.7	100.0	261
Kericho	62.8	58.7	2.8	0.0	2.9	22.1	28.4	0.6	1.4	0.0	0.5	0.0	0.0	4.1	2.9	1.2	0.0	37.2	100.0	198
Bomet	55.0	51.3	4.4	0.0	1.5	19.1	24.5	1.8	0.0	0.0	0.0	0.0	0.0	3.7	3.3	0.4	0.0	45.0	100.0	242
Kakamega	67.0	66.1	2.5	0.0	4.0	18.4	35.9	4.8	0.1	0.3	0.0	0.0	0.0	0.9	0.9	0.0	0.0	33.0	100.0	406
Vihiga	62.8	61.3	4.6	0.0	1.8	15.2	35.3	2.8	1.0	0.0	0.0	0.6	0.0	1.5	0.7	0.8	0.0	37.2	100.0	118
Bungoma	57.9	55.8	2.7	0.0	1.7	20.6	25.6	1.7	1.1	0.0	1.2	1.3	0.0	2.1	1.6	0.0	0.5	42.1	100.0	349
Busia	59.1	56.4	2.2	0.0	1.6	14.5	31.7	2.8	0.7	1.0	0.0	1.8	0.0	2.6	2.6	0.0	0.0	40.9	100.0	212
Siaya	51.6	51.3	2.3	0.3	1.5	22.1	20.6	1.2	1.3	0.0	0.7	1.4	0.0	0.3	0.2	0.2	0.0	48.4	100.0	206
Kisumu	74.0	66.1	3.5	0.0	1.5	23.3	27.8	4.2	4.3	0.4	0.0	1.2	0.0	7.9	7.1	0.2	0.6	26.0	100.0	296
Homa Bay	68.9	65.3	1.8	0.0	0.9	21.3	32.4	3.3	3.8	0.7	0.5	0.6	0.0	3.6	2.7	0.8	0.0	31.1	100.0	279
Migori	60.5	57.0	3.7	0.0	2.0	13.7	32.0	2.5	2.1	0.0	0.0	1.1	0.0	3.4	2.8	0.3	0.3	39.5	100.0	314
Kisii	71.3	68.7	3.0	0.0	2.6	31.0	24.0	6.2	1.6	0.0	0.0	0.5	0.0	2.5	1.8	0.7	0.0	28.7	100.0	253
Nyamira	69.0	59.0	3.3	0.0	0.6	30.5	20.9	3.2	0.5	0.0	0.0	0.0	0.0	10.0	4.4	2.9	2.8	31.0	100.0	123
Nairobi City	61.5	54.6	0.0	0.0	4.0	15.2	19.4	12.4	2.7	0.0	0.0	0.9	0.0	6.9	5.0	1.3	0.6	38.5	100.0	1,920
Total	60.7	56.1	1.6	0.0	3.5	18.7	22.8	7.1	1.4	0.1	0.2	0.7	0.0	4.5	3.2	1.0	0.3	39.3	100.0	11,672

Note: If more than one method is used, only the most effective method is considered in this tabulation.
IUD = intrauterine device
LAM = lactational amenorrhoea method
SDM = Standard Days Method

Table 7.5 Timing of sterilisation

Percent distribution of sterilised women age 15–49 by age at time of sterilisation and median age at sterilisation, according to number of years since the operation, Kenya Mini DHS 2025–26

Years since operation	Age at time of sterilisation						Total	Number of women	Median age[1]
	<25	25–29	30–34	35–39	40–44	45–49			
<2	(0.0)	(1.4)	(10.4)	(35.4)	(29.7)	(23.1)	100.0	28	(36.6)
2–3	(0.0)	(4.8)	(16.8)	(50.6)	(17.7)	(10.1)	100.0	48	(36.7)
4–5	(1.0)	(4.5)	(31.9)	(34.3)	(25.9)	(2.4)	100.0	34	(34.8)
6–7	(0.0)	(6.4)	(37.1)	(49.0)	(7.5)	(0.0)	100.0	28	(35.2)
8–9	*	*	*	*	*	*	100.0	12	*
10+	7.4	38.3	37.8	16.5	0.0	0.0	100.0	92	a
Total	3.1	18.1	29.6	32.3	11.9	5.0	100.0	241	33.7

Note: Figures in parentheses are based on 25–49 unweighted cases. An asterisk indicates a figure is based on fewer than 25 unweighted cases and has been suppressed.
a = not calculated due to censoring
[1] Median age at sterilisation is calculated only for women sterilised before age 40 to avoid problems of censoring.

Table 7.6 Use of subcutaneous depot-medroxyprogesterone acetate/Sayana Press

Percentage of current injectable users age 15–49 using DMPA-SC/Sayana Press; and among women using DMPA-SC/Sayana Press, percent distribution by the person administering the injection the last time, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Percentage using DMPA-SC/ Sayana Press	Number of women using injectables	Among DMPA-SC/Sayana Press users, person administering injection the last time:			
			Self-injection	Health care provider	Total	Number of women
Age						
15–19	21.9	62	*	*	100.0	14
20–24	21.9	403	0.6	99.4	100.0	88
25–29	14.4	652	2.8	97.2	100.0	94
30–34	15.9	562	3.2	96.8	100.0	89
35–39	12.4	550	5.4	94.6	100.0	68
40–44	17.4	283	0.0	100.0	100.0	49
45–49	14.3	172	(10.9)	(89.1)	100.0	25
Residence						
Urban	11.8	1,040	1.0	99.0	100.0	122
Rural	18.5	1,644	3.8	96.2	100.0	305
Education[1]						
No education	20.4	83	*	*	100.0	17
Primary	15.2	1,095	2.5	97.5	100.0	166
Secondary	15.7	1,136	2.3	97.7	100.0	178
More than secondary	17.8	370	3.4	96.6	100.0	66
Wealth quintile						
Lowest	16.8	432	3.2	96.8	100.0	73
Second	20.8	590	4.3	95.7	100.0	123
Middle	18.4	613	3.0	97.0	100.0	113
Fourth	13.1	694	2.2	97.8	100.0	91
Highest	7.9	355	(0.0)	(100.0)	100.0	28
Total	15.9	2,684	3.0	97.0	100.0	427

Notes: DMPA-SC is a self-injectable contraceptive; its brand name is Sayana Press. Figures in parentheses are based on 25–49 unweighted cases. An asterisk indicates that a figure is based on fewer than 25 unweighted cases and has been suppressed.
DMPA-SC = subcutaneous depot-medroxyprogesterone acetate
[1] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 7.7 Use of emergency contraception

Percentage of women age 15–49 who used emergency contraception in the past 12 months, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Percentage who used emergency contraception	Number of women
Age		
15–19	1.7	3,916
20–24	8.2	3,653
25–29	6.7	3,732
30–34	4.8	3,108
35–39	3.5	2,989
40–44	2.6	2,048
45–49	1.9	1,784
Residence		
Urban	5.7	9,829
Rural	3.4	11,402
Education[1]		
No education	0.4	1,161
Primary	2.8	6,053
Secondary	4.3	9,849
More than secondary	8.6	4,167
Wealth quintile		
Lowest	1.6	3,312
Second	3.1	3,852
Middle	4.3	4,133
Fourth	5.9	4,800
Highest	6.3	5,134
Total	4.5	21,231

[1] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 7.7C Use of emergency contraception: County level

Percentage of women age 15–49 who used emergency contraception in the past 12 months, according to county, Kenya Mini DHS 2025–26

County	Percentage who used emergency contraception	Number of women
Mombasa	4.2	803
Kwale	0.4	374
Kilifi	7.7	608
Tana River	1.2	137
Lamu	0.3	54
Taita Taveta	6.5	153
Garissa	0.5	168
Wajir	0.5	130
Mandera	0.3	123
Marsabit	2.8	142
Isiolo	2.2	76
Meru	2.0	530
Tharaka-Nithi	4.7	147
Embu	6.0	239
Kitui	1.6	412
Machakos	10.2	677
Makueni	3.0	350
Nyandarua	3.1	223
Nyeri	5.7	223
Kirinyaga	7.0	258
Murang'a	3.5	377
Kiambu	2.4	1,689
Turkana	0.7	178
West Pokot	1.3	225
Samburu	2.1	98
Trans Nzoia	1.8	370
Uasin Gishu	7.2	667
Elgeyo-Marakwet	7.2	178
Nandi	3.4	361
Baringo	4.5	187
Laikipia	2.1	195
Nakuru	6.1	1,078
Narok	5.3	428
Kajiado	8.1	467
Kericho	8.5	387
Bomet	4.6	480
Kakamega	4.1	665
Vihiga	5.3	233
Bungoma	1.7	625
Busia	2.1	373
Siaya	0.6	370
Kisumu	6.5	551
Homa Bay	3.9	485
Migori	4.8	509
Kisii	5.4	439
Nyamira	3.5	238
Nairobi City	5.4	3,550
Total	4.5	21,231

Table 7.8 Knowledge of fertile period

Percent distribution of rhythm users, SDM users, and all women age 15–49, according to knowledge of the fertile period during the ovulatory cycle, Kenya Mini DHS 2025–26

Perceived fertile period	Users of rhythm method	Users of SDM	All women
Just before her menstrual period begins	13.1	(19.3)	14.4
During her menstrual period	0.4	(0.0)	1.3
Right after her menstrual period has ended	33.3	(41.9)	30.2
Halfway between two menstrual periods	46.5	(33.7)	34.6
Other	0.0	(0.0)	0.1
No specific time	2.8	(5.1)	7.8
Don't know	3.8	(0.0)	11.5
Missing	0.0	(0.0)	0.0
Total	100.0	100.0	100.0
Number of women	563	26	21,231

Note: Figures in parentheses are based on 25–49 unweighted cases.
SDM = Standard Days Method

Table 7.9 Knowledge of fertile period by age

Percentage of women age 15–49 with correct knowledge of the fertile period during the ovulatory cycle, according to age, Kenya Mini DHS 2025–26

Age	Percentage with correct knowledge of the fertile period	Number of women
15–19	22.0	3,916
20–24	35.4	3,653
25–29	40.6	3,732
30–34	38.5	3,108
35–39	35.1	2,989
40–44	36.6	2,048
45–49	38.3	1,784
Total	34.6	21,231

Note: Correct knowledge of the fertile period is defined as 'halfway between two menstrual periods'.

Table 7.10 Source of modern contraceptive methods

Percent distribution of modern contraceptive users age 15–49 by most recent source of method, according to specific method, Kenya Mini DHS 2025–26

Source	Female sterili-sation	IUD	Inject-ables	Implants	Pills	Male condoms	Emergency contra-ception	Other modern method[1]	Total
Public sector	**77.8**	**76.2**	**64.7**	**87.6**	**32.8**	**9.8**	**2.6**	**(23.9)**	**67.5**
Government hospital	62.1	54.4	22.9	39.1	17.8	4.3	0.0	(11.2)	30.6
Government health centre	10.3	12.4	14.5	19.5	6.8	1.8	1.7	(6.0)	14.5
Government dispensary	5.4	8.8	27.0	28.5	7.6	3.1	0.9	(5.5)	22.0
Mobile clinic	0.0	0.4	0.3	0.3	0.5	0.5	0.0	(0.0)	0.3
Community health worker/ fieldworker	0.0	0.2	0.0	0.2	0.1	0.1	0.0	(1.1)	0.1
Private medical sector (non-NGO)	**16.6**	**22.5**	**35.1**	**11.4**	**65.5**	**64.1**	**91.2**	**(16.8)**	**29.7**
Private hospital	16.6	18.2	12.4	7.4	7.1	1.8	0.0	(0.9)	9.5
Private clinic	0.0	3.5	9.7	2.6	4.8	1.6	0.0	(0.0)	5.0
Pharmacy	0.0	0.8	12.4	1.2	53.6	60.0	91.2	(12.7)	14.9
Private doctor	0.0	0.1	0.2	0.1	0.0	0.2	0.0	(0.0)	0.1
Mobile clinic	0.0	0.0	0.2	0.0	0.0	0.2	0.0	(3.1)	0.1
Community health worker/ fieldworker	0.0	0.0	0.2	0.0	0.0	0.3	0.0	(0.0)	0.1
Private medical sector (NGO)	**1.5**	**0.7**	**0.0**	**0.4**	**0.0**	**0.0**	**2.2**	**(1.9)**	**0.3**
NGO hospital	1.1	0.0	0.0	0.1	0.0	0.0	0.0	(1.9)	0.1
NGO clinic	0.4	0.7	0.0	0.2	0.0	0.0	2.2	(0.0)	0.2
FBO medical sector	**3.8**	**0.6**	**0.1**	**0.4**	**0.0**	**0.1**	**0.0**	**(1.3)**	**0.4**
FBO/mission clinic	3.8	0.6	0.1	0.4	0.0	0.1	0.0	(1.3)	0.3
Other source	**0.3**	**0.0**	**0.1**	**0.2**	**1.5**	**22.5**	**4.0**	**(55.7)**	**1.9**
Shop	0.0	0.0	0.1	0.0	1.5	19.3	3.2	(5.5)	1.4
Friends/relatives	0.0	0.0	0.0	0.2	0.0	2.9	0.8	(20.7)	0.4
Other	0.3	0.0	0.0	0.0	0.0	0.3	0.0	(29.5)	0.2
Don't know	0.0	0.0	0.0	0.0	0.1	3.5	0.0	(0.4)	0.2
Total	100.0	100.0	100.0	99.9	100.0	100.0	100.0	100.0	100.0
Number of women	241	493	2,684	3,402	959	503	60	35	8,378

Notes: Total includes other modern methods but excludes lactational amenorrhoea method. Figures in parentheses are based on 25–49 unweighted cases.
FBO = faith-based organisation
IUD = intrauterine device
NGO = nongovernmental organisation
SDM = Standard Days Method
[1] Male sterilisation, female condoms, SDM, lactational amenorrhoea method, and other modern methods

Table 7.11 Informed choice

Among current users of selected modern methods age 15–49 who started the last episode of use within the 5 years preceding the survey, percentage who were informed about possible side effects or problems of that method, percentage who were informed about what to do if they experienced side effects, percentage who were informed about other methods they could use, percentage who were informed of all three types of information, and percentage who were informed that they could switch to another method if they wanted or needed to, according to method and initial source, Kenya Mini DHS 2025–26

	Among women who started last episode of modern contraceptive method use within 5 years preceding the survey:					
Method/source	Percentage who were informed about side effects or problems of method used	Percentage who were informed about what to do if side effects experienced	Percentage who were informed of other methods that could be used	Percentage who received all three types of information (Method Information Index)[1]	Percentage who were informed that they could switch to another method if they wanted or needed to	Number of women
Method						
Female sterilisation	55.3	53.2	50.9	40.7	na	91
Intrauterine device (IUD)	77.8	82.9	85.5	73.8	89.2	274
Injectables	56.0	58.6	65.8	48.9	70.5	2,218
Implants	68.7	71.4	73.9	61.7	79.4	2,901
Pills	50.5	46.8	61.7	42.3	65.3	778
Initial source of method[2]						
Public sector	66.7	69.7	73.8	59.7	78.1	4,588
Government hospital	72.8	75.9	78.1	65.2	80.5	2,011
Government health centre	62.4	66.0	71.9	57.5	77.9	985
Government dispensary	61.3	64.1	69.7	54.1	75.1	1,566
Mobile clinic	*	*	*	*	*	21
Community health worker/ fieldworker	*	*	*	*	*	5
Private medical sector (non-NGO)	49.5	48.3	58.2	41.5	61.9	1,614
Private hospital	64.9	64.1	70.2	58.9	73.6	587
Private clinic	53.7	55.7	60.2	45.8	63.8	325
Pharmacy	34.2	30.9	47.0	24.4	50.9	691
Private doctor	*	*	*	*	*	4
Mobile clinic	*	*	*	*	*	3
Community health worker/ fieldworker	*	*	*	*	*	4
Private medical sector (NGO)	*	*	*	*	*	14
NGO hospital	*	*	*	*	*	6
NGO clinic	*	*	*	*	*	8
FBO medical sector	(65.6)	(64.8)	(65.5)	(64.8)	(67.5)	24
FBO/mission clinic	*	*	*	*	*	20
FBO/mobile clinic	*	*	*	*	*	1
Other NGO medical sector	*	*	*	*	*	2
Other source	*	*	*	*	*	20
Shop	*	*	*	*	*	14
Friends/relatives	*	*	*	*	*	6
Other	*	*	*	*	*	1
Don't know	*	*	*	*	*	1
Total	62.1	64.1	69.7	55.0	73.8	6,262

Notes: Table includes users of only the methods listed individually. Figures in parentheses are based on 25–49 unweighted cases. An asterisk indicates that a figure is based on fewer than 25 unweighted cases and has been suppressed.
na = not applicable
FBO = faith-based organisation
NGO = nongovernmental organisation
[1] The Method Information Index is the percentage who were informed about (1) side effects or problems of method used, (2) what to do if side effects are experienced, and (3) other methods that could be used.
[2] Source at start of current episode of use

Table 7.12 Twelve-month contraceptive discontinuation rates

Among episodes of contraceptive use experienced within the 5 years preceding the survey, percentage of episodes discontinued within 12 months, according to reason for discontinuation and specific method, Kenya Mini DHS 2025–26

Method	Method failure	Desire to become pregnant	Other fertility-related reasons[1]	Changes in menstrual bleeding	Other side effects/ health concerns	Wanted more effective method	Other method-related reasons[2]	Husband/ partner disapproved	Other reasons[3]	Any reason[4]	Switched to another method[5]	Number of episodes of use[6]
IUD	0.5	3.7	0.4	0.4	4.9	0.4	0.2	0.1	0.2	11.0	2.3	546
Injectables	1.5	7.4	3.2	4.6	11.2	2.3	5.1	0.5	0.8	36.5	6.0	4,926
Implants	0.5	2.3	0.3	2.0	6.9	0.3	1.3	0.1	0.3	13.9	2.1	4,818
Pills	5.7	4.8	5.0	2.3	8.5	5.6	3.5	0.1	2.4	37.9	10.7	1,777
Male condoms	1.7	2.2	21.4	0.1	1.0	4.7	2.2	1.4	1.1	35.8	5.4	748
Withdrawal	5.4	3.3	8.1	0.0	0.3	7.3	1.2	0.7	0.6	26.9	7.1	319
Other[7]	5.1	3.3	4.9	1.7	1.0	12.0	3.3	0.4	1.7	33.3	12.5	1,478
All methods	2.1	4.5	3.5	2.7	7.4	3.2	3.0	0.3	0.9	27.6	5.8	14,610

Note: Figures are based on life table calculations using information on episodes of use that occurred 3–62 months preceding the survey.
[1] Includes infrequent sex/husband away, difficult to get pregnant/menopausal, and marital dissolution/separation
[2] Includes lack of access/too far, costs too much, and inconvenient to use
[3] Includes up to God/fatalistic and other reasons
[4] Reasons for discontinuation are mutually exclusive and add to the total given in this column.
[5] A woman is considered to have switched to another method if she used a different method in the month following discontinuation or if she gave 'wanted a more effective method' as the reason for discontinuation and started another method within 2 months of discontinuation.
[6] All episodes of use that occurred within the 5 years preceding the survey are included. Episodes of use include both episodes that were discontinued and episodes that were not discontinued during the period of observation.
[7] Emergency contraception, lactational amenorrhoea method, female condoms, rhythm method, Standard Days Method, other modern methods, and other traditional methods

Table 7.13 Reasons for discontinuation

Percent distribution of discontinuations of contraceptive methods in the 5 years preceding the survey by main reason stated for discontinuation, according to specific method, Kenya Mini DHS 2025–26

Reason	IUD	Injectables	Implants	Pills	Male condoms	Emergency contraception	Rhythm	Withdrawal	Other[1]	All methods
Became pregnant while using	3.2	6.6	2.5	14.9	9.3	8.5	34.4	26.3	10.2	8.7
Wanted to become pregnant	42.7	27.0	25.9	22.9	9.4	4.4	26.8	14.7	9.3	24.5
Husband/partner disapproved	1.0	0.9	0.7	0.1	4.6	1.6	1.1	5.8	0.2	1.0
Wanted a more effective method	3.9	5.8	3.3	11.4	17.0	14.4	18.6	22.4	45.3	8.8
Changes in menstrual bleeding	3.4	10.3	9.4	4.7	0.6	12.6	0.4	0.0	4.8	7.7
Other side effects/health concerns	26.8	26.0	33.7	20.9	3.0	11.1	0.0	0.4	1.6	23.5
Lack of access/too far	0.8	1.9	0.7	1.5	1.6	2.1	0.0	0.0	0.9	1.3
Costs too much	0.2	1.1	0.0	0.6	0.2	0.9	0.0	0.0	0.5	0.6
Inconvenient to use	0.7	1.9	0.9	5.0	5.1	5.9	3.0	4.0	6.3	2.5
Up to God/fatalistic	0.5	0.6	0.1	0.2	0.2	1.1	1.2	0.6	0.3	0.4
Difficult to get pregnant/ menopausal	1.1	0.3	0.2	1.0	0.1	0.0	0.6	0.0	0.0	0.4
Infrequent sex/husband away	2.8	7.7	3.0	10.9	44.6	37.2	9.9	19.9	10.5	9.5
Marital dissolution/ separation	0.7	0.7	0.8	0.5	2.2	0.0	0.2	3.6	0.5	0.8
Method expired	8.9	7.7	16.6	0.5	0.0	0.0	0.0	0.1	5.5	7.9
Other	2.9	1.0	1.3	1.8	0.7	0.0	1.5	0.4	2.4	1.3
Don't know	0.0	0.1	0.1	0.3	0.6	0.4	0.2	0.7	0.1	0.2
Missing	0.4	0.4	0.7	2.8	0.9	0.0	2.0	1.1	1.8	1.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Number of discontinuations	220	3,542	2,368	1,338	433	122	424	190	278	8,916

IUD = intrauterine device
[1] Male sterilisation, female condoms, Standard Days Method, lactational amenorrhoea method, other modern methods, and other traditional methods

Table 7.14.1 Need and demand for family planning among currently married women

Percentage of currently married women age 15–49 with unmet need for family planning, percentage with met need for family planning, total demand for family planning, percentage of the demand for family planning that is satisfied, and percentage of the demand for family planning that is satisfied by modern methods, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Unmet need for family planning			Met need for family planning (currently using)			Total demand for family planning[1]			Number of women	Percentage of demand satisfied[2]	Percentage of demand satisfied by modern methods[3]
	For spacing	For limiting	Total	For spacing	For limiting	Total	For spacing	For limiting	Total			
Age												
15–19	18.9	0.9	19.8	39.6	2.3	41.9	58.5	3.2	61.7	231	67.9	64.6
20–24	11.1	1.5	12.7	51.6	5.7	57.2	62.7	7.2	69.9	1,584	81.9	74.2
25–29	11.2	3.3	14.5	47.1	14.7	61.7	58.2	18.0	76.2	2,570	81.0	77.0
30–34	6.3	5.4	11.8	33.5	29.8	63.3	39.9	35.2	75.1	2,334	84.3	79.0
35–39	5.8	11.1	16.9	21.0	45.1	66.1	26.8	56.2	83.0	2,216	79.6	74.2
40–44	3.2	15.9	19.1	9.3	54.6	63.8	12.5	70.4	82.9	1,478	77.0	68.6
45–49	2.5	18.4	20.8	2.8	45.1	47.9	5.3	63.4	68.7	1,259	69.7	62.2
Residence												
Urban	7.7	7.4	15.1	35.8	24.5	60.4	43.6	31.9	75.4	5,174	80.0	72.4
Rural	7.1	8.8	15.9	25.9	35.0	60.9	33.0	43.7	76.8	6,498	79.3	74.6
Education[4]												
No education	15.5	13.5	29.0	10.1	14.1	24.1	25.6	27.6	53.1	896	45.4	40.2
Primary	6.0	10.7	16.7	22.8	39.4	62.2	28.8	50.1	78.9	4,288	78.8	74.1
Secondary	7.2	5.6	12.8	37.1	27.5	64.7	44.3	33.1	77.4	4,207	83.5	77.7
More than secondary	7.2	5.9	13.1	39.9	24.9	64.7	47.1	30.8	77.9	2,280	83.1	74.5
Wealth quintile												
Lowest	10.1	10.3	20.4	22.8	25.5	48.4	32.9	35.8	68.7	1,993	70.4	65.8
Second	6.2	8.4	14.6	27.9	37.4	65.2	34.1	45.7	79.8	2,107	81.7	78.3
Middle	6.6	8.6	15.2	27.8	35.9	63.8	34.4	44.6	79.0	2,230	80.7	76.1
Fourth	6.7	6.6	13.3	32.8	29.9	62.7	39.5	36.5	75.9	2,626	82.5	75.9
Highest	7.7	7.5	15.2	37.3	24.3	61.6	45.0	31.7	76.8	2,716	80.3	71.0
Total	7.4	8.1	15.5	30.3	30.3	60.7	37.7	38.5	76.2	11,672	79.6	73.7

Note: Numbers in this table correspond to the revised definition of unmet need described by Bradley et al. (2012).
[1] Total demand is the sum of unmet need and met need.
[2] Percentage of demand satisfied is met need divided by total demand.
[3] Modern methods include female sterilisation, male sterilisation, intrauterine devices, injectables, implants, pills, male condoms, female condoms, emergency contraception, Standard Days Method, lactational amenorrhoea method, and other modern methods.
[4] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 7.14.1C Need and demand for family planning among currently married women: County level

Percentage of currently married women age 15–49 with unmet need for family planning, percentage with met need for family planning, total demand for family planning, percentage of the demand for family planning that is satisfied, and percentage of the demand for family planning that is satisfied by modern methods, according to county, Kenya Mini DHS 2025–26

	Unmet need for family planning			Met need for family planning (currently using)			Total demand for family planning[1]					
County	For spacing	For limiting	Total	For spacing	For limiting	Total	For spacing	For limiting	Total	Number of women	Percentage of demand satisfied[2]	Percentage of demand satisfied by modern methods[3]
Mombasa	10.5	12.8	23.4	28.8	18.7	47.5	39.3	31.6	70.8	444	67.0	62.5
Kwale	7.6	11.0	18.6	30.2	15.6	45.8	37.8	26.6	64.3	205	71.1	70.7
Kilifi	7.9	6.1	14.1	42.2	20.9	63.0	50.1	27.0	77.1	302	81.8	74.2
Tana River	11.9	17.4	29.3	13.9	7.9	21.8	25.7	25.4	51.1	88	42.7	42.7
Lamu	11.1	13.9	24.9	20.7	19.3	39.9	31.7	33.1	64.9	27	61.6	61.0
Taita Taveta	3.9	7.2	11.1	30.0	42.1	72.2	33.9	49.4	83.3	85	86.7	78.1
Garissa	9.7	6.7	16.4	4.1	1.3	5.4	13.9	8.0	21.8	90	24.7	22.7
Wajir	15.8	11.9	27.7	0.8	0.0	0.8	16.5	11.9	28.4	72	2.6	2.6
Mandera	24.4	0.4	24.7	0.3	0.0	0.3	24.7	0.4	25.0	77	1.3	1.3
Marsabit	17.8	5.7	23.5	28.8	6.1	35.0	46.6	11.8	58.5	77	59.8	52.0
Isiolo	12.0	12.2	24.2	21.6	13.3	34.9	33.6	25.5	59.1	36	59.0	54.7
Meru	2.8	5.2	8.0	27.9	48.6	76.4	30.7	53.8	84.5	308	90.5	84.6
Tharaka-Nithi	5.5	4.9	10.5	27.1	42.1	69.1	32.6	47.0	79.6	80	86.8	81.5
Embu	3.8	2.8	6.6	22.7	50.7	73.4	26.4	53.5	79.9	129	91.8	87.2
Kitui	5.6	10.3	15.9	19.6	44.7	64.3	25.2	55.0	80.2	251	80.2	78.0
Machakos	6.1	2.2	8.4	32.7	42.3	75.1	38.9	44.6	83.4	420	90.0	81.6
Makueni	2.2	1.6	3.8	36.0	43.5	79.6	38.3	45.1	83.4	192	95.4	85.9
Nyandarua	7.0	8.2	15.2	26.6	42.0	68.6	33.6	50.2	83.8	122	81.8	78.3
Nyeri	4.9	14.4	19.2	20.5	40.9	61.5	25.4	55.3	80.7	104	76.2	75.2
Kirinyaga	3.0	4.0	6.9	29.0	47.8	76.7	31.9	51.7	83.7	141	91.7	85.1
Murang'a	1.8	5.0	6.8	25.9	52.5	78.4	27.6	57.5	85.2	182	92.0	88.7
Kiambu	9.9	8.0	17.9	35.6	26.0	61.6	45.5	34.0	79.6	819	77.4	72.0
Turkana	25.6	8.3	33.9	15.8	5.6	21.4	41.4	13.8	55.2	116	38.7	30.8
West Pokot	21.6	19.9	41.5	9.6	8.7	18.3	31.2	28.6	59.8	145	30.6	29.4
Samburu	14.7	6.7	21.4	25.8	18.8	44.6	40.5	25.5	66.0	58	67.5	56.6
Trans Nzoia	8.6	11.0	19.7	24.7	32.1	56.7	33.3	43.1	76.4	197	74.3	71.1
Uasin Gishu	9.4	10.0	19.4	30.9	32.2	63.1	40.3	42.2	82.5	335	76.5	68.5
Elgeyo-Marakwet	2.8	10.5	13.4	27.7	41.6	69.3	30.5	52.1	82.6	92	83.8	76.6
Nandi	7.9	12.1	20.0	21.9	34.1	55.9	29.8	46.1	75.9	188	73.7	73.1
Baringo	9.3	4.3	13.6	32.5	34.1	66.6	41.8	38.4	80.2	109	83.0	59.5
Laikipia	2.8	13.0	15.8	23.9	38.5	62.5	26.7	51.6	78.3	105	79.9	71.3
Nakuru	5.8	6.7	12.5	32.9	32.5	65.4	38.7	39.2	77.9	614	84.0	79.9
Narok	8.1	10.2	18.3	28.7	31.2	59.8	36.8	41.3	78.1	283	76.6	70.8
Kajiado	4.7	2.1	6.9	36.1	27.2	63.3	40.8	29.4	70.2	261	90.2	76.7
Kericho	6.3	6.8	13.1	28.7	34.1	62.8	35.0	40.9	75.9	198	82.7	77.3
Bomet	9.0	11.4	20.5	23.2	31.9	55.0	32.2	43.3	75.5	242	72.9	68.0
Kakamega	5.4	9.1	14.6	26.1	40.9	67.0	31.6	50.0	81.6	406	82.1	81.0
Vihiga	6.2	12.9	19.1	25.7	37.1	62.8	31.8	50.0	81.9	118	76.7	74.9
Bungoma	10.5	8.8	19.3	25.0	32.9	57.9	35.5	41.7	77.2	349	75.0	72.3
Busia	8.0	12.8	20.8	28.1	30.9	59.1	36.1	43.7	79.8	212	74.0	70.7
Siaya	8.0	14.7	22.6	25.1	26.6	51.6	33.1	41.2	74.3	206	69.5	69.1
Kisumu	6.8	4.8	11.6	34.8	39.2	74.0	41.7	44.0	85.7	296	86.4	77.2
Homa Bay	8.3	4.7	13.1	39.1	29.7	68.9	47.5	34.5	81.9	279	84.1	79.7
Migori	10.0	6.7	16.7	35.5	25.0	60.5	45.4	31.8	77.2	314	78.3	73.9
Kisii	2.3	10.0	12.3	31.4	39.9	71.3	33.6	49.9	83.6	253	85.3	82.2
Nyamira	5.2	6.7	11.9	20.6	48.4	69.0	25.7	55.2	80.9	123	85.3	72.9
Nairobi	5.1	7.7	12.8	38.4	23.2	61.5	43.5	30.8	74.4	1,920	82.8	73.5
Total	7.4	8.1	15.5	30.3	30.3	60.7	37.7	38.5	76.2	11,672	79.6	73.7

Note: Numbers in this table correspond to the revised definition of unmet need described by Bradley et al. (2012).
[1] Total demand is the sum of unmet need and met need.
[2] Percentage of demand satisfied is met need divided by total demand.
[3] Modern methods include female sterilisation, male sterilisation, intrauterine devices, injectables, implants, pills, male condoms, female condoms, emergency contraception, Standard Days Method, lactational amenorrhoea method, and other modern methods.

Table 7.14.2 Need and demand for family planning among all women and sexually active unmarried women

Percentage of all women and sexually active unmarried women age 15–49 with unmet need for family planning, percentage with met need for family planning, total demand for family planning, percentage of the demand for family planning that is satisfied, and percentage of the demand for family planning that is satisfied by modern methods, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Unmet need for family planning			Met need for family planning (currently using)			Total demand for family planning[1]			Number of women	Percentage of demand satisfied[2]	Percentage of demand satisfied by modern methods[3]
	For spacing	For limiting	Total	For spacing	For limiting	Total	For spacing	For limiting	Total			
ALL WOMEN												
Age												
15–19	4.1	0.4	4.5	7.6	0.4	8.0	11.7	0.8	12.5	3,916	63.7	56.8
20–24	8.9	1.1	10.0	36.6	4.2	40.8	45.5	5.3	50.9	3,653	80.3	70.2
25–29	9.1	2.7	11.9	41.7	14.6	56.3	50.9	17.3	68.2	3,732	82.6	77.0
30–34	5.4	5.2	10.6	29.6	27.7	57.3	35.0	32.9	67.9	3,108	84.4	78.8
35–39	4.7	9.0	13.7	17.9	40.8	58.7	22.6	49.8	72.4	2,989	81.1	75.3
40–44	2.7	12.0	14.7	7.5	47.2	54.7	10.3	59.1	69.4	2,048	78.8	70.7
45–49	1.7	14.1	15.9	2.1	37.7	39.8	3.8	51.9	55.7	1,784	71.5	64.3
Residence												
Urban	5.4	4.7	10.1	25.9	17.4	43.3	31.3	22.1	53.3	9,829	81.1	72.3
Rural	6.1	5.5	11.6	20.1	23.9	44.0	26.2	29.4	55.6	11,402	79.2	73.9
Education[4]												
No education	12.9	10.8	23.7	8.2	12.3	20.5	21.1	23.1	44.2	1,161	46.4	41.1
Primary	5.1	8.4	13.5	19.5	34.9	54.4	24.6	43.3	67.9	6,053	80.2	75.7
Secondary	5.2	2.9	8.0	22.3	15.1	37.5	27.5	18.0	45.5	9,849	82.3	75.5
More than secondary	6.1	4.1	10.2	32.7	16.5	49.2	38.8	20.6	59.5	4,167	82.8	71.6
Wealth quintile												
Lowest	7.9	6.7	14.7	17.3	19.1	36.5	25.3	25.9	51.2	3,312	71.3	66.7
Second	5.7	5.1	10.8	21.2	24.7	45.9	26.9	29.8	56.7	3,852	81.0	77.1
Middle	5.6	5.2	10.8	21.4	23.8	45.3	27.0	29.0	56.1	4,133	80.7	75.1
Fourth	5.0	4.4	9.4	25.2	21.7	46.9	30.2	26.1	56.3	4,800	83.3	75.2
Highest	5.2	4.7	9.9	26.3	16.0	42.3	31.5	20.7	52.3	5,134	81.0	70.5
Total	5.8	5.1	10.9	22.8	20.9	43.7	28.5	26.0	54.5	21,231	80.1	73.2
SEXUALLY ACTIVE UNMARRIED WOMEN[5]												
Age												
15–19	44.2	2.8	47.0	45.4	1.8	47.3	89.6	4.6	94.2	120	50.2	36.4
20–24	32.1	1.9	34.0	55.1	4.7	59.8	87.2	6.6	93.7	279	63.8	52.4
25–29	16.0	5.8	21.8	44.7	28.1	72.8	60.7	33.9	94.6	216	77.0	67.0
30–34	8.4	21.9	30.3	25.8	31.6	57.4	34.2	53.5	87.7	138	65.4	64.7
35–39	4.0	15.9	20.0	14.4	50.8	65.2	18.5	66.7	85.2	136	76.5	65.5
40–44	9.7	11.0	20.7	7.7	52.8	60.5	17.4	63.8	81.2	81	74.5	70.8
45–49	0.0	40.2	40.2	1.7	41.2	42.9	1.7	81.4	83.1	49	51.6	47.7
Residence												
Urban	16.0	11.1	27.1	38.0	26.6	64.6	53.9	37.8	91.7	544	70.4	59.0
Rural	24.2	8.6	32.8	33.8	22.5	56.3	58.0	31.1	89.1	475	63.2	57.0
Education[4]												
No education	(32.4)	(8.1)	(40.5)	(4.0)	(21.6)	(25.5)	(36.4)	(29.7)	(66.1)	31	(38.7)	(36.8)
Primary	11.0	12.0	22.9	21.5	43.3	64.8	32.5	55.2	87.7	271	73.9	71.9
Secondary	23.2	7.4	30.6	38.8	22.6	61.4	62.0	29.9	92.0	430	66.7	57.2
More than secondary	21.7	12.2	33.8	49.0	10.7	59.7	70.7	22.9	93.5	287	63.8	48.7
Wealth quintile												
Lowest	24.9	10.6	35.5	26.4	26.5	53.0	51.3	37.1	88.4	135	59.9	58.0
Second	21.1	6.0	27.2	32.6	27.5	60.1	53.7	33.6	87.3	165	68.9	64.6
Middle	22.4	8.3	30.7	36.7	21.8	58.5	59.1	30.1	89.2	203	65.6	58.4
Fourth	17.4	8.9	26.4	37.5	27.2	64.7	54.9	36.1	91.1	266	71.1	59.0
Highest	16.7	14.7	31.3	41.3	21.6	62.9	57.9	36.3	94.2	250	66.7	52.8
Total	19.8	10.0	29.8	36.0	24.7	60.7	55.8	34.7	90.5	1,018	67.1	58.1

Notes: Numbers in this table correspond to the revised definition of unmet need described by Bradley et al. (2012). Figures in parentheses are based on 25–49 unweighted cases.

[1] Total demand is the sum of unmet need and met need.

[2] Percentage of demand satisfied is met need divided by total demand.

[3] Modern methods include female sterilisation, male sterilisation, intrauterine devices, injectables, implants, pills, male condoms, female condoms, emergency contraception, Standard Days Method, lactational amenorrhoea method, and other modern methods.

[4] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

[5] Had sexual intercourse within 30 days preceding the survey

Table 7.14.2C Need and demand for family planning among all women: County level

Percentage of all women age 15–49 with unmet need for family planning, percentage with met need for family planning, total demand for family planning, percentage of the demand for family planning that is satisfied, and percentage of the demand for family planning that is satisfied by modern methods, according to county, Kenya Mini DHS 2025–26

	Unmet need for family planning			Met need for family planning (currently using)			Total demand for family planning[1]					
County	For spacing	For limiting	Total	For spacing	For limiting	Total	For spacing	For limiting	Total	Number of women	Percentage of demand satisfied[2]	Percentage of demand satisfied by modern methods[3]
Kwale	5.7	6.5	12.2	20.4	10.3	30.7	26.1	16.8	42.9	374	71.6	70.7
Kilifi	5.4	3.2	8.6	29.5	12.7	42.3	34.9	15.9	50.9	608	83.1	73.0
Tana River	8.8	11.2	20.0	11.7	5.6	17.4	20.5	16.9	37.4	137	46.4	45.0
Lamu	7.2	8.3	15.5	12.5	12.9	25.5	19.7	21.3	41.0	54	62.1	61.7
Taita Taveta	3.1	4.0	7.1	24.3	28.1	52.4	27.4	32.1	59.5	153	88.0	77.3
Garissa	5.8	3.6	9.4	2.4	1.6	4.1	8.2	5.2	13.5	168	30.1	28.4
Wajir	8.7	6.6	15.3	0.4	0.0	0.4	9.1	6.6	15.7	130	2.6	2.6
Mandera	15.8	0.5	16.2	0.2	0.0	0.2	16.0	0.5	16.5	123	1.3	1.3
Marsabit	9.9	3.1	13.0	18.3	4.1	22.4	28.2	7.2	35.4	142	63.3	55.4
Isiolo	6.9	6.3	13.1	13.5	10.4	23.9	20.4	16.6	37.0	76	64.5	61.2
Meru	2.1	3.0	5.1	20.1	33.5	53.7	22.2	36.6	58.8	530	91.3	86.0
Tharaka-Nithi	3.4	3.5	6.8	22.1	27.6	49.6	25.4	31.0	56.5	147	87.9	81.0
Embu	4.1	1.8	5.9	18.7	33.4	52.1	22.8	35.3	58.0	239	89.8	83.7
Kitui	7.0	6.5	13.5	16.6	31.1	47.7	23.5	37.6	61.2	412	78.0	75.5
Machakos	4.8	1.6	6.4	32.1	31.2	63.3	36.9	32.8	69.7	677	90.8	77.7
Makueni	3.4	0.9	4.3	29.5	31.7	61.3	32.9	32.6	65.5	350	93.5	80.1
Nyandarua	4.3	5.5	9.8	18.2	28.6	46.8	22.5	34.1	56.7	223	82.7	78.7
Nyeri	4.2	7.5	11.6	14.9	25.2	40.1	19.1	32.6	51.8	223	77.5	76.8
Kirinyaga	2.7	2.4	5.1	22.5	34.5	57.0	25.2	36.8	62.1	258	91.9	83.2
Murang'a	2.4	2.6	5.0	17.5	29.4	46.9	19.9	32.0	51.9	377	90.3	86.6
Kiambu	6.3	5.2	11.5	23.3	15.4	38.7	29.6	20.6	50.3	1,689	77.1	71.3
Turkana	17.5	6.1	23.6	14.0	3.6	17.6	31.5	9.7	41.2	178	42.8	34.8
West Pokot	18.9	13.9	32.8	10.0	6.5	16.6	29.0	20.4	49.4	225	33.5	32.0
Samburu	11.3	4.0	15.3	22.9	14.7	37.7	34.2	18.7	52.9	98	71.1	57.3
Trans Nzoia	6.8	6.5	13.2	16.9	20.6	37.5	23.7	27.0	50.7	370	73.9	70.4
Uasin Gishu	7.9	5.8	13.7	23.9	20.9	44.8	31.7	26.7	58.5	667	76.6	68.8
Elgeyo-Marakwet	4.7	6.4	11.1	23.2	26.0	49.2	27.9	32.4	60.3	178	81.6	72.3
Nandi	8.0	8.6	16.6	13.7	22.9	36.6	21.7	31.5	53.2	361	68.7	68.3
Baringo	9.4	2.8	12.2	24.8	24.9	49.7	34.2	27.7	61.9	187	80.3	59.3
Laikipia	2.4	8.4	10.8	17.2	24.7	41.9	19.6	33.0	52.6	195	79.6	70.1
Nakuru	4.6	5.0	9.6	23.0	23.5	46.5	27.6	28.5	56.1	1,078	82.9	78.5
Narok	8.1	7.2	15.3	21.2	23.4	44.7	29.3	30.6	60.0	428	74.5	69.0
Kajiado	4.9	2.2	7.0	27.1	19.4	46.5	31.9	21.6	53.6	467	86.9	72.3
Kericho	5.1	3.7	8.7	24.3	23.1	47.4	29.3	26.8	56.1	387	84.4	78.9
Bomet	6.9	6.3	13.3	17.0	18.7	35.8	24.0	25.1	49.1	480	73.0	63.9
Kakamega	5.0	5.6	10.6	23.4	27.7	51.1	28.4	33.3	61.7	665	82.9	81.3
Vihiga	5.2	6.9	12.0	19.8	22.2	41.9	24.9	29.0	54.0	233	77.7	73.6
Bungoma	8.3	5.1	13.5	17.2	21.9	39.2	25.5	27.1	52.6	625	74.4	71.8
Busia	6.4	7.7	14.0	20.3	20.7	41.0	26.6	28.4	55.0	373	74.5	70.9
Siaya	7.5	8.7	16.2	17.4	20.7	38.2	25.0	29.4	54.4	370	70.2	69.4
Kisumu	5.0	2.9	7.9	30.5	28.1	58.6	35.5	31.0	66.5	551	88.1	80.3
Homa Bay	6.4	3.4	9.8	33.6	21.0	54.7	40.0	24.5	64.5	485	84.8	78.5
Migori	9.1	4.8	13.8	31.3	18.8	50.1	40.3	23.6	63.9	509	78.3	73.5
Kisii	2.0	6.6	8.6	24.0	27.3	51.3	26.0	33.8	59.9	439	85.7	81.5
Nyamira	5.4	3.5	8.9	19.7	30.4	50.1	25.1	33.9	59.0	238	84.9	71.7
Nairobi City	3.9	5.1	9.0	27.0	18.1	45.1	30.9	23.2	54.1	3,550	83.3	73.5
Total	5.8	5.1	10.9	22.8	20.9	43.7	28.5	26.0	54.5	21,231	80.1	73.2

Note: Numbers in this table correspond to the revised definition of unmet need described by Bradley et al. (2012). Too few unweighted cases were available to generate precise estimates for sexually active unmarried women by county, so these values are not shown.

[1] Total demand is the sum of unmet need and met need.

[2] Percentage of demand satisfied is met need divided by total demand.

[3] Modern methods include female sterilisation, male sterilisation, intrauterine devices, injectables, implants, pills, male condoms, female condoms, emergency contraception, Standard Days Method, lactational amenorrhoea method, and other modern methods.

Table 7.15 Future use of contraception

Percent distribution of currently married women age 15–49 who are not using a contraceptive method by intention to use one in the future, according to number of living children, Kenya Mini DHS 2025–26

	Number of living children[1]					
Intention to use in the future	0	1	2	3	4+	Total
Intends to use	35.6	52.3	57.1	47.2	31.5	44.3
Unsure	6.3	5.0	1.9	3.0	2.1	3.1
Does not intend to use	58.1	42.7	41.0	49.8	66.4	52.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Number of women	340	920	1,006	785	1,541	4,593

[1] Includes current pregnancies

Table 7.16 Exposure to family planning messages

Percentage of women age 15–49 who heard or saw family planning messages through specific sources in the past 12 months, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Radio	Television	News-paper/ magazine	Mobile phone	Social media[1]	Poster/ leaflet/ brochure	Outdoor sign or billboard	Com-munity meeting or event	Friends/ peers	Internet	None of these 10 media sources	Number of women
Age												
15–19	35.5	33.2	11.5	9.7	31.5	28.9	22.0	29.0	64.2	27.6	18.6	3,916
20–24	53.4	48.2	17.5	24.5	64.2	45.4	36.8	44.5	80.1	59.3	6.2	3,653
25–29	57.2	55.6	19.7	29.2	65.2	55.7	40.8	52.8	87.3	60.6	4.8	3,732
30–34	61.6	54.7	17.8	28.0	57.4	54.3	42.0	56.5	85.0	55.4	5.3	3,108
35–39	64.1	51.5	15.1	23.8	43.8	52.2	38.3	61.1	84.3	40.2	5.0	2,989
40–44	60.7	49.0	16.6	20.1	38.8	51.7	35.5	58.4	83.0	35.9	6.6	2,048
45–49	64.9	48.6	15.6	19.9	31.1	49.6	35.1	61.8	82.5	30.4	7.1	1,784
Residence												
Urban	50.9	58.3	19.2	26.2	65.2	50.8	40.8	47.0	78.7	61.1	7.9	9,829
Rural	58.8	39.6	13.7	18.7	35.5	44.4	30.8	52.5	81.5	32.5	8.2	11,402
Education[2]												
No education	28.8	12.5	2.9	6.5	13.7	16.2	12.0	44.7	66.6	12.8	24.4	1,161
Primary	60.5	40.8	10.6	18.5	29.2	44.3	29.9	55.8	81.2	25.9	7.3	6,053
Secondary	53.3	48.1	15.7	21.9	51.0	45.7	33.9	45.3	78.1	46.9	8.6	9,849
More than secondary	59.0	69.5	29.5	32.6	84.1	64.6	53.8	54.2	87.7	81.0	3.3	4,167
Wealth quintile												
Lowest	50.9	16.3	8.0	11.6	16.0	34.4	22.1	50.6	76.7	15.0	13.2	3,312
Second	61.3	34.1	11.6	17.2	30.4	42.6	29.6	52.6	82.0	26.8	7.3	3,852
Middle	59.0	51.2	15.8	21.5	45.9	49.3	34.3	51.0	80.6	41.9	7.4	4,133
Fourth	58.0	62.8	20.6	27.6	62.9	52.6	39.6	51.4	82.7	57.8	6.4	4,800
Highest	47.5	63.4	21.4	28.3	74.6	52.9	45.6	45.5	78.5	71.7	7.5	5,134
Total	55.1	48.2	16.3	22.2	49.2	47.4	35.5	50.0	80.2	45.7	8.1	21,231

[1] Social media includes platforms such as Facebook, Twitter, or Instagram.
[2] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 7.16C Exposure to family planning messages: County level

Percentage of women age 15–49 who heard or saw family planning messages through specific sources in the past 12 months, according to county, Kenya Mini DHS 2025–26

County	Radio	Television	News-paper/ magazine	Mobile phone	Social media[1]	Poster/ leaflet/ brochure	Outdoor sign or billboard	Com-munity meeting or event	Friends/ peers	Internet	None of these 10 media sources	Number of women
Mombasa	51.4	53.2	15.9	16.4	62.0	54.2	53.1	41.0	68.3	58.4	10.1	803
Kwale	67.7	40.2	13.4	10.6	35.6	44.5	13.5	69.5	90.3	36.2	4.8	374
Kilifi	50.9	47.2	35.5	43.9	52.3	54.0	45.2	66.9	80.0	51.8	9.6	608
Tana River	53.9	35.8	4.5	2.1	19.7	39.8	33.1	62.9	86.2	34.9	3.7	137
Lamu	26.8	26.9	5.4	13.9	31.0	14.1	3.1	27.1	59.3	30.0	33.7	54
Taita Taveta	57.6	45.9	16.3	15.0	54.2	51.8	41.7	55.0	86.2	53.9	4.2	153
Garissa	19.9	16.6	12.3	14.2	33.3	15.6	16.2	32.1	51.1	30.9	38.6	168
Wajir	0.0	5.3	1.4	1.7	27.4	0.3	0.0	9.9	83.4	22.0	15.3	130
Mandera	1.4	5.8	2.6	3.3	11.5	3.1	2.4	19.3	38.1	8.3	55.9	123
Marsabit	20.8	13.9	3.8	7.3	24.3	13.4	5.0	38.5	72.1	20.5	20.5	142
Isiolo	33.2	33.1	6.3	12.9	32.7	23.7	16.9	37.3	70.2	25.8	19.4	76
Meru	45.3	35.5	7.2	15.7	38.6	43.6	30.0	39.1	77.4	32.2	10.2	530
Tharaka-Nithi	48.5	35.9	15.3	14.0	39.8	38.9	18.4	36.9	69.2	37.8	16.5	147
Embu	47.0	46.6	10.6	28.5	43.7	23.6	17.0	34.0	80.8	47.8	5.6	239
Kitui	59.0	24.9	11.3	18.0	31.9	36.8	47.8	52.5	84.6	29.2	5.6	412
Machakos	62.0	64.0	19.1	28.6	59.5	58.8	46.4	51.6	85.7	43.7	3.7	677
Makueni	63.1	32.0	10.2	15.7	47.4	60.2	20.7	44.1	89.5	34.8	2.4	350
Nyandarua	56.0	63.1	14.0	19.2	45.5	21.9	37.8	55.8	85.9	49.8	9.4	223
Nyeri	53.7	60.1	19.5	28.8	57.2	60.9	37.2	40.7	75.5	56.3	9.8	223
Kirinyaga	42.5	49.3	13.2	18.2	53.7	59.3	29.6	40.9	84.6	54.5	3.4	258
Murang'a	57.9	58.2	15.1	13.7	51.0	56.0	38.6	58.9	93.1	50.3	1.2	377
Kiambu	61.6	73.9	20.6	26.9	74.7	62.4	62.1	56.5	84.9	76.9	4.0	1,689
Turkana	44.3	21.8	9.6	13.9	27.1	30.5	25.4	63.0	76.6	21.6	14.8	178
West Pokot	54.2	21.9	12.9	24.8	22.8	38.3	50.6	36.5	66.1	20.1	14.8	225
Samburu	59.5	29.2	8.9	6.1	27.3	38.8	13.7	58.1	88.7	14.2	5.0	98
Trans Nzoia	61.5	53.6	18.5	14.6	37.5	53.3	30.4	44.0	72.6	39.5	11.0	370
Uasin Gishu	49.5	49.9	18.6	12.5	54.6	36.4	31.1	35.7	81.4	50.9	9.1	667
Elgeyo-Marakwet	70.5	47.3	37.3	32.0	47.2	48.5	47.6	69.5	87.1	48.4	7.2	178
Nandi	54.7	40.9	29.8	31.3	39.7	39.9	37.4	37.3	56.0	38.2	24.0	361
Baringo	67.2	44.2	25.0	10.4	38.8	54.4	44.8	55.3	83.2	34.1	8.9	187
Laikipia	44.1	52.7	13.1	8.1	35.1	57.8	41.7	18.6	53.5	19.8	10.8	195
Nakuru	57.6	55.3	19.1	20.5	49.9	51.9	32.7	55.1	80.6	49.7	6.8	1,078
Narok	73.4	39.9	16.4	29.6	37.8	38.0	16.7	48.4	88.1	35.6	5.0	428
Kajiado	59.1	54.7	19.0	35.1	57.3	42.4	33.6	53.3	90.0	59.0	3.3	467
Kericho	71.8	45.0	24.4	31.1	45.1	64.6	28.3	50.4	89.5	39.8	2.1	387
Bomet	39.5	18.8	10.4	2.5	29.6	33.4	6.9	53.7	95.0	28.6	3.8	480
Kakamega	54.4	36.5	6.2	17.0	33.2	33.5	29.8	36.7	72.6	31.7	9.8	665
Vihiga	58.7	39.2	12.5	19.3	42.5	43.8	33.7	36.3	80.4	32.4	7.1	233
Bungoma	77.6	44.7	19.6	27.6	38.9	48.7	37.3	57.5	89.4	35.5	3.3	625
Busia	56.4	44.8	13.9	21.9	33.9	46.4	14.6	60.5	82.7	31.7	5.2	373
Siaya	79.7	53.0	11.5	20.3	47.1	51.4	59.9	73.6	90.3	50.0	3.4	370
Kisumu	62.3	56.8	15.2	25.6	48.1	61.4	44.3	52.4	79.4	43.9	5.7	551
Homa Bay	81.9	52.4	11.0	17.1	43.0	58.0	53.3	64.0	92.7	28.9	1.0	485
Migori	69.0	63.1	18.4	36.2	46.5	54.9	38.8	78.3	91.5	33.3	1.8	509
Kisii	60.9	38.0	10.5	31.0	43.5	55.9	23.2	77.8	92.8	45.8	3.0	439
Nyamira	76.0	53.8	18.1	19.1	30.1	43.5	35.6	56.4	74.7	28.3	8.5	238
Nairobi City	41.8	49.9	15.9	25.0	62.1	44.5	30.8	41.8	73.0	55.4	11.3	3,550
Total	55.1	48.2	16.3	22.2	49.2	47.4	35.5	50.0	80.2	45.7	8.1	21,231

[1] Social media includes platforms such as Facebook, Twitter, or Instagram.

Table 7.17 Contact of non-users with family planning providers

Among women age 15–49 who are not using contraception, percentage who during the past 12 months were visited by a fieldworker who discussed family planning, percentage who visited a health facility and discussed family planning, percentage who visited a health facility but did not discuss family planning, and percentage who did not discuss family planning either with a fieldworker or at a health facility, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Percentage of women who were visited by a fieldworker who discussed family planning	Percentage of women who visited a health facility in the past 12 months and who:		Percentage of women who did not discuss family planning either with a fieldworker or at a health facility	Number of women
		Discussed family planning	Did not discuss family planning		
Age					
15–19	2.0	5.0	33.0	93.6	3,605
20–24	4.8	18.4	37.9	79.5	2,162
25–29	6.1	27.4	44.1	71.1	1,630
30–34	7.2	28.4	40.0	68.9	1,327
35–39	10.2	19.7	42.5	73.8	1,233
40–44	7.6	15.5	41.9	81.2	928
45–49	6.7	10.8	45.0	85.1	1,074
Residence					
Urban	4.1	14.9	42.0	82.9	5,577
Rural	6.5	16.8	36.2	80.3	6,383
Education[1]					
No education	8.5	14.0	44.8	81.6	923
Primary	7.4	20.0	38.8	76.2	2,760
Secondary	4.1	12.5	35.7	85.5	6,161
More than secondary	5.1	21.3	46.1	76.4	2,116
Wealth quintile					
Lowest	7.4	16.6	34.5	80.2	2,103
Second	5.6	18.5	34.1	79.6	2,083
Middle	5.9	15.8	38.0	80.8	2,262
Fourth	4.9	16.9	39.0	80.3	2,549
Highest	3.7	12.8	46.1	85.2	2,961
Total	5.4	15.9	38.9	81.5	11,959

[1] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 7.17C Contact of non-users with family planning providers: County level

Among women age 15–49 who are not using contraception, percentage who during the past 12 months were visited by a fieldworker who discussed family planning, percentage who visited a health facility and discussed family planning, percentage who visited a health facility but did not discuss family planning, and percentage who did not discuss family planning either with a fieldworker or at a health facility, according to county, Kenya Mini DHS 2025–26

County	Percentage of women who were visited by a fieldworker who discussed family planning	Percentage of women who visited a health facility in the past 12 months and who:		Percentage of women who did not discuss family planning either with a fieldworker or at a health facility	Number of women
		Discussed family planning	Did not discuss family planning		
Mombasa	6.9	16.2	32.0	79.6	537
Kwale	10.1	15.8	50.6	77.9	259
Kilifi	3.4	22.7	27.2	75.7	351
Tana River	13.3	22.5	51.7	69.8	113
Lamu	1.5	15.4	61.3	84.6	40
Taita Taveta	3.8	16.7	46.6	81.0	73
Garissa	9.7	9.5	38.1	86.6	161
Wajir	0.6	5.7	35.0	94.2	130
Mandera	7.7	7.5	8.9	91.1	123
Marsabit	9.4	11.3	57.9	84.9	110
Isiolo	1.0	11.3	27.0	88.3	58
Meru	3.2	4.6	44.3	92.2	246
Tharaka-Nithi	3.9	13.2	51.1	85.1	74
Embu	1.2	11.4	40.7	87.4	114
Kitui	3.7	15.8	50.4	81.9	215
Machakos	5.3	17.1	51.1	79.6	249
Makueni	7.6	16.1	38.9	78.3	136
Nyandarua	2.6	7.0	49.6	90.4	119
Nyeri	5.5	13.4	53.4	82.3	133
Kirinyaga	2.9	13.8	52.6	84.5	111
Murang'a	6.4	15.7	37.4	80.5	200
Kiambu	0.8	12.5	33.1	86.7	1,035
Turkana	31.2	27.3	45.5	60.1	147
West Pokot	3.9	15.8	21.8	82.2	188
Samburu	7.9	9.3	48.4	84.9	61
Trans Nzoia	2.1	8.8	37.5	89.4	232
Uasin Gishu	2.8	17.4	38.8	80.7	368
Elgeyo-Marakwet	12.1	21.7	22.4	77.5	90
Nandi	9.1	21.2	27.3	75.8	229
Baringo	7.1	18.2	44.8	79.6	94
Laikipia	1.0	23.1	38.0	76.4	113
Nakuru	5.2	12.9	32.0	83.9	577
Narok	1.0	34.0	34.0	65.4	237
Kajiado	5.5	20.2	35.8	77.3	250
Kericho	1.5	14.6	52.2	84.8	204
Bomet	0.0	6.1	18.3	93.9	308
Kakamega	4.1	18.6	30.9	80.5	325
Vihiga	11.9	15.4	29.6	78.9	135
Bungoma	4.3	22.1	36.0	75.6	380
Busia	14.0	17.5	46.8	74.4	220
Siaya	7.1	20.3	20.1	78.4	229
Kisumu	8.4	25.8	32.5	72.0	228
Homa Bay	16.6	23.4	34.1	70.4	220
Migori	12.4	31.4	39.9	64.4	254
Kisii	4.8	20.0	28.0	79.8	214
Nyamira	3.3	6.5	27.6	91.1	119
Nairobi City	3.9	12.8	53.1	85.1	1,950
Total	5.4	15.9	38.9	81.5	11,959

INFANT AND CHILD MORTALITY 8

Key Findings

- ***Under-5 mortality:*** In the 5 years before the survey, the under-5 mortality rate was 37 deaths per 1,000 live births.
- ***Neonatal mortality:*** In the 5 years before the survey, the neonatal mortality rate was 19 deaths per 1,000 live births.
- ***Infant mortality:*** The infant mortality rate was 30 deaths per 1,000 live births in the 5 years before the survey.
- ***Trends:*** Between 2003 and 2025–26, the under-5 mortality rate declined from 115 to 37 deaths per 1,000 live births.
- ***High-risk fertility:*** Forty percent of births in the 5 years preceding the survey were in an avoidable high-risk category; 26% were in a single risk category, and 13% were in multiple risk categories.

Information on infant and child mortality is an important indicator of a country's socioeconomic development and its people's quality of life. It can also help identify children who may be at higher risk of death and lead to strategies for reducing this risk, such as promoting birth spacing.

This chapter presents information on levels, trends, and differentials in perinatal, neonatal, infant, and under-5 mortality rates. It also examines biodemographic factors and fertility behaviours that increase mortality risks for infants and children. The information was collected as part of a retrospective pregnancy history in which female respondents listed all of the children to whom they had given birth, along with each child's date of birth, survivorship status, and current age or age at death.

The quality of mortality estimates calculated from pregnancy histories depends on each mother's ability to recall all of the children she has birthed, as well as their birth dates and ages at death. Potential data quality problems include the following:

- The selective omission from pregnancy histories of those births that did not survive, which can result in underestimation of childhood mortality
- The displacement of birth dates, which could distort mortality trends: This can occur if an interviewer knowingly records a birth as occurring in a different year than the one in which it occurred. This could happen if an interviewer is trying to cut down on his or her overall workload, because live births that occurred in the 3 years before the interview are subject to a lengthy set of additional questions.
- The misreporting of age at death: Misreporting the child's age at death may distort the age pattern of mortality, especially if the net effect of the age misreporting is the transfer of deaths from one age bracket to another.

- Misassumptions about female adult mortality: Any method of measuring childhood mortality that relies on mothers' reports (for example, birth histories) assumes that female adult mortality is not high or, if it is high, that there is little or no correlation between the mortality risks of mothers and their children.

Appendix C presents selected indicators of quality for the mortality data on which estimates in this chapter are based (see Tables C.5 and C.6).

8.1 Infant and Child Mortality

Neonatal mortality: The probability of dying within the first month of life

Post-neonatal mortality: The probability of dying between the first month of life and the first birthday (computed as the difference between infant and neonatal mortality)

Infant mortality: The probability of dying between birth and the first birthday

Child mortality: The probability of dying between the first and fifth birthday


Under-5 mortality: The probability of dying between birth and the fifth birthday

During the 5 years preceding the Kenya Mini Demographic and Health Survey (KMDHS), the neonatal mortality rate was 19 deaths per 1,000 live births, the infant mortality rate was 30 deaths per 1,000 live births, and the under-5 mortality rate was 37 deaths per 1,000 live births (**Table 8.1**). Neonatal deaths accounted for two-thirds of all infant deaths.

Trends: Childhood mortality peaked during the 5-year period preceding the 2003 Kenya Demographic and Health Survey (KDHS) and has declined steadily since then, with relatively little change since the 5-year period preceding the 2022 KDHS. The under-5 mortality rate declined from 115 deaths per 1,000 live births in the 5 years before the 2003 KDHS to 37 deaths per 1,000 live births in the 5 years before the 2025–26 KMDHS. Similarly, the infant mortality rate decreased from 77 to 30 deaths per 1,000 live births over the same period, while the neonatal mortality rate declined from 33 to 19 deaths per 1,000 live births (**Figure 8.1**).

Figure 8.1 **Trends in early childhood mortality rates**

Deaths per 1,000 live births in the 5-year period before the survey



Childhood mortality has declined steadily since the 5-year period preceding the 2003 KDHS and has remained relatively stable since the 5-year period preceding the 2022 KDHS. The under-5 mortality rate declined from 115 deaths per 1,000 live births in the 5 years before the 2003 survey to 37 deaths per 1,000 live births in the 5 years before the 2025–26 survey. Over the same period, the infant mortality rate declined from 77 to 30 deaths per 1,000 live births, and the neonatal mortality rate declined from 33 to 19 deaths per 1,000 live births.

Table 8.2 presents early childhood mortality rates according to background characteristics for the 5-year period before the 2025–26 KMDHS. **Table 8.3** presents data on the relationships between child mortality and additional characteristics for the 10-year period preceding the 2025–26 survey. A 10-year reference period was used to improve the robustness of the estimates.

Patterns by background characteristics

- The neonatal mortality rate in the 5 years preceding the 2025–26 KMDHS was higher in rural areas (21 deaths per 1,000 live births) than in urban areas (17 deaths per 1,000 live births) (**Table 8.2**).
- Mortality rates in the 5 years preceding the survey were higher for male children than for female children across all indicators.
- Rural areas had a higher infant mortality rate (31 deaths per 1,000 live births) than urban areas (28 deaths per 1,000 live births).
- First births had a higher under-5 mortality rate in the 10 years preceding the survey (41 deaths per 1,000 live births) than did middle-order births (that is, second- and third-order births) (35 deaths per 1,000 live births) (**Table 8.3**).
- Births with birth intervals of less than 2 years had a higher under-5 mortality rate in the 10 years preceding the survey (41 deaths per 1,000 live births) than did those with birth intervals of 3 years (27 deaths per 1,000 live births) (**Table 8.3**).

8.2 PERINATAL MORTALITY

Perinatal mortality rate

Perinatal deaths comprise stillbirths (pregnancy losses occurring after 28 weeks of gestation) and early neonatal deaths (deaths of live births within the first 7 days of life). The perinatal mortality rate is calculated as the number of perinatal deaths per 1,000 pregnancies lasting 28 or more weeks.

Sample: Number of pregnancies lasting 28 or more weeks among women age 15–49 in the 5 years before the survey

In 2014, the Every Newborn Action Plan, a global multi-partner movement to end preventable maternal and newborn deaths and stillbirths, set a target for a national stillbirth rate of 12 or fewer stillbirths per 1,000 births in all countries by 2030 (World Health Organization and the United Nations Children's Fund 2014).

Perinatal mortality, comprised of both stillbirths and early neonatal deaths, is an important indicator of the quality of antenatal and perinatal care. In the 5 years preceding the 2025–26 KMDHS, 146 recorded stillbirths equate to a rate of 14 stillbirths per 1,000 pregnancies of 28 or more weeks. The 175 early neonatal deaths reported during the same period equate to a rate of 17 early neonatal deaths per 1,000 live births. This results in a perinatal mortality rate of 30 perinatal deaths per 1,000 pregnancies of 28 or more weeks. The perinatal mortality rate is higher in urban areas (33 perinatal deaths per 1,000 pregnancies of 28 or more weeks) than in rural areas (28 perinatal deaths per 1,000 pregnancies of 28 or more weeks) (**Table 8.4**).

Patterns by background characteristics

- The perinatal mortality rate decreases as the interval between the two most recent pregnancies increases, from 39 deaths per 1,000 pregnancies of at least 28 weeks for intervals of less than 15 months to 28 deaths per 1,000 pregnancies of at least 28 weeks for intervals of 39 months or more (**Table 8.4**).
- The perinatal mortality rate decreases with increasing level of maternal education. It is highest (47 deaths per 1,000 pregnancies of at least 28 weeks) among mothers with no education and lowest (23 deaths per 1,000 pregnancies of at least 28 weeks) among mothers with secondary education. The perinatal mortality

rate is 29 deaths per 1,000 pregnancies of at least 28 weeks among mothers with more than secondary education.

- The perinatal mortality rate varies with wealth, ranging from 25 deaths per 1,000 pregnancies of 28 weeks or more in the lowest wealth quintile to 35 deaths per 1,000 such pregnancies in the highest wealth quintile.

8.3 High-risk Fertility Behaviour

The survival of infants and children depends in part on the demographic and biological characteristics of their mothers. Typically, the probability of dying in infancy is much greater among children born to mothers younger than age 18 or older than age 34, children born after a short birth interval (less than 24 months), and children born to mothers of high parity (more than three children).

Table 8.5 presents the percentage distribution of children born in the 5 years preceding the 2025–26 KMDHS by category of elevated mortality risk (along with risk ratios), as well as the percentage distribution of currently married women by category of risk if they were to conceive a child at the time of the survey.

- Thirty-five percent of births in the 5 years preceding the survey were not in any high-risk category (**Table 8.5**).
- Twenty-six percent of births were in an unavoidable high-risk category (that is, first-order births to women age 18–34).
- Forty percent of births were in at least one avoidable high-risk category, including 13% that were in multiple high-risk categories (**Table 8.5**).
- Seventy percent of currently married women would be in an avoidable high-risk category if they were to conceive at the time of the survey; 29% would be in a single high-risk category and 40% in multiple high-risk categories.
- The risk ratio denotes the relationship between risk factors and mortality. The risk of dying is 1.29 times higher for a child in any avoidable high-risk category than it is for a child in no high-risk category.

List of Tables

For more information on infant and child mortality, see the following tables:

- **Table 8.1 Early childhood mortality rates**
- **Table 8.2 Five-year early childhood mortality rates according to background characteristics**
- **Table 8.3 Ten-year early childhood mortality rates according to additional characteristics**
- **Table 8.4 Perinatal mortality**
- **Table 8.5 High-risk fertility behaviour**

Table 8.1 Early childhood mortality rates

Neonatal, post-neonatal, infant, child, and under-5 mortality rates for the 15-year period preceding the survey, by 5-year intervals, Kenya Mini DHS 2025–26

Years preceding the survey	Approximate calendar year	Neonatal mortality (NN)	Post-neonatal mortality (PNN)[1]	Infant mortality ($_1q_0$)	Child mortality ($_4q_1$)	Under-5 mortality ($_5q_0$)
0–4	2021–2025	19 (CI: 15–23)	11 (CI: 8–14)	30 (CI: 25–35)	8 (CI: 5–10)	37 (CI: 31–43)
5–9	2015–2020	16 (CI: 13–20)	12 (CI: 8–15)	28 (CI: 23–33)	10 (CI: 6–14)	38 (CI: 32–44)
10–14	2010–2014	19 (CI: 15–24)	10 (CI: 7–14)	29 (CI: 24–35)	9 (CI: 7–11)	38 (CI: 32–44)

CI = confidence interval
[1] Computed as the difference between the infant and neonatal mortality rates

Table 8.2 Five-year early childhood mortality rates according to background characteristics

Neonatal, post-neonatal, infant, child, and under-5 mortality rates for the 5-year period preceding the survey, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Neonatal mortality (NN)	Postneonatal mortality (PNN)[1]	Infant mortality ($_1q_0$)	Child mortality ($_4q_1$)	Under-5 mortality ($_5q_0$)
Child's sex					
Male	24	12	36	9	45
Female	14	10	24	6	30
Residence					
Urban	17	11	28	9	37
Rural	21	11	31	7	38
Total	19	11	30	8	37

[1] Computed as the difference between the infant and neonatal mortality rates

Table 8.3 Ten-year early childhood mortality rates according to additional characteristics

Neonatal, post-neonatal, infant, child, and under-5 mortality rates for the 10-year period preceding the survey, according to additional characteristics, Kenya Mini DHS 2025–26

Characteristic	Neonatal mortality (NN)	Post-neonatal mortality (PNN)[1]	Infant mortality ($_1q_0$)	Child mortality ($_4q_1$)	Under-5 mortality ($_5q_0$)
Mother's age at birth					
<20	19	12	31	17	47
20–29	18	12	30	7	37
30–39	16	10	26	8	34
40–49	22	10	32	(10)	(41)
Birth order					
1	21	10	31	10	41
2–3	15	12	27	9	35
4–6	19	12	30	8	38
7+	17	11	28	5	33
Previous birth interval[2]					
<2 years	16	15	31	10	41
2 years	16	11	27	6	33
3 years	14	5	19	8	27
4+ years	17	13	31	8	39
Mother's education[3]					
No education	14	8	22	6	28
Primary	19	12	31	12	43
Secondary	19	14	33	8	40
More than secondary	15	6	22	4	25
Wealth quintile					
Lowest	17	12	28	7	35
Second	19	14	33	8	41
Middle	21	8	29	10	39
Fourth	14	16	30	13	43
Highest	18	6	25	6	30

Note: Figures in parentheses are based on 25–49 unweighted cases.
[1] Computed as the difference between the infant and neonatal mortality rates
[2] Excludes first-order births
[3] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 8.4 Perinatal mortality

Number of stillbirths, number of early neonatal deaths, stillbirth rate, early neonatal death rate, perinatal mortality rate, and the ratio of stillbirths to early neonatal deaths for the 5-year period preceding the survey, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Number of stillbirths[1]	Number of early neonatal deaths[2]	Stillbirth rate[3]	Early neonatal death rate[4]	Perinatal mortality rate[5]	Number of pregnancies of 28+ weeks duration[6]	Ratio of stillbirths to early neonatal deaths
Mother's age at birth							
<20	15	23	12	18	30	1,288	0.7
20–29	85	92	14	16	30	5,965	0.9
30–39	44	53	14	18	32	3,065	0.8
40–49	2	6	5	18	23	348	0.3
Previous pregnancy interval in months[7]							
First pregnancy	27	54	9	18	27	2,972	0.5
<15	30	29	20	19	39	1,523	1.0
15–26	35	18	21	11	31	1,717	2.0
27–38	18	17	16	15	31	1,142	1.0
39+	36	57	11	17	28	3,311	0.6
Residence							
Urban	83	70	18	15	33	4,684	1.2
Rural	63	105	11	18	28	5,981	0.6
Mother's education[8]							
No education	34	14	33	14	47	1,018	2.4
Primary	58	58	18	18	35	3,274	1.0
Secondary	28	69	7	16	23	4,249	0.4
More than secondary	27	34	13	16	29	2,124	0.8
Wealth quintile							
Lowest	26	32	11	14	25	2,275	0.8
Second	20	35	10	18	28	1,971	0.6
Middle	23	42	12	22	34	1,912	0.6
Fourth	34	34	14	15	28	2,409	1.0
Highest	44	31	21	15	35	2,099	1.4
Total	146	175	14	17	30	10,665	0.8

Note: Respondents may choose to report the duration of their pregnancy in either weeks or months.
[1] Stillbirths are foetal deaths in pregnancies lasting 28 or more weeks. When pregnancy duration is reported in months, stillbirths are foetal deaths in pregnancies lasting 7 or more months.
[2] Early neonatal deaths are deaths at age 0–6 days among live-born children.
[3] Stillbirth rate is the number of stillbirths divided by the number of pregnancies lasting 28 or more weeks, expressed per 1,000.
[4] Early neonatal death rate is the number of early neonatal deaths divided by the number of live births, expressed per 1,000.
[5] Perinatal mortality rate is the sum of the number of stillbirths and early neonatal deaths divided by the number of pregnancies lasting 28 or more weeks, expressed per 1,000.
[6] Includes pregnancies lasting 7 or more months when duration of pregnancy is reported in months
[7] Previous pregnancy interval is the interval between the date of delivery or termination of the prior pregnancy and the date of conception of the pregnancy of interest. Pregnancy interval categories correspond to birth interval categories of <24 months, 24–35 months, 36–47 months, and 48+ months assuming a pregnancy duration of 9 months.
[8] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 8.5 High-risk fertility behaviour

Percent distribution of children born in the 5 years preceding the survey by category of elevated risk of mortality and the risk ratio, and percent distribution of currently married women by category of risk if they were to conceive a child at the time of the survey, Kenya Mini DHS 2025–26

Risk category	Births in the 5 years preceding the survey		Percentage of currently married women[1]
	Percentage of births	Risk ratio	
No high-risk category	34.6	1.00	25.4[a]
Unavoidable high-risk category			
First-order births between age 18 and age 34	25.9	1.31	5.0
Any avoidable high-risk category	39.5	1.29	69.6
Single high-risk category			
Mother's age <18 only	4.0	1.00	0.1
Mother's age >34 only	3.6	2.06	8.4
Birth interval <24 months only	5.5	1.20	8.4
Birth order >3 only	13.3	1.10	12.4
Subtotal	26.4	1.24	29.3
Multiple high-risk categories			
Age <18 and birth interval <24 months[2]	0.2	(3.28)	0.1
Age >34 and birth interval <24 months	0.2	*	0.2
Age >34 and birth order >3	8.6	1.38	31.3
Age >34 and birth interval <24 months and birth order >3	1.1	1.22	3.2
Birth interval <24 months and birth order >3	2.9	1.41	5.6
Subtotal	13.1	1.39	40.3
Total	100.0	na	100.0
Subtotal by individual avoidable high-risk category			
Mother's age <18	4.3	1.13	0.2
Mother's age >34	13.5	1.53	43.0
Birth interval <24 months	9.9	1.29	17.5
Birth order >3	25.9	1.23	52.4
Number of births/women	10,519	na	11,672

Notes: Risk ratio is the ratio of the proportion dead among births in a specific high-risk category to the proportion dead among births not in any high-risk category. Figures in parentheses are based on 25–49 unweighted cases. An asterisk indicates that a figure is based on fewer than 25 unweighted cases and has been suppressed.
na = not applicable
[1] Women are assigned to risk categories according to the status they would have at the birth of a child if they were to conceive at the time of the survey: current age younger than 17 years and 3 months or older than 34 years and 2 months, latest birth less than 15 months ago, or latest birth being of order 3 or higher.
[2] Includes the category 'age <18 and birth order >3'
[a] Includes sterilised women

MATERNAL AND NEWBORN HEALTH CARE 9

Key Findings

- ***Antenatal care from a skilled provider:*** Nearly all (98%) of women age 15–49 received antenatal care (ANC) from a skilled provider for the most recent live birth or stillbirth in the 2 years before the survey.
- ***Timing and number of antenatal care visits:*** 72% of the women had at least four ANC visits during their most recent pregnancy, including 9% of women who had eight or more visits. Among women who received ANC, the median number of months they were pregnant at the first visit was 4.6.
- ***Components of antenatal care:*** 84% of women who received ANC were asked about vaginal bleeding, 87% were counselled about breastfeeding, and 90% were counselled about a maternal diet.
- ***Delivery care services:*** 91% of live births and stillbirths in the 2 years before the survey were delivered in health facilities, and 92% were assisted by skilled providers.
- ***Postnatal health checks within 48 hours:*** 84% of mothers and 86% of live newborns received their first postnatal health check during the first 2 days after birth.

Health care services during pregnancy, childbirth, and the postdelivery period are important for the survival and well-being of both the mother and the infant. ANC provided by skilled health professionals is essential for ensuring positive health outcomes for pregnant women, adolescent girls, and their babies. According to national guidelines and maternal and newborn health standards for the provision of quality care, ANC offers an opportunity to prevent, detect, and manage conditions that contribute to maternal and newborn morbidity and mortality. Quality intrapartum care further promotes safe and respectful childbirth by providing women with continuous, supportive, and woman-centred care in clinically and psychologically safe environments. In Kenya, about 1.4 million women become pregnant annually, with most experiencing vaginal births. Caesarean section rates (CSRs) have increased notably over the past three decades, and medically indicated caesarean deliveries remain critical in reducing maternal and neonatal complications and mortality.

Routine essential newborn care and timely postnatal care (PNC) within the first 48 hours after delivery are vital for the survival and well-being of both mothers and their newborns, as this period carries the highest risk of complications and mortality. Early postnatal checks facilitate prompt identification and management of danger signs and support breastfeeding, thermal care, and overall maternal and newborn health outcomes. In addition, breast and cervical cancers continue to pose substantial public health challenges among women. Early screening and preventive measures, particularly human papillomavirus (HPV) vaccination, are important strategies for preventing cervical cancer, reducing future disease burden, and lowering the costs of long-term treatment and care.

The first portion of this chapter presents information on ANC providers, number and timing of ANC visits, and various components of care. The middle portion focuses on childbirth and postnatal care including information on place of delivery, assistance during delivery, caesarean deliveries, and postnatal health checks for mothers and newborns. The final portion covers issues that affect women's health, regardless of their maternal status, including whether they have been examined for breast or cervical cancer, HPV vaccination coverage, problems accessing health care, and distance to the nearest health facility.

9.1 Antenatal Care Coverage

9.1.1 Skilled Providers

Antenatal care from a skilled provider

Pregnancy care received from a skilled provider such as a doctor, nurse/midwife, or clinical officer


Sample: Women age 15–49 who had a live birth or stillbirth in the 2 years before the survey

Nearly all (98%) of women age 15–49 who had a live birth or stillbirth in the 2 years before the survey received ANC from a skilled provider at least once during pregnancy for the most recent live birth. Only 2% received no ANC during pregnancy for their most recent live birth (**Table 9.1**).

Trends: The percentage of women who received ANC from a skilled provider at least once for their most recent live birth steadily increased from 87% in the 2003 Kenya Demographic and Health Survey (KDHS) to 98% in the 2022 KDHS and then remained unchanged in the 2025–26 Kenya Mini Demographic and Health Survey (KMDHS) (**Figure 9.1**).

Figure 9.1 **Trends in antenatal care coverage**

Percentage of women age 15–49 who had a live birth in the 2 years before the survey (for the most recent birth)



Patterns by background characteristics

- Ten percent of women with no education received no ANC services for their most recent live birth, compared with 1% of women with at least a primary education (**Table 9.1**).
- The percentage of women who received no ANC services for their most recent live birth is highest in Garissa (23%), Mandera (19%), and Wajir (18%) Counties (**Table 9.1C**).

9.1.2 Timing and Number of Antenatal Care Visits

Seventy-two percent of women age 15–49 had at least four ANC visits during their most recent pregnancy resulting in a live birth in the 2 years preceding the survey, including 10% of women who had eight or more visits (**Table 9.2**). Thirty-six percent of women had their first ANC visit during the first trimester, 53% had their first ANC visit between the fourth and sixth month of pregnancy, and 9% did not have their first ANC visit until the seventh month or later. Those who received ANC were a median of 4.7 months pregnant at their first visit.

Trends: Among women who had a live birth in the 2 years before the survey, the percentage who received ANC during the first trimester increased steadily from 10% in the 2003 KDHS to 36% in the 2025–26 KMDHS. Similarly, the percentage of women who had four or more ANC visits decreased from 49% to 44% between the 2003 and 2008–09 surveys and then increased steadily to 72% in the 2025–26 survey (**Figure 9.1**).

Patterns by background characteristics

- The percentage of women who had four or more ANC visits for their most recent live birth is higher in urban areas (80%) than in rural areas (67%) (**Table 9.2**).
- The percentage of women who had their first ANC visit when they were less than 4 months pregnant increases with level of education, from 24% among those with no education to 52% among those with more than a secondary education.
- At the county level, the percentage of women who had four or more ANC visits for their most recent live birth is highest in Makueni (89%); Mombasa (86%); and Kiambu, Kwale, and Nairobi City (84% each). It is lowest in Mandera (22%), West Pokot (37%), and Garissa (38%) (**Table 9.2C**).

9.2 Components of Antenatal Care

Components of antenatal care

Specific ANC services performed by a health care provider include measuring blood pressure, taking a urine sample, taking a blood sample, listening for the baby's heartbeat, counselling about a maternal diet, counselling about breastfeeding, and asking about vaginal bleeding.

Sample—quality of care indicator: Women age 15–49 who had a live birth or stillbirth in the 2 years before the survey and attended at least one ANC visit.

Sample—population-based indicator: All women age 15–49 who had a live birth or stillbirth in the 2 years before the survey

The ability of ANC providers to effectively identify issues that could adversely affect pregnancy outcomes is dictated in large part by the ANC services offered by the health care provider.

The following interventions and tests are recommended at each ANC contact:

- Measuring blood pressure: Taking a woman's blood pressure at each ANC visit is essential for monitoring for gestational hypertension or pre-eclampsia.
- Conducting urine and blood tests: These tests assess signs of infection or other diseases and conditions that could negatively affect a woman or her baby during or after pregnancy.
- Listening to the baby's heartbeat: This can confirm that the foetus is alive and reassure the mother.
- Counselling on maternal nutrition, specifically on healthy eating during pregnancy and breastfeeding: These counselling messages promote healthy weight gain during pregnancy and can help the pregnant woman initiate breastfeeding early.

- Asking about vaginal bleeding: Light bleeding or spotting is common, especially during the first few months of a pregnancy. Heavy bleeding may be a sign of something more serious; a pregnant woman experiencing heavy bleeding should visit a health care provider.


In the 2025–26 KMDHS, data collected on components of ANC were tabulated in two ways. **Table 9.3.1** presents the percentage of women with a live birth or stillbirth in the 2 years before the survey who reported that they had at least one ANC visit and received specific ANC services. This tabulation is a measure of the quality of the ANC services received. **Table 9.3.2** presents the percentage of all women with a live birth or stillbirth in the past 2 years who received specific ANC services, regardless of whether they reported an ANC visit. This tabulation is a measure of coverage of key ANC interventions among the population of women who need them.

Nearly all women who received ANC for their most recent live birth in the 2 years before the survey had their blood pressure measured (98%), blood sample taken (98%), urine sample taken (96%), and baby's heartbeat checked (98%) (**Table 9.3.1** and **Table 9.3.1C**).

Among women age 15–49 who had a live birth in the 2 years preceding the survey and received ANC for their most recent pregnancy, 84% reported that they were asked about vaginal bleeding, 87% were counselled about breastfeeding, and 89% were counselled about a maternal diet (**Table 9.3.1** and **Figure 9.2**). A similar pattern was found among all women who had a live birth or stillbirth in the 2 years preceding the survey (**Table 9.3.2** and **Figure 9.2**).

***Figure 9.2* Components of antenatal care**

Percentage of women age 15–49 who had a live birth in the 2 years before the survey and received specific antenatal services



Trends: Among women who received ANC for their most recent live birth in the 2 years before the survey, the percentage who had a urine sample taken increased from 49% in the 2003 KDHS to 96% in the 2025–26 KMDHS. During the same period, the percentage who had a blood sample taken increased from 56% to 98%, while the percentage who had their blood pressure measured increased from 81% to 98%.

Patterns by background characteristics

- Among all women age 15–49 who received ANC for their most recent live birth in the 2 years preceding the survey, little variability was found across background characteristics in the percentages who had their blood pressure measured and their baby's heartbeat assessed during ANC, with coverage exceeding 95% in nearly all groups.

- Among all women age 15–49 who received ANC for their most recent live birth in the 2 years preceding the survey, those with no education had much lower levels of coverage for several ANC components when compared with those with more than a secondary education: urine sample taken (84% versus 98%), counselling on breastfeeding (73% versus 91%), and being asked about vaginal bleeding: (71% versus 90%) (**Table 9.3.1**).

- Urban women generally receive more complete ANC than rural women. The urban-rural differences are modest for clinical checks but more noticeable for counselling and assessment. For example, 93% of urban women were counselled about a maternal diet compared with 87% of rural women, and 87% of urban women were asked about vaginal bleeding compared with 81% of rural women (**Table 9.3.2**).

- Women in the lowest wealth quintile have lower levels of coverage than women in the highest quintile for most ANC services, most noticeably counselling on a maternal diet (81% versus 95%), counselling on breastfeeding (79% versus 92%), and being asked about vaginal bleeding (76% versus 89%).

- Relatively little county-level variation exists in receipt of routine clinical ANC services. The percentage of women whose blood pressure was measured ranges from 83% in West Pokot to 100% in several counties. The percentage whose baby's heartbeat was checked ranges from 80% in Mandera to 100% in several counties, with most counties reporting coverage above 95% (**Table 9.3.2C**).

- Greater county-level variation is observed for counselling services and assessment of danger signs. The percentage of women counselled about breastfeeding ranges from 44% in Samburu to 100% in Kilifi, while the percentage asked about vaginal bleeding ranges from 42% in Bomet to 99% in Kilifi (**Table 9.3.2C**).

9.2.1 Deworming and Iron-containing Supplementation During Pregnancy

Pregnant women have higher micronutrient needs and are at risk of micronutrient deficiencies including iron deficiency, which is a primary cause of anaemia. Severe anaemia can place the mother and the baby in danger through increased risk of blood loss during labour and increased risk of preterm delivery, low birth weight, and perinatal mortality (Haider et al. 2013). Providing iron tablets or syrups and/or iron-containing micronutrient supplements to pregnant women can help address maternal anaemia (World Health Organization [WHO] 2016).

Regardless of their ANC status, women age 15–49 with a live birth or stillbirth in the 2 years before the survey were asked if they took iron-containing supplements and deworming medication during their most recent pregnancy. Overall, 89% of these women took some form of iron supplementation, and 27% took deworming medication during their most recent pregnancy. Among women who took some form of iron supplementation, 26% took iron supplements for 180 days or more and 38% took them for 90–179 days (**Table 9.4**).

Patterns by background characteristics

- The percentage of women who took iron-containing supplements during pregnancy for their most recent live birth increases with level of education, from 70% among those with no education to 95% among those with more than a secondary education.

- The percentage of women who took iron-containing supplements during pregnancy is higher in urban areas (91%) than in rural areas (87%).

- By county, the percentage of women who took any iron-containing supplements during pregnancy for their most recent live birth is high in most counties; only Nandi, Mandera, Wajir, and Garissa have coverage below 75%. However, the percentage of women who took iron supplements for more than 180 days is substantially lower. The lowest percentages are in Wajir (<1%), Mandera (<1%), and Meru (4%), while the highest are in Makueni (49%), Migori (42%), and Kilifi (40%) (**Table 9.4C**).

9.2.2 Source of Iron-containing Supplements

Information on sources of iron-containing supplements can increase understanding of the distribution patterns of the supplements.

Among women age 15–49 with a live birth in the 2 years preceding the survey who took some form of iron-containing supplementation during their most recent pregnancy, 76% received or purchased the supplements from a public-sector facility, with 36% obtaining them from a government hospital. Twenty-five percent of women received their supplements from the private sector (**Table 9.5**).

9.3 Protection against Neonatal Tetanus

Protection against neonatal tetanus

The number of tetanus toxoid injections needed to protect a baby from neonatal tetanus depends on the mother's vaccinations. A birth is protected against neonatal tetanus if the mother has received any of the following:

- Two tetanus toxoid injections during the pregnancy
- Two or more injections, the last one within 3 years of the birth
- Three or more injections, the last one within 5 years of the birth
- Four or more injections, the last one within 10 years of the birth
- Five or more injections at any time prior to the birth

Sample: Women age 15–49 with a live birth in the 2 years before the survey

Tetanus toxoid injections are given during pregnancy to prevent neonatal tetanus, a major cause of early infant death in many countries. Neonatal tetanus is often caused by failure to implement hygienic procedures during delivery.

Forty-four percent of women age 15–49 received two or more tetanus injections for their most recent live birth in the 2 years before the survey. Overall, 79% of women with a live birth in the 2 years before the survey received sufficient tetanus toxoid injections to protect their babies against neonatal tetanus (**Table 9.6**).

Trends: The percentage of live births protected against neonatal tetanus increased from 70% in the 2008–09 KDHS to 74% in the 2014 KDHS and then further to 75% in the 2022 KDHS and 79% in the 2025–26 KMDHS.

Patterns by background characteristics

- The percentage of women whose most recent live birth was protected from tetanus increases with maternal level of education, from 64% among women with no education to 89% among those with more than a secondary education.
- The percentage of women whose most recent live birth was protected against neonatal tetanus is slightly higher in urban areas (82%) than in rural areas (76%).
- The percentage of women whose most recent live birth was protected against neonatal tetanus is highest in Kilifi, Makueni, and Murang'a (95% each); Vihiga (94%); and Kajiado (93%) Counties. It is lowest in Garissa (50%), Siaya (53%), Baringo (58%), and Isiolo (59%) Counties (**Table 9.6C**).

9.4 DELIVERY SERVICES

9.4.1 Institutional Deliveries


Institutional deliveries
Deliveries that occur in a health facility.
Sample: All live births and/or stillbirths in the 2 years before the survey

Overall, 91% of live births in the 2 years before the survey occurred in a health facility and 9% occurred at home. Sixty-six percent of the live births occurred in a public-sector health facility, 22% occurred in the non-governmental organisation private medical sector, and 3% occurred in other private medical-sector facilities (**Table 9.7**).

Trends: The percentage of live births that occurred in a health facility more than doubled from 39% in the 2003 KDHS to 88% in the 2022 KDHS, increasing further to 91% in the 2025–26 KMDHS (**Figure 9.3**).

Figure 9.3 **Trends in health facility births**

Percentage of live births in the 2 years before the survey by place of delivery




Patterns by background characteristics

- Delivery occurred in a health facility for 95% of births to women who had four or more ANC visits compared with only 41% of births to women who had no ANC visits (**Table 9.7**). Fifty-eight percent of the births to women with no ANC visits occurred at home.
- In the 2 years before the survey, 96% of first-order births were delivered in a health facility, compared with 67% of sixth- or higher-order births.
- Ninety-nine percent of live births in urban areas occurred in a health facility compared with 86% of live births in rural areas.
- The percentage of deliveries in a health facility increases with maternal level of education, from 56% for mothers with no education to 100% for mothers with more than a secondary education.
- The percentage of deliveries occurring in a health facility ranges from 47% in Garissa County to 100% in Nairobi City, Nyandarua, and Nyeri Counties (**Table 9.7C** and **Map 9.1**).

Map 9.1 Health facility births by county

Percentage of live births in the 2 years before the survey that were delivered in a health facility



9.4.2 Delivery by Caesarean Section

A caesarean section is a surgical procedure in which a foetus is delivered through incisions made in the abdomen and uterus. Over the past three decades, CSRs have increased globally, raising public health concerns due to associated maternal and neonatal risks, inequities in access, and cost implications. When performed for appropriate medical indications, caesarean sections can substantially reduce maternal and neonatal morbidity and mortality. The CSR is defined as the number of caesarean deliveries expressed as a percentage of total live births. The internationally recommended optimal CSR is between 10% and 15%; rates below 5% may indicate inadequate access to emergency obstetric care, while rates above 15% suggest possible overuse for non-medical reasons. Evidence indicates that a CSR of up to around 10% is associated with reductions in maternal and neonatal mortality, but a rate beyond this threshold provides no additional health benefits (WHO 2015). The WHO further recommends that caesarean sections be performed only when medically necessary, and supports the use of the Robson Classification System for monitoring and evaluating CSRs (WHO 2017c).

Twenty percent of live births in the 2 years before the survey were delivered by caesarean section (**Table 9.8**).


Trends: The percentage of births delivered via caesarean section have increased five-fold since 2003, from 4% in the 2003 KDHS to 17% in the 2022 KDHS and further to 20% in the 2025–26 KMDHS.

Patterns by background characteristics

- The percentage of caesarean deliveries is twice as high in urban areas (29%) as in rural areas (14%).
- Forty-one percent of live births to women in the highest wealth quintile were delivered by caesarean section compared with 9% of live births to women in the lowest wealth quintile.
- The percentage of live births delivered by caesarean section decreases with increasing birth order, from 24% among first-order births to 7% among sixth- or higher-order births. It also increases with the number of ANC visits, from 2% among women with no ANC visits to 22% among women with four or more visits.
- In 21 counties, CSRs exceed the recommended upper limit of 15%. The highest rates are in Kiambu (39%), Nyeri (37%), Murang'a (36%), Nairobi City (31%), and Makueni (30%). In contrast, the lowest CSRs are in Kwale (9%), Tana River (9%), Nakuru (8%), Garissa (8%), Kilifi (7%), Trans Nzoia (6%), Wajir (5%), and Mandera (2%) (**Table 9.8C** and **Map 9.2**).

Map 9.2 **Caesarean section rates by county**

Percentage of live births in the 2 years before the survey that were delivered via caesarean section



9.4.3 Skilled Assistance during Delivery

Skilled assistance during delivery

Births delivered with the assistance of doctors, nurse/midwives, or registered clinical officers.

Sample: All live births and/or stillbirths in the 2 years before the survey

According to the Kenya National Guidelines on Quality Obstetric and Perinatal Care (Ministry of Health [MoH] 2024b), a skilled birth attendant is a health worker such as a doctor, nurse, midwife, or registered clinical officer with specialisation in reproductive health. The skills, attitudes, and behaviours of skilled birth attendants within an enabling work environment impact the quality of care provided and facilitate favourable outcomes for both mothers and their newborns (MoH 2024b; MoH 2022b).


Obstetric care from a health professional during delivery is recognised as a critical element in managing complications that may arise during childbirth and in reducing maternal and neonatal mortality. Newborns without complications should be kept in skin-to-skin contact with their mothers during the first hour after birth to prevent hypothermia and promote breastfeeding.

Overall, 92% of live births that occurred in the 2 years before the survey were assisted by a skilled provider. More than half of the births were assisted by a doctor (54%), while 38% were assisted by a nurse, midwife, or clinical officer (**Table 9.9** and **Figure 9.4**).

Among most recent live births that occurred in the 2 years before the survey, 70% had skin-to-skin contact immediately after birth.

Figure 9.4 **Assistance during delivery**


Percent distribution of births in the 2 years before the survey



Trends: The percentage of live births assisted by a skilled provider increased from 41% in the 2003 KDHS to 89% in the 2022 KDHS and further to 92% in the 2025–26 KMDHS (**Figure 9.5**).

Figure 9.5 **Trends in delivery assistance**

Percentage of live births in the 2 years before the survey by type of assistance




Patterns by background characteristics

- Ninety-nine percent of live births in urban areas are delivered by a skilled provider compared with 87% of live births in rural areas.
- The percentage of live births delivered by a skilled provider increases with wealth, from 72% in the lowest wealth quintile to more than 99% in the highest wealth quintile (**Table 9.9**).
- The percentage of live births with immediate skin-to-skin contact declines with increasing birth order, from 73% among first births to 63% among births of order six and above (**Table 9.9**).

- At the county level, the percentages of live births delivered by a skilled provider are lowest in Garissa (49%), Wajir (56%), Turkana (56%), and Mandera (59%) and highest in Nairobi City, Nyandarua, and Nyeri (100% each) (**Table 9.9C** and **Map 9.3**).

Map 9.3 Skilled delivery assistance by county

Percentage of live births in the 2 years before the survey that were delivered by a skilled provider



Duration of Stay at Health Facility

Following an uncomplicated vaginal birth in a health facility, the Kenya National Guidelines on Quality Obstetric and Perinatal Care (MoH 2024b) recommend that healthy mothers and newborns remain in the facility for at least 24 hours after delivery to allow for adequate monitoring and PNC. For women who undergo caesarean section, discharge may be considered once the mother is clinically stable and can adequately care for herself and her newborn. In many cases, patients are ready for discharge by the second or third postoperative day.

Sixty percent of women who had a vaginal live birth in a health facility in the 2 years preceding the survey reported a stay of 1–2 days after their most recent birth (in line with the recommendation for an uncomplicated vaginal birth), while 6% stayed for less than 6 hours and 22% for more than 3 days. Eighty-seven percent of women who delivered by caesarean section stayed in a health facility for 3 or more days, as recommended, while 2% stayed for less than 6 hours and 10% for 1–2 days (**Table 9.10**).

9.5 POSTNATAL CARE

9.5.1 Postnatal Health Check for Mothers

A substantial proportion of maternal and neonatal deaths occur within the first 48 hours after delivery, making timely postnatal care essential for both mothers and their newborns (WHO 2022). Prompt postnatal care supports the early identification and management of complications related to childbirth while providing mothers with critical information on self-care, breastfeeding, newborn care, and danger signs. The national guidelines recommend that all women and newborns receive postnatal care within the first 2 days after birth to improve maternal and neonatal survival outcomes (WHO 2022; MoH 2024b).

Vaginal bleeding, uterine contractions, fundal height, temperature, and heart rate (pulse) should be routinely assessed in all postpartum women for the first 24 hours, starting from the first hour after birth. Blood pressure should be measured shortly after birth. If normal, a second blood pressure measurement should be taken within 6 hours. Urine void should be documented within 6 hours.

Among women age 15–49 with a live birth or stillbirth in the 2 years before the survey, 85% with a live birth and 74% with a stillbirth had a postnatal check during the first 2 days after birth. However, 12% of mothers did not receive a postnatal check during the first 2 days after birth (**Table 9.11**).

Trends: The percentage of women with a live birth who had a postnatal check during the first 2 days after birth increased from 31% in the 2008–09 KDHS to 78% in the 2022 KDHS and further to 85% in the 2025–26 KMDHS.

Patterns by background characteristics

- Ninety percent of women with a live birth who delivered in a health facility received a postnatal check during the first 2 days after birth, compared with 33% of those who delivered elsewhere (**Table 9.11**).
- The percentage of women with a live birth who received a postnatal check during the first 2 days after birth increases with wealth, from 70% in the lowest wealth quintile to 93% in the highest wealth quintile.
- The percentage of women with a live birth who received a postnatal check during the first 2 days after birth is highest in Migori (97%), Nyeri (97%), Murang'a (96%), Makueni (96%), and Nairobi City (95%) Counties and lowest in Wajir (51%), Mandera (41%), and Garissa (33%) Counties (**Table 9.11C**).

Type of Provider for Mothers

Eighty-three percent of women with a live birth in the 2 years preceding the survey received their first postnatal health check during the 2 days after the most recent live birth from a skilled provider (doctor, nurse, midwife, or clinical officer), while 15% did not receive a postnatal check during the first 2 days after birth (**Table 9.12**).

At the county level, the percentage of women who received their first postnatal health check during the 2 days after the most recent live birth from a skilled provider is highest in Migori (97%), Nyeri (97%), Murang'a (97%), and Makueni (96%). The percentage is lowest in Wajir (44%), Mandera (40%), and Garissa (33%) (**Table 9.12C**).

Content of Care for Mothers

Seventy-eight percent of women with a live birth in the 2 years before the survey had their blood pressure checked, 76% were informed about vaginal bleeding, and 68% were counselled about family planning by a health care provider during the first 2 days after birth.

Overall, 61% of women had all three checks performed by a health care provider in the first 2 days after a live birth (**Table 9.13**).

Patterns by background characteristics

- The percentage of women with a live birth in the 2 years before the survey who received all three checks increases with maternal level of education, from 31% among women with no education to 74% among those with more than a secondary education.
- The percentage of women with a live birth in the 2 years before the survey who received all three checks generally increases with wealth, from 45% in the lowest wealth quintile to 69% in the fourth wealth quintile and 68% in the highest wealth quintile.
- Counties with the highest percentage of women who had all three checks in the first 2 days after birth are Kilifi (91%), Murang'a (88%), Mombasa (83%), Makueni (83%), and Kiambu (81%). Those with the lowest percentage are Garissa (5%), Mandera (13%), West Pokot (20%), and Samburu (20%) (**Table 9.13C**).

9.5.2 Postnatal Health Check for Newborns

Eighty-six percent of newborns born in the 2 years preceding the survey received a postnatal check during the first 2 days after birth. However, 13% of newborns did not receive any postnatal check (**Table 9.14**).

Patterns by background characteristics

- Ninety percent of newborns delivered in a health facility received a postnatal check after birth, compared with 39% of newborns delivered outside a health facility.
- The percentage of newborns with a postnatal check within the first 2 days after birth increases with maternal level of education, ranging from 60% among women with no education to 90% among those with more than a secondary education (**Table 9.14**).
- The percentage of newborns who received a postnatal check during the first 2 days after birth ranges from 31% in Garissa County to 98% in Makueni County (**Table 9.14C**).

Type of Provider for Newborns

Eighty-four percent of newborns born in the 2 years before the survey received their first postnatal health check during the 2 days after birth from a skilled provider (doctor, nurse, midwife, or clinical officer). Two percent of newborns received their first postnatal check from a traditional health attendant (**Table 9.15**).

The percentage of newborns who received their first postnatal health check during the 2 days after birth from a traditional birth attendant is highest in Wajir (18%), Turkana (11%), Narok (11%), and Tana River (10%) Counties (**Table 9.15C**).

Content of Care for Newborns

Breastfeeding counselling provides timely guidance and support to mothers during the early postnatal period to facilitate exclusive breastfeeding practices. Face-to-face counselling is particularly effective, as it enables direct observation of the infant's positioning and latch, allowing health care providers to offer individualised, practical support that improves breastfeeding technique and outcomes (WHO 2018).

The five signal functions performed during the first 2 days after birth refer to essential postnatal newborn care interventions that indicate whether a health facility is providing basic, life-saving newborn care during the immediate postnatal period. They are used to assess the availability and quality of essential newborn care services within 48 hours after birth, which is the critical window in which most newborn deaths occur.

Results of the 2025–26 KMDHS show that weighing the newborn (91%) is the most commonly performed intervention, while counselling mothers on recognising newborn danger signs and knowing when to seek immediate medical attention is the least reported (75%). In addition, 85% of newborns had their umbilical cord examined, 82% had their temperature measured, 85% received support for breastfeeding, and 84% were observed while breastfeeding (**Table 9.16**).

Sixty-nine percent of newborns had all five signal functions performed during the first 2 days after birth.

Patterns by background characteristics

- The percentage of newborns with the five signal functions performed during the first 2 days after birth is higher in urban areas (78%) than in rural areas (62%).
- The percentage of newborns with the five signal functions performed during the first 2 days after birth increases with maternal level of education, from 32% for mothers with no education to 81% for mothers with more than a secondary education.
- The percentage of newborns with the five signal functions performed during the first 2 days after birth is highest in Kilifi (93%), Mombasa (91%), and Makueni (91%) Counties. It is lowest in Garissa (5%), West Pokot (16%), and Mandera (17%) Counties (**Table 9.16C**).

9.5.3 Postnatal Health Checks for Mothers and Newborns

Overall, both the mother and her newborn received a postnatal health check during the first 2 days after delivery for 81% of the most recent live births in the 2 years before the survey. For 11% of the live births, neither the mother nor her newborn received a postnatal check (**Table 9.17**).


Patterns by background characteristics

- The mother received a postnatal check during the first 2 days after delivery for 90% of births that occurred in a health facility, compared with 33% of births that occurred outside a health facility. A similar pattern was observed for the newborns; 90% of newborns who were delivered in a health facility received a postnatal check during the first 2 days after delivery, compared with 39% who were delivered outside a health facility (**Table 9.17** and **Figure 9.6**).

- The percentage of most recent live births in the 2 years preceding the survey for which both the mother and the newborn received a postnatal check within 2 days of delivery is higher in urban areas (85%) than in rural areas (78%) (**Table 9.17**).

Figure 9.6 **Postnatal health checks for mothers and newborns**

Percentage of most recent live births in the 2 years before the survey for which both the mother and the newborn received a postnatal check during the first 2 days after birth by place of delivery



- The percentage of most recent live births in the 2 years before the survey for which both the mother and the newborn received a postnatal check during the first 2 days after delivery is highest in Nyeri (97%), Makueni (96%), Migori (95%), and Murang'a (94%) Counties and lowest in Wajir (50%), Mandera (35%), and Garissa (30%) Counties (**Table 9.17C**).

9.6 Breast and Cervical Cancer Examinations

Breast cancer examination

Women were asked if a doctor or other health care provider examined their breasts to check for cancer. The examination could be either a clinical breast exam, in which the health care provider feels for lumps or other changes, or the use of medical equipment to image breast tissue, such as a mammogram.

Cervical cancer examination

To check for cervical cancer, a health care worker uses a brush or swab to collect a sample from a woman's cervix. The sample is then sent to a laboratory for testing. This test is called a Pap smear or HPV test. Another method is visual inspection with acetic acid, in which a health care worker puts vinegar on the cervix and checks for a reaction. Women were asked if a doctor or other health care provider ever tested them for cervical cancer. Information on the type of screening test was not collected.

Sample: Women age 15–49

Early detection of cancer is among the most effective strategies for improving treatment outcomes and survival. Detecting cancer at an early stage enables timely initiation of treatment, slows disease progression, and increases the chances of successful management and recovery.

Breast cancer screening involves examining the breast tissue to identify any abnormal lumps or growths. This examination may be conducted by a health care professional through clinical breast examination/imaging, or by individuals through breast self-examination. When an abnormality is detected, a tissue sample may be surgically removed and analysed in a laboratory to confirm the presence of cancer cells.

Cervical cancer screening involves identifying abnormal changes in the cervix using methods such as visual inspection or laboratory-based tests. If precancerous or abnormal lesions are detected, the affected tissue is appropriately treated or removed to prevent progression to cervical cancer.

Sixteen percent of women age 15–49 were ever examined by a doctor or health care worker for breast cancer, while 19% were tested for cervical cancer (**Table 9.18**).


Patterns by background characteristics

- Women in urban areas are more likely to be tested for both breast and cervical cancers (19% and 21%, respectively) than are women in rural areas (13% and 17%, respectively) (**Figure 9.7**).

- The percentage of women examined for breast cancer increases with wealth, from 9% among women in the lowest wealth quintile to 21% among those in the highest wealth quintile. The percentage of women undergoing cervical cancer testing also increases with wealth, from 11% among women in the lowest quintile to 24% among those in the highest quintile.

Figure 9.7 **Breast and cervical cancer exams by residence**

Percentage of women age 15–19 who were ever examined by a health care worker for breast or cervical cancer




- Counties with the lowest percentages of women age 15–49 ever examined for breast cancer are West Pokot (4%), Tana River (3%), Marsabit (3%), Garissa (2%), Wajir (<1%), and Mandera (<1%) (**Table 9.18C**).

- The percentages ever screened for cervical cancer are also lowest in Tana River (3%), Garissa (3%), West Pokot (3%), Wajir (1%), and Mandera (<1%) (**Map 9.4**).

Map 9.4 Cervical cancer screening by county

Percentage of women age 15–49 who have ever been screened for cervical cancer



9.7 Problems in Accessing Health Care

Problems in accessing health care

Women were asked whether each of the following factors is a serious problem in seeking medical advice or treatment for themselves when sick:

- Getting permission to go to the doctor
- Getting money for advice or treatment
- Distance to a health facility
- Not wanting to go alone

Sample: Women age 15–49

Fifty-seven percent of women age 15–49 have at least one serious problem accessing health care for themselves when they are sick. The two most common problems are obtaining money for treatment (50%) and distance to a health facility (23%) (**Table 9.19**).

Patterns by background characteristics

- The percentage of women who experience at least one serious problem accessing health care for themselves when sick increases with their number of living children, from 50% among those with no children to 75% among those with five or more children.
- The percentage of women who experience at least one serious problem accessing health care for themselves when sick decreases with increasing wealth, from 81% among those in the lowest wealth quintile to 34% among those in the highest wealth quintile.
- The highest percentages of women who experience at least one serious problem accessing health care for themselves when sick are in Lamu (85%), West Pokot (81%), Wajir (78%), Homabay (79%), and Trans Nzoia (77%) Counties. The lowest percentages are in Machakos (38%), Murang'a (37%), Kiambu (37%), Kericho (30%), and Elgeyo-Marakwet (25%) Counties (**Table 9.19C**).

9.8 Distance and Means of Transport to the Nearest Health Facility

Fifty-nine percent of women age 15–49 report that the travel time from their home to the nearest health facility is less than 30 minutes. Four percent report a travel time of 2 hours or more. Nearly 8 in 10 women (77%) use non-motorised transportation to travel to the nearest health facility (**Table 9.20**).

Turkana (26%), Kitui (24%), Baringo (20%), Samburu (17%), and Kisii (11%) Counties have the highest percentages of women who report a travel time of 2 hours or more to the nearest health facility (**Table 9.20C**).

9.9 Vaccination against Human Papillomavirus

Vaccination against human papillomavirus

HPV is a common sexually transmitted infection that can cause cervical cancer. The HPV vaccine is a preventive measure, offering protection against the strains most often linked to cervical cancer. In Kenya, girls age 10–14 are routinely offered the vaccine, administered as an injection in the upper arm.

Women age 15–49 were asked whether they had ever received a vaccination against HPV. Those who reported any HPV vaccination were additionally asked about the number of doses of HPV vaccine received and the source of vaccination.

Sample: Women age 15–19 and women age 20–49

Overall, 59% of women age 15–19 have received at least one dose of the HPV vaccine, and 48% have ever received an HPV vaccination card (**Table 9.21.1**). Twenty percent of women age 15–19 have received the first dose of the HPV vaccine, 39% have received two doses, and 41% have not received any HPV vaccination.

Eight percent of women age 20–49 have received at least one dose of the HPV vaccine; 4% have received one dose, and 4% have received two doses. Ninety-two percent have not received any HPV vaccination (**Table 9.21.2**).

Patterns by background characteristics

- The percentage of women age 15–19 who have received any HPV vaccination declines with age, from 75% among women age 15 to 35% among women age 19 (**Table 9.21.1**).

- The percentage of women age 15–19 who have received any HPV vaccination is higher in rural areas (63%) than in urban areas (51%) (**Table 9.21.1**).

- The percentage of women age 15–19 who have received at least one dose of the HPV vaccine is highest in Kitui (88%), Kilifi (83%), Vihiga (81%), and Kericho (80%) Counties and lowest in Mandera (1%), Garissa (8%), and Wajir (9%) Counties (**Table 9.21.1C**).

- The percentage of women age 20–49 who have received at least one dose of the HPV vaccine is highest in Baringo (32%) and Kilifi (31%) Counties and lowest in Mandera County (<1%) (**Table 9.21.2C**).

Source of Human Papillomavirus Vaccination

Among women age 15–49 who have received any HPV vaccination, 86% received their most recent HPV vaccination at school and 12% from a public health facility (**Table 9.22**).

Patterns by background characteristics

- Among women age 20–49 who have received any HPV vaccination, the percentage vaccinated at school decreases with increasing age, from 92% among those age 20–24 to 17% among those age 40–44.

- Women in rural areas (88%) are more likely than women in urban areas (83%) to have been vaccinated at school.

- HPV vaccination through schools is most common in Tan River (99%), Lamu (98%), Isiolo (97%), and Kwale (97%) Counties, while immunisation through public health facilities is most common in Taita Taveta (29%), Kericho (23%), and Makueni (20%) Counties (**Table 9.22C**).

List of Tables

For more information on maternal and newborn health care, see the following tables:

- **Table 9.1 Antenatal care**
- **Table 9.1C Antenatal care: County level**
- **Table 9.2 Number of antenatal care visits and timing of first visit**
- **Table 9.2C Number of antenatal care visits and timing of first visit: County level**
- **Table 9.3.1 Components of antenatal care among women receiving antenatal care**
- **Table 9.3.1C Components of antenatal care among women receiving antenatal care: County level**
- **Table 9.3.2 Components of antenatal care among all women**
- **Table 9.3.2C Components of antenatal care among all women: County level**
- **Table 9.4 Deworming and iron-containing supplementation during pregnancy**
- **Table 9.4C Deworming and iron-containing supplementation during pregnancy: County level**
- **Table 9.5 Source of iron-containing supplements**
- **Table 9.6 Tetanus toxoid injections**
- **Table 9.6C Tetanus toxoid injections: County level**
- **Table 9.7 Place of delivery**
- **Table 9.7C Place of delivery: County level**
- **Table 9.8 Caesarean section**
- **Table 9.8C Caesarean section: County level**
- **Table 9.9 Assistance during delivery**
- **Table 9.9C Assistance during delivery: County level**
- **Table 9.10 Duration of stay in a health facility after birth**
- **Table 9.11 Timing of first postnatal check for the mother**
- **Table 9.11C Timing of first postnatal check for the mother: County level**
- **Table 9.12 Type of provider of first postnatal check for the mother**
- **Table 9.12C Type of provider of first postnatal check for the mother: County level**
- **Table 9.13 Content of postnatal care for the mother**
- **Table 9.13C Content of postnatal care for the mother: County level**
- **Table 9.14 Timing of first postnatal check for the newborn**
- **Table 9.14C Timing of first postnatal check for the newborn: County level**
- **Table 9.15 Type of provider of first postnatal check for the newborn**
- **Table 9.15C Type of provider of first postnatal check for the newborn: County level**
- **Table 9.16 Content of postnatal care for the newborn**
- **Table 9.16C Content of postnatal care for the newborn: County level**
- **Table 9.17 Postnatal checks for mothers and newborns**
- **Table 9.17C Postnatal checks for mothers and newborns: County level**
- **Table 9.18 Examinations for breast and cervical cancers**
- **Table 9.18C Examinations for breast and cervical cancers: County level**
- **Table 9.19 Problems in accessing health care**
- **Table 9.19C Problems in accessing health care: County level**
- **Table 9.20 Distance from a health care facility**
- **Table 9.20C Distance from a health care facility: County level**
- **Table 9.21.1 Human papillomavirus vaccination coverage among women age 15–19**
- **Table 9.21.1C Human papillomavirus vaccination coverage among women age 15–19: County level**
- **Table 9.21.2 Human papillomavirus vaccination coverage among women age 20–49**
- **Table 9.21.2C Human papillomavirus vaccination coverage among women age 20–49: County level**
- **Table 9.22 Source of human papillomavirus vaccinations**
- **Table 9.22C Source of human papillomavirus vaccinations: County level**

Table 9.1 Antenatal care

Percent distribution of women age 15–49 who had a live birth and/or stillbirth in the 2 years preceding the survey by antenatal care (ANC) provider during pregnancy for the most recent birth or stillbirth, and percentage receiving ANC from a skilled provider for the most recent birth or stillbirth, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	ANC provider				No ANC	Total	Percentage receiving ANC from a skilled provider[1]	Number of women
	Doctor	Nurse/ midwife/ clinical officer	Community health worker/ fieldworker	Traditional birth attendant				
				LIVE BIRTHS				
Age at birth								
<20	43.5	54.1	0.0	0.1	2.3	100.0	97.6	492
20–34	48.3	49.9	0.0	0.0	1.8	100.0	98.2	2,943
35–49	50.5	46.2	0.1	0.0	3.2	100.0	96.7	548
Birth order[2]								
1	49.9	48.5	0.0	0.0	1.6	100.0	98.4	1,269
2–3	49.9	48.6	0.0	0.0	1.5	100.0	98.5	1,724
4–5	46.3	52.2	0.0	0.1	1.4	100.0	98.4	685
6+	33.6	58.0	0.2	0.0	8.2	100.0	91.6	305
Residence								
Urban	55.9	42.7	0.0	0.0	1.4	100.0	98.6	1,698
Rural	42.1	55.3	0.0	0.1	2.5	100.0	97.4	2,285
Education[3]								
No education	31.8	57.6	0.2	0.2	10.3	100.0	89.3	350
Primary	49.5	48.9	0.0	0.1	1.4	100.0	98.5	1,076
Secondary	47.4	51.1	0.0	0.0	1.4	100.0	98.6	1,756
More than secondary	54.3	45.2	0.0	0.0	0.5	100.0	99.5	800
Wealth quintile								
Lowest	36.5	58.2	0.1	0.2	5.1	100.0	94.6	836
Second	42.6	56.2	0.0	0.0	1.2	100.0	98.8	745
Middle	45.8	53.2	0.0	0.0	1.0	100.0	99.0	720
Fourth	50.3	47.8	0.0	0.0	1.9	100.0	98.1	914
Highest	65.0	34.4	0.0	0.0	0.5	100.0	99.5	768
Total	48.0	49.9	0.0	0.0	2.0	100.0	97.9	3,983
				STILLBIRTHS				
Total	57.5	33.4	0.0	0.0	9.1	100.0	90.9	69
				LIVE BIRTHS AND STILLBIRTHS[4]				
Total	48.1	49.7	0.0	0.0	2.1	100.0	97.8	4,035

Notes: If more than one source of ANC was mentioned, only the provider with the highest qualifications is considered in this tabulation. Stillbirths are foetal deaths in pregnancies lasting 28 or more weeks. When pregnancy duration is reported in months, stillbirths are foetal deaths in pregnancies lasting 7 or more months.

[1] Skilled provider is a doctor, nurse, midwife, or clinical officer.

[2] Birth order refers to the order of the birth among the respondent's live births.

[3] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

[4] For women who had both a live birth and a stillbirth in the 2 years preceding the survey, data are tabulated for the most recent birth only.

Table 9.1C Antenatal care: County level

Percent distribution of women age 15–49 who had a live birth and/or stillbirth in the 2 years preceding the survey by antenatal care (ANC) provider during pregnancy for the most recent birth or stillbirth, and percentage receiving ANC from a skilled provider for the most recent live birth or stillbirth, according to county, Kenya Mini DHS 2025–26

County	ANC provider				No ANC	Total	Percentage receiving ANC from a skilled provider[1]	Number of women
	Doctor	Nurse/ midwife/ clinical officer	Community health worker/ fieldworker	Traditional birth attendant				
				LIVE BIRTHS				
Mombasa	58.3	40.1	0.0	0.0	1.5	100.0	98.5	127
Kwale	98.9	1.1	0.0	0.0	0.0	100.0	100.0	71
Kilifi	44.2	55.8	0.0	0.0	0.0	100.0	100.0	102
Tana River	1.1	97.5	0.0	0.0	1.4	100.0	98.6	40
Lamu	10.7	87.2	0.0	0.0	2.1	100.0	97.9	10
Taita Taveta	65.6	31.2	0.0	0.0	3.2	100.0	96.8	30
Garissa	36.0	40.8	0.0	0.0	23.2	100.0	76.8	42
Wajir	34.2	47.5	0.0	0.0	18.3	100.0	81.7	36
Mandera	3.4	75.4	0.0	1.9	19.3	100.0	78.8	39
Marsabit	26.6	66.6	1.1	0.0	5.7	100.0	93.2	35
Isiolo	52.5	47.5	0.0	0.0	0.0	100.0	100.0	16
Meru	56.1	41.4	0.0	0.0	2.5	100.0	97.5	72
Tharaka-Nithi	(78.8)	(18.1)	(0.0)	(0.0)	(3.1)	100.0	(96.9)	22
Embu	(44.8)	(55.2)	(0.0)	(0.0)	(0.0)	100.0	(100.0)	27
Kitui	83.8	14.8	0.0	0.0	1.4	100.0	98.6	64
Machakos	26.2	73.8	0.0	0.0	0.0	100.0	100.0	156
Makueni	36.3	63.7	0.0	0.0	0.0	100.0	100.0	59
Nyandarua	65.3	32.7	0.0	0.0	2.0	100.0	98.0	48
Nyeri	76.0	24.0	0.0	0.0	0.0	100.0	100.0	42
Kirinyaga	(67.9)	(30.3)	(0.0)	(0.0)	(1.9)	100.0	(98.1)	37
Murang'a	62.5	36.0	0.0	0.0	1.5	100.0	98.5	59
Kiambu	83.4	14.7	0.0	0.0	1.8	100.0	98.2	269
Turkana	28.0	71.0	1.0	0.0	0.0	100.0	99.0	62
West Pokot	10.5	81.5	0.0	0.0	8.1	100.0	91.9	73
Samburu	16.3	74.9	0.0	0.0	8.8	100.0	91.2	31
Trans Nzoia	30.7	67.7	0.0	0.0	1.6	100.0	98.4	57
Uasin Gishu	37.9	62.1	0.0	0.0	0.0	100.0	100.0	123
Elgeyo-Marakwet	11.1	85.4	0.0	0.0	3.5	100.0	96.5	32
Nandi	39.7	60.3	0.0	0.0	0.0	100.0	100.0	75
Baringo	15.1	81.6	0.0	0.0	3.3	100.0	96.7	53
Laikipia	37.1	57.1	0.0	0.0	5.8	100.0	94.2	50
Nakuru	68.8	31.2	0.0	0.0	0.0	100.0	100.0	222
Narok	27.4	67.4	0.0	1.1	4.0	100.0	94.8	87
Kajiado	53.5	45.7	0.0	0.0	0.8	100.0	99.2	123
Kericho	18.5	81.5	0.0	0.0	0.0	100.0	100.0	68
Bomet	33.6	66.4	0.0	0.0	0.0	100.0	100.0	89
Kakamega	46.1	52.1	0.0	0.0	1.8	100.0	98.2	126
Vihiga	40.0	60.0	0.0	0.0	0.0	100.0	100.0	38
Bungoma	57.0	42.2	0.0	0.0	0.8	100.0	99.2	128
Busia	68.9	29.8	0.0	0.0	1.2	100.0	98.8	78
Siaya	45.3	53.9	0.0	0.0	0.8	100.0	99.2	53
Kisumu	35.6	62.7	0.0	0.0	1.7	100.0	98.3	101
Homa Bay	30.7	69.3	0.0	0.0	0.0	100.0	100.0	81
Migori	24.1	75.9	0.0	0.0	0.0	100.0	100.0	131
Kisii	9.0	89.4	0.0	0.0	1.6	100.0	98.4	84
Nyamira	17.7	81.2	0.0	0.0	1.1	100.0	98.9	40
Nairobi City	60.7	37.2	0.0	0.0	2.1	100.0	97.9	577
Total	48.0	49.9	0.0	0.0	2.0	100.0	97.9	3,983
				STILLBIRTHS				
Total	57.5	33.4	0.0	0.0	9.1	100.0	90.9	69
				LIVE BIRTHS AND STILLBIRTHS[2]				
Total	48.1	49.7	0.0	0.0	2.1	100.0	97.8	4,035

Notes: If more than one source of ANC was mentioned, only the provider with the highest qualifications is considered in this tabulation. Stillbirths are foetal deaths in pregnancies lasting 28 or more weeks. When pregnancy duration is reported in months, stillbirths are foetal deaths in pregnancies lasting 7 or more months. Figures in parentheses are based on 25–49 unweighted cases.

[1] Skilled provider is a doctor, nurse, midwife, or clinical officer.

[2] For women who had both a live birth and a stillbirth in the 2 years preceding the survey, data are tabulated for the most recent birth only.

Table 9.2 Number of antenatal care visits and timing of first visit

Percent distribution of women age 15–49 who had a live birth and/or a stillbirth in the 2 years preceding the survey, by number of antenatal care (ANC) visits during pregnancy for the most recent live birth or stillbirth, and by the timing of the first visit; and among women with ANC, median months pregnant at first visit, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Number of ANC visits						Total	4+ ANC visits	Number of months pregnant at time of first ANC visit					Total	Number of women	Median months pregnant at first visit (for those with ANC)	Number of women with ANC
	None	1	2–3	4–7	8+	Don't know			No ANC	<4	4–6	7+	Don't know				
LIVE BIRTHS																	
Age at birth																	
<20	2.3	3.5	27.6	61.9	4.7	0.0	100.0	66.6	2.3	26.1	59.3	12.3	0.0	100.0	492	5.2	481
20–34	1.8	2.4	21.1	64.6	10.0	0.2	100.0	74.6	1.8	38.6	52.0	7.6	0.0	100.0	2,943	4.5	2,891
35–49	3.2	2.3	28.4	54.6	11.2	0.3	100.0	65.8	3.2	31.4	51.0	14.4	0.0	100.0	548	5.0	530
Birth order[1]																	
1	1.6	2.6	20.3	65.2	10.3	0.1	100.0	75.5	1.6	39.3	52.4	6.8	0.0	100.0	1,269	4.5	1,249
2–3	1.5	1.8	21.7	65.3	9.5	0.2	100.0	74.8	1.5	38.7	50.8	9.0	0.0	100.0	1,724	4.5	1,698
4–5	1.4	3.4	25.9	58.1	11.0	0.1	100.0	69.1	1.4	31.2	55.7	11.5	0.1	100.0	685	5.0	675
6+	8.2	4.6	33.4	50.8	2.6	0.4	100.0	53.4	8.2	18.5	58.8	14.5	0.0	100.0	305	5.4	280
Residence																	
Urban	1.4	1.9	16.5	67.3	12.7	0.1	100.0	80.1	1.4	41.9	49.0	7.6	0.0	100.0	1,698	4.4	1,674
Rural	2.5	3.0	27.7	59.7	7.1	0.2	100.0	66.7	2.5	31.7	55.5	10.3	0.0	100.0	2,285	4.8	2,229
Education[2]																	
No education	10.3	7.4	30.1	46.3	6.0	0.0	100.0	52.3	10.3	24.0	54.5	11.2	0.0	100.0	350	5.1	314
Primary	1.4	2.0	29.0	60.1	7.3	0.2	100.0	67.4	1.4	28.0	58.3	12.3	0.1	100.0	1,076	5.0	1,061
Secondary	1.4	2.9	22.3	65.9	7.3	0.2	100.0	73.2	1.4	36.3	53.4	8.9	0.0	100.0	1,756	4.7	1,731
More than secondary	0.5	0.3	12.9	67.4	18.8	0.1	100.0	86.2	0.5	51.6	43.3	4.6	0.0	100.0	800	3.9	796
Wealth quintile																	
Lowest	5.1	4.9	31.5	55.8	2.7	0.1	100.0	58.5	5.1	25.3	58.1	11.6	0.0	100.0	836	5.1	793
Second	1.2	2.9	29.1	60.1	6.1	0.5	100.0	66.2	1.2	30.6	56.8	11.2	0.1	100.0	745	4.9	736
Middle	1.0	1.8	23.0	63.5	10.5	0.2	100.0	74.0	1.0	31.6	56.5	10.9	0.0	100.0	720	4.9	713
Fourth	1.9	1.3	19.3	67.7	9.7	0.1	100.0	77.4	1.9	37.6	52.7	7.7	0.0	100.0	914	4.6	896
Highest	0.5	1.7	11.7	67.2	18.9	0.0	100.0	86.1	0.5	55.4	39.6	4.5	0.0	100.0	768	3.8	764
Total	2.0	2.5	22.9	62.9	9.5	0.2	100.0	72.4	2.0	36.0	52.8	9.1	0.0	100.0	3,983	4.7	3,902
STILLBIRTHS																	
Total	9.1	0.8	18.9	69.8	1.4	0.0	100.0	71.2	9.1	35.4	49.9	5.6	0.0	100.0	69	4.3	62
LIVE BIRTHS AND STILLBIRTHS[3]																	
Total	2.1	2.5	22.8	63.0	9.4	0.2	100.0	72.3	2.1	36.1	52.6	9.1	0.0	100.0	4,035	4.6	3,948

[1] Birth order refers to the order of the birth among the respondent's live births.
[2] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.
[3] For women who had both a live birth and a stillbirth in the 2 years preceding the survey, data are tabulated for the most recent birth only.

Table 9.2C Number of antenatal care visits and timing of first visit: County level

Percent distribution of women age 15–49 who had a live birth and/or a stillbirth in the 2 years preceding the survey, by number of antenatal care (ANC) visits during pregnancy for the most recent live birth or stillbirth, and by the timing of the first visit; and among women with ANC, median months pregnant at first visit, according to county, Kenya Mini DHS 2025–26

	Number of ANC visits							4+ ANC visits	Number of months pregnant at time of first ANC visit						Number of women	Median months pregnant at first visit (for those with ANC)	Number of women with ANC
County	None	1	2–3	4–7	8+	Don't know	Total		No ANC	<4	4–6	7+	Don't know	Total			
LIVE BIRTHS																	
Mombasa	1.5	2.6	10.2	74.7	10.9	0.0	100.0	85.6	1.5	31.8	54.6	12.0	0.0	100.0	127	4.8	125
Kwale	0.0	0.7	14.9	77.7	6.7	0.0	100.0	84.4	0.0	37.7	56.9	5.4	0.0	100.0	71	4.4	71
Kilifi	0.0	1.7	14.9	74.3	9.1	0.0	100.0	83.4	0.0	40.8	51.5	7.7	0.0	100.0	102	4.5	102
Tana River	1.4	0.8	22.7	70.8	4.3	0.0	100.0	75.1	1.4	26.8	63.2	8.7	0.0	100.0	40	4.8	40
Lamu	2.1	2.9	31.4	61.7	1.9	0.0	100.0	63.6	2.1	26.2	56.8	14.8	0.0	100.0	10	5.3	10
Taita Taveta	3.2	2.9	19.9	61.6	11.9	0.5	100.0	73.5	3.2	44.3	44.5	8.0	0.0	100.0	30	4.2	29
Garissa	23.2	7.3	31.4	33.7	4.4	0.0	100.0	38.1	23.2	31.1	40.4	5.3	0.0	100.0	42	4.6	32
Wajir	18.3	0.5	21.6	59.6	0.0	0.0	100.0	59.6	18.3	7.2	70.7	3.8	0.0	100.0	36	4.9	30
Mandera	19.3	9.1	49.6	22.0	0.0	0.0	100.0	22.0	19.3	20.2	52.6	7.9	0.0	100.0	39	4.9	32
Marsabit	5.7	1.1	16.0	75.0	2.2	0.0	100.0	77.2	5.7	33.9	52.7	7.7	0.0	100.0	35	4.4	33
Isiolo	0.0	2.0	33.6	58.9	5.5	0.0	100.0	64.4	0.0	27.3	57.2	15.2	0.3	100.0	16	4.8	16
Meru	2.5	0.0	50.0	43.2	4.3	0.0	100.0	47.5	2.5	25.4	50.6	21.5	0.0	100.0	72	5.5	70
Tharaka-Nithi	(3.1)	(0.0)	(19.0)	(73.4)	(4.6)	(0.0)	100.0	(78.0)	(3.1)	(26.3)	(64.9)	(5.8)	(0.0)	100.0	22	(4.7)	21
Embu	(0.0)	(2.3)	(31.8)	(50.6)	(15.3)	(0.0)	100.0	(65.9)	(0.0)	(45.8)	(43.3)	(10.9)	(0.0)	100.0	27	(4.6)	27
Kitui	1.4	3.1	13.1	75.1	7.3	0.0	100.0	82.4	1.4	31.1	64.8	2.7	0.0	100.0	64	4.6	63
Machakos	0.0	0.7	26.8	64.6	7.9	0.0	100.0	72.6	0.0	38.4	54.9	6.7	0.0	100.0	156	4.8	156
Makueni	0.0	0.0	10.9	65.3	23.8	0.0	100.0	89.1	0.0	41.9	54.3	3.8	0.0	100.0	59	4.4	59
Nyandarua	2.0	0.7	39.1	47.9	10.3	0.0	100.0	58.2	2.0	28.8	63.0	6.2	0.0	100.0	48	4.9	47
Nyeri	0.0	5.1	14.8	70.7	9.4	0.0	100.0	80.1	0.0	42.5	51.6	5.9	0.0	100.0	42	4.3	42
Kirinyaga	(1.9)	(0.0)	(16.7)	(70.0)	(11.5)	(0.0)	100.0	(81.5)	(1.9)	(55.2)	(33.1)	(9.8)	(0.0)	100.0	37	(3.8)	36
Murang'a	1.5	0.0	30.7	58.5	9.3	0.0	100.0	67.8	1.5	30.7	54.4	13.4	0.0	100.0	59	5.0	58
Kiambu	1.8	0.3	13.6	68.4	15.8	0.0	100.0	84.3	1.8	56.2	35.4	6.6	0.0	100.0	269	3.6	264
Turkana	0.0	11.7	37.4	50.9	0.0	0.0	100.0	50.9	0.0	26.1	65.0	8.8	0.0	100.0	62	5.0	62
West Pokot	8.1	11.0	43.6	36.1	1.3	0.0	100.0	37.4	8.1	12.0	65.2	14.8	0.0	100.0	73	5.8	67
Samburu	8.8	2.1	23.8	59.8	5.6	0.0	100.0	65.3	8.8	36.0	47.4	7.8	0.0	100.0	31	4.5	28
Trans Nzoia	1.6	0.0	38.1	49.0	11.4	0.0	100.0	60.3	1.6	33.2	56.2	9.0	0.0	100.0	57	4.8	56
Uasin Gishu	0.0	0.8	17.2	67.5	14.5	0.0	100.0	82.0	0.0	43.4	50.2	6.4	0.0	100.0	123	4.3	123
Elgeyo-Marakwet	3.5	5.4	37.4	45.9	7.8	0.0	100.0	53.6	3.5	32.1	51.0	13.4	0.0	100.0	32	5.2	31
Nandi	0.0	2.3	34.6	57.5	4.4	1.2	100.0	61.9	0.0	27.3	66.1	5.4	1.2	100.0	75	5.1	75
Baringo	3.3	3.3	31.2	58.3	3.2	0.8	100.0	61.5	3.3	16.3	71.2	9.3	0.0	100.0	53	5.3	51
Laikipia	5.8	6.9	22.2	59.3	5.7	0.0	100.0	65.0	5.8	26.4	53.8	14.1	0.0	100.0	50	4.8	47
Nakuru	0.0	2.6	23.1	68.3	6.0	0.0	100.0	74.3	0.0	34.8	53.6	11.6	0.0	100.0	222	4.8	222
Narok	4.0	2.2	37.0	53.7	3.0	0.0	100.0	56.7	4.0	14.4	63.2	18.4	0.0	100.0	87	5.3	83
Kajiado	0.8	1.5	22.2	66.1	9.4	0.0	100.0	75.6	0.8	39.2	51.0	9.0	0.0	100.0	123	4.6	122
Kericho	0.0	10.2	34.7	53.3	1.8	0.0	100.0	55.1	0.0	14.9	62.6	22.5	0.0	100.0	68	5.5	68
Bomet	0.0	1.0	29.8	58.6	10.6	0.0	100.0	69.2	0.0	31.7	50.6	17.7	0.0	100.0	89	5.1	89
Kakamega	1.8	4.1	18.6	70.0	5.5	0.0	100.0	75.5	1.8	40.3	54.1	3.8	0.0	100.0	126	4.4	124
Vihiga	0.0	1.9	18.9	60.5	15.8	2.9	100.0	76.3	0.0	67.5	23.1	9.4	0.0	100.0	38	3.4	38
Bungoma	0.8	0.8	25.3	62.6	10.4	0.0	100.0	73.1	0.8	38.4	52.5	8.3	0.0	100.0	128	4.5	127
Busia	1.2	1.8	15.9	68.5	9.9	2.6	100.0	78.5	1.2	37.1	56.1	5.6	0.0	100.0	78	4.7	77
Siaya	0.8	1.5	27.5	64.3	5.9	0.0	100.0	70.2	0.8	35.9	49.8	13.5	0.0	100.0	53	4.6	52
Kisumu	1.7	2.3	27.3	56.9	10.7	1.1	100.0	67.6	1.7	39.1	52.3	6.9	0.0	100.0	101	4.4	99
Homa Bay	0.0	1.4	25.0	62.1	11.5	0.0	100.0	73.6	0.0	45.4	51.3	3.3	0.0	100.0	81	4.3	81
Migori	0.0	4.7	24.9	61.2	9.3	0.0	100.0	70.4	0.0	32.8	54.1	13.1	0.0	100.0	131	4.9	131
Kisii	1.6	4.5	36.0	54.4	3.4	0.0	100.0	57.8	1.6	27.6	55.7	15.0	0.0	100.0	84	5.2	82
Nyamira	1.1	2.6	27.5	58.7	8.7	1.3	100.0	67.4	1.1	25.1	65.9	7.9	0.0	100.0	40	5.1	39
Nairobi City	2.1	2.2	12.0	68.2	15.4	0.0	100.0	83.7	2.1	40.8	50.2	7.0	0.0	100.0	577	4.5	565
Total	2.0	2.5	22.9	62.9	9.5	0.2	100.0	72.4	2.0	36.0	52.8	9.1	0.0	100.0	3,983	4.7	3,902
STILLBIRTHS																	
Total	9.1	0.8	18.9	69.8	1.4	0.0	100.0	71.2	9.1	35.4	49.9	5.6	0.0	100.0	69	4.3	62
LIVE BIRTHS AND STILLBIRTHS[1]																	
Total	2.1	2.5	22.8	63.0	9.4	0.2	100.0	72.3	2.1	36.1	52.6	9.1	0.0	100.0	4,035	4.6	3,948

Note: Figures in parentheses are based on 25–49 unweighted cases.

[1] For women who had both a live birth and a stillbirth in the 2 years preceding the survey, data are tabulated for the most recent birth only.

Table 9.3.1 Components of antenatal care among women receiving antenatal care

Among women age 15–49 receiving antenatal care (ANC) for the most recent live birth and/or stillbirth in the 2 years preceding the survey, percentage receiving specific antenatal services from a health care provider, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Among women who received ANC for their most recent live birth or stillbirth in the past 2 years, percentage who received specific services from a health care provider during ANC							Number of women with ANC for their most recent live birth and/or stillbirth in the past 2 years
	Blood pressure measured	Urine sample taken	Blood sample taken	Baby's heartbeat listened for	Counselled about maternal diet	Counselled about breastfeeding	Asked about vaginal bleeding	
LIVE BIRTHS								
Age at birth								
<20	98.0	91.8	94.4	96.4	83.0	84.0	75.5	481
20–34	98.4	97.0	97.9	97.9	90.7	87.7	85.5	2,891
35–49	98.5	97.0	98.0	98.0	88.3	88.3	83.2	530
Birth order[1]								
1	98.3	96.2	97.5	98.1	89.3	86.6	83.0	1,249
2–3	98.5	97.5	98.1	98.1	91.5	90.1	86.5	1,698
4–5	98.5	96.5	97.4	97.9	89.3	85.7	83.7	675
6+	96.9	90.2	94.0	94.0	77.7	77.5	73.4	280
Residence								
Urban	99.4	98.2	98.9	98.8	93.3	91.3	87.4	1,674
Rural	97.5	95.0	96.4	96.9	86.6	84.4	81.4	2,229
Education[2]								
No education	95.5	84.2	90.5	92.8	77.5	73.3	71.4	314
Primary	97.7	96.1	97.2	97.0	88.1	87.1	80.3	1,061
Secondary	98.9	97.9	98.5	98.3	90.6	88.3	85.5	1,731
More than secondary	99.2	98.2	98.5	99.5	93.3	91.1	90.2	796
Wealth quintile								
Lowest	96.2	91.1	94.2	95.8	80.6	79.0	75.9	793
Second	97.8	95.9	96.6	96.1	89.2	85.5	80.3	736
Middle	98.8	98.6	98.8	98.5	90.1	90.6	88.2	713
Fourth	99.5	98.6	99.0	98.7	92.5	89.5	86.1	896
Highest	99.3	97.8	98.8	99.7	94.5	92.2	89.3	764
Total	98.3	96.4	97.5	97.8	89.4	87.3	83.9	3,902
STILLBIRTHS								
Total	100.0	98.8	98.3	100.0	95.0	74.0	77.2	62
LIVE BIRTHS AND STILLBIRTHS[3]								
Total	98.4	96.4	97.5	97.8	89.5	87.1	83.8	3,948

Note: The denominator for this table includes all women with a birth in the 2 years preceding the survey who received ANC for this birth.
[1] Birth order refers to the order of the birth among the respondent's live births.
[2] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.
[3] For women who had both a live birth and a stillbirth in the 2 years preceding the survey, data are tabulated for the most recent birth only.

Table 9.3.1C Components of antenatal care among women receiving antenatal care: County level

Among women age 15–49 receiving antenatal care (ANC) for the most recent live birth and/or stillbirth in the 2 years preceding the survey, percentage receiving specific antenatal services from a health care provider, according to county, Kenya Mini DHS 2025–26

County	Among women who received ANC for their most recent live birth or stillbirth in the past 2 years, percentage who received specific services from a health care provider during ANC							Number of women with ANC for their most recent live birth and/or stillbirth in the past 2 years
	Blood pressure measured	Urine sample taken	Blood sample taken	Baby's heartbeat listened for	Counselled about maternal diet	Counselled about breastfeeding	Asked about vaginal bleeding	
LIVE BIRTHS								
Mombasa	100.0	99.0	100.0	100.0	97.6	97.2	95.7	125
Kwale	97.9	98.8	98.8	100.0	98.1	96.6	91.6	71
Kilifi	100.0	100.0	100.0	100.0	98.7	100.0	99.1	102
Tana River	100.0	100.0	99.4	98.0	86.3	84.5	88.3	40
Lamu	100.0	100.0	97.7	100.0	93.5	92.3	96.0	10
Taita Taveta	100.0	95.4	100.0	97.8	97.8	92.0	87.4	29
Garissa	95.8	92.3	93.5	90.5	74.5	66.6	59.2	32
Wajir	98.9	76.4	76.8	88.0	56.6	62.7	54.7	30
Mandera	93.5	90.2	85.0	80.3	72.7	67.2	75.0	32
Marsabit	100.0	96.3	95.4	92.9	96.6	88.1	91.8	33
Isiolo	98.8	99.7	100.0	93.1	75.7	68.9	80.8	16
Meru	99.1	97.4	95.9	100.0	91.8	79.9	79.1	70
Tharaka-Nithi	(94.7)	(97.2)	(100.0)	(100.0)	(89.3)	(77.9)	(69.9)	21
Embu	(97.9)	(97.9)	(97.9)	(97.9)	(93.2)	(97.9)	(91.2)	27
Kitui	98.7	98.7	98.7	97.0	92.0	93.8	95.6	63
Machakos	98.0	97.3	97.3	96.4	92.9	81.9	87.0	156
Makueni	100.0	100.0	98.4	99.0	95.1	78.2	94.3	59
Nyandarua	98.3	100.0	100.0	100.0	90.1	88.2	73.2	47
Nyeri	100.0	100.0	100.0	98.4	89.7	84.5	83.6	42
Kirinyaga	(100.0)	(96.4)	(100.0)	(97.4)	(98.3)	(97.2)	(94.0)	36
Murang'a	100.0	100.0	100.0	100.0	97.4	91.4	98.1	58
Kiambu	99.6	99.6	99.6	99.6	99.6	95.0	95.1	264
Turkana	95.3	66.4	83.1	96.2	69.5	67.2	59.3	62
West Pokot	82.5	78.3	87.2	89.6	81.2	80.0	72.1	67
Samburu	96.9	88.7	91.9	95.4	74.9	44.0	51.1	28
Trans Nzoia	99.2	98.1	100.0	98.4	91.5	92.7	88.0	56
Uasin Gishu	96.2	99.2	100.0	99.2	82.8	73.9	86.1	123
Elgeyo-Marakwet	100.0	97.1	98.5	87.0	91.0	75.3	79.0	31
Nandi	94.7	94.8	94.4	94.5	88.2	92.2	87.1	75
Baringo	95.3	93.6	95.9	100.0	84.6	87.2	85.5	51
Laikipia	100.0	100.0	100.0	100.0	91.0	90.6	86.0	47
Nakuru	99.1	96.1	98.4	97.5	91.4	86.9	88.6	222
Narok	97.4	97.6	97.8	98.6	82.8	75.7	77.3	83
Kajiado	98.5	97.4	99.2	100.0	97.2	86.8	90.8	122
Kericho	100.0	100.0	100.0	100.0	84.4	83.4	79.4	68
Bomet	100.0	100.0	98.2	100.0	65.5	66.9	41.7	89
Kakamega	98.0	91.7	95.0	97.2	85.6	89.2	69.3	124
Vihiga	97.2	97.1	100.0	100.0	89.9	94.9	82.6	38
Bungoma	96.1	97.1	97.2	95.4	83.8	84.4	81.0	127
Busia	97.8	99.1	97.4	99.1	93.1	91.9	93.7	77
Siaya	100.0	97.3	97.0	96.2	91.0	97.9	93.1	52
Kisumu	97.3	96.1	94.9	94.0	87.9	93.8	83.1	99
Homa Bay	97.3	91.4	93.7	97.4	88.4	89.2	81.6	81
Migori	98.4	95.7	98.0	96.5	80.5	80.1	85.1	131
Kisii	98.6	98.6	100.0	100.0	90.4	88.7	91.5	82
Nyamira	100.0	97.4	100.0	98.5	83.9	86.2	75.6	39
Nairobi City	100.0	97.3	98.4	98.9	92.1	94.9	81.9	565
Total	98.3	96.4	97.5	97.8	89.4	87.3	83.9	3,902
STILLBIRTHS								
Total	100.0	98.8	98.3	100.0	95.0	74.0	77.2	62
LIVE BIRTHS AND STILLBIRTHS[1]								
Total	98.4	96.4	97.5	97.8	89.5	87.1	83.8	3,948

Notes: The denominator for this table includes all women with a birth in the 2 years preceding the survey who received ANC for this birth. Figures in parentheses are based on 25–49 unweighted cases.

[1] Birth order refers to the order of the birth among the respondent's live births.

Table 9.3.2 Components of antenatal care among all women

Among all women age 15–49 with a live birth and/or stillbirth in the 2 years preceding the survey, percentage receiving specific antenatal services from a health care provider for their most recent live birth and/or stillbirth, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Blood pressure measured	Urine sample taken	Blood sample taken	Baby's heartbeat listened for	Counselled about maternal diet	Counselled about breastfeeding	Asked about vaginal bleeding	Number of women with a live birth and/or stillbirth in the past 2 years
LIVE BIRTHS								
Age at birth								
<20	95.7	89.7	92.2	94.2	81.1	82.0	73.7	492
20–34	96.6	95.3	96.2	96.2	89.1	86.2	84.0	2,943
35–49	95.4	93.9	94.9	94.9	85.5	85.5	80.6	548
Birth order[1]								
1	96.8	94.7	96.0	96.5	87.9	85.3	81.7	1,269
2–3	97.0	96.0	96.6	96.6	90.2	88.8	85.2	1,724
4–5	97.1	95.1	96.0	96.5	88.0	84.4	82.5	685
6+	88.9	82.8	86.3	86.3	71.3	71.2	67.4	305
Residence								
Urban	98.0	96.8	97.5	97.4	91.9	90.0	86.1	1,698
Rural	95.1	92.7	94.0	94.6	84.4	82.3	79.4	2,285
Education[2]								
No education	85.7	75.6	81.2	83.3	69.6	65.7	64.1	350
Primary	96.3	94.7	95.8	95.7	86.8	85.9	79.2	1,076
Secondary	97.5	96.5	97.1	96.9	89.3	87.0	84.3	1,756
More than secondary	98.7	97.7	98.0	99.0	92.8	90.7	89.8	800
Wealth quintile								
Lowest	91.3	86.5	89.4	90.9	76.6	75.0	72.0	836
Second	96.6	94.7	95.4	94.9	88.1	84.4	79.3	745
Middle	97.8	97.6	97.8	97.5	89.2	89.7	87.3	720
Fourth	97.6	96.7	97.1	96.8	90.7	87.8	84.4	914
Highest	98.7	97.3	98.3	99.1	94.0	91.7	88.9	768
Total	96.4	94.4	95.5	95.8	87.6	85.6	82.2	3,983
STILLBIRTHS								
Total	90.9	89.7	89.3	90.9	86.3	67.3	70.2	69
LIVE BIRTHS AND STILLBIRTHS[3]								
Total	96.2	94.3	95.4	95.7	87.6	85.2	82.0	4,035

Note: The denominator for this table includes all women with a birth in the 2 years preceding the survey, whether or not they received ANC for that birth.

[1] Birth order refers to the order of the birth among the respondent's live births.

[2] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

[3] For women who had both a live birth and a stillbirth in the 2 years preceding the survey, data are tabulated for the most recent birth only.

Table 9.3.2C Components of antenatal care among all women: County level

Among all women age 15–49 with a live birth and/or stillbirth in the 2 years preceding the survey, percentage receiving specific antenatal services from a health care provider for their most recent live birth and/or stillbirth, according to county, Kenya Mini DHS 2025–26

County	Blood pressure measured	Urine sample taken	Blood sample taken	Baby's heartbeat listened for	Counselled about maternal diet	Counselled about breastfeeding	Asked about vaginal bleeding	Number of women with a live birth and/or stillbirth in the past 2 years
LIVE BIRTHS								
Mombasa	98.5	97.5	98.5	98.5	96.2	95.7	94.2	127
Kwale	97.9	98.8	98.8	100.0	98.1	96.6	91.6	71
Kilifi	100.0	100.0	100.0	100.0	98.7	100.0	99.1	102
Tana River	98.6	98.6	98.0	96.6	85.1	83.4	87.1	40
Lamu	97.9	97.9	95.7	97.9	91.6	90.4	94.0	10
Taita Taveta	96.8	92.3	96.8	94.7	94.7	89.1	84.6	30
Garissa	73.6	70.9	71.8	69.5	57.2	51.2	45.5	42
Wajir	80.8	62.4	62.8	71.9	46.2	51.2	44.7	36
Mandera	75.4	72.8	68.6	64.8	58.7	54.2	60.5	39
Marsabit	94.3	90.8	90.0	87.6	91.1	83.1	86.6	35
Isiolo	98.8	99.7	100.0	93.1	75.7	68.9	80.8	16
Meru	96.6	95.0	93.6	97.5	89.5	77.9	77.2	72
Tharaka-Nithi	(91.8)	(94.3)	(96.9)	(96.9)	(86.6)	(75.5)	(67.8)	22
Embu	(97.9)	(97.9)	(97.9)	(97.9)	(93.2)	(97.9)	(91.2)	27
Kitui	97.2	97.2	97.2	95.6	90.7	92.5	94.2	64
Machakos	98.0	97.3	97.3	96.4	92.9	81.9	87.0	156
Makueni	100.0	100.0	98.4	99.0	95.1	78.2	94.3	59
Nyandarua	96.3	98.0	98.0	98.0	88.3	86.5	71.7	48
Nyeri	100.0	100.0	100.0	98.4	89.7	84.5	83.6	42
Kirinyaga	(98.1)	(94.7)	(98.1)	(95.6)	(96.4)	(95.4)	(92.2)	37
Murang'a	98.5	98.5	98.5	98.5	96.0	90.1	96.6	59
Kiambu	97.8	97.8	97.8	97.8	97.8	93.3	93.4	269
Turkana	95.3	66.4	83.1	96.2	69.5	67.2	59.3	62
West Pokot	75.8	72.0	80.1	82.4	74.6	73.5	66.3	73
Samburu	88.4	80.9	83.8	87.0	68.3	40.2	46.7	31
Trans Nzoia	97.6	96.5	98.4	96.8	90.1	91.2	86.6	57
Uasin Gishu	96.2	99.2	100.0	99.2	82.8	73.9	86.1	123
Elgeyo-Marakwet	96.5	93.7	95.1	83.9	87.8	72.7	76.3	32
Nandi	94.7	94.8	94.4	94.5	88.2	92.2	87.1	75
Baringo	92.2	90.5	92.8	96.7	81.9	84.4	82.7	53
Laikipia	94.2	94.2	94.2	94.2	85.7	85.3	80.9	50
Nakuru	99.1	96.1	98.4	97.5	91.4	86.9	88.6	222
Narok	93.5	93.7	93.9	94.6	79.5	72.7	74.2	87
Kajiado	97.7	96.7	98.5	99.2	96.4	86.1	90.1	123
Kericho	100.0	100.0	100.0	100.0	84.4	83.4	79.4	68
Bomet	100.0	100.0	98.2	100.0	65.5	66.9	41.7	89
Kakamega	96.3	90.1	93.3	95.4	84.1	87.6	68.1	126
Vihiga	97.2	97.1	100.0	100.0	89.9	94.9	82.6	38
Bungoma	95.3	96.3	96.4	94.6	83.1	83.7	80.3	128
Busia	96.6	97.9	96.2	97.9	91.9	90.8	92.6	78
Siaya	99.2	96.5	96.2	95.4	90.3	97.2	92.4	53
Kisumu	95.7	94.4	93.3	92.4	86.4	92.1	81.7	101
Homa Bay	97.3	91.4	93.7	97.4	88.4	89.2	81.6	81
Migori	98.4	95.7	98.0	96.5	80.5	80.1	85.1	131
Kisii	96.9	96.9	98.4	98.4	88.9	87.2	90.0	84
Nyamira	98.9	96.3	98.9	97.4	83.0	85.3	74.8	40
Nairobi City	97.9	95.3	96.3	96.9	90.2	92.9	80.1	577
Total	96.4	94.4	95.5	95.8	87.6	85.6	82.2	3,983
STILLBIRTHS								
Total	90.9	89.7	89.3	90.9	86.3	67.3	70.2	69
LIVE BIRTHS AND STILLBIRTHS[1]								
Total	96.2	94.3	95.4	95.7	87.6	85.2	82.0	4,035

Notes: The denominator for this table includes all women with a birth in the 2 years preceding the survey, whether or not they received ANC for that birth. Figures in parentheses are based on 25–49 unweighted cases.
[1] For women who had both a live birth and a stillbirth in the 2 years preceding the survey, data are tabulated for the most recent birth only.

Table 9.4 Deworming and iron-containing supplementation during pregnancy

Among women age 15–49 with a live birth and/or stillbirth in the 2 years preceding the survey, percentage who took deworming medication during the pregnancy for the most recent birth, percentage who took any iron-containing supplements during the pregnancy for the most recent live birth or stillbirth, and percent distribution of the number of days during which the women took iron-containing supplements during the pregnancy for the most recent live birth or stillbirth, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Among women with a live birth and/or stillbirth in the past 2 years, percentage who during pregnancy for the most recent live birth or stillbirth:		Number of days during which women with a live birth and/or stillbirth in the past 2 years took iron-containing supplements[1] during pregnancy for the most recent live birth or stillbirth							Number of women with a live birth and/or stillbirth in the past 2 years
	Took deworming medication	Took any iron-containing supplements[1]	None	<60	60–89	90–179	180+	Don't know	Total	
LIVE BIRTHS										
Age at birth										
<20	17.1	86.9	13.1	21.2	11.5	36.9	16.5	0.7	100.0	492
20–34	27.3	89.4	10.6	13.7	9.4	38.1	27.6	0.5	100.0	2,943
35–49	33.0	87.7	12.3	12.9	12.1	37.2	25.1	0.5	100.0	548
Birth order[2]										
1	20.4	91.2	8.8	14.6	10.3	35.9	30.0	0.3	100.0	1,269
2–3	30.3	89.1	10.9	13.4	8.8	39.1	27.3	0.6	100.0	1,724
4–5	31.2	89.5	10.5	16.7	10.9	40.3	20.5	1.1	100.0	685
6+	24.1	76.2	23.8	16.2	13.9	32.9	13.2	0.2	100.0	305
Residence										
Urban	32.8	90.8	9.2	12.8	9.4	35.9	32.2	0.6	100.0	1,698
Rural	22.4	87.4	12.6	15.8	10.6	39.3	21.2	0.5	100.0	2,285
Education[3]										
No education	22.8	70.0	30.0	20.1	11.4	30.8	6.5	1.2	100.0	350
Primary	27.8	87.2	12.8	16.3	11.2	39.4	19.5	0.8	100.0	1,076
Secondary	25.1	90.8	9.2	14.2	9.3	39.7	27.1	0.5	100.0	1,756
More than secondary	31.0	95.1	4.9	10.5	9.5	34.6	40.4	0.1	100.0	800
Wealth quintile										
Lowest	23.0	81.9	18.1	18.5	11.2	38.0	13.5	0.7	100.0	836
Second	22.0	88.5	11.5	15.8	11.8	40.7	19.7	0.5	100.0	745
Middle	25.8	89.5	10.5	14.5	11.8	37.8	25.0	0.5	100.0	720
Fourth	30.5	92.2	7.8	13.9	10.2	38.0	29.8	0.3	100.0	914
Highest	32.3	92.3	7.7	9.9	5.3	34.7	41.6	0.8	100.0	768
Total	26.8	88.9	11.1	14.5	10.1	37.8	25.9	0.5	100.0	3,983
STILLBIRTHS										
Total	12.4	79.0	21.0	5.8	15.3	46.9	10.5	0.4	100.0	69
LIVE BIRTHS AND STILLBIRTHS[4]										
Total	26.7	88.9	11.1	14.4	10.1	38.1	25.7	0.5	100.0	4,035

Notes: Stillbirths are foetal deaths in pregnancies lasting 28 or more weeks. When pregnancy duration is reported in months, stillbirths are foetal deaths in pregnancies lasting 7 or more months.

[1] Iron tablets, iron syrups, or iron and folic acid supplements (IFAS)

[2] Birth order refers to the order of the birth among the respondent's live births.

[3] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

[4] For women who had both a live birth and a stillbirth in the 2 years preceding the survey, data are tabulated for the most recent birth only.

Table 9.4C Deworming and iron-containing supplementation during pregnancy: County level

Among women age 15–49 with a live birth and/or stillbirth in the 2 years preceding the survey, percentage who took deworming medication during the pregnancy for the most recent birth, percentage who took any iron-containing supplements during the pregnancy for the most recent live birth or stillbirth, and percent distribution of the number of days during which the women took iron-containing supplements during the pregnancy for the most recent live birth or stillbirth, according to county, Kenya Mini DHS 2025–26

County	Among women with a live birth and/or stillbirth in the past 2 years, percentage who during pregnancy for the most recent live birth or stillbirth:		Number of days during which women with a live birth and/or stillbirth in the past 2 years took iron-containing supplements[1] during pregnancy for the most recent live birth or stillbirth							Number of women with a live birth and/or stillbirth in the past 2 years
	Took deworming medication	Took any iron-containing supplements[1]	None	<60	60–89	90–179	180+	Don't know	Total	
LIVE BIRTHS										
Mombasa	59.6	96.3	3.7	29.9	13.0	29.0	22.8	1.5	100.0	127
Kwale	55.7	96.1	3.9	16.4	15.5	49.6	14.6	0.0	100.0	71
Kilifi	68.8	97.0	3.0	7.0	6.3	43.4	40.2	0.0	100.0	102
Tana River	50.2	87.0	13.0	4.9	12.0	52.9	17.2	0.0	100.0	40
Lamu	32.9	97.6	2.4	5.8	16.5	52.2	23.1	0.0	100.0	10
Taita Taveta	35.7	88.8	11.2	14.1	1.7	43.6	29.4	0.0	100.0	30
Garissa	12.3	48.7	51.3	17.1	12.8	14.0	4.8	0.0	100.0	42
Wajir	42.6	67.3	32.7	35.1	7.3	24.9	0.0	0.0	100.0	36
Mandera	3.1	64.9	35.1	20.3	21.7	22.5	0.3	0.0	100.0	39
Marsabit	12.7	89.9	10.1	20.2	15.8	45.1	8.9	0.0	100.0	35
Isiolo	12.2	83.3	16.7	30.0	5.5	41.5	6.4	0.0	100.0	16
Meru	18.4	84.6	15.4	32.5	21.6	25.2	3.8	1.5	100.0	72
Tharaka-Nithi	(18.0)	(89.4)	(10.6)	(35.7)	(3.9)	(26.9)	(23.0)	(0.0)	100.0	22
Embu	(50.0)	(97.7)	(2.3)	(9.7)	(22.1)	(28.6)	(34.8)	(2.5)	100.0	27
Kitui	16.7	92.5	7.5	7.8	5.1	55.7	23.8	0.0	100.0	64
Machakos	29.6	90.8	9.2	6.8	11.1	40.2	32.8	0.0	100.0	156
Makueni	17.9	96.1	3.9	4.8	4.1	38.2	48.9	0.0	100.0	59
Nyandarua	36.0	93.8	6.2	7.2	18.7	45.5	22.4	0.0	100.0	48
Nyeri	42.5	90.6	9.4	7.6	9.4	40.5	33.2	0.0	100.0	42
Kirinyaga	(54.0)	(96.7)	(3.3)	(8.6)	(5.2)	(38.3)	(44.7)	(0.0)	100.0	37
Murang'a	23.5	94.4	5.6	11.2	15.4	44.3	23.5	0.0	100.0	59
Kiambu	30.1	95.0	5.0	9.8	10.2	35.2	39.2	0.6	100.0	269
Turkana	26.7	82.3	17.7	9.1	11.4	45.2	16.6	0.0	100.0	62
West Pokot	15.5	75.5	24.5	28.3	13.1	23.3	10.1	0.6	100.0	73
Samburu	7.9	83.2	16.8	18.9	9.7	42.4	12.2	0.0	100.0	31
Trans Nzoia	6.1	77.2	22.8	29.8	5.2	28.1	14.0	0.0	100.0	57
Uasin Gishu	19.0	91.3	8.7	11.5	8.5	44.2	27.1	0.0	100.0	123
Elgeyo-Marakwet	12.4	91.3	8.7	62.4	0.5	15.2	11.7	1.5	100.0	32
Nandi	12.3	69.7	30.3	20.2	10.2	20.3	19.0	0.0	100.0	75
Baringo	42.6	91.8	8.2	19.5	11.2	51.4	8.3	1.4	100.0	53
Laikipia	11.1	89.3	10.7	10.1	6.8	54.6	16.6	1.2	100.0	50
Nakuru	8.5	95.4	4.6	9.8	10.3	45.0	30.2	0.0	100.0	222
Narok	13.1	89.1	10.9	20.6	13.4	40.6	14.5	0.0	100.0	87
Kajiado	18.3	94.1	5.9	13.0	9.9	39.3	31.9	0.0	100.0	123
Kericho	8.4	87.9	12.1	23.7	22.3	29.6	12.3	0.0	100.0	68
Bomet	6.6	97.3	2.7	18.9	13.7	45.7	18.9	0.0	100.0	89
Kakamega	21.5	88.2	11.8	12.6	8.8	46.0	20.8	0.0	100.0	126
Vihiga	10.1	94.8	5.2	7.2	4.6	48.1	27.6	7.3	100.0	38
Bungoma	15.5	79.4	20.6	17.2	6.3	34.8	21.1	0.0	100.0	128
Busia	26.6	90.3	9.7	15.6	6.7	35.0	26.3	6.7	100.0	78
Siaya	28.8	84.0	16.0	35.9	10.0	20.4	17.7	0.0	100.0	53
Kisumu	30.3	98.0	2.0	12.7	9.9	53.0	22.5	0.0	100.0	101
Homa Bay	33.8	94.1	5.9	10.3	7.2	46.2	30.3	0.0	100.0	81
Migori	53.4	90.1	9.9	9.2	5.6	32.8	42.4	0.0	100.0	131
Kisii	18.3	92.2	7.8	11.3	13.6	47.9	19.4	0.0	100.0	84
Nyamira	11.3	80.6	19.4	3.9	8.3	38.1	30.3	0.0	100.0	40
Nairobi City	30.5	84.9	15.1	10.9	8.1	31.6	33.3	1.0	100.0	577
Total	26.8	88.9	11.1	14.5	10.1	37.8	25.9	0.5	100.0	3,983
STILLBIRTHS										
Total	12.4	79.0	21.0	5.8	15.3	46.9	10.5	0.4	100.0	69
LIVE BIRTHS AND STILLBIRTHS[2]										
Total	26.7	88.9	11.1	14.4	10.1	38.1	25.7	0.5	100.0	4,035

Note: Figures in parentheses are based on 25–49 unweighted cases.
[1] Iron tablets, iron syrups, or iron and folic acid supplements (IFAS)
[2] For women who had both a live birth and a stillbirth in the 2 years preceding the survey, data are tabulated for the most recent birth only.

Table 9.5 Source of iron-containing supplements

Among women age 15–49 who had a live birth and/or stillbirth in the 2 years preceding the survey who were given or bought iron-containing supplements during the pregnancy for the most recent live birth or stillbirth, percentage who obtained supplements, according to source, Kenya Mini DHS 2025–26

	Percentage who obtained iron-containing supplements[1] from each source		
Source	Live births	Stillbirths	Live births and stillbirths[2]
Public sector	75.9	79.6	75.9
Government hospital	35.8	37.9	35.8
Government health centre	18.2	32.8	18.4
Government dispensary	23.4	8.8	23.2
Private medical sector (non-NGO)	24.7	20.4	24.6
Private hospital	13.1	10.0	13.0
Private clinic	1.8	3.0	1.8
Pharmacy	10.7	7.4	10.6
Private medical sector (NGO)	0.2	0.0	0.2
Private medical sector (FBO)	1.8	0.0	1.8
FBO/mission hospital	1.6	0.0	1.5
FBO/mission clinic	0.2	0.0	0.2
Other private sector	0.6	0.0	0.6
Shop	0.5	0.0	0.5
Number of women	3,570	54	3,618

Note: Supplements may have been obtained from more than one source.
NGO = nongovernmental organisation
FBO = faith-based organisation
[1] Iron tablets, iron syrups, or iron and folic acid supplements (IFAS)
[2] For women who had both a live birth and a stillbirth in the 2 years preceding the survey, data are tabulated for the most recent birth only.

Table 9.6 Tetanus toxoid injections

Among women age 15–49 with a live birth in the 2 years preceding the survey, percentage receiving two or more tetanus toxoid injections during the pregnancy for the most recent live birth and percentage whose last live birth was protected against neonatal tetanus, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Percentage receiving two or more injections during the pregnancy for the last live birth	Percentage whose most recent live birth was protected against neonatal tetanus[1]	Number of women
Age at birth			
<20	60.2	71.3	492
20–34	44.3	80.9	2,943
35–49	24.6	71.9	548
Birth order[2]			
1	72.0	78.8	1,269
2–3	34.1	81.3	1,724
4–5	25.1	78.0	685
6+	20.2	62.3	305
Residence			
Urban	47.9	82.0	1,698
Rural	40.3	75.9	2,285
Education[3]			
No education	23.2	64.1	350
Primary	30.9	74.8	1,076
Secondary	49.8	78.9	1,756
More than secondary	55.9	88.9	800
Wealth quintile			
Lowest	32.7	70.3	836
Second	40.8	75.9	745
Middle	46.1	78.6	720
Fourth	45.5	79.6	914
Highest	53.3	88.6	768
Total	43.6	78.5	3,983

[1] Includes women with two injections during the pregnancy for the most recent live birth, two or more injections (the last within 3 years of the most recent live birth), three or more injections (the last within 5 years of the most recent live birth), four or more injections (the last within 10 years of the most recent live birth), or five or more injections at any time prior to the most recent live birth

[2] Birth order refers to the order of the birth among the respondent's live births.

[3] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 9.6C Tetanus toxoid injections: County level

Among women age 15–49 with a live birth in the 2 years preceding the survey, percentage receiving two or more tetanus toxoid injections during the pregnancy for the most recent live birth and percentage whose last live birth was protected against neonatal tetanus, according to county, Kenya Mini DHS 2025–26

County	Percentage receiving two or more injections during the pregnancy for the last live birth	Percentage whose most recent live birth was protected against neonatal tetanus[1]	Number of women
Mombasa	51.2	74.0	127
Kwale	57.3	73.6	71
Kilifi	64.2	94.7	102
Tana River	26.7	78.6	40
Lamu	43.5	79.2	10
Taita Taveta	40.9	88.6	30
Garissa	8.7	49.9	42
Wajir	25.8	68.8	36
Mandera	13.6	65.4	39
Marsabit	24.9	69.1	35
Isiolo	23.1	58.9	16
Meru	35.6	77.9	72
Tharaka-Nithi	(58.0)	(78.0)	22
Embu	(68.1)	(95.7)	27
Kitui	48.5	85.8	64
Machakos	31.3	73.8	156
Makueni	50.1	94.5	59
Nyandarua	41.7	81.1	48
Nyeri	57.0	89.1	42
Kirinyaga	(39.4)	(73.5)	37
Murang'a	61.1	94.7	59
Kiambu	48.8	81.7	269
Turkana	19.7	79.9	62
West Pokot	12.7	64.3	73
Samburu	23.6	63.1	31
Trans Nzoia	27.7	68.0	57
Uasin Gishu	43.5	75.5	123
Elgeyo-Marakwet	36.6	64.6	32
Nandi	47.0	69.7	75
Baringo	34.1	57.5	53
Laikipia	52.4	81.8	50
Nakuru	51.6	78.1	222
Narok	23.1	77.5	87
Kajiado	47.3	92.8	123
Kericho	40.4	86.3	68
Bomet	30.1	65.3	89
Kakamega	42.3	87.3	126
Vihiga	36.0	93.6	38
Bungoma	41.9	79.3	128
Busia	43.1	69.4	78
Siaya	37.8	53.1	53
Kisumu	57.0	84.2	101
Homa Bay	54.0	80.1	81
Migori	42.4	61.9	131
Kisii	47.6	71.4	84
Nyamira	47.0	71.0	40
Nairobi City	49.9	86.3	577
Total	43.6	78.5	3,983

Note: Figures in parentheses are based on 25–49 unweighted cases.
[1] Includes women with two injections during the pregnancy for the most recent live birth, two or more injections (the last within 3 years of the most recent live birth), three or more injections (the last within 5 years of the most recent live birth), four or more injections (the last within 10 years of the most recent live birth), or five or more injections at any time prior to the most recent live birth

Table 9.7 Place of delivery

Percent distribution of live births and/or stillbirths in the 2 years preceding the survey by place of delivery, and percentage delivered in a health facility, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Health facility								
	Public sector	Private medical sector (non-NGO)	Private medical sector (NGO)	Private medical sector (FBO)	Home	Other	Total	Percentage delivered in a health facility	Number of births
				LIVE BIRTHS					
Mother's age at birth									
<20	76.2	13.0	0.0	2.2	7.8	0.8	100.0	91.4	501
20–34	65.5	24.0	0.3	2.3	7.7	0.2	100.0	92.1	3,053
35–49	60.5	21.0	0.0	4.2	14.0	0.3	100.0	85.7	562
Birth order[1]									
1	71.5	21.8	0.5	2.5	3.5	0.2	100.0	96.2	1,306
2–3	65.6	26.0	0.1	2.3	5.8	0.2	100.0	93.9	1,787
4–5	61.7	20.0	0.0	3.5	14.6	0.2	100.0	85.2	705
6+	56.8	8.9	0.0	1.8	31.6	1.0	100.0	67.4	318
Antenatal care visits[2]									
None	24.3	14.8	0.0	1.4	58.0	1.5	100.0	40.5	80
1–3	67.0	13.6	0.2	2.9	15.7	0.6	100.0	83.7	1,012
4+	66.4	26.0	0.2	2.5	4.6	0.2	100.0	95.2	2,884
Don't know/missing	*	*	*	*	*	*	100.0	*	7
Residence									
Urban	62.9	33.8	0.4	1.6	1.2	0.1	100.0	98.7	1,755
Rural	68.5	13.7	0.0	3.3	14.1	0.4	100.0	85.5	2,360
Mother's education[3]									
No education	42.1	13.0	0.1	0.6	43.9	0.2	100.0	55.8	363
Primary	70.2	14.2	0.2	2.7	12.2	0.5	100.0	87.3	1,111
Secondary	74.8	18.9	0.4	2.5	3.1	0.3	100.0	96.6	1,817
More than secondary	51.8	44.6	0.0	3.3	0.3	0.0	100.0	99.7	825
Wealth quintile									
Lowest	62.8	5.6	0.0	1.8	29.1	0.6	100.0	70.3	873
Second	74.4	14.9	0.0	2.4	7.7	0.5	100.0	91.7	770
Middle	76.4	15.8	0.2	3.9	3.6	0.1	100.0	96.3	739
Fourth	71.4	24.5	0.1	2.5	1.3	0.2	100.0	98.5	945
Highest	45.5	51.3	0.7	2.4	0.1	0.0	100.0	99.9	788
Total	66.1	22.3	0.2	2.5	8.6	0.3	100.0	91.1	4,115
				STILLBIRTHS					
Total	56.1	23.0	0.0	1.5	19.4	0.0	100.0	80.6	69
				LIVE BIRTHS AND STILLBIRTHS[4]					
Total	65.9	22.3	0.2	2.5	8.8	0.3	100.0	90.9	4,184

Note: An asterisk indicates that a figure is based on fewer than 25 unweighted cases and has been suppressed.
NGO = nongovernmental organisation
FBO = faith-based organisation
[1] Birth order refers to the order of the birth among the respondent's live births.
[2] Includes only the most recent birth in the 2 years preceding the survey
[3] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.
[4] For women who had both a live birth and a stillbirth in the 2 years preceding the survey, data are tabulated for the most recent birth only.

Table 9.7C Place of delivery: County level

Percent distribution of live births and/or stillbirths in the 2 years preceding the survey by place of delivery, and percentage delivered in a health facility, according to county, Kenya Mini DHS 2025–26

County	Health facility								
	Public sector	Private medical sector (non-NGO)	Private medical sector (NGO)	Private medical sector (FBO)	Home	Other	Total	Percentage delivered in a health facility	Number of births
LIVE BIRTHS									
Mombasa	71.0	25.5	1.1	0.0	2.4	0.0	100.0	97.6	131
Kwale	85.9	6.1	0.0	0.0	8.0	0.0	100.0	92.0	71
Kilifi	74.4	14.4	0.0	0.0	11.2	0.0	100.0	88.8	105
Tana River	70.1	4.9	0.0	0.0	25.0	0.0	100.0	75.0	42
Lamu	75.1	4.5	0.0	0.0	20.5	0.0	100.0	79.5	11
Taita Taveta	75.3	14.1	0.0	0.0	8.7	2.0	100.0	89.3	31
Garissa	39.1	7.1	1.0	0.0	52.9	0.0	100.0	47.1	46
Wajir	51.2	0.9	0.0	0.0	47.8	0.0	100.0	52.2	40
Mandera	55.9	2.8	0.0	0.0	41.3	0.0	100.0	58.7	42
Marsabit	61.3	14.3	0.0	6.2	18.1	0.0	100.0	81.9	35
Isiolo	69.2	18.0	0.0	2.0	10.8	0.0	100.0	89.2	17
Meru	55.3	18.9	0.0	14.5	11.3	0.0	100.0	88.7	72
Tharaka-Nithi	(59.0)	(16.8)	(0.0)	(11.9)	(10.3)	(1.9)	100.0	(87.8)	24
Embu	(57.9)	(32.6)	(0.0)	(7.6)	(2.0)	(0.0)	100.0	(98.0)	29
Kitui	82.7	9.5	0.0	0.0	6.3	1.5	100.0	92.2	66
Machakos	78.4	19.0	0.0	0.0	2.6	0.0	100.0	97.4	159
Makueni	80.9	14.6	1.6	0.0	1.5	1.5	100.0	97.1	62
Nyandarua	84.4	14.9	0.0	0.7	0.0	0.0	100.0	100.0	48
Nyeri	77.9	10.7	0.0	11.5	0.0	0.0	100.0	100.0	42
Kirinyaga	(65.3)	(30.3)	(0.0)	(4.5)	(0.0)	(0.0)	100.0	(100.0)	38
Murang'a	78.8	13.9	0.0	4.0	3.3	0.0	100.0	96.7	60
Kiambu	47.4	50.8	0.0	0.0	1.7	0.0	100.0	98.3	272
Turkana	47.0	2.7	0.0	2.1	47.3	1.0	100.0	51.8	63
West Pokot	63.4	3.5	0.0	0.0	33.2	0.0	100.0	66.8	76
Samburu	59.6	5.7	0.0	3.0	31.1	0.6	100.0	68.2	33
Trans Nzoia	43.1	35.2	0.6	6.9	14.2	0.0	100.0	85.8	59
Uasin Gishu	87.1	11.9	0.0	0.0	1.0	0.0	100.0	99.0	124
Elgeyo-Marakwet	79.1	4.4	0.0	11.6	4.9	0.0	100.0	95.1	33
Nandi	68.3	17.7	0.0	3.1	10.9	0.0	100.0	89.1	77
Baringo	61.0	11.3	0.0	9.6	15.9	2.2	100.0	81.9	55
Laikipia	59.3	18.5	0.0	4.8	17.4	0.0	100.0	82.6	51
Nakuru	72.7	23.6	0.0	0.0	3.7	0.0	100.0	96.3	224
Narok	65.0	5.7	0.0	2.1	27.2	0.0	100.0	72.8	91
Kajiado	49.3	31.9	0.0	4.0	12.5	2.2	100.0	85.3	130
Kericho	70.9	14.2	0.0	5.7	9.2	0.0	100.0	90.8	70
Bomet	68.6	16.3	0.0	11.2	2.9	1.0	100.0	96.0	89
Kakamega	77.2	10.6	0.0	4.8	7.4	0.0	100.0	92.6	130
Vihiga	80.1	6.8	0.0	6.4	6.8	0.0	100.0	93.2	38
Bungoma	66.7	20.7	0.0	2.2	9.6	0.8	100.0	89.6	131
Busia	62.2	27.2	0.0	2.2	7.1	1.2	100.0	91.6	78
Siaya	81.8	10.6	0.0	0.0	7.6	0.0	100.0	92.4	55
Kisumu	71.2	24.5	0.0	0.0	3.3	1.0	100.0	95.7	106
Homa Bay	75.7	12.0	0.0	3.0	9.3	0.0	100.0	90.7	86
Migori	65.7	24.0	0.0	3.4	6.5	0.3	100.0	93.2	139
Kisii	74.0	16.7	0.0	3.9	5.5	0.0	100.0	94.5	84
Nyamira	85.3	10.3	0.0	2.1	2.3	0.0	100.0	97.7	42
Nairobi City	56.8	40.5	0.9	1.8	0.0	0.0	100.0	100.0	611
Total	66.1	22.3	0.2	2.5	8.6	0.3	100.0	91.1	4,115
STILLBIRTHS									
Total	56.1	23.0	0.0	1.5	19.4	0.0	100.0	80.6	69
LIVE BIRTHS AND STILLBIRTHS[1]									
Total	65.9	22.3	0.2	2.5	8.8	0.3	100.0	90.9	4,184

Note: Figures in parentheses are based on 25–49 unweighted cases.
NGO = nongovernmental organisation
FBO = faith-based organisation
[1] For women who had both a live birth and a stillbirth in the 2 years preceding the survey, data are tabulated for the most recent birth only.

Table 9.8 Caesarean section

Percentage of live births and/or stillbirths in the 2 years preceding the survey delivered by caesarean section, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Percentage delivered by caesarean section	Number of births
LIVE BIRTHS		
Mother's age at birth		
<20	11.5	501
20–34	20.9	3,053
35–49	24.8	562
Birth order[1]		
1	23.9	1,306
2–3	21.6	1,787
4–5	16.0	705
6+	7.2	318
Antenatal care visits[2]		
None	1.7	80
1–3	14.7	1,012
4+	22.2	2,884
Don't know/missing	*	7
Place of delivery		
Health facility	*	3,749
Public sector	18.2	2,719
Private medical sector (non-NGO)	33.1	917
Private medical sector (NGO)	*	9
Private medical sector (FBO)	29.2	104
Residence		
Urban	29.1	1,755
Rural	13.7	2,360
Mother's education[3]		
No education	14.2	363
Primary	12.4	1,111
Secondary	19.5	1,817
More than secondary	35.2	825
Wealth quintile		
Lowest	8.8	873
Second	12.7	770
Middle	18.1	739
Fourth	21.4	945
Highest	41.0	788
Total	20.3	4,115
STILLBIRTHS		
Total	29.5	69
LIVE BIRTHS AND STILLBIRTHS[4]		
Total	20.4	4,184

Notes: The question on caesarean section is asked only of women who delivered in a health facility. In this table, it is assumed that women who did not give birth in a health facility did not receive a caesarean section. Stillbirths are foetal deaths in pregnancies lasting 28 or more weeks. When pregnancy duration is reported in months, stillbirths are foetal deaths in pregnancies lasting 7 or more months. An asterisk indicates that a figure is based on fewer than 25 unweighted cases and has been suppressed.
NGO = nongovernmental organisation
FBO = faith-based organisation
[1] Birth order refers to the order of the birth among the respondent's live births.
[2] Includes only the most recent birth in the 2 years preceding the survey
[3] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.
[4] For women who had both a live birth and a stillbirth in the 2 years preceding the survey, data are tabulated for the most recent birth only.

Table 9.8C Caesarean section: County level

Percentage of live births and/or stillbirths in the 2 years preceding the survey delivered by caesarean section, according to county, Kenya Mini DHS 2025–26

County	Percentage delivered by caesarean section	Number of births
LIVE BIRTHS		
Mombasa	23.9	131
Kwale	9.3	71
Kilifi	8.7	105
Tana River	8.5	42
Lamu	27.1	11
Taita Taveta	15.4	31
Garissa	8.1	46
Wajir	4.7	40
Mandera	1.9	42
Marsabit	10.6	35
Isiolo	19.8	17
Meru	24.8	72
Tharaka-Nithi	(33.7)	24
Embu	(43.0)	29
Kitui	19.3	66
Machakos	22.1	159
Makueni	30.2	62
Nyandarua	23.0	48
Nyeri	36.9	42
Kirinyaga	(37.5)	38
Murang'a	35.6	60
Kiambu	38.6	272
Turkana	10.0	63
West Pokot	11.2	76
Samburu	9.8	33
Trans Nzoia	5.8	59
Uasin Gishu	10.4	124
Elgeyo-Marakwet	12.2	33
Nandi	20.1	77
Baringo	14.4	55
Laikipia	16.9	51
Nakuru	8.5	224
Narok	13.6	91
Kajiado	26.3	130
Kericho	21.5	70
Bomet	14.4	89
Kakamega	13.5	130
Vihiga	13.5	38
Bungoma	13.2	131
Busia	12.8	78
Siaya	13.2	55
Kisumu	22.2	106
Homa Bay	19.0	86
Migori	14.3	139
Kisii	21.5	84
Nyamira	10.1	42
Nairobi City	30.7	611
Total	20.3	4,115
STILLBIRTHS		
Total	29.5	69
LIVE BIRTHS AND STILLBIRTHS[1]		
Total	20.4	4,184

Notes: The question on caesarean section is asked only of women who delivered in a health facility. In this table, it is assumed that women who did not give birth in a health facility did not receive a caesarean section. Stillbirths are foetal deaths in pregnancies lasting 28 or more weeks. When pregnancy duration is reported in months, stillbirths are foetal deaths in pregnancies lasting 7 or more months. Figures in parentheses are based on 25–49 unweighted cases.

[1] For women who had both a live birth and a stillbirth in the 2 years preceding the survey, data are tabulated for the most recent birth only.

Table 9.9 Assistance during delivery

Percent distribution of live births and/or stillbirths in the 2 years preceding the survey by person providing assistance during delivery, and percentage assisted by a skilled provider; and among live births in the 2 years preceding the survey, percentage with skin-to-skin contact immediately after birth, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Person providing assistance during delivery					Total	Percentage delivered by a skilled provider[1]	Number of live births and/or stillbirths	Among most recent live births[5]: Percentage with skin-to-skin contact immediately after birth	Among most recent live births[5]: Number of live births
	Doctor	Nurse/ midwife/ clinical officer	Traditional birth attendant	Relative/ friend/other	No one					
LIVE BIRTHS										
Mother's age at birth										
<20	48.3	43.9	5.6	1.9	0.3	100.0	92.2	501	73.2	492
20–34	55.3	37.5	4.2	2.2	0.8	100.0	92.8	3,053	70.7	2,943
35–49	52.9	33.7	8.6	2.8	2.0	100.0	86.6	562	65.4	548
Birth order[2]										
1	58.4	38.2	2.2	1.0	0.2	100.0	96.7	1,306	72.5	1,269
2–3	56.5	37.7	3.5	1.8	0.5	100.0	94.3	1,787	70.2	1,724
4–5	48.7	38.0	8.1	3.3	1.9	100.0	86.7	705	69.6	685
6+	35.3	34.8	18.0	7.5	4.4	100.0	70.1	318	62.6	305
Antenatal care visits[3]										
None	28.8	14.6	42.7	10.2	3.6	100.0	43.4	80	50.4	80
1–3	42.3	43.1	9.1	3.8	1.6	100.0	85.5	1,012	67.4	1,012
4+	59.2	36.3	2.4	1.5	0.6	100.0	95.6	2,884	71.9	2,884
Don't know/missing	*	*	*	*	*	100.0	*	7	*	7
Place of delivery										
Health facility	59.2	40.6	0.0	0.0	0.1	100.0	99.9	3,749	74.4	3,630
Public sector	55.5	44.4	0.1	0.1	0.1	100.0	99.8	2,719	77.1	2,618
Private medical sector (non-NGO)	69.8	30.2	0.0	0.0	0.0	100.0	100.0	917	68.1	899
Private medical sector (NGO)	*	*	*	*	*	100.0	*	9	*	9
Private medical sector (FBO)	62.6	37.4	0.0	0.0	0.0	100.0	100.0	104	66.0	104
Elsewhere	2.2	7.8	55.6	24.5	9.9	100.0	10.0	366	27.2	352
Residence										
Urban	62.8	36.1	0.5	0.3	0.3	100.0	98.9	1,755	70.9	1,698
Rural	47.7	38.9	8.3	3.7	1.4	100.0	86.6	2,360	69.7	2,285
Mother's education[4]										
No education	30.7	28.4	28.5	10.4	2.0	100.0	59.1	363	47.5	350
Primary	50.1	38.2	6.6	2.9	2.2	100.0	88.3	1,111	70.8	1,076
Secondary	56.6	40.4	1.5	1.2	0.3	100.0	97.0	1,817	75.2	1,756
More than secondary	64.6	35.2	0.2	0.0	0.0	100.0	99.8	825	68.6	800
Wealth quintile										
Lowest	38.6	33.5	17.8	7.6	2.5	100.0	72.1	873	61.4	836
Second	50.8	42.1	4.4	1.5	1.2	100.0	92.9	770	74.6	745
Middle	52.3	44.8	1.3	1.3	0.4	100.0	97.0	739	78.4	720
Fourth	56.8	41.8	0.6	0.4	0.4	100.0	98.6	945	72.8	914
Highest	73.3	26.6	0.1	0.1	0.0	100.0	99.9	788	65.0	768
Total	54.2	37.7	5.0	2.2	0.9	100.0	91.9	4,115	70.3	3,983
STILLBIRTHS										
Total	56.2	24.4	4.1	14.4	0.9	100.0	80.6	69	na	na
LIVE BIRTHS AND STILLBIRTHS[5]										
Total	54.2	37.5	5.0	2.4	0.9	100.0	91.7	4,184	na	na

Notes: If the respondent mentioned more than one person attending during delivery, only the most qualified person is considered in this tabulation. An asterisk indicates that a figure is based on fewer than 25 unweighted cases and has been suppressed.
NGO = nongovernmental organisation
FBO = faith-based organisation
na = not applicable
[1] Skilled provider is a doctor, nurse, midwife, or clinical officer.
[2] Birth order refers to the order of the birth among the respondent's live births.
[3] Includes only the most recent birth in the 2 years preceding the survey
[4] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.
[5] For women who had both a live birth and a stillbirth in the 2 years preceding the survey, data are tabulated for the most recent birth only.

Table 9.9C Assistance during delivery: County level

Percent distribution of live births and/or stillbirths in the 2 years preceding the survey by person providing assistance during delivery, and percentage assisted by a skilled provider; and among live births in the 2 years preceding the survey, percentage with skin-to-skin contact immediately after birth, according to county, Kenya Mini DHS 2025–26

	Person providing assistance during delivery								Among most recent live births[2]:	
County	Doctor	Nurse/ midwife/ clinical officer	Traditional birth attendant	Relative/ friend/other	No one	Total	Percentage delivered by a skilled provider[1]	Number of live births and/or stillbirths	Percentage with skin-to-skin contact immediately after birth	Number of live births
LIVE BIRTHS										
Mombasa	76.8	22.1	0.0	1.1	0.0	100.0	98.9	131	78.1	127
Kwale	85.7	6.3	1.8	2.5	3.7	100.0	92.0	71	70.9	71
Kilifi	50.5	38.1	7.4	0.8	3.2	100.0	88.6	105	84.8	102
Tana River	11.6	64.1	22.1	2.3	0.0	100.0	75.6	42	76.6	40
Lamu	39.4	42.2	5.5	10.9	2.1	100.0	81.6	11	62.8	10
Taita Taveta	64.8	26.5	3.0	3.6	2.1	100.0	91.3	31	66.5	30
Garissa	31.3	17.5	31.8	19.4	0.0	100.0	48.8	46	34.7	42
Wajir	22.0	33.6	43.1	1.3	0.0	100.0	55.6	40	24.4	36
Mandera	2.1	57.2	40.7	0.0	0.0	100.0	59.3	42	83.9	39
Marsabit	45.5	40.4	13.2	0.9	0.0	100.0	85.9	35	83.6	35
Isiolo	34.5	55.0	5.1	4.0	1.4	100.0	89.5	17	74.2	16
Meru	64.7	26.4	4.9	3.9	0.0	100.0	91.1	72	47.5	72
Tharaka-Nithi	(68.8)	(20.9)	(2.8)	(7.5)	(0.0)	100.0	(89.7)	24	(50.1)	22
Embu	(65.4)	(34.6)	(0.0)	(0.0)	(0.0)	100.0	(100.0)	29	(56.6)	27
Kitui	82.7	9.5	1.6	4.7	1.5	100.0	92.2	66	78.9	64
Machakos	41.3	57.6	1.1	0.0	0.0	100.0	98.9	159	69.0	156
Makueni	72.0	26.5	0.0	1.5	0.0	100.0	98.5	62	71.8	59
Nyandarua	81.1	18.9	0.0	0.0	0.0	100.0	100.0	48	70.6	48
Nyeri	58.6	41.4	0.0	0.0	0.0	100.0	100.0	42	67.1	42
Kirinyaga	(82.4)	(17.6)	(0.0)	(0.0)	(0.0)	100.0	(100.0)	38	(50.8)	37
Murang'a	62.4	34.3	0.0	3.3	0.0	100.0	96.7	60	83.9	59
Kiambu	78.1	20.5	0.6	0.8	0.0	100.0	98.6	272	73.9	269
Turkana	29.6	26.3	15.2	26.6	2.4	100.0	55.9	63	54.5	62
West Pokot	20.4	52.7	27.0	0.0	0.0	100.0	73.0	76	89.9	73
Samburu	29.3	38.9	12.7	16.5	2.6	100.0	68.2	33	49.1	31
Trans Nzoia	41.6	52.3	3.1	3.0	0.0	100.0	93.9	59	73.9	57
Uasin Gishu	46.6	52.4	1.0	0.0	0.0	100.0	99.0	124	71.2	123
Elgeyo-Marakwet	22.7	72.4	1.5	1.5	1.9	100.0	95.1	33	84.1	32
Nandi	45.6	44.6	4.1	3.4	2.3	100.0	90.2	77	59.2	75
Baringo	36.8	45.1	12.5	4.7	1.0	100.0	81.9	55	58.9	53
Laikipia	55.4	28.7	8.6	3.0	4.2	100.0	84.1	51	71.5	50
Nakuru	65.1	31.8	0.8	0.0	2.3	100.0	96.9	224	82.4	222
Narok	26.8	46.5	22.0	3.3	1.5	100.0	73.3	91	30.5	87
Kajiado	57.9	28.0	9.7	4.1	0.5	100.0	85.8	130	51.5	123
Kericho	18.6	73.1	2.5	3.2	2.6	100.0	91.7	70	49.8	68
Bomet	60.5	35.5	0.0	1.9	2.0	100.0	96.0	89	63.0	89
Kakamega	55.6	36.6	2.2	5.6	0.0	100.0	92.2	130	81.4	126
Vihiga	51.4	40.0	1.8	5.6	1.2	100.0	91.4	38	84.3	38
Bungoma	64.1	25.5	5.9	1.9	2.5	100.0	89.6	131	86.2	128
Busia	62.0	30.8	6.0	0.0	1.2	100.0	92.8	78	76.2	78
Siaya	54.3	38.1	5.8	0.0	1.7	100.0	92.4	55	81.9	53
Kisumu	47.2	49.5	2.4	0.0	0.9	100.0	96.7	106	71.5	101
Homa Bay	20.8	70.0	7.3	0.0	1.9	100.0	90.7	86	73.1	81
Migori	42.9	50.6	2.9	1.0	2.6	100.0	93.5	139	82.3	131
Kisii	26.7	67.8	1.6	3.8	0.0	100.0	94.5	84	71.2	84
Nyamira	45.2	52.5	1.3	1.0	0.0	100.0	97.7	42	82.9	40
Nairobi City	64.8	35.2	0.0	0.0	0.0	100.0	100.0	611	68.2	577
Total	54.2	37.7	5.0	2.2	0.9	100.0	91.9	4,115	70.3	3,983
STILLBIRTHS										
Total	56.2	24.4	4.1	14.4	0.9	100.0	80.6	69	na	na
LIVE BIRTHS AND STILLBIRTHS[2]										
Total	54.2	37.5	5.0	2.4	0.9	100.0	91.7	4,184	na	na

Notes: If the respondent mentioned more than one person attending during delivery, only the most qualified person is considered in this tabulation. Figures in parentheses are based on 25–49 unweighted cases.
na = not applicable
[1] Skilled provider is a doctor, nurse, midwife, or clinical officer.
[2] For women who had both a live birth and a stillbirth in the 2 years preceding the survey, data are tabulated for the most recent birth only.

Table 9.10 Duration of stay in a health facility after birth

Among women with a live birth and/or stillbirth in the 2 years preceding the survey who delivered their most recent live birth in a health facility, percent distribution by duration of stay in the health facility following their most recent live birth, according to type of delivery, Kenya Mini DHS 2025–26

Type of delivery	<6 hours	6–11 hours	12–23 hours	1–2 days	3+ days	Missing	Total	Number of women
LIVE BIRTHS								
Vaginal birth	6.4	6.4	5.2	59.5	22.5	0.0	100.0	2,837
Caesarean section	2.5	0.3	0.2	9.9	87.1	0.0	100.0	794
STILLBIRTHS								
Vaginal birth	(7.9)	(4.4)	(0.0)	(73.5)	(14.2)	(0.0)	100.0	35
Caesarean section	*	*	*	*	*	*	100.0	20
LIVE BIRTHS AND STILLBIRTHS[1]								
Vaginal birth	6.4	6.4	5.1	59.6	22.4	0.0	100.0	2,868
Caesarean section	2.4	0.3	0.2	9.8	87.2	0.0	100.0	802

Note: Figures in parentheses are based on 25–49 unweighted cases. An asterisk indicates that a figure is based on fewer than 25 unweighted cases and has been suppressed.

[1] For women who had both a live birth and a stillbirth in the 2 years preceding the survey, data are tabulated for the most recent birth only.

Table 9.11 Timing of first postnatal check for the mother

Among women age 15–49 with a live birth and/or stillbirth in the 2 years preceding the survey, percent distribution of the mother's first postnatal check for the most recent live birth or stillbirth by time after delivery, and percentage who received a postnatal check in the first 2 days after giving birth, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Time after delivery of mother's first postnatal check[1]						No postnatal check[2]	Total	Percentage of women with a postnatal check during the first 2 days after birth[1]	Number of women
	<4 hours	4–23 hours	1–2 days	3–6 days	7–41 days	Don't know/ missing				
					LIVE BIRTHS					
Age at birth										
<20	58.7	14.9	7.2	0.8	1.2	0.6	16.5	100.0	80.8	492
20–34	62.3	15.7	7.8	1.0	1.9	0.4	10.9	100.0	85.8	2,943
35–49	63.8	10.5	7.4	1.4	1.8	0.4	14.7	100.0	81.7	548
Birth order[3]										
1	61.2	18.2	7.5	1.1	1.9	0.5	9.6	100.0	86.8	1,269
2–3	63.2	14.9	8.6	0.8	1.7	0.4	10.4	100.0	86.7	1,724
4–5	65.1	11.2	7.5	1.3	1.2	0.4	13.4	100.0	83.7	685
6+	53.0	9.2	3.4	0.9	3.6	0.1	29.8	100.0	65.6	305
Place of delivery										
Health facility	66.2	15.6	7.8	0.8	1.5	0.5	7.6	100.0	89.6	3,630
Elsewhere	19.8	6.9	6.6	3.2	4.9	0.0	58.6	100.0	33.3	352
Residence										
Urban	66.5	14.9	8.2	1.0	1.8	0.6	7.0	100.0	89.6	1,698
Rural	58.7	14.8	7.3	1.0	1.9	0.3	16.0	100.0	80.9	2,285
Education[4]										
No education	43.1	9.4	7.6	0.4	2.0	0.3	37.1	100.0	60.1	350
Primary	61.4	13.0	6.2	1.4	2.5	0.5	15.0	100.0	80.6	1,076
Secondary	64.1	17.2	7.2	1.0	1.6	0.4	8.5	100.0	88.5	1,756
More than secondary	66.8	14.7	10.6	0.8	1.3	0.5	5.3	100.0	92.0	800
Wealth quintile										
Lowest	51.5	13.3	4.7	0.9	1.7	0.3	27.7	100.0	69.5	836
Second	61.2	15.0	8.0	1.8	2.9	0.4	10.7	100.0	84.2	745
Middle	66.6	13.2	9.0	0.3	1.3	0.5	9.1	100.0	88.8	720
Fourth	65.2	15.8	7.6	1.8	2.0	0.6	6.9	100.0	88.7	914
Highest	66.3	16.9	9.3	0.1	1.3	0.3	5.7	100.0	92.6	768
Total	62.1	14.9	7.7	1.0	1.8	0.4	12.1	100.0	84.6	3,983
					STILLBIRTHS					
Total	49.9	5.8	18.4	3.1	14.4	0.0	8.5	100.0	74.0	69
					LIVE BIRTHS AND STILLBIRTHS[5]					
Total	62.0	14.8	7.6	1.0	2.1	0.4	12.1	100.0	84.4	4,035

[1] Includes women who received a check from a doctor, nurse/midwife, auxiliary midwife, community health worker/fieldworker, or traditional birth attendant

[2] Includes women who received a check after 41 days

[3] Birth order refers to the order of the birth among the respondent's live births.

[4] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

[5] For women who had both a live birth and a stillbirth in the 2 years preceding the survey, data are tabulated for the most recent birth only.

Table 9.11C Timing of first postnatal check for the mother: County level

Among women age 15–49 with a live birth and/or stillbirth in the 2 years preceding the survey, percent distribution of the mother's first postnatal check for the most recent live birth or stillbirth by time after delivery, and percentage who received a postnatal check in the first 2 days after giving birth, according to county, Kenya Mini DHS 2025–26

County	Time after delivery of mother's first postnatal check[1]								Percentage of women with a postnatal check during the first 2 days after birth[1]	Number of women
	<4 hours	4–23 hours	1–2 days	3–6 days	7–41 days	Don't know/ missing	No postnatal check[2]	Total		
LIVE BIRTHS										
Mombasa	79.1	4.6	8.4	0.0	1.4	4.1	2.4	100.0	92.1	127
Kwale	57.7	29.4	0.7	0.0	1.0	1.2	9.9	100.0	87.8	71
Kilifi	76.0	10.3	5.6	1.0	2.8	0.0	4.3	100.0	91.9	102
Tana River	75.8	11.0	3.7	0.0	5.0	0.0	4.4	100.0	90.6	40
Lamu	70.4	5.5	3.5	1.3	3.2	0.0	16.2	100.0	79.4	10
Taita Taveta	51.4	18.2	12.2	4.5	1.3	0.0	12.3	100.0	81.9	30
Garissa	20.5	9.8	2.4	0.0	0.3	0.0	67.0	100.0	32.7	42
Wajir	42.9	8.3	0.0	0.0	0.0	0.0	48.8	100.0	51.2	36
Mandera	32.4	6.4	1.8	0.0	0.0	0.0	59.4	100.0	40.6	39
Marsabit	64.2	16.9	2.4	0.0	2.4	0.9	13.4	100.0	83.4	35
Isiolo	72.5	12.0	0.0	0.0	0.0	0.0	15.5	100.0	84.5	16
Meru	57.8	12.8	10.2	2.4	0.7	0.0	16.0	100.0	80.8	72
Tharaka-Nithi	(33.9)	(22.6)	(16.2)	(0.0)	(3.5)	(0.0)	(23.8)	100.0	(72.7)	22
Embu	(84.7)	(3.0)	(3.2)	(0.0)	(0.0)	(0.0)	(9.2)	100.0	(90.8)	27
Kitui	51.6	28.2	5.5	0.0	1.7	0.0	13.0	100.0	85.3	64
Machakos	65.8	19.6	3.6	3.5	0.0	2.3	5.2	100.0	89.0	156
Makueni	56.5	23.4	15.7	0.0	0.0	0.0	4.4	100.0	95.6	59
Nyandarua	52.4	13.6	11.6	1.3	5.4	0.0	15.6	100.0	77.7	48
Nyeri	65.3	12.8	18.4	0.0	3.5	0.0	0.0	100.0	96.5	42
Kirinyaga	(63.1)	(20.9)	(3.1)	(1.7)	(3.7)	(2.1)	(5.4)	100.0	(87.1)	37
Murang'a	75.4	13.3	7.1	0.7	0.0	0.0	3.5	100.0	95.8	59
Kiambu	61.9	6.8	12.4	1.8	5.5	0.0	11.6	100.0	81.1	269
Turkana	48.4	11.3	2.4	1.8	3.5	1.3	31.3	100.0	62.0	62
West Pokot	55.4	1.4	6.2	0.0	0.3	0.0	36.7	100.0	63.0	73
Samburu	48.3	9.8	5.1	0.0	0.0	0.0	36.9	100.0	63.1	31
Trans Nzoia	48.3	18.9	2.2	2.2	4.6	1.5	22.3	100.0	69.4	57
Uasin Gishu	60.5	13.9	12.2	1.0	0.7	1.0	10.7	100.0	86.7	123
Elgeyo-Marakwet	67.9	15.5	7.0	0.3	3.4	0.0	5.8	100.0	90.5	32
Nandi	38.5	13.4	20.2	1.8	6.9	0.0	19.1	100.0	72.2	75
Baringo	77.1	4.5	1.6	0.0	0.0	1.4	15.4	100.0	83.2	53
Laikipia	61.5	13.6	7.2	0.0	7.0	1.6	9.2	100.0	82.2	50
Nakuru	61.4	19.7	7.8	1.0	1.7	0.0	8.4	100.0	88.9	222
Narok	56.5	8.9	5.1	0.0	0.0	0.0	29.5	100.0	70.5	87
Kajiado	43.1	25.5	5.4	1.3	2.3	0.0	22.4	100.0	74.1	123
Kericho	61.5	15.2	1.9	1.2	2.6	0.0	17.6	100.0	78.6	68
Bomet	67.9	16.3	7.9	0.0	0.0	0.0	8.0	100.0	92.0	89
Kakamega	56.5	16.2	5.7	4.8	4.2	0.0	12.6	100.0	78.4	126
Vihiga	37.7	36.0	15.3	3.3	0.0	0.9	6.9	100.0	88.9	38
Bungoma	53.1	19.9	10.9	0.4	0.8	0.0	14.9	100.0	83.9	128
Busia	70.0	10.3	11.0	0.0	0.9	0.0	7.8	100.0	91.3	78
Siaya	50.9	19.1	16.0	0.0	1.9	0.0	12.1	100.0	86.1	53
Kisumu	63.7	21.0	4.4	0.8	2.7	0.8	6.6	100.0	89.1	101
Homa Bay	68.0	17.3	6.9	0.0	2.0	0.0	5.9	100.0	92.2	81
Migori	67.1	22.2	8.0	0.0	0.7	0.0	1.9	100.0	97.3	131
Kisii	80.6	8.6	3.2	1.4	0.0	0.0	6.3	100.0	92.3	84
Nyamira	51.1	13.5	10.1	0.0	2.2	1.5	21.5	100.0	74.8	40
Nairobi City	73.1	13.7	7.7	0.8	0.6	0.0	4.1	100.0	94.5	577
Total	62.1	14.9	7.7	1.0	1.8	0.4	12.1	100.0	84.6	3,983
STILLBIRTHS										
Total	49.9	5.8	18.4	3.1	14.4	0.0	8.5	100.0	74.0	69
LIVE BIRTHS AND STILLBIRTHS[3]										
Total	62.0	14.8	7.6	1.0	2.1	0.4	12.1	100.0	84.4	4,035

Note: Figures in parentheses are based on 25–49 unweighted cases.
[1] Includes women who received a check from a doctor, nurse/midwife, auxiliary midwife, community health worker/fieldworker, or traditional birth attendant
[2] Includes women who received a check after 41 days
[3] For women who had both a live birth and a stillbirth in the 2 years preceding the survey, data are tabulated for the most recent birth only.

Table 9.12 Type of provider of first postnatal check for the mother

Among women age 15–49 with a live birth and/or stillbirth in the 2 years preceding the survey, percent distribution by type of provider of the mother's first postnatal health check during the 2 days after the most recent live birth, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Type of health provider of mother's first postnatal check			No postnatal check during the first 2 days after birth	Total	Number of women
	Doctor/nurse/ midwife/ clinical officer	Community health worker/fieldworker	Traditional birth attendant			
			LIVE BIRTHS			
Age at birth						
<20	78.9	0.4	1.5	19.2	100.0	492
20–34	84.6	0.2	1.0	14.2	100.0	2,943
35–49	78.9	0.1	2.7	18.3	100.0	548
Birth order[1]						
1	86.1	0.1	0.6	13.2	100.0	1,269
2–3	85.7	0.2	0.8	13.3	100.0	1,724
4–5	81.4	0.2	2.1	16.3	100.0	685
6+	59.8	0.2	5.6	34.4	100.0	305
Place of delivery						
Health facility	89.4	0.1	0.1	10.4	100.0	3,630
Elsewhere	18.3	1.4	13.6	66.7	100.0	352
Residence						
Urban	89.4	0.0	0.2	10.4	100.0	1,698
Rural	78.4	0.3	2.1	19.1	100.0	2,285
Education[2]						
No education	53.0	0.7	6.4	39.9	100.0	350
Primary	78.5	0.2	1.9	19.4	100.0	1,076
Secondary	87.9	0.2	0.4	11.5	100.0	1,756
More than secondary	91.9	0.0	0.2	8.0	100.0	800
Wealth quintile						
Lowest	64.5	0.6	4.4	30.5	100.0	836
Second	83.0	0.2	1.0	15.8	100.0	745
Middle	87.9	0.2	0.6	11.2	100.0	720
Fourth	88.3	0.0	0.4	11.3	100.0	914
Highest	92.6	0.0	0.0	7.4	100.0	768
Total	83.1	0.2	1.3	15.4	100.0	3,983
			STILLBIRTHS			
Total	72.3	0.0	1.7	26.0	100.0	69
			LIVE BIRTHS AND STILLBIRTHS[3]			
Total	82.9	0.2	1.3	15.6	100.0	4,035

[1] Birth order refers to the order of the birth among the respondent's live births.
[2] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.
[3] For women who had both a live birth and a stillbirth in the 2 years preceding the survey, data are tabulated for the most recent birth only.

Table 9.12C Type of provider of first postnatal check for the mother: County level

Among women age 15–49 with a live birth and/or stillbirth in the 2 years preceding the survey, percent distribution by type of provider of the mother's first postnatal health check during the 2 days after the most recent live birth, according to county, Kenya Mini DHS 2025–26

County	Type of health provider of mother's first postnatal check				No postnatal check during the first 2 days after birth	Total	Number of women
	Doctor/nurse/ midwife/ clinical officer	Community health worker/ fieldworker	Traditional birth attendant	Other			
			LIVE BIRTHS				
Mombasa	92.1	0.0	0.0	0.0	7.9	100.0	127
Kwale	87.8	0.0	0.0	0.0	12.2	100.0	71
Kilifi	90.5	0.0	1.4	0.0	8.1	100.0	102
Tana River	78.4	0.0	12.2	0.0	9.4	100.0	40
Lamu	79.4	0.0	0.0	0.0	20.6	100.0	10
Taita Taveta	78.7	0.0	3.2	0.0	18.1	100.0	30
Garissa	32.7	0.0	0.0	0.0	67.3	100.0	42
Wajir	43.8	0.0	7.4	0.0	48.8	100.0	36
Mandera	40.3	0.0	0.3	0.0	59.4	100.0	39
Marsabit	80.0	0.0	3.4	0.0	16.6	100.0	35
Isiolo	82.2	0.0	2.3	0.0	15.5	100.0	16
Meru	75.9	0.0	4.9	0.0	19.2	100.0	72
Tharaka-Nithi	(72.7)	(0.0)	(0.0)	(0.0)	(27.3)	100.0	22
Embu	(90.8)	(0.0)	(0.0)	(0.0)	(9.2)	100.0	27
Kitui	83.6	0.0	1.7	0.0	14.7	100.0	64
Machakos	89.0	0.0	0.0	0.0	11.0	100.0	156
Makueni	95.6	0.0	0.0	0.0	4.4	100.0	59
Nyandarua	77.7	0.0	0.0	0.0	22.3	100.0	48
Nyeri	96.5	0.0	0.0	0.0	3.5	100.0	42
Kirinyaga	(87.1)	(0.0)	(0.0)	(0.0)	(12.9)	100.0	37
Murang'a	95.8	0.0	0.0	0.0	4.2	100.0	59
Kiambu	81.1	0.0	0.0	0.0	18.9	100.0	269
Turkana	52.3	3.6	6.2	0.0	38.0	100.0	62
West Pokot	53.3	0.0	9.8	0.0	37.0	100.0	73
Samburu	61.7	0.0	1.5	0.0	36.9	100.0	31
Trans Nzoia	66.2	0.0	3.2	0.0	30.6	100.0	57
Uasin Gishu	86.7	0.0	0.0	0.0	13.3	100.0	123
Elgeyo-Marakwet	89.0	0.0	1.5	0.0	9.5	100.0	32
Nandi	72.2	0.0	0.0	0.0	27.8	100.0	75
Baringo	75.8	1.3	6.2	0.0	16.8	100.0	53
Laikipia	80.7	0.0	1.6	0.0	17.8	100.0	50
Nakuru	88.9	0.0	0.0	0.0	11.1	100.0	222
Narok	61.6	0.0	8.8	0.0	29.5	100.0	87
Kajiado	73.6	0.0	0.5	0.0	25.9	100.0	123
Kericho	77.4	0.0	1.2	0.0	21.4	100.0	68
Bomet	91.2	0.9	0.0	0.0	8.0	100.0	89
Kakamega	78.4	0.0	0.0	0.0	21.6	100.0	126
Vihiga	88.9	0.0	0.0	0.0	11.1	100.0	38
Bungoma	82.2	0.9	0.8	0.0	16.1	100.0	128
Busia	88.1	0.0	3.2	0.0	8.7	100.0	78
Siaya	81.6	1.7	2.8	0.0	13.9	100.0	53
Kisumu	89.1	0.0	0.0	0.0	10.9	100.0	101
Homa Bay	87.2	2.2	2.7	0.0	7.8	100.0	81
Migori	96.5	0.3	0.5	0.0	2.7	100.0	131
Kisii	90.7	0.0	1.6	0.0	7.7	100.0	84
Nyamira	74.8	0.0	0.0	0.0	25.2	100.0	40
Nairobi City	94.5	0.0	0.0	0.0	5.5	100.0	577
Total	83.1	0.2	1.3	0.0	15.4	100.0	3,983
			STILLBIRTHS				
Total	72.3	0.0	1.7	0.0	26.0	100.0	69
			LIVE BIRTHS AND STILLBIRTHS[1]				
Total	82.9	0.2	1.3	0.0	15.6	100.0	4,035

Note: Figures in parentheses are based on 25–49 unweighted cases.
[1] For women who had both a live birth and a stillbirth in the 2 years preceding the survey, data are tabulated for the most recent birth only.

Table 9.13 Content of postnatal care for the mother

Among women age 15–49 with a live birth and/or stillbirth in the 2 years preceding the survey, percentage for whom selected checks were performed during the first 2 days after the most recent birth, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Percentage for whom, during the first 2 days after the most recent birth, any health care provider:			Percentage with all three checks performed in the first 2 days after birth	Number of women
	Measured blood pressure	Discussed vaginal bleeding	Discussed family planning		
			LIVE BIRTHS		
Age at birth					
<20	71.5	70.6	62.5	53.0	492
20–34	79.5	76.2	68.6	62.7	2,943
35–49	78.9	77.6	69.3	61.9	548
Birth order[1]					
1	79.0	75.3	65.6	56.8	1,269
2–3	82.5	79.0	72.4	66.9	1,724
4–5	77.3	77.1	69.3	64.3	685
6+	55.4	55.9	49.1	43.0	305
Place of delivery					
Health facility	83.2	80.2	72.4	65.5	3,526
Public sector	81.6	79.2	72.6	65.0	2,618
Private medical sector (non-NGO)	87.6	83.3	72.4	67.5	899
Private medical sector (NGO)	*	*	*	*	9
Private medical sector (FBO)	89.8	77.0	61.6	56.8	104
Elsewhere	27.2	30.6	24.7	21.4	352
Residence					
Urban	85.6	83.2	74.1	67.7	1,698
Rural	73.1	70.1	63.4	56.7	2,285
Mother's education[2]					
No education	44.9	42.1	32.7	31.1	350
Primary	74.7	72.6	64.6	58.1	1,076
Secondary	82.8	79.1	71.5	63.7	1,756
More than secondary	88.5	87.0	80.0	74.1	800
Wealth quintile					
Lowest	59.4	57.3	50.7	45.2	836
Second	79.2	76.2	68.1	61.8	745
Middle	82.3	76.6	73.4	64.3	720
Fourth	85.5	83.9	75.0	68.5	914
Highest	86.3	84.5	73.0	67.6	768
Total	78.4	75.7	67.9	61.4	3,983
			STILLBIRTHS		
Total	91.6	78.8	66.3	65.9	69
			LIVE BIRTHS AND STILLBIRTHS[3]		
Total	78.6	75.7	67.8	61.4	4,035

Note: An asterisk indicates that a figure is based on fewer than 25 unweighted cases and has been suppressed.
NGO = nongovernmental organisation
FBO = faith-based organisation
[1] Birth order refers to the order of the birth among the respondent's live births.
[2] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.
[3] For women who had both a live birth and a stillbirth in the 2 years preceding the survey, data are tabulated for the most recent birth only.

Table 9.13C Content of postnatal care for the mother: County level

Among women age 15–49 with a live birth and/or stillbirth in the 2 years preceding the survey, percentage for whom selected checks were performed during the first 2 days after the most recent birth, according to county, Kenya Mini DHS 2025–26

County	Percentage for whom, during the first 2 days after the most recent birth, any health care provider:			Percentage with all three checks performed in the first 2 days after birth	Number of women
	Measured blood pressure	Discussed vaginal bleeding	Discussed family planning		
			LIVE BIRTHS		
Mombasa	93.0	87.7	90.1	83.4	127
Kwale	88.3	80.1	82.6	70.4	71
Kilifi	97.7	94.9	91.6	90.6	102
Tana River	72.4	73.7	54.9	51.9	40
Lamu	82.0	81.0	73.8	71.5	10
Taita Taveta	79.1	86.4	74.8	68.2	30
Garissa	8.2	6.2	5.0	4.6	42
Wajir	36.9	38.5	31.4	30.3	36
Mandera	32.2	25.9	13.3	13.3	39
Marsabit	80.4	78.7	58.8	58.8	35
Isiolo	76.4	64.8	53.5	50.2	16
Meru	74.8	66.1	51.1	46.4	72
Tharaka-Nithi	(60.3)	(37.3)	(36.7)	(28.0)	22
Embu	(79.0)	(76.0)	(78.2)	(73.1)	27
Kitui	79.2	72.7	77.5	68.1	64
Machakos	84.3	81.1	76.1	66.6	156
Makueni	92.6	93.1	94.1	82.7	59
Nyandarua	88.5	72.0	72.5	57.1	48
Nyeri	96.5	72.2	59.9	51.0	42
Kirinyaga	(91.9)	(90.8)	(71.4)	(67.4)	37
Murang'a	95.5	94.8	89.9	87.9	59
Kiambu	87.6	85.7	89.1	81.0	269
Turkana	34.0	37.2	29.4	24.2	62
West Pokot	21.8	20.7	19.6	19.6	73
Samburu	50.3	46.3	23.2	19.6	31
Trans Nzoia	79.4	76.4	71.7	66.3	57
Uasin Gishu	82.3	79.0	72.9	64.8	123
Elgeyo-Marakwet	56.3	56.4	51.6	47.1	32
Nandi	82.6	81.9	81.3	77.0	75
Baringo	57.8	61.1	34.7	29.6	53
Laikipia	79.5	78.3	70.8	68.3	50
Nakuru	76.1	77.6	67.2	58.1	222
Narok	63.2	67.5	57.7	51.0	87
Kajiado	76.0	74.4	66.1	63.9	123
Kericho	64.7	59.8	43.0	39.1	68
Bomet	91.9	66.8	55.2	47.4	89
Kakamega	88.2	85.8	84.5	74.0	126
Vihiga	87.0	85.2	72.4	67.5	38
Bungoma	74.9	73.5	64.6	59.6	128
Busia	90.3	90.4	78.0	74.6	78
Siaya	82.3	84.8	86.0	79.0	53
Kisumu	79.1	73.5	72.5	56.3	101
Homa Bay	78.3	79.6	69.8	66.6	81
Migori	70.7	72.9	67.1	54.1	131
Kisii	92.9	92.5	77.5	75.4	84
Nyamira	42.7	36.9	31.2	20.6	40
Nairobi City	87.1	85.0	68.1	62.4	577
Total	78.4	75.7	67.9	61.4	3,983
			STILLBIRTHS		
Total	91.6	78.8	66.3	65.9	69
			LIVE BIRTHS AND STILLBIRTHS[1]		
Total	78.6	75.7	67.8	61.4	4,035

Note: Figures in parentheses are based on 25–49 unweighted cases.
[1] For women who had both a live birth and a stillbirth in the 2 years preceding the survey, data are tabulated for the most recent birth only.

Table 9.14 Timing of first postnatal check for the newborn

Percent distribution of most recent live births in the 2 years preceding the survey by time after birth of first postnatal check, and percentage of those births with a postnatal check during the first 2 days after birth, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Time after delivery of newborn's first postnatal check[1]						No postnatal check[2]	Total	Percentage of births with a postnatal check during the first 2 days after birth[1]	Number of births
	<1 hour	1–3 hours	4–23 hours	1–2 days	3–6 days	Don't know				
Mother's age at birth										
<20	18.3	45.9	14.5	3.8	1.4	0.5	15.5	100.0	82.5	492
20–34	23.3	42.3	13.3	7.2	0.9	0.6	12.4	100.0	86.1	2,943
35–49	21.9	46.0	9.9	7.3	1.3	0.5	13.2	100.0	85.1	548
Birth order[3]										
1	23.0	43.6	15.7	5.3	1.0	0.6	10.8	100.0	87.5	1,269
2–3	24.0	41.4	13.2	8.8	0.8	0.4	11.4	100.0	87.4	1,724
4–5	23.0	46.3	9.5	6.2	1.6	0.8	12.7	100.0	85.0	685
6+	10.9	45.4	8.4	3.2	1.3	0.5	30.2	100.0	67.9	305
Place of delivery										
Health facility	23.8	46.0	13.5	6.7	0.8	0.6	8.6	100.0	90.1	3,630
Elsewhere	8.6	14.7	7.4	8.1	4.0	0.3	57.0	100.0	38.8	352
Residence										
Urban	26.3	41.6	12.7	8.0	1.0	0.5	10.0	100.0	88.6	1,698
Rural	19.6	44.5	13.2	5.9	1.1	0.6	15.0	100.0	83.3	2,285
Mother's education[4]										
No education	9.3	33.7	8.8	8.0	0.6	0.6	39.0	100.0	59.8	350
Primary	23.0	45.0	10.2	5.4	1.3	0.4	14.8	100.0	83.5	1,076
Secondary	23.9	44.1	15.2	6.8	0.8	0.8	8.5	100.0	89.9	1,756
More than secondary	24.4	43.3	13.9	8.3	1.4	0.3	8.4	100.0	89.9	800
Wealth quintile										
Lowest	14.1	40.3	11.7	4.6	1.2	0.9	27.2	100.0	70.7	836
Second	22.6	47.4	12.3	5.4	1.2	0.9	10.2	100.0	87.7	745
Middle	23.1	48.5	13.8	5.9	0.6	0.4	7.8	100.0	91.3	720
Fourth	23.3	44.9	13.0	8.5	1.5	0.5	8.3	100.0	89.6	914
Highest	30.0	35.6	14.3	9.4	0.6	0.0	10.2	100.0	89.3	768
Total	22.5	43.3	13.0	6.8	1.0	0.5	12.9	100.0	85.5	3,983

[1] Includes newborns who received a check from a doctor, nurse, midwife, clinical officer, community health worker/fieldworker, or traditional birth attendant

[2] Includes newborns who received a check after the first week of life

[3] Birth order refers to the order of the birth among the respondent's live births.

[4] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 9.14C Timing of first postnatal check for the newborn: County level

Percent distribution of most recent live births in the 2 years preceding the survey by time after birth of first postnatal check, and percentage of births with a postnatal check during the first 2 days after birth, according to county, Kenya Mini DHS 2025–26

County	Time after delivery of newborn's first postnatal check[1]						No postnatal check[2]	Total	Percentage of births with a postnatal check during the first 2 days after birth[1]	Number of births
	<1 hour	1–3 hours	4–23 hours	1–2 days	3–6 days	Don't know				
Mombasa	19.9	58.4	2.6	9.8	1.5	1.4	6.4	100.0	90.7	127
Kwale	19.8	52.0	19.4	0.8	2.2	1.0	4.8	100.0	91.9	71
Kilifi	53.3	23.2	7.6	7.8	1.8	0.0	6.4	100.0	91.8	102
Tana River	15.9	59.6	9.3	4.9	0.0	0.0	10.3	100.0	89.7	40
Lamu	25.2	49.7	4.0	4.5	1.3	0.0	15.3	100.0	83.4	10
Taita Taveta	33.9	27.6	15.3	10.1	0.0	2.9	10.2	100.0	87.0	30
Garissa	0.0	20.9	9.1	1.1	1.8	0.0	67.2	100.0	31.0	42
Wajir	0.0	52.8	8.7	1.5	0.0	0.0	37.1	100.0	62.9	36
Mandera	0.0	32.1	8.0	1.9	0.0	0.0	57.9	100.0	42.1	39
Marsabit	5.8	70.3	7.4	0.9	0.0	0.0	15.6	100.0	84.4	35
Isiolo	5.5	75.7	2.8	0.6	0.0	0.0	15.5	100.0	84.5	16
Meru	17.5	49.7	10.8	7.0	0.0	0.0	15.1	100.0	84.9	72
Tharaka-Nithi	(20.2)	(23.2)	(24.0)	(21.4)	(0.0)	(0.0)	(11.2)	100.0	(88.8)	22
Embu	(75.4)	(9.8)	(3.0)	(2.5)	(3.0)	(0.0)	(6.4)	100.0	(90.6)	27
Kitui	2.1	46.4	27.1	8.4	0.0	0.0	15.9	100.0	84.1	64
Machakos	24.2	49.9	12.6	6.0	2.8	2.3	2.3	100.0	92.7	156
Makueni	24.2	40.3	18.2	15.5	0.0	0.0	1.9	100.0	98.1	59
Nyandarua	19.1	55.9	9.1	11.8	0.0	0.0	4.1	100.0	95.9	48
Nyeri	8.2	59.0	12.8	16.5	0.0	0.0	3.5	100.0	96.5	42
Kirinyaga	(9.4)	(62.1)	(17.4)	(5.6)	(3.5)	(2.1)	(0.0)	100.0	(94.4)	37
Murang'a	5.4	70.1	14.5	5.9	0.0	0.8	3.4	100.0	95.9	59
Kiambu	36.0	32.3	4.9	11.7	2.5	0.0	12.6	100.0	84.9	269
Turkana	2.3	45.5	11.6	3.8	1.8	2.2	32.8	100.0	63.2	62
West Pokot	0.0	49.4	1.4	7.1	1.1	0.0	41.0	100.0	57.9	73
Samburu	14.5	35.0	11.2	4.8	0.0	0.0	34.6	100.0	65.4	31
Trans Nzoia	17.7	44.0	17.0	2.8	0.6	0.0	17.9	100.0	81.5	57
Uasin Gishu	22.3	43.4	15.8	9.3	0.8	0.0	8.6	100.0	90.7	123
Elgeyo-Marakwet	21.5	59.2	5.9	5.3	0.3	0.0	7.8	100.0	91.9	32
Nandi	12.4	24.4	10.7	12.0	5.3	0.0	35.1	100.0	59.6	75
Baringo	51.6	25.2	4.4	3.0	1.0	0.0	14.9	100.0	84.1	53
Laikipia	2.2	66.4	12.6	1.3	0.0	2.3	15.1	100.0	82.6	50
Nakuru	15.9	49.3	19.5	6.9	0.0	0.0	8.3	100.0	91.7	222
Narok	27.8	37.5	7.9	3.9	0.0	0.0	22.9	100.0	77.1	87
Kajiado	8.9	49.3	18.3	6.5	0.4	0.0	16.6	100.0	82.9	123
Kericho	17.7	57.6	17.2	1.9	0.0	0.0	5.6	100.0	94.4	68
Bomet	4.1	64.8	16.1	6.8	0.0	1.1	7.1	100.0	91.8	89
Kakamega	39.2	31.0	11.9	3.4	3.7	0.0	10.7	100.0	85.6	126
Vihiga	15.8	28.5	29.0	12.7	2.7	5.6	5.8	100.0	86.0	38
Bungoma	26.7	33.2	17.5	8.6	0.8	0.0	13.2	100.0	86.0	128
Busia	65.0	14.1	10.8	5.7	0.0	0.0	4.5	100.0	95.5	78
Siaya	20.4	38.8	16.8	10.9	0.0	0.0	13.1	100.0	86.9	53
Kisumu	37.3	33.1	17.8	2.5	1.2	1.8	6.3	100.0	90.7	101
Homa Bay	13.8	52.6	16.6	6.1	1.1	0.0	9.7	100.0	89.1	81
Migori	7.9	62.3	20.7	3.5	0.0	0.7	4.8	100.0	94.5	131
Kisii	21.2	55.8	6.0	3.8	1.4	5.1	6.7	100.0	86.9	84
Nyamira	5.0	56.4	14.9	4.1	0.0	3.0	16.5	100.0	80.5	40
Nairobi City	29.2	36.3	13.6	7.4	0.6	0.0	12.9	100.0	86.5	577
Total	22.5	43.3	13.0	6.8	1.0	0.5	12.9	100.0	85.5	3,983

Note: Figures in parentheses are based on 25–49 unweighted cases.
[1] Includes newborns who received a check from a doctor, nurse, midwife, clinical officer, community health worker/fieldworker, or traditional birth attendant
[2] Includes newborns who received a check after the first week of life

Table 9.15 Type of provider of first postnatal check for the newborn

Percent distribution of most recent live births in the 2 years preceding the survey by type of provider of the newborn's first postnatal health check during the 2 days after the most recent live birth, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Type of health provider of newborn's first postnatal check			No postnatal check during the first 2 days after birth	Total	Number of births
	Doctor/nurse/ midwife/ clinical officer	Community health worker/ fieldworker	Traditional birth attendant			
Mother's age at birth						
<20	80.5	0.4	1.7	17.5	100.0	492
20–34	84.6	0.2	1.3	13.9	100.0	2,943
35–49	82.1	0.2	2.9	14.9	100.0	548
Birth order[1]						
1	86.8	0.1	0.6	12.5	100.0	1,269
2–3	85.9	0.2	1.2	12.6	100.0	1,724
4–5	82.3	0.2	2.5	15.0	100.0	685
6+	61.9	0.3	5.7	32.1	100.0	305
Place of delivery						
Health facility	89.9	0.1	0.1	9.9	100.0	3,630
Elsewhere	20.2	1.6	17.0	61.2	100.0	352
Residence						
Urban	88.6	0.0	0.0	11.4	100.0	1,698
Rural	80.2	0.4	2.8	16.7	100.0	2,285
Mother's education[2]						
No education	50.6	0.5	8.7	40.2	100.0	350
Primary	81.4	0.2	1.9	16.5	100.0	1,076
Secondary	89.0	0.2	0.7	10.1	100.0	1,756
More than secondary	89.9	0.0	0.0	10.1	100.0	800
Wealth quintile						
Lowest	64.6	0.5	5.7	29.3	100.0	836
Second	85.8	0.3	1.6	12.3	100.0	745
Middle	90.4	0.2	0.6	8.7	100.0	720
Fourth	89.6	0.0	0.0	10.4	100.0	914
Highest	89.3	0.0	0.0	10.7	100.0	768
Total	83.7	0.2	1.6	14.5	100.0	3,983

[1] Birth order refers to the order of the birth among the respondent's live births.
[2] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 9.15C Type of provider of first postnatal check for the newborn: County level

Percent distribution of most recent live births in the 2 years preceding the survey by type of provider of the newborn's first postnatal health check during the 2 days after the most recent live birth, according to county, Kenya Mini DHS 2025–26

County	Type of health provider of newborn's first postnatal check			No postnatal check during the first 2 days after birth	Total	Number of births
	Doctor/nurse/ midwife/ clinical officer	Community health worker/ fieldworker	Traditional birth attendant			
Mombasa	90.7	0.0	0.0	9.3	100.0	127
Kwale	91.9	0.0	0.0	8.1	100.0	71
Kilifi	90.7	0.0	1.2	8.2	100.0	102
Tana River	79.9	0.0	9.8	10.3	100.0	40
Lamu	80.9	0.0	2.4	16.6	100.0	10
Taita Taveta	87.0	0.0	0.0	13.0	100.0	30
Garissa	31.0	0.0	0.0	69.0	100.0	42
Wajir	45.0	0.0	17.9	37.1	100.0	36
Mandera	40.4	0.0	1.7	57.9	100.0	39
Marsabit	79.9	0.0	4.5	15.6	100.0	35
Isiolo	82.2	0.0	2.3	15.5	100.0	16
Meru	80.0	0.0	4.9	15.1	100.0	72
Tharaka-Nithi	(83.0)	(2.7)	(3.1)	(11.2)	100.0	22
Embu	(90.6)	(0.0)	(0.0)	(9.4)	100.0	27
Kitui	84.1	0.0	0.0	15.9	100.0	64
Machakos	92.7	0.0	0.0	7.3	100.0	156
Makueni	98.1	0.0	0.0	1.9	100.0	59
Nyandarua	95.9	0.0	0.0	4.1	100.0	48
Nyeri	96.5	0.0	0.0	3.5	100.0	42
Kirinyaga	(94.4)	(0.0)	(0.0)	(5.6)	100.0	37
Murang'a	95.9	0.0	0.0	4.1	100.0	59
Kiambu	84.9	0.0	0.0	15.1	100.0	269
Turkana	49.0	3.6	10.6	36.8	100.0	62
West Pokot	48.8	0.0	9.2	42.1	100.0	73
Samburu	65.0	0.0	0.4	34.6	100.0	31
Trans Nzoia	78.3	0.0	3.2	18.5	100.0	57
Uasin Gishu	90.7	0.0	0.0	9.3	100.0	123
Elgeyo-Marakwet	90.4	1.6	0.0	8.1	100.0	32
Nandi	59.6	0.0	0.0	40.4	100.0	75
Baringo	77.8	0.3	6.0	15.9	100.0	53
Laikipia	81.0	0.0	1.6	17.4	100.0	50
Nakuru	91.7	0.0	0.0	8.3	100.0	222
Narok	66.6	0.0	10.5	22.9	100.0	87
Kajiado	82.1	0.0	0.8	17.1	100.0	123
Kericho	90.4	0.0	4.0	5.6	100.0	68
Bomet	91.0	0.9	0.0	8.2	100.0	89
Kakamega	84.5	0.0	1.0	14.4	100.0	126
Vihiga	84.2	1.8	0.0	14.0	100.0	38
Bungoma	84.2	0.0	1.8	14.0	100.0	128
Busia	89.9	0.0	5.6	4.5	100.0	78
Siaya	83.2	1.7	2.0	13.1	100.0	53
Kisumu	90.7	0.0	0.0	9.3	100.0	101
Homa Bay	84.2	2.2	2.7	10.9	100.0	81
Migori	94.2	0.3	0.0	5.5	100.0	131
Kisii	85.2	0.0	1.6	13.1	100.0	84
Nyamira	80.5	0.0	0.0	19.5	100.0	40
Nairobi City	86.5	0.0	0.0	13.5	100.0	577
Total	83.7	0.2	1.6	14.5	100.0	3,983

Note: Figures in parentheses are based on 25–49 unweighted cases.

Table 9.16 Content of postnatal care for the newborn

Among most recent live births in the 2 years preceding the survey, percentage for whom selected functions were performed during the first 2 days after the birth and percentage with at least two signal functions performed during the first 2 days after the birth, according to background characteristics, Kenya Mini DHS 2025–26

	Percentage of most recent live births for whom a health care provider performed selected functions during the first 2 days after the birth									
Background characteristic	Examined the cord	Measured temper-ature	Told mother how to recognise if the baby needs immediate medical attention	Counselled mother on breast-feeding	Observed breast-feeding	Both counselled mother on breast-feeding and observed breast-feeding	Weighed[1]	Showed how to clean the cord	Percent-age with all five[2] signal functions performed during the first 2 days after birth	Number of births
Mother's age at birth										
<20	81.0	78.9	71.4	83.4	81.2	79.1	89.6	74.5	62.7	492
20–34	86.2	82.6	76.2	85.5	84.9	82.8	92.5	81.5	70.8	2,943
35–49	84.1	79.0	71.2	80.8	82.1	77.7	86.9	77.2	63.2	548
Birth order[3]										
1	86.9	85.2	77.1	89.9	88.6	87.1	95.7	83.4	70.8	1,269
2–3	87.4	84.4	77.9	86.3	85.6	83.7	94.2	82.5	72.9	1,724
4–5	84.2	78.3	71.7	80.2	81.9	76.9	87.0	77.1	65.7	685
6+	68.7	58.6	55.7	62.6	61.5	58.2	67.6	58.4	43.2	305
Place of delivery										
Health facility	88.8	86.3	79.0	88.8	88.5	86.2	97.8	84.1	74.0	3,630
Elsewhere	49.3	33.8	32.7	41.0	38.6	35.5	25.1	37.9	14.7	352
Residence										
Urban	90.9	89.9	81.9	91.1	90.2	88.4	98.3	88.7	78.1	1,698
Rural	81.1	75.4	69.6	79.7	79.6	76.7	86.2	73.5	61.8	2,285
Mother's education[4]										
No education	56.3	48.4	42.5	52.0	49.5	47.3	53.9	49.8	32.0	350
Primary	84.2	79.0	70.9	81.0	81.1	77.8	88.1	77.0	63.1	1,076
Secondary	88.6	85.8	79.1	89.9	89.4	87.4	97.0	83.7	73.8	1,756
More than secondary	92.2	90.5	85.1	91.9	91.6	89.3	99.8	89.2	81.3	800
Wealth quintile										
Lowest	70.6	60.4	57.0	65.6	65.2	61.7	70.8	60.8	46.9	836
Second	84.6	81.5	72.9	84.3	84.7	81.3	92.5	76.3	66.2	745
Middle	86.7	85.0	79.2	89.0	88.5	86.7	96.7	82.3	72.6	720
Fourth	91.0	90.8	82.5	91.8	90.4	89.9	98.1	88.6	78.2	914
Highest	93.8	90.8	83.0	92.6	92.3	89.2	99.6	92.1	80.1	768
Total	85.3	81.6	74.9	84.6	84.1	81.7	91.4	80.0	68.7	3,983

[1] Captures newborns who were weighed 'at birth'; may exclude some newborns who were weighed during the first 2 days after birth
[2] As described by Moran et al. (2013), the five signal functions for assessing the content of postnatal care for newborns are (1) examining the umbilical cord, (2) measuring temperature, (3) observing and/or counselling on breastfeeding, (4) telling the mother about danger signs/how to recognise if the baby needs immediate attention, and (5) weighing.
[3] Birth order refers to the order of the birth among the respondent's live births.
[4] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 9.16C Content of postnatal care for the newborn: County level

Among most recent live births in the 2 years preceding the survey, percentage for whom selected functions were performed during the first 2 days after the birth and percentage with at least two signal functions performed during the first 2 days after birth, according to county, Kenya Mini DHS 2025–26

	Percentage of most recent live births for whom a health care provider performed selected functions during the first 2 days after the birth									
County	Examined the cord	Measured temper-ature	Told mother how to recognise if the baby needs immediate medical attention	Counselled mother on breast-feeding	Observed breast-feeding	Both counselled mother on breast-feeding and observed breast-feeding	Weighed[1]	Showed how to clean the cord	Percent-age with five[2] signal functions performed during the first 2 days after birth	Number of births
Mombasa	95.6	97.1	96.7	94.5	95.6	94.5	92.3	94.6	90.5	127
Kwale	92.9	91.6	76.3	92.7	93.9	91.0	98.0	88.8	72.8	71
Kilifi	96.6	96.9	93.7	96.6	94.5	94.5	97.8	91.3	92.7	102
Tana River	83.9	71.3	61.0	76.5	37.3	36.3	71.6	74.0	41.7	40
Lamu	87.6	84.5	84.0	86.3	87.6	86.3	91.0	87.6	79.8	10
Taita Taveta	94.1	86.5	79.8	89.9	85.2	81.5	94.2	87.7	69.3	30
Garissa	8.3	9.1	8.1	6.9	6.9	6.9	47.3	7.5	5.4	42
Wajir	49.0	38.5	41.0	47.3	46.0	44.8	31.6	44.5	23.8	36
Mandera	56.3	42.4	23.6	22.1	18.1	17.1	55.9	21.2	16.6	39
Marsabit	86.8	85.8	77.4	81.6	83.7	81.6	81.2	76.8	68.7	35
Isiolo	76.5	67.6	56.5	69.1	65.7	63.2	83.9	71.7	51.3	16
Meru	86.5	73.5	65.2	81.6	80.6	77.0	86.5	70.2	54.2	72
Tharaka-Nithi	(80.9)	(84.1)	(63.5)	(82.0)	(80.3)	(75.9)	(88.8)	(72.1)	(53.2)	22
Embu	(89.0)	(91.2)	(86.0)	(88.5)	(88.5)	(88.5)	(97.9)	(91.5)	(81.2)	27
Kitui	90.7	92.1	85.3	92.1	93.8	92.1	89.0	90.7	76.3	64
Machakos	89.3	89.5	74.4	90.8	90.3	88.8	97.7	91.0	68.6	156
Makueni	98.1	94.0	96.7	97.7	95.8	95.8	98.4	99.0	90.9	59
Nyandarua	92.2	87.5	80.5	96.4	98.1	96.4	100.0	89.4	70.1	48
Nyeri	96.8	86.1	79.8	98.4	96.1	96.1	100.0	96.1	74.7	42
Kirinyaga	(98.3)	(97.9)	(93.1)	(93.0)	(96.5)	(93.0)	(100.0)	(91.2)	(91.0)	37
Murang'a	96.2	93.6	88.8	92.5	91.9	90.0	98.5	94.4	86.7	59
Kiambu	95.4	92.4	88.1	95.7	93.9	93.1	98.5	94.5	85.0	269
Turkana	61.2	42.2	53.7	59.4	60.0	56.1	46.7	55.8	28.8	62
West Pokot	27.3	19.9	21.0	26.0	26.3	24.6	66.3	25.4	16.3	73
Samburu	54.3	49.1	41.5	56.7	60.7	55.6	63.1	38.7	29.7	31
Trans Nzoia	85.6	76.3	80.9	90.2	89.4	89.4	90.5	75.2	70.1	57
Uasin Gishu	90.3	82.6	78.4	91.7	91.0	90.2	99.0	79.0	70.6	123
Elgeyo-Marakwet	61.5	58.3	57.1	64.1	60.8	59.3	94.0	58.1	48.0	32
Nandi	89.5	87.2	87.2	87.3	88.5	87.3	93.2	81.6	77.2	75
Baringo	70.3	67.6	64.3	65.2	68.8	65.2	79.7	67.5	58.4	53
Laikipia	83.1	81.5	78.0	78.9	79.6	78.2	86.3	78.4	76.8	50
Nakuru	85.6	82.2	78.6	85.5	85.2	83.7	97.4	81.9	74.8	222
Narok	75.9	55.4	63.1	72.1	75.3	68.4	72.8	55.3	44.1	87
Kajiado	84.2	78.7	72.1	78.3	83.4	78.3	85.0	79.8	67.9	123
Kericho	80.4	73.0	71.9	76.9	75.3	71.6	92.9	65.3	62.2	68
Bomet	90.4	70.7	59.6	80.9	84.8	77.9	94.1	61.7	50.3	89
Kakamega	87.9	85.8	72.0	89.6	89.4	86.2	95.0	83.2	68.3	126
Vihiga	88.2	87.8	88.5	96.5	90.4	90.4	97.1	84.3	81.8	38
Bungoma	79.9	76.4	61.8	77.2	83.8	75.3	91.3	74.0	57.0	128
Busia	90.5	90.1	85.1	90.8	93.0	89.8	95.2	91.2	75.5	78
Siaya	97.2	91.0	92.9	98.1	98.1	98.1	85.4	91.2	79.0	53
Kisumu	88.8	87.1	69.2	92.6	89.4	86.4	94.9	76.5	62.8	101
Homa Bay	88.3	85.0	80.0	87.1	85.9	84.5	91.8	81.9	78.9	81
Migori	83.3	81.9	65.8	86.5	84.8	79.6	92.6	76.1	61.6	131
Kisii	93.2	90.8	89.7	95.9	90.8	90.8	97.4	90.1	86.8	84
Nyamira	42.5	38.3	32.8	47.3	44.0	42.7	96.7	34.2	26.3	40
Nairobi City	92.5	94.3	82.2	93.6	92.9	90.6	100.0	92.5	79.1	577
Total	85.3	81.6	74.9	84.6	84.1	81.7	91.4	80.0	68.7	3,983

Note: Figures in parentheses are based on 25–49 unweighted cases.
[1] Captures newborns who were weighed 'at birth'; may exclude some newborns who were weighed during the first 2 days after birth
[2] As described by Moran et al. (2013), the five signal functions for assessing the content of postnatal care for newborns are (1) examining the umbilical cord, (2) measuring temperature, (3) observing and/or counselling on breastfeeding, (4) telling the mother about danger signs/how to recognise if the baby needs immediate attention, and (5) weighing.

Table 9.17 Postnatal health checks for mothers and newborns

Among most recent live births in the 2 years preceding the survey, percentage for which the mother age 15–49 received a postnatal check during the first 2 days after birth, percentage for which the newborn received a postnatal check during the first 2 days after birth, percentage for which both the mother and the newborn received a postnatal check, and percentage for which neither the mother nor the newborn received a postnatal check, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Percentage who received a postnatal check[1] during the first 2 days after birth				Number of births
	Mother	Newborn	Both mother and newborn	Neither mother nor newborn[2]	
Mother's age at birth					
<20	80.8	82.5	77.4	14.0	492
20–34	85.8	86.1	81.3	9.5	2,943
35–49	81.7	85.1	80.0	13.1	548
Birth order[3]					
1	86.8	87.5	83.0	8.6	1,269
2–3	86.7	87.4	82.4	8.4	1,724
4–5	83.7	85.0	80.1	11.5	685
6+	65.6	67.9	62.3	28.7	305
Place of delivery					
Health facility	89.6	90.1	85.6	5.9	3,630
Public sector	88.6	90.1	85.0	6.3	2,618
Private medical sector (non-NGO)	92.5	89.6	87.3	5.1	899
Private medical sector (NGO)	*	*	*	*	9
Private medical sector (FBO)	87.2	93.4	85.1	4.4	104
Elsewhere	33.3	38.8	29.9	57.9	352
Residence					
Urban	89.6	88.6	84.5	6.3	1,698
Rural	80.9	83.3	77.8	13.7	2,285
Mother's education[4]					
No education	60.1	59.8	53.9	34.1	350
Primary	80.6	83.5	77.5	13.4	1,076
Secondary	88.5	89.9	85.5	7.0	1,756
More than secondary	92.0	89.9	86.0	4.1	800
Wealth quintile					
Lowest	69.5	70.7	65.4	25.2	836
Second	84.2	87.7	81.3	9.4	745
Middle	88.8	91.3	86.6	6.5	720
Fourth	88.7	89.6	83.7	5.4	914
Highest	92.6	89.3	87.4	5.6	768
Total	84.6	85.5	80.7	10.5	3,983

Note: An asterisk indicates that a figure is based on fewer than 25 unweighted cases and has been suppressed.
NGO = nongovernmental organisation
FBO = faith-based organisation
[1] Includes checks from a doctor, nurse, midwife, clinical officer, community health worker/fieldworker, or traditional birth attendant
[2] Includes checks after the first 2 days or by other persons
[3] Birth order refers to the order of the birth among the respondent's live births.
[4] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 9.17C Postnatal health checks for mothers and newborns: County level

Among most recent live births in the 2 years preceding the survey, percentage for which the mother age 15–49 received a postnatal check during the first 2 days after birth, percentage for which the newborn received a postnatal check during the first 2 days after birth, percentage for which both the mother and the newborn received a postnatal check, and percentage for which neither the mother nor the newborn received a postnatal check, according to county, Kenya Mini DHS 2025–26

	Percentage who received a postnatal check[1] during the first 2 days after birth				
County	Mother	Newborn	Both mother and newborn	Neither mother nor newborn[2]	Number of births
Mombasa	92.1	90.7	84.2	1.4	127
Kwale	87.8	91.9	84.4	4.6	71
Kilifi	91.9	91.8	87.1	3.4	102
Tana River	90.6	89.7	87.3	7.1	40
Lamu	79.4	83.4	78.1	15.4	10
Taita Taveta	81.9	87.0	77.3	8.5	30
Garissa	32.7	31.0	29.7	65.9	42
Wajir	51.2	62.9	49.7	35.6	36
Mandera	40.6	42.1	35.2	52.5	39
Marsabit	83.4	84.4	78.3	10.6	35
Isiolo	84.5	84.5	84.0	14.9	16
Meru	80.8	84.9	79.1	13.3	72
Tharaka-Nithi	(72.7)	(88.8)	(69.8)	(8.3)	22
Embu	(90.8)	(90.6)	(87.9)	(6.4)	27
Kitui	85.3	84.1	82.1	12.7	64
Machakos	89.0	92.7	87.4	5.7	156
Makueni	95.6	98.1	95.6	1.9	59
Nyandarua	77.7	95.9	77.7	4.1	48
Nyeri	96.5	96.5	96.5	3.5	42
Kirinyaga	(87.1)	(94.4)	(83.6)	(2.1)	37
Murang'a	95.8	95.9	93.9	2.1	59
Kiambu	81.1	84.9	80.3	14.3	269
Turkana	62.0	63.2	58.3	33.2	62
West Pokot	63.0	57.9	55.4	34.4	73
Samburu	63.1	65.4	57.9	29.3	31
Trans Nzoia	69.4	81.5	67.8	16.9	57
Uasin Gishu	86.7	90.7	84.2	6.8	123
Elgeyo-Marakwet	90.5	91.9	89.0	6.7	32
Nandi	72.2	59.6	51.5	19.7	75
Baringo	83.2	84.1	77.9	10.5	53
Laikipia	82.2	82.6	80.2	15.3	50
Nakuru	88.9	91.7	88.1	7.5	222
Narok	70.5	77.1	68.9	21.4	87
Kajiado	74.1	82.9	73.6	16.6	123
Kericho	78.6	94.4	77.7	4.6	68
Bomet	92.0	91.8	91.0	7.1	89
Kakamega	78.4	85.6	76.8	12.9	126
Vihiga	88.9	86.0	82.2	7.3	38
Bungoma	83.9	86.0	82.6	12.8	128
Busia	91.3	95.5	90.1	3.3	78
Siaya	86.1	86.9	82.2	9.3	53
Kisumu	89.1	90.7	86.3	6.5	101
Homa Bay	92.2	89.1	88.2	6.9	81
Migori	97.3	94.5	94.5	2.7	131
Kisii	92.3	86.9	84.4	5.2	84
Nyamira	74.8	80.5	73.3	18.0	40
Nairobi City	94.5	86.5	85.1	4.1	577
Total	84.6	85.5	80.7	10.5	3,983

Note: Figures in parentheses are based on 25–49 unweighted cases.
[1] Includes checks from a doctor, nurse, midwife, clinical officer, community health worker/fieldworker, or traditional birth attendant
[2] Includes checks after the first 2 days or by other persons

Table 9.18 Examinations for breast and cervical cancers

Percentage of women age 15–49 ever examined by a doctor or health care worker for breast cancer and percentage ever tested by a doctor or health care worker for cervical cancer, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Percentage examined for breast cancer	Percentage tested for cervical cancer	Number of women
Age			
15–29	9.7	10.0	11,301
30–49	22.8	29.0	9,930
30–34	18.7	23.2	3,108
35–39	22.5	26.2	2,989
40–44	26.2	35.2	2,048
45–49	26.5	36.5	1,784
30–44	22.0	27.4	8,146
40–49	26.3	35.8	3,833
Number of living children			
0	7.5	7.8	6,186
1–2	18.6	21.3	7,685
3–4	22.0	27.7	4,983
5+	15.1	21.2	2,376
Marital status			
Never married	8.6	9.2	6,983
Married or living together	18.9	22.5	11,672
Divorced/separated/ widowed	21.4	28.7	2,576
Residence			
Urban	18.8	20.9	9,829
Rural	13.2	17.1	11,402
Education[1]			
No education	9.6	10.4	1,161
Primary	17.3	22.3	6,053
Secondary	12.0	14.4	9,849
More than secondary	24.2	26.9	4,167
Wealth quintile			
Lowest	9.3	11.2	3,312
Second	12.6	16.7	3,852
Middle	15.4	18.6	4,133
Fourth	17.5	21.1	4,800
Highest	21.0	23.6	5,134
Total	15.8	18.9	21,231

[1] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 9.18C Examinations for breast and cervical cancers: County level

Percentage of women age 15–49 ever examined by a doctor or health care worker for breast cancer and percentage ever tested by a doctor or health care worker for cervical cancer, according to county, Kenya Mini DHS 2025–26

County	Percentage examined for breast cancer	Percentage tested for cervical cancer	Number of women
Mombasa	20.2	17.2	803
Kwale	7.7	9.8	374
Kilifi	21.6	20.2	608
Tana River	2.6	3.4	137
Lamu	6.8	12.5	54
Taita Taveta	17.2	20.5	153
Garissa	1.9	2.7	168
Wajir	0.2	0.7	130
Mandera	0.0	0.2	123
Marsabit	3.2	4.2	142
Isiolo	11.7	13.1	76
Meru	14.0	16.2	530
Tharaka-Nithi	16.9	20.2	147
Embu	22.5	25.3	239
Kitui	16.0	20.3	412
Machakos	14.5	20.8	677
Makueni	14.9	17.9	350
Nyandarua	16.6	21.9	223
Nyeri	25.6	30.2	223
Kirinyaga	28.4	27.6	258
Murang'a	16.0	16.1	377
Kiambu	19.7	23.7	1,689
Turkana	11.1	13.6	178
West Pokot	4.0	2.6	225
Samburu	8.1	10.8	98
Trans Nzoia	14.5	16.1	370
Uasin Gishu	14.7	17.9	667
Elgeyo-Marakwet	10.3	15.2	178
Nandi	14.1	17.1	361
Baringo	8.5	9.0	187
Laikipia	19.0	23.0	195
Nakuru	21.1	22.6	1,078
Narok	13.7	12.3	428
Kajiado	21.0	15.2	467
Kericho	14.3	12.8	387
Bomet	9.6	8.8	480
Kakamega	12.3	18.1	665
Vihiga	10.3	16.1	233
Bungoma	9.8	12.9	625
Busia	15.7	21.0	373
Siaya	10.3	29.2	370
Kisumu	19.5	29.6	551
Homa Bay	17.1	23.1	485
Migori	14.8	27.4	509
Kisii	16.6	22.4	439
Nyamira	13.2	16.4	238
Nairobi City	17.5	20.5	3,550
Total	15.8	18.9	21,231

Table 9.19 Problems in accessing health care

Percentage of women age 15–49 who reported that they have serious problems accessing health care for themselves when they are sick, by type of problem, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Problem in accessing health care							
	Getting permission to go for treatment	Getting money for treatment	Distance to health facility	Not wanting to go alone	Religion	Culture	At least one problem accessing health care	Number of women
Age								
15–19	3.5	44.3	22.5	15.4	0.6	1.8	55.2	3,916
20–34	3.4	47.7	20.0	7.4	0.8	1.6	53.5	10,493
35–49	3.7	57.2	27.5	7.5	0.8	1.2	62.6	6,822
Number of living children								
0	3.3	41.4	17.8	12.6	0.8	1.9	50.3	6,186
1–2	3.4	47.2	19.8	7.3	0.6	1.2	53.3	7,685
3–4	3.3	55.9	26.1	7.2	0.7	1.3	61.5	4,983
5+	4.9	69.9	39.3	8.5	1.2	2.0	74.7	2,376
Marital status								
Never married	3.3	43.7	18.7	11.7	0.8	1.8	52.1	6,983
Married or living together	3.6	51.9	25.0	7.6	0.7	1.4	57.9	11,672
Divorced/separated/ widowed	3.9	59.5	24.4	7.3	0.8	1.2	64.3	2,576
Residence								
Urban	3.2	41.3	11.2	7.7	0.7	1.5	46.9	9,829
Rural	3.8	57.7	32.9	10.0	0.8	1.5	65.3	11,402
Education[1]								
No education	6.4	65.4	43.2	11.3	0.6	1.7	74.2	1,161
Primary	4.7	65.0	31.2	8.5	0.9	1.7	70.6	6,053
Secondary	3.0	48.4	20.8	10.4	0.7	1.3	55.3	9,849
More than secondary	2.3	28.3	10.2	5.6	0.7	1.7	35.1	4,167
Wealth quintile								
Lowest	4.7	73.7	49.3	11.8	1.0	2.2	80.9	3,312
Second	3.9	63.4	32.8	10.3	0.8	1.4	70.4	3,852
Middle	4.0	53.0	23.4	9.0	0.7	0.9	59.8	4,133
Fourth	3.6	44.7	13.9	7.7	0.7	1.6	51.3	4,800
Highest	2.1	27.6	6.3	7.1	0.6	1.5	33.5	5,134
Total	3.5	50.1	22.9	8.9	0.7	1.5	56.7	21,231

[1] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 9.19C Problems in accessing health care: County level

Percentage of women age 15–49 who reported that they have serious problems accessing health care for themselves when they are sick, by type of problem, according to county, Kenya Mini DHS 2025–26

	Problem in accessing health care							
County	Getting permission to go for treatment	Getting money for treatment	Distance to health facility	Not wanting to go alone	Religion	Culture	At least one problem accessing health care	Number of women
Mombasa	5.6	41.6	11.7	5.4	0.6	2.4	46.8	803
Kwale	4.2	60.7	36.7	5.7	0.4	1.5	67.5	374
Kilifi	3.2	63.0	32.3	8.6	1.3	2.7	71.6	608
Tana River	0.2	64.1	29.7	5.9	0.0	1.8	74.4	137
Lamu	3.0	80.5	28.2	14.2	1.7	2.0	84.9	54
Taita Taveta	2.5	50.4	25.3	13.8	0.8	0.7	58.7	153
Garissa	8.9	45.8	32.1	19.2	0.7	2.8	58.7	168
Wajir	0.4	76.6	39.8	3.2	1.0	3.0	77.9	130
Mandera	2.6	51.9	44.1	33.8	1.2	4.1	56.8	123
Marsabit	0.7	71.1	38.9	3.0	0.1	0.3	74.4	142
Isiolo	3.1	55.3	24.2	7.2	0.5	1.3	59.6	76
Meru	2.2	65.6	37.6	16.3	0.2	1.1	74.7	530
Tharaka-Nithi	8.1	63.1	45.6	29.8	3.4	0.8	74.7	147
Embu	8.6	45.5	24.4	10.8	0.9	3.6	55.2	239
Kitui	5.9	71.0	43.9	6.2	1.7	0.8	75.4	412
Machakos	1.2	25.7	23.4	19.4	0.0	0.2	38.0	677
Makueni	2.0	43.5	33.6	14.5	0.3	2.4	54.1	350
Nyandarua	7.0	58.4	38.7	11.7	1.3	2.6	69.3	223
Nyeri	2.5	39.9	15.5	16.7	0.8	0.6	50.1	223
Kirinyaga	7.9	46.6	19.2	7.2	1.0	1.7	54.6	258
Murang'a	0.1	31.7	14.5	4.4	0.0	0.6	36.6	377
Kiambu	0.7	31.7	8.6	4.4	0.2	0.9	36.6	1,689
Turkana	4.4	65.2	53.8	8.9	1.6	2.5	73.5	178
West Pokot	9.6	75.2	48.4	12.9	1.1	1.7	80.6	225
Samburu	3.1	54.1	30.8	10.1	0.3	3.9	64.9	98
Trans Nzoia	1.4	71.7	16.0	8.5	0.4	1.0	76.7	370
Uasin Gishu	3.0	39.0	7.9	3.5	0.2	0.2	41.9	667
Elgeyo-Marakwet	1.0	20.0	7.2	2.6	0.1	0.9	25.0	178
Nandi	1.6	34.2	16.1	5.2	0.4	1.2	40.5	361
Baringo	3.7	52.0	44.5	5.6	0.0	1.1	63.5	187
Laikipia	4.3	24.5	22.4	9.2	0.8	1.0	38.5	195
Nakuru	6.1	52.7	26.1	10.8	0.5	1.5	60.0	1,078
Narok	6.9	60.8	35.6	6.9	1.3	2.4	67.0	428
Kajiado	3.1	33.2	15.1	5.5	1.0	1.6	39.5	467
Kericho	2.4	24.3	9.4	9.1	0.7	0.9	30.0	387
Bomet	1.5	56.5	35.0	3.3	1.5	0.4	60.1	480
Kakamega	4.2	68.8	28.7	10.2	2.0	2.8	72.7	665
Vihiga	0.2	42.7	19.4	4.6	0.2	0.7	46.2	233
Bungoma	2.0	78.3	34.6	11.2	0.5	2.4	82.6	625
Busia	1.7	60.3	33.3	10.8	0.6	0.0	69.1	373
Siaya	2.1	61.2	32.3	2.9	0.6	1.8	68.6	370
Kisumu	13.3	60.6	28.7	16.9	0.7	0.8	70.7	551
Homa Bay	3.1	70.0	41.7	12.2	0.6	1.4	78.9	485
Migori	2.8	47.2	10.6	7.3	0.9	1.3	52.8	509
Kisii	2.0	71.1	30.8	10.4	0.3	1.1	75.7	439
Nyamira	5.8	63.7	24.3	3.9	2.0	3.1	71.2	238
Nairobi City	3.3	43.6	11.1	8.1	0.9	1.8	49.6	3,550
Total	3.5	50.1	22.9	8.9	0.7	1.5	56.7	21,231

Table 9.20 Distance from a health care facility

Percent distribution of women age 15–49 by travel time to nearest health facility and by means of transport to nearest health facility, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Travel time to nearest health facility					Means of transport to nearest health facility				Number of women
	<30 minutes	30–59 minutes	60–119 minutes	≥2 hours	Total	Motorised[1]	Not motorised[2]	Other	Total	
Age										
15–19	53.8	28.5	13.6	4.1	100.0	21.0	79.0	0.0	100.0	3,916
20–34	62.9	24.1	9.9	3.1	100.0	24.1	75.9	0.0	100.0	10,493
35–49	57.2	26.3	12.3	4.2	100.0	23.7	76.3	0.0	100.0	6,822
Accessing health care										
Distance to health facility is a problem	24.7	33.1	29.3	12.9	100.0	30.0	70.0	0.0	100.0	4,858
Distance to health facility is not a problem	69.7	23.4	6.0	0.9	100.0	21.4	78.6	0.0	100.0	16,373
Means of transport to nearest health facility										
Motorised[1]	56.1	29.3	12.1	2.5	100.0	na	na	na	na	4,964
Not motorised[2]	60.4	24.5	11.1	4.0	100.0	na	na	na	na	16,263
Other	*	*	*	*	100.0	na	na	na	na	4
Residence										
Urban	77.8	18.1	3.5	0.6	100.0	20.7	79.3	0.0	100.0	9,829
Rural	43.5	32.1	18.1	6.2	100.0	25.7	74.3	0.0	100.0	11,402
Education[3]										
No education	44.2	25.5	16.7	13.6	100.0	26.8	73.2	0.0	100.0	1,161
Primary	49.2	30.4	15.2	5.2	100.0	22.9	77.0	0.0	100.0	6,053
Secondary	59.0	26.8	11.4	2.8	100.0	22.2	77.8	0.0	100.0	9,849
More than secondary	79.5	15.9	4.1	0.5	100.0	25.8	74.2	0.0	100.0	4,167
Wealth quintile										
Lowest	29.8	30.9	25.3	13.9	100.0	22.2	77.8	0.0	100.0	3,312
Second	42.2	34.6	18.9	4.3	100.0	24.0	76.0	0.0	100.0	3,852
Middle	53.7	32.1	12.0	2.2	100.0	24.3	75.7	0.0	100.0	4,133
Fourth	70.6	23.8	4.9	0.7	100.0	21.9	78.1	0.0	100.0	4,800
Highest	85.5	12.0	2.2	0.3	100.0	24.3	75.6	0.0	100.0	5,134
Total	59.4	25.6	11.4	3.6	100.0	23.4	76.6	0.0	100.0	21,231

Note: An asterisk indicates that a figure is based on fewer than 25 unweighted cases and has been suppressed.
na = not applicable
[1] Includes car/truck, tuk tuk, public bus, motorcycle/scooter, and boat with motor
[2] Includes animal-drawn cart, bicycle, boat without motor, and walking
[3] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 9.20C Distance from a health care facility: County level

Percent distribution of women age 15–49 by travel time to nearest health facility and by means of transport to nearest health facility, according to county, Kenya Mini DHS 2025–26

	Travel time to nearest health facility					Means of transport to nearest health facility				
County	<30 minutes	30–59 minutes	60–119 minutes	≥2 hours	Total	Motorised[1]	Not motorised[2]	Other	Total	Number of women
Mombasa	70.1	23.9	4.9	1.1	100.0	38.6	61.4	0.0	100.0	803
Kwale	45.6	28.7	18.9	6.8	100.0	19.1	80.9	0.0	100.0	374
Kilifi	47.7	27.4	15.5	9.5	100.0	12.7	87.3	0.0	100.0	608
Tana River	74.2	18.1	6.7	0.9	100.0	31.2	68.8	0.0	100.0	137
Lamu	62.5	26.9	6.9	3.7	100.0	17.4	82.6	0.0	100.0	54
Taita Taveta	58.4	25.4	13.5	2.7	100.0	23.1	76.9	0.0	100.0	153
Garissa	61.7	27.9	9.6	0.8	100.0	44.9	55.1	0.0	100.0	168
Wajir	45.3	35.8	8.9	10.0	100.0	16.3	83.7	0.0	100.0	130
Mandera	50.1	28.7	14.5	6.7	100.0	20.8	79.2	0.0	100.0	123
Marsabit	62.6	27.2	4.6	5.5	100.0	20.1	79.9	0.0	100.0	142
Isiolo	64.1	17.3	12.8	5.7	100.0	5.8	94.2	0.0	100.0	76
Meru	54.1	26.4	15.6	3.8	100.0	36.0	64.0	0.0	100.0	530
Tharaka-Nithi	32.3	27.8	26.0	13.8	100.0	23.6	76.4	0.0	100.0	147
Embu	62.2	25.0	8.7	4.1	100.0	21.7	78.3	0.0	100.0	239
Kitui	26.2	25.1	24.6	24.2	100.0	19.0	81.0	0.0	100.0	412
Machakos	50.8	24.8	18.1	6.3	100.0	24.9	75.1	0.0	100.0	677
Makueni	45.9	40.6	11.9	1.6	100.0	21.6	78.4	0.0	100.0	350
Nyandarua	44.6	36.5	15.8	3.2	100.0	30.7	69.3	0.0	100.0	223
Nyeri	62.0	32.2	5.1	0.8	100.0	6.5	93.5	0.0	100.0	223
Kirinyaga	53.3	38.2	7.5	1.0	100.0	18.4	81.0	0.5	100.0	258
Murang'a	52.6	36.2	9.3	1.9	100.0	26.9	73.1	0.0	100.0	377
Kiambu	74.6	19.4	5.9	0.1	100.0	21.8	78.2	0.0	100.0	1,689
Turkana	31.0	24.6	18.0	26.4	100.0	27.3	72.7	0.0	100.0	178
West Pokot	54.2	30.2	8.4	7.2	100.0	36.7	63.3	0.0	100.0	225
Samburu	26.9	29.8	26.3	17.0	100.0	8.3	91.4	0.3	100.0	98
Trans Nzoia	60.9	30.8	7.4	0.9	100.0	43.3	56.7	0.0	100.0	370
Uasin Gishu	71.0	25.0	4.0	0.0	100.0	36.9	62.9	0.2	100.0	667
Elgeyo-Marakwet	42.3	38.8	17.3	1.6	100.0	16.9	83.1	0.0	100.0	178
Nandi	60.4	25.9	11.3	2.3	100.0	24.1	75.9	0.0	100.0	361
Baringo	30.3	24.1	25.3	20.3	100.0	8.9	91.1	0.0	100.0	187
Laikipia	41.2	30.8	24.9	3.2	100.0	24.2	75.8	0.0	100.0	195
Nakuru	59.7	27.8	10.3	2.2	100.0	21.8	78.2	0.0	100.0	1,078
Narok	34.1	24.3	33.8	7.8	100.0	39.1	60.9	0.0	100.0	428
Kajiado	65.4	21.0	9.4	4.1	100.0	21.0	79.0	0.0	100.0	467
Kericho	60.8	29.3	9.6	0.3	100.0	12.4	87.6	0.0	100.0	387
Bomet	47.9	39.5	12.5	0.2	100.0	8.9	91.1	0.0	100.0	480
Kakamega	50.4	35.3	12.3	1.9	100.0	24.7	75.3	0.0	100.0	665
Vihiga	36.3	32.8	28.6	2.3	100.0	27.3	72.7	0.0	100.0	233
Bungoma	42.4	30.4	22.4	4.8	100.0	37.0	63.0	0.0	100.0	625
Busia	62.8	23.0	12.8	1.4	100.0	32.0	67.7	0.2	100.0	373
Siaya	49.4	31.2	17.8	1.6	100.0	29.0	71.0	0.0	100.0	370
Kisumu	61.5	23.3	13.3	1.9	100.0	29.2	70.8	0.0	100.0	551
Homa Bay	35.4	35.0	20.5	9.2	100.0	19.3	80.7	0.0	100.0	485
Migori	66.9	24.0	8.2	0.9	100.0	25.7	74.3	0.0	100.0	509
Kisii	39.1	27.2	23.0	10.8	100.0	27.7	72.3	0.0	100.0	439
Nyamira	45.7	33.4	16.5	4.4	100.0	13.2	86.8	0.0	100.0	238
Nairobi City	82.9	14.9	1.5	0.7	100.0	14.7	85.3	0.0	100.0	3,550
Total	59.4	25.6	11.4	3.6	100.0	23.4	76.6	0.0	100.0	21,231

[1] Includes car/truck, tuk tuk, public bus, motorcycle/scooter, and boat with motor
[2] Includes animal-drawn cart, bicycle, boat without motor, and walking

Table 9.21.1 Human papillomavirus vaccination coverage among women age 15–19

Percent distribution of women age 15–19 by number of doses of HPV vaccine received, percentage who have received any HPV vaccination, and percentage who ever received an HPV vaccination card, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Number of doses of HPV vaccine received					Percentage who received any HPV vaccination[2]	Percentage who ever received an HPV vaccination card[3]	Number of women
	None[1]	1	2	Don't know number of doses	Total			
Age								
15–17	30.3	20.2	49.4	0.1	100.0	69.7	58.4	2,433
15	24.7	19.6	55.6	0.1	100.0	75.3	67.9	771
16	29.4	18.2	52.3	0.1	100.0	70.6	59.1	809
17	36.1	22.7	41.1	0.1	100.0	63.9	49.2	853
18	52.8	20.9	26.3	0.0	100.0	47.2	33.5	776
19	64.6	15.7	19.3	0.4	100.0	35.4	26.4	707
Residence								
Urban	48.7	15.3	36.0	0.0	100.0	51.3	41.5	1,398
Rural	36.6	21.8	41.3	0.2	100.0	63.4	51.2	2,519
Wealth quintile								
Lowest	39.3	23.1	37.4	0.1	100.0	60.7	48.1	771
Second	33.3	20.0	46.5	0.3	100.0	66.7	54.4	905
Middle	38.4	22.6	38.7	0.3	100.0	61.6	51.0	827
Fourth	48.0	17.3	34.7	0.0	100.0	52.0	43.7	651
Highest	48.4	13.8	37.8	0.0	100.0	51.6	39.2	762
Total	40.9	19.5	39.4	0.1	100.0	59.1	47.7	3,916

HPV = human papillomavirus
[1] Includes women who don't know if they received an HPV vaccination
[2] Includes women who don't know how many doses of HPV vaccine they received
[3] Women who did not receive an HPV vaccination or did not know if they received an HPV vaccination are assumed not to have received an HPV vaccination card.

Table 9.21.1C Human papillomavirus vaccination coverage among women age 15–19: County level

Percent distribution of women age 15–19 by number of doses of HPV vaccine received, percentage who have received any HPV vaccination, and percentage who ever received an HPV vaccination card, according to county, Kenya Mini DHS 2025–26

	Number of doses of HPV vaccine received					Percentage who received any HPV vaccination[2]	Percentage who ever received an HPV vaccination card[3]	Number of women
County	None[1]	1	2	Don't know number of doses	Total			
Mombasa	61.8	13.0	25.2	0.0	100.0	38.2	37.8	96
Kwale	40.1	21.9	37.0	0.9	100.0	59.9	38.1	94
Kilifi	16.6	19.2	63.7	0.6	100.0	83.4	62.0	137
Tana River	45.6	22.5	31.9	0.0	100.0	54.4	39.0	35
Lamu	49.4	9.4	41.2	0.0	100.0	50.6	39.8	14
Taita Taveta	35.8	21.5	42.7	0.0	100.0	64.2	48.9	25
Garissa	91.9	8.1	0.0	0.0	100.0	8.1	7.5	45
Wajir	90.7	7.4	1.9	0.0	100.0	9.3	3.0	38
Mandera	99.2	0.8	0.0	0.0	100.0	0.8	0.8	31
Marsabit	43.2	31.7	25.1	0.0	100.0	56.8	44.5	40
Isiolo	66.7	24.4	8.9	0.0	100.0	33.3	24.7	18
Meru	36.5	47.7	15.8	0.0	100.0	63.5	54.4	103
Tharaka-Nithi	34.6	20.8	44.6	0.0	100.0	65.4	49.7	27
Embu	58.6	9.0	32.4	0.0	100.0	41.4	42.9	42
Kitui	12.3	6.9	80.8	0.0	100.0	87.7	74.3	82
Machakos	43.2	20.2	36.6	0.0	100.0	56.8	51.5	98
Makueni	47.1	18.8	34.1	0.0	100.0	52.9	39.2	64
Nyandarua	(15.5)	(9.8)	(72.4)	(2.2)	100.0	(84.5)	(62.7)	36
Nyeri	(42.3)	(19.4)	(38.3)	(0.0)	100.0	(57.7)	(57.7)	33
Kirinyaga	(29.1)	(11.9)	(59.1)	(0.0)	100.0	(70.9)	(55.7)	30
Murang'a	24.4	19.1	56.5	0.0	100.0	75.6	64.4	77
Kiambu	44.0	12.5	43.5	0.0	100.0	56.0	52.3	245
Turkana	74.7	17.0	8.3	0.0	100.0	25.3	20.5	37
West Pokot	53.1	32.3	14.6	0.0	100.0	46.9	43.7	52
Samburu	44.7	24.3	31.0	0.0	100.0	55.3	47.6	22
Trans Nzoia	49.1	15.3	35.6	0.0	100.0	50.9	43.0	84
Uasin Gishu	35.6	17.9	46.5	0.0	100.0	64.4	60.6	112
Elgeyo-Marakwet	40.1	14.5	45.4	0.0	100.0	59.9	54.5	35
Nandi	55.5	7.9	36.6	0.0	100.0	44.5	40.1	86
Baringo	27.9	9.9	62.2	0.0	100.0	72.1	36.0	39
Laikipia	25.2	20.2	54.6	0.0	100.0	74.8	68.8	32
Nakuru	20.9	32.3	46.9	0.0	100.0	79.1	69.8	188
Narok	52.1	22.8	25.1	0.0	100.0	47.9	34.3	78
Kajiado	40.7	22.9	36.5	0.0	100.0	59.3	51.8	65
Kericho	19.6	22.5	57.6	0.4	100.0	80.4	63.3	70
Bomet	30.2	25.9	42.7	1.3	100.0	69.8	53.9	106
Kakamega	31.8	28.1	39.0	1.1	100.0	68.2	61.7	139
Vihiga	19.4	20.7	60.0	0.0	100.0	80.6	66.2	66
Bungoma	44.2	16.5	39.3	0.0	100.0	55.8	41.8	158
Busia	31.9	26.2	41.9	0.0	100.0	68.1	42.5	87
Siaya	36.3	20.9	42.8	0.0	100.0	63.7	58.4	76
Kisumu	32.8	21.4	45.7	0.0	100.0	67.2	61.5	106
Homa Bay	38.5	17.7	43.9	0.0	100.0	61.5	33.6	131
Migori	37.1	26.6	36.3	0.0	100.0	62.9	45.1	96
Kisii	52.2	20.5	27.3	0.0	100.0	47.8	38.4	85
Nyamira	29.2	25.5	45.3	0.0	100.0	70.8	62.6	61
Nairobi City	53.0	14.1	32.9	0.0	100.0	47.0	33.4	498
Total	40.9	19.5	39.4	0.1	100.0	59.1	47.7	3,916

Note: Figures in parentheses are based on 25–49 unweighted cases.
HPV = human papillomavirus
[1] Includes women who don't know if they received an HPV vaccination
[2] Includes women who don't know how many doses of HPV vaccine they received
[3] Women who did not receive an HPV vaccination or did not know if they received an HPV vaccination are assumed not to have received an HPV vaccination card.

Table 9.21.2 Human papillomavirus vaccination coverage among women age 20–49

Percent distribution of women age 20–49 by number of doses of HPV vaccine received, and percentage who have received any HPV vaccination, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Number of doses of HPV vaccine received					Percentage who received any HPV vaccination[2]	Number of women
	None[1]	1	2	Don't know	Total		
Age							
20–24	80.4	10.3	9.2	0.1	100.0	19.6	3,653
25–29	92.4	3.8	3.5	0.3	100.0	7.6	3,732
30–34	95.9	1.7	2.3	0.2	100.0	4.1	3,108
35–39	95.9	1.7	2.4	0.0	100.0	4.1	2,989
40–44	96.7	1.4	1.9	0.0	100.0	3.3	2,048
45–49	98.3	0.6	1.1	0.0	100.0	1.7	1,784
Residence							
Urban	92.1	4.2	3.6	0.2	100.0	7.9	8,432
Rural	92.3	3.5	4.1	0.1	100.0	7.7	8,883
Wealth quintile							
Lowest	92.2	3.0	4.6	0.1	100.0	7.8	2,541
Second	91.3	3.9	4.8	0.0	100.0	8.7	2,947
Middle	92.4	3.9	3.6	0.1	100.0	7.6	3,306
Fourth	91.3	4.7	3.9	0.1	100.0	8.7	4,149
Highest	93.5	3.4	2.9	0.2	100.0	6.5	4,372
Total	92.2	3.8	3.8	0.1	100.0	7.8	17,315

HPV = human papillomavirus
[1] Includes women who don't know if they received an HPV vaccination
[2] Includes women who don't know how many doses of HPV vaccine they received

Table 9.21.2C Human papillomavirus vaccination coverage among women age 20–49: County level

Percent distribution of women age 20–49 by number of doses of HPV vaccine received, and percentage who have received any HPV vaccination, according to county, Kenya Mini DHS 2025–26

County	Number of doses of HPV vaccine received					Percentage who received any HPV vaccination[2]	Number of women
	None[1]	1	2	Don't know	Total		
Mombasa	96.4	2.1	1.3	0.3	100.0	3.6	707
Kwale	98.9	1.1	0.0	0.0	100.0	1.1	281
Kilifi	69.0	6.7	24.0	0.3	100.0	31.0	471
Tana River	96.3	0.8	2.9	0.0	100.0	3.7	102
Lamu	93.3	4.3	2.3	0.0	100.0	6.7	40
Taita Taveta	90.2	5.5	4.4	0.0	100.0	9.8	128
Garissa	96.7	1.7	1.6	0.0	100.0	3.3	123
Wajir	98.7	1.1	0.1	0.0	100.0	1.3	92
Mandera	100.0	0.0	0.0	0.0	100.0	0.0	92
Marsabit	94.7	3.9	1.5	0.0	100.0	5.3	102
Isiolo	92.5	6.3	1.2	0.0	100.0	7.5	58
Meru	94.0	3.5	2.5	0.0	100.0	6.0	427
Tharaka-Nithi	94.6	3.3	1.6	0.5	100.0	5.4	120
Embu	99.2	0.7	0.1	0.0	100.0	0.8	197
Kitui	91.3	0.3	8.2	0.1	100.0	8.7	330
Machakos	91.0	5.9	3.1	0.0	100.0	9.0	579
Makueni	96.0	2.7	1.2	0.0	100.0	4.0	286
Nyandarua	94.7	3.7	1.6	0.0	100.0	5.3	187
Nyeri	98.6	0.6	0.4	0.4	100.0	1.4	189
Kirinyaga	97.1	1.7	1.2	0.0	100.0	2.9	228
Murang'a	95.1	2.2	2.7	0.0	100.0	4.9	300
Kiambu	98.2	1.0	0.8	0.0	100.0	1.8	1,444
Turkana	94.2	3.2	2.5	0.0	100.0	5.8	141
West Pokot	89.6	8.4	2.0	0.0	100.0	10.4	174
Samburu	85.0	8.1	6.9	0.0	100.0	15.0	76
Trans Nzoia	91.3	5.7	3.0	0.0	100.0	8.7	286
Uasin Gishu	92.1	4.2	3.4	0.2	100.0	7.9	556
Elgeyo-Marakwet	97.7	1.3	1.0	0.0	100.0	2.3	142
Nandi	95.6	1.4	3.0	0.0	100.0	4.4	275
Baringo	68.1	7.1	23.9	0.9	100.0	31.9	148
Laikipia	90.6	4.6	4.7	0.0	100.0	9.4	164
Nakuru	91.8	3.5	4.7	0.0	100.0	8.2	891
Narok	93.5	4.9	1.5	0.0	100.0	6.5	350
Kajiado	90.3	5.0	4.7	0.0	100.0	9.7	403
Kericho	85.2	6.6	7.9	0.3	100.0	14.8	317
Bomet	95.8	2.5	1.7	0.0	100.0	4.2	374
Kakamega	92.0	4.9	2.8	0.4	100.0	8.0	526
Vihiga	88.8	8.3	2.9	0.0	100.0	11.2	167
Bungoma	93.2	4.4	2.4	0.0	100.0	6.8	467
Busia	93.8	3.9	2.2	0.0	100.0	6.2	286
Siaya	98.9	0.0	1.1	0.0	100.0	1.1	294
Kisumu	92.9	2.0	5.1	0.0	100.0	7.1	445
Homa Bay	71.0	12.0	16.7	0.3	100.0	29.0	354
Migori	94.9	2.2	2.9	0.0	100.0	5.1	413
Kisii	98.1	1.8	0.1	0.0	100.0	1.9	354
Nyamira	97.5	1.4	1.1	0.0	100.0	2.5	177
Nairobi City	90.5	5.5	3.8	0.3	100.0	9.5	3,052
Total	92.2	3.8	3.8	0.1	100.0	7.8	17,315

HPV = human papillomavirus
[1] Includes women who don't know if they received an HPV vaccination
[2] Includes women who don't know how many doses of HPV vaccine they received

Table 9.22 Source of human papillomavirus vaccinations

Among women age 15–49 who received any HPV vaccination, percent distribution by source of most recent HPV vaccination, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Place where most recent HPV vaccination was received					Total	Number of women who received any HPV vaccination
	Public health facility	Private health facility	NGO health facility	School	Other		
Age							
15–19	7.8	0.6	0.0	91.2	0.4	100.0	2,313
15	7.9	0.8	0.0	91.1	0.2	100.0	580
16	10.7	0.9	0.0	88.4	0.0	100.0	571
17	6.2	0.7	0.0	92.0	1.1	100.0	545
18	6.7	0.2	0.0	92.9	0.2	100.0	366
19	6.6	0.0	0.0	93.3	0.2	100.0	250
20–49	18.0	2.4	0.9	76.5	2.1	100.0	1,347
20–24	6.7	1.2	0.1	91.8	0.2	100.0	717
25–29	22.8	1.7	0.3	74.4	0.8	100.0	282
30–34	34.1	1.3	5.5	56.2	2.9	100.0	128
35–39	32.2	7.7	0.4	53.8	4.9	100.0	122
40–44	51.8	9.4	1.0	17.0	20.9	100.0	67
45–49	(41.3)	(5.4)	(7.4)	(42.6)	(3.2)	100.0	31
HPV vaccination card[1]							
Ever received card	8.7	0.7	0.0	90.1	0.5	100.0	1,827
Never received card	4.4	0.4	0.0	95.1	0.1	100.0	486
Number of doses received							
1	10.1	1.4	0.7	86.6	1.3	100.0	1,425
2	12.6	1.2	0.1	85.3	0.9	100.0	2,210
Residence							
Urban	13.0	1.7	0.7	82.9	1.6	100.0	1,385
Rural	10.7	1.0	0.1	87.5	0.7	100.0	2,275
Wealth quintile							
Lowest	7.8	0.3	0.0	91.2	0.6	100.0	666
Second	9.0	1.0	0.0	89.4	0.6	100.0	860
Middle	13.7	0.8	0.1	84.9	0.6	100.0	760
Fourth	14.8	0.6	0.0	83.5	1.1	100.0	697
Highest	13.0	3.8	1.7	79.0	2.4	100.0	677
Total	11.6	1.3	0.3	85.8	1.0	100.0	3,660

Notes: Figures in parentheses are based on 25–49 unweighted cases. Total includes 25 women who didn't know the number of HPV vaccine doses received.
HPV = human papillomavirus
NGO = nongovernmental organisation
[1] Women age 20–49 were not asked about vaccination cards.

Table 9.22C Source of human papillomavirus vaccinations: County level

Among women age 15–49 who received any HPV vaccination, percent distribution by source of most recent HPV vaccination, according to county, Kenya Mini DHS 2025–26

County	Place where most recent HPV vaccination was received					Total	Number of women who received any HPV vaccination
	Public health facility	Private health facility	NGO health facility	School	Other		
Mombasa	(19.6)	(2.1)	(3.7)	(72.6)	(2.1)	100.0	62
Kwale	2.9	0.0	0.0	97.1	0.0	100.0	59
Kilifi	5.8	0.4	0.0	93.8	0.0	100.0	260
Tana River	1.1	0.0	0.0	98.9	0.0	100.0	23
Lamu	0.0	2.3	0.0	97.7	0.0	100.0	10
Taita Taveta	29.3	0.0	0.0	70.7	0.0	100.0	29
Garissa	(0.0)	(0.0)	(0.0)	(100.0)	(0.0)	100.0	8
Wajir	*	*	*	*	*	100.0	5
Mandera	*	*	*	*	*	100.0	0
Marsabit	5.1	0.0	0.0	94.9	0.0	100.0	28
Isiolo	3.0	0.0	0.0	97.0	0.0	100.0	10
Meru	8.0	3.9	0.0	82.6	5.6	100.0	91
Tharaka-Nithi	(28.7)	(7.0)	(0.0)	(64.3)	(0.0)	100.0	24
Embu	(25.0)	(0.0)	(0.0)	(75.0)	(0.0)	100.0	19
Kitui	5.2	0.4	0.0	94.4	0.0	100.0	100
Machakos	12.9	1.3	0.0	85.8	0.0	100.0	108
Makueni	20.3	0.0	0.0	79.7	0.0	100.0	45
Nyandarua	(6.7)	(4.8)	(0.0)	(88.5)	(0.0)	100.0	41
Nyeri	(29.3)	(4.3)	(0.0)	(66.4)	(0.0)	100.0	22
Kirinyaga	(22.5)	(6.7)	(0.0)	(65.1)	(5.7)	100.0	28
Murang'a	12.0	1.0	0.0	87.0	0.0	100.0	73
Kiambu	14.0	4.6	0.0	79.6	1.8	100.0	163
Turkana	(17.4)	(4.1)	(0.0)	(78.6)	(0.0)	100.0	17
West Pokot	5.7	0.0	0.0	94.3	0.0	100.0	42
Samburu	13.3	0.0	0.0	85.1	1.6	100.0	23
Trans Nzoia	3.8	0.7	0.0	94.2	1.3	100.0	68
Uasin Gishu	17.2	2.1	0.0	80.7	0.0	100.0	115
Elgeyo-Marakwet	16.7	2.6	0.0	80.7	0.0	100.0	24
Nandi	12.1	1.6	0.0	86.2	0.0	100.0	50
Baringo	16.8	0.9	0.0	82.3	0.0	100.0	75
Laikipia	18.6	0.0	1.7	79.7	0.0	100.0	39
Nakuru	7.8	0.0	0.0	89.5	2.7	100.0	222
Narok	5.4	0.0	0.0	94.6	0.0	100.0	60
Kajiado	6.7	2.7	0.0	88.2	2.4	100.0	78
Kericho	23.3	0.9	0.0	74.8	1.0	100.0	103
Bomet	13.8	0.0	0.0	86.2	0.0	100.0	90
Kakamega	8.5	2.1	0.0	89.4	0.0	100.0	137
Vihiga	14.7	1.1	0.0	83.3	0.8	100.0	72
Bungoma	5.1	1.1	0.0	93.8	0.0	100.0	120
Busia	11.0	0.0	0.0	88.1	1.0	100.0	77
Siaya	12.5	0.0	0.0	86.9	0.7	100.0	52
Kisumu	7.5	1.1	1.0	90.5	0.0	100.0	103
Homa Bay	4.4	1.2	0.7	93.1	0.0	100.0	184
Migori	13.8	1.6	0.0	83.3	1.3	100.0	81
Kisii	(12.2)	(0.0)	(0.0)	(87.8)	(0.0)	100.0	47
Nyamira	9.6	1.1	0.0	87.9	1.4	100.0	48
Nairobi	16.4	0.9	1.3	78.8	2.5	100.0	525
Total	11.6	1.3	0.3	85.8	1.0	100.0	3,660

Notes: Figures in parentheses are based on 25–49 unweighted cases. An asterisk indicates that a figure is based on fewer than 25 unweighted cases and has been suppressed.
HPV = human papillomavirus
NGO = nongovernmental organisation

CHILD HEALTH 10

Key Findings

- ***Birth weight:*** Among children born within the previous 2 years, 9% were born with a low birth weight (<2.5 kg).
- ***Vaccinations:*** By the time of the survey, 80% of children age 12–23 months were fully vaccinated against all basic antigens and 46% were fully vaccinated according to the national schedule.
- ***Symptoms of acute respiratory infection:*** Advice or treatment was sought for 75% of children under age 5 who had symptoms of acute respiratory infection in the 2 weeks before the survey.
- ***Fever:*** Advice or treatment was sought for 68% of children under age 5 who had a fever in the 2 weeks before the survey.
- ***Diarrhoea:*** Advice or treatment was sought for 58% of children under age 5 who had diarrhoea 2 weeks before the survey. Sixty-three percent of children with diarrhoea received oral rehydration therapy, 26% received oral rehydration salts and zinc, and 23% received no treatment.
- ***Early childhood development:*** Overall, 85% of children age 3–23 months and 83% of children age 24–59 months are developmentally on track.

Information on child health and survival can help policymakers and programme managers assess the efficacy of current strategies, formulate appropriate interventions to prevent deaths from childhood illnesses, and improve the health of children in Kenya.

This chapter presents information on birth weight and vaccination status for young children. It also examines the prevalence of and care-seeking behaviours for three common childhood illnesses: symptoms of acute respiratory infection (ARI), fever, and diarrhoea. In addition, it presents data on early childhood development indicators, providing insight into whether children in Kenya are on track in terms of health, learning, and psychosocial well-being. These indicators are also relevant for monitoring progress towards Sustainable Development Goal Target 4.2 on early childhood development outcomes.

10.1 CHILD'S SIZE AND BIRTH WEIGHT

Low birth weight

Percentage of births with a reported birth weight below 2.5 kg regardless of gestational age

Sample: Live births in the 2 years before the survey that have a reported birth weight, from either a written record or the mother's report

Birth weight is an important indicator of a child's health and survival. Babies born with low birth weight are more vulnerable to illness, death, and developmental challenges later in life. A baby is considered to have low birth weight if he or she is born weighing less than 2,500 grams (2.5 kg).

In the 2025–26 Kenya Mini Demographic and Health Survey (KMDHS), birth weight was obtained from either written records or mothers' recall for 91% of children born during the 2 years preceding the survey. Nine percent of these babies had a low birth weight. Overall, 3% of the babies were reported to be very small at birth, 11% were reported to be smaller than average, and 83% were reported to be average size or larger (**Table 10.1**).

Patterns by background characteristics

- Children born to mothers younger than age 20 were more likely to have low birth weight (15%) than those born to mothers age 20–34 (8%) or age 35–49 years (9%).
- A higher percentage of children born in rural areas (10%) than of children born in rural areas (8%) had low birth weight.
- Low birth weight was more prevalent among children from households in the lowest wealth quintile (14%) than among those from households in the highest wealth quintile (8%).
- Low birth weight was most common in Kwale County (24%), followed by Tana River and Makueni Counties (15% each), while Machakos County recorded the lowest prevalence of low birth weight (2%) (**Table 10.1C**).

10.2 Vaccination of Children

Universal immunisation of children against common vaccine-preventable diseases is crucial for reducing infant and child morbidity and mortality. In Kenya, routine childhood vaccines include the Bacille Calmette-Guérin (BCG) vaccine against tuberculosis; oral polio vaccine (OPV) and inactivated polio vaccine (IPV); pentavalent diphtheria, pertussis, tetanus, hepatitis B, and *Haemophilus influenzae* type b (DPT-HepB-Hib) vaccine; pneumococcal conjugate vaccine (PCV); rotavirus vaccine (RV), and measles-rubella (MR) vaccine. The yellow fever vaccine is also routinely administered in selected high-risk counties including Baringo, Elgeyo/Marakwet, West Pokot, and Turkana.

Information on vaccination coverage in the 2025–26 KMDHS was obtained from two sources: written vaccination records (vaccination cards or other health documents) and mothers' recall. For each living child born in the 3 years before the survey, the mother was asked to show the interviewer the vaccination card or other document used for recording the child's immunisations. If such documentation was available, the interviewer copied the dates of each vaccine received. If a vaccine was not recorded as administered, the mother was asked to recall whether that particular vaccine had been given. If the mother could not present documentation of vaccination, she was asked to recall whether the child had received the BCG, OPV or IPV, DPT-HepB-Hib, PCV, RV, MR, and yellow fever vaccines. If she indicated that the child had received any of the multi-dose vaccines, she was asked the number of doses the child had received.

10.2.1 Vaccination Card Ownership and Availability

Vaccination cards are a critical tool in ensuring that a child receives all recommended vaccinations on schedule. Ninety-eight percent of both children age 12–23 months and those age 24–35 months had ever had a vaccination card or other document on which their vaccinations were recorded (**Table 10.2**). However, not all mothers could produce their child's vaccination card at the time of the interview. Sixty-nine percent of children

age 12–23 months and 56% of children age 24–35 months had vaccination cards available at the time of the interview.

Patterns by background characteristics

- The percentage of children age 12–23 months whose vaccination card was observed is higher in rural areas (74%) than in urban areas (62%).
- Similarly, the percentage of children age 24–35 months whose vaccination card was observed is higher in rural areas (65%) than in urban areas (44%).

10.2.2 Basic Antigen Coverage

Fully vaccinated: basic antigens

Percentage of children who received specific vaccines at any time before the survey (according to a vaccination card or the mother's report). To have received all basic antigens, a child must have received at least:

- One dose of BCG vaccine, which protects against tuberculosis
- Three doses of polio vaccine given as OPV, IPV, or a combination of OPV and IPV
- Three doses of DPT-containing vaccine, which protects against diphtheria, pertussis (whooping cough), and tetanus
- One dose of measles-containing vaccine given as MR

Sample: Children age 12–23 months and children age 24–35 months


Historically, an important measure of vaccination coverage has been the percentage of children receiving all 'basic' antigens. Children are considered fully vaccinated against all basic antigens if they have received the BCG vaccine, three doses each of a polio vaccine (excluding polio given at birth) and a DPT-containing vaccine, and a single dose of a measles-containing vaccine. In Kenya, the BCG vaccine is usually given at birth or at first clinic contact, while the polio and DPT-containing vaccines are given at approximately age 6, 10, and 14 weeks. The first measles-containing vaccination should be given at or soon after age 9 months.

Among children age 12–23 months, 80% are fully vaccinated with basic antigens, including 73% who were fully vaccinated with basic antigens by the appropriate age per recommendations within the National Vaccines and Immunisation Program. Among children age 24–35 months, 85% are fully vaccinated against all basic antigens, including 78% who were fully vaccinated by the appropriate age (**Table 10.3**).

Trends: Vaccination coverage with all basic antigens has increased substantially over time, rising from 57% in the 2003 Kenya Demographic and Health Survey (KDHS) to 77% in the 2008–09 KDHS. It then increased more gradually to 79% in 2014 and 80% in 2022, where it remained in 2025–26. Over the same period, the percentage of children receiving no vaccinations declined steadily, dropping from 7% in 2003 to 3% in 2008–09, then to 2% in 2014 and 2022, and further to 1% in 2025–26 (**Figure 10.1**).

Figure 10.1 **Trends in basic antigen coverage in children**

Percentage of children age 12–23 months who received all basic antigens at any time before the survey



Patterns by background characteristics

- Vaccination coverage with all basic antigens is higher among girls age 12–23 months than among boys of the same age (81% versus 78%) (**Table 10.4**).
- Children are more likely to be fully vaccinated against basic antigens if their mothers have more than a secondary education (87%) than if their mothers have no education (62%).
- A higher percentage of children in the highest wealth quintile (91%) than of children in the lowest wealth quintile (65%) are fully vaccinated against basic antigens.

10.2.3 National Schedule Coverage

Fully vaccinated according to national schedule: age 12–23 months

Percentage of children who received specific vaccines at any time before the survey (according to a vaccination card or the mother's report). To be fully vaccinated according to the national schedule, a child must receive the following:

- One dose of BCG vaccine
- Birth dose of OPV
- Three doses of OPV and one dose of IPV
- Three doses of DPT-HepB-Hib vaccine
- Three doses of pneumococcal conjugate vaccine
- Three doses of rotavirus vaccine
- One dose of a measles-containing vaccine given as MR

Sample: Children age 12–23 months

Fully vaccinated according to national schedule: age 24–35 months

Percentage of children who received specific vaccines at any time before the survey (according to a vaccination card or the mother's report). To be fully vaccinated according to the national schedule, a child must receive all the vaccinations listed above plus the following:

- A second dose of MR

Sample: Children age 24–35 months

A second measure of vaccination coverage is the percentages of children age 12–23 months and children age 24–35 months who are fully vaccinated according to the national schedule. In this report, a child age 12–23 months is considered fully vaccinated according to the national schedule if the child has received all basic antigens as well as a birth dose of OPV, a dose of IPV, three doses of the pneumococcal vaccine, and three doses of the rotavirus vaccine, reflecting the updated national immunisation schedule introduced in 2023. Children age 24–35 months are considered fully vaccinations according to the national schedule if they have received a second dose of the MR vaccine in addition to all relevant vaccinations for children age 12–23 months.

Of note, a review of vaccination cards used in the pilot field practice of the main training showed that some private facilities were administering Hexaxim, a hexavalent vaccine containing DPT, HepB, Hib, and IPV in place of the pentavalent vaccine and, in some cases, as a substitute for OPV. Under Kenya's national immunisation schedule, children are expected to receive at least three doses of OPV and one dose of IPV. This variation in administration and recording created challenges in classifying children's vaccination status, because some children immunised in private facilities received both OPV and IPV while others received IPV only. As a result, some children may not have met the definition of being fully vaccinated according to the national schedule but would have met the definition for full basic antigen coverage.

To improve measurement of these vaccinations, the vaccination card section and survey programme logic were revised to capture additional IPV doses recorded as Hexaxim. Interviewers were instructed to record OPV 0–3 and IPV as usual and, where applicable, to enter any additional IPV or Hexaxim doses shown on the card. This guidance was provided to all research assistants, and secondary editors were instructed to pay particular attention to vaccination cards from private facilities. However, public facility cards and mother-child booklets do not include a field for recording the vaccine brand name. Therefore, if Hexaxim was administered, it may not have been identified from the card unless the additional IPV doses were explicitly recorded, making accurate classification dependent on how the provider documented the vaccination.

Results of the 2025–26 KMDHS show that 46% of children age 12–23 months are fully vaccinated according to the national schedule, with 42% being vaccinated by the appropriate age. Similarly, 38% of children age 24–35 months are fully vaccinated according to the national schedule, with 33% being vaccinated by the appropriate age (**Table 10.3**).


Among children age 12–23 months, 98% have received a BCG vaccine, with 97% being vaccinated with BCG by the appropriate age. Coverage is highest for the first doses of the DPT-HepB-Hib (98%), pneumococcal (98%), and rotavirus (96%) vaccines. Ninety-four percent of children age 24–35 months had received the first dose of MR, but only 72% had received the second dose (**Figure 10.2**).

Patterns by background characteristics

- Vaccination coverage according to the national schedule is slightly higher among boys age 12–23 months (48%) than among girls age 12–23 months (43%). Among children age 24–35 months, coverage is equal for boys and girls (38% each) (**Table 10.4**).
- Children whose mothers have more than a secondary education are more likely to be fully vaccinated according to the national schedule than are children whose mothers have no education, and this is true among both children age 12–23 months (49% versus 23%) and children age 24–35 months (44% versus 18%).

Figure 10.2 **Childhood vaccinations**

Percentage of children age 12–23 months who received specific vaccines at any time before the survey



Source of Vaccinations

The main source of childhood vaccinations is the public medical sector, with 83% of children age 12–23 months and 85% of children age 24–35 months receiving their vaccinations from this sector. The non-nongovernmental organisation (non-NGO) private sector is the second most common source of childhood vaccination, at 16% for children age 12–23 months and 13% for children age 24–35 months (**Table 10.5**).

10.3 Symptoms of Acute Respiratory Infection and Care-seeking Behaviour

ARIs are defined as bacterial or viral infections of the respiratory tract leading to breathing difficulties, fever, and other complications. ARIs, which include pneumonia, are the leading cause of morbidity and mortality among children under age 5 in Kenya. Treatment with antibiotics can prevent many deaths caused by ARIs. Indoor air pollution has been documented as a risk factor for ARIs and is associated with low socioeconomic status and overcrowding.

Care seeking for symptoms of acute respiratory infection

Children with symptoms of ARI for whom advice or treatment was sought. ARI symptoms consist of short, rapid breathing that is chest-related and/or difficult breathing that is chest related.

Sample: Children under age 5 with symptoms of ARI in the 2 weeks before the survey

Among children under age 5, 2% exhibited symptoms of ARI in the 2 weeks before the survey. Among the children with ARI symptoms, advice or treatment was sought for 75%, and 50% received treatment or advice on the same or next day (**Table 10.6.1**).

10.3.1 Treatment for Symptoms

Among children under age 5 with symptoms of ARI, 53% received antibiotics; 59% were treated with other medicines such as aspirin, paracetamol, or ibuprofen; 9% were given artemisinin-based combination therapy; and 25% were given no treatment (**Table 10.6.2**).

10.3.2 Source of Advice or Treatment

For children under age 5 with symptoms of ARI for whom advice or treatment was sought, the primary source of care was the public medical sector (56%), followed by the private medical sector (mainly non-NGO providers) (45%). Government health centres were the leading source in the public sector (24%), while pharmacies (30%) were the main source in the private sector (**Table 10.7**).

10.4 Fever and Care-seeking Behaviour

Fever is a key symptom of malaria and other acute infections in young children. In Kenya, malaria is one of the leading causes of morbidity and mortality in children under age 5 and is most prevalent in the lake and coastal endemic regions of the country.

Care seeking for fever

Children with fever for whom advice or treatment was sought.

Sample: Children under age 5 with a fever in the 2 weeks before the survey

In the 2 weeks preceding the survey, mothers reported that 14% of children under age 5 had experienced fever. Advice or treatment was sought for 68% of these children, and care was sought on the same day or the following day for 44% of them. Among children under age 5 with fever, 36% received antibiotics (**Table 10.8**).

Patterns by background characteristics

- Among children under age 5, the percentage with fever for whom advice or treatment was sought in the 2 weeks before the survey was higher in urban areas (70%) than in rural areas (66%). The percentage of with fever who for whom advice or treatment was sought the same or next day was also higher in urban areas (47%) than in rural areas (41%).
- Care seeking for fever for children under age 5 was more common when mothers had more than a secondary education than when they had no education, both for any advice or treatment (74% versus 63%) and for prompt care seeking the same or next day (52% versus 38%).

10.5 Diarrhoeal Disease

Diarrhoeal disease remains an important cause of morbidity and mortality among young children in Kenya. It accounts for approximately 7% of all deaths among children under age 5, despite the availability of a simple treatment solution (United Nations Children's Fund [UNICEF] 2021). Oral rehydration therapy (ORT) and supplemental zinc, combined with continued feeding, are the recommended interventions for treating diarrhoea. ORT can be provided as increased fluids (especially increased breastfeeding), as fluid prepared from a packet of oral rehydration salts (ORS), or as government-recommended homemade fluids. Zinc has been shown to reduce the severity and duration of diarrhoea, and it is recommended that all children with diarrhoea receive a 5-day course.

10.5.1 Care-seeking Behaviour

Care seeking for diarrhoea

Children with diarrhoea for whom advice or treatment was sought.

Sample: Children under age 5 with diarrhoea in the 2 weeks before the survey


Eleven percent of children under age 5 had a diarrhoeal episode in the 2 weeks before the survey. Advice or treatment was sought for 58% of the children (**Table 10.9**).

Patterns by background characteristics

- Diarrhoea prevalence is highest among children age 6–11 months (19%) and decreases with age to 5% among those age 48–59 months (**Figure 10.3**).
- The percentage of children with diarrhoea in the past 2 weeks was slightly higher in urban areas (13%) than in rural areas (10%). Care seeking for diarrhoea was also slightly higher in urban areas than in rural areas (59% versus 57%).
- Care seeking for diarrhoea increases with level of maternal education, from 62% among children whose mothers have no education to 68% among those whose mothers have more than a secondary education.

Figure 10.3 **Diarrhoea prevalence by age**

Percentage of children under age 5 who had diarrhoea in the 2 weeks before the survey



- Care seeking for diarrhoea also increases with wealth, from 54% among children in the lowest wealth quintile to 64% among those in the highest wealth quintile.

10.5.2 Feeding Practices

Appropriate feeding practices

Children with diarrhoea who were given more liquids than usual and at least as much food as usual.

Sample: Children under age 5 with diarrhoea in the 2 weeks before the survey


Without enough liquids and food, diarrhoea can quickly lead to dehydration, weakness, poor nutrition, and serious illness. Giving more liquids and food than usual and continuing breastfeeding help prevent dehydration, weight loss, and malnutrition.

Twenty-six percent of children under age 5 with diarrhoea in the 2 weeks before the survey were given more liquids than usual, as recommended, and 44% were given the same amount of liquids as usual. However, 28% of the children with diarrhoea received somewhat less or much less liquid than usual, and 2% received no liquids at all (**Table 10.10**).

Forty-two percent of children with diarrhoea were given at least the same amount of food as usual, as recommended. Forty-four percent were given less food than usual, while 5% received no food (**Figure 10.4**).

Figure 10.4 **Feeding practices during diarrhoea**

Percentage of children under age 5 with diarrhoea in the 2 weeks before the survey who received more or less liquids and foods than usual



Patterns by background characteristics

- Among children under age 5 with diarrhoea in the 2 weeks before the survey, a higher percentage in urban areas (28%) than in rural areas (24%) were given more liquids than usual during a diarrhoeal episode (**Table 10.10**).
- Children whose mothers had more than a secondary education were more likely than children whose mothers had no education to receive at least the same amount of liquids as usual during diarrhoea (78% versus 67%).
- The percentage of children given at least the same amount of liquids as usual during diarrhoea generally increased with wealth, from 63% among children in the lowest wealth quintile to 80% among those in the highest wealth quintile.

10.5.3 Oral Rehydration Therapy, Zinc, Continued Feeding, and Other Treatments

Oral rehydration therapy

Children with diarrhoea who were given increased fluids, fluids prepared from a packet of ORS, or government-recommended homemade fluids.

Sample: Children under age 5 with diarrhoea in the 2 weeks before the survey

ORT is a simple and effective way to reduce dehydration caused by diarrhoea. Using a combination of ORS, zinc, continued feeding, and other supportive treatments is important because each addresses a different danger caused by diarrhoea. ORS prevents and treats dehydration, and zinc shortens the duration of diarrhoea and reduces the severity of symptoms. Continued feeding and breastfeeding prevent malnutrition and weight loss.


Sixty-three percent of children with diarrhoea in the 2 weeks before the survey received some form of ORT; 41% received fluids from ORS packets, and 22% received recommended homemade fluids. Forty-six percent received ORT and continued feeding as recommended. Thirty-six percent of the children were given zinc alone, 26% were given zinc and ORS, 10% were given antibiotics, and 23% received no treatment (**Table 10.11** and **Figure 10.5**)

Source of Advice or Treatment for Diarrhoea

Among children under age 5 with diarrhoea for whom advice or treatment was sought, it was mainly sought from the public medical sector (55%), followed by the private medical sector (non-NGO) (41%) (**Table 10.12**).

***Figure 10.5* Treatment of diarrhoea**

Percentage of children under age 5 with diarrhoea in the 2 weeks before the survey who received specific treatments




10.6 Childhood Illness and Care-seeking Behaviour

ARI, fever, and dehydration from diarrhoea are major causes of childhood morbidity and mortality in developing countries (WHO 2024). Prompt medical attention when a child develops symptoms of these illnesses is crucial in reducing child deaths.

Two percent of children under age 5 were reported to have symptoms of ARI. Fever and diarrhoea were reported among 14% and 11% of the children, respectively (**Figure 10.6**).

Treatment or advice was sought for 73% of children with symptoms of ARI, 68% of children with fever, and 58% of children with diarrhoea (**Figure 10.6**).

***Figure 10.6* Symptoms of childhood illness and care seeking**



10.7 Early Childhood Development

Early childhood development (ECD) refers to the process of cognitive, physical, language, temperament, socioemotional and motor development of children that starts at the time of conception until 8 years of age (WHO 2020). These progressions represent distinct yet interconnected domains of development.

Nurturing and supporting progress in a holistic manner is key to ensuring that children have the best chance to thrive and reach their full potential. Physical growth, literacy and numeracy skills, socioemotional development, and learning readiness set the trajectory for lifelong health, learning, and well-being (Shonkoff and Phillips 2000).

The Kenya Newborn and Child Health Strategic Plan 2022–2026 identifies early childhood development (ECD) as a national priority, underscoring the need for reliable data to guide investments. The first years of life, from pregnancy to age three, are the most critical for human development, shaping lifelong health, learning, and productivity with lasting effects across generations. Supporting children to reach their full potential is both a human right and a driver of sustainable development.

The 2025–26 KMDHS used the WHO Global Scales for Early Development (GSED) tool and the UNICEF Early Childhood Development Index 2030 (ECDI2030). Both instruments were administered as part of the Woman's Questionnaire. Respondents were asked questions for each biological child living with them age 3–23 months for the GSED and age 24–59 months for the ECDI2030. The questions focused on the children's behaviours in everyday situations and on the skills and knowledge they had acquired, reflecting the increasing difficulty of skills acquired as children grow. Data generated by the GSED and the ECDI2030 can be used to inform government efforts to improve developmental outcomes among children.

The GSED includes 48 items, organised into different 20-item age-specific sections based on the child's age. The tool provides a holistic assessment of child development across key domains, including motor, language, cognitive, socio-emotional, and adaptive development. These are grouped into three broad domains: motor, language/cognitive, and social skills: Each domain comprises a set of three core subdomains:

- ***Motor subdomains:*** gross motor development, fine motor development, and self-care
- ***Language/cognitive subdomains:*** expressive language, literacy, numeracy, prewriting, and executive functioning
- ***Social skills subdomains:*** emotional skills, social skills, internalising behaviour, and externalising behaviour

The ECDI2030 includes 20 items organised according to the three domains of health, learning, and psychosocial well-being. Each domain includes a set of core subdomains:

- ***Health subdomains:*** gross motor development, fine motor development, and self-care
- ***Learning subdomains:*** expressive language, literacy, numeracy, prewriting, and executive functioning
- ***Psychosocial well-being subdomains:*** emotional skills, social skills, internalising behaviour, and externalising behaviour

The ECDI2030 is not designed for reporting individual domains separately. It is used to produce a single summary score that captures the interlinked developmental concepts embedded in the three domains cited in Sustainable Development Goal Indicator 4.2.1.

Children developmentally on track according to the Global Scales for Early Development

The percentage of children who have achieved the minimum number of GSED milestones expected for their age group as follows:

- 3–6 months: at least 7 milestones
- 6–11 months: at least 9 milestones
- 12–17 months: at least 11 milestones
- 18–23 months: at least 13 milestones

Sample: Children age 3–23 months

Children developmentally on track according to the Early Childhood Development Index 2030

Percentage of children who have achieved the minimum number of ECDI2030 milestones expected for their age group as follows:

- 24–29 months: at least 7 milestones
- 30–35 months: at least 9 milestones
- 36–41 months: at least 11 milestones
- 42–47 months: at least 13 milestones
- 48–59 months: at least 15 milestones

Sample: Children age 24–59 months


Overall, 85% of children age 3–23 months and 83% of children age 24–59 months are developmentally on track in health, learning, and psychosocial well-being (**Tables 10.13** and **10.14**).

Eighty-four percent of children age 3–59 months are developmentally on track according to the combined GSED and ECDI2030 metrics (**Table 10.15**).

Patterns by background characteristics

- A higher percentage of girls than boys are developmentally on track among those age 3–23 months (86% versus 84%) and those age 24–59 months (85% versus 80%) (**Tables 10.13** and **10.14**).
- Children in urban areas are more likely than those in rural areas to be developmentally on track (88% versus 83% among those age 3–23 months and 87% versus 80% among those age 24–59 months).
- The percentage of children who are developmentally on track declines with age in both cohorts, from 90% at age 3–11 months to 81% at age 12–23 months, and from 87% at age 24–35 months to 79% at age 48–59 months (**Figure 10.7**).

Figure 10.7 **Early childhood development**




- The likelihood that children are developmentally on track increases with level of maternal education, from 70% among children age 3–23 months and 64% among those age 24–59 months whose mothers have no education to 90% and 92%, respectively, among those whose mothers have more than a secondary education.

- At the county level, the percentage of children who are developmentally on track ranges from 52% in West Pokot to 97% in Kakamega among those age 3–23 months, and from 51% in Turkana to 97% in Trans Nzoia among those age 24–59 months (**Table 10.13C** and **10.14C**).

- When looking at the two metrics combined, the percentage of children age 3–59 months who are developmentally on track ranges from lows of 54% in West Pokot County and 56% in Turkana County to highs of 95% in Trans Nzoia County and 93% in both Busia and Nyeri Counties (**Table 10.15C** and **Map 10.1**).

Map 10.1 Early childhood development

Percentage of children under age 3–59 months who are developmentally on track according to ECDI and GSED (combined)



LIST OF TABLES

For more information on child health, see the following tables:

- **Table 10.1 Child's size and weight at birth**
- **Table 10.1C Child's size and weight at birth: County level**
- **Table 10.2 Possession and observation of vaccination cards according to background characteristics**
- **Table 10.3 Vaccinations by source of documentation**
- **Table 10.4 Vaccinations by background characteristics**
- **Table 10.5 Source of vaccinations**
- **Table 10.6.1 Children with symptoms of acute respiratory infection and care seeking for symptoms**
- **Table 10.6.2 Treatment for children with symptoms of acute respiratory infection**
- **Table 10.7 Source of advice or treatment for children with symptoms of acute respiratory infection**
- **Table 10.8 Children with fever and care seeking for fever**
- **Table 10.9 Children with diarrhoea and care seeking for diarrhoea**
- **Table 10.10 Feeding practices during diarrhoea**
- **Table 10.11 Oral rehydration salts, zinc, continued feeding, and other treatments for diarrhoea**
- **Table 10.12 Source of advice or treatment for children with diarrhoea**
- **Table 10.13 Global Scales of Early Development**
- **Table 10.13C Global Scales of Early Development: County level**
- **Table 10.14 Early Childhood Development Index 2030**
- **Table 10.14C Early Childhood Development Index 2030: County level**
- **Table 10.15 Combined Early Childhood Development Index 2030 and Global Scales of Early Development**
- **Table 10.15C Combined Early Childhood Development Index 2030 and Global Scales of Early Development: County level**

Table 10.1 Child's size and weight at birth

Percent distribution of live births in the 2 years preceding the survey by mother's estimate of baby's size at birth; percentage of live births in the 2 years preceding the survey that have a reported birth weight by source of information (written record or mother's report); and among live births in the 2 years preceding the survey with a reported birth weight, percentage with birth weight less than 2.5 kg, according to background characteristics, Kenya Mini DHS 2025–26

	Percent distribution of births by size of baby at birth based on mother's estimate					Percentage of births that have a reported birth weight according to:				Among births with a reported birth weight[1]:	
Background characteristic	Very small	Smaller than average	Average or larger	Don't know	Total	Written record	Mother's report	Either	Number of births	Percent-age less than 2.5 kg	Number of births
Mother's age at birth											
<20	3.6	14.4	79.6	2.3	100.0	35.0	54.3	89.4	501	14.8	447
20–34	3.5	10.3	84.2	2.0	100.0	36.4	56.0	92.5	3,053	8.1	2,822
35–49	2.9	14.4	80.1	2.6	100.0	37.6	49.0	86.6	562	9.2	486
Birth order											
1	3.8	11.9	82.9	1.4	100.0	36.5	59.0	95.5	1,306	9.2	1,247
2–3	2.6	11.0	85.2	1.2	100.0	39.5	54.8	94.3	1,787	9.0	1,685
4–5	4.2	11.9	81.1	2.8	100.0	34.3	52.1	86.4	705	9.5	609
6+	4.5	10.1	76.2	9.1	100.0	23.5	44.3	67.7	317	8.1	215
Residence											
Urban	2.4	12.3	84.7	0.7	100.0	39.5	58.7	98.2	1,755	7.6	1,724
Rural	4.2	10.7	81.9	3.2	100.0	34.1	52.0	86.1	2,360	10.3	2,033
Mother's education[2]											
No education	4.8	11.3	71.8	12.1	100.0	17.5	36.4	53.8	363	11.2	195
Primary	5.3	13.4	78.8	2.4	100.0	38.3	49.6	87.9	1,111	12.7	976
Secondary	2.4	10.8	86.2	0.7	100.0	36.9	60.1	97.0	1,817	7.6	1,762
More than secondary	2.6	9.9	86.8	0.6	100.0	41.4	58.4	99.8	825	7.5	823
Wealth quintile											
Lowest	6.0	12.1	75.2	6.8	100.0	27.6	43.4	71.0	873	13.7	620
Second	4.7	12.1	82.3	0.9	100.0	32.8	59.5	92.3	770	11.1	711
Middle	2.1	9.4	86.8	1.8	100.0	41.2	55.6	96.8	739	7.2	715
Fourth	2.7	10.0	86.9	0.4	100.0	39.1	58.9	98.0	945	7.1	926
Highest	1.6	13.3	84.4	0.6	100.0	42.1	57.5	99.6	788	7.7	785
Total	3.4	11.4	83.1	2.1	100.0	36.4	54.9	91.3	4,115	9.1	3,756

[1] Based on either a written record or the mother's recall
[2] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 10.1C Child's size and weight at birth: County level

Percent distribution of live births in the 2 years preceding the survey by mother's estimate of baby's size at birth; percentage of live births in the 2 years preceding the survey that have a reported birth weight by source of information (written record or mother's report); and among live births in the 2 years preceding the survey with a reported birth weight, percentage with birth weight less than 2.5 kg, according to county, Kenya Mini DHS 2025–26

	Percent distribution of births by size of baby at birth based on mother's estimate					Percentage of births that have a reported birth weight according to:				Among births with a reported birth weight[1]:	
County	Very small	Smaller than average	Average or larger	Don't know	Total	Written record	Mother's report	Either	Number of births	Percent-age less than 2.5 kg	Number of births
Mombasa	7.3	6.2	83.4	3.1	100.0	33.0	58.0	91.0	131	5.4	119
Kwale	10.8	9.4	79.8	0.0	100.0	37.7	60.4	98.1	71	24.4	70
Kilifi	3.8	15.4	80.8	0.0	100.0	25.8	72.1	97.9	105	13.0	103
Tana River	11.7	18.4	69.8	0.0	100.0	23.7	47.5	71.2	42	15.2	30
Lamu	3.3	8.3	87.0	1.3	100.0	55.7	35.4	91.1	11	13.4	10
Taita Taveta	5.3	4.4	90.3	0.0	100.0	58.6	34.3	92.9	31	9.2	29
Garissa	11.3	6.3	55.1	27.4	100.0	19.1	26.0	45.1	46	7.2	21
Wajir	8.2	6.9	65.0	19.9	100.0	16.9	15.1	32.0	40	(25.8)	13
Mandera	0.3	5.4	89.8	4.5	100.0	11.7	44.9	56.6	42	4.0	24
Marsabit	0.0	12.2	76.7	11.0	100.0	18.8	62.6	81.4	35	12.0	28
Isiolo	1.1	15.4	82.4	1.2	100.0	3.5	80.8	84.3	17	11.3	14
Meru	6.0	4.9	89.0	0.0	100.0	32.2	54.3	86.5	72	9.6	62
Tharaka-Nithi	(11.3)	(9.8)	(78.9)	(0.0)	100.0	(25.3)	(64.4)	(89.7)	24	(17.8)	21
Embu	(0.0)	(17.0)	(83.0)	(0.0)	100.0	(85.7)	(12.3)	(98.0)	29	(8.4)	28
Kitui	2.6	9.3	85.1	3.0	100.0	32.5	56.8	89.3	66	13.1	59
Machakos	0.0	18.9	81.1	0.0	100.0	27.5	70.2	97.7	159	2.1	155
Makueni	4.7	11.8	83.5	0.0	100.0	65.7	32.9	98.5	62	15.2	61
Nyandarua	0.0	10.8	89.2	0.0	100.0	63.1	36.9	100.0	48	6.7	48
Nyeri	9.6	2.8	87.6	0.0	100.0	1.4	98.6	100.0	42	5.3	42
Kirinyaga	(1.0)	(9.3)	(89.7)	(0.0)	100.0	(72.9)	(27.1)	(100.0)	38	(10.3)	38
Murang'a	1.2	12.8	86.0	0.0	100.0	75.3	23.2	98.5	60	9.7	59
Kiambu	4.4	14.0	78.9	2.6	100.0	27.4	71.1	98.6	272	5.1	268
Turkana	3.4	7.3	86.6	2.7	100.0	18.7	29.1	47.8	63	9.6	30
West Pokot	0.6	5.5	89.1	4.8	100.0	66.7	0.0	66.7	76	13.6	51
Samburu	5.7	10.8	82.7	0.8	100.0	25.5	38.2	63.6	33	12.8	21
Trans Nzoia	4.0	14.6	79.2	2.2	100.0	45.1	45.8	90.9	59	10.2	54
Uasin Gishu	1.4	8.4	90.2	0.0	100.0	15.6	83.4	99.0	124	9.3	122
Elgeyo-Marakwet	3.3	12.5	84.2	0.0	100.0	65.8	28.2	94.0	33	7.2	31
Nandi	3.3	10.0	85.6	1.1	100.0	59.4	32.8	92.3	77	9.6	71
Baringo	2.6	13.9	72.2	11.4	100.0	52.2	27.5	79.7	55	11.6	44
Laikipia	0.0	9.7	85.8	4.6	100.0	53.3	32.0	85.3	51	12.7	43
Nakuru	2.7	6.2	91.0	0.0	100.0	40.0	57.4	97.4	224	5.7	218
Narok	0.8	6.4	72.2	20.6	100.0	19.2	54.8	74.0	91	9.9	67
Kajiado	3.6	14.9	81.5	0.0	100.0	48.2	36.9	85.0	130	8.4	111
Kericho	4.6	5.0	90.4	0.0	100.0	42.9	50.2	93.2	70	9.0	65
Bomet	0.0	4.7	93.2	2.1	100.0	50.8	43.4	94.1	89	6.1	84
Kakamega	8.8	10.0	80.3	0.9	100.0	21.2	73.9	95.1	130	11.2	123
Vihiga	0.7	7.6	90.5	1.2	100.0	51.1	46.0	97.1	38	12.6	37
Bungoma	3.1	10.5	81.6	4.7	100.0	12.1	79.4	91.5	131	13.6	120
Busia	3.6	8.9	85.8	1.7	100.0	20.3	74.9	95.2	78	6.1	74
Siaya	6.4	3.4	89.6	0.6	100.0	4.7	81.4	86.1	55	10.2	48
Kisumu	3.0	12.0	84.1	0.9	100.0	25.1	70.0	95.1	106	11.1	100
Homa Bay	4.3	10.5	85.2	0.0	100.0	23.8	67.1	91.0	86	3.5	78
Migori	6.0	11.8	81.4	0.7	100.0	37.9	54.3	92.2	139	7.7	128
Kisii	6.7	24.1	69.2	0.0	100.0	31.7	65.7	97.4	84	12.1	81
Nyamira	0.0	15.5	83.9	0.6	100.0	12.5	84.4	96.9	42	9.2	40
Nairobi City	0.7	15.7	83.6	0.0	100.0	49.7	50.3	100.0	611	8.8	611
Total	3.4	11.4	83.1	2.1	100.0	36.4	54.9	91.3	4,115	9.1	3,756

Note: Figures in parentheses are based on 25–49 unweighted cases.
[1] Based on either a written record or the mother's recall

Table 10.2 Possession and observation of vaccination cards according to background characteristics

Percentages of children age 12–23 months and children age 24–35 months who ever had a vaccination card, and percentages with a vaccination card observed, according to background characteristics, Kenya Mini DHS 2025–26

	Children age 12–23 months			Children age 24–35 months		
Background characteristic	Percentage who ever had a vaccination card[1]	Percentage with a vaccination card seen[1]	Number of children	Percentage who ever had a vaccination card[1]	Percentage with a vaccination card seen[1]	Number of children
Sex						
Male	97.8	67.6	1,006	98.1	54.6	1,018
Female	98.6	70.7	929	98.5	56.8	988
Birth order						
1	98.0	68.9	608	98.0	52.5	599
2–3	98.7	69.1	850	99.2	56.0	902
4–5	98.2	70.1	334	97.9	60.0	343
6+	96.5	67.3	142	95.1	56.6	162
Residence						
Urban	98.3	61.7	784	99.1	44.0	906
Rural	98.2	74.1	1,150	97.6	65.3	1,100
Mother's education[2]						
No education	93.6	55.7	173	93.7	44.9	171
Primary	98.7	77.9	518	98.4	63.7	584
Secondary	98.2	65.8	858	98.6	55.4	819
More than secondary	99.6	70.7	385	99.3	49.5	432
Wealth quintile						
Lowest	96.2	72.4	413	95.8	63.7	394
Second	99.4	76.8	359	98.7	64.1	374
Middle	96.9	68.8	375	98.1	65.2	363
Fourth	99.5	63.4	432	99.1	44.7	449
Highest	99.1	64.6	356	99.5	44.3	427
Total	98.2	69.1	1,934	98.3	55.7	2,007

[1] Mother and child handbook or other home-based record

[2] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 10.3 Vaccinations by source of documentation

Percentages of children age 12–23 months and children age 24–35 months who received specific vaccines at any time before the survey, by source of information (vaccination card or mother's report), and percentage who received specific vaccines by the appropriate age, Kenya Mini DHS 2025–26

	Children age 12–23 months				Children age 24–35 months			
	Vaccinated at any time before survey				Vaccinated at any time before survey			
Vaccine	Vaccination card[1]	Mother's report	Either source (crude coverage)	Vaccinated by appro-priate age[2,3,4]	Vaccination card[1]	Mother's report	Either source (crude coverage)	Vaccinated by appro-priate age[2,3,4]
BCG	67.9	29.7	97.6	96.5	55.3	42.7	98.0	97.1
DPT-HepB-Hib								
1	68.7	29.1	97.9	97.6	55.6	41.8	97.4	97.1
2	68.2	26.3	94.5	93.8	55.4	38.5	93.9	93.4
3	65.5	24.3	89.9	88.9	54.5	36.2	90.6	89.4
Polio								
OPV 0 (birth dose)	60.6	27.4	87.9	87.2	51.9	38.3	90.2	89.0
OPV 1	68.7	29.2	97.9	97.6	55.6	40.6	96.2	95.8
OPV 2	67.2	27.2	94.4	93.8	55.2	38.1	93.3	92.7
OPV 3	62.5	11.5	74.0	71.9	52.8	16.0	68.8	67.9
IPV	57.8	28.5	86.4	84.8	50.8	41.1	91.9	90.6
Pneumococcal								
1	68.5	29.3	97.8	97.5	55.5	42.1	97.6	97.3
2	68.1	28.8	96.9	96.4	55.1	40.7	95.8	95.0
3	65.3	27.1	92.4	90.8	53.4	39.0	92.4	89.4
Rotavirus								
1	68.1	28.3	96.4	96.1	54.5	41.1	95.6	95.2
2	66.7	27.5	94.2	93.6	52.4	39.5	91.9	90.7
3	56.4	24.1	80.5	79.5	43.5	34.6	78.1	77.0
MR								
1	61.3	27.4	88.7	84.9	53.3	40.2	93.5	87.1
2	na	na	na	na	40.8	31.2	72.0	70.3
Yellow fever[5]	4.5	5.4	9.8	8.6	3.8	10.1	13.9	11.2
Fully vaccinated (basic antigens)[6]	57.8	21.7	79.5	73.2	52.3	32.8	85.1	77.6
Fully vaccinated (according to national schedule)[7]	37.8	7.8	45.6	41.5	29.0	9.3	38.3	32.6
No vaccinations	0.1	1.0	1.1	na	0.1	1.5	1.6	na
Number of children	1,337	598	1,934	1,934	1,117	889	2,007	2,007

BCG = Bacille Calmette-Guérin
DPT = diphtheria, pertussis, and tetanus
HepB = hepatitis B
Hib = *Haemophilus influenzae* type b
OPV = Oral polio vaccine
IPV = Inactivated polio vaccine
MR = Measles-rubella
na = Not applicable
[1] Mother and child handbook or other home-based record
[2] Received by age 12 months
[3] For children whose vaccination information is based on the mother's report, date of vaccination is not collected. The proportions of vaccinations given during the first and second years of life are assumed to be the same as for children with a written record of vaccination.
[4] Received by age 12 months for all vaccines except for second dose of MR, which should be received by age 24 months
[5] Only given to children in high-risk counties (currently Baringo, Elgeyo, Marakwet, West Pokot, and Turkana)
[6] BCG, three doses of DPT-HepB-Hib, three doses of polio vaccine (excluding polio vaccine given at birth), and one dose of MR
[7] BCG, three doses of DPT-HepB-Hib, four doses of OPV, one dose of IPV, three doses of pneumococcal vaccine, three doses of rotavirus vaccine, and one dose of MR; for children age 24–35 months, all of these plus a second dose of MR

Table 10.4 Vaccinations by background characteristics

Percentages of children age 12–23 months and children age 24–35 months who received specific vaccines at any time before the survey (according to a vaccination card or the mother's report), percentages fully vaccinated (basic antigens), percentages fully vaccinated according to national schedule, and percentages who received no vaccinations, according to background characteristics, Kenya Mini DHS 2025–26

	Children age 12–13 months																				Children age 24–35 months		
		DPT-HepB-Hib			Polio[1]					Pneumococcal			Rotavirus										
Background characteristic	BCG	1	2	3	OPV 0 (birth dose)	OPV 1	OPV 2	OPV 3	IPV	1	2	3	1	2	3	MR 1	Yellow fever[2]	Fully vaccinated (basic antigens)[3]	Fully vaccinated according to national schedule[4]	No vaccinations	Number of children	Measles 2	Fully vaccinated according to national schedule[5]	Number of children
Sex																								
Male	97.1	97.8	93.9	88.9	87.3	98.3	94.1	73.6	87.0	98.1	97.1	92.3	96.2	94.3	81.1	88.8	9.1	77.9	48.1	1.0	1,006	70.4	38.4	1,018
Female	98.1	97.9	95.2	90.9	88.7	97.6	94.8	74.4	85.7	97.5	96.7	92.4	96.7	94.2	79.9	88.7	10.6	81.3	42.9	1.2	929	73.6	38.3	988
Birth order																								
1	98.1	97.6	94.0	90.8	88.1	98.5	95.7	74.4	87.5	97.8	97.1	93.2	96.5	95.3	84.1	91.3	10.7	81.6	47.8	1.0	608	76.0	38.0	599
2–3	98.2	98.2	95.4	90.8	90.2	98.3	94.8	75.5	85.4	97.9	97.3	92.6	97.1	94.8	80.5	90.4	9.9	82.4	46.9	0.8	850	73.0	42.2	902
4–5	96.6	97.8	94.4	89.8	85.4	96.4	92.0	72.9	88.1	97.6	96.6	92.8	95.1	92.4	76.9	85.2	8.8	75.4	42.8	1.8	334	68.3	32.6	343
6+	94.5	96.9	91.5	80.5	79.6	96.7	92.5	66.2	82.6	97.6	95.0	86.7	94.7	90.3	73.4	75.7	8.2	62.8	35.1	1.8	142	59.4	30.2	162
Vaccination card[6]																								
Seen	98.2	99.4	98.7	94.8	87.7	99.5	97.3	90.4	83.7	99.1	98.5	94.5	98.6	96.5	81.6	88.7	6.5	83.7	54.7	0.1	1,337	73.3	52.1	1,117
Not seen or no longer has	98.2	96.2	86.7	80.4	90.8	96.2	89.8	38.7	94.3	96.7	95.3	89.5	93.5	91.0	79.8	90.7	18.1	71.7	26.7	1.5	563	72.2	21.6	855
Never had	65.3	62.8	59.6	51.1	51.7	67.0	59.8	14.0	61.6	65.1	61.4	56.2	59.3	58.1	47.6	55.9	6.6	44.0	3.0	33.0	35	24.8	8.4	35
Residence																								
Urban	98.5	98.4	94.9	91.6	91.1	98.9	95.9	74.7	85.2	98.3	98.2	94.2	97.2	96.5	84.7	91.8	9.7	82.5	46.6	0.7	784	75.4	38.6	906
Rural	97.0	97.5	94.2	88.7	85.7	97.3	93.4	73.5	87.1	97.5	96.1	91.1	95.9	92.7	77.6	86.6	9.9	77.5	45.0	1.4	1,150	69.2	38.1	1,100
Mother's education[7]																								
No education	92.7	94.2	89.7	80.6	66.8	92.4	86.8	59.9	80.0	94.7	92.0	82.4	91.6	85.7	66.9	72.9	7.3	61.8	22.9	4.2	173	41.6	18.2	171
Primary	95.7	96.8	94.1	87.9	82.9	97.1	91.3	72.7	83.7	96.7	95.5	91.1	96.1	92.5	78.4	81.8	7.1	71.1	42.9	1.7	518	66.9	36.0	584
Secondary	99.0	98.9	96.0	92.6	92.2	98.9	95.9	74.8	89.3	98.7	98.0	93.8	96.5	94.9	83.2	93.0	10.3	84.9	50.2	0.6	858	76.5	41.2	819
More than secondary	99.4	98.8	93.8	90.5	94.7	99.5	98.6	80.2	86.2	98.8	98.8	95.3	99.0	98.7	83.4	95.6	13.6	86.7	49.3	0.1	385	82.4	44.0	432

Continued…

Table 10.4—*Continued*

Background characteristic	Children age 12–13 months																					Children age 24–35 months		
		DPT-HepB-Hib			Polio[1]					Pneumococcal			Rotavirus											
	BCG	1	2	3	OPV 0 (birth dose)	OPV 1	OPV 2	OPV 3	IPV	1	2	3	1	2	3	MR 1	Yellow fever[2]	Fully vaccinated (basic antigens)[3]	Fully vaccinated according to national schedule[4]	No vaccinations	Number of children	Measles 2	Fully vaccinated according to national schedule[5]	Number of children
Wealth quintile																								
Lowest	94.2	96.0	91.1	83.8	74.1	95.3	88.0	66.6	81.0	95.6	92.5	84.8	92.3	87.7	70.2	76.7	10.4	65.2	31.7	2.7	413	57.2	28.9	394
Second	98.3	98.9	96.5	91.0	91.6	98.1	94.8	76.0	89.6	98.8	98.0	93.5	98.9	94.3	79.6	90.4	9.1	80.1	51.2	0.5	359	71.3	39.1	374
Middle	98.5	95.5	93.0	89.4	91.5	97.8	95.0	73.5	91.9	97.1	97.0	94.2	95.7	94.6	84.7	89.3	8.8	82.6	49.7	1.1	375	77.4	40.3	363
Fourth	97.5	99.1	95.2	91.1	87.8	98.7	95.8	74.1	84.7	97.9	97.6	93.4	96.4	95.9	81.9	92.0	11.4	80.6	45.7	0.9	432	78.5	45.8	449
Highest	100.0	100.0	97.1	94.7	96.7	100.0	99.2	80.9	85.5	100.0	100.0	96.7	99.5	99.2	87.3	96.3	9.3	90.7	51.7	0.0	356	74.8	36.9	427
Total	97.6	97.9	94.5	89.9	87.9	97.9	94.4	74.0	86.4	97.8	96.9	92.4	96.4	94.2	80.5	88.7	9.8	79.5	45.6	1.1	1,934	72.0	38.3	2,007

BCG = Bacille Calmette-Guérin
DPT = diphtheria, pertussis, and tetanus
HepB = hepatitis B
Hib = *Haemophilus influenzae* type b
OPV = oral polio vaccine
IPV = inactivated polio vaccine
MR = measles-rubella
[1] OPV 0 is the polio vaccination given at birth.
[2] Only required for children in high-risk counties (currently Baringo, Elgeyo, Marakwet, West Pokot, and Turkana)
[3] BCG, three doses of DPT-HepB-Hib, three doses of polio vaccine (excluding polio vaccine given at birth), and one dose of MR
[4] BCG, three doses of DPT-HepB-Hib, four doses of OPV, one dose of IPV, three doses of pneumococcal vaccine, three doses of rotavirus vaccine, and one dose of MR
[5] BCG, three doses of DPT-HepB-Hib, four doses of OPV, one dose of IPV, three doses of pneumococcal vaccine, three doses of rotavirus vaccine, and two doses of MR
[6] Mother and child handbook or other home-based record
[7] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 10.5 Source of vaccinations

Among children who received at least one vaccination, percent distributions of children age 12–23 months and children age 24–35 months by source of most vaccinations, according to background characteristics, Kenya Mini DHS 2025–26

	Children age 12–23 months who received at least one vaccination							Children age 24–35 months who received at least one vaccination						
	Source of most vaccinations							Source of most vaccinations						
Background characteristic	Public medical sector	Private medical sector (non-NGO)	Private medical sector (NGO)	Private medical sector (FBO)	Other	Total	Number of children	Public medical sector	Private medical sector (non-NGO)	Private medical sector (NGO)	Private medical sector (FBO)	Other	Total	Number of children
Sex														
Male	84.1	13.9	0.1	1.7	0.2	100.0	996	85.4	12.5	0.1	1.8	0.3	100.0	1,002
Female	81.4	17.6	0.0	0.8	0.2	100.0	918	84.4	13.4	0.0	2.2	0.1	100.0	972
Birth order														
1	86.5	11.3	0.0	1.9	0.4	100.0	602	84.4	14.0	0.1	1.5	0.0	100.0	596
2–3	77.7	21.5	0.1	0.7	0.0	100.0	844	81.6	15.8	0.0	2.5	0.2	100.0	888
4–5	85.5	12.9	0.0	1.5	0.1	100.0	328	90.8	6.6	0.0	2.2	0.4	100.0	335
6+	91.4	5.9	0.5	1.4	0.8	100.0	139	92.4	6.5	0.0	0.9	0.2	100.0	155
Residence														
Urban	71.9	27.3	0.1	0.7	0.1	100.0	779	77.5	21.3	0.1	1.1	0.0	100.0	899
Rural	90.3	7.7	0.1	1.6	0.3	100.0	1,135	91.1	5.9	0.0	2.7	0.3	100.0	1,076
Mother's education[1]														
No education	87.7	10.2	0.3	1.2	0.7	100.0	166	85.4	13.6	0.0	0.5	0.5	100.0	164
Primary	91.2	6.9	0.1	1.7	0.0	100.0	509	94.3	4.2	0.0	1.2	0.3	100.0	566
Secondary	83.8	14.7	0.0	1.4	0.1	100.0	854	85.6	11.4	0.0	3.0	0.0	100.0	813
More than secondary	67.4	31.8	0.0	0.4	0.5	100.0	385	71.0	26.9	0.1	1.7	0.2	100.0	432
Wealth quintile														
Lowest	94.5	3.7	0.0	1.5	0.3	100.0	402	95.7	2.5	0.0	1.6	0.3	100.0	381
Second	91.1	6.8	0.0	1.7	0.4	100.0	357	92.8	5.9	0.0	1.3	0.0	100.0	367
Middle	87.0	11.2	0.2	1.5	0.1	100.0	371	88.4	6.7	0.0	4.6	0.3	100.0	357
Fourth	80.5	18.1	0.0	1.3	0.1	100.0	428	87.8	10.4	0.1	1.4	0.3	100.0	443
Highest	59.6	40.0	0.1	0.3	0.0	100.0	356	62.4	36.1	0.0	1.5	0.0	100.0	427
Total	82.8	15.7	0.1	1.3	0.2	100.0	1,914	84.9	12.9	0.0	2.0	0.2	100.0	1,974

NGO = nongovernmental organisation
FBO = faith-based organisation
[1] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 10.6.1 Children with symptoms of acute respiratory infection and care seeking for symptoms

Among children under age 5, percentage who had symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey; and among children with symptoms of ARI in the 2 weeks preceding the survey, percentage for whom advice or treatment was sought, according to background characteristics, Kenya Mini DHS 2025–26

	Among children under age 5:		Among children under age 5 with symptoms of ARI:		
Background characteristic	Percentage with symptoms of ARI[1]	Number of children	Percentage for whom advice or treatment was sought[2]	Percentage for whom advice or treatment was sought the same or next day[2]	Number of children
Age in months					
<6	2.0	1,047	*	*	21
6–11	1.5	994	*	*	15
12–23	2.1	1,934	(77.1)	(50.7)	40
24–35	0.8	2,007	(70.3)	(43.3)	17
36–47	1.8	2,136	84.2	62.1	38
48–59	1.9	2,048	(65.5)	(30.1)	39
Sex					
Male	1.8	5,081	80.0	50.6	89
Female	1.6	5,086	68.3	49.8	80
Cooking fuels and technologies					
Clean fuel and technology[3]	1.2	3,401	*	*	42
Solid fuel[4]	1.8	6,549	75.6	51.9	120
Gasoline/diesel	*	19	*	*	0
Kerosene/paraffin	(3.6)	185	*	*	7
No food cooked in household	*	13	*	*	0
Residence					
Urban	1.4	4,444	(71.3)	(45.9)	61
Rural	1.9	5,722	76.3	52.6	108
Mother's education[5]					
No education	1.8	952	83.3	44.8	17
Primary	2.2	3,106	80.7	59.1	69
Secondary	1.5	4,055	66.9	48.2	60
More than secondary	1.1	2,053	*	*	23
Wealth quintile					
Lowest	2.2	2,185	76.0	42.8	47
Second	2.1	1,875	(72.5)	(59.2)	39
Middle	1.4	1,819	(73.0)	(51.7)	26
Fourth	1.6	2,271	(79.3)	(55.9)	37
Highest	1.0	2,016	*	*	19
Total	1.7	10,166	74.5	50.2	169

Notes: Figures in parentheses are based on 25–49 unweighted cases. An asterisk indicates that a figure is based on fewer than 25 unweighted cases and has been suppressed.

[1] Symptoms of ARI include short, rapid breathing that is chest related and/or difficult breathing that is chest related.

[2] Includes advice or treatment from the following sources: public medical sector, private medical sector (non-nongovernmental organisation), private medical sector (nongovernmental organisation), private medical sector (faith-based organisation), shop, market, and itinerant drug seller; excludes advice or treatment from a traditional practitioner

[3] Includes stove/cookers using electricity, liquefied petroleum gas (LPG) /natural gas/ biogas, solar, and alcohol/ethanol

[4] Includes coal/lignite, charcoal, wood, straw/shrubs/grass, agricultural crops, animal dung/waste, processed biomass (pellets) or woodchips, garbage/plastic, and sawdust

[5] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 10.6.2 Treatment for children with symptoms of acute respiratory infection

Among children under age 5 who had symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey, percentage given antibiotics and other treatments and percentage given no treatment, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Percentage of children with symptoms of ARI who were given:					Percentage given no treatment	Number of children with ARI
	Antibiotic	Other medicine[1]	Oxygen	Antimalarial (ACT)	Non-ACT antimalarial		
Age in months							
<6	*	*	*	*	*	*	21
6–11	*	*	*	*	*	*	15
12–23	(61.4)	(72.0)	(0.0)	(7.7)	(5.4)	(20.7)	40
24–35	(50.3)	(67.4)	(2.2)	(9.2)	(3.6)	(22.2)	17
36–47	60.3	76.0	0.9	13.4	1.6	9.7	38
48–59	(39.8)	(39.4)	(0.0)	(13.2)	(2.4)	(32.8)	39
Sex							
Male	67.7	71.5	0.0	10.4	6.4	12.5	89
Female	37.4	45.1	2.8	8.0	1.3	38.5	80
Residence							
Urban	(55.8)	(49.4)	(0.0)	(1.7)	(0.0)	(34.0)	61
Rural	51.9	64.4	2.1	13.6	6.2	19.6	108
Mother's education[2]							
No education	61.9	72.4	1.9	6.0	7.7	9.6	17
Primary	65.1	62.8	2.2	8.8	3.2	15.7	69
Secondary	36.9	45.6	0.0	12.1	5.4	43.4	60
More than secondary	*	*	*	*	*	*	23
Wealth quintile							
Lowest	56.0	65.3	0.7	6.0	10.8	20.9	47
Second	(40.5)	(57.2)	(4.0)	(16.4)	(1.8)	(32.0)	39
Middle	(44.4)	(67.2)	(0.0)	(21.1)	(3.6)	(14.9)	26
Fourth	(62.3)	(40.7)	(1.0)	(2.4)	(0.0)	(33.1)	37
Highest	*	*	*	*	*	*	19
Total	53.3	59.0	1.3	9.3	4.0	24.8	169

Notes: Figures in parentheses are based on 25–49 unweighted cases. An asterisk indicates that a figure is based on fewer than 25 unweighted cases and has been suppressed.
ACT = artemisinin-based combination therapy
[1] Other medicine includes aspirin, paracetamol, and ibuprofen.
[2] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 10.7 Source of advice or treatment for children with symptoms of acute respiratory infection

Percentage of children under age 5 with symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey for whom advice or treatment was sought from specific sources; and among children under age 5 with symptoms of ARI in the 2 weeks preceding the survey for whom advice or treatment was sought, percentage for whom advice or treatment was sought from specific sources, Kenya Mini DHS 2025–26

	Percentage for whom advice or treatment was sought from each source:	
Source	Among children with symptoms of ARI[1]	Among children with symptoms of ARI for whom advice or treatment was sought[1]
Public sector	41.9	56.3
Government hospital	15.4	20.7
Government health centre	17.5	23.6
Government dispensary	10.6	14.2
Private medical sector (non-NGO)	33.2	44.5
Private hospital	6.4	8.6
Private clinic	4.4	5.9
Pharmacy	22.4	30.0
Other private sector	0.5	0.7
Shop	0.4	0.5
Itinerant drug seller	0.1	0.1
Other	1.3	1.7
Number of children	169	126

Note: Advice or treatment for children with symptoms of ARI may have been sought from more than one source.
NGO = nongovernmental organisation
[1] Symptoms of ARI include short, rapid breathing that is chest related and/or difficult breathing that is chest related.

Table 10.8 Children with fever and care seeking for fever

Among children under age 5, percentage who had a fever in the 2 weeks preceding the survey; and among children with fever in the 2 weeks preceding the survey, percentage for whom advice or treatment was sought, percentage for whom advice or treatment was sought the same or next day following the onset of fever, and percentage who received antibiotics as treatment, according to background characteristics, Kenya Mini DHS 2025–26

	Among children under age 5:		Among children under age 5 with fever:			
Background characteristic	Percentage with fever	Number of children	Percentage for whom advice or treatment was sought[1]	Percentage for whom advice or treatment was sought the same or next day[1]	Percentage who took antibiotics	Number of children with fever
Age in months						
<6	10.2	1,047	67.7	52.0	28.9	107
6–11	17.1	994	56.4	37.2	29.9	170
12–23	17.3	1,934	71.6	47.7	39.6	335
24–35	15.2	2,007	72.3	54.1	35.3	304
36–47	14.2	2,136	62.7	29.1	41.6	303
48–59	9.0	2,048	72.3	46.5	32.5	185
Sex						
Male	13.7	5,081	67.4	44.1	36.0	697
Female	13.9	5,086	68.1	43.9	36.3	706
Residence						
Urban	14.6	4,444	70.4	47.0	38.2	649
Rural	13.2	5,722	65.5	41.4	34.4	754
Mother's education[2]						
No education	10.7	952	62.8	37.9	24.0	102
Primary	14.7	3,106	66.1	40.9	30.9	458
Secondary	13.6	4,055	66.8	43.6	36.5	550
More than secondary	14.3	2,053	73.9	51.6	48.1	293
Wealth quintile						
Lowest	12.3	2,185	62.0	33.0	33.3	269
Second	15.0	1,875	65.1	39.0	34.5	281
Middle	13.1	1,819	66.7	50.4	31.0	238
Fourth	13.5	2,271	73.3	46.0	32.9	307
Highest	15.3	2,016	70.6	51.2	47.4	308
Total	13.8	10,166	67.8	44.0	36.2	1,403

[1] Includes advice or treatment from the following sources: public medical sector, private medical sector (non-nongovernmental organisation), private medical sector (nongovernmental organisation), private medical sector (faith-based organisation), shop, market, and itinerant drug seller; excludes advice or treatment from a traditional practitioner

[2] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 10.9 Children with diarrhoea and care seeking for diarrhoea

Percentage of children under age 5 who had diarrhoea in the 2 weeks preceding the survey; and among children with diarrhoea in the 2 weeks preceding the survey, percentage for whom advice or treatment was sought, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Percentage with diarrhoea	Number of children	Among children under age 5 with diarrhoea: Percentage for whom advice or treatment was sought[1]	Among children under age 5 with diarrhoea: Number of children with diarrhoea
Age in months				
<6	11.0	1,047	44.1	115
6–11	19.4	994	55.5	192
12–23	17.5	1,934	54.7	338
24–35	11.3	2,007	67.9	227
36–47	8.5	2,136	64.1	181
48–59	4.5	2,048	53.3	91
Sex				
Male	11.8	5,081	58.0	600
Female	10.7	5,086	57.5	546
Source of drinking water[2]				
Improved	11.3	8,318	57.5	937
Unimproved	12.9	484	55.5	62
Surface	10.7	1,364	60.7	146
Type of toilet facility[3]				
Improved sanitation facility	11.7	7,321	58.5	856
Unimproved facility	10.6	1,980	53.2	210
Open defecation	9.3	865	61.7	80
Residence				
Urban	12.5	4,444	59.0	554
Rural	10.3	5,722	56.6	592
Mother's education[4]				
No education	7.3	952	62.1	69
Primary	12.3	3,106	56.1	381
Secondary	11.3	4,055	53.1	457
More than secondary	11.6	2,053	68.0	239
Wealth quintile				
Lowest	10.0	2,185	54.2	219
Second	11.0	1,875	55.8	206
Middle	11.5	1,819	56.2	210
Fourth	12.7	2,271	58.5	287
Highest	11.1	2,016	63.6	223
Total	11.3	10,166	57.8	1,146

Note: Advice or treatment for children with diarrhoea may have been sought from more than one source.

[1] Includes advice or treatment from the following sources: public medical sector, private medical sector (non-nongovernmental organisation), private medical sector (nongovernmental organisation), private medical sector (faith-based organisation), shop, market, and itinerant drug seller; excludes advice or treatment from a traditional practitioner

[2] See Table 13.1 for definition of categories.

[3] See Table 13.3 for definition of categories.

[4] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 10.10 Feeding practices during diarrhoea

Percent distribution of children under age 5 who had diarrhoea in the 2 weeks preceding the survey by amount of liquids and food given compared with normal practice, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Amount of liquids given							Amount of food given								Number of children with diarrhoea
	More	Same as usual	Somewhat less	Much less	None	Don't know/ missing	Total	More	Same as usual	Somewhat less	Much less	None	Never gave food	Don't know/ missing	Total	
Age in months																
<6	23.8	54.8	14.4	1.4	5.7	0.0	100.0	2.4	36.7	6.5	7.1	0.0	47.3	0.0	100.0	115
6–11	18.8	45.4	21.3	10.5	4.0	0.0	100.0	2.3	36.0	33.3	12.3	5.9	10.2	0.0	100.0	192
12–23	25.5	41.5	19.1	13.0	0.9	0.0	100.0	3.8	38.3	30.3	20.1	5.3	2.0	0.2	100.0	338
24–35	22.9	45.9	20.2	8.6	2.0	0.4	100.0	2.7	39.0	33.5	16.3	5.9	2.2	0.4	100.0	227
36–47	38.8	37.9	11.6	9.8	1.9	0.0	100.0	17.0	36.4	26.9	15.1	4.1	0.5	0.0	100.0	181
48–59	28.4	40.4	24.2	4.8	0.4	1.8	100.0	3.6	33.5	50.2	8.2	4.4	0.0	0.0	100.0	91
Sex																
Male	26.8	45.5	16.6	9.6	1.6	0.0	100.0	5.0	40.7	27.3	15.8	6.2	4.8	0.0	100.0	600
Female	25.2	41.8	20.5	9.1	3.0	0.5	100.0	5.5	33.4	33.1	14.1	3.1	10.6	0.3	100.0	546
Breastfeeding status[1]																
Breastfeeding	23.3	44.4	19.2	10.1	3.0	0.0	100.0	2.4	36.1	27.5	14.8	5.2	14.0	0.0	100.0	578
Not breastfeeding	22.7	47.0	19.3	9.1	1.5	0.3	100.0	4.1	41.1	30.9	17.3	4.3	1.7	0.5	100.0	295
Residence																
Urban	27.8	48.2	15.3	6.4	2.1	0.3	100.0	6.4	40.6	30.1	12.7	3.0	7.2	0.1	100.0	554
Rural	24.4	39.6	21.4	12.1	2.4	0.1	100.0	4.2	34.1	30.1	17.1	6.4	8.0	0.1	100.0	592
Mother's education[2]																
No education	31.4	35.9	19.2	12.6	0.9	0.0	100.0	9.2	28.8	32.5	14.5	9.1	5.9	0.0	100.0	69
Primary	23.8	41.1	21.4	11.5	2.3	0.0	100.0	5.0	33.4	34.1	14.7	4.5	8.2	0.2	100.0	381
Secondary	24.3	45.6	19.0	8.7	2.2	0.2	100.0	6.4	40.2	26.6	15.7	4.9	6.0	0.2	100.0	457
More than secondary	31.2	46.5	12.4	6.4	2.7	0.7	100.0	2.3	40.0	29.7	14.2	3.5	10.2	0.0	100.0	239
Wealth quintile																
Lowest	21.7	41.6	20.9	13.6	2.2	0.0	100.0	5.0	33.7	29.1	18.7	5.5	8.0	0.0	100.0	219
Second	21.1	41.1	25.5	10.8	1.5	0.0	100.0	3.1	35.6	32.3	16.4	5.6	7.0	0.0	100.0	206
Middle	29.6	37.6	19.2	11.7	1.9	0.0	100.0	3.3	39.0	29.8	15.5	7.0	5.1	0.3	100.0	210
Fourth	25.1	48.4	13.9	8.7	3.5	0.3	100.0	4.4	39.8	25.1	17.6	4.7	8.1	0.3	100.0	287
Highest	32.4	48.0	14.6	2.5	1.7	0.7	100.0	10.3	37.1	35.7	6.2	1.1	9.6	0.0	100.0	223
Total	26.0	43.7	18.4	9.4	2.3	0.2	100.0	5.2	37.2	30.1	15.0	4.7	7.6	0.1	100.0	1,146

Note: The recommendation is that children should be given more liquids than usual and at least as much food as usual during diarrhoea.
[1] Breastfeeding status is captured for children age 0–35 months only.
[2] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 10.11 Oral rehydration salts, zinc, continued feeding, and other treatments for diarrhoea

Among children under age 5 who had diarrhoea in the 2 weeks preceding the survey, percentage given fluids from an oral rehydration salts (ORS) packet or pre-packaged ORS fluid; zinc; ORS and zinc; ORS and continued feeding; ORS, zinc, and continued feeding; ORS or increased fluids; recommended homemade fluids (RHF); oral rehydration therapy (ORT); ORT and continued feeding; and other treatments, and percentage given no treatment, according to background characteristics, Kenya Mini DHS 2025–26

	Percentage of children with diarrhoea who were given:														
Background characteristic	Fluids from ORS packet or pre-packaged ORS fluid	Zinc	ORS and zinc	ORS and continued feeding[1]	ORS, zinc, and continued feeding[1]	ORS or increased fluids	RHF	ORT (ORS, RHF, or increased fluids)	ORT and continued feeding[1]	Antibiotic	Antimotility agent	Intravenous solution	Home remedy/ other	Percentage given no treatment	Number of children with diarrhoea
Age in months															
<6	14.2	20.5	7.4	8.8	4.0	35.3	17.6	47.3	21.1	15.7	4.5	1.9	4.3	34.9	115
6–11	32.5	35.5	22.8	21.0	14.0	43.3	15.4	49.5	36.3	9.1	1.4	0.3	9.3	32.1	192
12–23	45.2	36.5	30.2	29.6	19.2	59.2	18.6	64.8	45.4	10.8	4.0	0.0	13.6	23.8	338
24–35	50.3	44.0	32.6	38.0	24.9	60.9	26.6	68.8	52.2	9.5	3.2	0.5	10.8	17.3	227
36–47	45.3	32.1	19.7	35.6	14.4	63.7	25.0	71.9	55.7	9.4	3.1	0.7	4.9	15.6	181
48–59	50.8	44.7	39.7	45.7	37.4	67.3	37.6	71.1	62.2	5.8	1.3	1.1	7.6	19.0	91
Sex															
Male	43.6	36.3	27.9	30.3	19.0	57.0	22.4	64.8	45.8	9.4	2.2	0.2	9.5	23.6	600
Female	39.0	36.0	24.4	29.6	18.2	54.5	21.7	60.8	45.7	10.9	4.1	0.9	9.5	23.1	546
Residence															
Urban	41.1	37.4	24.0	32.6	18.4	57.7	19.3	63.0	48.8	7.8	2.2	0.2	7.5	22.8	554
Rural	41.8	35.1	28.3	27.5	18.8	54.0	24.7	62.7	42.9	12.3	3.9	0.9	11.4	23.9	592
Mother's education[2]															
No education	47.6	46.9	39.1	37.4	29.3	60.1	28.1	65.2	47.4	10.2	2.6	0.4	10.6	20.4	69
Primary	42.9	39.8	29.5	30.5	21.4	58.0	23.6	64.0	48.5	7.2	3.2	0.4	10.5	21.5	381
Secondary	38.2	33.0	25.2	27.4	16.6	52.4	21.2	61.7	45.2	12.5	3.6	0.8	8.9	25.4	457
More than secondary	43.4	33.4	19.3	31.8	15.2	57.7	19.6	62.5	41.8	10.3	2.0	0.3	8.8	23.2	239
Wealth quintile															
Lowest	42.2	37.8	30.6	30.0	21.6	54.3	24.7	63.5	43.5	8.4	4.1	0.3	11.1	26.9	219
Second	38.7	33.6	24.7	24.7	16.2	48.5	23.8	54.8	38.7	10.5	3.7	1.3	18.2	23.7	206
Middle	46.1	36.5	32.0	29.1	20.4	60.6	20.0	67.3	44.6	14.9	3.1	0.6	7.4	22.3	210
Fourth	36.4	38.6	25.0	24.4	17.9	52.2	25.9	61.3	46.6	10.4	1.7	0.3	8.6	21.8	287
Highest	45.3	33.6	19.6	42.7	17.2	64.2	15.0	67.7	54.4	6.8	3.3	0.3	3.2	22.5	223
Total	41.4	36.2	26.2	30.0	18.6	55.8	22.1	62.9	45.7	10.1	3.1	0.5	9.5	23.3	1,146

[1] Continued feeding includes children who were given more food than usual, the same amount of food as usual, or somewhat less food than usual during the diarrhoea episode.
[2] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 10.12 Source of advice or treatment for children with diarrhoea

Percentage of children under age 5 with diarrhoea in the 2 weeks preceding the survey for whom advice or treatment was sought from specific sources; among children under age 5 with diarrhoea in the 2 weeks preceding the survey for whom advice or treatment was sought, percentage for whom advice or treatment was sought from specific sources; among children with diarrhoea who received ORS, percentage for whom advice or treatment was sought from specific sources; and among children with diarrhoea who were given zinc tablets or syrups, percentage for whom advice or treatment was sought from specific sources, Kenya Mini DHS 2025–26

	Percentage for whom advice or treatment was sought from each source:			
Source	Among children with diarrhoea	Among children with diarrhoea for whom advice or treatment was sought	Among children with diarrhoea who received ORS[1]	Among children with diarrhoea who were given zinc
Public sector	**32.7**	**54.8**	**54.8**	**60.1**
Government hospital	14.8	24.8	21.9	28.2
Government health centre	5.0	8.3	9.8	8.1
Government dispensary	13.1	21.9	23.8	24.3
CHW/Fieldworker	0.1	0.2	0.1	0.2
Private medical sector (non-NGO)	**24.5**	**41.0**	**34.3**	**29.4**
Private hospital	9.1	15.2	14.0	14.0
Private clinic	4.0	6.6	5.6	6.5
Pharmacy	11.3	18.9	14.5	8.8
Private doctor	0.3	0.5	0.6	0.4
CHW/Fieldworker	0.1	0.2	0.2	0.2
Private medical sector (NGO)	0.4	0.6	0.4	0.3
Private medical sector (FBO)	**0.4**	**0.6**	**0.4**	**0.3**
FBO/Mission hospital	0.4	0.6	0.4	0.3
Other private sector	**2.2**	**3.7**	**1.7**	**2.6**
Shop	0.3	0.4	0.1	0.6
Traditional practitioner	2.0	3.3	1.6	2.0
Other	0.3	0.5	0.0	0.0
Number of children	1,146	683	475	415

Note: Advice or treatment for children with diarrhoea may have been sought from more than one source.
ORS = oral rehydration salts
CHW = community health worker
NGO = nongovernmental organisation
FBO = faith-based organisation
[1] Fluids from ORS packet or pre-packaged ORS fluid

Table 10.13 Global Scales of Early Development

Percentage of children age 3–23 months who are developmentally on track in health, learning, and psychosocial well-being according to the Global Scales of Early Development (GSED), Kenya Mini DHS 2025–26

Background characteristic	Percentage developmentally on track	Number of children age 3–23 months
Sex		
Male	84.1	1,701
Female	85.6	1,711
Residence		
Urban	87.9	1,437
Rural	82.6	1,975
Age in months		
3–11	89.6	1,524
12–23	81.1	1,888
Mother's education[1]		
No education	69.6	306
Primary	81.5	904
Secondary	87.4	1,467
More than secondary	90.3	734
Wealth quintile		
Lowest	74.5	741
Second	84.6	638
Middle	87.6	609
Fourth	88.8	729
Highest	89.5	696
Total	84.9	3,412

[1] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 10.13C Global Scales of Early Development: County level

Percentage of children age 3–23 months who are developmentally on track in health, learning, and psychosocial well-being according to the Global Scales of Early Development (GSED), Kenya Mini DHS 2025–26

County	Percentage developmentally on track	Number of children age 3–23 months
Mombasa	81.9	110
Kwale	92.8	64
Kilifi	84.8	90
Tana River	82.5	36
Lamu	88.0	9
Taita Taveta	(83.1)	25
Garissa	73.7	37
Wajir	64.1	36
Mandera	61.6	36
Marsabit	73.0	31
Isiolo	78.5	14
Meru	94.4	58
Tharaka-Nithi	(89.3)	18
Embu	(87.9)	21
Kitui	67.5	56
Machakos	73.2	141
Makueni	95.0	46
Nyandarua	82.2	45
Nyeri	(87.7)	34
Kirinyaga	(92.7)	33
Murang'a	93.7	51
Kiambu	95.8	211
Turkana	65.3	52
West Pokot	52.1	67
Samburu	66.9	27
Trans Nzoia	92.2	44
Uasin Gishu	87.4	109
Elgeyo-Marakwet	76.3	27
Nandi	77.9	66
Baringo	82.6	47
Laikipia	94.7	41
Nakuru	93.0	184
Narok	67.6	79
Kajiado	88.8	106
Kericho	88.1	54
Bomet	75.4	77
Kakamega	97.4	113
Vihiga	88.5	32
Bungoma	89.8	105
Busia	89.8	61
Siaya	95.6	45
Kisumu	89.2	87
Homa Bay	81.1	68
Migori	93.6	123
Kisii	90.0	69
Nyamira	74.9	35
Nairobi City	85.4	492
Total	84.9	3,412

Note: Figures in parentheses are based on 25–49 unweighted cases.

Table 10.14 Early Childhood Development Index 2030

Percentage of children age 24–59 months who are developmentally on track in health, learning, and psychosocial well-being according to the Early Childhood Development Index 2030 (ECDI2030),[1] Kenya Mini DHS 2025–26

Background characteristic	Percentage developmentally on track	Number of children age 24–59 months
Sex		
Male	80.3	2,862
Female	85.2	2,882
Residence		
Urban	86.7	2,482
Rural	79.8	3,262
Age in months		
24–35	87.1	1,863
36–47	82.4	2,001
48–59	78.8	1,881
Mother's education[2]		
No education	63.8	582
Primary	78.8	1,922
Secondary	86.5	2,105
More than secondary	92.4	1,135
Wealth quintile		
Lowest	69.8	1,291
Second	82.4	1,076
Middle	85.5	1,030
Fourth	87.5	1,226
Highest	90.5	1,120
Total	82.8	5,744

[1] Sustainable Development Goal Indicator 4.2.1

[2] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 10.14C Early Childhood Development Index 2030: County level

Percentage of children age 24–59 months who are developmentally on track in health, learning, and psychosocial well-being according to the Early Childhood Development Index 2030 (ECDI2030),[1] Kenya Mini DHS 2025–26

County	Percentage developmentally on track	Number of children age 24–59 months
Mombasa	85.8	186
Kwale	74.4	114
Kilifi	87.7	159
Tana River	69.2	55
Lamu	81.7	16
Taita Taveta	76.6	47
Garissa	62.2	72
Wajir	74.6	56
Mandera	62.9	75
Marsabit	58.2	62
Isiolo	59.1	25
Meru	84.7	143
Tharaka-Nithi	82.8	39
Embu	73.7	43
Kitui	58.9	108
Machakos	77.9	167
Makueni	84.8	82
Nyandarua	81.9	51
Nyeri	(96.5)	40
Kirinyaga	89.5	51
Murang'a	89.3	78
Kiambu	85.7	401
Turkana	50.9	96
West Pokot	55.2	100
Samburu	69.4	43
Trans Nzoia	96.5	85
Uasin Gishu	88.0	165
Elgeyo-Marakwet	85.5	52
Nandi	87.1	79
Baringo	78.3	64
Laikipia	83.4	47
Nakuru	87.9	299
Narok	70.8	142
Kajiado	88.4	132
Kericho	93.2	86
Bomet	85.1	134
Kakamega	82.5	179
Vihiga	85.9	55
Bungoma	86.5	159
Busia	94.6	83
Siaya	80.1	103
Kisumu	78.5	136
Homa Bay	84.0	152
Migori	90.5	174
Kisii	92.3	124
Nyamira	80.1	53
Nairobi City	89.1	934
Total	82.8	5,744

Note: Figures in parentheses are based on 25–49 unweighted cases.

[1] Sustainable Development Goal Indicator 4.2.1

Table 10.15 Combined Early Childhood Development Index 2030 and Global Scales of Early Development

Percentage of children age 3–59 months who are developmentally on track in health, learning, and psychosocial well-being according to the combined Early Childhood Development Index 2030 (ECDI2030)[1] and Global Scales of Early Development (GSED), Kenya Mini DHS 2025–26

Background characteristic	Percentage developmentally on track	Number of children age 3–59 months
Sex		
Male	81.7	4,563
Female	85.4	4,593
Residence		
Urban	87.1	3,918
Rural	80.9	5,238
Age in months		
3–11	89.6	1,524
12–23	81.1	1,888
24–35	87.1	1,863
36–47	82.4	2,001
48–59	78.8	1,881
Mother's education[2]		
No education	65.8	888
Primary	79.7	2,826
Secondary	86.8	3,572
More than secondary	91.5	1,869
Wealth quintile		
Lowest	71.5	2,032
Second	83.2	1,715
Middle	86.3	1,639
Fourth	88.0	1,954
Highest	90.1	1,816
Total	83.5	9,156

[1] Sustainable Development Goal Indicator 4.2.1

[2] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 10.15C Combined Early Childhood Development Index 2030 and Global Scales of Early Development: County level

Percentage of children age 3–59 months who are developmentally on track in health, learning, and psychosocial well-being being according to the combined Early Childhood Development Index 2030 (ECDI2030)[1] and Global Scales of Early Development (GSED), Kenya Mini DHS 2025–26

County	Percentage developmentally on track	Number of children age 3–59 months
Mombasa	84.4	297
Kwale	81.0	177
Kilifi	86.6	249
Tana River	74.5	92
Lamu	83.9	25
Taita Taveta	78.8	72
Garissa	66.1	109
Wajir	70.5	92
Mandera	62.5	111
Marsabit	63.2	93
Isiolo	66.2	39
Meru	87.5	201
Tharaka-Nithi	84.9	58
Embu	78.4	64
Kitui	61.8	164
Machakos	75.7	309
Makueni	88.5	128
Nyandarua	82.0	96
Nyeri	92.5	74
Kirinyaga	90.7	84
Murang'a	91.0	129
Kiambu	89.2	612
Turkana	55.9	148
West Pokot	53.9	167
Samburu	68.4	69
Trans Nzoia	95.0	129
Uasin Gishu	87.8	274
Elgeyo-Marakwet	82.3	79
Nandi	82.9	144
Baringo	80.1	110
Laikipia	88.7	88
Nakuru	89.8	483
Narok	69.7	221
Kajiado	88.6	238
Kericho	91.2	140
Bomet	81.6	210
Kakamega	88.3	291
Vihiga	86.8	87
Bungoma	87.8	263
Busia	92.6	144
Siaya	84.7	148
Kisumu	82.6	223
Homa Bay	83.1	221
Migori	91.8	298
Kisii	91.5	192
Nyamira	78.0	88
Nairobi City	87.8	1,426
Total	83.5	9,156

[1] Sustainable Development Goal Indicator 4.2.1

NUTRITION OF CHILDREN 11

Key Findings

- ***Breastfeeding:*** 97% of children born in the 2 years before the survey are breastfed, and 72% of children age 0–5 months are exclusively breastfed.
- ***Minimum acceptable diet:*** 32% of children age 6–23 months are fed a minimum acceptable diet.
- ***Minimum dietary diversity:*** 43% of children are fed with at least five out of eight food groups.
- ***Growth monitoring:*** 49% of children under age 5 had their weight and height measured by a health care provider in the 3 months before the survey.
- ***Vitamin A supplementation:*** 68% of children age 6–59 months received vitamin A supplements in the past 6 months.

Nutrition is the foundation for the health and development of children and adults. This chapter reports on infant and young child feeding (IYCF) practices as well as the key nutritional interventions of IYCF counselling, child growth monitoring, micronutrient supplementation, and deworming.

11.1 GROWTH MONITORING AND PROMOTION

Growth monitoring and promotion programmes use physical growth measurements to monitor children's nutritional status and help caregivers counsel and refer children whose growth appears abnormal (World Health Organization [WHO] 2017a; WHO 2013). An important component of growth monitoring is the regular measurement of children's weight, length/height, and/or mid-upper arm circumference (MUAC). Anthropometry is commonly used to measure child nutritional status.

Weight measured in the past 3 months

Percentage of children under age 5 who had their weight measured in the past 3 months.

Weight and height measured in the past 3 months

Percentage of children under age 5 who had their weight and height measured in the past 3 months.

Mid-upper arm circumference measured in the past 3 months

Percentage of children under age 5 who had their MUAC measured in the past 3 months.

Weight, height, and mid-upper arm circumference measured in the past 3 months

Percentage of children under age 5 who had their weight, height, and MUAC measured in the past 3 months.

Sample: Children under age 5

As shown in **Table 11.1**, 49% of children under age 5 had their weight and height measured by a health care provider in the 3 months preceding the 2025–26 Kenya Mini Demographic and Health Survey (KMDHS). This is an improvement from 37% reported in the 2022 Kenya Demographic and Health Survey (KDHS). Fifty-four percent of children had their weight measured, 50% had their height measured, and 28% had their MUAC measured. Twenty-eight percent had all three measurements (weight, height, and MUAC) taken. Comparison with the 2022 KDHS shows a remarkable improvement in growth monitoring and promotion among children under age 5 in 2025–26.

Patterns by background characteristics

- Seventy-five percent of children age 0–23 months, compared with 33% of children age 24–59 months, had their weight and height measured (**Table 11.1**).
- A slightly higher percentage of children in urban areas (52%) than in rural areas (47%) had their weight and height measured.
- The percentage of children whose weight and height were measured increases with mother's level of education, from 34% among children whose mothers have no education to 49% and 61% among children whose mothers have a secondary and more than a secondary education, respectively.
- The percentage of children whose weight and height were measured increases with wealth, from 43% among those in the lowest wealth quintile to 57% among those in the highest wealth quintile.
- The percentage of children whose weight, height, and MUAC were measured is highest in Kwale County (72%) and lowest in Mandera, Kajiado, and Kakamega Counties (5% each) (**Table 11.1C**).

11.2 Infant and Young Child Feeding Practices

Optimal IYCF practices are critical to the health and survival of young children. Recommended IYCF practices include early initiation of breastfeeding within the first hour after birth; exclusive breastfeeding for the first 2 days after birth; exclusive breastfeeding for the first 6 months of life; continued breastfeeding for at least 2 years; and the introduction of safe, appropriate, and adequate complementary foods at age 6 months (WHO and the United Nations Children's Fund [UNICEF] 2021). This section reports on these IYCF indicators for children under age 2.

11.2.1 Ever Breastfed, Early Initiation of Breastfeeding, and Exclusive Breastfeeding for the First 2 Days After Birth

Breastfeeding both supports children's growth and development and benefits mothers' health. Initiation of breastfeeding within the first hour of birth is important for both the mother and the child. The first breast milk contains colostrum, which is highly nutritious and has antibodies that protect the newborn from infections. Early initiation of breastfeeding also encourages bonding between the mother and her newborn, especially through skin-to-skin contact, which facilitates the production of breast milk. Because feeding newborns anything other than breast milk in the first 2 days after birth can delay early initiation of breastfeeding and interrupt exclusive breastfeeding, it is not recommended unless medically indicated (WHO and UNICEF 2021).

Ever breastfed

Percentage of children born in the 2 years preceding the survey who have ever been breastfed.

Early initiation of breastfeeding

Percentage of children born in the 2 years preceding the survey who were put to the breast within 1 hour of birth.

Exclusive breastfeeding for the first 2 days after birth

Percentage of children born in the 2 years preceding the survey who were fed exclusively with breast milk for the first 2 days after birth.

Sample: Children born in the 2 years preceding the survey

As shown in **Table 11.2**, 97% of children born in the 2 years preceding the survey were ever breastfed. Sixty-three percent of the children were put to the breast within 1 hour of birth, while 83% were exclusively breastfed for the first 2 days after birth.

Patterns by background characteristics

- Eighty-six percent of children whose mothers were counselled on breastfeeding during antenatal care visits were exclusively breastfed for the first 2 days after birth, compared with 32% of children whose mothers did not receive any antenatal care.
- Seventy-one percent of children whose mothers had a vaginal birth were put to the breast within 1 hour of birth, compared with 30% of children whose mothers had a caesarean section. Similarly, 86% of children whose mothers had a vaginal birth were exclusively breastfed for the first 2 days after birth, compared with 72% of children whose mothers had a caesarean birth.
- Children born to mothers in rural areas are more likely to be put to the breast within 1 hour of birth than are those born to mothers in urban areas (67% versus 57%).
- Seventy percent of children born to mothers in the lowest wealth quintile are put to the breast within 1 hour of birth, compared with 50% of children born to mothers in the highest wealth quintile.
- Sixty-seven percent of children born to mothers with no education are breastfed within 1 hour of birth, compared with 57% of children born to mothers with more than a secondary education.

11.2.2 Exclusive Breastfeeding and Mixed Milk Feeding

In their first 6 months, children should be exclusively breastfed (that is, given nothing but breast milk). Exclusive breastfeeding for 6 months lowers the risk of infections that can lead to diarrhoea and respiratory illnesses and provides all the nutrients and liquid an infant requires for optimal growth and development. The production of breast milk is modulated by the frequency and intensity of suckling. Therefore, mixed milk feeding, in which children are fed both breast milk and formula or animal milk within the first 6 months, reduces breast milk output. Mixed feeding in children under age 6 months can also increase their risk of diarrhoea, alter their intestinal microflora, and lead to early cessation of breastfeeding (WHO and UNICEF 2021).

Exclusive breastfeeding under 6 months

Percentage of children age 0–5 months who were fed exclusively with breast milk during the previous day.

Sample: Youngest children age 0–5 months living with the mother

Mixed milk feeding under 6 months

Percentage of children age 0–5 months who were fed both breast milk and formula and/or animal milk during the previous day.


Sample: Youngest children age 0–5 months living with the mother

Seventy-two percent of children age 0–5 months are exclusively breastfed (**Table 11.3**).

Table 11.4 and **Figure 11.1** show the pattern of how children are fed in the first 6 months. At age 0–1 month, 90% of children are exclusively breastfed, which is below the WHO recommendation of 100%. Among children age 0–1 month, 10% receive foods or liquids other than breast milk, including animal or formula milk (3%); plain water (3%); solid, semi-solid, or soft foods (2%); and non-milk liquids (1%). One percent of children age 0–1 month are not breastfed. Exclusive breastfeeding declines to 75% among children age 2–3 months, while 21% of children receive liquids or foods in addition to breast milk at this age. By age 4–5 months, exclusive breastfeeding further declines to 51%; 48% of children age 4–5 months receive liquids or foods in addition to breast milk, with 29% of them consuming solid, semi-solid, or soft foods.

Figure 11.1 **Infant feeding practices by age**


Percent distribution of youngest children age 0–5 months



Trends: The percentage of children age 0–5 months who are exclusively breastfed has improved over time, from 13% in the 2003 KDHS to 60% in the 2022 KDHS to 72% in the 2025–26 KMDHS (**Figure 11.2**).

Figure 11.2 **Trends of exclusive breastfeeding**

Percentage of children age 0–5 months exclusively breastfed



Patterns by background characteristics

- Seventy-four percent of female children age 0–5 months are exclusively breastfed compared with 69% of male children.
- The percentage of children age 0–5 months who are exclusively breastfed is lower among those born to mothers in the lowest wealth quintile (60%) than among those born to mothers in the highest wealth quintile (78%).

- The percentage of children age 0–5 months who are exclusively breastfed is higher in urban areas (76%) than in rural areas (68%).

11.2.3 Continued Breastfeeding and Bottle Feeding

Breastfeeding should continue for at least the first 2 years of life because breast milk lowers children's risk of illness, promotes their recovery during illness, and continues to be an important source of nutrients for healthy growth and development. Continued breastfeeding also has many health benefits for women, including reducing the risks of certain breast and ovarian cancers and the risk of diabetes. Feeding children from a bottle with a nipple is not recommended for children under age 2. Nipples on feeding bottles are susceptible to contamination and increase the risk of disease among children who use them (WHO and UNICEF 2021).

Continued breastfeeding 12–23 months

Percentage of children age 12–23 months who were fed breast milk during the previous day.

Sample: Children age 12–23 months

Bottle feeding

Percentage of children age 0–23 months who were fed from a bottle with a nipple during the previous day.

Sample: Children age 0–23 months

Among children age 12–23 months, 64% are currently breastfeeding. Thirty-seven percent of children age 0–23 months are fed using a bottle with a nipple.

Trends: The proportion of children age 12–23 months who are currently breastfeeding has remained unchanged, at 65% in the 2022 KDHS and 64% in the 2025–26 KMDHS. In contrast, bottle feeding among children age 0–23 months increased from 34% in 2022 to 37% in 2025–26.

Patterns by background characteristic

- Sixty-six percent of children age 12–23 months in rural areas are currently breastfeeding compared with 61% of those in urban areas.
- Forty-three percent of children age 0–23 months in urban areas are using a bottle with a nipple compared with 32% of those in rural areas.
- Fifty-one percent of children age 0–23 months who are in the highest wealth quintile are using a bottle with a nipple compared with 31% of those in the lowest wealth quintile.

11.2.4 Introduction of Complementary Foods

After the first 6 months, breast milk alone is no longer sufficient to meet all the nutritional needs of an infant. After 6 months, appropriate complementary foods should be introduced while breastfeeding continues until age 2 or older. Children are most vulnerable to becoming undernourished during the transition from exclusive breastfeeding to the addition of complementary family foods. During this time, it is important that children receive solid, semi-solid, or soft foods (Dewey 2003; WHO and UNICEF 2021).

Introduction of solid, semi-solid, or soft foods 6–8 months

Percentage of children age 6–8 months who were fed solid, semi-solid, or soft foods during the previous day.

Sample: Youngest children age 6–8 months living with the mother

Eighty-eight percent of children age 6–8 months have been introduced to solid, semi-solid, or soft foods (see **Table 11.9**).

11.2.5 Minimum Dietary Diversity, Minimum Meal Frequency, Minimum Milk Feeding Frequency, Minimum Acceptable Diet, and Egg and/or Flesh Food Consumption

Infants and young children should be fed a minimum acceptable diet, which means they are fed meals with appropriate frequency and variety of foods to meet their energy and nutrient needs. The minimum acceptable diet indicator is a combination of minimum dietary diversity and minimum meal frequency for breastfeeding children, and the same plus minimum milk feeding frequency for non-breastfed children.

Minimum dietary diversity is a proxy for adequate micronutrient density of foods. By consuming food from at least five out of eight food groups, a child has a high likelihood of consuming at least one animal source of food and at least one fruit or vegetable, in addition to a staple food such as grains, roots, or tubers. The eight food groups are breast milk; grains, roots, and tubers; legumes and nuts; dairy products (milk, yogurt, and cheese); flesh foods (meat, fish, poultry, and organ meat); eggs; vitamin A-rich fruits and vegetables; and other fruits and vegetables.

Minimum meal frequency is a proxy for meeting energy requirements. Breastfed children age 6–8 months are considered to be consuming the minimum meal frequency if they receive solid, semi-solid, or soft foods at least twice a day. Breastfed children age 9–23 months are considered to be consuming the minimum meal frequency if they receive solid, semi-solid, or soft foods at least three times a day. Non-breastfed children age 6–23 months are considered to be fed with a minimum meal frequency if they receive solid foods, semi-solid foods, soft foods, or milk feeds at least four times a day and if at least one of the meals is a solid, semi-solid, or soft food.

Minimum milk feeding frequency is a proxy for meeting the nutritional needs of non-breastfed children. Milk and milk products are important sources of nutrients. Non-breastfed children age 6–23 months are considered to be meeting the minimum milk feeding frequency if they receive at least two feeds of milk and/or milk products a day.

Egg and/or flesh food consumption by breastfed and non-breastfed children age 6–23 months increases children's energy, protein, and nutrient intake. Eggs, meat, fish, poultry, and organ meat are important sources of nutrients that support healthy child growth (WHO and UNICEF 2021).

Minimum dietary diversity 6–23 months

Percentage of children age 6–23 months who are fed a minimum of five out of the following eight food groups during the previous day: breast milk; grains, roots, and tubers; legumes and nuts; dairy products (milk, yogurt, and cheese); flesh foods (meat, fish, poultry, and organ meat); eggs; vitamin A-rich fruits and vegetables; and other fruits and vegetables

Minimum meal frequency 6–23 months

Percentage of children age 6–23 months who are fed solid, semi-solid, or soft foods (including milk feeds for non-breastfed children) the minimum recommended number of times or more during the previous day

Minimum milk feeding frequency 6–23 months

Percentage of non-breastfed children age 6–23 months who were given at least two milk feeds during the previous day

Minimum acceptable diet 6–23 months

Percentage of children age 6–23 months who were fed a minimum acceptable diet during the previous day. This indicator is a composite of children who have met minimum dietary diversity and minimum meal frequency, with the extra requirement that non-breastfed children have met minimum milk feeding frequency.

Sample: Youngest children age 6–23 months living with the mother

Egg and/or flesh food consumption 6–23 months

Percentage of children age 6–23 months who were fed egg and/or flesh food during the previous day


Sample: Youngest children age 6–23 months living with the mother

The percentages of children under age 2 who were given various liquids and foods the day before the survey were used to assess whether minimum feeding practice recommendations were being met (**Tables 11.5** and **11.6**).

Overall, 43% of children age 6–23 months meet minimum dietary diversity, and 67% meet minimum meal frequency; however, only 32% meet a minimum acceptable diet (**Table 11.7** and **Figure 11.3**).

Figure 11.3 **IYCF indicators on Minimum Acceptable Diet (MAD), Minimum Meal Frequency (MMF) and Minimum Dietary Diversity (MDD)**

Percentage of children age 6–23 months meeting feeding practice recommendations



Among breastfed children age 6–23 months, 47% meet minimum dietary diversity, 67% meet minimum meal frequency, and 36% meet a minimum acceptable diet. Among non-breastfed children age 6–23 months, 70% meet minimum meal frequency, and 54% meet minimum milk feeding frequency; however, only 29% meet minimum dietary diversity, and only 21% meet a minimum acceptable diet (**Table 11.7**).

Overall, 35% of children age 6–23 months consumed eggs and/or flesh foods the day preceding the survey (**Table 11.8**).


Trends: The trends for complementary feeding practices are mixed. Minimum dietary diversity increased from 37% in the 2022 KDHS to 43% in the 2025–26 KMDHS. Minimum meal frequency declined from 71% in the 2022 survey to 67% in the 2025–26 survey, while minimum acceptable diet remained relatively stable over the same period (31% in 2022 and 32% in 2025–26). The percentage of children consuming eggs and/or flesh foods increased from 28% in 2022 to 35% in 2025–26.

Patterns by background characteristics

- The percentage of children meeting dietary diversity increases from 35% among those age 6–11 months to 50% among those age 12–17 months, before declining to 44% among those age 18–23 months (**Table 11.7**).
- The percentage of children meeting a minimum acceptable diet is 28% among those age 6–11 months, 38% among those age 12–17 months, and 31% among those age 18–23 months.
- Thirty-five percent of female children meet a minimum acceptable diet compared with 30% of male children.
- Thirty-nine percent of children in urban areas meet a minimum acceptable diet compared with 28% of those in rural areas.
- Twelve percent of children whose mothers have no education meet a minimum acceptable diet compared with 47% of those whose mothers have more than a secondary education (**Figure 11.4**).
- Sixteen percent of children in the lowest wealth quintile meet a minimum acceptable diet compared with 44% of those in the fourth wealth quintile and 42% in the highest wealth quintile.
- The percentage of children consuming eggs and/or flesh foods increases from 24% among those age 6–11 months to 40% among those age 12–17 months and further to 42% among those age 18–23 months.

Figure 11.4 **IYCF indicators on Minimum Acceptable Diet (MAD) by mother's education**

Percentage of children age 6–23 months meeting feeding practice recommendations



- Forty-two percent of children in urban areas consume eggs and/or flesh foods compared with 30% of those in rural areas.
- Twenty-two percent of children whose mothers have no education consume eggs and/or flesh foods compared with 44% of those whose mothers have more than a secondary education.
- Twenty-one percent of children in the lowest wealth quintile consume eggs and/or flesh foods compared with 48% of those in the highest wealth quintile.

11.2.6 Sweet Beverage Consumption, Unhealthy Food Consumption, and Zero Vegetable or Fruit Consumption

Unhealthy IYCF practices should be avoided because they can replace nutritious foods. The consumption of sweet foods and beverages increases the risk of dental caries and obesity in childhood. In addition, too much salt in the diet increases the risk of noncommunicable diseases, and unhealthy fats and refined carbohydrates

contribute to unhealthy weight gain. Low consumption of vegetables and fruits leads to reduced nutrient intake, which can negatively impact the growth and development of young children and is associated with noncommunicable diseases later in life. The indicator for unhealthy food consumption defines foods that are high in sugar, salt, and/or unhealthy fats as 'sentinel unhealthy foods' (WHO and UNICEF 2021).

Sweet beverage consumption 6–23 months

Percentage of children age 6–23 months who were given a sweet beverage during the previous day.

Unhealthy food consumption 6–23 months

Percentage of children age 6–23 months who were fed sentinel unhealthy foods during the previous day.

Zero vegetable or fruit consumption 6–23 months

Percentage of children age 6–23 months who were not fed any vegetables or fruits during the previous day.

Sample: Youngest children age 6–23 months living with the mother


Forty-seven percent of the youngest children age 6–23 months living with their mother had a sweet beverage, 18% consumed unhealthy foods, and 23% consumed no vegetables or fruits the previous day (**Tables 11.7** and **11.8**).

Patterns by background characteristics

- Thirty-two percent of children age 6–11 months consume sweet beverages compared with 49% of those age 12–17 months and 61% of those age 18–23 months.
- The percentage of children age 6–23 months who consume no vegetables or fruits is highest among children whose mothers have no education and lowest among those whose mothers have more than a secondary education (55% versus 10%) (**Figure 11.5**).


Figure 11.5 **Unhealthy feeding practices among children age 6–23 months by mother's education**

Percentage of children age 6–23 months with unhealthy feeding practices



- Thirty-nine percent of children in the lowest wealth quintile consume sweet beverages compared with 53% of those in the fourth wealth quintile.
- Eight percent of children in the lowest wealth quintile consume unhealthy foods compared with 25% of those in the fourth wealth quintile.
- Sixteen percent of children in urban areas consume no vegetables and fruits compared with 27% of those in rural areas (**Figure 11.6**).

Figure 11.6 **Unhealthy feeding practices among children age 6–23 months by residence**



11.2.7 Infant and Young Child Feeding Indicators

Table 11.9 summarises all 17 WHO-UNICEF IYCF indicators and provides corresponding values from the 2025–26 KMDHS.

11.3 Infant and Young Child Feeding Counselling

IYCF counselling supports appropriate breastfeeding and complementary feeding practices (WHO 2018; Dewey 2003). Counselling is an interactive process that helps empower mothers and caregivers to follow the recommended IYCF practices. Counselling can take place in health facilities or in the community, and it is delivered by trained health providers, community health workers, and others in the community.

Mothers who received infant and young child feeding counselling in the past 6 months

Percentage of mothers with children age 6–23 months who received IYCF counselling from a health care provider or community health worker in the past 6 months.

Sample: Women age 15–49 whose youngest child age 6–23 months is living with them

Forty-three percent of women age 15–49 whose youngest child age 6–23 months is living with them were counselled by a health care provider in the previous 6 months about how or what to feed their child (**Table 11.10**).

Trends: The percentage of women age 15–49 who were counselled on how or what to feed their youngest child age 6–23 months increased from 25% in the 2022 KDHS to 43% in the 2025–26 KMDHS.

Patterns by background characteristics

- Forty-six percent of mothers age 20–29 received counselling compared with 34% of those age 40–49.
- Forty-eight percent of mothers in urban areas received counselling compared with 39% of those in rural areas.
- Twenty-one percent of mothers with no education received counselling compared with 52% of those with more than a secondary education.

- Thirty-two percent of mothers in the lowest wealth quintile received counselling compared with 51% of those in the highest wealth quintile.

11.4 Micronutrient Supplementation and Deworming

Micronutrient deficiency is a major contributor to childhood morbidity and mortality. Micronutrient deficiency can be caused by a lack of consumption of foods that supply vitamins and minerals, by infections, and by genetic abnormalities. Strategies to prevent or address micronutrient deficiency include agricultural approaches such as biofortification; food-based approaches, which can be complemented with food fortification; and, for specific life stages and population groups, direct micronutrient supplementation (UNICEF 2024; Ashraf 2025).

Iron is a micronutrient that plays an important role in numerous biological systems. Iron deficiency is one of the primary causes of anaemia. Interventions targeting iron deficiency and anaemia include periodically giving children iron tablets or syrups (and/or iron-containing micronutrient powders) (WHO 2011a; WHO 2016a; WHO 2016b).

Vitamin A is a micronutrient that supports the immune system and plays an important role in maintaining the body's epithelial tissues. Severe vitamin A deficiency can damage eyes, increase the severity of infections and illnesses such as measles, and slow recovery from illnesses. Vitamin A supplementation programmes help reduce vitamin A deficiency and mortality in children (WHO 2011b).

Soil-transmitted helminth infections can cause internal bleeding, inflammation, impaired nutrient absorption, diarrhoea, vomiting, and loss of appetite. Deworming programmes help reduce the burden of helminth infections (WHO 2017b).

Given iron tablets or syrups 6–59 months

Percentage of children age 6–59 months who were given iron tablets or syrups in the past 12 months.

Given multiple micronutrient powders 6–59 months

Percentage of children age 6–59 months who were given micronutrient powders in the past 12 months.

Given iron-containing supplements 6–59 months

Percentage of children age 6–59 months who were given iron-containing supplements (tablets, syrups, or micronutrient powders) in the past 12 months.

Sample: Children age 6–59 months

Given vitamin A supplements 6–59 months

Percentage of children age 6–59 months who were given vitamin A supplements in the past 6 months.

Sample: Children age 6–59 months

Given deworming medication 12–59 months

Percentage of children age 12–59 months who were given deworming medication in the past 6 months.

Sample: Children age 12–59 months

Overall, 20% of children age 6–59 months were given iron-containing tablets or syrups in the 12 months preceding the survey, 17% received micronutrient powders, and 24% received any iron-containing supplements. Sixty-eight percent of children age 6–59 months were given vitamin A supplements in the 6 months preceding the survey. In addition, 72% of children age 12–59 months were given deworming medication in the 6 months preceding the survey (**Table 11.11**).

Trends: The percentage of children age 6–59 months who received iron tablets or syrups in the 12 months preceding the survey has remained unchanged at 20% since the 2022 KDHS. However, the percentage who received vitamin A supplements increased from 64% in the 2022 KDHS to 68% in the 2025–26 KMDHS. The percentage of children age 12–59 months who received deworming medication increased from 66% in 2022 to 72% in 2025–26.

Patterns by background characteristics

- Seventy-six percent of children born to mothers age 15–19 received vitamin A supplements compared with 66% of those born to mothers age 40–49.
- Among children age 12–59 months, 66% in urban areas received deworming medication compared with 68% in rural areas.
- Eight counties reported vitamin A supplementation coverage of 80% or higher: Kitui (90%); Taita/Taveta (87%); Kwale (85%); Kisumu (84%); and Kirinyaga, Elgeyo/Marakwet, Homa Bay, and Baringo (81% each). Coverage was less than 50% in Lamu (49%), Wajir (14%), Mandera (17%), Garissa (34%), and Samburu (38%) (**Tables 11.11C**).
- Eleven counties reported deworming coverage of 80% or higher: Nyeri (94%), Vihiga (88%), Kirinyaga (87%), Makueni (87%), Kwale (87%), Kiambu (85%), Taita/Taveta (83%), Kakamega (83%), Embu (82%), Kisumu (82%), and Kericho (80%). Deworming coverage was below 50% in Samburu (41%), Garissa (26%), Wajir (6%), and Mandera (5%).

List of Tables

For more information on nutrition of children, see the following tables:

- **Table 11.1 Child growth monitoring**
- **Table 11.1C Child growth monitoring: County level**
- **Table 11.2 Early breastfeeding**
- **Table 11.3 Breastfeeding status according to age**
- **Table 11.4 Infant feeding practices by age**
- **Table 11.5 Liquids consumed by children in the day or night preceding the interview**
- **Table 11.6 Foods consumed by children in the day or night preceding the interview**
- **Table 11.7 Minimum dietary diversity, minimum meal frequency, and minimum acceptable diet among children**
- **Table 11.8 Egg and/or flesh food consumption and unhealthy feeding practices among children**
- **Table 11.9 Infant and young child feeding indicators**
- **Table 11.10 Infant and young child feeding counselling**
- **Table 11.11 Micronutrient supplementation and deworming among children**
- **Table 11.11C Micronutrient supplementation and deworming among children: County level**

Table 11.1 Child growth monitoring

Percentage of children under age 5 who had selected measurements performed by a health care provider in the 3 months preceding the survey, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Weight	Height	MUAC	Weight and height	Weight, height, and MUAC	Number of children
Age in months						
<6	88.3	80.1	34.2	80.0	34.1	1,047
6–11	89.4	84.4	44.8	83.2	43.6	994
12–23	71.8	67.4	36.5	67.2	35.6	1,934
24–35	45.3	41.6	27.5	41.0	26.4	2,007
36–47	35.9	31.6	21.9	31.4	21.1	2,136
48–59	30.5	26.6	17.4	25.8	16.6	2,048
0–23	80.5	75.0	38.0	74.6	37.2	3,975
24–59	37.2	33.2	22.2	32.7	21.3	6,191
Sex						
Male	54.8	50.1	28.8	49.7	27.9	5,081
Female	53.4	49.0	27.9	48.4	27.2	5,086
Mother's age						
15–19	51.8	45.2	26.3	45.0	24.7	1,205
20–29	54.5	49.9	28.2	49.5	27.6	5,698
30–39	54.5	50.6	29.3	49.8	28.2	2,916
40–49	52.7	49.8	30.5	49.4	29.4	347
Residence						
Urban	57.0	52.3	30.9	51.6	29.8	4,444
Rural	51.9	47.4	26.4	47.1	25.7	5,722
Mother's education[1]						
No education	37.1	33.9	24.8	33.8	24.0	952
Primary	49.5	46.6	28.6	46.1	27.6	3,106
Secondary	54.6	49.5	25.6	48.9	24.7	4,055
More than secondary	68.0	61.4	35.2	61.0	34.6	2,053
Wealth quintile						
Lowest	46.4	43.5	27.6	43.1	26.8	2,185
Second	51.7	47.0	25.9	46.6	25.0	1,875
Middle	53.8	48.3	24.9	47.8	23.8	1,819
Fourth	56.5	51.4	30.1	50.8	29.0	2,271
Highest	62.4	57.5	32.7	56.9	32.4	2,016
Total	54.1	49.5	28.4	49.0	27.5	10,166

Note: 'Height' refers to length (recumbent measurement) or height (standing measurement).
MUAC = mid-upper arm circumference
[1] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 11.1C Child growth monitoring: County level

Percentage of children under age 5 who had selected measurements performed by a health care provider in the 3 months preceding the survey, according to county, Kenya Mini DHS 2025–26

County	Weight	Height	MUAC	Weight and height	Weight, height, and MUAC	Number of children
Mombasa	59.7	58.2	38.8	57.0	37.5	329
Kwale	80.2	80.7	72.2	79.9	71.5	192
Kilifi	67.5	65.8	52.8	65.8	52.3	280
Tana River	37.9	36.4	34.6	36.4	33.6	96
Lamu	32.9	33.3	22.0	32.9	22.0	28
Taita/Taveta	65.6	63.1	33.9	62.4	31.5	80
Garissa	35.8	35.0	33.6	34.5	33.6	119
Wajir	15.1	15.7	15.8	15.0	14.0	96
Mandera	8.1	7.7	5.0	7.3	5.0	118
Marsabit	50.2	45.4	42.1	45.4	41.4	97
Isiolo	50.6	38.4	22.5	38.4	22.0	42
Meru	41.4	28.7	7.4	28.7	6.8	221
Tharaka-Nithi	47.7	43.1	22.0	43.1	21.1	65
Embu	64.3	58.8	20.9	58.8	18.9	75
Kitui	77.9	75.5	51.8	75.5	50.1	175
Machakos	50.4	38.5	13.0	38.5	13.0	339
Makueni	80.2	75.7	29.5	75.7	29.5	152
Nyandarua	57.7	55.7	9.2	52.9	8.1	102
Nyeri	65.0	60.0	15.1	60.0	15.1	84
Kirinyaga	65.4	56.1	15.8	56.1	15.8	91
Murang'a	61.0	46.1	13.4	46.1	13.4	138
Kiambu	52.6	49.9	31.3	48.9	31.3	714
Turkana	56.5	54.4	42.1	54.1	39.7	162
West Pokot	31.4	31.2	29.2	30.5	28.4	181
Samburu	42.6	35.8	22.5	35.5	21.5	75
Trans Nzoia	65.0	62.1	26.8	62.1	26.6	150
Uasin Gishu	59.1	55.8	33.0	55.8	32.0	304
Elgeyo/Marakwet	63.1	61.5	40.4	61.5	40.0	86
Nandi	71.1	62.7	49.9	62.7	49.9	161
Baringo	62.9	59.1	31.3	59.1	30.2	118
Laikipia	42.7	40.5	11.0	39.8	10.3	100
Nakuru	53.6	43.5	20.7	43.5	19.7	542
Narok	57.5	55.1	37.1	54.9	36.5	236
Kajiado	55.5	53.0	4.6	52.3	4.6	268
Kericho	47.0	39.9	20.3	39.9	20.1	159
Bomet	41.1	34.2	7.2	34.2	6.8	227
Kakamega	42.4	37.5	5.7	36.2	5.3	318
Vihiga	50.0	49.8	23.8	48.8	21.5	95
Bungoma	51.7	47.3	30.3	47.3	30.3	305
Busia	60.4	57.7	36.6	57.7	36.0	168
Siaya	33.2	32.7	10.6	32.7	10.6	161
Kisumu	62.1	61.0	32.9	59.8	29.5	256
Homa Bay	69.4	66.7	51.4	66.5	51.4	237
Migori	46.8	43.5	16.2	43.2	15.9	320
Kisii	34.6	33.2	17.8	32.2	17.8	208
Nyamira	52.4	31.5	18.8	29.8	16.9	96
Nairobi City	57.4	52.1	35.1	51.2	33.1	1,600
Total	54.1	49.5	28.4	49.0	27.5	10,166

Note: 'Height' refers to length (recumbent measurement) or height (standing measurement).
MUAC = mid-upper arm circumference

Table 11.2 Early breastfeeding

Percentage of children born in the past 2 years who were ever breastfed, percentage who were put to the breast within 1 hour of birth, and percentage who were exclusively breastfed for the first 2 days after birth, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Percentage ever breastfed	Percentage put to the breast within 1 hour of birth	Percentage exclusively breastfed for the first 2 days after birth[1]	Number of children born in the past 2 years
Sex				
Male	96.9	62.2	83.2	2,078
Female	97.0	63.3	83.2	2,037
Breastfeeding counselling during ANC[2]				
Counselled	97.3	62.7	86.2	3,408
Not counselled/don't know	96.5	63.4	84.2	495
Did not receive ANC	92.4	61.7	32.1	213
Assistance at delivery				
Health personnel[3]	97.1	62.4	83.3	3,781
Traditional birth attendant	95.5	65.8	79.9	205
Other	96.5	67.1	85.4	91
No one	(95.2)	(68.8)	(87.0)	38
Place of delivery				
Health facility	97.1	62.3	83.2	3,749
At home	96.0	67.3	82.9	353
Other	*	*	*	12
Type of delivery				
Vaginal birth	96.7	71.1	86.0	3,281
Caesarean section	98.2	29.9	72.0	834
Breastfeeding counselling during PNC[2,4]				
Counselled	97.8	61.8	86.5	3,368
Not counselled/don't know	94.2	70.3	83.2	615
Breastfeeding observation during PNC[2,4]				
Observed	97.9	61.9	86.9	3,349
Not observed/don't know	93.7	69.6	80.7	634
Residence				
Urban	97.3	56.7	81.7	1,755
Rural	96.7	67.3	84.3	2,360
Mother's education[5]				
No education	97.6	66.9	85.1	363
Primary	96.9	67.9	86.9	1,111
Secondary	96.6	62.0	81.8	1,817
More than secondary	97.7	55.6	80.5	825
Wealth quintile				
Lowest	97.3	70.0	86.0	873
Second	97.0	68.4	84.3	770
Middle	95.7	64.3	82.9	739
Fourth	97.7	61.2	83.5	945
Highest	96.8	49.6	78.9	788
Total	97.0	62.8	83.2	4,115

Note: Table is based on children born in the 2 years preceding the survey regardless of whether the children were living or dead at the time of interview.
ANC = antenatal care
PNC = postnatal care
[1] Children given nothing other than breast milk to eat or drink during the first 2 days after delivery
[2] Information available for the most recent live birth only
[3] Doctor, nurse/midwife, or auxiliary midwife
[4] Women were asked about counselling on breastfeeding from any health care provider in the first 2 days after their most recent live birth regardless of where they gave birth.
[5] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 11.3 Breastfeeding status according to age

Among youngest children age 0–5 months living with their mother, percentage exclusively breastfeeding and percentage receiving mixed milk feeding; among all children age 12–23 months, percentage currently breastfeeding; and among all children age 0–23 months, percentage using a bottle with a nipple, according to background characteristics, Kenya Mini DHS 2025–26

	Among youngest children age 0–5 months living with their mother:			Among all children age 12–23 months:		Among all children age 0–23 months:	
Background characteristic	Percentage exclusively breastfeeding	Percentage receiving mixed milk feeding[1]	Number of children	Percentage currently breastfeeding[2]	Number of children	Percentage using a bottle with a nipple	Number of children
Age in months							
0–1	90.1	2.9	327	na	na	3.3	334
2–3	75.3	8.5	334	na	na	16.8	348
4–5	50.7	14.8	355	na	na	27.8	364
6–11	na	na	na	na	na	49.6	994
12–15	na	na	na	84.4	645	48.0	645
16–19	na	na	na	64.2	690	40.1	690
20–23	na	na	na	41.2	599	35.4	599
Sex							
Male	69.1	9.6	486	61.5	1,006	36.6	1,993
Female	73.7	8.2	530	66.4	929	37.0	1,983
Residence							
Urban	75.8	7.1	458	61.0	784	43.4	1,686
Rural	68.0	10.4	558	65.8	1,150	31.9	2,289
Mother's education[3]							
No education	66.4	13.1	92	69.9	173	29.9	354
Primary	70.6	6.7	272	66.4	518	32.7	1,070
Secondary	71.9	9.0	438	60.8	858	36.5	1,739
More than secondary	74.0	9.7	214	64.5	385	45.7	813
Wealth quintile							
Lowest	60.1	14.0	203	67.2	413	31.3	850
Second	70.5	6.9	191	67.4	359	29.9	739
Middle	74.1	8.4	169	62.5	375	33.7	719
Fourth	74.3	6.6	257	62.9	432	37.9	892
Highest	78.3	9.0	196	58.8	356	51.0	776
Total	71.5	8.9	1,016	63.9	1,934	36.8	3,975

Note: Breastfeeding status refers to a '24-hour' period (yesterday during the day or night).
na = not applicable
[1] Received breast milk plus either animal milk (fresh, packaged, or powdered) or infant formula; excludes yogurt drinks because they are generally not fed as a substitute for breast milk; excludes soy and nut milks
[2] Corresponds to infant and young child feeding indicator 'Continued breastfeeding'
[3] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 11.4 Infant feeding practices by age

Percent distribution of youngest children age 0–5 months living with their mother by feeding category, according to age, Kenya Mini DHS 2025–26

Age in months	Breastmilk only (exclusively breastfed)	Breastmilk and plain water only	Breastmilk and non-milk liquids[1]	Breastmilk and animal milk and/or infant formula[2]	Breastmilk and solid, semi-solid, or soft foods[3]	Not breastfed	Unknown[4]	Total	Number of youngest children 0–5 months living with their mother
0–1	90.1	3.1	1.3	2.7	1.6	0.9	0.3	100.0	327
2–3	75.3	7.0	1.4	6.2	7.2	2.1	0.6	100.0	334
4–5	50.7	10.7	0.8	7.1	28.8	0.9	0.9	100.0	355
0–5	71.5	7.0	1.2	5.4	12.9	1.3	0.6	100.0	1,016

Notes: Breastfeeding status refers to a '24-hour' period (yesterday during the day or night). The categories of breastmilk only; breast milk and plain water only; breast milk and non-milk liquids; breastmilk and formula and/or animal milk; breastmilk and solid, semi-solid, or soft foods; and not breastfed are hierarchical and mutually exclusive. Thus, children who receive breastmilk and non-milk liquids; who do not receive breastmilk and formula and/or animal milk; and who do not receive any solid, semi-solid, or soft foods are classified in the non-milk liquids category even though they may also receive plain water. When combined with the percentages of children whose feeding category is classified as unknown due to 'don't know' responses, the percentages in each row add to 100%.

[1] Children fed breast milk plus non-milk liquids (e.g., juice, herbal tea, sweetened water, flavoured water). Children in this category may have also been fed plain water.

[2] Includes children fed breast milk plus animal milk, breast milk plus infant formula, or breast milk plus animal milk-based yogurt drinks. Children in this category may have also been fed non-milk liquids and/or plain water.

[3] Children fed breast milk plus solid, semi-solid, or soft foods from any food group (e.g., grains, meat, eggs, fruits, vegetables). Children in this category may have also been fed plain water; non-milk liquids; and/or animal milk, infant formula, or animal milk-based yogurt drinks.

[4] Not classified elsewhere due to 'don't know' responses

Table 11.5 Liquids consumed by children in the day or night preceding the interview

Percentage of youngest children under age 2 living with their mother by type of liquids consumed in the day or night preceding the interview, according to age and breastfeeding status, Kenya Mini DHS 2025–26

Age in months	Plain water	Infant formula[1]	Animal milk		Yogurt drinks		Fruit juice and fruit-flavoured drinks	Sodas, malt drinks, sports drinks, and energy drinks	Tea, coffee, and herbal drinks		Clear broth and clear soup	Other liquids		Number of youngest children under age 2 living with their mother
			Any	Sweet/ flavoured	Any	Sweet/ flavoured			Any	Sweetened		Any	Sweetened	
BREASTFEEDING CHILDREN														
0–1	4.8	1.8	1.1	0.2	0.0	0.0	0.6	0.2	0.6	0.0	0.0	0.2	0.0	324
2–3	12.6	3.7	5.0	0.8	0.0	0.0	1.0	0.0	1.2	0.4	0.5	0.6	0.4	327
4–5	33.0	0.6	14.3	2.4	0.0	0.0	1.6	0.1	1.8	1.1	3.9	1.5	0.0	351
6–8	85.7	0.6	47.8	9.1	0.0	0.0	10.7	0.4	12.4	9.1	18.8	3.6	0.8	529
9–11	89.8	3.8	54.0	11.1	0.0	0.0	13.3	1.1	20.7	14.6	24.5	6.3	0.7	432
12–17	92.1	0.4	59.7	12.1	0.0	0.0	12.4	3.1	41.4	30.0	20.9	8.1	0.6	768
18–23	94.3	2.2	54.2	14.7	0.0	0.0	13.5	8.2	54.8	41.4	22.7	9.6	0.5	457
0–5	17.2	2.0	7.0	1.2	0.0	0.0	1.1	0.1	1.2	0.5	1.5	0.8	0.1	1,003
6–11	87.6	2.0	50.6	10.0	0.0	0.0	11.9	0.7	16.1	11.6	21.4	4.8	0.7	961
12–23	92.9	1.1	57.6	13.1	0.0	0.0	12.8	5.0	46.4	34.3	21.6	8.6	0.6	1,224
6–23	90.6	1.5	54.5	11.7	0.0	0.0	12.4	3.1	33.1	24.3	21.5	6.9	0.6	2,186
Total	67.5	1.7	39.6	8.4	0.0	0.0	8.8	2.2	23.1	16.8	15.2	5.0	0.5	3,189
NON-BREASTFEEDING CHILDREN														
0–1	*	*	*	*	*	*	*	*	*	*	*	*	*	3
2–3	*	*	*	*	*	*	*	*	*	*	*	*	*	7
4–5	*	*	*	*	*	*	*	*	*	*	*	*	*	3
6–8	*	*	*	*	*	*	*	*	*	*	*	*	*	7
9–11	*	*	*	*	*	*	*	*	*	*	*	*	*	11
12–17	96.2	1.8	73.4	14.4	0.0	0.0	11.6	4.8	43.5	28.8	16.4	7.0	0.0	183
18–23	92.7	0.9	63.0	17.3	0.0	0.0	10.5	5.8	57.6	47.8	16.7	10.1	0.7	415
0–5	*	*	*	*	*	*	*	*	*	*	*	*	*	13
6–11	(89.5)	(4.3)	(65.8)	(38.7)	(0.0)	(0.0)	(5.8)	(10.2)	(17.3)	(11.2)	(26.7)	(8.8)	(6.1)	18
12–23	93.8	1.2	66.2	16.4	0.0	0.0	10.9	5.5	53.3	42.0	16.6	9.1	0.5	598
6–23	93.7	1.3	66.1	17.1	0.0	0.0	10.7	5.6	52.2	41.1	16.9	9.1	0.6	616
Total	92.6	1.7	65.0	16.8	0.0	0.0	10.9	5.5	51.1	40.2	16.5	9.1	0.6	629

Notes: Figures in parentheses are based on 25–49 unweighted cases. An asterisk indicates that a figure is based on fewer than 25 unweighted cases and has been suppressed.

[1] Infant formula includes NAN (Nestle), SMA, or S26.

Table 11.6 Foods consumed by children in the day or night preceding the interview

Percentage of youngest children under age 2 living with their mother by type of foods consumed in the day or night preceding the interview, according to age and breastfeeding status, Kenya Mini DHS 2025–26

	Solid or semi-solid foods												Number of youngest children under age 2 living with their mother
Age in months	Grains	White/ pale starchy roots, tubers, and plan-tains	Pulses (beans, peas, lentils, nuts, seeds)	Dairy products (milk, infant formula, yogurt, cheese)	Flesh foods (meat, fish, poultry, organ meat)	Eggs	Vitamin A-rich fruits and vege-tables	Other fruits and vege-tables	Insects and other small protein foods	Sweet foods[1]	Fried and salty foods[2]	Other solid, semi-solid, and soft foods	
BREASTFEEDING CHILDREN													
0–1	1.2	0.0	0.1	0.0	0.0	0.1	0.0	0.0	0.0	0.4	0.0	0.0	324
2–3	7.1	1.6	0.3	0.3	0.2	0.3	1.2	1.1	0.0	0.0	0.0	0.0	327
4–5	19.8	5.6	0.6	3.2	0.2	0.3	8.1	11.2	0.0	0.3	0.0	0.0	351
6–8	75.7	40.7	11.5	10.1	14.2	7.3	50.2	43.7	0.8	4.4	0.0	0.0	529
9–11	85.9	37.9	23.1	17.1	21.2	9.3	61.6	51.5	1.6	11.1	0.0	0.0	432
12–17	89.0	42.3	25.1	15.0	30.7	16.2	69.1	63.4	1.0	19.4	0.0	0.0	768
18–23	93.4	43.9	32.4	17.0	33.2	16.7	70.8	65.1	0.4	22.7	0.0	0.0	457
0–5	9.6	2.5	0.3	1.2	0.1	0.2	3.3	4.3	0.0	0.2	0.0	0.0	1,003
6–11	80.3	39.4	16.7	13.2	17.4	8.2	55.3	47.2	1.2	7.4	0.0	0.0	961
12–23	90.7	42.9	27.8	15.8	31.6	16.4	69.8	64.0	0.8	20.6	0.0	0.0	1,224
6–23	86.1	41.4	22.9	14.7	25.4	12.8	63.4	56.6	1.0	14.8	0.0	0.0	2,186
Total	62.1	29.2	15.8	10.4	17.4	8.8	44.5	40.2	0.7	10.2	0.0	0.0	3,189
NON-BREASTFEEDING CHILDREN													
0–1	*	*	*	*	*	*	*	*	*	*	*	*	3
2–3	*	*	*	*	*	*	*	*	*	*	*	*	7
4–5	*	*	*	*	*	*	*	*	*	*	*	*	3
6–8	*	*	*	*	*	*	*	*	*	*	*	*	7
9–11	*	*	*	*	*	*	*	*	*	*	*	*	11
12–17	92.3	48.6	30.6	21.8	25.8	11.4	66.5	59.7	1.3	22.5	0.0	0.0	183
18–23	90.1	40.4	32.1	15.7	34.9	7.3	74.1	68.1	0.6	29.7	0.0	0.0	415
0–5	*	*	*	*	*	*	*	*	*	*	*	*	13
6–11	(92.4)	(46.2)	(17.2)	(19.6)	(28.6)	(10.6)	(57.3)	(65.0)	(0.0)	(5.4)	(0.0)	(0.0)	18
12–23	90.8	42.9	31.7	17.6	32.1	8.6	71.8	65.5	0.8	27.5	0.0	0.0	598
6–23	90.8	43.0	31.2	17.6	32.0	8.6	71.4	65.5	0.8	26.9	0.0	0.0	616
Total	89.6	42.2	30.6	17.3	31.4	8.5	70.4	64.5	0.8	26.3	0.0	0.0	629

Notes: See Woman's Questionnaire for list of liquids and foods. Figures in parentheses are based on 25–49 unweighted cases. An asterisk indicates that a figure is based on fewer than 25 unweighted cases and has been suppressed.

[1] Sentinel sweet foods such as chocolates, candies, pastries, cakes, biscuits, ice cream, and popsicles

[2] Sentinel fried and salty foods such as chips, crisps, puffs, french fries, fried dough, and instant noodles

Table 11.7 Minimum dietary diversity, minimum meal frequency, and minimum acceptable diet among children

Percentage of youngest children age 6–23 months living with their mother who are fed a minimum acceptable diet based on breastfeeding status, number of food groups, and times they were fed during the day or night preceding the survey, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Among youngest breastfed children age 6–23 months living with their mother, percentage who meet:			Number of breast-fed children age 6–23 months	Among youngest non-breastfed children age 6–23 months living with their mother, percentage who meet:				Number of non-breast-fed children age 6–23 months	Among youngest children age 6–23 months living with their mother, percentage who meet:			Number of all children age 6–23 months
	Minimum dietary diversity[1]	Minimum meal frequency[2]	Minimum acceptable diet[3]		Minimum milk feeding frequency[4]	Minimum dietary diversity[1]	Minimum meal frequency[5]	Minimum acceptable diet[6]		Minimum dietary diversity[1]	Minimum meal frequency[7]	Minimum acceptable diet[8]	
Age in months													
6–11	34.9	68.8	28.3	961	(53.4)	(22.4)	(63.7)	(12.5)	18	34.7	68.7	28.1	979
6–8	31.8	77.7	29.3	529	*	*	*	*	7	31.5	77.4	29.0	536
9–11	38.6	58.0	27.2	432	*	*	*	*	11	38.4	58.3	26.9	443
12–17	54.9	64.9	40.8	768	66.8	30.2	71.7	24.7	183	50.2	66.2	37.7	951
18–23	57.5	64.2	42.7	457	48.0	28.2	70.0	19.1	415	43.5	66.9	31.5	871
Sex													
Male	44.3	64.5	33.3	1,081	52.5	33.2	66.1	21.2	343	41.6	64.9	30.4	1,424
Female	48.9	68.4	38.1	1,105	55.3	22.9	75.7	19.7	272	43.8	69.8	34.5	1,377
Residence													
Urban	52.5	70.7	42.8	892	54.6	36.5	72.2	28.1	264	48.8	71.1	39.4	1,156
Rural	42.6	63.5	30.9	1,294	53.0	22.7	68.9	14.9	352	38.4	64.7	27.5	1,646
Mother's education[9]													
No education	21.5	36.2	13.1	205	68.0	8.0	65.5	4.2	45	19.0	41.5	11.5	250
Primary	34.5	64.5	27.1	608	43.6	19.8	53.4	13.6	151	31.6	62.3	24.4	760
Secondary	52.3	70.7	39.7	931	51.3	30.2	74.7	19.6	296	47.0	71.7	34.8	1,227
More than secondary	63.1	74.2	49.8	441	66.7	43.2	82.3	37.2	123	58.8	76.0	47.1	565
Wealth quintile													
Lowest	26.8	51.3	17.8	498	56.6	13.4	58.4	8.6	114	24.3	52.6	16.1	611
Second	42.6	67.5	31.9	415	39.9	22.6	64.7	11.1	105	38.5	66.9	27.7	520
Middle	51.7	69.3	39.0	397	50.5	25.6	76.3	15.8	128	45.4	71.0	33.3	525
Fourth	60.8	73.5	48.7	454	56.1	35.9	77.1	26.7	139	55.0	74.3	43.5	593
Highest	54.0	73.1	43.6	421	63.0	42.0	72.2	36.7	131	51.2	72.9	42.0	552
Total	46.6	66.5	35.7	2,186	53.7	28.6	70.3	20.6	616	42.7	67.3	32.4	2,802

Notes: Figures in parentheses are based on 25–49 unweighted cases. An asterisk indicates that a figure is based on fewer than 25 unweighted cases and has been suppressed.

[1] Minimum dietary diversity is receiving foods from five or more of the following eight food groups: breast milk; grains, white/pale starchy roots, tubers, and plantains; beans, peas, lentils, nuts, and seeds; dairy products (tinned, powdered, or fresh animal milk; infant formula; yogurt; and cheese); flesh foods (meat, fish, poultry, and organ meat); eggs; vitamin A-rich fruits and vegetables; and other fruits and vegetables.

[2] For breastfed children, minimum meal frequency is receiving solid, semi-solid, or soft foods at least twice a day for infants age 6–8 months and at least three times a day for children age 9–23 months.

[3] For breastfed children, minimum acceptable diet is receiving minimum dietary diversity (footnote 1) and minimum meal frequency (footnote 2).

[4] For non-breastfed children, minimum milk feeding frequency is two or more feedings of infant formula; tinned, powdered, or fresh animal milk; or a yogurt drink or solid.

[5] For non-breastfed children, minimum meal frequency is receiving solid, semi-solid, or soft foods or milk feeds at least four times a day. At least one of the feeds must be a solid, semi-solid, or soft food.

[6] For non-breastfed children, minimum acceptable diet is receiving minimum dietary diversity (footnote 1), minimum milk feeding frequency (footnote 4), and minimum meal frequency (footnote 5).

[7] Minimum meal frequency is receiving the minimum recommended number of feeds per day according to age and breastfeeding status as defined in footnotes 2 and 5.

[8] Minimum acceptable diet is receiving minimum dietary diversity (footnote 1), minimum meal frequency (footnote 2 for breastfed children and footnote 5 for non-breastfed children), and minimum milk feeding frequency (footnote 4 for non-breastfed children).

[9] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 11.8 Egg and/or flesh food consumption and unhealthy feeding practices among children

Percentage of youngest children age 6–23 months living with their mother who consumed eggs and/or flesh foods, and percentage who experienced specific unhealthy feeding practices, during the day or night preceding the survey, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Eggs and/or flesh foods (meat, fish, poultry, organ meat)	Unhealthy feeding practices:			Number of youngest children age 6–23 months living with their mother
		Sweet beverage[1]	Unhealthy food[2]	No vegetables or fruits[3]	
Age in months					
6–11	23.7	31.8	7.4	32.4	979
6–8	20.2	27.3	4.4	38.3	536
9–11	28.1	37.3	10.9	25.2	443
12–17	39.7	48.5	20.0	18.7	951
18–23	41.9	61.4	26.1	15.6	871
Sex					
Male	32.5	46.2	17.4	21.3	1,424
Female	37.3	47.2	17.6	23.7	1,377
Breastfeeding status					
Breastfeeding	33.9	43.0	14.8	24.5	2,186
Not breastfeeding	38.0	59.5	26.9	15.5	616
Residence					
Urban	41.7	47.3	20.0	15.8	1,156
Rural	30.0	46.2	15.7	27.2	1,646
Mother's education[4]					
No education	21.7	41.0	7.9	55.1	250
Primary	28.8	44.8	14.4	29.1	760
Secondary	37.1	46.2	21.2	17.4	1,227
More than secondary	43.9	52.7	17.8	10.3	565
Wealth quintile					
Lowest	20.5	39.2	8.4	45.0	611
Second	28.9	47.4	15.6	21.9	520
Middle	36.3	49.3	19.8	14.6	525
Fourth	41.6	53.4	25.0	16.3	593
Highest	47.6	44.5	18.9	12.4	552
Total	34.8	46.7	17.5	22.5	2,802

[1] Sweet beverages include sweet/flavoured milk and yogurt drinks; sweet/flavoured soy milks and nut milks; fruit juice and fruit-flavoured drinks; chocolate-flavoured drinks; sodas, malt drinks, sports drinks, and energy drinks; sweetened tea, coffee, and herbal drinks; and other sweetened liquids.
[2] Sentinel unhealthy foods including sweet foods such as chocolates, candies, pastries, cakes, biscuits, ice cream, and ice lollies; and fried and salty foods such as crisps, chips, ngumu, mandaazi, samosas, bhajias, and indomie
[3] No vitamin A-rich fruits or vegetables and no other fruits or vegetables
[4] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 11.9 Infant and young child feeding indicators

Percentage of children fed according to various IYCF practices, Kenya Mini DHS 2025–26

IYCF		DHS-8	Percentage of children fed according to various IYCF practices		
#	Abbrev.	Table #	Indicator	Indicator definition and denominator	Value
1	EvBF	11.3	Ever breastfed[1]	Percentage of children born in the past 2 years who were ever breastfed	97.0
				Number of children born in the past 2 years	4,115
2	EIBF	11.3	Early initiation of breastfeeding[1]	Percentage of children born in the past 2 years who were put to the breast within 1 hour of birth	62.8
				Number of children born in the past 2 years	4,115
3	EBF2D	11.3	Exclusive breastfeeding for the first 2 days after birth[1]	Percentage of children born in the past 2 years who were fed exclusively with breast milk for the first 2 days after birth	83.2
				Number of children born in the past 2 years	4,115
4	EBF	11.4	Exclusive breastfeeding under 6 months	Percentage of children age 0–5 months who were fed exclusively with breast milk during the previous day	71.5
				Number of youngest children age 0–5 months living with the mother	1,016
5	MixMF	11.4	Mixed milk feeding under 6 months	Percentage of children age 0–5 months who were fed both breast milk and formula and/or animal milk during the previous day	8.9
				Number of youngest children age 0–5 months living with the mother	1,016
6	CBF	11.4	Continued breastfeeding 12–23 months	Percentage of children age 12–23 months who were fed breast milk during the previous day	63.9
				Number of children age 12–23 months	1,934
7	ISSSF	--	Introduction of solid, semi-solid, or soft foods 6–8 months	Percentage of children age 6–8 months who were fed solid, semi-solid, or soft foods during the previous day	87.8
				Number of youngest children age 6–8 months living with the mother	536
8	MDD	11.8	Minimum dietary diversity 6–23 months	Percentage of children age 6–23 months who were fed foods and beverages from at least five out of eight defined food groups during the previous day	42.7
				Number of youngest children age 6–23 months living with the mother	2,802
9	MMF	11.8	Minimum meal frequency 6–23 months	Percentage of children age 6–23 months who were fed solid, semi-solid, or soft foods (including milk feeds for non-breastfed children) the minimum recommended number of times or more during the previous day	67.3
				Number of youngest children age 6–23 months living with the mother	2,802
10	MMFF	11.8	Minimum milk feeding frequency for non-breastfed children 6–23 months	Percentage of non-breastfed children age 6–23 months who were given at least two milk feeds during the previous day	53.7
				Number of youngest children age 6–23 months living with the mother who were not breastfed	616
11	MAD	11.8	Minimum acceptable diet 6–23 months	Percentage of children age 6–23 months who were fed a minimum acceptable diet during the previous day	32.4
				Number of youngest children age 6–23 months living with the mother	2,802
12	EFF	11.9	Egg and/or flesh food consumption 6–23 months	Percentage of children age 6–23 months who were fed eggs and/or flesh foods during the previous day	34.8
				Number of youngest children age 6–23 months living with the mother	2,802
13	SWB	11.9	Sweet beverage consumption 6–23 months	Percentage of children age 6–23 months who were given a sweet beverage during the previous day	46.7
				Number of youngest children age 6–23 months living with the mother	2,802
14	UFC	11.9	Unhealthy food consumption 6–23 months	Percentage of children age 6–23 months who were fed selected sentinel unhealthy foods during the previous day	17.5
				Number of youngest children age 6–23 months living with the mother	2,802
15	ZVF	11.9	Zero vegetable or fruit consumption 6–23 months	Percentage of children age 6–23 months who were not fed any vegetables or fruits during the previous day	22.5
				Number of youngest children age 6–23 months living with the mother	2,802
16	BoF	11.4	Bottle feeding 0–23 months	Percentage of children age 0–23 months who were fed from a bottle with a nipple during the previous day	36.8
				Number of children age 0–23 months	3,975
17	--	11.5	Infant feeding area graph	Percent distribution of youngest children age 0–5 months living with the mother, by feeding category	
				Number of youngest children age 0–5 months living with the mother	

[1] Includes children born in the 2 years preceding the survey regardless of whether the children were living or dead at the time of interview.
IYCF = infant and young child feeding
DHS = Demographic and Health Survey

Table 11.10 Infant and young child feeding counselling

Among women age 15–49 whose youngest child age 6–23 months is living with them, percentage who talked with a health care provider or community health worker about how or what to feed their child in the past 6 months, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Counselled in past 6 months about how or what to feed their child	Number of women whose youngest child age 6–23 months is living with them
Child's age in months		
6–11	43.3	979
12–23	42.4	1,822
Child's sex		
Male	41.7	1,424
Female	43.7	1,377
Age		
15–19	41.6	199
20–29	45.5	1,585
30–39	39.1	891
40–49	34.2	126
Residence		
Urban	48.0	1,156
Rural	39.0	1,646
Education[1]		
No education	20.5	250
Primary	36.9	760
Secondary	46.5	1,227
More than secondary	52.0	565
Wealth quintile		
Lowest	31.5	611
Second	38.8	520
Middle	45.8	525
Fourth	47.5	593
Highest	50.6	552
Total	42.7	2,802

[1] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 11.11 Micronutrient supplementation and deworming among children

Among children age 6–59 months, percentages who were given iron tablets or syrups, micronutrient powders, and iron-containing supplements in the past 12 months and percentage who were given vitamin A supplements in the past 6 months; and among children age 12–59 months, percentage who were given deworming medication in the past 6 months, according to background characteristics, Kenya Mini DHS 2025–26

	Among children age 6–59 months:					Among children age 12–59 months:	
Background characteristic	Percentage given iron tablets or syrups in past 12 months[1]	Percentage given micronutrient powders in past 12 months[1,2]	Percentage given iron-containing supplements in past 12 months[1,3]	Percentage given vitamin A supplements in past 6 months[4]	Number of children	Percentage given deworming medication in past 6 months[1,5]	Number of children
Age in months							
6–8	14.2	9.6	16.3	64.1	541	na	na
9–11	17.6	13.5	19.8	80.1	454	na	na
12–17	18.7	15.2	21.0	79.8	985	59.0	985
18–23	18.2	15.2	21.5	82.3	950	72.9	950
24–35	21.3	18.0	25.6	70.5	2,007	73.9	2,007
36–47	24.4	19.2	27.6	64.6	2,136	72.5	2,136
48–59	19.7	16.0	23.2	57.0	2,048	73.7	2,048
6–23	17.6	13.9	20.1	77.7	2,929	65.8	1,934
24–59	21.8	17.8	25.5	64.0	6,191	73.3	6,191
Sex							
Male	20.6	17.0	24.2	68.1	4,581	71.6	4,094
Female	20.3	16.0	23.4	68.7	4,539	71.5	4,032
Breastfeeding status[6]							
Breastfeeding	16.6	12.9	19.0	77.0	2,434	64.9	1,461
Not breastfeeding	21.4	18.2	25.7	72.6	2,502	72.9	2,480
Mother's age							
15–19	20.0	16.4	24.8	75.5	282	68.5	195
20–29	20.8	16.5	23.8	68.7	4,667	70.8	4,083
30–39	19.6	16.5	23.3	67.9	3,483	72.6	3,191
40–49	22.6	17.4	25.8	65.7	687	71.6	657
Residence							
Urban	23.5	18.9	27.3	68.5	3,965	75.7	3,542
Rural	18.1	14.7	21.0	68.4	5,155	68.3	4,583
Mother's education[7]							
No education	11.8	10.7	13.7	43.2	858	38.7	771
Primary	22.5	18.0	25.2	66.9	2,829	69.9	2,554
Secondary	20.5	15.6	23.8	72.9	3,604	74.7	3,175
More than secondary	21.2	18.8	26.3	73.8	1,828	83.4	1,625
Wealth quintile							
Lowest	16.9	13.9	19.0	60.0	1,977	57.3	1,747
Second	20.6	16.6	23.5	70.6	1,681	69.7	1,496
Middle	19.7	16.2	22.9	73.5	1,647	74.1	1,475
Fourth	20.8	16.0	24.6	74.9	2,005	79.2	1,811
Highest	24.4	20.2	29.2	63.7	1,809	77.8	1,596
Total	20.4	16.5	23.8	68.4	9,119	71.5	8,125

na = not applicable
[1] Based on mother's recall
[2] Locally available and distributed micronutrient powders
[3] Iron-containing supplements include tablets, syrups, and micronutrient powders.
[4] Based on both mother's recall and the vaccination card (where available)
[5] Deworming for intestinal parasites is commonly done for helminths and schistosomiasis.
[6] Information available for children age 0–35 months only
[7] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 11.11C Micronutrient supplementation and deworming among children: County level

Among children age 6–59 months, percentages who were given iron tablets or syrups, micronutrient powders, and iron-containing supplements in the past 12 months and percentage who were given vitamin A supplements in the past 6 months; and among children age 12–59 months, percentage who were given deworming medication in the past 6 months, according to county, Kenya Mini DHS 2025–26

	Among children age 6–59 months:					Among children age 12–59 months:	
County	Percentage given iron tablets or syrups in past 12 months[1]	Percentage given micronutrient powders in past 12 months[1,2]	Percentage given iron-containing supplements in past 12 months[1,3]	Percentage given vitamin A supplements in past 6 months[4]	Number of children	Percentage given deworming medication in past 6 months[1,5]	Number of children
Mombasa	34.7	30.7	35.3	67.0	304	75.0	263
Kwale	29.5	17.2	32.0	85.3	177	87.4	157
Kilifi	41.4	34.7	42.8	68.2	254	69.6	231
Tana River	2.3	1.7	3.0	62.6	89	72.0	77
Lamu	11.4	8.2	13.0	49.4	25	71.5	22
Taita/Taveta	17.0	11.7	19.5	86.7	74	83.4	66
Garissa	10.3	8.6	11.6	34.2	106	26.2	97
Wajir	0.2	1.1	1.1	13.9	88	6.0	76
Mandera	7.4	5.3	7.4	17.2	107	4.7	97
Marsabit	14.0	19.9	20.5	75.1	90	67.1	79
Isiolo	7.1	0.3	7.1	66.8	39	65.8	34
Meru	19.1	4.9	20.4	60.1	202	65.5	184
Tharaka-Nithi	9.4	4.2	11.6	49.6	56	50.2	50
Embu	25.0	12.2	27.9	68.2	68	81.5	62
Kitui	22.2	19.0	23.4	90.3	157	77.5	149
Machakos	9.1	7.7	10.3	76.9	305	72.1	277
Makueni	18.4	12.4	20.6	76.3	128	86.9	114
Nyandarua	11.4	10.2	17.2	66.1	91	77.6	79
Nyeri	13.0	15.4	20.8	79.0	74	93.5	63
Kirinyaga	21.2	9.3	23.7	80.0	84	86.5	75
Murang'a	13.3	4.2	13.9	76.7	126	79.4	111
Kiambu	24.8	25.1	28.8	68.2	618	84.8	556
Turkana	37.1	33.5	40.5	68.6	147	63.2	134
West Pokot	22.6	21.8	24.8	70.3	165	56.4	140
Samburu	10.2	12.7	16.0	37.5	68	41.3	57
Trans Nzoia	23.2	13.7	24.6	75.4	130	71.5	114
Uasin Gishu	16.6	21.1	24.7	74.7	282	67.7	243
Elgeyo/Marakwet	24.0	25.3	29.1	80.4	79	77.9	70
Nandi	22.4	14.3	27.0	78.0	141	75.1	127
Baringo	5.0	3.3	7.0	80.8	104	76.9	90
Laikipia	13.2	15.7	18.6	60.4	90	65.7	78
Nakuru	12.9	8.3	17.2	63.8	488	71.9	440
Narok	17.5	12.9	18.5	62.9	218	58.7	192
Kajiado	7.9	4.9	9.3	73.7	227	69.4	198
Kericho	10.0	7.3	13.1	53.5	139	80.0	128
Bomet	5.3	4.4	5.7	64.0	209	61.8	190
Kakamega	19.3	12.4	25.8	71.1	287	83.1	255
Vihiga	27.8	25.3	32.8	75.7	83	87.5	72
Bungoma	20.8	20.0	24.7	76.9	274	75.9	240
Busia	32.6	25.0	35.5	77.6	147	64.7	131
Siaya	21.2	16.4	22.9	57.5	145	71.0	128
Kisumu	23.9	11.8	26.1	83.5	227	82.3	203
Homa Bay	47.3	45.0	50.3	84.9	220	71.0	196
Migori	8.3	8.6	13.5	58.8	289	65.2	254
Kisii	31.7	30.8	34.1	72.7	193	61.8	165
Nyamira	29.4	28.9	32.7	72.0	87	71.0	78
Nairobi City	22.8	16.6	27.7	65.8	1,423	75.9	1,281
Total	20.4	16.5	23.8	68.4	9,119	71.5	8,125

[1] Based on mother's recall
[2] Locally available and distributed micronutrient powders
[3] Iron-containing supplements include tablets, syrups, and micronutrient powders.
[4] Based on both mother's recall and the vaccination card (where available)
[5] Deworming for intestinal parasites is commonly done for helminths and schistosomiasis.

MALARIA 12

Key Findings

- ***Ownership of insecticide-treated nets (ITNs):*** 52% of households have at least one mosquito net, 41% have at least one ITN), and 25% have at least one ITN for every two persons.
- ***Source of ITNs:*** 71% of ITNs are obtained from mass distribution campaigns, 12% from antenatal care visits, and 9% from shops/markets.
- ***Use of ITNs by children:*** 42% of children under age 5 slept under an ITN the night before the survey.
- ***Use of ITNs by pregnant women:*** 39% of pregnant women slept under an ITN the night before the survey.
- ***Use of intermittent preventive treatment in pregnancy (IPTp) by women:*** In areas where IPTp is implemented, 77% of women with a recent live birth and/or stillbirth received at least one dose of sulfadoxine-pyrimethamine (SP)/Fansidar during their last pregnancy, while 41% received three or more doses.
- ***Types of antimalarials used:*** 74% of children under age 5 with fever in the 2 weeks preceding the survey took artemisinin-combination therapy (ACT).

This chapter presents data that are useful for assessing the implementation of malaria control strategies, including the availability and use of mosquito nets, prophylactic use of antimalarial drugs among pregnant women, care seeking and therapeutic use of antimalarial drugs among children with fever, and the prevalence of anaemia and malaria among children under age 5.

According to the *World Malaria Report 2025* (World Health Organization [WHO] 2025), an estimated 282 million malaria cases occurred globally in 2024. Kenya remains among the highest-burden countries, contributing approximately 1.5% of global malaria cases.

Malaria control efforts are aligned with Sustainable Development Goal 3.3, which aims to end the malaria epidemic by 2030 (United Nations 2015). In support of this goal, the WHO Global Technical Strategy for Malaria 2016–2030 established the following milestones:

- Reduce malaria incidence and mortality rates by at least 75% by 2025, relative to 2015 levels
- Achieve at least a 90% reduction in incidence and mortality rates by 2030
- Eliminate malaria in at least 35 countries by 2030
- Stratify malaria in Kenya

To support targeted malaria interventions, the National Malaria Control Programme developed an analytical framework for stratification mapping using 5-year passive surveillance data (2019–2023), environmental and ecological factors, and Kenya Demographic and Health Survey (KDHS) data on health-seeking behaviour (Ministry of Health 2024a). Using principal component analysis, multiple correlated indicators were combined to classify Kenya's 47 counties into the following seven malaria burden strata for improved planning and resource allocation:

- **No transmission:** Nairobi was uniquely classified as a no transmission area due to historically negligible local malaria transmission, although imported cases may still occur.
- **Very low burden:** Counties in this stratum recorded a median adjusted malaria incidence of 2 cases per 1,000 population, indicating very limited malaria transmission. These areas experience sporadic and localised cases, often associated with imported infections, ecological suitability in limited pockets, or seasonal transmission. The counties in this category are Embu, Kirinyaga, Kitui, Laikipia, Lamu, Machakos, Makueni, Mandera, Murang'a, Nyandarua, Nyeri, and Wajir.
- **Low burden:** Counties in the low burden stratum had a median incidence of 11 cases per 1,000 population, characterised by low but ongoing seasonal or focal malaria transmission. Transmission is generally unstable, seasonal, or focal, with potential outbreaks under favourable environmental conditions. Counties in this category are Bomet, Garissa, Isiolo, Kajiado, Kiambu, Marsabit, Meru, Nakuru, Taita/Taveta, and Tharaka-Nithi.
- **Moderate burden:** Counties classified under the moderate burden stratum recorded a median adjusted incidence of 54 cases per 1,000 population, indicating sustained but moderate malaria transmission. Transmission patterns can vary geographically and seasonally, with some areas experiencing periodic increases in cases linked to climatic and ecological factors. The counties in this category are Elgeyo/Marakwet, Mombasa, Narok, Nyamira, Samburu, Tana River, and Uasin Gishu.
- **Moderate to high burden:** Counties in this stratum recorded a median adjusted incidence of 95 cases per 1,000 population, reflecting relatively intense and persistent malaria transmission. Transmission is often stable or seasonal, with substantial morbidity among vulnerable populations. Counties in this category are Baringo, Kericho, Kilifi, Kisii, Nandi, and Trans Nzoia.
- **High burden:** Counties in the high burden stratum had a median adjusted incidence of 474 cases per 1,000 population, indicating substantial and sustained transmission. Malaria remains a major public health problem in these areas, contributing substantially to outpatient visits, hospitalisations, and mortality, especially among children under age 5 and pregnant women. Counties in this category are Bungoma, Homa Bay, Kwale, Turkana, and West Pokot.
- **Very high burden:** Counties in this stratum recorded the highest malaria burden, with a median adjusted incidence of 748 cases per 1,000 population. Transmission is intense and perennial, with substantial impact on health systems and socioeconomic well-being. Counties in this category are Busia, Kakamega, Kisumu, Migori, Siaya, and Vihiga.

Mass Distribution of Insecticide-treated Nets

Insecticide-treated nets (ITNs) are a key malaria prevention intervention in Kenya, particularly in areas with sustained or high malaria transmission. During 2023/2024, the Ministry of Health implemented its most recent mass distribution campaign, targeting 24 counties. The campaign aimed to distribute approximately 15.3 million long-lasting insecticidal nets (LLINs) to approximately 27 million people at risk of malaria. The targeted counties were Baringo, Bungoma, Busia, Homa Bay, Kakamega, Kericho, Kilifi, Kirinyaga, Kisii, Kisumu, Kwale, Lamu, Migori, Mombasa, Nandi, Narok, Nyamira, Siaya, Taita/Taveta, Tana River, Trans Nzoia, Turkana, Vihiga, and West Pokot. However, the mass distribution campaign covered only selected sub-counties in Baringo, Kericho, Kirinyaga, and Nandi.

In addition to mass distribution campaigns every 3 years, Kenya routinely distributes ITNs through health facilities in the same counties. Routine distribution complements mass distribution by ensuring continued access to ITNs between mass distribution campaigns and by reaching vulnerable population groups through routine contact with health services. Through routine distribution, ITNs are provided to pregnant women through antenatal care (ANC) services and to children under age 1 through child welfare clinic/immunisation services. This approach is particularly important for ensuring that pregnant women and young children can

access an ITN even during periods when their county or sub-county is not covered by a mass distribution campaign.

The timing and geographic coverage of the 2023/2024 mass distribution campaign provide important context for interpreting findings on ITN ownership, access, and use from the 2025–26 Kenya Mini Demographic and Health Survey (KMDHS). Counties that received LLINs through the mass campaign may have had greater access to ITNs than counties that were not included in the campaign, although access may also have been influenced by routine distribution through health facilities and other channels. Consequently, differences in ITN-related indicators across counties should not be interpreted solely in terms of malaria endemicity.

Routine distribution is also important for maintaining ITN coverage over time, as it provides an opportunity for eligible pregnant women and young children to receive nets without having to wait for the next mass distribution campaign. It therefore serves as a complementary mechanism for sustaining access to ITNs following mass distribution.

County-level findings on ITN ownership, access, and use should be considered alongside both the malaria endemicity classification and the recent history of LLIN distribution. The households in counties that had recently received nets may have reported higher levels of ITN ownership or access. Conversely, lower coverage in a county should be interpreted in light of whether the county was included in the recent mass distribution campaign and the timing of that distribution relative to the survey.

12.1 Ownership of Insecticide-treated Nets

Ownership of insecticide-treated nets
Households that have at least one insecticide-treated net (ITN). An ITN is a factory-treated net that does not require any further treatment.
Sample: Households


Full household ITN coverage
Percentage of households with at least one ITN for every two people.
Sample: Households (with at least one person who stayed in the household the night before the survey)

Fifty-two percent of households have at least one mosquito net, while 41% have at least one ITN. On average, there are 0.8 ITNs per household (**Table 12.1**).

Twenty-five percent of households have at least one ITN for every two people who stayed in the household the night preceding the survey. Sixteen percent of households have at least one ITN but not enough ITNs for all household members, and 59% have no ITNs (**Figure 12.1**).

Figure 12.1 **Household ownership of ITNs**


Percent distribution of households



Trends: At the national level, ITN ownership rose sharply from 8% in the 2003 KDHS to 56% in the 2008–09 KDHS. It then remained relatively high through 2015 (59% in 2014 and 63% in 2015) before declining to 49% in the 2020 Kenya Malaria Indicator Survey (KMIS), recovering slightly to 54% in the 2022 KDHS and then falling again to 41% in the 2025–26 KMDHS (**Figure 12.2**).

Figure 12.2 **Trends in household ownership of ITNs**

Percentage of households owning at least one insecticide-treated net (ITN)




Note: The definition of an ITN in surveys conducted prior to the 2014 KDHS included nets that had been soaked with insecticides within the past 12 months.

Patterns by background characteristics

- Ownership of at least one ITN is higher in rural areas (51%) than in urban areas (28%).
- The percentage of households that own at least one ITN is highest in very high malaria burden areas (76%) and lowest in the no transmission area (15%).
- Ownership of at least one ITN per two people (full household coverage) is most common in high malaria burden areas (49%).
- ITN ownership is highest among households in the second wealth quintile (54%) and lowest among those in the highest wealth quintile (25%).
- By county, ownership of at least one ITN is highest in Turkana (84%), Vihiga (82%), Trans Nzoia (80%), and Migori (80%) and lowest in Nyeri (3%), Garissa (8%), and Nyandarua (9%) (**Table 12.1C** and **Map 12.1**).

Map 12.1 ITN ownership by county

Percentage of households with at least one ITN




Source of Nets

The majority of ITNs (71%) are obtained from the mass distribution campaigns. Fewer ITNs are acquired during ANC visits (12%), from shops or markets (9%), from child welfare clinic or immunisation visits (3%), and from government facilities (2%). An additional 1% are obtained from other sources (**Figure 12.3**).

Figure 12.3 Source of ITNs

Percent distribution of ITNs in households



Patterns by background characteristics

- A higher percentage of households in rural areas (77%) than of those in urban areas (57%) obtain their ITNs through mass campaigns. Shops/markets are the most common source of ITNs in the no transmission area (40%) and in very low burden areas (30%). However, in very low burden areas, ITNs are also largely sourced from ANC visits (26%) and mass campaigns (28%).
- Eighty percent of households in the lowest wealth quintile compared with only 48% of households in the highest wealth quintile obtain their ITNs through mass campaigns.

- By county, the highest percentages of households obtaining their ITNs through mass campaigns are in Busia and Nyamira (93% each), the highest percentages obtaining them from shops/markets are in Wajir (98%) and Mandera (84%), and the highest percentages obtaining them from ANC visits are in Murang'a (53%) and Elgeyo/Marakwet (49%) (**Table 12.2C**).

12.2 HOUSEHOLD ACCESS TO AND USE OF ITNS

Access to an ITN

Percentage of the population that could sleep under an ITN if each ITN in the household were used by up to two people.

Sample: De facto household population

Use of ITNs

Percentage of the population that slept under an ITN the night before the survey.

Sample: De facto household population

ITNs act as both physical and chemical barriers against mosquitoes. By reducing the vector population, ITNs help reduce malaria risk at the community level as well as among individuals who use them.


Access to an ITN is measured by the proportion of the population that could sleep under an ITN if each ITN in the household is used by up to two people. Comparing ITN access and ITN use indicators can help programmes identify any behavioural gaps in which available ITNs are not being used. If the difference between these indicators is substantial, the ITN programme may need to design an appropriate intervention that focuses on changing behaviour and identifying the main drivers of or barriers to ITN use. This analysis helps programme managers determine whether they need to achieve higher ITN coverage, promote ITN use through behavioural change communication, or both.

Thirty-six percent of the de facto population has access to an ITN (**Table 12.3**), whereas 32% of the de facto population slept under an ITN the night before the survey (**Table 12.3**, **Table 12.4**, and **Figure 12.4**).

Seven in 10 people in households with at least one ITN slept under an ITN the night before the survey. Overall, 78% of ITNs were used the night before the survey (**Tables 12.4** and **12.5**).

***Figure 12.4* Trends in ITN access and use**

Percentage of the household population that have access to an ITN and percentage of the population that slept under an ITN the night before the survey



Note: The definition of an ITN in surveys conducted prior to the 2014 KDHS included nets that had been soaked with insecticides within the past 12 months.

Trends: The percentage of the de facto population with access to an ITN increased from 5% in the 2003 KDHS to a peak of 53% in the 2015 KMIS, but has since declined to 36% in the 2025–26 KMDHS (**Figure 12.4**). Similarly, the percentage of the population that slept under an ITN the night before the survey increased from 5% in 2003 to 48% in 2015 and then declined to 33% in 2025–26.

The behavioural gap between access and use remained relatively small throughout the same period, ranging from 0 percentage points in the 2003 KDHS to 7 percentage points in the 2008–09 and 2022 KDHS surveys. It has narrowed to 4 percentage points in the 2025–26 KMDHS, suggesting that most available ITNs are being


used (**Figure 12.4**). The narrow gap of 4 percentage points between ITN access (36%) and use (32%) demonstrates a strong community acceptance.

Patterns by background characteristics

- Access to and use of ITNs is higher in rural areas (43% and 39%, respectively) than in urban areas (25% and 22%, respectively) (**Tables 12.3** and **12.4**)
- Use of ITNs is higher in rural (39%) than urban (22%) areas (**Table 12.4**).
- Access to and use of ITNs substantially increases by malaria burden, from 11% and 9%, respectively, in no transmission zones to 64% and 58%, respectively, in very high burden zones.
- However, among households that own at least one ITN, use is much higher in general and varies much more by burden area (ranging from 61% in low burden areas to 76% in moderate to high burden areas).
- By county, ITN access is highest among those living in Vihiga (75%) and in Trans Nzoia, Turkana, and Nyamira (70% each) and lowest among those living in Nyeri (3%), Garissa (4%), Nyandarua (7%), and Samburu (7%) (**Table 12.3C**).
- ITN use is highest among those living in Trans Nzoia (67%), Kilifi (66%), Tana River (65%), Vihiga (65%), and Turkana (65%) Counties and lowest among those living in Nyeri (2%), Garissa (4%), and Marsabit (8%) Counties (**Table 12.4C** and **Map 12.2**).

Map 12.2 ITN use by county

Percent of the household population who slept under an ITN the night before the survey by county



12.3 USE OF ITNS BY CHILDREN AND PREGNANT WOMEN

Forty-two percent of children under age 5 and 39% of pregnant women age 15–49 slept under an ITN the night before the survey (**Table 12.6**, **Table 12.6C**, **Table 12.7**, and **Figure 12.5**).

ITN use is higher in households with at least one ITN: 79% of children under age 5 and 82% of pregnant women in households with at least one ITN slept under an ITN the night before the survey.

Trends: The use of ITNs among children under age 5 in households with at least one ITN increased from 69% in the 2008–09 KDHS to 79% in the 2015 KMIS. It then declined to 77% in the 2022 KDHS, rebounding to 79% in the 2025–26 KMDHS. Similarly, the percentage of pregnant women who reported sleeping under an ITN the night before the survey in households with at least one ITN grew from 77% in 2008–09 to 82% in 2015, fell to 75% in 2022, and then returned to 82% in 2025–26 (**Figure 12.6**).

***Figure 12.5* ITN use**

Percentage who slept under an ITN the night before the survey





***Figure 12.6* Trends in ITN use among pregnant women and children under age 5**

Percentage of pregnant women and children under 5 who slept under an ITN the night before the survey in households with at least one ITN



Note: The definition of an ITN in surveys conducted prior to the 2014 KDHS included nets that had been soaked with insecticides within the past 12 months.

Patterns by background characteristics


- Use of ITNs by children under age 5 in households with at least one ITN decreases from 83% among children younger than 12 months of age to 75% among those 48–59 months of age.
- Among households with at least one ITN, ITN use among children under age 5 is high across all malaria burden zones (ranging from 76% to 83%) and is highest in the high burden zone (83%). Similarly, among households with at least one ITN, ITN use among pregnant women is high across all malaria burden zones (ranging from 75% to 87%) and is highest in the low burden zone (87%).

12.4 Reasons Mosquito Nets Were Not Used

Overall, 22% of ITNs were not used by anyone on the night preceding the survey, a decrease of 6% compared with the 2022 KDHS. Additionally, 48% of respondents reported that the reason for not using the ITNs was that they were extra nets or were reserved for future use, an 11 percentage-point decrease from the 2022 survey. This suggests a positive behavioural shift towards using nets immediately upon receipt (**Table 12.8**, **Table 12.8C**, and **Figure 12.7**).

Figure 12.7 **Reason ITN was not used**

Among ITNs that were not used the night before the survey, main reason the net was not used (percentage)



Patterns by background characteristics

- Households in low burden areas had the highest percentage of unused ITNs (26%), followed by those in very high burden areas (22%).
- The main reason for ITN non-use varies by malaria burden. In the no transmission area, the most common reason for non-use is the absence of mosquitoes or malaria (though this is based on very few cases). In very low burden areas, the main reason is that the usual user did not sleep in the household the night before the survey. In all areas from low to very high burden, the most common reason for non-use was that the ITN was an extra net or was being saved for later.

12.5 Malaria in Pregnancy

Intermittent preventive treatment (IPTp) during pregnancy

Percentage of women who took at least three doses of sulfadoxine-pyrimethamine (SP)/Fansidar during their last pregnancy.

Sample: Women age 15–49 with a live birth or a stillbirth in the 2 years before the survey

Malaria infection during pregnancy is a major public health problem in Kenya, with substantial risks for the mother, her foetus, and the neonate. Intermittent preventive treatment in pregnancy (IPTp) is a full therapeutic course of antimalarial medicine given to pregnant women at routine ANC visits to prevent malaria. IPTp helps prevent maternal malaria episodes, maternal and foetal anaemia, placental parasitaemia, low birth weight, and neonatal mortality.

The WHO recommends **at least three doses** of sulfadoxine-pyrimethamine (SP) during pregnancy, administered at each scheduled ANC visit starting in the second trimester. In Kenya, IPTp is given to pregnant women in 15 full-coverage counties and four fringe areas. The full-coverage counties are Mombasa, Kilifi, Kwale, Taita/Taveta, Tana River, Lamu, Bungoma, Kakamega, Busia, Vihiga, Siaya, Kisumu, Migori, Homabay, and Turkana.

Twenty-nine percent of women with a live birth and/or stillbirth in the 2 years before the survey reported receiving one or more doses of SP/Fansidar during the pregnancy that resulted in the last live birth or stillbirth, while 21% received two or more doses. Nationally, 14% of women received three or more doses of SP/Fansidar (**Tables 12.9** and **12.9C**).


Notably, in areas where IPTp is implemented, 77% of women with a live birth and/or stillbirth in the 2 years before the survey reported receiving one or more doses of SP/Fansidar during the pregnancy that resulted in the last live birth or stillbirth, while 61% received two or more doses and 41% received three or more doses of SP/Fansidar (**Table 12.9**).

The national IPTp coverage of 14% (IPTp 3+) masks the reality that in targeted implementation zones, coverage of three or more doses of SP/Fansidar reaches 41%. However, this still falls short of the national target of 80%, indicating both geographic and systemic challenges in scaling this intervention

Trends: Among women with a live birth and/or stillbirth in the 2 years before the survey, the percentage who reported receiving at least three doses of SP/Fansidar during their last pregnancy fluctuated from 7% in the 2008–09 KDHS to 23% in the 2015 KMIS to 14% in the 2025–26 KMDHS (**Figure 12.8**).

Figure 12.8 **Trends in IPTp use by pregnant women**

Percentage of women with a live birth in the 2 years before the survey who received at least 1, 2, or 3 doses of SP/Fansidar



Patterns by background characteristics

- In the counties where IPTp is implemented, 77% of women received one or more doses of SP/Fansidar, 61% received two or more doses, and 41% received the recommended three or more doses during the pregnancy that resulted in their most recent live birth or stillbirth.
- A higher percentage of women in rural areas (17%) than in urban areas (10%) received the recommended three or more doses of SP/Fansidar during the pregnancy that resulted in their most recent live birth or stillbirth.
- The percentage of women who received three or more doses of SP/Fansidar during the pregnancy that resulted in their most recent live birth or stillbirth increases with malaria burden, from 4% in the no transmission area to 45% in very high burden areas.

12.6 Case Management of Malaria in Children

Care seeking for children under age 5 with a fever

Percentage of children under age 5 with a fever in the 2 weeks before the survey for whom advice or treatment was sought from a health provider, a health facility, or a pharmacy.

Sample: Children under age 5 with a fever in the 2 weeks before the survey

Diagnosis of malaria in children under age 5 with a fever

Percentage of children under age 5 with a fever in the 2 weeks before the survey who had blood taken from a finger or heel for testing. This is a proxy measure of diagnostic testing for malaria.

Sample: Children under age 5 with a fever in the 2 weeks before the survey

Artemisinin-based combination therapy (ACT) for children under age 5 with a fever

Percentage of children under age 5 with a fever in the 2 weeks before the survey who received artemisinin-based combination therapy (ACT).

Sample: Children under age 5 with a fever in the 2 weeks before the survey who took any antimalarial drug

Fourteen percent of children under age 5 had fever in the 2 weeks before the survey. Of these children, 68% were taken for advice or treatment, including 44% who were taken for advice or treatment the same or following day. Thirty percent of children under age 5 with fever had blood taken from their finger or heel for testing, whereas 16% were diagnosed with malaria by a health care provider (**Table 12.10**).


Among the children with fever in the 2 weeks before the survey who were taken for advice or treatment, 51% were taken to the public medical sector, including 22% who visited government dispensaries; 49% were taken to the private medical sector, including 22% who visited a pharmacy (**Table 12.11**).

Seventy-four percent of the children under age 5 with fever in the 2 weeks preceding the survey took artemisinin-based combination therapy (ACT), most commonly artemether-lumefantrine (64%) (**Table 12.12**).

Trends: Among children under age 5 with fever in the 2 weeks before the survey, the percentage who received ACT declined from 91% in the 2020 KMIS to 84% in the 2022 KDHS and further to 74% in the 2025–26 KMDHS (**Figure 12.9**).

Figure 12.9 **Trend in ACT use by children with fever**

Among children with recent fever who took an antimalarial, percentage who received ACT



Patterns by background characteristics

- The percentage of children under age 5 with fever who were diagnosed with malaria by a health care provider is higher in rural areas (23%) than in urban areas (8%).
- The percentage of children who sought advice or treatment on the same or next day after fever onset is slightly higher in urban areas (47%) than in rural areas (41%).
- Thirty-eight percent of children whose mothers have no education sought advice or treatment on the same or next day after fever onset compared with 52% of those whose mothers have a secondary or higher education.
- A higher percentage of children in rural areas (76%) than in urban areas (68%) took ACT.

List of Tables

For more information on malaria, see the following tables:

- **Table 12.1 Household possession of mosquito nets**
- **Table 12.1C Household possession of mosquito nets by county**
- **Table 12.2 Source of mosquito nets**
- **Table 12.2C Source of mosquito nets by county**
- **Table 12.3 Access to an insecticide-treated net**
- **Table 12.3C Access to an insecticide-treated net by county**
- **Table 12.4 Use of mosquito nets by persons in the household**
- **Table 12.4C Use of mosquito nets by persons in the household by county**
- **Table 12.5 Use of existing**
- **Table 12.5C Use of existing by county**
- **Table 12.6 Use of mosquito nets by children**
- **Table 12.6C Use of mosquito nets by children by county**
- **Table 12.7 Use of mosquito nets by pregnant women**
- **Table 12.8 Main reason mosquito net was not used the night before the survey**
- **Table 12.8C Main reason mosquito net was not used the night before the survey by county**
- **Table 12.9 Use of Intermittent Preventive Treatment (IPTp) by women during pregnancy**
- **Table 12.9C Use of Intermittent Preventive Treatment (IPTp) by women during pregnancy by county**
- **Table 12.10 Children with fever and careseeking, prompt treatment, and diagnosis**
- **Table 12.11 Source of advice or treatment for children with fever**
- **Table 12.12 Type of antimalarial drugs used**

Table 12.1 Household possession of mosquito nets

Percentage of households with at least one mosquito net (treated or untreated) and insecticide-treated net (ITN); average number of nets and ITNs per household; and percentage of households with at least one net and ITN per two persons who stayed in the household last night, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Percentage of households with at least one mosquito net		Average number of nets per household		Number of households	Percentage of households with at least one net for every two persons who stayed in the household last night[1]		Number of households with at least one person who stayed in the household last night
	Any mosquito net	Insecticide-treated mosquito net (ITN)[2]	Any mosquito net	Insecticide-treated mosquito net (ITN)[2]		Any mosquito net	Insecticide-treated mosquito net (ITN)[2]	
Residence								
Urban	47.5	28.4	0.8	0.5	11,975	31.4	17.8	11,813
Rural	56.0	50.8	1.2	1.1	14,405	35.0	31.0	14,340
Malaria burden								
No transmission	42.8	15.2	0.6	0.2	3,992	25.5	8.6	3,896
Very low burden	35.3	24.3	0.6	0.4	4,891	23.1	14.6	4,850
Low burden	37.4	25.4	0.7	0.4	6,452	23.0	14.7	6,431
Moderate burden	58.2	50.5	1.2	1.0	2,981	37.3	30.3	2,950
Moderate to high burden	68.5	65.0	1.5	1.4	2,809	47.4	44.0	2,790
High burden	78.4	73.8	1.8	1.6	2,114	47.1	42.8	2,108
Very high burden	82.2	76.0	1.8	1.6	3,141	55.1	49.4	3,127
Wealth quintile								
Lowest	52.2	48.3	1.0	0.9	4,299	26.4	24.3	4,279
Second	57.5	54.1	1.2	1.1	4,719	34.5	32.4	4,698
Middle	53.4	46.6	1.1	1.0	5,346	35.9	30.4	5,304
Fourth	50.8	34.6	0.9	0.6	6,192	35.4	23.1	6,124
Highest	47.9	24.8	0.9	0.5	5,825	33.2	16.8	5,748
Total	52.1	40.6	1.0	0.8	26,380	33.4	25.1	26,153

[1] De facto household members
[2] An insecticide-treated net (ITN) is a factory-treated net that does not require any further treatment. In KDHS and KMIS surveys conducted prior to 2020, this was known as a long-lasting insecticidal net (LLIN).

Table 12.1C Household possession of mosquito nets by county

Percentage of households with at least one mosquito net (treated or untreated) and insecticide-treated net (ITN); average number of nets and ITNs per household; and percentage of households with at least one net and ITN per two persons who stayed in the household last night, according to county, Kenya Mini DHS 2025–26

County	Percentage of households with at least one mosquito net		Average number of nets per household		Number of households	Percentage of households with at least one net for every two persons who stayed in the household last night[1]		Number of households with at least one person who stayed in the household last night
	Any mosquito net	Insecticide-treated mosquito net (ITN)[2]	Any mosquito net	Insecticide-treated mosquito net (ITN)[2]		Any mosquito net	Insecticide-treated mosquito net (ITN)[2]	
Mombasa	66.2	52.9	1.2	0.9	866	43.7	30.1	853
Kwale	78.1	70.5	1.9	1.7	318	46.9	39.7	317
Kilifi	79.9	73.9	1.7	1.6	699	58.5	53.1	693
Tana River	87.2	74.9	1.9	1.6	143	50.6	39.6	143
Lamu	71.0	68.0	1.5	1.4	62	50.2	47.9	62
Taita/Taveta	84.5	76.1	2.0	1.8	206	68.2	58.4	204
Garissa	78.0	8.4	2.2	0.1	145	46.6	3.0	145
Wajir	64.0	25.5	1.7	0.5	126	31.2	6.7	125
Mandera	64.3	17.5	1.6	0.3	137	28.5	4.8	136
Marsabit	49.6	31.7	0.8	0.5	138	16.7	8.4	137
Isiolo	49.2	29.8	0.9	0.5	87	26.9	12.8	87
Meru	29.5	24.1	0.5	0.4	840	16.7	13.1	832
Tharaka-Nithi	37.0	31.6	0.7	0.6	209	23.7	18.6	208
Embu	31.2	24.2	0.5	0.4	385	21.3	15.5	378
Kitui	37.8	30.5	0.7	0.6	526	23.7	17.6	520
Machakos	39.2	24.2	0.6	0.3	1,021	22.6	10.7	1,016
Makueni	49.5	35.3	0.9	0.6	506	34.9	21.2	506
Nyandarua	13.7	8.9	0.2	0.1	341	8.0	4.0	339
Nyeri	13.3	3.4	0.2	0.0	441	6.5	1.4	437
Kirinyaga	49.8	44.6	1.0	0.9	482	44.2	39.5	471
Murang'a	25.7	18.8	0.5	0.3	571	18.1	11.1	568
Kiambu	30.0	16.8	0.5	0.2	2,295	19.5	10.0	2,295
Turkana	88.1	84.4	2.4	2.2	232	55.0	51.5	231
West Pokot	68.2	65.4	1.5	1.5	251	31.1	28.8	249
Samburu	22.7	12.9	0.3	0.2	109	5.8	2.3	109
Trans Nzoia	84.8	80.4	2.0	1.9	393	61.1	56.8	393
Uasin Gishu	51.3	42.0	1.0	0.8	794	32.9	25.5	780
Elgeyo/Marakwet	35.2	35.0	0.5	0.5	209	17.2	17.1	207
Nandi	57.0	53.8	1.1	1.0	414	36.5	32.3	414
Baringo	52.3	50.5	1.1	1.0	242	31.7	30.2	242
Laikipia	18.5	15.4	0.3	0.3	293	11.7	9.9	292
Nakuru	27.6	16.6	0.4	0.2	1,442	17.0	9.9	1,435
Narok	53.0	51.3	1.2	1.1	576	32.6	31.0	574
Kajiado	44.7	34.5	0.7	0.5	653	26.8	19.2	651
Kericho	73.3	69.6	1.5	1.4	464	48.5	45.4	455
Bomet	70.8	64.8	1.4	1.4	437	40.1	35.3	437
Kakamega	81.1	74.0	1.7	1.6	850	54.5	49.3	847
Vihiga	83.9	81.7	2.0	1.9	267	62.6	60.2	267
Bungoma	79.5	76.6	1.8	1.7	746	53.1	50.0	744
Busia	80.9	78.6	1.9	1.9	429	56.9	55.0	426
Siaya	71.2	68.7	1.3	1.2	482	44.1	41.7	479
Kisumu	86.8	76.6	1.8	1.5	615	58.3	47.9	609
Homa Bay	77.7	71.3	1.4	1.3	568	43.2	37.5	567
Migori	89.3	80.2	2.1	1.9	499	57.3	48.4	499
Kisii	55.4	54.6	1.2	1.1	597	38.5	37.4	593
Nyamira	79.6	78.2	1.9	1.9	284	59.5	57.8	284
Nairobi City	42.8	15.2	0.6	0.2	3,992	25.5	8.6	3,896
Total	52.1	40.6	1.0	0.8	26,380	33.4	25.1	26,153

[1] De facto household members

[2] An insecticide-treated net (ITN) is a factory-treated net that does not require any further treatment. In KDHS and KMIS surveys conducted prior to 2020, this was known as a long-lasting insecticidal net (LLIN).

Table 12.2 Source of mosquito nets

Percent distribution of insecticide treated nets (ITNs), non-ITNs, and all mosquito nets by source of net, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Mass distribution campaign	ANC visit	Child welfare visit/ immunisation visit	Government health facility	Private health facility	Pharmacy	Shop/ market	Community health worker	Religious institution	School	Other	Don't know/ missing	Total	Number of mosquito nets
							ITNs[1]							
Residence														
Urban	57.2	14.8	2.9	2.0	0.3	0.5	19.6	0.1	0.0	0.0	2.2	0.3	100.0	5,672
Rural	76.7	11.2	3.2	1.4	0.2	0.4	5.5	0.1	0.0	0.2	0.9	0.1	100.0	15,368
Malaria burden														
No transmission	37.1	12.1	0.6	6.2	0.0	0.0	39.7	0.5	0.0	0.0	3.8	0.0	100.0	843
Very low burden	28.2	26.1	8.5	4.3	0.6	1.2	30.1	0.0	0.0	0.0	0.7	0.3	100.0	2,021
Low burden	42.1	21.0	6.7	1.8	0.9	2.0	23.3	0.0	0.0	0.0	1.7	0.5	100.0	2,798
Moderate burden	79.2	11.9	2.1	0.8	0.1	0.0	4.8	0.1	0.0	0.1	1.0	0.0	100.0	2,958
Moderate to high burden	84.3	9.4	2.0	0.7	0.0	0.1	2.5	0.0	0.0	0.0	0.6	0.2	100.0	3,867
High burden	84.8	8.1	1.9	1.3	0.5	0.1	1.1	0.0	0.0	0.6	1.6	0.0	100.0	3,459
Very high burden	87.0	6.9	1.8	0.8	0.0	0.1	1.6	0.4	0.0	0.0	1.2	0.1	100.0	5,093
Wealth quintile														
Lowest	80.1	11.6	2.7	0.8	0.0	0.1	2.9	0.1	0.0	0.1	1.4	0.0	100.0	3,937
Second	80.2	10.9	3.2	1.1	0.1	0.2	3.2	0.2	0.0	0.1	0.7	0.1	100.0	5,245
Middle	75.2	11.9	3.3	1.6	0.3	0.4	5.7	0.1	0.0	0.2	1.0	0.1	100.0	5,118
Fourth	62.8	14.4	3.8	1.3	0.4	0.8	14.1	0.3	0.0	0.1	1.7	0.2	100.0	4,001
Highest	47.7	12.7	2.2	3.8	0.6	0.8	29.9	0.0	0.0	0.0	1.8	0.5	100.0	2,738
Total	71.4	12.2	3.1	1.5	0.3	0.4	9.3	0.1	0.0	0.1	1.3	0.2	100.0	21,040
							NON-ITNs							
Total	na	na	na	na	0.0	0.3	96.7	0.0	0.2	0.2	1.9	0.7	100.0	5,655
							ALL MOSQUITO NETS							
Total	56.3	9.6	2.5	1.2	0.2	0.4	27.8	0.1	0.1	0.2	1.4	0.3	100.0	26,695

na = not applicable
ANC = Antenatal care
[1] An insecticide-treated net (ITN) is a factory-treated net that does not require any further treatment. In KDHS and KMIS surveys conducted prior to 2020, this was known as a long-lasting insecticidal net (LLIN).

Table 12.2C Source of mosquito nets by county

Percent distribution of insecticide treated nets (ITNs), non-ITNs, and all mosquito nets by source of net, according to county, Kenya Mini DHS 2025–26

County	Mass distribution campaign	ANC visit	Child welfare visit/ immunisation visit	Government health facility	Private health facility	Pharmacy	Shop/ market	Community health worker	Religious institution	School	Other	Don't know/ missing	Total	Number of mosquito nets
							ITNs[1]							
Mombasa	77.2	13.4	0.8	0.2	0.0	0.0	6.2	0.2	0.0	0.0	1.9	0.0	100.0	793
Kwale	87.5	10.4	0.0	0.5	0.1	0.0	0.2	0.0	0.1	0.0	1.3	0.0	100.0	542
Kilifi	87.2	5.0	1.0	1.4	0.0	0.0	4.8	0.0	0.0	0.1	0.6	0.0	100.0	1,103
Tana River	88.8	8.5	1.6	0.3	0.0	0.0	0.6	0.0	0.0	0.0	0.3	0.0	100.0	233
Lamu	83.8	4.1	4.8	0.8	0.0	0.0	5.7	0.0	0.0	0.0	0.8	0.0	100.0	88
Taita/Taveta	85.2	10.4	2.5	1.2	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.2	100.0	370
Garissa	14.5	9.7	0.0	2.4	1.1	4.0	68.3	0.0	0.0	0.0	0.0	0.0	100.0	21
Wajir	0.3	0.0	1.7	0.0	0.0	0.0	97.6	0.0	0.0	0.4	0.0	0.0	100.0	59
Mandera	9.4	3.9	0.0	1.5	0.0	1.5	83.7	0.0	0.0	0.0	0.0	0.0	100.0	46
Marsabit	62.5	16.1	6.5	0.5	0.0	0.0	13.4	0.0	0.0	0.0	1.1	0.0	100.0	69
Isiolo	21.0	24.0	11.9	1.5	0.6	1.0	39.6	0.0	0.0	0.0	0.4	0.0	100.0	46
Meru	11.0	26.0	22.4	1.5	2.1	8.2	27.8	0.0	0.0	0.0	0.8	0.2	100.0	337
Tharaka-Nithi	27.0	20.2	19.8	6.5	1.9	2.4	17.3	0.0	0.0	0.7	3.3	0.9	100.0	121
Embu	8.3	32.0	14.3	1.8	1.8	0.0	40.3	0.0	0.0	0.0	1.4	0.0	100.0	152
Kitui	15.7	34.5	12.5	4.2	0.0	3.5	29.6	0.0	0.0	0.0	0.0	0.0	100.0	301
Machakos	19.8	34.1	8.4	1.3	0.0	0.8	35.3	0.0	0.0	0.0	0.0	0.4	100.0	341
Makueni	8.2	35.8	20.8	12.5	0.4	1.2	18.4	0.0	0.3	0.0	1.9	0.4	100.0	285
Nyandarua	29.1	14.7	6.0	2.1	0.0	0.0	42.9	0.0	0.0	0.0	5.2	0.0	100.0	45
Nyeri	(33.2)	(11.4)	(25.8)	(5.4)	(10.7)	(0.0)	(13.5)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	100.0	19
Kirinyaga	66.7	9.4	2.0	4.0	0.7	0.4	15.8	0.0	0.0	0.0	0.2	0.7	100.0	429
Murang'a	11.7	52.5	1.4	5.4	1.3	0.4	26.7	0.0	0.0	0.0	0.6	0.0	100.0	182
Kiambu	13.5	31.5	3.1	1.3	0.2	1.7	44.4	0.0	0.0	0.0	2.8	1.6	100.0	557
Turkana	82.9	7.8	1.6	1.1	0.0	0.0	1.2	0.0	0.1	0.0	5.4	0.0	100.0	522
West Pokot	89.8	9.2	0.8	0.0	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.0	100.0	369
Samburu	44.2	14.2	8.5	0.0	0.0	1.6	27.4	0.0	0.0	0.0	1.6	2.5	100.0	19
Trans Nzoia	91.5	5.2	0.7	0.6	0.0	0.2	0.9	0.0	0.2	0.1	0.6	0.0	100.0	751
Uasin Gishu	66.2	18.5	5.2	1.1	0.2	0.0	7.1	0.0	0.0	0.6	1.1	0.0	100.0	610
Elgeyo/Marakwet	13.3	49.1	5.0	10.2	0.6	0.4	20.5	0.0	0.0	0.0	0.9	0.1	100.0	112
Nandi	62.9	34.8	1.1	0.0	0.0	0.1	0.5	0.0	0.0	0.0	0.0	0.6	100.0	429
Baringo	64.7	11.5	8.7	2.6	0.0	0.0	9.4	0.0	0.0	0.0	0.9	2.2	100.0	253
Laikipia	17.8	8.4	0.0	2.7	1.6	5.5	62.5	0.0	0.0	0.0	1.5	0.0	100.0	75
Nakuru	44.4	16.9	3.6	5.9	3.7	2.4	18.7	0.0	0.0	0.0	3.8	0.6	100.0	351
Narok	90.9	5.4	1.4	0.1	0.0	0.0	1.7	0.0	0.0	0.0	0.5	0.0	100.0	654
Kajiado	10.4	23.1	10.8	1.0	0.0	1.2	50.5	0.0	0.0	0.0	3.0	0.0	100.0	333
Kericho	86.3	7.8	3.0	0.4	0.0	0.0	1.7	0.0	0.0	0.0	0.9	0.0	100.0	651
Bomet	79.3	17.3	0.6	0.1	0.0	0.1	2.4	0.0	0.0	0.0	0.0	0.0	100.0	593
Kakamega	88.9	4.4	1.4	0.9	0.0	0.1	0.5	1.7	0.0	0.1	2.1	0.0	100.0	1,332
Vihiga	92.0	6.1	0.8	0.0	0.0	0.0	0.6	0.0	0.0	0.0	0.4	0.0	100.0	517
Bungoma	91.7	5.1	1.7	0.2	0.0	0.0	0.4	0.0	0.0	0.0	1.0	0.0	100.0	1,294
Busia	92.5	2.4	0.3	0.0	0.0	0.0	4.4	0.0	0.0	0.0	0.5	0.0	100.0	796
Siaya	88.2	8.1	2.1	0.0	0.0	0.3	1.3	0.0	0.0	0.0	0.1	0.0	100.0	573
Kisumu	87.4	8.1	3.0	0.6	0.1	0.1	0.5	0.0	0.0	0.0	0.1	0.1	100.0	930
Homa Bay	69.7	11.5	4.3	4.6	2.1	0.4	3.5	0.0	0.1	2.7	1.0	0.0	100.0	732
Migori	75.9	12.8	2.6	2.2	0.0	0.1	3.0	0.0	0.0	0.0	3.0	0.3	100.0	945
Kisii	90.8	5.7	2.1	0.0	0.2	0.3	0.3	0.0	0.0	0.0	0.5	0.1	100.0	681
Nyamira	93.4	3.5	0.4	0.5	0.0	0.0	1.5	0.0	0.0	0.0	0.6	0.0	100.0	537
Nairobi City	37.1	12.1	0.6	6.2	0.0	0.0	39.7	0.5	0.0	0.0	3.8	0.0	100.0	843
Total	71.4	12.2	3.1	1.5	0.3	0.4	9.3	0.1	0.0	0.1	1.3	0.2	100.0	21,040
							NON-ITNs							
Total	na	na	na	na	0.0	0.3	96.7	0.0	0.2	0.2	1.9	0.7	100.0	5,655
							ALL MOSQUITO NETS							
Total	56.3	9.6	2.5	1.2	0.2	0.4	27.8	0.1	0.1	0.2	1.4	0.3	100.0	26,695

Note: Figures in parentheses are based on 25–49 unweighted cases.
na = not applicable
ANC = Antenatal care
[1] An insecticide-treated net (ITN) is a factory-treated net that does not require any further treatment. In KDHS and KMIS surveys conducted prior to 2020, this was known as a long-lasting insecticidal net (LLIN).

Table 12.3 Access to an insecticide-treated net (ITN)

Percentage of the de facto population with access to an ITN in the household, by background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Percentage of the de facto population with access to an ITN[1,2]	Number of persons
Residence		
Urban	24.6	35,243
Rural	43.3	54,867
Malaria burden		
No transmission	11.0	12,318
Very low burden	20.2	15,361
Low burden	21.5	20,239
Moderate burden	42.6	10,681
Moderate to high burden	56.3	10,399
High burden	60.6	9,101
Very high burden	63.8	12,011
Wealth quintile		
Lowest	36.6	17,874
Second	46.2	18,045
Middle	42.9	18,092
Fourth	32.6	18,060
Highest	21.5	18,039
Total	36.0	90,110

[1] An insecticide-treated net (ITN) is a factory-treated net that does not require any further treatment. In KDHS and KMIS surveys conducted prior to 2020, this was known as a long-lasting insecticidal net (LLIN).
[2] Percentage of the de facto household population who could sleep under an ITN if each ITN in the household were used by up to two people

Table 12.3C Access to an insecticide-treated net (ITN) by county

Percentage of the de facto population with access to an ITN in the household, by county, Kenya Mini DHS 2025–26

County	Percentage of the de facto population with access to an ITN[1,2]	Number of persons
Mombasa	43.7	2,887
Kwale	57.7	1,547
Kilifi	65.2	2,569
Tana River	58.8	665
Lamu	56.2	249
Taita/Taveta	69.1	671
Garissa	4.3	821
Wajir	14.5	763
Mandera	10.2	793
Marsabit	19.9	651
Isiolo	22.1	358
Meru	20.8	2,611
Tharaka-Nithi	26.9	666
Embu	23.8	1,014
Kitui	22.5	2,012
Machakos	18.8	2,961
Makueni	28.7	1,604
Nyandarua	7.3	1,044
Nyeri	3.0	1,107
Kirinyaga	47.3	1,151
Murang'a	16.8	1,762
Kiambu	14.6	6,099
Turkana	69.8	1,120
West Pokot	52.6	1,209
Samburu	6.8	534
Trans Nzoia	70.4	1,566
Uasin Gishu	37.2	2,543
Elgeyo/Marakwet	22.6	837
Nandi	45.0	1,494
Baringo	40.8	978
Laikipia	12.7	901
Nakuru	12.1	4,462
Narok	46.5	2,194
Kajiado	27.5	1,982
Kericho	61.5	1,671
Bomet	49.0	1,918
Kakamega	62.6	3,094
Vihiga	75.1	1,026
Bungoma	65.8	3,034
Busia	68.4	1,698
Siaya	50.9	1,773
Kisumu	64.0	2,240
Homa Bay	55.1	2,192
Migori	67.1	2,181
Kisii	46.2	2,121
Nyamira	69.6	1,020
Nairobi City	11.0	12,318
Total	36.0	90,110

[1] An insecticide-treated net (ITN) is a factory-treated net that does not require any further treatment. In KDHS and KMIS surveys conducted prior to 2020, this was known as a long-lasting insecticidal net (LLIN).
[2] Percentage of the de facto household population who could sleep under an ITN if each ITN in the household were used by up to two people

Table 12.4 Use of mosquito nets by persons in the household

Percentage of the de facto household population who slept under a mosquito net (treated or untreated) and under an insecticide-treated net (ITN) the night before the survey; and among the de facto household population in households with at least one ITN, percentage who slept under an ITN the night before the survey, according to background characteristics, Kenya Mini DHS 2025–26

	Household population			Household population in households with at least one ITN[1]	
Background characteristic	Percentage who slept under any mosquito net last night	Percentage who slept under an ITN[1] last night	Number of persons	Percentage who slept under an ITN[1] last night	Number of persons
Age					
<5	53.0	41.7	10,881	79.4	5,711
5–14	38.4	31.4	22,563	63.7	11,114
15–34	36.2	27.5	30,393	63.3	13,197
35–49	44.8	35.0	14,082	78.9	6,245
50+	42.7	34.7	12,121	78.8	5,343
Don't know/ Missing	38.3	23.0	71	(51.5)	32
Sex					
Male	38.1	30.1	44,730	66.4	20,272
Female	43.8	34.5	45,380	73.3	21,370
Residence					
Urban	38.1	22.1	35,243	69.8	11,142
Rural	42.8	38.9	54,867	70.0	30,500
Malaria burden					
No transmission	29.4	9.2	12,318	62.0	1,827
Very low burden	26.5	17.3	15,361	63.5	4,176
Low burden	27.4	17.7	20,239	61.0	5,884
Moderate burden	45.0	39.2	10,681	71.3	5,862
Moderate to high burden	54.7	52.0	10,399	75.7	7,142
High burden	60.3	56.5	9,101	71.8	7,161
Very high burden	64.1	58.4	12,011	73.2	9,590
Wealth quintile					
Lowest	37.8	34.8	17,874	67.0	9,281
Second	44.7	42.3	18,045	71.1	10,735
Middle	44.1	38.3	18,092	72.1	9,607
Fourth	42.0	28.9	18,060	71.4	7,309
Highest	36.3	17.4	18,039	66.4	4,711
Total	41.0	32.3	90,110	69.9	41,642

Note: Figures in parentheses are based on 25–49 unweighted cases.
[1] An insecticide-treated net (ITN) is a factory-treated net that does not require any further treatment. In KDHS and KMIS surveys conducted prior to 2020, this was known as a long-lasting insecticidal net (LLIN).

Table 12.4C Use of mosquito nets by persons in the household by county

Percentage of the de facto household population who slept under a mosquito net (treated or untreated) and under an insecticide-treated net (ITN) the night before the survey; and among the de facto household population in households with at least one ITN, percentage who slept under an ITN the night before the survey, according to county, Kenya Mini DHS 2025–26

	Household population			Household population in households with at least one ITN[1]	
County	Percentage who slept under any mosquito net last night	Percentage who slept under an ITN[1] last night	Number of persons	Percentage who slept under an ITN[1] last night	Number of persons
Mombasa	52.8	41.7	2,887	73.4	1,641
Kwale	63.8	58.0	1,547	78.3	1,146
Kilifi	69.6	65.8	2,569	84.8	1,995
Tana River	76.1	65.3	665	83.3	522
Lamu	54.0	52.0	249	79.0	164
Taita/Taveta	59.6	51.4	671	64.7	533
Garissa	58.0	3.7	821	49.9	61
Wajir	39.2	11.7	763	44.8	199
Mandera	39.0	8.8	793	43.1	162
Marsabit	17.3	8.4	651	25.2	216
Isiolo	39.2	23.4	358	67.1	125
Meru	21.5	17.5	2,611	61.0	751
Tharaka-Nithi	26.7	22.9	666	63.5	240
Embu	23.9	19.3	1,014	65.3	300
Kitui	20.8	17.2	2,012	55.6	624
Machakos	31.4	17.0	2,961	60.2	836
Makueni	34.5	24.4	1,604	63.3	618
Nyandarua	8.6	5.8	1,044	53.3	113
Nyeri	9.1	2.4	1,107	51.3	52
Kirinyaga	47.3	43.2	1,151	86.2	576
Murang'a	19.0	14.2	1,762	65.8	380
Kiambu	22.9	11.9	6,099	61.5	1,178
Turkana	69.3	64.9	1,120	73.5	988
West Pokot	47.2	45.2	1,209	60.4	904
Samburu	10.1	5.6	534	40.8	73
Trans Nzoia	71.1	67.3	1,566	81.7	1,291
Uasin Gishu	39.5	32.7	2,543	67.5	1,232
Elgeyo/Marakwet	20.1	20.0	837	59.7	281
Nandi	40.7	37.7	1,494	64.9	868
Baringo	36.4	34.9	978	64.3	530
Laikipia	13.4	10.4	901	61.6	153
Nakuru	17.5	9.9	4,462	58.6	755
Narok	40.2	39.1	2,194	66.4	1,292
Kajiado	33.9	26.3	1,982	69.3	752
Kericho	52.3	49.3	1,671	64.7	1,273
Bomet	43.5	40.6	1,918	61.2	1,272
Kakamega	60.5	55.4	3,094	70.1	2,447
Vihiga	67.1	64.6	1,026	74.7	887
Bungoma	61.7	59.3	3,034	72.7	2,474
Busia	65.3	63.5	1,698	76.8	1,402
Siaya	51.7	49.2	1,773	72.4	1,205
Kisumu	71.4	61.4	2,240	76.6	1,795
Homa Bay	58.5	53.4	2,192	71.1	1,649
Migori	69.2	60.2	2,181	70.8	1,854
Kisii	44.6	44.0	2,121	78.7	1,185
Nyamira	65.4	64.2	1,020	79.7	821
Nairobi City	29.4	9.2	12,318	62.0	1,827
Total	41.0	32.3	90,110	69.9	41,642

[1] An insecticide-treated net (ITN) is a factory-treated net that does not require any further treatment. In KDHS and KMIS surveys conducted prior to 2020, this was known as a long-lasting insecticidal net (LLIN).

Table 12.5 Use of existing ITNs

Percentage of insecticide-treated nets (ITNs) that were used by anyone the night before the survey, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Percentage of existing ITNs[1] used last night	Number of ITNs[1]
Residence		
Urban	77.7	5,672
Rural	78.3	15,368
Malaria burden		
No transmission	81.3	843
Very low burden	78.1	2,021
Low burden	74.4	2,798
Moderate burden	78.8	2,958
Moderate to high burden	79.7	3,867
High burden	78.7	3,459
Very high burden	77.6	5,093
Wealth quintile		
Lowest	80.0	3,937
Second	80.3	5,245
Middle	79.1	5,118
Fourth	75.7	4,001
Highest	73.0	2,738
Total	78.1	21,040

[1] An insecticide-treated net (ITN) is a factory-treated net that does not require any further treatment. In KDHS and KMIS surveys conducted prior to 2020, this was known as a long-lasting insecticidal net (LLIN).

Table 12.5C Use of existing ITNs by county

Percentage of insecticide-treated nets (ITNs) that were used by anyone the night before the survey, according to county, Kenya Mini DHS 2025–26

County	Percentage of existing ITNs[1] used last night	Number of ITNs[1]
Mombasa	78.7	793
Kwale	88.7	542
Kilifi	82.6	1,103
Tana River	91.8	233
Lamu	84.2	88
Taita Taveta	53.7	370
Garissa	82.7	21
Wajir	74.7	59
Mandera	90.3	46
Marsabit	43.1	69
Isiolo	88.4	46
Meru	83.4	337
Tharaka-Nithi	72.7	121
Embu	78.3	152
Kitui	72.8	301
Machakos	78.8	341
Makueni	76.6	285
Nyandarua	74.1	45
Nyeri	(82.8)	19
Kirinyaga	78.9	429
Murang'a	81.1	182
Kiambu	76.5	557
Turkana	74.3	522
West Pokot	66.3	369
Samburu	66.4	19
Trans Nzoia	79.5	751
Uasin Gishu	79.1	610
Elgeyo-Marakwet	78.3	112
Nandi	79.1	429
Baringo	73.7	253
Laikipia	79.4	75
Nakuru	73.0	351
Narok	76.4	654
Kajiado	80.9	333
Kericho	74.9	651
Bomet	80.2	593
Kakamega	74.4	1,332
Vihiga	78.0	517
Bungoma	76.6	1,294
Busia	78.6	796
Siaya	87.7	573
Kisumu	80.8	930
Homa Bay	84.6	732
Migori	71.9	945
Kisii	82.5	681
Nyamira	76.6	537
Nairobi City	81.3	843
Total	78.1	21,040

Note: Figures in parentheses are based on 25–49 unweighted cases.

[1] An insecticide-treated net (ITN) is a factory-treated net that does not require any further treatment. In KDHS and KMIS surveys conducted prior to 2020, this was known as a long-lasting insecticidal net (LLIN).

Table 12.6 Use of mosquito nets by children

Percentage of children under age 5 who slept under a mosquito net (treated or untreated) and under an insecticide-treated net (ITN) the night before the survey; and among children under age 5 in households with at least one ITN, percentage who slept under an ITN the night before the survey, according to background characteristics, Kenya Mini DHS 2025–26

	Children under age 5 in all households			Children under age 5 in households with at least one ITN[1]	
Background characteristic	Percentage who slept under any mosquito net last night	Percentage who slept under an ITN[1] last night	Number of children	Percentage who slept under an ITN[1] last night	Number of children
Age in months					
<12	62.1	44.2	2,004	83.3	1,063
12–23	55.4	45.0	2,009	82.9	1,090
24–35	53.9	42.9	2,151	80.1	1,151
36–47	48.9	39.5	2,394	76.3	1,239
48–59	46.7	37.9	2,323	75.3	1,168
Sex					
Male	52.7	41.4	5,444	80.0	2,821
Female	53.4	41.9	5,437	78.9	2,890
Residence					
Urban	51.1	30.3	4,400	80.2	1,664
Rural	54.4	49.4	6,481	79.1	4,047
Malaria burden					
No transmission	40.3	13.2	1,546	(76.9)	265
Very low burden	40.5	26.6	1,648	75.6	580
Low burden	39.9	27.2	2,478	75.6	891
Moderate burden	57.0	50.6	1,292	79.4	823
Moderate to high burden	63.0	60.1	1,192	82.1	872
High burden	73.6	69.4	1,227	82.6	1,032
Very high burden	73.7	66.7	1,497	80.0	1,248
Wealth quintile					
Lowest	47.9	43.9	2,500	76.5	1,434
Second	58.1	55.1	2,095	82.1	1,406
Middle	56.8	48.6	1,991	81.0	1,195
Fourth	54.7	37.7	2,239	81.6	1,034
Highest	48.7	23.1	2,055	73.7	643
Total	53.0	41.7	10,881	79.4	5,711

Note: Figures in parentheses are based on 25–49 unweighted cases.
Table is based on children who stayed in the household the night before the interview.
[1] An insecticide-treated net (ITN) is a factory-treated net that does not require any further treatment. In KDHS and KMIS surveys conducted prior to 2020, this was known as a long-lasting insecticidal net (LLIN).

Table 12.6C Use of mosquito nets by children by county

Percentage of children under age 5 who slept under a mosquito net (treated or untreated) and under an insecticide-treated net (ITN) the night before the survey; and among children under age 5 in households with at least one ITN, percentage who slept under an ITN the night before the survey, according to county, Kenya Mini DHS 2025–26

	Children under age 5 in all households			Children under age 5 in households with at least one ITN[1]	
County	Percentage who slept under any mosquito net last night	Percentage who slept under an ITN[1] last night	Number of children	Percentage who slept under an ITN[1] last night	Number of children
Mombasa	68.1	59.4	337	79.9	250
Kwale	70.3	63.4	200	80.0	159
Kilifi	74.7	71.5	290	88.0	236
Tana River	84.2	71.8	104	89.2	84
Lamu	58.6	57.4	31	84.9	21
Taita/Taveta	66.1	56.9	85	68.2	71
Garissa	66.6	5.1	126	(69.2)	9
Wajir	46.4	15.4	107	59.8	28
Mandera	45.5	10.8	134	54.7	27
Marsabit	27.2	12.3	102	33.3	37
Isiolo	50.7	30.9	47	73.8	19
Meru	41.2	34.1	269	88.5	103
Tharaka-Nithi	43.5	38.1	69	85.6	31
Embu	49.9	45.5	83	84.8	45
Kitui	32.2	30.8	205	74.1	85
Machakos	48.4	24.4	355	76.0	114
Makueni	47.3	38.3	170	68.8	95
Nyandarua	14.2	10.7	107	*	15
Nyeri	21.0	7.0	87	*	10
Kirinyaga	62.1	54.8	102	90.3	62
Murang'a	38.7	30.5	163	86.1	58
Kiambu	37.0	23.2	685	82.0	193
Turkana	88.0	81.3	184	88.0	170
West Pokot	60.7	58.3	201	73.9	159
Samburu	15.2	10.0	87	59.4	15
Trans Nzoia	78.5	73.4	176	87.9	147
Uasin Gishu	55.0	44.6	302	79.4	170
Elgeyo/Marakwet	35.4	35.3	102	73.6	49
Nandi	53.4	49.3	175	79.4	109
Baringo	42.2	40.3	127	70.1	73
Laikipia	18.0	13.5	104	(63.5)	22
Nakuru	24.2	14.6	577	68.2	124
Narok	47.4	46.8	253	71.0	167
Kajiado	48.8	40.4	286	87.0	133
Kericho	64.9	61.5	175	75.6	142
Bomet	53.3	49.7	233	68.1	170
Kakamega	70.2	63.2	370	74.7	313
Vihiga	72.3	68.7	124	77.2	110
Bungoma	75.5	72.9	379	83.3	332
Busia	76.1	73.1	196	83.9	171
Siaya	57.6	56.1	194	79.1	138
Kisumu	78.3	68.9	264	82.1	221
Homa Bay	73.1	69.2	263	85.6	212
Migori	81.9	70.2	349	83.1	294
Kisii	54.3	53.9	249	81.5	165
Nyamira	78.1	75.7	106	90.8	89
Nairobi City	40.3	13.2	1,546	(76.9)	265
Total	53.0	41.7	10,881	79.4	5,711

Note: Table is based on children who stayed in the household the night before the interview.
Figures in parentheses are based on 25–49 unweighted cases. An asterisk indicates that a figure is based on fewer than 25 unweighted cases and has been suppressed.
[1] An insecticide-treated net (ITN) is a factory-treated net that does not require any further treatment. In KDHS and KMIS surveys conducted prior to 2020, this was known as a long-lasting insecticidal net (LLIN).

Table 12.7 Use of mosquito nets by pregnant women

Percentage of pregnant women age 15–49 who slept under a mosquito net (treated or untreated) and under an insecticide-treated net (ITN) the night before the survey; and among pregnant women age 15–49 in households with at least one ITN, percentage who slept under an ITN the night before the survey, according to background characteristics, Kenya Mini DHS 2025–26

	Among pregnant women age 15–49 in all households			Among pregnant women age 15–49 in households with at least one ITN[1]	
Background characteristic	Percentage who slept under any mosquito net last night	Percentage who slept under an ITN[1] last night	Number of pregnant women	Percentage who slept under an ITN[1] last night	Number of pregnant women
Residence					
Urban	44.6	28.3	443	80.1	157
Rural	55.1	48.1	544	82.2	318
Malaria burden					
No transmission	(28.1)	(15.3)	127	*	30
Very low burden	39.6	24.3	147	74.6	48
Low burden	40.6	28.4	229	87.2	75
Moderate burden	55.6	48.3	143	84.5	82
Moderate to high burden	69.7	59.7	86	84.3	61
High burden	71.5	64.5	125	85.2	94
Very high burden	62.9	50.5	131	77.6	85
Education[2]					
No education	47.3	26.1	65	69.2	25
Primary	51.0	44.3	247	81.3	135
Secondary	49.5	39.7	492	84.1	233
More than secondary	53.2	35.7	183	78.3	84
Wealth quintile					
Lowest	48.0	42.2	205	78.6	110
Second	55.6	50.6	173	81.0	108
Middle	58.7	45.6	192	85.1	103
Fourth	50.4	35.9	224	90.5	89
Highest	40.0	23.4	194	(69.4)	66
Total	50.4	39.2	988	81.5	475

Note: Figures in parentheses are based on 25–49 unweighted cases. An asterisk indicates that a figure is based on fewer than 25 unweighted cases and has been suppressed.
Table is based on women who stayed in the household the night before the interview.
[1] An insecticide-treated net (ITN) is a factory-treated net that does not require any further treatment. In KDHS and KMIS surveys conducted prior to 2020, this was known as a long-lasting insecticidal net (LLIN).
[2] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level

Table 12.8 Main reason mosquito net was not used the night before the survey

Among ITNs, non-ITNs, and all mosquito nets, percentage that were not used by anyone the night before the survey, and among mosquito nets that were not used by anyone the night before the survey, percent distribution by the main reason each net was not used, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Percentage of nets not used the night before the survey	Total number of mosquito nets	Main reason each net was not used the night before the survey															Number of mosquito nets not used the night before the survey
			Too hot	Don't like net shape, colour, and/or size	Don't like smell	Unable to hang net	Slept outdoors	Usual user didn't sleep in household last night	No mosquitoes/ no malaria	Extra net/ saving for later	Net too small/ short	Net brought bedbugs	Net Too old/ torn/ damaged	Net Is too dirty	No place to hang net	Other	Total	
			ITNs[1]															
Residence																		
Urban	22.2	5,672	9.7	0.8	2.9	3.6	1.1	14.9	18.2	36.4	0.1	0.2	5.2	1.9	3.3	1.7	100.0	1,262
Rural	21.6	15,368	4.7	0.3	1.0	2.1	0.6	16.5	7.7	52.1	0.4	0.9	7.1	2.8	2.9	0.8	100.0	3,316
Malaria burden																		
No transmission	18.7	843	(4.7)	(0.0)	(7.2)	(4.8)	(0.0)	(18.4)	(49.7)	(5.8)	(0.0)	(0.0)	(7.3)	(0.0)	(0.0)	(2.1)	100.0	158
Very low burden	21.8	2,021	12.3	0.5	0.7	4.4	0.8	28.0	14.5	23.1	0.4	0.0	10.7	3.6	0.2	0.8	100.0	440
Low burden	25.5	2,798	7.0	0.8	2.6	2.4	1.2	12.6	26.1	31.3	0.3	0.0	6.1	4.8	3.2	1.7	100.0	713
Moderate burden	20.9	2,958	9.7	0.6	2.2	2.6	1.7	16.1	8.4	47.1	0.7	0.1	2.9	1.9	5.5	0.5	100.0	617
Moderate to high burden	20.2	3,867	3.8	0.7	1.6	1.2	0.3	17.6	4.4	55.6	0.4	0.1	7.4	2.0	2.9	2.0	100.0	782
High burden	21.3	3,459	5.2	0.2	0.9	2.3	1.0	9.2	7.9	59.2	0.0	1.0	4.9	1.8	5.7	0.7	100.0	736
Very high burden	22.2	5,093	3.3	0.4	0.3	2.6	0.2	16.7	1.1	61.1	0.2	2.2	7.8	2.3	1.2	0.6	100.0	1,132
Wealth quintile																		
Lowest	19.9	3,937	5.2	0.4	1.1	2.7	0.4	9.1	10.3	50.5	0.2	1.9	8.4	3.2	6.2	0.4	100.0	784
Second	19.6	5,245	5.3	0.3	0.6	1.7	0.5	12.6	6.3	55.9	0.7	1.1	8.8	2.4	2.8	1.2	100.0	1,027
Middle	20.8	5,118	4.8	0.6	1.6	2.4	0.7	16.8	7.5	51.8	0.2	0.6	6.8	2.6	2.2	1.2	100.0	1,065
Fourth	24.3	4,001	8.0	0.7	1.9	3.2	1.2	20.5	11.0	40.3	0.2	0.2	5.8	3.1	2.5	1.5	100.0	973
Highest	26.6	2,738	7.2	0.4	2.6	2.8	0.9	21.5	20.9	37.6	0.2	0.0	2.2	1.3	1.6	0.8	100.0	729
Total	21.8	21,040	6.0	0.5	1.5	2.5	0.8	16.1	10.6	47.8	0.3	0.7	6.6	2.5	3.0	1.1	100.0	4,578
			NON-ITNs															
Total	19.5	5,655	11.9	0.4	0.1	2.4	0.7	20.9	25.0	21.4	0.1	0.0	6.9	5.3	3.3	1.5	100.0	1,101
			ALL MOSQUITO NETS															
Total	21.3	26,695	7.2	0.5	1.2	2.5	0.7	17.0	13.4	42.7	0.3	0.6	6.7	3.1	3.0	1.2	100.0	5,679

Note: Figures in parentheses are based on 25–49 unweighted cases.

[1] An insecticide-treated net (ITN) is a factory-treated net that does not require any further treatment. In KDHS and KMIS surveys conducted prior to 2020, this was known as a long-lasting insecticidal net (LLIN).

Table 12.8C Main reason mosquito net was not used the night before the survey by county

Among ITNs, non-ITNs, and all mosquito nets, percentage that were not used by anyone the night before the survey, and among mosquito nets that were not used by anyone the night before the survey, percent distribution by the main reason each net was not used, according to county, Kenya Mini DHS 2025–26

County	Percentage of nets not used the night before the survey	Total number of mosquito nets	Main reason each net was not used the night before the survey															Number of mosquito nets not used the night before the survey
			Too hot	Don't like net shape, colour, and/or size	Don't like smell	Unable to hang net	Slept outdoors	Usual user didn't sleep in household last night	No mosquitoes/ no malaria	Extra net/ saving for later	Net too small/ short	Net brought bedbugs	Net too old/ torn/ damaged	Net is too dirty	No place to hang net	Other	Total	
ITNs[1]																		
Mombasa	21.3	793	26.7	1.6	3.4	0.0	1.3	11.7	6.8	42.2	0.0	0.0	2.5	2.3	1.6	0.0	100.0	169
Kwale	11.2	542	4.8	0.0	0.0	0.0	6.7	18.6	9.1	51.5	0.0	4.3	0.0	5.0	0.0	0.0	100.0	61
Kilifi	17.4	1,103	12.4	0.6	1.3	0.8	0.0	11.2	1.5	60.8	0.4	0.0	11.1	0.0	0.0	0.0	100.0	191
Tana River	8.2	233	0.6	0.0	0.0	0.0	6.0	27.7	11.7	48.8	0.0	0.0	0.0	5.2	0.0	0.0	100.0	19
Lamu	15.8	88	26.2	0.0	0.0	2.9	0.0	20.1	2.9	25.7	0.0	0.0	17.8	1.8	0.8	1.8	100.0	14
Taita/Taveta	46.2	370	14.3	1.1	0.0	0.8	0.0	10.7	12.5	48.8	0.9	0.0	1.3	7.0	2.3	0.4	100.0	171
Garissa	17.3	21	*	*	*	*	*	*	*	*	*	*	*	*	*	*	100.0	4
Wajir	25.3	59	8.2	0.0	0.0	0.0	0.0	2.5	44.0	0.0	0.0	0.0	35.5	9.9	0.0	0.0	100.0	15
Mandera	9.7	46	*	*	*	*	*	*	*	*	*	*	*	*	*	*	100.0	4
Marsabit	56.9	69	4.1	0.0	0.0	0.8	0.0	2.9	72.8	5.8	0.0	0.0	4.9	7.0	1.8	0.0	100.0	39
Isiolo	11.6	46	(0.0)	(0.0)	(5.5)	(0.0)	(7.8)	(11.3)	(9.7)	(28.3)	(0.0)	(0.0)	(23.5)	(13.9)	(0.0)	(0.0)	100.0	5
Meru	16.6	337	(1.3)	(0.9)	(2.2)	(0.0)	(6.5)	(19.5)	(22.7)	(27.9)	(0.0)	(0.0)	(8.3)	(10.8)	(0.0)	(0.0)	100.0	56
Tharaka-Nithi	27.3	121	9.8	0.0	1.8	3.5	0.0	31.1	25.6	14.1	1.6	0.0	5.9	3.9	0.0	2.7	100.0	33
Embu	21.7	152	15.8	0.0	0.0	2.6	0.0	21.5	25.3	18.4	0.0	0.0	11.7	1.6	2.4	0.5	100.0	33
Kitui	27.2	301	26.6	0.0	0.0	0.0	0.0	32.6	8.3	15.8	0.0	0.0	16.7	0.0	0.0	0.0	100.0	82
Machakos	21.2	341	(8.9)	(0.0)	(0.0)	(21.0)	(2.9)	(26.1)	(8.0)	(21.9)	(1.9)	(0.0)	(7.7)	(1.5)	(0.0)	(0.0)	100.0	72
Makueni	23.4	285	18.9	0.0	0.0	1.3	0.0	30.4	15.5	22.2	0.0	0.0	5.1	6.6	0.0	0.0	100.0	67
Nyandarua	23.5	45	*	*	*	*	*	*	*	*	*	*	*	*	*	*	100.0	11
Nyeri	(17.2)	19	*	*	*	*	*	*	*	*	*	*	*	*	*	*	100.0	3
Kirinyaga	21.1	429	0.7	1.0	3.4	0.0	1.0	44.4	1.7	43.2	0.0	0.0	3.1	0.6	0.0	0.8	100.0	91
Murang'a	18.4	182	(3.4)	(2.1)	(0.0)	(2.6)	(0.0)	(7.0)	(35.1)	(21.0)	(0.0)	(0.0)	(13.3)	(14.5)	(0.0)	(1.1)	100.0	33
Kiambu	23.5	557	7.1	0.0	0.0	0.0	0.0	8.4	51.8	10.1	0.0	0.0	8.4	3.6	4.5	6.1	100.0	131
Turkana	25.7	522	0.7	0.0	0.0	0.0	2.6	10.4	3.7	81.2	0.0	0.0	1.4	0.0	0.0	0.0	100.0	134
West Pokot	33.7	369	11.0	0.7	1.9	2.8	0.0	7.3	22.6	27.2	0.0	0.8	5.7	2.3	17.1	0.6	100.0	125
Samburu	33.6	19	(31.8)	(0.0)	(2.2)	(5.3)	(0.0)	(7.6)	(33.0)	(5.9)	(0.0)	(0.0)	(11.9)	(2.2)	(0.0)	(0.0)	100.0	6
Trans Nzoia	20.5	751	2.1	0.4	2.6	1.9	0.0	20.3	0.8	60.6	0.0	0.0	3.8	1.7	5.4	0.3	100.0	154
Uasin Gishu	19.5	610	0.0	0.8	6.1	4.6	2.0	25.7	15.6	37.5	0.0	0.0	2.3	3.4	1.9	0.0	100.0	119
Elgeyo/ Marakwet	21.7	112	8.9	0.0	0.4	13.4	1.1	9.3	21.3	29.7	0.0	1.9	8.2	1.6	0.0	4.1	100.0	24
Nandi	20.9	429	2.4	1.0	3.1	0.9	0.3	7.6	4.6	59.1	2.8	0.0	8.7	2.7	6.7	0.0	100.0	89
Baringo	25.6	253	0.0	0.0	1.5	1.1	0.0	29.1	13.9	37.2	0.0	1.3	4.8	1.6	3.9	5.5	100.0	65
Laikipia	20.6	75	(0.0)	(3.5)	(0.0)	(0.0)	(0.0)	(23.8)	(36.8)	(14.3)	(2.3)	(0.0)	(12.6)	(6.6)	(0.0)	(0.0)	100.0	15
Nakuru	27.0	351	(3.9)	(2.3)	(13.6)	(12.9)	(2.0)	(14.2)	(20.6)	(19.3)	(0.0)	(0.0)	(0.0)	(1.8)	(8.1)	(1.4)	100.0	95
Narok	23.6	654	3.5	0.0	0.0	1.2	0.0	4.0	7.3	65.9	1.7	0.0	1.4	0.0	14.6	0.4	100.0	154
Kajiado	18.7	333	(6.9)	(1.3)	(0.0)	(0.0)	(4.1)	(14.2)	(13.3)	(23.5)	(0.0)	(0.0)	(20.1)	(8.3)	(6.6)	(1.7)	100.0	62
Kericho	25.1	651	0.5	0.0	0.7	2.1	0.0	19.7	7.4	48.1	0.0	0.0	8.5	2.4	3.7	6.9	100.0	163
Bomet	19.6	593	0.8	0.0	2.9	1.6	0.0	12.2	16.3	58.7	0.0	0.0	6.8	0.0	0.6	0.0	100.0	116
Kakamega	25.6	1,332	2.5	0.4	1.0	2.6	0.0	15.6	0.3	55.8	0.0	3.4	15.6	2.0	0.4	0.4	100.0	341
Vihiga	22.0	517	3.4	0.9	0.4	1.5	0.0	15.4	1.1	69.2	0.0	0.4	2.9	3.5	0.5	0.8	100.0	114
Bungoma	23.4	1,294	4.2	0.2	0.4	2.1	0.0	8.6	6.4	63.6	0.0	0.7	4.4	2.4	5.9	1.2	100.0	303
Busia	20.6	796	5.0	0.5	0.0	0.0	0.0	21.9	2.8	54.7	0.0	6.1	6.5	1.5	0.5	0.5	100.0	164
Siaya	12.2	573	0.5	0.0	0.0	3.4	3.9	14.8	1.5	64.8	1.1	0.0	4.6	0.0	4.1	1.3	100.0	70
Kisumu	19.2	930	8.2	0.0	0.0	5.9	0.0	19.3	1.5	57.0	0.0	0.8	2.9	2.9	1.3	0.2	100.0	179
Homa Bay	15.4	732	7.2	0.0	2.6	6.4	0.0	6.6	0.0	60.3	0.0	1.5	11.9	0.0	2.6	0.9	100.0	113
Migori	28.1	945	0.8	0.4	0.0	2.0	0.0	14.3	0.7	70.4	0.7	0.4	4.7	2.9	2.0	0.8	100.0	265
Kisii	17.5	681	0.0	2.1	1.0	0.0	1.8	23.2	4.1	58.3	0.0	0.0	4.6	4.9	0.0	0.0	100.0	119
Nyamira	23.4	537	4.3	0.0	0.4	4.0	3.4	27.6	0.9	45.0	1.5	0.0	4.9	1.8	5.2	1.0	100.0	126
Nairobi City	18.7	843	(4.7)	(0.0)	(7.2)	(4.8)	(0.0)	(18.4)	(49.7)	(5.8)	(0.0)	(0.0)	(7.3)	(0.0)	(0.0)	(2.1)	100.0	158
Total	21.8	21,040	6.0	0.5	1.5	2.5	0.8	16.1	10.6	47.8	0.3	0.7	6.6	2.5	3.0	1.1	100.0	4,578
NON-ITNs																		
Total	19.5	5,655	11.9	0.4	0.1	2.4	0.7	20.9	25.0	21.4	0.1	0.0	6.9	5.3	3.3	1.5	100.0	1,101
ALL MOSQUITO NETS																		
Total	21.3	26,695	7.2	0.5	1.2	2.5	0.t7	17.0	13.4	42.7	0.3	0.6	6.7	3.1	3.0	1.2	100.0	5,679

Note: Figures in parentheses are based on 25–49 unweighted cases. An asterisk indicates that a figure is based on fewer than 25 unweighted cases and has been suppressed.
[1] An insecticide-treated net (ITN) is a factory-treated net that does not require any further treatment. In KDHS and KMIS surveys conducted prior to 2020, this was known as a long-lasting insecticidal net (LLIN).

Table 12.9 Use of Intermittent Preventive Treatment (IPTp) by women during pregnancy

Percentage of women age 15–49 with a live birth and/or a stillbirth in the 2 years preceding the survey who, during the pregnancy that resulted in the last live birth or stillbirth, received one or more doses of SP/Fansidar, received two or more doses of SP/Fansidar, and received three or more doses of SP/Fansidar, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Percentage who received one or more doses of SP/Fansidar	Percentage who received two or more doses of SP/Fansidar	Percentage who received three or more doses of SP/Fansidar	Number of women with a live birth and/or a stillbirth in the 2 years preceding the survey
LIVE BIRTHS				
Birth order[1]				
1	28.1	21.5	14.6	1,269
2–3	27.6	20.1	12.0	1,724
4–5	31.4	22.4	15.3	685
6+	33.2	26.0	18.2	304
Residence				
Urban	24.8	18.5	10.4	1,698
Rural	31.8	23.6	16.5	2,285
Malaria burden				
No transmission	16.0	9.5	3.9	577
Very low burden	7.5	3.5	2.2	627
Low burden	7.0	4.3	2.0	919
Moderate burden	28.6	21.7	11.2	480
Moderate to high burden	29.8	23.5	17.4	439
High burden	60.3	46.3	32.2	415
Very high burden	81.0	63.8	44.5	527
IPT Implementation				
Areas where IPTp is implemented	76.5	60.7	41.4	1,178
Areas where IPTp is not implemented	8.8	4.9	2.3	2,805
Education[2]				
No education	17.1	11.1	6.8	350
Primary	33.3	25.3	17.0	1,076
Secondary	30.3	22.2	14.1	1,756
More than secondary	24.9	19.0	12.2	800
Wealth quintile				
Lowest	30.0	21.8	14.9	836
Second	39.9	29.8	19.6	745
Middle	31.6	25.1	18.0	720
Fourth	23.9	17.3	12.0	914
Highest	20.2	14.4	5.6	768
Total	28.8	21.4	13.9	3,983
STILLBIRTHS				
Total	17.9	17.9	6.3	38
LIVE BIRTHS AND STILLBIRTHS[3]				
Total	28.7	21.4	13.8	4,021

Note: Stillbirths are foetal deaths in pregnancies lasting 28 or more weeks. When pregnancy duration is reported in months, stillbirths are foetal deaths in pregnancies lasting 7 or more months.
[1] Birth order refers to the order of the births among the respondent's live births.
[2] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level
[3] For women who had both a live birth and a stillbirth in the 2 years preceding the survey, data are tabulated for the most recent birth only.

Table 12.9C Use of Intermittent Preventive Treatment (IPTp) by women during pregnancy by county

Percentage of women age 15–49 with a live birth and/or a stillbirth in the 2 years preceding the survey who, during the pregnancy that resulted in the last live birth or stillbirth, received one or more doses of SP/Fansidar, received two or more doses of SP/Fansidar, and received three or more doses of SP/Fansidar, according to county, Kenya Mini DHS 2025–26

County	Percentage who received one or more doses of SP/Fansidar	Percentage who received two or more doses of SP/Fansidar	Percentage who received three or more doses of SP/Fansidar	Number of women with a live birth and/or a stillbirth in the 2 years preceding the survey
LIVE BIRTHS				
Mombasa	72.1	65.4	32.8	127
Kwale	84.5	75.3	51.6	71
Kilifi	92.5	79.6	63.7	102
Tana River	41.9	14.2	4.6	40
Lamu	54.7	41.4	25.4	10
Taita/Taveta	78.1	55.6	34.0	30
Garissa	8.1	2.3	0.8	42
Wajir	0.0	0.0	0.0	36
Mandera	2.8	1.6	0.4	39
Marsabit	4.1	4.1	3.0	35
Isiolo	4.0	1.7	1.7	16
Meru	0.0	0.0	0.0	72
Tharaka-Nithi	8.2	2.7	0.0	22
Embu	11.7	6.5	4.0	27
Kitui	15.1	8.5	5.6	64
Machakos	9.5	3.3	2.2	156
Makueni	8.1	3.0	1.6	59
Nyandarua	4.8	0.0	0.0	48
Nyeri	2.8	0.0	0.0	42
Kirinyaga	8.7	7.2	5.7	37
Murang'a	0.0	0.0	0.0	59
Kiambu	4.0	3.1	2.0	269
Turkana	38.4	9.8	2.0	62
West Pokot	10.8	6.6	2.1	73
Samburu	4.8	1.0	1.0	31
Trans Nzoia	7.5	3.0	0.0	57
Uasin Gishu	7.3	3.7	3.0	123
Elgeyo/Marakwet	9.5	8.1	6.1	32
Nandi	20.9	15.1	9.4	75
Baringo	8.5	2.6	0.0	53
Laikipia	2.8	1.2	0.0	50
Nakuru	2.9	0.0	0.0	222
Narok	3.9	2.1	2.1	87
Kajiado	12.8	9.4	0.9	123
Kericho	9.7	9.1	4.4	68
Bomet	1.3	0.2	0.1	89
Kakamega	84.3	67.7	41.9	126
Vihiga	74.5	67.0	62.3	38
Bungoma	66.8	51.8	39.0	128
Busia	72.0	58.2	46.9	78
Siaya	67.1	57.0	37.0	53
Kisumu	92.9	68.2	50.0	101
Homa Bay	90.8	76.1	54.9	81
Migori	81.4	61.8	39.3	131
Kisii	5.8	1.2	1.2	84
Nyamira	29.6	14.3	5.9	40
Nairobi City	16.0	9.5	3.9	577
Total	28.8	21.4	13.9	3,983
STILLBIRTHS				
Total	17.9	17.9	6.3	38
LIVE BIRTHS AND STILLBIRTHS[1]				
Total	28.7	21.4	13.8	4,021

[1] For women who had both a live birth and a stillbirth in the 2 years preceding the survey, data are tabulated for the most recent birth only.

Table 12.10 Children with fever and careseeking, prompt treatment, and diagnosis

Percentage of children under age 5 with fever in the 2 weeks preceding the survey; and among children under age 5 with fever, percentage for whom advice or treatment was sought, percentage for whom advice or treatment was sought the same or next day following the onset of fever, percentage who had blood taken from a finger or heel for testing, and percentage who were diagnosed with malaria by a healthcare provider, according to background characteristics, Kenya Mini DHS 2025–26

	Children under age 5		Children under age 5 with fever				
Background characteristic	Percentage with fever in the 2 weeks preceding the survey	Number of children	Percentage for whom advice or treatment was sought[1]	Percentage for whom advice or treatment was sought the same or next day	Percentage who had blood taken from a finger or heel for testing	Percentage who were diagnosed with malaria by a healthcare provider	Number of children
Age in months							
<12	13.5	2,041	60.8	42.9	18.5	7.2	276
12–23	17.3	1,934	71.6	47.7	31.6	15.9	335
24–35	15.2	2,007	72.3	54.1	32.2	17.5	304
36–47	14.2	2,136	62.7	29.1	29.6	18.1	303
48–59	9.0	2,048	72.3	46.5	37.4	26.6	185
Sex							
Male	13.7	5,081	67.4	44.1	30.3	18.6	697
Female	13.9	5,086	68.1	43.9	28.7	14.3	706
Residence							
Urban	14.6	4,444	70.4	47.0	27.9	8.3	649
Rural	13.2	5,722	65.5	41.4	30.9	23.4	754
Malaria burden							
No transmission	18.6	1,600	(69.0)	(47.0)	(22.6)	(2.1)	297
Very low burden	13.3	1,497	69.3	49.5	20.2	9.6	199
Low burden	9.1	2,375	64.1	41.8	22.2	7.0	216
Moderate burden	10.9	1,222	67.7	39.9	24.1	6.8	133
Moderate to high burden	10.8	1,075	76.6	48.0	33.7	18.4	116
High burden	17.4	1,077	62.4	34.6	53.6	43.0	188
Very high burden	19.2	1,318	68.2	45.3	34.2	31.3	253
Mother's education[2]							
No education	10.7	952	62.8	37.9	34.8	28.2	102
Primary	14.7	3,106	66.1	40.9	32.5	20.2	458
Secondary	13.6	4,055	66.8	43.6	25.3	15.4	550
More than secondary	14.3	2,053	73.9	51.6	30.9	8.5	293
Wealth quintile							
Lowest	12.3	2,185	62.0	33.0	29.3	22.0	269
Second	15.0	1,875	65.1	39.0	31.4	26.2	281
Middle	13.1	1,819	66.7	50.4	27.0	19.3	238
Fourth	13.5	2,271	73.3	46.0	28.2	13.7	307
Highest	15.3	2,016	70.6	51.2	31.0	3.2	308
Total	13.8	10,166	67.8	44.0	29.5	16.4	1,403

Note: Figures in parentheses are based on 25–49 unweighted cases.
[1] Includes advice or treatment from the following sources: public medical sector, private medical sector, NGO medical sector, FBO medical sector, shop, market, and itinerant drug seller. Excludes advice or treatment from a traditional practitioner.
[2] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level

Table 12.11 Source of advice or treatment for children with fever

Percentage of children under age 5 with fever in the 2 weeks preceding the survey for whom advice or treatment was sought from specific sources; and among children under age 5 with fever in the 2 weeks preceding the survey for whom advice or treatment was sought, percentage for whom advice or treatment was sought from specific sources, Kenya Mini DHS 2025–26

	Percentage for whom advice or treatment was sought from each source:	
Source	Among children with fever	Among children with fever for whom advice or treatment was sought
Public sector	**34.6**	**51.0**
Government hospital	13.7	20.2
Government health centre	6.1	9.0
Government dispensary	14.8	21.8
Mobile clinic	0.1	0.1
CHW/Fieldworker	0.3	0.4
Private medical sector (non-NGO)	**33.0**	**48.6**
Private hospital	12.7	18.7
Private clinic	4.9	7.2
Pharmacy	14.8	21.9
Private doctor	0.2	0.4
CHW/Fieldworker	0.6	0.8
Private medical sector (FBO)	**0.4**	**0.6**
FBO/Mission hospital	0.2	0.3
FBO/Mission clinic	0.1	0.1
FBO/Mobile clinic	0.1	0.2
Other private sector	**0.5**	**0.8**
Shop	0.4	0.6
Traditional practitioner	0.1	0.1
Itinerant drug seller	0.1	0.1
Other	0.3	0.4
Number of children	1,403	952

Note: Advice or treatment for children with fever may have been sought from more than one source.
CHW = Community health worker
NGO = Non-governmental organisation
FBO = Faith-based organisation

Table 12.12 Type of antimalarial drugs used

Among children under age 5 with fever in the 2 weeks preceding the survey who took any antimalarial medication, percentage who took specific antimalarial drugs, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Any ACT	Specific ACT			Percentage of children who took:								Number of children with fever who took anti-malarial drug
		AL	DHAP	Other	SP/ Fansidar	Chloro-quine	Amodia-quine	Quinine pills	Quinine injection	Arte-sunate rectal	Arte-sunate injection	Other anti-malarial	
Age in months													
<6	*	*	*	*	*	*	*	*	*	*	*	*	3
6–11	*	*	*	*	*	*	*	*	*	*	*	*	26
12–23	77.3	65.6	11.7	2.1	7.4	13.5	2.2	0.0	6.9	1.4	7.6	2.7	52
24–35	74.9	57.7	5.1	12.0	10.0	2.9	1.2	2.1	8.4	1.5	3.5	0.0	53
36–47	79.4	72.1	6.9	6.0	12.9	15.6	7.6	2.2	2.6	0.8	0.0	0.8	63
48–59	78.3	73.0	2.8	2.6	3.6	5.9	8.7	0.0	3.6	0.0	4.1	0.0	49
Sex													
Male	76.0	66.3	9.8	2.2	10.8	11.2	4.9	0.0	5.5	0.6	3.2	1.5	129
Female	72.1	62.3	2.9	8.3	6.7	17.0	3.9	2.2	3.7	1.1	3.1	0.0	116
Residence													
Urban	68.3	56.4	6.5	10.8	5.7	26.0	4.9	1.4	4.4	1.1	0.8	0.0	65
Rural	76.3	67.3	6.6	3.0	10.0	9.6	4.2	0.9	4.8	0.7	4.0	1.0	180
Malaria burden													
No transmission	*	*	*	*	*	*	*	*	*	*	*	*	22
Very low burden	(53.5)	(38.7)	(14.9)	(0.0)	(7.0)	(21.3)	(7.6)	(0.8)	(3.8)	(5.6)	(0.6)	(2.9)	27
Low burden	*	*	*	*	*	*	*	*	*	*	*	*	17
Moderate burden	*	*	*	*	*	*	*	*	*	*	*	*	7
Moderate to high burden	*	*	*	*	*	*	*	*	*	*	*	*	18
High burden	81.7	73.2	5.8	2.7	9.7	1.9	3.6	2.2	4.1	0.0	2.4	0.0	66
Very high burden	89.0	81.9	4.0	4.4	4.8	4.0	3.1	1.0	8.2	0.0	4.9	0.0	89
Mother's education[1]													
No education	(69.5)	(68.2)	(0.0)	(1.3)	(3.3)	(9.5)	(5.2)	(3.9)	(10.6)	(0.0)	(6.4)	(0.0)	20
Primary	75.4	63.0	6.1	10.3	11.9	13.5	6.2	0.8	6.4	0.0	4.4	0.6	104
Secondary	69.9	60.3	8.4	1.2	7.5	18.2	1.9	0.0	2.9	2.2	1.1	0.9	92
More than secondary	*	*	*	*	*	*	*	*	*	*	*	*	30
Wealth quintile													
Lowest	74.4	65.2	4.2	5.0	7.9	8.9	1.6	2.7	2.8	0.8	3.9	0.0	61
Second	81.8	66.6	3.8	11.4	11.1	5.2	3.0	0.0	6.1	0.0	4.3	0.8	79
Middle	(74.5)	(70.0)	(7.3)	(0.0)	(12.7)	(16.5)	(2.8)	(0.0)	(3.3)	(2.0)	(5.1)	(0.0)	40
Fourth	(76.8)	(66.5)	(17.4)	(0.0)	(1.2)	(20.7)	(14.6)	(0.0)	(5.8)	(0.0)	(0.0)	(1.8)	43
Highest	*	*	*	*	*	*	*	*	*	*	*	*	22
Total	74.1	64.4	6.5	5.1	8.9	13.9	4.4	1.0	4.6	0.8	3.2	0.8	245

Note: Figures in parentheses are based on 25–49 unweighted cases. An asterisk indicates that a figure is based on fewer than 25 unweighted cases and has been suppressed.
ACT = Artemisinin-based combination therapy
AL = Artemether-lumefantrine
DHAP = Dihydroartemisinin-piperaquine
SP = Sulfadoxine pyrimethamine
[1] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level

HOUSEHOLD WATER AND SANITATION 13

Key Findings

- ***Drinking water sources:*** Nationally, 85% of households have an improved source of drinking water, with higher coverage in urban areas (98%) than in rural areas (73%).
- ***Drinking water service ladder:*** 73% of the population has access to at least basic drinking water services.
- ***Sanitation:*** 76% of households have access to an improved sanitation facility.
- ***Sanitation service ladder:*** 45% of the population has access to at least basic sanitation services, 28% have limited services, and 7% practice open defecation.
- ***Management of household excreta:*** 71% of the household population has access to facilities that are connected to a sewer, has excreta safely disposed of on site, or has excreta removed for treatment off site.
- ***Handwashing:*** Nationally, 53% of the population has a basic handwashing facility (60% in urban areas and 49% in rural areas).
- ***Menstrual health and hygiene:*** 97% of women were able to wash and change in privacy and used appropriate materials to collect or absorb menstrual blood during their last period.

The extent to which households can access and use safe drinking water and sanitation facilities and can engage in hygienic practices profoundly affects the health, safety, and overall well-being of the population. This chapter presents information on sources of drinking water, types of sanitation facilities, disposal of excreta (including disposal of young children's stools), handwashing, and menstrual health and hygiene.

13.1 DRINKING WATER SOURCES, AVAILABILITY, AND TREATMENT

Improved sources of drinking water

Include piped water, public taps, standpipes, tube wells, boreholes, protected dug wells and springs, rainwater, water delivered via a tanker truck or a cart with a small tank, and bottled water.

Sample: Households and de jure population

Households that use unimproved sources of drinking water are at a higher risk of waterborne diseases and contamination. Therefore, increasing access to improved sources of drinking water is crucial for improving public health.

Results of the 2025–26 Kenya Mini Demographic and Health Survey (KMDHS) show that nationally, 85% of households have an improved source of drinking water. The percentage of households with access to an improved source of drinking water is higher in urban areas (98%) than in rural areas (73%) (**Table 13.1**).


The main improved sources of drinking water for urban households are piped water in the dwelling, yard, or plot (41%); bottled water (19%); and public tap or standpipe (14%). Water piped into the dwelling, yard, or plot (16%) is also the most common source in rural households, followed by a tube well or borehole (14%), protected spring water (11%), and rainwater (9%). Twenty percent of rural households, compared with only 1% of urban households, rely on unprotected surface water.

Nationally, 57% of households have water on their premises; the percentage is much higher in urban areas (78%) than in rural areas (40%). Twelve percent of households (2% in urban areas and 20% in rural areas) spend more than 30 minutes (round trip) to obtain drinking water.

Trends: The percentage of households with improved sources of drinking water increased from 45% in the 2003 Kenya Demographic and Health Survey (KDHS) to 85% in the 2025–26 KMDHS. The change was more pronounced in rural areas than in urban areas, increasing by 39 percentage points in rural areas (from 34% in 2003 to 73% in 2025–26) and by 19 percentage points in urban areas (from 79% in 2003 to 98% in 2025–26) (**Figure 13.1**).

Figure 13.1 **Household with improved source of drinking water service by residence**

Percentage of households with an improved source of drinking water by residence



Drinking Water Service Ladder

Drinking water service ladder

Safely managed

Drinking water from an improved water source that is located on the premises, available when needed, and free from faecal and priority chemical contamination.

Basic

Drinking water from an improved source, provided either water is on the premises or round-trip collection time is 30 minutes or less.

Limited

Drinking water from an improved source, and round-trip collection time is more than 30 minutes.

Unimproved

Drinking water from an unprotected dug well or unprotected spring.

Surface water

Drinking water directly from a river, dam, lake, pond, stream, canal, or irrigation canal.


Sample: De jure population

Building off the classification of drinking water sources as improved or unimproved, the Joint Monitoring Programme for Water Supply, Sanitation and Hygiene (JMP) has devised a five-rung drinking water service ladder to benchmark and compare progress towards achieving Sustainable Development Goal (SDG) targets (WHO/UNICEF JMP 2018). The 2025–26 KMDHS captured information on four out of the five rungs; because drinking water was not tested for faecal or chemical contamination during the survey, safely managed and basic drinking water services cannot be distinguished and are grouped together in **Tables 13.2** and **13.2C** as 'at least basic service'.

Nationally, 73% of the household population has access to at least basic drinking water services, 8% have limited services, and 5% use unimproved sources. The remaining 14% use surface water for drinking (**Figure 13.2**).

Figure 13.2 Household population Drinking water service by residence

Percent distribution of de jure population by drinking water service ladder




Patterns by background characteristics

- A higher percentage of the urban population (95%) than of the rural population (59%) has access to at least basic drinking water services. Twenty-two percent of households in rural areas use surface water for drinking (**Table 13.2**).
- The percentage of the population with access to at least basic drinking water services increases with household wealth, from 40% in the lowest wealth quintile to 98% in the highest wealth quintile.
- By county, the percentage of the population with at least basic service is lowest in Narok (26%), followed by Kitui (31%) and Samburu (33%). The percentages of the population with at least basic service are highest in Nairobi City, Mombasa, and Kiambu (97% each), followed by Uasin Gishu (90%), Nyandarua (89%), and Nyeri (89%) (**Table 13.2C** and **Map 13.1**).

Map 13.1 At least basic service for drinking water by county

Percentage of household population with at least basic service for drinking water



13.2 SANITATION

Improved sanitation facilities

Include flush/pour flush toilets that flush water and waste to a piped sewer system, septic tank, pit latrine, or unknown destination; ventilated improved pit (VIP) latrines; pit latrines with slabs; and composting toilets.


Sample: Households and de jure population

Three-quarters of households (76%) have access to an improved sanitation facility (**Table 13.3**). The percentage of households with improved sanitation facilities is higher in urban areas (93%) than in rural areas (63%). Nevertheless, open defecation is still practiced by 5% of households, including 9% of rural households and less than 1% of urban households.

Trends: The percentage of households with unimproved sanitation facilities decreased from 64% in the 2003 KDHS to 19% in the 2025–26 KMDHS. During the same period, the percentage of households with improved sanitation facilities increased from 19% to 76% (**Figure 13.3**).

***Figure 13.3* Household with improved source of drinking water service**

Percentage of households with improved and unimproved sanitation facilities



Patterns by background characteristics

- Among households with access to improved sanitation facilities, only 18% have a flush/pour flush toilet that flushes water to a piped sewer system. However, a flush pour toilet that flushes to a piped sewer system is the most common sanitation facility in urban areas (38%), followed by a pit latrine with a slab (21%) and a flush/pour flush toilet that flushes water to a septic tank (16%). In rural areas, the most common type of improved sanitation facility is a pit latrine with a slab (44%), followed by a ventilated improved pit latrine (13%).

- Sanitation facilities are located within the dwelling in 23% of households, including 38% of urban households and 9% of rural households. Almost three-quarters of households (72%) have toilet facilities within their yards/plots.

13.2.1 Sanitation Service Ladder

Sanitation service ladder

Safely managed

Use of improved facilities that are not shared with other households and where excreta are safely disposed of in situ or transported and treated off-site.

Basic

Use of improved facilities that are not shared with other households.

Limited

Use of improved facilities shared by two or more households.

Unimproved

Use of pit latrines without a slab or platform, hanging latrines, or bucket latrines.

Open defecation

Disposal of human faeces in fields, forests, bushes, open bodies of water, beaches, or other open spaces or with solid waste.

Sample: De jure population

The JMP has also devised a five-rung sanitation service ladder to benchmark and compare progress towards achieving SDG targets related to sanitation. The 2025–26 KMDHS captured information about all five rungs. However, for those households whose excreta are taken off-site, it is not possible to know if they are treated appropriately; therefore, safely managed and basic sanitation services are grouped together in **Table 13.4** as 'at least basic service'.

Overall, 45% of the population has access to at least basic sanitation services. Twenty-eight percent have limited services, and 7% practice open defecation (**Figure 13.4**).


Trends: The percentage of households with basic sanitation services increased from 10% in the 2003 KDHS to 45% in the 2025–26 KMDHS.

Patterns by background characteristics

- The percentage of the population with at least basic sanitation services is higher in urban areas (51%) than in rural areas (41%). Open defecation is still practiced in rural areas (11%) but very rarely in urban areas (<1%) (**Table 13.4**).

Figure 13.4 **Household population sanitation service by residence**

Percent distribution of de jure population by sanitation service ladder



- Access to at least basic sanitation services is least common in the lowest wealth quintile (16%) and most common in the highest wealth quintile (81%).
- By county, the highest percentage of the population with at least basic sanitation services is in Makueni (71%), followed by Lamu (68%) and Kericho (62%). The lowest percentage is in Samburu (14%), followed by Bomet and Turkana (15% each) (**Table 13.4C**).
- The highest percentage of the population practicing open defecation is in Turkana County (71%), followed by Samburu (58%), Wajir (39%), and Kwale (35%) Counties.

13.2.2 Removal and Disposal of Excreta

Disposal of excreta from on-site facilities

Excreta safely disposed of in situ

Includes septic tanks and latrines in which waste is buried in a covered pit, waste is never emptied, and it is unknown if waste is ever emptied.

Excreta disposed of unsafely

Includes septic tanks and latrines in which waste is emptied to uncovered pits, open ground, water bodies, or other locations.

Excreta removed for treatment

Includes septic tanks and latrines in which waste is removed by a service provider to a treatment plant or unknown location or is removed by a non-service provider to an unknown location.

Appropriate management of excreta

Household is connected to a sewer, has excreta safely disposed of on-site, or has excreta removed for treatment off-site.

Sample: De jure population with on-site sanitation facilities (septic tanks, pit latrines, and composting toilets)

Information on the disposal of excreta from sanitation facilities that are not connected to a sewer system is essential for assessing the percentage of the population using safely managed sanitation services.


Among the household population with improved on-site sanitation facilities, 87% have never emptied their septic tank or latrine, 3% have had the waste removed by a service provider but do not know the removal location, 2% have had their waste removed by a service provider to a treatment plant, and 1% have buried the waste in a covered pit. Overall, 94% of the household population with on-site sanitation facilities have had excreta safely disposed of in situ, 5% have had excreta removed for treatment, and less than 1% dispose of excreta unsafely (**Tables 13.5** and **13.5C**).

Tables 13.6 and **13.6C** shows the percent distribution of the household population by management of household excreta. Overall, 71% of the household population appropriately manages excreta. Households with appropriate management include those with facilities connected to a sewer (15%), those for which excreta are safely disposed of on-site (52%), and those for which excreta is removed for treatment off-site (3%). However, 28% of the population lacks appropriate management of household excreta (**Figure 13.5**).

Trends: The percentage of the household population with appropriate management of excreta increased from 66% in the 2022 KDHS to 71% in the 2025–26 KMDHS.

Figure 13.5 **Appropriate management of household excreta**

Percent distribution of household population by management of household excreta




Patterns by background characteristics

- Appropriate management of excreta is less common among the rural population (59%) than among the urban population (88%) (**Table 13.6**).
- The percentage of the population that appropriately manages excreta increases with wealth, from 31% in the lowest wealth quintile to 93% in the highest wealth quintile (**Figure 13.6**).

Figure 13.6 **Appropriate management of household excreta by wealth**

Percentage of household population connected to sewer or using improved on-site sanitation facilities with safe disposal of excreta on site or removal of excreta for off-site treatment



13.3 Disposal of Children's Stools

Appropriate disposal of children's stools

The child's last stools were put or rinsed into a toilet or latrine, or the child used a toilet or latrine.

Sample: Youngest children under age 2 living with their mother

Fifty percent of children under age 2 living with their mother have their last stools disposed of appropriately (**Tables 13.7** and **13.7C**).

Patterns by background characteristics

- The percentage of children whose stools are disposed of appropriately ranges from 58% among children age 18–23 months to 42% among children age 0–1 month.
- Sixty-five percent of children in rural areas have their stools disposed of appropriately compared with 29% of children in urban areas.
- Appropriate disposal of children's stools is more common among children in the lowest wealth quintile (46%) and the second wealth quintile (75%) than among those in the fourth wealth quintile (43%) and the highest wealth quintile (20%).

13.4 Handwashing

Handwashing facilities

Basic

Availability of a handwashing facility on the premises with soap and water.

Limited

Availability of a handwashing facility on the premises without soap and water.

Sample: De jure population for whom a place for handwashing was observed or with no place for handwashing in dwelling, yard, or plot; excludes the de jure population for whom permission to see the facility was not granted

Handwashing is an important step in monitoring hygiene and preventing the spread of disease. Rather than asking direct questions on the practice of handwashing, which can be subject to overreporting, interviewers asked to see the place where members of the household most often washed their hands.

A place for washing hands was observed for 87% of the de jure population (**Table 13.8**). Of the observed handwashing places, 28% were in a fixed location and 59% were mobile. Additional information on the distribution of handwashing facilities is provided in Appendix C (see Table C.8). According to definitions of handwashing facilities developed by the JMP, 53% of the population has a basic handwashing facility and 36% has a limited handwashing facility.

Patterns by background characteristics

- The percentage of the de jure population with a basic handwashing facility is higher in urban areas (60%) than in rural areas (49%).
- The percentage of the de jure population with a basic handwashing facility increases with wealth, from 39% in the lowest wealth quintile to 75% in the highest wealth quintile.
- By county, Busia has the highest percentage of the de jure population with access to a basic handwashing facility (94%), followed by Trans Nzoia (91%), Kwale (83%), and Tharaka Nithi (83%). Wajir has the lowest percentage (15%), followed by Taita Taveta (17%) and Garissa (18%) (**Table 13.8C**).

13.5 MENSTRUAL HYGIENE

Appropriate menstrual hygiene materials

Reusable sanitary pads, disposable sanitary pads, tampons, menstrual cups, cloths, toilet paper, and/or cotton wool.

Sample: Women age 15–49 with a menstrual period in the last year

Privacy and use of appropriate menstrual hygiene materials

Percentage of women who were able to wash and change in privacy and who used appropriate materials during their last menstruation.

Sample: Women age 15–49 with a menstrual period in the last year who were home during their last menstrual period


Among women age 15–49 with a menstrual period in the past year, the most commonly used material for collecting or absorbing blood from the most recent menstrual period was disposable sanitary pads (94%), followed by cloth (4%). Almost all women (98%) who were at home during their last menstrual period were able to wash and change in privacy, and 97% used appropriate materials to collect or absorb menstrual blood (**Table 13.9**).

Patterns by background characteristics

- The percentage of women who were able to wash in private, change in private, and use appropriate menstrual hygiene materials during their last menstruation is highest among women with a secondary education and those with more than a secondary education (98% each). It is lowest among those with no education (89%) (**Figure 13.7**).
- Twenty-two percent of women with no education, compared with less than 1% of those with more than a secondary education, used cloth to collect or absorb blood from their most recent menstrual period.

Figure 13.7 **Menstrual hygiene by education**

Percentage of women who were able to wash and change in privacy and who used appropriate materials during last menstruation



- Most women in the highest and fourth wealth quintiles (96% and 97%, respectively) used disposable sanitary towels to collect or absorb blood from their most recent menstrual period, compared with 85% of women in the lowest wealth quintile.
- At the county level, using cloth to collect or absorb menstrual blood from the most recent period is most common in Mandera (55%), followed by Garissa (43%) and Wajir (29%) (**Table 13.9C**).

List of Tables

For more information on water and sanitation characteristics, see the following tables:

- **Table 13.1 Household drinking water**
- **Table 13.2 Drinking water service ladder**
- **Table 13.2C Drinking water service ladder: County level**
- **Table 13.3 Household sanitation facilities**
- **Table 13.4 Sanitation service ladder**
- **Table 13.4C Sanitation service ladder: County level**
- **Table 13.5 Emptying and removal of waste from on-site sanitation facilities**
- **Table 13.5C Emptying and removal of waste from on-site sanitation facilities: County level**
- **Table 13.6 Management of household excreta**
- **Table 13.6C Management of household excreta: County level**
- **Table 13.7 Disposal of children's stools**
- **Table 13.7C Disposal of children's stools: County level**
- **Table 13.8 Handwashing**
- **Table 13.8C Handwashing: County level**
- **Table 13.9 Menstrual hygiene**
- **Table 13.9C Menstrual hygiene: County level**

Table 13.1 Household drinking water

Percent distribution of households and de jure population by source of drinking water and by time to obtain drinking water, according to residence, Kenya Mini DHS 2025–26

	Households			Population		
	Residence			Residence		
Characteristic	Urban	Rural	Total	Urban	Rural	Total
Source of drinking water						
Improved source	98.2	73.3	84.6	98.0	70.6	81.3
Piped into dwelling/yard/plot	41.2	16.1	27.5	40.4	13.5	24.0
Piped to neighbour	9.0	5.2	6.9	9.2	4.7	6.4
Public tap/standpipe	13.9	7.2	10.3	13.4	7.6	9.9
Tube well or borehole	3.9	13.8	9.3	4.0	14.8	10.5
Protected dug well	1.6	7.8	5.0	1.9	8.1	5.7
Protected spring	1.6	11.4	6.9	1.7	11.6	7.7
Rainwater	2.7	8.8	6.1	3.0	7.7	5.9
Tanker truck/cart with small tank	4.9	1.8	3.2	5.6	2.0	3.4
Bottled water	19.3	1.1	9.4	18.7	0.7	7.8
Unimproved source	0.6	6.6	3.9	0.7	7.3	4.7
Unprotected dug well	0.2	2.3	1.4	0.3	2.7	1.7
Unprotected spring	0.2	4.2	2.4	0.3	4.6	2.9
Other	0.2	0.1	0.1	0.2	0.1	0.1
Surface water	1.1	20.1	11.5	1.3	22.1	14.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Time to obtain drinking water (round trip)						
Water on premises[1]	77.6	39.9	57.0	77.4	35.6	51.9
30 minutes or less	20.0	39.9	30.8	19.2	41.3	32.7
More than 30 minutes	2.1	20.1	11.9	3.0	23.0	15.1
Don't know	0.3	0.1	0.2	0.4	0.1	0.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Number of households/ population	11,975	14,405	26,380	36,354	56,592	92,946

[1] Includes water piped to a neighbour and water with a reported round-trip collection time of zero minutes

Table 13.2 Drinking water service ladder

Percent distribution of de jure population by drinking water service ladder, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	At least basic service[1]	Limited service[2]	Unimproved[3]	Surface water	Total	Number of persons
Residence						
Urban	95.2	2.8	0.7	1.3	100.0	36,354
Rural	59.3	11.3	7.3	22.1	100.0	56,592
Wealth quintile						
Lowest	39.8	17.2	8.9	34.1	100.0	18,594
Second	58.6	11.4	7.7	22.3	100.0	18,589
Middle	76.4	7.0	5.3	11.4	100.0	18,584
Fourth	93.6	2.8	1.7	1.9	100.0	18,589
Highest	98.4	1.4	0.2	0.1	100.0	18,589
Total	73.4	8.0	4.7	14.0	100.0	92,946

Note: Service ladder concept/definitions based on the WHO/UNICEF Joint Monitoring Programme for Water Supply, Sanitation and Hygiene (JMP).
[1] Defined as drinking water from an improved source, provided that either water is on the premises or round-trip collection time is 30 minutes or less; includes safely managed drinking water, which is not shown separately
[2] Drinking water from an improved source, provided round-trip collection time is more than 30 minutes or is unknown
[3] Drinking water from an unprotected dug well or unprotected spring

Table 13.2C Drinking water service ladder: County level

Percent distribution of de jure population by drinking water service ladder, according to county, Kenya Mini DHS 2025–26

County	At least basic service[1]	Limited service[2]	Unimproved[3]	Surface water	Total	Number of persons
Mombasa	96.5	2.6	0.9	0.0	100.0	2,969
Kwale	58.7	9.8	4.3	27.2	100.0	1,586
Kilifi	79.9	6.0	1.0	13.0	100.0	2,666
Tana River	55.3	1.9	0.8	42.0	100.0	721
Lamu	78.7	11.9	9.0	0.4	100.0	261
Taita Taveta	76.6	3.0	5.9	14.6	100.0	691
Garissa	60.1	16.7	1.8	21.5	100.0	852
Wajir	56.0	39.1	0.9	3.9	100.0	850
Mandera	64.1	12.5	0.6	22.8	100.0	812
Marsabit	44.2	37.2	3.6	15.0	100.0	748
Isiolo	80.5	7.3	1.3	10.9	100.0	377
Meru	82.2	10.1	1.5	6.2	100.0	2,693
Tharaka-Nithi	54.8	3.1	1.4	40.8	100.0	704
Embu	72.2	5.5	8.0	14.3	100.0	1,035
Kitui	31.3	24.9	3.7	40.1	100.0	2,122
Machakos	69.2	11.7	5.6	13.5	100.0	2,993
Makueni	55.9	13.2	8.3	22.5	100.0	1,665
Nyandarua	89.1	3.0	1.1	6.8	100.0	1,060
Nyeri	88.7	0.6	2.1	8.6	100.0	1,148
Kirinyaga	84.7	0.6	0.9	13.8	100.0	1,210
Murang'a	82.5	0.1	9.6	7.8	100.0	1,708
Kiambu	96.9	0.3	1.5	1.3	100.0	6,076
Turkana	44.3	27.5	18.3	9.9	100.0	1,183
West Pokot	40.5	8.5	13.0	37.9	100.0	1,259
Samburu	32.6	11.7	11.0	44.7	100.0	554
Trans Nzoia	60.9	2.9	7.0	29.2	100.0	1,568
Uasin Gishu	89.7	0.1	6.8	3.3	100.0	2,677
Elgeyo-Marakwet	71.8	1.8	0.0	26.4	100.0	848
Nandi	63.9	0.8	5.0	30.3	100.0	1,550
Baringo	38.3	5.9	16.2	39.6	100.0	1,012
Laikipia	68.3	6.2	11.7	13.8	100.0	931
Nakuru	80.9	4.8	1.6	12.7	100.0	4,659
Narok	26.1	4.7	2.4	66.8	100.0	2,265
Kajiado	77.9	14.2	1.0	6.9	100.0	1,991
Kericho	57.1	4.6	19.0	19.3	100.0	1,777
Bomet	38.9	3.0	9.0	49.1	100.0	1,963
Kakamega	74.7	13.1	11.3	0.9	100.0	3,162
Vihiga	83.8	11.0	1.9	3.3	100.0	1,055
Bungoma	67.0	22.9	7.8	2.2	100.0	3,052
Busia	68.0	10.1	4.3	17.6	100.0	1,758
Siaya	58.9	10.8	4.4	25.9	100.0	1,804
Kisumu	81.4	7.5	6.9	4.2	100.0	2,293
Homa Bay	62.7	3.7	4.3	29.3	100.0	2,246
Migori	66.2	7.4	5.6	20.8	100.0	2,240
Kisii	66.0	19.9	10.5	3.6	100.0	2,210
Nyamira	46.4	27.2	18.3	8.1	100.0	1,050
Nairobi City	97.3	2.7	0.0	0.0	100.0	12,891
Total	73.4	8.0	4.7	14.0	100.0	92,946

Note: Service ladder concept/definitions based on the WHO/UNICEF Joint Monitoring Programme for Water Supply, Sanitation and Hygiene (JMP).
[1] Defined as drinking water from an improved source, provided that either water is on the premises or round-trip collection time is 30 minutes or less; includes safely managed drinking water, which is not shown separately.
[2] Drinking water from an improved source, provided round-trip collection time is more than 30 minutes or is unknown
[3] Drinking water from an unprotected dug well or unprotected spring

Table 13.3 Household sanitation facilities

Percent distribution of households and de jure population by type of toilet/latrine facilities, percent distribution of households and de jure population with a toilet/latrine facility by location of the facility, according to residence, Kenya Mini DHS 2025–26

	Households			Population		
	Residence			Residence		
Type and location of toilet/latrine facility	Urban	Rural	Total	Urban	Rural	Total
Improved sanitation facility	92.9	62.7	76.4	92.7	59.1	72.3
Flush/pour flush to piped sewer system	37.8	0.6	17.5	38.5	0.4	15.3
Flush/pour flush to septic tank	15.8	2.0	8.2	15.1	1.5	6.8
Flush/pour flush to pit latrine	8.4	2.9	5.4	8.5	2.8	5.0
Flush/pour flush, don't know to where	3.8	0.1	1.8	3.5	0.1	1.4
Ventilated improved pit (VIP) latrine	6.7	12.8	10.0	6.5	11.8	9.7
Pit latrine with slab	20.5	44.2	33.4	20.5	42.4	33.8
Composting toilet	0.0	0.2	0.1	0.0	0.2	0.1
Unimproved sanitation facility	6.8	28.5	18.6	6.9	29.8	20.8
Flush/pour flush not to sewer/septic tank/pit latrine	1.4	0.1	0.7	1.3	0.1	0.6
Pit latrine without slab/open pit	4.5	28.0	17.3	4.8	29.1	19.6
Bucket	0.8	0.0	0.4	0.7	0.0	0.3
Hanging toilet/hanging latrine	0.0	0.4	0.2	0.0	0.4	0.3
Other	0.1	0.0	0.1	0.1	0.0	0.1
Open defecation (no facility/bush/field)	0.3	8.8	4.9	0.4	11.1	6.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Number of households/ population	11,975	14,405	26,380	36,354	56,592	92,946
Location of toilet facility						
In own dwelling	37.6	9.3	22.8	40.1	8.8	21.9
In own yard/plot	58.6	84.2	72.0	56.2	85.0	72.9
Elsewhere	3.8	6.5	5.2	3.7	6.2	5.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Number of households/ population with a toilet/latrine facility	11,942	13,136	25,078	36,213	50,305	86,518

Table 13.4 Sanitation service ladder

Percent distribution of de jure population by type of sanitation service, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	At least basic service[1]	Limited service[2]	Unimproved[3]	Open defecation	Total	Number of persons
Residence						
Urban	50.8	41.9	6.9	0.4	100.0	36,354
Rural	40.9	18.2	29.8	11.1	100.0	56,592
Wealth quintile						
Lowest	15.8	15.5	37.1	31.7	100.0	18,594
Second	37.8	21.2	38.6	2.3	100.0	18,589
Middle	48.6	31.2	19.7	0.5	100.0	18,584
Fourth	40.6	52.6	6.8	0.0	100.0	18,589
Highest	81.1	17.0	1.9	0.0	100.0	18,589
Total	44.8	27.5	20.8	6.9	100.0	92,946

Note: Service ladder concept/definitions based on the WHO/UNICEF Joint Monitoring Programme for Water Supply, Sanitation and Hygiene (JMP).

[1] Defined as use of improved facilities that are not shared with other households; includes safely managed sanitation service, which is not shown separately

[2] Defined as use of improved facilities shared by 2 or more households

[3] Use of flush/pour flush toilet not to sewer, septic tank, or pit latrine, pit latrines without a slab/open pit, hanging toilets/latrines, or buckets

Table 13.4C Sanitation service ladder: County level

Percent distribution of de jure population by type of sanitation service, according to county, Kenya Mini DHS 2025–26

County	At least basic service[1]	Limited service[2]	Unimproved[3]	Open defecation	Total	Number of persons
Mombasa	50.2	42.3	7.1	0.4	100.0	2,969
Kwale	30.1	18.4	16.3	35.2	100.0	1,586
Kilifi	35.7	37.5	6.0	20.8	100.0	2,666
Tana River	26.2	21.9	21.5	30.4	100.0	721
Lamu	67.6	11.4	6.1	14.8	100.0	261
Taita Taveta	50.5	23.3	22.6	3.5	100.0	691
Garissa	28.4	30.6	8.0	33.0	100.0	852
Wajir	34.3	13.8	13.2	38.7	100.0	850
Mandera	34.6	29.3	9.7	26.5	100.0	812
Marsabit	28.3	39.0	14.4	18.2	100.0	748
Isiolo	25.6	50.8	10.4	13.3	100.0	377
Meru	54.2	20.4	20.2	5.2	100.0	2,693
Tharaka-Nithi	55.6	19.8	23.9	0.6	100.0	704
Embu	49.0	29.4	21.6	0.0	100.0	1,035
Kitui	50.0	11.7	22.7	15.6	100.0	2,122
Machakos	49.0	42.7	8.2	0.1	100.0	2,993
Makueni	70.7	24.5	4.7	0.0	100.0	1,665
Nyandarua	45.5	20.5	34.0	0.1	100.0	1,060
Nyeri	41.6	27.2	31.2	0.0	100.0	1,148
Kirinyaga	58.2	31.0	10.7	0.1	100.0	1,210
Murang'a	56.4	36.9	6.7	0.0	100.0	1,708
Kiambu	55.1	32.5	12.4	0.0	100.0	6,076
Turkana	14.8	11.0	3.0	71.2	100.0	1,183
West Pokot	47.5	18.7	1.2	32.6	100.0	1,259
Samburu	13.8	15.4	12.8	57.9	100.0	554
Trans Nzoia	52.1	12.3	35.5	0.1	100.0	1,568
Uasin Gishu	51.0	20.9	27.6	0.5	100.0	2,677
Elgeyo-Marakwet	47.5	11.5	39.0	2.0	100.0	848
Nandi	57.7	21.0	19.0	2.4	100.0	1,550
Baringo	40.1	11.4	21.9	26.7	100.0	1,012
Laikipia	39.1	13.3	38.6	9.0	100.0	931
Nakuru	35.5	36.7	23.6	4.2	100.0	4,659
Narok	22.4	17.8	43.7	16.1	100.0	2,265
Kajiado	39.6	37.2	7.8	15.3	100.0	1,991
Kericho	61.8	33.5	4.3	0.4	100.0	1,777
Bomet	14.8	9.0	76.1	0.1	100.0	1,963
Kakamega	50.1	24.7	25.2	0.0	100.0	3,162
Vihiga	50.7	24.5	18.6	6.2	100.0	1,055
Bungoma	43.3	11.5	45.2	0.0	100.0	3,052
Busia	42.8	27.6	28.2	1.4	100.0	1,758
Siaya	35.0	19.0	39.4	6.7	100.0	1,804
Kisumu	36.3	35.8	23.9	4.0	100.0	2,293
Homa Bay	19.5	15.7	53.5	11.2	100.0	2,246
Migori	34.6	31.2	29.5	4.7	100.0	2,240
Kisii	37.6	12.8	49.6	0.0	100.0	2,210
Nyamira	41.8	11.5	46.7	0.0	100.0	1,050
Nairobi City	56.1	39.8	4.1	0.0	100.0	12,891
Total	44.8	27.5	20.8	6.9	100.0	92,946

Note: Service ladder concept/definitions based on the WHO/UNICEF Joint Monitoring Programme for Water Supply, Sanitation and Hygiene (JMP).
[1] Defined as use of improved facilities that are not shared with other households. Includes safely managed sanitation service, which is not shown separately
[2] Defined as use of improved facilities shared by 2 or more households
[3] Use of flush/pour flush toilet not to sewer, septic tank, or pit latrine, pit latrines without a slab/open pit, hanging toilets/latrines, or buckets

Table 13.5 Emptying and removal of wastes from on-site sanitation facilities

Percent distribution of de jure population in households with septic tanks and improved latrines by method of emptying and removal, and percentage of the population with on-site sanitation facilities for which excreta was safely disposed of in situ, percentage with on-site sanitation facilities for which excreta was disposed of unsafely, and percentage with on-site sanitation facilities for which excreta was removed for treatment, according to background characteristics, Kenya Mini DHS 2025–26

	Percent distribution by method of emptying and disposal of waste from septic tanks or other on-site sanitation facilities									Percentage of population with on-site sanitation facilities for which:			
Background characteristic	Removed by a service provider to treatment plant	Removed by a service provider, don't know to where	Buried in a covered pit	Taken to un-covered pit, open ground, water body, or else-where	Other	Don't know where wastes are taken	Never emptied	Don't know if ever emptied	Total	Excreta are safely disposed of in situ[1]	Excreta are disposed of unsafely[2]	Excreta are removed for treat-ment[3]	Number of persons with improved on-site sanitation facilities
Sanitation facility type													
Flush to septic tank	5.3	10.9	0.6	1.7	1.2	0.8	57.9	21.6	100.0	80.1	2.9	17.0	6,361
Latrines and other improved facilities	1.0	2.3	1.1	0.3	0.1	0.5	91.2	3.6	100.0	95.9	0.4	3.7	45,238
Flush to pit latrine	0.9	5.7	2.2	0.4	0.1	1.2	78.7	10.8	100.0	91.8	0.5	7.8	4,656
Ventilated improved pit latrine	1.7	2.4	1.4	0.3	0.2	0.6	90.1	3.3	100.0	94.8	0.4	4.8	9,026
Pit latrine with slab	0.8	1.7	0.8	0.3	0.1	0.3	93.3	2.7	100.0	96.8	0.3	2.9	31,423
Composting toilet	3.8	0.0	0.0	0.0	2.3	0.0	87.0	6.9	100.0	93.9	2.3	3.8	133
Residence													
Urban	3.5	8.4	1.3	1.0	0.5	1.1	70.3	13.9	100.0	85.5	1.5	13.0	18,412
Rural	0.4	0.6	0.9	0.2	0.1	0.2	96.3	1.4	100.0	98.6	0.2	1.2	33,188
Wealth quintile													
Lowest	0.0	0.0	0.5	0.1	0.1	0.1	97.1	2.1	100.0	99.7	0.2	0.1	5,794
Second	0.2	0.3	0.9	0.2	0.1	0.0	96.4	1.9	100.0	99.2	0.2	0.6	10,909
Middle	0.7	1.5	1.3	0.2	0.0	0.6	92.1	3.5	100.0	96.9	0.3	2.8	14,068
Fourth	2.4	5.5	1.1	1.0	0.0	0.7	81.4	7.9	100.0	90.4	1.0	8.6	13,266
Highest	4.5	10.0	1.0	0.5	1.2	1.0	66.5	15.3	100.0	82.7	1.8	15.5	7,562
Total	1.5	3.4	1.0	0.4	0.2	0.5	87.1	5.9	100.0	93.9	0.7	5.4	51,600

Notes: On-site sanitation facilities are those in which the excreta are stored in a septic tank, pit latrine, or composting toilet.
In situ- on site
[1] Includes septic tanks and latrines in which wastes have been buried in a covered pit, that have never been emptied, and don't know if they have ever been emptied
[2] Includes septic tanks and latrines in which wastes have been emptied to uncovered pits, open ground, water bodies, or other locations
[3] Includes septic tanks and latrines in which wastes have been removed by a service provider to a treatment plant or unknown location or have been removed by a non-service provider to an unknown location

Table 13.5C Emptying and removal of waste from on-site sanitation facilities: County level

Percent distribution of de jure population in households with septic tanks or other on-site sanitation facilities by method of emptying and removal; and among population with on-site sanitation facilities, percentage for which excreta are safely disposed of in situ, percentage for which excreta are disposed of unsafely, and percentage for which excreta are removed for treatment, according to county, Kenya Mini DHS 2025–26

	Percent distribution by method of emptying and disposal of waste from septic tanks or other on-site sanitation facilities									Percentage of population with on-site sanitation facilities for which:			
County	Removed by a service provider to treatment plant	Removed by a service provider, don't know to where	Buried in a covered pit	Taken to un-covered pit, open ground, water body, or else-where	Other	Don't know where wastes are taken	Never emptied	Don't know if ever emptied	Total	Excreta are safely disposed of in situ[1]	Excreta are disposed of un-safely[2]	Excreta are removed for treat-ment[3]	Number of persons with improved on-site sanitation facilities
Mombasa	0.3	1.3	1.4	0.0	0.8	1.1	67.5	27.6	100.0	96.5	0.8	2.7	2,595
Kwale	0.0	0.0	0.9	0.0	0.7	0.0	90.5	7.8	100.0	99.3	0.7	0.0	769
Kilifi	0.0	1.4	3.6	0.9	0.0	0.8	92.7	0.6	100.0	96.9	0.9	2.2	1,846
Tana River	0.0	0.9	1.5	0.3	0.5	0.0	96.8	0.0	100.0	98.4	0.7	0.9	347
Lamu	0.0	0.0	0.0	0.0	0.0	0.0	99.9	0.1	100.0	100.0	0.0	0.0	201
Taita Taveta	0.6	5.6	0.1	2.0	0.0	0.0	88.6	3.1	100.0	91.7	2.0	6.2	509
Garissa	0.0	3.1	0.7	0.0	0.0	0.6	87.9	7.7	100.0	96.2	0.0	3.8	483
Wajir	0.0	6.2	0.0	0.0	0.0	0.8	91.8	1.2	100.0	93.0	0.0	7.0	387
Mandera	0.0	0.0	0.0	0.0	0.0	0.0	99.8	0.2	100.0	100.0	0.0	0.0	518
Marsabit	0.0	0.0	0.0	0.0	0.0	0.0	96.2	3.8	100.0	100.0	0.0	0.0	500
Isiolo	0.0	1.8	4.4	2.0	0.0	0.0	89.5	2.3	100.0	96.2	2.0	1.8	265
Meru	1.8	1.7	0.0	0.0	0.0	0.1	92.0	4.4	100.0	96.4	0.0	3.6	1,990
Tharaka-Nithi	0.3	0.5	0.0	0.0	0.1	0.0	98.4	0.7	100.0	99.2	0.1	0.7	521
Embu	0.0	1.0	0.3	0.0	0.5	3.1	90.7	4.4	100.0	95.4	0.5	4.1	620
Kitui	0.0	0.7	0.5	0.2	0.0	0.8	94.8	3.0	100.0	98.3	0.2	1.5	1,308
Machakos	0.3	6.0	4.5	0.2	0.0	0.0	82.6	6.3	100.0	93.5	0.2	6.3	2,228
Makueni	0.7	0.5	0.0	0.0	0.0	0.4	91.6	6.8	100.0	98.4	0.0	1.6	1,581
Nyandarua	0.0	6.5	1.3	0.3	0.0	0.7	87.4	3.9	100.0	92.6	0.3	7.2	693
Nyeri	0.4	1.0	1.6	0.4	0.0	0.0	95.9	0.8	100.0	98.2	0.4	1.4	774
Kirinyaga	0.2	1.9	0.1	0.0	0.2	0.0	90.6	6.9	100.0	97.6	0.2	2.1	1,058
Murang'a	2.2	0.6	0.0	0.3	0.2	0.2	94.1	2.4	100.0	96.5	0.5	3.0	1,555
Kiambu	2.0	9.0	0.0	0.0	0.0	0.0	77.5	11.5	100.0	89.0	0.0	11.0	2,504
Turkana	0.0	2.4	0.0	0.0	0.0	0.0	93.6	4.1	100.0	97.6	0.0	2.4	304
West Pokot	1.3	0.1	0.0	0.0	0.0	0.1	98.4	0.0	100.0	98.5	0.0	1.5	831
Samburu	0.0	1.3	0.0	0.0	0.0	0.1	95.3	3.3	100.0	98.6	0.0	1.4	160
Trans Nzoia	0.0	0.4	1.2	0.3	0.0	0.2	97.6	0.4	100.0	99.1	0.3	0.6	996
Uasin Gishu	5.8	3.5	0.0	0.0	0.0	2.0	80.9	7.7	100.0	88.6	0.0	11.4	1,591
Elgeyo-Marakwet	0.2	0.0	0.0	0.0	0.0	0.0	96.8	3.0	100.0	99.8	0.0	0.2	499
Nandi	0.0	0.0	0.0	0.0	0.0	0.0	99.7	0.2	100.0	100.0	0.0	0.0	1,186
Baringo	0.5	0.0	0.0	0.0	0.0	0.0	97.6	1.8	100.0	99.5	0.0	0.5	518
Laikipia	0.7	1.4	1.3	0.0	0.6	4.7	87.9	3.3	100.0	92.5	0.6	6.8	375
Nakuru	3.3	9.4	0.1	0.0	0.0	0.4	81.1	5.7	100.0	86.9	0.0	13.1	2,908
Narok	0.7	3.6	0.2	0.1	0.0	0.1	88.4	6.9	100.0	95.5	0.1	4.4	899
Kajiado	9.4	12.4	0.0	0.2	0.3	1.6	54.5	21.5	100.0	76.0	0.5	23.5	1,403
Kericho	1.1	0.5	0.0	0.0	0.0	0.1	96.2	2.1	100.0	98.3	0.0	1.7	1,650
Bomet	0.5	0.0	0.0	0.0	0.0	0.0	99.5	0.0	100.0	99.5	0.0	0.5	463
Kakamega	0.4	0.5	0.3	0.0	0.0	0.0	97.4	1.3	100.0	99.0	0.0	1.0	2,286
Vihiga	0.0	0.4	0.0	0.0	0.0	0.0	99.3	0.2	100.0	99.6	0.0	0.4	793
Bungoma	0.0	0.3	0.4	0.2	0.0	0.0	98.2	0.9	100.0	99.4	0.2	0.4	1,629
Busia	0.5	2.6	0.0	0.4	0.0	0.3	92.7	3.5	100.0	96.2	0.4	3.4	1,211
Siaya	0.3	0.2	0.2	0.0	0.0	0.7	98.7	0.0	100.0	98.9	0.0	1.1	929
Kisumu	7.7	11.3	7.3	2.0	1.0	1.7	65.4	3.6	100.0	76.3	3.0	20.7	1,583
Homa Bay	1.0	2.6	7.2	0.5	0.0	0.0	86.7	2.0	100.0	95.9	0.5	3.6	777
Migori	4.0	0.2	4.3	1.6	0.0	0.0	88.7	1.2	100.0	94.2	1.6	4.2	1,432
Kisii	0.0	0.0	0.0	0.0	0.0	0.2	99.7	0.1	100.0	99.8	0.0	0.2	1,104
Nyamira	0.0	0.8	0.0	0.0	0.0	0.0	98.7	0.6	100.0	99.2	0.0	0.8	557
Nairobi City	2.6	12.0	0.0	4.7	2.5	1.6	59.7	16.9	100.0	76.6	7.2	16.2	2,266
Total	1.5	3.4	1.0	0.4	0.2	0.5	87.1	5.9	100.0	93.9	0.7	5.4	51,600

Note: On-site sanitation facilities are those in which the excreta are stored in a septic tank, pit latrine, or composting toilet.
In situ refers to on-site disposal.
[1] Includes septic tanks and latrines in which wastes have been buried in a covered pit, that have never been emptied, and don't know if they have ever been emptied
[2] Includes septic tanks and latrines in which wastes have been emptied to uncovered pits, open ground, water bodies, or other locations
[3] Includes septic tanks and latrines in which wastes have been removed by a service provider to a treatment plant or unknown location or have been removed by a non-service provider to an unknown location

Table 13.6 Management of household excreta

Percent distribution of de jure population by management of excreta from household sanitation facilities, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Connected to sewer	Using improved on-site sanitation facilities			Using improved sanitation facilities, on-site status unknown	Using un-improved sanitation facilities	Practicing open defecation	Total	Percent-age connected to sewer, safely disposed of on-site, or removed for treatment off-site	Number of persons
		Safe disposal in situ of excreta from on-site sanitation facilities	Unsafe disposal of excreta from on-site sanitation facilities	Removal of excreta for treatment off-site						
Residence										
Urban	38.5	43.3	0.8	6.6	3.5	6.9	0.4	100.0	88.4	36,354
Rural	0.4	57.8	0.1	0.7	0.1	29.8	11.1	100.0	58.9	56,592
Wealth quintile										
Lowest	0.0	31.1	0.1	0.0	0.1	37.1	31.7	100.0	31.1	18,594
Second	0.3	58.2	0.1	0.3	0.0	38.6	2.3	100.0	58.9	18,589
Middle	3.1	73.3	0.2	2.1	1.0	19.7	0.5	100.0	78.6	18,584
Fourth	20.5	64.5	0.7	6.1	1.2	6.8	0.0	100.0	91.2	18,589
Highest	52.7	33.7	0.7	6.3	4.7	1.9	0.0	100.0	92.7	18,589
Total	15.3	52.2	0.4	3.0	1.4	20.8	6.9	100.0	70.5	92,946

Note: On-site sanitation facilities are those in which the excreta are stored in a septic tank, pit latrine, or composting toilet.
In situ refers to on-site disposal.

Table 13.6C Management of household excreta: County level

Percent distribution of de jure population by management of excreta from household sanitation facilities, according to county, Kenya Mini DHS 2025–26

County	Connected to sewer	Using improved on-site sanitation facilities: Safe disposal in situ of excreta from on-site sanitation facilities	Using improved on-site sanitation facilities: Unsafe disposal of excreta from on-site sanitation facilities	Using improved on-site sanitation facilities: Removal of excreta for treatment off-site	Using improved sanitation facilities, on-site status unknown	Using un-improved sanitation facilities	Practicing open defecation	Total	Percent-age connected to sewer, safely disposed of on-site, or removed for treatment off-site	Number of persons
Mombasa	4.7	84.4	0.7	2.4	0.4	7.1	0.4	100.0	91.4	2,969
Kwale	0.0	48.2	0.4	0.0	0.0	16.3	35.2	100.0	48.2	1,586
Kilifi	3.9	67.1	0.6	1.5	0.0	6.0	20.8	100.0	72.5	2,666
Tana River	0.0	47.3	0.4	0.4	0.0	21.5	30.4	100.0	47.8	721
Lamu	1.8	77.0	0.0	0.0	0.3	6.1	14.8	100.0	78.8	261
Taita Taveta	0.1	67.6	1.5	4.6	0.0	22.6	3.5	100.0	72.3	691
Garissa	2.3	54.5	0.0	2.1	0.0	8.0	33.0	100.0	59.0	852
Wajir	2.6	42.4	0.0	3.2	0.0	13.2	38.7	100.0	48.1	850
Mandera	0.1	63.8	0.0	0.0	0.0	9.7	26.5	100.0	63.9	812
Marsabit	0.5	66.9	0.0	0.0	0.0	14.4	18.2	100.0	67.4	748
Isiolo	6.1	67.6	1.4	1.3	0.0	10.4	13.3	100.0	74.9	377
Meru	0.6	71.2	0.0	2.7	0.1	20.2	5.2	100.0	74.5	2,693
Tharaka-Nithi	1.3	73.4	0.1	0.5	0.2	23.9	0.6	100.0	75.2	704
Embu	16.8	57.1	0.3	2.5	1.8	21.6	0.0	100.0	76.4	1,035
Kitui	0.0	60.6	0.1	0.9	0.1	22.7	15.6	100.0	61.5	2,122
Machakos	15.0	69.6	0.2	4.7	2.3	8.2	0.1	100.0	89.2	2,993
Makueni	0.3	93.4	0.0	1.5	0.0	4.7	0.0	100.0	95.2	1,665
Nyandarua	0.1	60.5	0.2	4.7	0.5	34.0	0.1	100.0	65.3	1,060
Nyeri	1.3	66.2	0.3	0.9	0.1	31.2	0.0	100.0	68.5	1,148
Kirinyaga	1.4	85.3	0.2	1.9	0.5	10.7	0.1	100.0	88.6	1,210
Murang'a	1.2	87.8	0.5	2.7	1.1	6.7	0.0	100.0	91.8	1,708
Kiambu	40.5	36.7	0.0	4.5	5.9	12.4	0.0	100.0	81.7	6,076
Turkana	0.1	25.1	0.0	0.6	0.0	3.0	71.2	100.0	25.8	1,183
West Pokot	0.2	65.0	0.0	1.0	0.0	1.2	32.6	100.0	66.2	1,259
Samburu	0.3	28.5	0.0	0.4	0.1	12.8	57.9	100.0	29.2	554
Trans Nzoia	0.9	63.0	0.2	0.4	0.0	35.5	0.1	100.0	64.2	1,568
Uasin Gishu	11.7	52.7	0.0	6.8	0.7	27.6	0.5	100.0	71.2	2,677
Elgeyo-Marakwet	0.1	58.7	0.0	0.1	0.0	39.0	2.0	100.0	58.9	848
Nandi	2.1	76.5	0.0	0.0	0.0	19.0	2.4	100.0	78.6	1,550
Baringo	0.3	50.9	0.0	0.3	0.0	21.9	26.7	100.0	51.5	1,012
Laikipia	9.3	37.3	0.3	2.7	2.9	38.6	9.0	100.0	49.3	931
Nakuru	8.9	54.2	0.0	8.2	0.8	23.6	4.2	100.0	71.4	4,659
Narok	0.6	37.9	0.1	1.7	0.0	43.7	16.1	100.0	40.2	2,265
Kajiado	3.8	53.6	0.3	16.6	2.6	7.8	15.3	100.0	73.9	1,991
Kericho	2.0	91.3	0.0	1.5	0.5	4.3	0.4	100.0	94.8	1,777
Bomet	0.2	23.5	0.0	0.1	0.0	76.1	0.1	100.0	23.8	1,963
Kakamega	2.6	71.6	0.0	0.7	0.0	25.2	0.0	100.0	74.8	3,162
Vihiga	0.0	74.9	0.0	0.3	0.0	18.6	6.2	100.0	75.2	1,055
Bungoma	1.4	53.1	0.1	0.2	0.0	45.2	0.0	100.0	54.7	3,052
Busia	1.4	66.2	0.3	2.3	0.1	28.2	1.4	100.0	70.0	1,758
Siaya	2.3	50.9	0.0	0.6	0.1	39.4	6.7	100.0	53.8	1,804
Kisumu	3.1	52.6	2.1	14.3	0.0	23.9	4.0	100.0	70.0	2,293
Homa Bay	0.6	33.2	0.2	1.2	0.0	53.5	11.2	100.0	35.1	2,246
Migori	1.9	60.2	1.0	2.7	0.0	29.5	4.7	100.0	64.8	2,240
Kisii	0.4	49.9	0.0	0.1	0.0	49.6	0.0	100.0	50.4	2,210
Nyamira	0.3	52.6	0.0	0.4	0.0	46.7	0.0	100.0	53.3	1,050
Nairobi City	73.2	13.5	1.3	2.9	5.1	4.1	0.0	100.0	89.5	12,891
Total	15.3	52.2	0.4	3.0	1.4	20.8	6.9	100.0	70.5	92,946

Note: On-site sanitation facilities are those in which the excreta are stored in a septic tank, pit latrine, or composting toilet.
In situ refers to on-site disposal.

Table 13.7 Disposal of children's stools

Percent distribution of youngest children under age 2 living with the mother by the manner of disposal of the child's last faecal matter, and percentage of children whose stools are disposed of appropriately, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Manner of disposal of children's stools							Total	Percentage of children whose stools are disposed of appro-priately[1]	Number of children
	Child used toilet or latrine	Put/rinsed into toilet or latrine	Buried	Put/rinsed into drain or ditch	Thrown into garbage	Left in the open	Other			
Age of child in months										
0–1	0.4	41.6	4.7	5.1	44.6	2.7	0.9	100.0	42.0	327
2–3	0.7	46.8	3.6	3.6	40.4	3.0	1.9	100.0	47.5	334
4–5	0.7	43.0	2.8	5.5	45.1	2.5	0.4	100.0	43.7	355
6–8	1.0	42.8	3.4	4.6	44.7	3.3	0.3	100.0	43.7	536
9–11	0.7	52.7	4.7	2.9	34.3	3.6	1.0	100.0	53.5	443
12–17	1.9	48.0	3.0	7.4	35.0	4.0	0.8	100.0	49.9	951
18–23	2.6	55.1	3.5	5.0	29.2	4.4	0.3	100.0	57.7	871
6–23	1.8	49.9	3.5	5.4	34.9	3.9	0.6	100.0	51.7	2,802
Type of toilet facility[2]										
Improved sanitation facility	1.6	47.5	1.7	3.1	44.6	1.0	0.6	100.0	49.1	2,771
Unimproved facility	1.5	67.7	2.2	9.7	15.4	2.2	1.4	100.0	69.2	722
Open defecation	0.0	12.2	22.0	13.8	22.4	29.0	0.5	100.0	12.2	325
Residence										
Urban	1.5	27.9	1.0	2.5	66.7	0.2	0.2	100.0	29.4	1,614
Rural	1.4	63.2	5.4	7.2	15.5	6.1	1.1	100.0	64.7	2,204
Mother's education[3]										
No education	3.1	21.4	17.2	6.9	33.3	17.8	0.3	100.0	24.5	341
Primary	1.1	56.0	3.8	7.1	27.8	3.7	0.6	100.0	57.0	1,032
Secondary	1.4	52.6	1.4	5.0	36.5	2.3	0.8	100.0	54.0	1,666
More than secondary	1.3	40.7	1.7	2.6	52.9	0.1	0.7	100.0	42.1	779
Wealth quintile										
Lowest	1.4	44.8	11.6	10.2	16.9	13.6	1.5	100.0	46.2	815
Second	0.8	74.6	2.3	6.3	12.9	2.7	0.5	100.0	75.4	711
Middle	1.9	65.8	1.4	4.6	24.3	1.0	1.1	100.0	67.7	695
Fourth	1.6	41.6	0.7	4.1	51.4	0.1	0.4	100.0	43.3	850
Highest	1.5	18.5	1.2	0.6	78.1	0.0	0.1	100.0	20.0	748
Total	1.4	48.3	3.5	5.2	37.2	3.6	0.7	100.0	49.8	3,818

[1] Children's stools are considered disposed of appropriately if the child used a toilet or latrine or if the faecal matter was put/rinsed into a toilet or latrine.
[2] See Table 13.3 for definitions of categories.
[3] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 13.7C Disposal of children's stools: County level

Percent distribution of youngest children under age 2 living with the mother by the manner of disposal of the child's last faecal matter, and percentage of children whose stools are disposed of appropriately, according to county, Kenya Mini DHS 2025–26

	Manner of disposal of children's stools										
County	Child used toilet or latrine	Put/rinsed into toilet or latrine	Buried	Put/rinsed into drain or ditch	Thrown into garbage	Left in the open	Other	Don't know	Total	Percentage of children whose stools are disposed of appropriately[1]	Number of children
Mombasa	0.0	31.3	1.5	0.0	67.2	0.0	0.0	0.0	100.0	31.3	120
Kwale	5.8	50.4	4.9	13.3	24.3	1.3	0.0	0.0	100.0	56.2	69
Kilifi	2.9	48.1	13.8	6.7	23.2	5.2	0.0	0.0	100.0	51.0	97
Tana River	0.0	69.5	0.0	2.4	6.9	21.2	0.0	0.0	100.0	69.5	38
Lamu	2.3	31.7	26.4	0.0	39.6	0.0	0.0	0.0	100.0	34.0	10
Taita Taveta	0.0	68.0	6.7	0.0	25.2	0.0	0.0	0.0	100.0	68.0	27
Garissa	0.0	11.6	24.3	17.4	7.1	39.6	0.0	0.0	100.0	11.6	40
Wajir	0.0	2.1	66.0	0.3	30.2	1.4	0.0	0.0	100.0	2.1	36
Mandera	0.0	47.6	23.9	0.4	22.9	5.1	0.0	0.0	100.0	47.6	38
Marsabit	0.0	58.4	0.0	10.9	22.9	7.9	0.0	0.0	100.0	58.4	34
Isiolo	0.0	15.4	3.1	4.8	53.5	23.1	0.0	0.0	100.0	15.4	16
Meru	2.5	55.2	7.4	1.7	33.2	0.0	0.0	0.0	100.0	57.7	72
Tharaka-Nithi	(0.0)	(95.9)	(0.0)	(0.0)	(1.7)	(0.0)	(2.4)	(0.0)	100.0	(95.9)	22
Embu	(6.9)	(60.5)	(0.0)	(0.0)	(32.5)	(0.0)	(0.0)	(0.0)	100.0	(67.5)	25
Kitui	0.0	60.5	1.7	1.3	28.6	6.2	1.7	0.0	100.0	60.5	63
Machakos	5.3	43.9	0.0	3.3	46.4	0.0	1.2	0.0	100.0	49.1	151
Makueni	1.6	82.9	3.3	1.5	8.7	1.9	0.0	0.0	100.0	84.5	56
Nyandarua	0.0	74.2	2.2	0.0	21.8	0.0	1.8	0.0	100.0	74.2	48
Nyeri	(5.5)	(53.3)	(1.9)	(11.3)	(27.9)	(0.0)	(0.0)	(0.0)	100.0	(58.8)	39
Kirinyaga	(0.0)	(70.2)	(0.0)	(0.0)	(29.8)	(0.0)	(0.0)	(0.0)	100.0	(70.2)	36
Murang'a	0.0	59.3	0.0	0.0	40.7	0.0	0.0	0.0	100.0	59.3	57
Kiambu	1.8	19.9	0.4	7.1	70.3	0.0	0.5	0.0	100.0	21.7	253
Turkana	0.0	18.1	22.0	0.2	54.0	5.7	0.0	0.0	100.0	18.1	59
West Pokot	0.0	46.6	0.0	12.4	26.6	14.4	0.0	0.0	100.0	46.6	72
Samburu	0.0	19.0	0.0	23.1	13.6	40.7	3.7	0.0	100.0	19.0	30
Trans Nzoia	1.6	65.1	0.0	4.8	28.5	0.0	0.0	0.0	100.0	66.7	56
Uasin Gishu	3.0	24.6	0.0	8.0	60.6	2.3	1.5	0.0	100.0	27.6	119
Elgeyo-Marakwet	7.4	86.8	5.8	0.0	0.0	0.0	0.0	0.0	100.0	94.2	32
Nandi	3.8	69.8	9.0	5.3	11.3	0.9	0.0	0.0	100.0	73.6	73
Baringo	0.0	54.9	1.3	5.6	7.2	31.0	0.0	0.0	100.0	54.9	52
Laikipia	11.6	49.0	4.9	3.9	21.7	8.9	0.0	0.0	100.0	60.6	47
Nakuru	0.0	67.2	0.0	1.9	26.0	3.8	1.2	0.0	100.0	67.2	202
Narok	0.0	29.3	0.0	26.4	38.7	4.2	1.4	0.0	100.0	29.3	84
Kajiado	0.0	22.2	6.3	1.6	62.3	6.1	1.6	0.0	100.0	22.2	120
Kericho	1.5	78.2	0.0	1.5	2.7	16.1	0.0	0.0	100.0	79.7	66
Bomet	0.0	53.2	8.6	37.9	0.4	0.0	0.0	0.0	100.0	53.2	88
Kakamega	2.3	87.7	1.1	2.3	4.5	1.0	1.1	0.0	100.0	90.0	118
Vihiga	0.0	98.4	0.0	0.0	1.6	0.0	0.0	0.0	100.0	98.4	38
Bungoma	0.0	88.1	0.0	0.8	1.9	0.0	9.1	0.0	100.0	88.1	123
Busia	0.0	79.6	0.9	7.4	10.8	1.2	0.0	0.0	100.0	79.6	73
Siaya	1.8	62.0	6.0	9.2	15.5	5.5	0.0	0.0	100.0	63.8	52
Kisumu	0.0	51.9	2.1	5.5	40.5	0.0	0.0	0.0	100.0	51.9	96
Homa Bay	0.0	39.7	5.2	7.6	37.3	10.2	0.0	0.0	100.0	39.7	73
Migori	0.3	83.0	2.9	2.9	9.0	1.5	0.3	0.0	100.0	83.3	128
Kisii	0.0	95.4	0.0	1.3	1.8	1.5	0.0	0.0	100.0	95.4	78
Nyamira	0.0	94.3	0.0	0.0	4.4	0.0	1.3	0.0	100.0	94.3	38
Nairobi City	1.6	12.9	0.7	2.2	82.5	0.0	0.0	0.0	100.0	14.6	553
Total	1.4	48.3	3.5	5.2	37.2	3.6	0.7	0.0	100.0	49.8	3,818

Note: Figures in parentheses are based on 25–49 unweighted cases.
[1] Children's stools are considered disposed of appropriately if the child used a toilet or latrine or if the faecal matter was put/rinsed into a toilet or latrine.

Table 13.8 Handwashing

Percentage of the de jure population for whom the place most often used for washing hands was observed, by whether the location was fixed or mobile, and total percentage of the de jure population for whom the place for handwashing was observed; and among the de jure population for whom the place for handwashing was observed, percentage with water available, percentage with soap available, and percentage with a cleansing agent other than soap available; percentage of the de jure population with a basic handwashing facility and percentage with a limited handwashing facility, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Percentage of de jure population for whom place for washing hands was observed			Number of persons	Place for handwashing observed and:			Number of persons for whom place for hand-washing was observed	Percent-age of the de jure population with a basic hand-washing facility[3]	Percent-age of the de jure population with a limited hand-washing facility[4]	Number of persons for whom a place for hand-washing was observed or with no place for hand-washing in the dwelling, yard, or plot
	Place for hand-washing was a fixed place	Place for hand-washing was mobile	Total		Water available	Soap available[1]	Cleansing agent other than soap available[2]				
Residence											
Urban	45.6	42.9	88.6	36,354	88.8	69.5	0.2	32,202	59.5	31.9	35,219
Rural	16.0	69.7	85.7	56,592	85.5	58.0	1.1	48,486	49.2	37.7	55,763
Wealth quintile											
Lowest	2.2	80.6	82.8	18,594	77.3	48.2	2.8	15,400	39.2	44.9	18,308
Second	7.6	75.3	82.9	18,589	86.5	57.6	0.3	15,403	47.6	36.7	18,276
Middle	20.0	66.1	86.1	18,584	88.8	60.0	0.3	16,004	51.1	36.6	18,257
Fourth	32.4	56.1	88.5	18,589	89.5	62.3	0.1	16,451	53.6	37.2	18,110
Highest	75.8	18.0	93.8	18,589	91.3	82.3	0.1	17,431	74.7	21.9	18,033
Total	27.6	59.2	86.8	92,946	86.8	62.6	0.7	80,688	53.2	35.5	90,983

[1] Soap includes soap or detergent in bar, liquid, powder, or paste form.
[2] Cleansing agents other than soap include locally available materials such as ash, mud, or sand.
[3] The availability of a handwashing facility on premises with soap and water
[4] The availability of a handwashing facility on premises without soap and/or water

Table 13.8C Handwashing: County level

Percentage of the de jure population for whom the place most often used for washing hands was observed, by whether the location was fixed or mobile, and total percentage of the de jure population for whom the place for handwashing was observed; and among the de jure population for whom the place for handwashing was observed, percentage with water available, percentage with soap available, and percentage with a cleansing agent other than soap available; percentage of the de jure population with a basic handwashing facility and percentage with a limited handwashing facility, according to county, Kenya Mini DHS 2025–26

County	Percentage of de jure population for whom place for washing hands was observed			Number of persons	Place for handwashing observed and:			Number of persons for whom place for handwashing was observed	Percentage of the de jure population with a basic handwashing facility[3]	Percentage of the de jure population with a limited handwashing facility[4]	Number of persons for whom a place for handwashing was observed or with no place for handwashing in the dwelling, yard, or plot
	Place for handwashing was a fixed place	Place for handwashing was mobile	Total		Water available	Soap available[1]	Cleansing agent other than soap available[2]				
Mombasa	25.9	33.1	59.1	2,969	99.4	49.1	0.0	1,754	29.1	30.8	2,926
Kwale	8.7	88.1	96.8	1,586	89.5	88.0	0.0	1,535	82.8	14.3	1,581
Kilifi	12.1	87.9	100.0	2,666	81.0	51.4	0.0	2,666	46.8	53.2	2,666
Tana River	4.6	95.4	100.0	721	86.6	21.9	0.0	721	21.9	78.1	721
Lamu	15.9	47.3	63.3	261	97.5	40.5	1.5	165	26.0	38.1	258
Taita Taveta	33.7	43.0	76.7	691	67.8	22.5	0.0	530	17.3	59.7	688
Garissa	6.0	44.8	50.8	852	98.6	35.8	1.8	433	18.2	32.6	852
Wajir	7.2	66.2	73.5	850	95.8	20.7	3.4	625	15.2	58.2	850
Mandera	7.0	91.1	98.1	812	93.3	54.8	4.3	797	54.4	44.9	802
Marsabit	6.9	69.0	75.9	748	71.0	58.2	5.5	568	41.1	35.2	744
Isiolo	21.0	32.6	53.6	377	98.3	64.4	0.1	202	34.3	19.3	377
Meru	28.9	65.8	94.6	2,693	84.3	41.1	0.0	2,549	40.9	58.7	2,560
Tharaka-Nithi	28.3	71.5	99.7	704	97.3	83.0	0.1	702	82.8	17.0	704
Embu	45.4	50.2	95.6	1,035	95.0	40.1	0.0	989	38.7	58.2	1,020
Kitui	13.9	86.1	100.0	2,122	52.2	23.5	0.0	2,122	22.9	77.1	2,122
Machakos	23.9	43.8	67.8	2,993	86.4	60.0	0.0	2,029	40.4	28.6	2,941
Makueni	23.9	72.6	96.5	1,665	78.9	31.2	0.4	1,605	29.4	67.0	1,665
Nyandarua	36.0	63.8	99.7	1,060	70.1	33.1	0.0	1,058	31.6	68.3	1,058
Nyeri	51.1	35.8	86.9	1,148	99.2	90.1	0.3	998	79.0	9.0	1,134
Kirinyaga	52.5	38.1	90.5	1,210	93.4	35.1	0.4	1,096	34.0	64.3	1,115
Murang'a	47.3	50.1	97.4	1,708	91.5	73.0	0.0	1,664	70.1	29.6	1,669
Kiambu	72.5	18.7	91.2	6,076	88.6	54.0	0.1	5,543	52.6	46.4	5,596
Turkana	11.0	86.4	97.4	1,183	82.8	57.2	23.1	1,152	55.6	41.9	1,181
West Pokot	9.6	64.3	73.9	1,259	60.3	41.3	6.2	931	29.8	45.5	1,236
Samburu	10.9	55.8	66.7	554	94.8	70.0	2.9	370	46.9	20.7	547
Trans Nzoia	12.2	85.3	97.5	1,568	99.6	93.7	0.0	1,528	91.0	6.5	1,567
Uasin Gishu	33.8	65.3	99.1	2,677	89.8	44.5	0.3	2,654	43.9	55.3	2,676
Elgeyo-Marakwet	27.5	71.4	98.9	848	98.0	61.8	4.6	839	61.1	37.8	848
Nandi	28.3	62.0	90.3	1,550	94.8	60.3	0.0	1,399	56.6	38.7	1,468
Baringo	10.7	63.8	74.5	1,012	80.4	53.7	1.2	754	40.2	34.9	1,004
Laikipia	38.1	43.9	82.0	931	89.3	38.0	0.1	763	30.6	51.4	931
Nakuru	26.1	67.3	93.4	4,659	90.8	72.1	0.3	4,349	65.9	28.4	4,614
Narok	4.3	80.2	84.6	2,265	94.1	89.0	1.2	1,916	72.9	11.7	2,265
Kajiado	32.0	67.2	99.1	1,991	89.9	82.3	0.1	1,974	81.3	17.9	1,991
Kericho	22.8	75.5	98.3	1,777	84.3	72.3	0.0	1,746	70.5	28.1	1,771
Bomet	11.6	37.8	49.4	1,963	81.3	77.6	0.0	969	38.4	11.3	1,951
Kakamega	15.6	75.5	91.2	3,162	78.6	27.5	0.0	2,882	23.0	68.1	3,162
Vihiga	7.1	33.4	40.5	1,055	97.2	90.2	0.0	427	37.2	4.6	1,022
Bungoma	6.9	87.8	94.8	3,052	99.7	78.1	0.0	2,892	74.1	21.1	3,038
Busia	10.5	78.7	89.1	1,758	99.1	94.2	0.0	1,567	94.2	5.8	1,567
Siaya	7.0	46.6	53.6	1,804	65.0	50.5	0.0	967	24.9	29.7	1,769
Kisumu	16.8	78.7	95.5	2,293	61.8	59.2	0.2	2,189	49.1	50.9	2,189
Homa Bay	9.7	90.3	100.0	2,246	98.4	73.7	0.2	2,246	73.6	26.4	2,246
Migori	15.7	39.9	55.6	2,240	98.3	92.7	0.0	1,246	51.7	4.3	2,224
Kisii	5.4	62.2	67.6	2,210	98.8	88.0	0.0	1,493	60.1	8.2	2,188
Nyamira	5.1	94.9	100.0	1,050	93.5	66.4	0.2	1,050	66.3	33.7	1,050
Nairobi City	52.8	40.6	93.3	12,891	84.9	77.5	0.2	12,033	65.8	31.0	12,430
Total	27.6	59.2	86.8	92,946	86.8	62.6	0.7	80,688	53.2	35.5	90,983

[1] Soap includes soap or detergent in bar, liquid, powder, or paste form.
[2] Cleansing agents other than soap include locally available materials such as ash, mud, or sand.
[3] The availability of a handwashing facility on premises with soap and water
[4] The availability of a handwashing facility on premises without soap and/or water

Table 13.9 Menstrual hygiene

Among women age 15–49 whose most recent menstrual period was in the last year, percentage who used specified materials to collect or absorb blood from the most recent menstrual period; and among women age 15–49 whose most recent menstrual period was in the last year and were at home during their last menstrual period, percentage who were able to wash and change in privacy while at home and percentage who were able to both wash and change in privacy and who used appropriate materials during their last menstruation, according to background characteristics, Kenya Mini DHS 2025–26

	Among women whose most recent menstrual period was in the last year, percentage who used the specified materials to collect or absorb blood from most recent menstrual period											Among women whose most recent menstrual period was in the last year and who were at home during their last menstrual period		
Background characteristic	Re-usable sanitary pads	Dis-posable sanitary pads	Tam-pons	Men-strual cup	Cloth	Toilet paper	Cotton wool	Under-wear only	Other	Nothing	Number of women	Percent-age able to wash and change in privacy	Percent-age able to wash and change in privacy and who used appro-priate materials during last men-stru-ation[1]	Number of women
Age														
15–19	1.7	96.7	0.2	0.0	2.5	0.2	0.2	0.5	0.0	0.1	3,726	97.1	96.8	3,680
20–24	1.6	95.7	2.0	0.2	2.0	0.3	0.5	0.8	0.0	0.3	3,269	97.9	97.1	3,252
25–29	1.5	94.9	2.6	0.4	2.3	0.1	0.6	0.9	0.0	0.1	3,156	98.0	97.2	3,143
30–34	1.5	93.1	2.7	0.1	3.7	0.2	1.1	0.8	0.0	1.0	2,678	97.2	96.0	2,669
35–39	1.1	92.2	1.2	0.0	6.1	0.2	0.6	1.4	0.0	0.8	2,614	97.6	96.1	2,600
40–44	1.1	92.4	1.4	0.0	6.5	0.4	0.3	1.6	0.0	0.7	1,817	97.4	95.9	1,807
45–49	2.0	91.2	0.8	0.1	5.5	0.4	1.8	1.6	0.0	0.7	1,288	97.9	96.0	1,287
Residence														
Urban	1.4	95.8	2.9	0.3	1.8	0.2	0.7	0.3	0.0	0.4	8,862	98.0	97.6	8,814
Rural	1.6	92.9	0.4	0.0	5.3	0.3	0.6	1.6	0.0	0.5	9,686	97.1	95.6	9,625
Education[2]														
No education	1.3	75.9	0.1	0.0	22.4	0.0	0.2	5.0	0.0	1.8	901	92.5	88.8	899
Primary	1.5	91.6	0.3	0.0	6.2	0.4	1.3	1.8	0.0	0.9	5,026	96.8	94.8	5,008
Secondary	1.6	96.8	1.2	0.1	1.7	0.2	0.5	0.5	0.0	0.1	8,764	98.0	97.6	8,702
More than secondary	1.3	96.1	4.7	0.6	0.5	0.2	0.2	0.0	0.0	0.3	3,858	98.7	98.3	3,830
Wealth quintile														
Lowest	1.4	84.6	0.0	0.0	12.7	0.2	0.8	3.5	0.1	1.2	2,695	95.1	91.9	2,687
Second	1.6	94.2	0.1	0.0	4.7	0.4	0.7	1.6	0.0	0.3	3,295	97.5	96.2	3,280
Middle	1.7	96.0	0.6	0.0	2.5	0.4	0.8	0.5	0.0	0.3	3,573	97.4	96.9	3,550
Fourth	1.5	96.6	1.2	0.0	1.4	0.2	0.8	0.2	0.0	0.2	4,253	98.1	98.0	4,226
Highest	1.3	96.3	4.7	0.6	0.6	0.1	0.2	0.2	0.0	0.5	4,731	98.6	97.9	4,697
Total	1.5	94.2	1.6	0.1	3.6	0.2	0.6	1.0	0.0	0.5	18,548	97.6	96.6	18,439

[1] Reusable sanitary pads, disposable sanitary pads, tampons, menstrual cup, cloth, and/or cotton wool

[2] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 13.9C Menstrual hygiene: County level

Among women age 15–49 whose most recent menstrual period was in the last year, percentage who used specified materials to collect or absorb blood from the most recent menstrual period; and among women age 15–49 whose most recent menstrual period was in the last year and were at home during their last menstrual period, percentage who were able to wash and change in privacy while at home and percentage who were able to both wash and change in privacy and who used appropriate materials during their last menstruation, according to county, Kenya Mini DHS 2025–26

	Among women whose most recent menstrual period was in the last year, percentage who used the specified materials to collect or absorb blood from most recent menstrual period											Among women whose most recent menstrual period was in the last year and who were at home during their last menstrual period:		
County	Re-usable sanitary pads	Dis-posable sanitary pads	Tam-pons	Men-strual cup	Cloth	Toilet paper	Cotton wool	Under-wear only	Other	Nothing	Number of women	Percent-age able to wash and change in privacy	Percent-age able to wash and change in privacy and who used appro-priate materials during last men-struation[1]	Number of women
Mombasa	0.3	98.0	1.9	0.0	2.4	0.1	1.3	0.0	0.0	0.0	753	97.1	97.1	744
Kwale	0.2	84.2	0.0	0.2	19.7	0.2	1.1	0.1	0.0	0.4	321	99.6	99.0	321
Kilifi	1.4	94.1	0.3	0.0	3.6	0.0	0.7	0.2	0.0	0.0	554	99.4	99.2	554
Tana River	1.9	86.5	0.0	0.0	11.7	0.0	0.4	0.9	0.0	0.0	118	100.0	99.1	118
Lamu	1.5	98.3	1.2	0.0	0.3	0.0	0.0	0.0	0.0	0.3	48	99.2	99.0	48
Taita Taveta	0.8	96.5	0.4	0.0	3.1	0.0	0.3	0.0	0.0	0.0	135	99.6	99.6	135
Garissa	0.6	63.6	0.0	0.0	42.5	0.3	0.0	6.7	0.0	1.3	140	98.8	92.7	140
Wajir	0.3	74.8	0.1	0.0	28.8	0.0	0.0	1.3	0.0	2.3	112	96.7	93.0	112
Mandera	3.0	46.3	0.6	0.0	55.3	0.0	2.7	3.4	0.0	0.0	112	57.6	57.4	111
Marsabit	0.4	94.1	0.0	0.0	8.0	0.0	0.0	2.9	0.0	0.0	120	98.3	97.4	118
Isiolo	3.1	93.1	0.0	0.0	4.3	0.0	0.0	1.5	0.0	0.4	62	97.5	96.5	62
Meru	0.6	97.9	1.7	0.0	0.0	0.0	1.1	0.8	0.0	0.0	460	93.3	92.5	453
Tharaka-Nithi	0.5	96.9	1.1	0.0	1.0	0.2	1.3	0.0	0.0	1.4	129	94.0	92.5	129
Embu	2.0	97.6	2.4	0.0	0.5	0.0	0.0	0.0	0.0	0.1	209	98.4	98.3	209
Kitui	1.0	98.5	0.9	0.0	0.3	0.3	0.0	0.0	0.0	0.2	365	99.1	99.1	362
Machakos	2.3	95.3	0.8	0.0	0.2	2.0	1.2	0.4	0.2	0.4	579	97.7	96.8	579
Makueni	0.0	98.5	3.5	0.2	1.6	0.0	0.3	0.0	0.0	0.0	280	98.9	98.9	270
Nyandarua	0.5	98.5	0.3	0.0	0.9	0.0	0.0	0.0	0.0	0.6	203	97.2	96.5	203
Nyeri	0.4	97.5	1.9	0.0	0.0	0.5	0.8	0.0	0.0	0.0	197	97.2	97.2	197
Kirinyaga	5.2	94.5	0.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	224	99.6	99.6	222
Murang'a	0.9	98.2	2.5	0.0	0.7	0.7	1.0	0.0	0.0	0.0	325	100.0	100.0	325
Kiambu	0.0	96.9	5.1	0.5	0.2	0.0	0.6	0.0	0.0	0.2	1,569	98.3	98.3	1,556
Turkana	4.2	86.0	0.6	0.0	9.4	0.0	0.0	6.5	0.0	0.8	110	96.5	95.3	108
West Pokot	0.2	93.9	0.0	0.0	0.4	0.0	0.0	5.7	0.0	0.8	173	97.2	91.5	173
Samburu	2.2	90.2	0.4	0.0	3.3	0.0	0.0	4.5	0.0	0.4	70	95.4	92.4	70
Trans Nzoia	2.4	97.5	0.0	0.0	0.6	0.0	0.0	0.6	0.0	0.0	315	99.6	99.0	314
Uasin Gishu	0.5	98.6	1.4	0.0	2.5	0.0	0.0	1.4	0.0	0.0	595	98.5	98.2	585
Elgeyo-Marakwet	2.7	93.9	0.7	0.3	4.5	0.6	0.0	1.2	0.0	0.0	152	98.0	97.0	151
Nandi	1.4	96.0	0.6	0.0	3.5	0.0	0.6	0.0	0.0	0.5	304	87.5	87.0	301
Baringo	3.1	87.6	0.0	0.0	9.3	0.0	2.4	3.6	0.0	1.9	162	97.1	95.5	162
Laikipia	51.8	45.7	0.0	0.0	1.8	0.0	1.1	0.0	0.0	0.0	165	99.1	99.1	165
Nakuru	0.9	95.8	1.3	0.0	2.1	0.0	0.4	0.4	0.0	0.6	946	97.6	96.5	946
Narok	0.0	93.6	0.9	0.2	4.8	1.7	0.0	2.9	0.3	0.5	364	91.7	90.3	362
Kajiado	0.5	96.9	3.4	0.0	1.3	0.0	0.2	0.7	0.0	0.0	405	97.8	97.4	404
Kericho	0.9	96.9	0.2	0.0	1.7	0.0	0.3	1.1	0.0	0.3	348	99.5	98.1	346
Bomet	0.0	94.8	0.0	0.0	6.1	0.0	0.4	2.2	0.0	0.0	395	100.0	98.1	395
Kakamega	0.8	89.8	0.5	0.0	6.4	1.1	0.6	3.6	0.0	1.9	535	99.4	94.3	532
Vihiga	1.7	93.8	0.0	0.0	4.9	0.0	0.7	1.0	0.0	0.3	197	99.8	98.8	197
Bungoma	0.4	90.6	0.3	0.0	5.0	0.0	0.9	5.1	0.2	2.0	564	99.7	94.3	561
Busia	1.3	94.7	0.6	0.0	3.4	0.0	0.0	0.5	0.0	0.3	329	99.4	98.6	328
Siaya	0.9	97.5	0.0	0.0	1.1	0.0	0.0	0.0	0.0	0.8	316	97.8	97.5	314
Kisumu	3.8	94.8	0.5	0.0	6.3	1.1	1.3	2.0	0.0	0.0	489	95.9	95.5	471
Homa Bay	2.0	94.7	0.2	0.1	7.4	0.2	0.9	1.0	0.0	1.1	431	98.7	97.0	431
Migori	1.1	95.4	0.3	0.0	3.4	0.0	0.5	1.1	0.0	0.0	418	97.7	97.4	413
Kisii	0.3	93.3	0.0	0.0	3.1	0.0	1.5	1.5	0.0	0.3	379	97.6	96.1	378
Nyamira	0.8	90.2	0.6	0.0	7.8	0.0	0.9	2.1	0.0	0.1	198	88.9	87.7	198
Nairobi City	1.0	96.1	3.1	0.5	1.4	0.2	0.8	0.2	0.0	0.7	3,173	98.3	97.6	3,166
Total	1.5	94.2	1.6	0.1	3.6	0.2	0.6	1.0	0.0	0.5	18,548	97.6	96.6	18,439

[1] Reusable sanitary pads, disposable sanitary pads, tampons, menstrual cup, cloth, and/or cotton wool

GENDER-BASED VIOLENCE 14

Key Findings

- ***Experience of technology-facilitated gender-based violence (TFGBV) from any perpetrator:*** 26% of women age 15–49 have ever experienced TFGBV from any perpetrator.
- ***Experience of various forms of TFGBV:*** 19% of women age 15–49 reported ever experiencing online control. Other common forms of TFGBV ever experienced include online harassment (3%) and hacking (5%).
- ***Experience of TFGBV by any intimate partner:*** 27% of women age 15–49 who have ever had a husband or intimate partner have ever experienced TFGBV committed by a current, most recent, or previous husband or intimate partner; 22% experienced TFGBV in the 12 months preceding the survey.
- ***Perpetrators of TFGBV:*** Among women age 15–49 who have ever been married or have ever had an intimate partner, the most common perpetrator of TFGBV is a current husband or intimate partner (61%), followed by a former husband or intimate partner (22%). Among never-married women and those who have never had an intimate partner, the most common perpetrators of TFGBV are strangers (45%), online attackers (23%), and friends/acquaintances (20%).

Gender-based violence (GBV) against women is widely recognised as a violation of human rights and a major public health concern due to its health, social, and demographic consequences. As access to digital technologies has expanded, new forms of violence have emerged through the use of online platforms and communication technologies. Technology-facilitated gender-based violence (TFGBV) is defined as an act of violence perpetrated by one or more individuals that is committed, assisted, aggravated, and amplified in part or fully by the use of information and communication technologies or digital media against a person on the basis of gender (United Nations Population Fund [UNFPA] 2021; UNFPA n.d.; UN Women 2025). TFGBV can include online harassment, stalking, exploitation, and other forms of abuse that may contribute to or exacerbate offline violence.

Kenya has established a range of legal and policy frameworks to address GBV, including cyber-enabled forms of violence, and has ratified several international agreements aimed at protecting women and girls from violence and discrimination. Despite these efforts, concerns remain regarding the prevalence and consequences of TFGBV and the need for evidence to inform prevention and response strategies. The 2025 *Report of the Technical Working Group on GBV Including Femicide* (Republic of Kenya 2025) indicated that many cases of homicide of women and girls, and of sexual and physical harm directed towards women and young girls, are preceded by online and offline harassment using digital tools. Nationally representative data are therefore important for monitoring the extent of TFGBV and assessing perceptions of safety and risk among women and girls.

Historically, The Demographic and Health Surveys (DHS) Program has only collected detailed information on intimate partner violence experienced by ever-married women, defined as women who are currently married or living with a man as if married and women who were formerly married or lived with a man as if married. However, for the 2025–26 Kenya Mini DHS (KMDHS), a domestic violence questionnaire module was revised to focus on experiences of TFGBV among women age 15–49.

The module of questions on TFGBV was administered in accordance with the World Health Organization's (WHO's) guidelines on the ethical collection of information on GBV (WHO 2001). Among the 22,507 women age 15–49 interviewed in the 2025–26 KMDHS, only one woman per household was eligible to complete the module on TFGBV. Among the eligible women, 15,614 (69%) were selected and interviewed; 951 women (4%) were selected but not interviewed due to lack of privacy or for other reasons. Specifically constructed weights were used to adjust for the selection of only one respondent per household and to ensure that the subsample was nationally representative.

This chapter presents findings for women age 15–49 on their experience of different forms of TFGBV. The different forms were purposively drawn based on the specific harm and impact they have on survivors as guided by the UNFPA framework (UNFPA n.d.). Sextortion (blackmailing with intimate images), online grooming, and image-based sexual abuse exploit women and girls; doxxing and online impersonation can damage reputations and cause social harm; cyberstalking, gendered hate speech, and online mobbing cause psychological and emotional harm; tracking and controlling a person's digital devices cause economic and livelihood harm; and nonconsensual exposure to surveillance threatens the physical safety of women and girls.

The chapter also provides information on forms of controlling behaviours, and on the perceived risks of violence and safety among women and girls.

14.1 Measurement of Technology-facilitated Gender-based Violence

Terminology for this chapter

Husband: A man with whom a woman is married or living with as if married.

Intimate partner: A man with whom a never-married woman is in a relationship that involves physical and/or emotional intimacy and for which the relationship is or has the expectation of being long lasting. For this chapter, an intimate partner is not a husband or a man a woman is living with, a boyfriend with whom her relationship is casual, or a man with whom she has a one-time encounter.

Husband/intimate partner: The current husband for currently married women; the most recent husband for divorced, separated, or widowed women; the current intimate partner for never-married women who currently have an intimate partner; and the most recent intimate partner for never-married women who do not currently have an intimate partner but had one in the past.

Boyfriend: A man with whom the woman has a causal relationship and whom she did not mention as an intimate partner

Technology-facilitated gender-based violence: An act of violence perpetrated by one or more individuals that is committed, assisted, aggravated, and amplified in part or fully by the use of information and communication technologies or digital media against a person on the basis of gender (UNFPA 2023). In this survey, TFGBV included one or more of the following actions from any partner:

a. **Revenge pornography:** Shared your sexually explicit messages, images, or videos with other people
b. **Online harassment:** Sent you threatening messages via the internet, text messages, instant messages, social media, or other digital tools/digital platforms or called you repeatedly to harass you
c. **Doxxing:** Shared your private or personal information (address, phone number, email) with malicious intent
d. **Blackmail/sextortion:** Threatened to share sensitive information, images, audio, or videos unless certain financial, sexual, or emotional demands were met
e. **Hacking:** Gained unauthorised access to your personal devices, social media, or emails to track, control, or harm you
f. **Cyberbullying:** Used digital tools (internet, text messages, instant messages, phone calls, social media, or any other digital tools) to spread false, distorted, or exaggerated information about you or publicly humiliate you
g. **Cyberstalking:** Used technology or digital devices to monitor and stalk you in real time or historically with malicious intent
h. **Online sexual harassment or cyber sexual abuse:** Sent you unwanted sexually explicit messages, images, audio, or videos
i. **Online control:** One or more of the following actions from any intimate partner:
    - Insists/insisted on reading call logs/messages/emails from your phone/tablet/laptop or other digital tools without your consent
    - Insists/insisted you give him passwords, pins, or patterns for your phone, tablet, laptop, email, Facebook, Instagram, WhatsApp, Snapchat, or other digital platforms
    - Uses/used mobile, internet, or other technology including phone calls, Facetime, video calling, or tracking software to check on you or track where you are/were in ways that make/made you feel controlled

In the 2025–26 KMDHS, information was obtained from women age 15–49 on their experience of TFGBV committed by any perpetrator, including a current and former spouse or other intimate partner. To capture intimate-partner TGFBV, ever-married women were asked about their experience of violence committed by their current and former spouses/live-in partners and, if applicable, never-married women were asked about their experience of violence committed by their current and former intimate partners. More specifically, intimate-partner violence was measured by asking women if their current or former spouse/intimate partner ever perpetrated any of the actions outlined above (a–i).

14.2 Experience of Technology-facilitated Gender-based Violence by Any Perpetrator

Technology-facilitated gender-based violence by any perpetrator

Percentage of women who have experienced any TFGBV committed by a husband, intimate partner, or anyone else in their lifetime and in the 12 months before the survey.

Sample: Women age 15–49


Twenty-six percent of women age 15–49 have ever experienced TGFBV from any perpetrator. Twenty-one percent have experienced such violence often or sometimes in the 12 months preceding the survey (**Table 14.1**).

Patterns by background characteristics

- The percentage of women who have ever experienced TFGBV is lowest among women age 15–19 (10%) and highest among women age 20–24 (37%).
- Forty-one percent of divorced, separated, or widowed women have ever experienced TFGBV, compared with 27% of women who are married or currently living with a partner and 19% of never-married women.
- At the county level, the percentage of women who have ever experienced TFGBV from any perpetrator is highest in Nairobi City (33%), followed by Murang'a and Baringo (32% each) (**Table 14.1C** and **Map 14.1**).
- The experience of TFGBV among women in the 12 months preceding the survey was most prevalent in Trans Nzoia County (29%), followed by Machakos and Kericho Counties (28% each).

Map 14.1 Experience of various forms of TFGBV by any perpetrator by county

Percentage of women age 15–49 who have ever experienced TFGBV by any perpetrator



County Code	County	%
13	Tharaka-Nithi	29
14	Embu	27
16	Machakos	31
17	Makueni	22
18	Nyandarua	31
19	Nyeri	31
20	Kirinyaga	30
21	Murang'a	32
22	Kiambu	29
27	Uasin Gishu	31
28	Elgeyo-Marakwet	10
29	Nandi	13
32	Nakuru	25
35	Kericho	30
36	Bomet	25
37	Kakamega	25
38	Vihiga	21
42	Kisumu	31
45	Kisii	17
46	Nyamira	21
47	Nairobi	33

14.2.1 Experience of Various Forms of Technology-facilitated Gender-based Violence by Any Perpetrator

The most common form of TFGBV ever experienced by women age 15–49 by any perpetrator is online control (19%), followed by online harassment (5%), hacking (5%), and online sexual harassment/sexual abuse (3%) (**Figure 14.1** and **Table 14.2**).


Online control (17%) and online harassment (3%) are the most common forms of TFGBV that women age 15–49 experienced in the 12 months preceding the survey.

Patterns by background characteristics

- Hacking is especially pronounced among women age 20–29 (8%) and among women with more than a secondary education (5%), while online harassment is most common among divorced, separated, or widowed women (7%) and among never-married women who have ever had an intimate partner (6%) (**Table 14.3**).
- At the county level, any form of TFGBV is highest in Trans Nzoia (29%), followed by Kericho and Machakos (28% each), and lowest in Wajir and Mandera (2% each). Online control is the most common form of TFGBV across all counties, reaching as high as 25% in Machakos. The next most common forms vary by county: Online harassment is reported in Bomet (8%), while online sexual harassment or cyber sexual abuse is reported in Nyeri (8%) (**Table 14.3C**).

Figure 14.1 **Experience of various forms of technology-facilitated gender-based violence by any perpetrator**

Percentage of women age 15–49 who have ever experienced technology facilitated gender-based violence by any perpetrator



14.2.2 Experience of Technology-facilitated Gender-based Violence by Any Intimate Partner

Technology-facilitated gender-based violence by any intimate partner

Percentage of women who have experienced any TFGBV committed by a current, most recent, or previous husband or intimate partner ever and in the 12 months before the survey

Sample: Women age 15–49

Twenty-seven percent of women age 15–49 who have ever had a husband or intimate partner have ever experienced TFGBV committed by a current, most recent, or previous husband or intimate partner, while 22% had experienced such violence in the 12 months preceding the survey (**Table 14.4**).

Patterns by background characteristics

- Among women who have ever had a husband or intimate partner, the percentage who have ever experienced TFGBV committed by an intimate partner is higher among women age 20–24 (37%) and those age 25–29 (33%) than among younger women age 15–19 (22%), women age 30–39 (25%), or those age 40–49 (17%)**.**
- The percentage of women who have ever experienced TFGBV by any intimate partner is highest in Homa Bay (36%) and Baringo (35%) Counties (**Table 14.4C**).
- The percentage of women who experienced TFGBV by any intimate partner in the 12 months preceding the survey was highest in Trans Nzoia (31%) County and in Tharaka-Nithi, Machakos, and Busia Counties (30% each) and lowest in Mandera and Wajir Counties (3% each).

14.2.3 Perpetrators of Technology-facilitated Gender-Based Violence

The most commonly reported perpetrators of TFGBV among women who have ever been married or have ever had an intimate partner are current husbands or intimate partners (61%), former husbands or intimate partners (22%), and strangers (13%) (**Table 14.5**).

Among never-married women and those who have never had an intimate partner, the most common perpetrators of TFGBV are strangers (45%), online attackers (23%), and friends or acquaintances (20%).

14.3 Forms of Controlling Behaviours

Controlling behaviour

Percentage of women whose current or most recent husband or intimate partner demonstrates one or more of the following controlling behaviours: is jealous or angry if she talks to other men; wrongly accuses her of being unfaithful; does not permit her to meet her female friends; tries to limit her contact with her family; insists on knowing where she is at all times; insists on reading call logs/messages/emails from her phone/tablet/laptop or other digital tools without her consent; insists on her giving him passwords, pins, or patterns for her phone, tablet, laptop, email, or digital platforms; and uses mobile, internet, or other technology including phone calls, Facetime, video calling, or tracking software to check on her in ways that makes her feel controlled.

Sample: Women age 15–49 who have ever had a husband or an intimate partner

Prevalence of Controlling Behaviours

Controlling behaviours exhibited by a husband or male intimate partner towards his wife or intimate partner are important early warning signs that may escalate into violence within a relationship. Given that the concentration of such behaviours carries greater significance than any single act, attention is also given to the percentage of women whose partners display at least three of the behaviours specified above.

Among women age 15–49 who have ever had a husband or intimate partner, 61% have ever experienced any controlling behaviours from their current or most recent husband or intimate partner, and 54% experienced any controlling behaviours in the 12 months preceding the survey. Twenty-seven percent of the women have ever experienced at least three forms of controlling behaviours, including 24% who experienced them in the past 12 months (**Tables 14.6** and **14.7**).


Fifty percent of women age 15–49 report that their current or most recent husband or intimate partner has ever been jealous or angry if they talked with other men, and 36% report that their current or most recent husband or intimate partner has ever insisted on knowing where they are at all times (**Figure 14.2**).

Patterns by background characteristics

- Thirty-four percent of women age 20–24 report that their husband or intimate partner has ever displayed three or more controlling behaviours, compared with 22% of women age 15–19 and 22% of those age 40–49 (**Table 14.6**).
- Thirty-nine percent of women who are divorced, separated, or widowed report that their husband or intimate partner has exhibited three or more controlling behaviours, compared with 24% of women who are married or living with their partner. Similarly, 28% of women who have never been married but have had an intimate partner report that their intimate partner has ever displayed three or more controlling behaviours.

***Figure 14.2* Forms of controlling behaviours**

Percentage of women age 15–49 who have ever had a husband or intimate partner who have experienced controlling behaviours



- At the county level, the highest percentages of women who report that their husband or intimate partner has ever displayed three or more controlling behaviours are in Kwale, Homa Bay, and Trans Nzoia (39% each); Tharaka-Nithi and Vihiga (38% each); and Kisumu (37%). In contrast, the lowest percentages of women who report that their husband or intimate partner has ever displayed three or more controlling behaviours are in Wajir and Mandera (4% each) (**Table 14.7C**).

14.4 Experience of Technology-facilitated Gender-based Violence by Non-intimate Partners According to Platform

Experience of technology-facilitated gender-based violence according to platform

Percentage of women who have experienced TFGBV from a non-intimate partner ever and in the 12 months preceding the survey by platform.

Note: Online control questions were excluded in the calculation of this indicator due to a skip error.


Social media: Facebook, Instagram, and other social media platforms (for example, Threads, TikTok, YouTube, reddit)

Messaging apps: Text message/iMessage, WhatsApp, and other messaging apps (for example, Signal, Telegram)

Sample: Women age 15–49 who have experienced TFGBV committed by any perpetrator

The lifetime experience of TFGBV among women age 15–49 varies according to platform, with social media and messaging apps emerging as the most common channels. Overall, 67% of women report having ever experienced TFGBV on social media, with Facebook reported as the most common platform (59%). Messaging apps are the next most common source (47%); text message/iMessage (27%) and WhatsApp (26%) are the most popular, with only 2% of women reporting experiencing TFGBV through other messaging apps (**Figure 4.3** and **Table 14.8**).

Figure 14.3 **Experience of technology-facilitated gender-based violence by platform**



Patterns by background characteristics

- Women's experience of TFGBV varies by age and platform, with younger women reporting higher exposure, particularly on social media. Seventy-seven percent of younger women age 15–19 and 72% of those age 20–24 report experiencing TFGBV on any social media platform, compared with 50% of women age 40–49. However, the experience of TFGBV through messaging apps is most common among women age 40–49 (54%) and least common among those age 15–19 (30%) (**Table 14.9**).

- Exposure to TFGBV on social media platforms increases with wealth, from 45% among women in the lowest wealth quintile to 74% among those in the highest wealth quintile. In contrast, women in the lowest wealth quintile report greater exposure to TFGBV through messaging apps (62%) than do those in the highest wealth quintile (39%).

14.5 Community Perceptions on Violence against Women and Girls

Information on perceptions of harm against women and girls were collected through a structured survey module administered to individual respondents at the community level. The target respondents were any

eligible adults selected within sampled households, who were then asked to reflect on prevailing attitudes, beliefs, and practices within their community. The questions were intentionally framed to capture community-level perceptions rather than solely personal experiences, thereby providing insight into social norms and collective behaviours.

The module consisted of a set of standardised questions designed to assess perceptions related to women's safety and exposure to violence. The questions covered the following key thematic areas:

- Perceived groups of women and girls at heightened risk of violence
- Perceived levels of safety for women and girls within the community
- Expected community responses to suspected risks of severe violence
- Prevailing attitudes regarding the acceptability of violence, including victim blaming and norms surrounding domestic violence as a private or public issue

14.5.1 Perceived Risks of Violence and Safety for Women and Girls

Perceived risks of violence and safety for women and girls

Percentage distribution of household respondents who perceive certain groups of women and girls to be at risk of violence; and

Percentage distribution of households by perception of safety of women and girls in the community according to background characteristics.

Sample: All sampled households

Household respondents were asked whether certain groups of women or girls are at a higher risk of violence than other groups. Eighteen percent of the household respondents report perceiving that certain groups of women and girls are at risk of violence, although perceptions of risk vary according to background characteristics.

Household respondents were also asked about how safe they feel women and girls are within their community. Ninety percent of household respondents perceive that women and girls are safe, somewhat safe, or very safe in the community (**Table 14.10**).

Patterns by background characteristics

- Nineteen percent of female household respondents perceive that certain groups of women and girls are at risk of violence, compared with 17% of their male counterparts. In terms of household perceptions on the safety of women and girls in the community, a higher percentage of male respondents (13%) than female respondents (13%) report that women and girls are 'very safe', and a higher percentage of female respondents (11%) than male respondents report that women and girls are 'not very safe' (**Table 14.10**).
- The perception that certain groups of women and girls are at a higher risk of violence is more common among household respondents with more than a secondary education (21%) than among those with no education (17%).
- Perceived risk of violence against women and girls is most common in Turkana (39%), Bungoma (38%), Homa Bay (37%), and Narok (36%) Counties and lowest in Kilifi, Baringo, Kitui, Wajir, Kwale, Vihiga

and Taita Taveta Counties (ranging from 4% to 7% each), indicating wide geographic disparities in perceived vulnerability (**Table 14.10C**).

- Perceptions of safety also vary by county. Mandera (53%), Nandi (41%), and Wajir (34%) Counties have the highest percentages of household respondents indicating that women and girls are 'very safe' in the community.

14.5.2 Perceived Community Response to Killing of Women and Girls

Household respondents were also asked about what action they would take if they suspected that a woman or girl was at risk of being killed. Most household respondents (60%) said they would report to National Government Administrative Officers (NGAOs) if they suspected a woman or a girl was at a high risk of being killed, while 48% indicated that they would report it to the police. Only 6% of household respondents indicated that they would do nothing (**Table 14.11**).


Patterns by background characteristics

- If they suspected a woman or girl was at risk of being killed, most older household respondents would report it to NGAOs (ranging from 56% among those age 30–34 to 75% among those age 65 and older), while most younger respondents prefer to report it to the police (ranging from 56% among those age 15–19 to 52% among those age 25–29) (**Table 14.11**).

- The percentage of household respondents who said they would take no action if they suspected that a woman or girl was at risk of being killed is generally low across background characteristics. However, the percentage indicating they would do nothing increases with both wealth (**Figure 14.4**) and level of education. Household respondents in the highest wealth quintile (9%) and those with more than a secondary education (8%) are more likely to take no action than are household respondents in the lowest wealth quintile (4%) and those with no education (4%).

***Figure 14.4* Perceived community response to femicide by household wealth**

Percentage distribution of household respondents according to action they would take if they suspected that a woman or a girl was at risk of being killed



- More than 90% of people in Marsabit, Baringo, and Tana River Counties prefer reporting to NGAOs. Preference for reporting to the police is most prevalent in Taita/Taveta County (74%), while preference for mobilising the community to rescue a woman or girl at risk of being killed is most common in Mandera (69%) and Murang'a (64%) Counties (**Table 14.11C**).

- The percentage of household respondents who said they would take no action if they suspected a woman or girl was at risk of being killed is generally low across counties. However, the highest percentages are in Samburu and Nyamira (14% each), followed by Meru (12%) and then Machakos, Nairobi City, and Kakamega (11% each). In contrast, the percentage reporting that they would do nothing is extremely low

in several counties, including Marsabit, Tana River, Wajir, Mandera, Turkana, and Elgeyo-Marakwet (<1% each).

14.5.3 Justification for Violence against Women or Girls

Household respondents were asked whether any social, cultural, or religious beliefs justify violence against women or girls. Eight percent of household respondents believe that some existing socio-cultural and religious beliefs justify such violence (**Table 14.12**).

At the county level, the percentage of household respondents who believe that socio-cultural or religious beliefs justify violence against women and girls varies widely. The highest percentages are in Samburu (25%), Migori (24%), and Mandera (22%), and the lowest percentages are in Wajir (<1%), Kitui (2%), and Machakos (2%) (**Table 14.12C**).

14.5.4 Perceptions of Victim Blaming and Privacy of Domestic Violence

Victim blaming

Victim blaming occurs when service providers or community members make excuses (verbally or otherwise, and intentionally or unintentionally) that tend to shift the responsibility of domestic violence from the perpetrator to the survivor or victim. Such action may be subtle or loud and often includes statements like 'She provoked the perpetrator' or 'Why was she loitering at night?'. Victim blaming causes psychologic harm to survivors and discourages them from reporting.

Household respondents were asked whether they have heard of people blaming women or girls for the violence they experience, Overall, 36% of household respondents have heard of people blaming the victim.

Household respondents were also asked whether they believe domestic violence is a private matter, and 38% of household respondents believe that it is (**Table 14.13**).

Patterns by background characteristics

- Awareness of victim blaming is slightly more common among female respondents than male respondents (37% versus 34%), but men are more likely than women to view domestic violence as a private matter (41% versus 35%).
- Awareness of victim blaming generally increases with age, from 29% among those age 15–19 to 40% among those age 50–54, before declining somewhat at older ages. The perception that domestic violence is a private matter is least common among younger respondents age 15–19 (32%) and most common among respondents age 60–64 and those age 65 or older (41% each).
- Awareness of victim blaming increases with level of education, from 28% among those with no education to 41% among those with more than a secondary education. In contrast, the belief that domestic violence is a private matter generally declines with level of education, from 40% among those with no education and 41% among those with a primary education to 28% among those with more than a secondary education.
- A similar inverse pattern appears by wealth. Household respondents in the highest wealth quintile are more likely to have heard of victim blaming (38%) but less likely to say domestic violence is private (31%) than are household respondents in the lowest wealth quintile (32% and 41%, respectively).

- At the county level, the percentages of household respondents who are aware of victim blaming are highest in Trans Nzoia (59%), Samburu (57%), Kisii (54%), and Turkana (51%) and lowest in Wajir (1%), Tana River (7%), and Baringo (7%) (**Table 14.13C**). The percentage of household respondents who perceive domestic violence to be a private matter varies widely across counties. The highest percentages are in Mandera (74%), Tharaka-Nithi (68%), Nyamira (62%), and Kwale (61%), and the lowest percentages are in Marsabit (5%), Tana River (6%), and Kitui (13%).

List of Tables

For more information on gender-based violence, see the following tables:

- **Table 14.1 Experience of technology-facilitated gender-based violence by any perpetrator**
- **Table 14.1C Experience of technology-facilitated gender-based violence by any perpetrator: County level**
- **Table 14.2 Experience of various forms of technology-facilitated gender-based violence by any perpetrator**
- **Table 14.3 Experience of various forms of technology-facilitated gender-based violence by any perpetrator according to background characteristics**
- **Table 14.3C Experience of various forms of technology-facilitated gender-based violence by any perpetrator: County level**
- **Table 14.4 Experience of technology-facilitated gender-based violence by any intimate partner**
- **Table 14.4C Experience of technology-facilitated gender-based violence by any intimate partner: County level**
- **Table 14.5 Persons committing technology-facilitated gender-based violence**
- **Table 14.6 Forms of controlling behaviours**
- **Table 14.7 Controlling behaviours of husband/intimate partner by background characteristics**
- **Table 14.7C Controlling behaviours of husband/intimate partner: County level**
- **Table 14.8 Experience of technology-facilitated gender-based violence by platform**
- **Table 14.9 Experience of technology-facilitated gender-based violence by platform and background characteristics**
- **Table 14.10 Perceived risk of violence and safety for women and girls**
- **Table 14.10C Perceived risk of violence and safety for women and girls: County level**
- **Table 14.11 Perceived community response to femicide**
- **Table 14.11C Perceived community response to femicide: County level**
- **Table 14.12 Perceived socio-cultural and religious beliefs on justification of violence against women and girls**
- **Table 14.12C Perceived socio-cultural and religious beliefs on justification of violence against women and girls: County level**
- **Table 14.13 Perceptions of victim blaming and privacy of domestic violence**
- **Table 14.13C Perceptions of victim blaming and privacy of domestic violence: County level**

Table 14.1 Experience of technology-facilitated gender-based violence by any perpetrator

Percentage of women age 15–49 who have ever experienced technology-facilitated gender-based violence (TFGBV) by any perpetrator since age 15 and percentage who have experienced TFGBV by any perpetrator in the 12 months preceding the survey, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Percentage of women who have experienced TFGBV by any perpetrator	Percentage who have experienced TFGBV in the past 12 months			Number of women
	Ever[1]	Often	Sometimes	Often or sometimes	
Age					
15–19	10.3	4.0	4.6	8.6	2,866
20–24	36.7	15.2	17.4	32.7	2,692
25–29	35.4	15.3	14.6	29.9	2,746
30–39	27.8	11.3	10.3	21.6	4,496
40–49	19.6	5.3	7.0	12.2	2,815
Residence					
Urban	30.1	10.7	12.3	22.9	7,096
Rural	22.6	9.9	9.3	19.2	8,518
Marital status					
Never married	19.2	6.1	9.6	15.7	5,099
Never had intimate partner	4.1	0.6	1.9	2.5	2,753
Ever had intimate partner	36.9	12.5	18.6	31.1	2,346
Ever married	29.3	12.3	11.2	23.4	10,515
Married/living together	26.8	11.4	11.3	22.7	8,612
Divorced/separated/ widowed	40.5	16.2	10.5	26.8	1,903
Education[2]					
No education	8.4	4.6	2.4	7.0	822
Primary	24.6	11.2	8.4	19.6	4,508
Secondary	26.2	10.1	11.0	21.1	7,231
More than secondary	32.2	10.7	15.3	26.0	3,054
Wealth quintile					
Lowest	16.1	7.9	6.5	14.3	2,477
Second	24.1	11.4	9.1	20.4	2,881
Middle	26.2	11.2	10.2	21.4	3,053
Fourth	30.6	12.2	13.4	25.6	3,508
Highest	29.6	8.3	12.4	20.7	3,695
Total	26.0	10.3	10.6	20.9	15,614

[1] Includes TFGBV in the past 12 months. For women who were married or living together before age 15 and reported violence only by their husband, and for never-married women who had an intimate partner before age 15 and reported violence only by their intimate partner, the violence could have occurred before age 15.

[2] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 14.1C Experience of technology-facilitated gender-based violence by any perpetrator: County level

Percentage of women age 15–49 who have ever experienced technology-facilitated gender-based violence (TFGBV) by any perpetrator since age 15 and percentage who have experienced TFGBV by any perpetrator in the 12 months preceding the survey, according to county, Kenya Mini DHS 2025–26

County	Percentage of women who have experienced TFGBV by any perpetrator	Percentage who have experienced TFGBV in the past 12 months			Number of women
	Ever[1]	Often	Sometimes	Often or sometimes	
Mombasa	18.7	9.8	7.6	17.5	556
Kwale	28.4	14.4	8.4	22.8	253
Kilifi	11.8	8.1	3.2	11.3	469
Tana River	13.9	1.2	10.4	11.6	96
Lamu	7.9	2.8	3.0	5.8	41
Taita Taveta	28.0	9.3	12.6	21.9	107
Garissa	7.4	2.9	4.1	7.0	127
Wajir	1.9	0.8	1.0	1.9	94
Mandera	1.9	1.4	0.6	1.9	96
Marsabit	10.9	3.7	4.7	8.5	98
Isiolo	14.7	5.8	4.1	9.9	58
Meru	21.9	5.7	10.5	16.1	388
Tharaka-Nithi	29.4	17.3	8.9	26.1	100
Embu	27.0	14.1	11.9	26.0	179
Kitui	24.0	10.2	11.9	22.1	319
Machakos	31.4	16.9	10.9	27.8	517
Makueni	21.9	5.2	11.1	16.3	259
Nyandarua	30.8	6.5	14.8	21.3	148
Nyeri	30.5	16.2	9.9	26.2	152
Kirinyaga	30.2	8.1	11.7	19.8	197
Murang'a	31.9	4.2	17.3	21.5	285
Kiambu	29.0	11.0	11.4	22.4	1,316
Turkana	28.1	11.3	9.7	21.0	134
West Pokot	13.4	4.3	9.1	13.4	172
Samburu	27.6	11.4	13.1	24.5	74
Trans Nzoia	30.7	19.6	9.0	28.6	289
Uasin Gishu	31.3	12.3	14.7	27.0	498
Elgeyo-Marakwet	10.0	6.3	3.5	9.8	133
Nandi	13.3	3.3	7.8	11.1	269
Baringo	32.1	8.0	16.7	24.6	131
Laikipia	24.7	16.9	5.3	22.2	142
Nakuru	25.0	9.8	9.8	19.6	809
Narok	26.4	14.0	9.4	23.4	334
Kajiado	26.2	13.5	11.0	24.6	325
Kericho	30.1	11.0	17.0	28.1	296
Bomet	24.9	5.2	16.7	21.9	349
Kakamega	24.9	13.1	9.2	22.3	501
Vihiga	20.9	3.4	9.0	12.4	169
Bungoma	27.6	13.9	11.0	24.9	489
Busia	25.9	19.3	4.9	24.2	275
Siaya	19.9	7.8	12.1	19.9	282
Kisumu	30.7	15.4	11.1	26.5	408
Homa Bay	29.6	15.3	7.7	23.0	382
Migori	27.0	12.4	10.9	23.3	383
Kisii	17.4	11.3	5.6	16.9	324
Nyamira	21.2	15.7	4.1	19.8	180
Nairobi City	33.3	7.1	14.1	21.2	2,412
Total	26.0	10.3	10.6	20.9	15,614

[1] Includes TFGBV in the past 12 months. For women who were married or living together before age 15 and reported violence only by their husband, and for never-married women who had an intimate partner before age 15 and reported violence only by their intimate partner, the violence could have occurred before age 15.

Table 14.2 Experience of various forms of technology-facilitated gender-based violence by any perpetrator

Percentage of women age 15–49 who have ever experienced technology-facilitated gender-based violence (TFGBV) by any perpetrator and percentage of women who have experienced TFGBV by any perpetrator in the 12 months preceding the survey, according to form of TFGBV, Kenya Mini DHS 2025–26

Form of TFGBV	Percentage of women who have experienced TFGBV by any perpetrator	Percentage who have experienced TFGBV in the past 12 months			Number of women
	Ever[1]	Often	Sometimes	Often or sometimes	
Revenge pornography	1.0	0.2	0.5	0.6	15,614
Online harassment	4.7	0.8	2.3	3.1	15,614
Doxxing	1.8	0.4	0.8	1.2	15,614
Blackmail/sextortion	1.7	0.4	0.9	1.3	15,614
Hacking	4.6	0.7	1.7	2.4	15,614
Cyberbullying	2.2	0.6	0.8	1.3	15,614
Cyberstalking	1.9	0.5	0.8	1.3	15,614
Online sexual harassment or cyber sexual abuse	3.0	0.4	1.8	2.2	15,614
Online control	19.2	9.0	7.6	16.6	15,614
TFGBV from previous partner	1.5	0.2	0.3	0.5	15,614
Total	26.0	10.3	10.6	20.9	15,614

[1] Includes TFGBV in the past 12 months. For women who were married or living together before age 15 and reported violence only by their husband, and for never-married women who had an intimate partner before age 15 and reported violence only by their intimate partner, the violence could have occurred before age 15.

Table 14.3 Experience of various forms of technology-facilitated gender-based violence by any perpetrator according to background characteristics

Percentage of women age 15–49 who experienced technology-facilitated gender-based violence (TFGBV) by any perpetrator in the 12 months preceding the survey by form of TFGBV, according to background characteristics, Kenya Mini DHS 2025–26

	Form of TFGBV											
Background characteristic	Any form of TFGBV	Revenge porno-graphy	Online harass-ment	Doxxing	Blackmail/ sextortion	Hacking	Cyber-bullying	Cyber-stalking	Online sexual harass-ment or cyber sexual abuse	Online control	TFGBV from previous partner	Number of women
Age												
15–19	8.6	0.4	2.5	0.6	1.2	0.8	1.0	0.4	1.5	5.6	0.1	2,866
20–24	32.7	1.0	4.3	1.4	1.9	3.9	1.4	2.6	4.2	25.7	0.8	2,692
25–29	29.9	0.7	4.3	1.7	1.8	4.4	1.8	2.1	2.6	24.8	0.8	2,746
30–39	21.6	0.8	3.2	1.2	1.1	2.1	1.6	1.4	2.0	17.5	0.6	4,496
40–49	12.2	0.4	1.2	0.9	0.6	0.8	0.8	0.4	0.8	9.7	0.4	2,815
Residence												
Urban	22.9	0.8	3.6	1.6	1.5	2.6	1.7	1.8	2.9	17.6	0.7	7,096
Rural	19.2	0.5	2.7	0.8	1.1	2.1	1.1	1.0	1.5	15.8	0.4	8,518
Marital status												
Never married	15.7	0.6	3.2	0.8	1.2	2.1	1.2	1.4	3.3	10.2	0.0	5,099
Never had intimate partner	2.5	0.2	1.0	0.2	0.5	0.6	0.3	0.3	1.2	0.0	0.0	2,753
Ever had intimate partner	31.1	1.1	5.8	1.6	2.0	3.9	2.1	2.7	5.9	22.2	0.0	2,346
Ever married	23.4	0.7	3.0	1.3	1.3	2.5	1.4	1.3	1.6	19.7	0.8	10,515
Married/living together	22.7	0.6	2.3	1.0	0.9	2.2	1.0	1.0	1.3	19.6	0.7	8,612
Divorced/separated/ widowed	26.8	1.2	6.6	3.0	3.0	3.7	3.6	2.8	2.9	20.4	1.1	1,903
Education[1]												
No education	7.0	0.4	0.5	0.6	0.7	0.6	0.4	0.2	0.1	6.3	0.2	822
Primary	19.6	0.5	2.7	1.1	1.4	1.5	1.5	0.5	1.0	16.8	0.7	4,508
Secondary	21.1	0.7	3.2	1.2	1.2	2.2	1.0	1.3	2.2	16.9	0.4	7,231
More than secondary	26.0	0.8	4.1	1.4	1.4	4.5	2.0	2.9	4.3	18.2	0.7	3,054
Wealth quintile												
Lowest	14.3	0.2	1.8	0.4	1.0	0.9	1.0	0.4	0.5	12.3	0.2	2,477
Second	20.4	0.6	3.1	0.9	1.1	1.7	0.9	0.9	1.1	17.6	0.5	2,881
Middle	21.4	0.9	3.1	1.4	1.8	2.7	1.8	1.3	2.3	17.4	0.6	3,053
Fourth	25.6	1.0	3.7	1.4	1.2	3.2	1.4	1.7	3.1	20.1	0.8	3,508
Highest	20.7	0.5	3.3	1.5	1.3	2.7	1.5	2.0	3.2	14.8	0.5	3,695
Total	20.9	0.6	3.1	1.2	1.3	2.4	1.3	1.3	2.2	16.6	0.5	15,614

[1] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 14.3C Experience of various forms technology-facilitated gender-based violence by any perpetrator: County level

Percentage of women age 15–49 who have experienced technology-facilitated gender-based violence (TFGBV) by any perpetrator and percentage of women who have experienced TFGBV by any perpetrator in the 12 months preceding the survey by form of TFGBV, according to county, Kenya Mini DHS 2025–26

	Form of TFGBV											
County	Any form of TFGBV	Revenge porno-graphy	Online harass-ment	Doxxing	Blackmail/ sextortion	Hacking	Cyber-bullying	Cyber-stalking	Online sexual harass-ment or cyber sexual abuse	Online control	TFGBV from previous partner	Number of Women
Mombasa	17.5	1.5	1.1	2.4	1.5	0.9	1.4	1.4	0.4	15.7	0.9	556
Kwale	22.8	0.4	0.6	0.7	0.0	2.0	0.9	0.6	1.1	21.0	0.0	253
Kilifi	11.3	0.0	0.7	0.0	0.0	0.2	0.0	0.2	0.0	10.8	0.0	469
Tana River	11.6	0.0	0.0	0.0	0.0	0.2	0.0	0.1	0.5	11.2	0.0	96
Lamu	5.8	0.0	0.3	0.7	0.0	0.0	0.0	0.0	0.2	4.9	0.0	41
Taita Taveta	21.9	0.4	3.5	1.3	1.1	2.8	2.1	1.9	4.1	18.8	0.0	107
Garissa	7.0	1.0	1.5	1.5	0.8	1.1	1.1	0.0	0.0	5.0	0.5	127
Wajir	1.9	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.9	0.0	94
Mandera	1.9	0.0	0.0	0.0	0.0	0.2	0.0	0.0	0.0	1.8	0.0	96
Marsabit	8.5	0.7	0.6	0.7	0.6	0.7	0.0	0.0	0.4	7.9	0.0	98
Isiolo	9.9	0.2	2.1	1.2	0.4	0.4	0.6	0.8	0.0	9.2	0.4	58
Meru	16.1	0.7	2.7	0.4	0.3	0.9	1.7	0.2	3.1	10.3	0.0	388
Tharaka-Nithi	26.1	0.0	4.1	1.1	0.6	2.2	4.3	1.7	1.2	24.5	0.0	100
Embu	26.0	0.6	1.4	2.0	1.9	4.8	1.4	1.8	3.4	20.8	0.7	179
Kitui	22.1	0.9	3.9	0.3	3.3	2.4	1.6	1.2	1.9	17.9	0.5	319
Machakos	27.8	2.1	3.1	2.7	1.0	4.1	1.9	3.2	2.4	25.0	0.5	517
Makueni	16.3	0.3	2.4	0.2	1.5	2.7	0.5	0.5	2.5	11.4	0.5	259
Nyandarua	21.3	2.3	1.6	0.7	0.0	2.1	1.5	1.2	5.4	15.0	0.3	148
Nyeri	26.2	2.3	3.4	3.0	2.4	7.2	2.8	4.6	7.9	14.8	1.4	152
Kirinyaga	19.8	0.0	3.1	0.4	0.2	3.6	1.3	0.3	1.4	16.9	0.0	197
Murang'a	21.5	0.2	6.5	0.0	0.0	2.7	0.7	0.4	3.6	17.5	0.6	285
Kiambu	22.4	0.3	2.5	0.5	1.0	1.5	1.2	1.7	3.9	15.1	0.0	1,316
Turkana	21.0	0.2	2.7	3.0	1.4	3.3	0.7	0.7	2.2	17.0	1.0	134
West Pokot	13.4	0.9	0.9	0.7	1.1	0.6	0.4	0.5	0.8	10.6	2.0	172
Samburu	24.5	0.1	4.0	2.7	2.3	5.5	3.7	2.1	0.9	19.2	0.3	74
Trans Nzoia	28.6	0.0	2.2	0.0	0.6	5.3	0.3	1.5	1.0	24.2	0.0	289
Uasin Gishu	27.0	0.8	7.1	1.7	2.2	6.4	2.5	3.0	3.2	19.7	2.2	498
Elgeyo-Marakwet	9.8	0.0	0.5	0.0	0.0	0.0	0.1	0.0	0.0	9.8	0.0	133
Nandi	11.1	0.4	1.1	0.5	0.7	1.9	0.7	1.1	0.1	8.1	0.5	269
Baringo	24.6	0.8	6.4	2.1	3.1	1.6	2.8	1.3	2.9	19.5	0.0	131
Laikipia	22.2	0.0	0.2	0.0	0.0	1.2	0.3	0.6	0.3	20.9	0.0	142
Nakuru	19.6	0.6	4.9	1.6	2.9	3.7	1.6	1.3	1.3	15.5	0.4	809
Narok	23.4	0.0	2.7	0.0	1.0	1.3	0.3	0.4	2.3	21.7	0.0	334
Kajiado	24.6	0.2	4.6	2.2	0.9	0.9	4.0	3.7	2.6	18.9	0.6	325
Kericho	28.1	0.5	5.8	0.4	2.3	5.5	1.9	1.3	2.9	20.9	0.2	296
Bomet	21.9	0.9	7.9	0.0	2.0	6.8	0.0	0.8	5.0	13.5	0.0	349
Kakamega	22.3	0.8	3.8	2.5	0.4	1.5	0.0	1.3	1.5	16.7	0.0	501
Vihiga	12.4	0.4	0.7	1.4	0.7	1.8	1.4	0.8	2.9	8.3	0.0	169
Bungoma	24.9	0.7	2.2	1.0	0.7	1.9	1.1	1.3	1.6	21.7	0.7	489
Busia	24.2	1.0	2.2	2.3	1.4	1.0	3.4	0.7	0.7	22.0	0.0	275
Siaya	19.9	0.6	2.3	0.9	0.6	2.0	1.0	1.7	1.1	15.5	0.0	282
Kisumu	26.5	3.5	7.1	2.5	2.1	2.6	4.4	2.8	3.1	21.2	1.2	408
Homa Bay	23.0	0.9	1.7	0.7	2.2	1.6	0.6	1.0	0.1	19.2	2.9	382
Migori	23.3	0.5	4.9	1.7	2.4	2.4	1.5	0.9	0.9	20.9	0.8	383
Kisii	16.9	0.0	2.6	0.0	0.3	1.3	0.0	2.4	0.5	14.8	0.2	324
Nyamira	19.8	2.0	4.9	2.3	3.5	2.4	2.3	1.4	2.6	15.4	1.1	180
Nairobi City	21.2	0.3	2.5	1.4	1.3	2.0	1.4	1.1	2.9	16.4	0.7	2,412
Total	20.9	0.6	3.1	1.2	1.3	2.4	1.3	1.3	2.2	16.6	0.5	15,614

Table 14.4 Experience of technology-facilitated gender-based violence by any intimate partner

Among women age 15–49 who have ever had a husband or intimate partner, percentage who have ever experienced technology-facilitated gender-based violence (TFGBV) committed by their current, most recent, or previous husband/intimate partner and percentage who have experienced TFGBV committed by their current, most recent, or previous husband/intimate partner in the 12 months preceding the survey, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Percentage of women who have ever experienced TFGBV by any intimate partner	Percentage of women who have experienced TFGBV by any intimate partner in the past 12 months			Number of women
	Ever[1]	Often	Sometimes	Often or sometimes	
Age					
15–19	21.6	11.7	9.0	20.7	853
20–24	37.0	17.5	17.0	34.5	2,217
25–29	32.7	15.4	12.6	27.9	2,598
30–39	24.4	10.9	8.2	19.1	4,413
40–49	16.9	4.9	5.9	10.9	2,779
Residence					
Urban	28.8	12.0	11.1	23.0	5,880
Rural	24.5	11.5	9.4	20.9	6,981
Marital status					
Never married	28.6	11.4	13.5	24.8	2,346
Ever had intimate partner	28.6	11.4	13.5	24.8	2,346
Ever married	25.9	11.8	9.5	21.2	10,515
Married/living together	23.6	11.1	9.8	20.9	8,612
Divorced/separated/ widowed	36.4	14.8	7.9	22.7	1,903
Education[2]					
No education	8.3	5.1	2.6	7.7	749
Primary	23.7	11.2	7.7	18.9	4,313
Secondary	30.7	13.7	12.4	26.1	5,079
More than secondary	27.8	10.5	12.1	22.6	2,720
Wealth quintile					
Lowest	18.8	9.6	7.2	16.9	1,996
Second	27.2	13.2	9.9	23.1	2,352
Middle	27.5	13.0	9.5	22.5	2,532
Fourth	30.5	13.7	12.1	25.9	2,964
Highest	26.0	8.8	11.0	19.8	3,017
Total	26.4	11.7	10.2	21.9	12,861

Note: The term husband includes a partner with whom a woman is living as if married. Husband/intimate partner refers to the current husband for currently married women; the most recent husband for divorced, separated, or widowed women; and the current intimate partner for never-married women who currently have an intimate partner.

[1] Includes TFGBV in the past 12 months. For women who were married or living together before age 15 and reported violence only by their husband, and for never-married women who had an intimate partner before age 15 and reported violence only by their intimate partner, the violence could have occurred before age 15.

[2] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 14.4C Experience of technology-facilitated gender-based violence by any intimate partner: County level

Among women age 15–49 who have ever had a husband or intimate partner, percentage who have ever experienced technology-facilitated gender-based violence (TFGBV) committed by their current, most recent, or previous husband/intimate partner and percentage who have experienced TFGBV committed by their current, most recent, or previous husband/intimate partner in the 12 months preceding the survey, according to county, Kenya Mini DHS 2025–26

	Percentage of women who have ever experienced TFGBV by any intimate partner	Percentage of women who have experienced TFGBV by any intimate partner in the past 12 months			
County	Ever[1]	Often	Sometimes	Often or sometimes	Number of women
Mombasa	22.4	11.8	8.7	20.5	452
Kwale	29.2	17.1	7.9	25.1	212
Kilifi	15.2	10.8	3.7	14.5	350
Tana River	15.2	1.5	12.2	13.8	78
Lamu	10.0	4.0	3.0	7.1	28
Taita Taveta	26.8	10.1	11.7	21.8	98
Garissa	9.0	3.9	4.9	8.8	94
Wajir	2.9	1.3	1.6	2.9	61
Mandera	2.6	2.0	0.6	2.6	65
Marsabit	14.8	5.3	6.1	11.4	69
Isiolo	18.9	7.4	4.8	12.3	45
Meru	20.9	5.3	8.1	13.4	337
Tharaka-Nithi	33.5	20.6	9.6	30.2	81
Embu	27.5	14.8	12.4	27.1	143
Kitui	27.4	11.9	13.2	25.1	267
Machakos	32.0	18.8	11.1	29.8	450
Makueni	21.2	4.8	11.0	15.8	215
Nyandarua	28.4	6.1	13.7	19.8	127
Nyeri	24.0	13.9	8.2	22.1	126
Kirinyaga	30.1	8.8	10.4	19.1	176
Murang'a	32.3	4.5	16.8	21.2	246
Kiambu	26.2	12.6	7.3	19.9	1,055
Turkana	28.4	12.5	9.3	21.8	108
West Pokot	16.2	4.9	11.2	16.2	140
Samburu	29.5	12.6	13.0	25.6	65
Trans Nzoia	32.0	23.5	7.1	30.6	230
Uasin Gishu	33.0	14.0	14.9	28.8	381
Elgeyo-Marakwet	12.7	8.0	4.4	12.4	104
Nandi	14.4	3.8	8.5	12.3	194
Baringo	34.7	9.3	18.6	27.9	108
Laikipia	28.3	20.2	5.6	25.8	117
Nakuru	25.6	11.3	9.4	20.7	693
Narok	28.5	16.1	9.5	25.6	286
Kajiado	27.4	13.0	13.2	26.2	254
Kericho	30.0	11.9	15.7	27.5	253
Bomet	23.0	6.5	12.6	19.1	280
Kakamega	25.2	13.5	8.5	22.0	440
Vihiga	25.7	3.9	9.2	13.1	128
Bungoma	29.8	15.9	11.5	27.4	418
Busia	31.8	23.5	6.6	30.1	217
Siaya	20.6	8.7	12.0	20.6	223
Kisumu	32.6	18.2	10.0	28.2	347
Homa Bay	35.9	18.5	8.8	27.3	312
Migori	28.8	14.1	11.4	25.5	335
Kisii	19.1	12.9	5.7	18.6	284
Nyamira	20.4	16.0	3.8	19.8	149
Nairobi City	29.5	8.1	13.1	21.2	2,019
Total	26.4	11.7	10.2	21.9	12,861

Note: The term husband includes a partner with whom a woman is living as if married. Husband/intimate partner refers to the current husband for currently married women; the most recent husband for divorced, separated, or widowed women; and the current intimate partner for never-married women who currently have an intimate partner.

[1] Includes TFGBV in the past 12 months. For women who were married or living together before age 15 and reported violence only by their husband, and for never-married women who had an intimate partner before age 15 and reported violence only by their intimate partner, the violence could have occurred before age 15.

Table 14.5 Persons committing technology-facilitated gender-based violence

Among women age 15–49 who have experienced technology-facilitated gender-based violence, percentage who report specific persons who committed the violence, according to the respondent's partnership status, Kenya Mini DHS 2025–26

	Partnership status		
Person	Ever married/ever had intimate partner	Never married/never had intimate partner	Total
Current husband/partner	60.6	na	59.0
Former husband/partner	22.3	na	21.7
Father/stepfather	0.1	0.6	0.1
Mother/stepmother	0.2	0.0	0.2
Brother/stepbrother	0.1	0.0	0.1
Sister/stepsister	0.1	0.0	0.1
Other relative	0.7	2.8	0.8
Current boyfriend	1.3	0.0	1.2
Former boyfriend	7.9	9.3	7.9
In-law	0.4	na	0.4
Own friend/acquaintance	3.4	19.6	3.8
Family friend	0.6	2.1	0.6
Schoolmate/classmate	0.5	6.1	0.7
Employer/someone at work	0.5	0.0	0.5
Priest/religious leader	0.2	0.0	0.2
Stranger	12.7	45.3	13.6
Online attacker	5.2	23.2	5.7
Coach/mentor	0.0	0.0	0.0
Other	0.2	0.0	0.2
Number of women	3,945	112	4,057

na = Not applicable

Table 14.6 Forms of controlling behaviours

Percentage of women age 15–49 who have ever had a husband or intimate partner who have experienced controlling behaviours ever or in the 12 months preceding the survey, perpetrated by a husband or intimate partner, Kenya Mini DHS 2025–26

			Frequency in the past 12 months	
Controlling behaviour	Ever experienced	Experienced in the past 12 months	Often	Sometimes
Any controlling behaviour	61.3	54.3	31.0	23.2
Is jealous or angry if she talks to other men	50.1	43.2	19.1	24.1
Wrongly accuses her of being unfaithful	20.3	16.5	6.7	9.8
Does not permit her to meet her female friends	14.7	12.4	5.6	6.7
Tries to limit her contact with her family	8.0	6.3	2.8	3.5
Insists on knowing where she is at all times	36.0	32.0	19.0	13.0
Insists on reading call logs/messages/emails without consent	18.8	16.2	8.6	7.6
Insists she gives him passwords to phone/digital platforms	15.5	13.3	7.5	5.8
Uses technology to track her in ways that make her feel controlled	5.6	4.8	2.4	2.3
At least three forms of controlling behaviours	27.1	24.1	18.7	5.4
Number of women ever married or never married who ever had an intimate partner	12,861	12,861	12,861	12,861

Note: The term husband includes a partner with whom a woman is living as if married. Husband/intimate partner refers to the current husband for currently married women; the most recent husband for divorced, separated, or widowed women; the current intimate partner for never-married women who currently have an intimate partner; and the most recent intimate partner for never-married women who do not currently have an intimate partner but had one in the past.

Table 14.7 Controlling behaviours of husband/intimate partner by background characteristics

Percentage of women age 15–49 who have ever had a husband or intimate partner whose husbands/intimate partners have ever demonstrated specific types of controlling behaviours, according to background characteristics, Kenya Mini DHS 2025–26

	Percentage of women whose husband/intimate partner:										
Background characteristic	Is jealous or angry if she talks to other men	Wrongly accuses her of being unfaithful	Does not permit her to meet her female friends	Tries to limit her contact with her family	Insists on knowing where she is at all times	Insists on reading call logs/ messages/ emails without consent	Insists she gives him passwords to phone/ digital platforms	Uses technology to track her in ways that make her feel controlled	Displays 3 or more of the specific behaviours	Displays none of the specific behaviours	Number of women who ever had a husband/ intimate partner
Age											
15–19	48.8	18.0	8.8	3.5	39.2	15.1	13.9	5.1	22.0	36.6	853
20–24	58.0	18.6	15.5	5.7	44.0	26.4	22.9	8.9	34.0	28.2	2,217
25–29	53.1	21.3	16.3	7.9	36.7	22.6	20.4	7.2	30.4	35.3	2,598
30–39	49.4	20.7	15.3	9.7	34.1	17.8	14.1	4.6	25.7	41.0	4,413
40–49	42.7	20.6	13.3	8.6	31.0	11.9	7.6	3.3	22.2	47.4	2,779
Religion											
Catholic	52.9	20.3	15.4	9.5	38.9	20.9	16.9	6.7	28.8	35.6	2,204
Protestant	50.6	21.0	15.1	8.3	36.0	18.6	15.3	5.7	26.8	38.0	5,329
Evangelical churches	51.1	20.4	13.9	7.8	35.2	17.8	16.0	5.2	26.8	39.2	2,451
African-instituted churches	54.2	23.0	14.8	7.6	38.1	23.2	16.8	5.5	31.5	35.4	1,215
Orthodox	43.6	23.2	5.8	3.5	33.3	8.7	5.6	4.5	30.9	45.2	51
Other Christian	41.4	20.1	16.1	7.8	34.2	15.1	18.8	2.7	28.6	42.9	416
Islam	38.3	12.1	12.6	5.1	31.0	13.0	9.0	5.0	18.1	49.8	947
Hindu	*	*	*	*	*	*	*	*	*	*	1
Traditionist	35.7	20.0	6.7	2.9	9.6	13.2	7.3	0.0	18.9	60.5	31
No religion/atheist	48.3	21.1	14.5	4.8	33.0	23.5	16.9	7.1	30.3	40.7	213
Other	*	*	*	*	*	*	*	*	*	*	2
Residence											
Urban	51.5	19.8	14.9	7.7	36.5	19.4	17.0	6.8	27.9	37.8	5,880
Rural	49.0	20.7	14.5	8.2	35.6	18.3	14.2	4.6	26.4	39.5	6,981
Marital status											
Never married	55.2	19.5	8.9	3.3	41.4	18.9	17.6	7.8	27.5	32.0	2,346
Currently has intimate partner	55.9	18.3	7.8	2.6	42.8	19.0	18.8	8.1	27.3	29.0	1,535
Had intimate partner	53.9	21.7	10.9	4.6	38.8	18.7	15.4	7.3	27.9	37.6	811
Ever married	49.0	20.5	16.0	9.0	34.8	18.8	15.0	5.1	27.0	40.2	10,515
Married/living together	47.6	18.0	13.6	6.8	33.1	16.8	13.4	4.3	24.3	41.6	8,612
Divorced/separated/ widowed	55.6	31.5	26.8	19.0	42.2	27.8	22.0	8.6	39.0	33.8	1,903
Education[1]											
No education	29.5	14.0	8.6	5.2	21.9	6.6	3.8	1.3	13.1	61.7	749
Primary	50.7	23.5	16.3	9.9	35.9	17.7	13.3	4.3	27.5	39.4	4,313
Secondary	53.8	20.8	15.6	8.1	38.6	22.1	19.2	6.1	30.3	34.6	5,079
More than secondary	48.2	15.9	12.1	5.5	35.3	17.7	15.3	8.0	24.2	39.1	2,720
Wealth quintile											
Lowest	45.2	21.6	14.2	9.1	35.3	14.8	10.2	2.3	25.2	43.5	1,996
Second	51.3	21.3	16.3	8.2	36.8	20.7	15.8	4.5	28.8	37.4	2,352
Middle	52.6	21.8	16.0	9.5	36.1	20.2	16.4	6.6	28.3	36.0	2,532
Fourth	53.3	22.2	15.5	9.1	38.2	21.8	19.5	7.1	29.7	36.6	2,964
Highest	47.4	15.5	11.8	4.7	33.6	15.9	13.9	6.3	23.3	40.9	3,017
Total	50.1	20.3	14.7	8.0	36.0	18.8	15.5	5.6	27.1	38.7	12,861

Notes: The term husband includes a partner with whom a woman is living as if married. Husband/intimate partner refers to the current husband for currently married women; the most recent husband for divorced, separated, or widowed women; and the current intimate partner for never-married women who currently have an intimate partner. An asterisk indicates that a figure is based on fewer than 25 unweighted cases and has been suppressed.

[1] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 14.7C Controlling behaviours of husband/intimate partner: County level

Percentage of women age 15–49 who have ever had a husband or intimate partner whose husbands/intimate partners have ever demonstrated specific types of controlling behaviours, according to county, Kenya Mini DHS 2025–26

	Percentage of women whose husband/intimate partner:										
County	Is jealous or angry if she talks to other men	Wrongly accuses her of being unfaithful	Does not permit her to meet her female friends	Tries to limit her contact with her family	Insists on knowing where she is at all times	Insists on reading call logs/ messages/ emails without consent	Insists she gives him passwords to phone/ digital platforms	Uses technology to track her in ways that make her feel controlled	Displays 3 or more of the specific behaviours	Displays none of the specific behaviours	Number of women who ever had a husband/ intimate partner
Mombasa	50.1	14.6	10.5	5.7	29.9	16.7	12.4	7.0	22.9	43.6	452
Kwale	60.6	25.7	14.5	8.9	59.7	26.3	18.5	7.5	38.9	24.4	212
Kilifi	37.8	16.5	9.1	6.0	29.3	13.6	8.7	6.9	18.8	56.1	350
Tana River	69.1	6.1	4.3	0.7	64.1	14.2	6.3	0.4	17.8	16.8	78
Lamu	34.8	12.9	3.4	4.7	19.0	9.5	3.2	1.1	13.4	63.2	28
Taita Taveta	60.4	24.0	14.9	7.9	41.3	19.9	15.7	5.2	31.9	32.4	98
Garissa	25.7	8.0	10.8	5.4	22.3	6.7	3.8	2.4	12.2	68.8	94
Wajir	3.5	1.4	7.7	1.4	24.5	2.9	1.4	0.2	4.0	74.4	61
Mandera	12.4	1.1	8.4	1.0	15.2	2.2	1.7	0.3	4.4	77.2	65
Marsabit	41.4	16.9	11.3	2.9	28.9	11.6	10.3	2.7	22.4	54.5	69
Isiolo	58.4	18.3	16.4	3.8	31.2	13.0	10.7	2.6	25.4	36.0	45
Meru	55.3	16.0	11.2	6.4	37.4	12.7	11.2	6.0	21.5	35.9	337
Tharaka-Nithi	54.4	30.3	11.9	15.4	61.8	31.0	15.6	7.0	37.8	20.6	81
Embu	47.2	16.3	17.0	9.0	48.7	22.8	17.2	2.7	30.0	34.4	143
Kitui	55.8	12.3	11.2	6.8	42.2	21.8	15.3	3.5	28.5	39.0	267
Machakos	52.9	22.8	22.6	9.6	43.1	23.2	24.5	10.0	34.7	36.9	450
Makueni	38.2	13.7	11.9	5.7	24.5	14.2	12.4	1.9	18.1	50.2	215
Nyandarua	60.3	20.6	18.6	11.4	36.8	16.6	15.5	4.9	26.2	27.2	127
Nyeri	45.4	20.2	17.0	4.3	40.3	16.2	11.4	6.0	22.1	36.4	126
Kirinyaga	59.7	23.3	15.4	10.2	43.5	22.4	20.6	5.8	32.3	31.2	176
Murang'a	48.7	24.6	16.9	7.8	47.2	27.6	18.1	7.8	29.3	40.5	246
Kiambu	39.6	12.7	7.3	4.4	23.3	20.5	15.3	4.7	23.1	54.3	1,055
Turkana	46.2	16.7	15.0	3.3	35.9	18.4	12.4	7.3	26.0	40.3	108
West Pokot	35.6	18.7	16.9	3.7	26.8	10.7	10.8	2.6	21.7	56.2	140
Samburu	58.6	34.5	16.9	12.2	42.8	20.7	17.0	3.7	33.9	27.4	65
Trans Nzoia	57.7	25.5	20.7	11.5	54.9	26.0	18.7	7.3	38.8	27.4	230
Uasin Gishu	58.2	22.7	13.3	8.6	28.9	23.5	17.2	9.9	25.4	33.5	381
Elgeyo-Marakwet	46.7	13.3	8.1	1.8	24.7	9.6	10.4	1.3	16.4	48.2	104
Nandi	26.0	17.2	11.2	10.9	17.7	8.9	8.0	2.9	15.4	61.3	194
Baringo	65.8	21.1	12.1	8.3	42.2	22.9	18.9	3.9	35.8	27.2	108
Laikipia	47.9	22.9	25.8	7.2	39.0	25.1	21.1	4.1	33.9	44.3	117
Nakuru	47.7	25.2	17.6	10.1	33.0	16.9	15.4	7.6	25.4	35.4	693
Narok	46.9	27.5	23.6	9.1	36.7	17.9	20.7	6.1	34.5	41.9	286
Kajiado	57.3	17.7	8.5	8.1	37.7	19.1	15.6	8.8	29.4	37.6	254
Kericho	54.9	20.7	16.3	8.9	28.5	21.9	17.0	6.6	25.3	34.1	253
Bomet	41.5	5.5	10.5	1.6	15.4	18.6	3.2	2.9	12.7	49.2	280
Kakamega	56.9	28.0	16.5	13.3	40.0	17.6	14.3	3.4	31.9	31.6	440
Vihiga	59.2	30.9	22.3	10.7	50.7	21.1	17.7	9.4	37.7	28.1	128
Bungoma	59.1	24.0	17.1	13.4	43.9	24.3	16.8	3.6	35.0	27.9	418
Busia	55.3	28.6	13.6	7.2	39.5	24.9	20.0	5.2	31.7	33.3	217
Siaya	41.9	20.2	12.0	8.6	27.8	13.5	15.3	4.9	22.6	45.7	223
Kisumu	51.3	30.4	17.8	16.3	48.8	22.3	18.5	8.7	37.0	34.2	347
Homa Bay	64.7	32.1	15.9	5.8	52.4	23.0	27.6	4.8	38.9	22.8	312
Migori	41.1	19.3	9.8	6.8	25.9	20.8	15.8	5.3	23.9	44.1	335
Kisii	43.7	21.7	12.1	6.8	30.5	16.7	13.1	1.5	25.0	49.7	284
Nyamira	42.4	17.6	14.7	9.7	20.0	15.7	13.1	8.1	22.2	48.8	149
Nairobi City	56.0	20.4	18.4	8.1	39.9	16.8	16.1	5.5	27.6	29.8	2,019
Total	50.1	20.3	14.7	8.0	36.0	18.8	15.5	5.6	27.1	38.7	12,861

Note: The term husband includes a partner with whom a woman is living as if married. Husband/intimate partner refers to the current husband for currently married women; the most recent husband for divorced, separated, or widowed women; and the current intimate partner for never-married women who currently have an intimate partner.

Table 14.8 Experience of technology-facilitated gender-based violence by platform

Percentage of women age 15–49 who have ever experienced technology-facilitated gender-based violence (TFGBV) and percentage who have experienced TFGBV in the 12 months preceding the survey, according to the platform from which they experienced such violence, Kenya Mini DHS 2025–26

Platform	Percentage of women who have experienced TFGBV by any perpetrator	Percentage who have experienced TFGBV in the past 12 months			Number of women
	Ever[1]	Often	Sometimes	Often or sometimes	
Social media	67.3	7.1	33.9	41.0	1,259
Facebook	59.2	6.6	32.0	38.6	1,259
Instagram	8.0	0.6	3.1	3.8	1,259
Other social media[2]	3.9	1.2	0.7	1.9	1,259
Messaging apps	46.5	8.4	22.4	30.8	1,259
Text message/iMessage	27.0	5.2	11.5	16.6	1,259
WhatsApp	26.0	4.3	14.0	18.3	1,259
Other messaging apps[3]	1.8	0.6	0.6	1.2	1,259
Other[4]	1.5	0.0	0.7	0.7	1,259

[1] Includes TFGBV in the past 12 months. For women who were married or living together before age 15 and reported violence only by their husband, and for never-married women who had an intimate partner before age 15 and reported violence only by their intimate partner, the violence could have occurred before age 15.
[2] Other social media include X (formerly Twitter), TikTok, YouTube, and other social media mentioned by the respondent.
[3] Other messaging apps include Signal, Telegram, and other messaging apps mentioned by respondent.
[4] Other includes email and other platforms mentioned by the respondent.

Table 14.9 Experience of technology-facilitated gender-based violence by platform and background characteristics

Percentage of women age 15–49 who have experienced technology-facilitated gender-based violence by platform from which they experienced such violence, according to background characteristics, Kenya Mini DHS 2025–26

	Social media				Messaging apps					
Background characteristic	Facebook	Instagram	Other social media[1]	Any social media	Text message/ iMessage	WhatsApp	Other messaging apps[2]	Any messaging app	Other[3]	Number of women
Age										
15–19	54.1	25.9	0.2	76.5	21.8	8.6	2.7	30.3	0.5	150
20–24	61.1	13.7	3.2	72.2	15.9	32.1	1.3	44.2	0.6	335
25–29	63.5	4.2	6.4	68.5	32.9	32.6	2.2	51.1	0.8	284
30–39	61.9	1.3	3.4	64.4	29.8	25.4	2.2	49.3	1.6	358
40–49	43.9	0.0	5.9	49.8	41.0	17.7	0.1	53.5	5.8	132
Residence										
Urban	60.7	12.0	4.8	72.2	22.7	28.2	2.3	42.9	1.1	779
Rural	56.9	1.6	2.4	59.4	34.0	22.5	0.9	52.4	2.1	480
Marital status										
Never married	57.0	18.7	3.5	72.6	17.5	25.1	1.4	39.2	0.5	476
Never had intimate partner	50.6	37.7	3.6	80.3	13.3	14.5	1.9	23.7	0.6	112
Ever had intimate partner	59.0	12.8	3.4	70.2	18.8	28.4	1.2	43.9	0.5	363
Ever married	60.6	1.6	4.2	64.1	32.8	26.6	2.0	51.0	2.0	784
Married/living together	64.1	1.7	2.9	66.6	31.4	25.3	1.1	49.1	2.2	571
Divorced/separated/ widowed	51.1	1.5	7.6	57.4	36.5	30.1	4.4	56.1	1.6	213
Education[4]										
No education	*	*	*	*	*	*	*	*	*	17
Primary	49.2	0.0	2.5	50.7	46.9	23.9	0.1	59.3	1.9	223
Secondary	63.9	10.1	2.1	72.8	23.8	21.9	1.6	40.6	0.6	599
More than secondary	59.0	9.7	7.3	69.8	20.5	32.7	3.0	48.0	1.5	420
Wealth quintile										
Lowest	44.7	0.0	0.0	44.7	55.2	10.1	0.1	62.1	7.4	72
Second	57.0	1.8	2.0	60.2	40.2	25.0	2.1	57.5	0.8	134
Middle	64.4	0.6	1.9	65.1	28.1	24.3	1.6	47.2	1.2	245
Fourth	63.5	3.9	3.1	66.5	29.0	27.4	2.3	48.9	0.5	326
Highest	56.5	17.6	6.6	74.3	17.2	28.7	1.7	39.1	1.5	482
Total	59.2	8.0	3.9	67.3	27.0	26.0	1.8	46.5	1.5	1,259

Notes: Online control questions are excluded from this table. An asterisk indicates that a figure is based on fewer than 25 unweighted cases and has been suppressed.
[1] Other social media include X (formerly Twitter), TikTok, YouTube, and other social media mentioned by the respondent.
[2] Other messaging apps include Signal, Telegram, and other messaging apps mentioned by respondent.
[3] Other includes email and other platforms mentioned by the respondent.
[4] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 14.10 Perceived risk of violence and safety for women and girls

Percent distribution of households that perceive that certain groups of women and girls are at risk of violence and percent distribution of households by perception of safety of women and girls in the community, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Risk of violence			Perceived safety of women in the community					
	Yes	No	Total	Very safe	Safe	Somewhat safe	Not very safe	Total	Number of households
Sex									
Male	17.0	83.0	100.0	12.6	55.4	23.3	8.6	100.0	9,067
Female	19.0	81.0	100.0	10.6	53.1	25.4	11.0	100.0	17,313
Age									
15–19	11.5	88.5	100.0	12.8	58.4	22.1	6.7	100.0	1,065
20–24	16.0	84.0	100.0	14.0	54.0	23.6	8.4	100.0	3,032
25–29	17.8	82.2	100.0	12.5	53.6	23.8	10.1	100.0	4,047
30–34	17.5	82.5	100.0	10.2	55.2	25.4	9.2	100.0	3,491
35–39	18.7	81.3	100.0	10.2	52.3	26.7	10.7	100.0	3,393
40–44	19.9	80.1	100.0	11.9	51.5	26.7	9.9	100.0	2,466
45–49	20.9	79.1	100.0	9.6	52.5	25.4	12.5	100.0	2,048
50–54	21.2	78.8	100.0	9.7	50.2	27.0	13.1	100.0	1,980
55–59	18.3	81.7	100.0	10.8	53.8	24.4	11.1	100.0	1,224
60–64	19.2	80.8	100.0	10.6	54.6	23.4	11.4	100.0	1,127
65+	19.0	81.0	100.0	11.1	58.6	21.0	9.4	100.0	2,499
Education[1]									
No education	17.2	82.8	100.0	15.6	58.7	19.1	6.6	100.0	2,346
Primary	17.8	82.2	100.0	10.3	54.9	24.6	10.2	100.0	9,347
Secondary	17.6	82.4	100.0	10.4	53.5	25.5	10.6	100.0	9,178
More than secondary	20.7	79.3	100.0	12.7	50.5	25.8	11.0	100.0	5,494
Residence									
Urban	17.3	82.7	100.0	10.7	52.6	25.1	11.7	100.0	11,975
Rural	19.1	80.9	100.0	11.8	55.0	24.3	8.9	100.0	14,405
Wealth quintile									
Lowest	18.0	82.0	100.0	12.2	55.4	24.3	8.1	100.0	4,299
Second	20.4	79.6	100.0	11.0	54.1	25.0	10.0	100.0	4,719
Middle	17.2	82.8	100.0	11.0	54.3	24.8	9.8	100.0	5,346
Fourth	17.5	82.5	100.0	10.0	52.0	27.0	11.1	100.0	6,192
Highest	18.7	81.3	100.0	12.5	54.2	22.1	11.2	100.0	5,825
Total	18.3	81.7	100.0	11.3	53.9	24.7	10.2	100.0	26,380

Notes: Because this question was administered in the Household Questionnaire, the sex, age, and education variables refer to those of the household respondent. Total includes eight households that were missing information on the age of the household respondent and 15 households that were missing information on the education level of the household respondent.

[1] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 14.10C Perceived risk of violence and safety for women and girls: County level

Percent distribution of households that perceive that certain groups of women and girls are at risk of violence and percent distribution of households by perception of safety of women and girls in the community, according to county, Kenya Mini DHS 2025–26

	Risk of violence			Perceived safety of women in the community					
County	Yes	No	Total	Very safe	Safe	Somewhat safe	Not very safe	Total	Number of households
Mombasa	14.6	85.4	100.0	8.3	57.8	25.9	7.9	100.0	866
Kwale	6.5	93.5	100.0	12.1	71.8	11.7	4.5	100.0	318
Kilifi	3.8	96.2	100.0	22.5	64.3	10.4	2.8	100.0	699
Tana River	10.6	89.4	100.0	14.7	72.0	12.6	0.7	100.0	143
Lamu	12.2	87.8	100.0	25.6	48.9	18.2	7.4	100.0	62
Taita Taveta	7.4	92.6	100.0	26.3	40.4	28.0	5.3	100.0	206
Garissa	11.5	88.5	100.0	35.2	51.3	8.4	5.2	100.0	145
Wajir	5.9	94.1	100.0	34.2	65.7	0.0	0.1	100.0	126
Mandera	21.7	78.3	100.0	52.8	19.4	22.9	4.9	100.0	137
Marsabit	14.1	85.9	100.0	2.5	88.7	6.7	2.0	100.0	138
Isiolo	16.7	83.3	100.0	5.4	66.2	20.3	8.0	100.0	87
Meru	20.0	80.0	100.0	10.2	48.5	29.0	12.2	100.0	840
Tharaka-Nithi	31.3	68.7	100.0	6.0	55.2	23.3	15.5	100.0	209
Embu	24.7	75.3	100.0	8.0	48.9	20.1	23.1	100.0	385
Kitui	5.8	94.2	100.0	9.4	75.3	13.5	1.8	100.0	526
Machakos	14.2	85.8	100.0	8.5	70.1	16.4	4.9	100.0	1,021
Makueni	22.5	77.5	100.0	8.4	53.9	31.8	5.9	100.0	506
Nyandarua	10.2	89.8	100.0	5.0	66.5	20.5	8.0	100.0	341
Nyeri	19.0	81.0	100.0	6.7	67.8	15.4	10.1	100.0	441
Kirinyaga	14.1	85.9	100.0	5.4	59.3	26.0	9.3	100.0	482
Murang'a	26.1	73.9	100.0	8.7	62.1	21.4	7.8	100.0	571
Kiambu	8.4	91.6	100.0	18.3	60.9	15.5	5.3	100.0	2,295
Turkana	39.1	60.9	100.0	2.4	36.2	38.3	23.1	100.0	232
West Pokot	16.7	83.3	100.0	11.1	61.5	25.3	2.1	100.0	251
Samburu	28.4	71.6	100.0	8.0	45.1	36.4	10.5	100.0	109
Trans Nzoia	21.1	78.9	100.0	4.1	48.9	32.8	14.2	100.0	393
Uasin Gishu	18.2	81.8	100.0	4.0	63.8	24.4	7.8	100.0	794
Elgeyo-Marakwet	14.8	85.2	100.0	15.4	65.6	13.6	5.4	100.0	209
Nandi	7.7	92.3	100.0	41.2	45.5	9.7	3.6	100.0	414
Baringo	5.6	94.4	100.0	7.3	76.5	15.3	1.0	100.0	242
Laikipia	9.9	90.1	100.0	15.4	66.8	10.2	7.7	100.0	293
Nakuru	16.7	83.3	100.0	14.4	40.0	31.9	13.8	100.0	1,442
Narok	35.7	64.3	100.0	14.3	50.5	27.8	7.4	100.0	576
Kajiado	12.1	87.9	100.0	17.5	42.2	32.5	7.8	100.0	653
Kericho	19.5	80.5	100.0	3.9	61.1	27.2	7.8	100.0	464
Bomet	18.8	81.2	100.0	23.3	50.7	20.0	6.0	100.0	437
Kakamega	15.5	84.5	100.0	10.0	37.7	43.7	8.6	100.0	850
Vihiga	7.3	92.7	100.0	15.4	51.6	28.7	4.3	100.0	267
Bungoma	37.7	62.3	100.0	2.2	42.7	32.4	22.7	100.0	746
Busia	20.8	79.2	100.0	10.7	53.3	17.9	18.1	100.0	429
Siaya	9.4	90.6	100.0	1.9	60.6	32.1	5.4	100.0	482
Kisumu	25.0	75.0	100.0	9.0	35.7	41.1	14.2	100.0	615
Homa Bay	36.7	63.3	100.0	16.1	45.1	30.6	8.3	100.0	568
Migori	26.5	73.5	100.0	5.0	53.3	36.0	5.7	100.0	499
Kisii	25.9	74.1	100.0	5.6	56.2	19.6	18.6	100.0	597
Nyamira	21.7	78.3	100.0	1.9	57.9	24.1	16.1	100.0	284
Nairobi City	22.9	77.1	100.0	7.7	47.9	27.6	16.8	100.0	3,992
Total	18.3	81.7	100.0	11.3	53.9	24.7	10.2	100.0	26,380

Note: Because this question was administered in the Household Questionnaire, the sex, age, and education variables refer to those of the household respondent.

Table 14.11 Perceived community response to femicide

Percentage of households by action they would take if they suspected that a woman or a girl was at risk of being killed, according to background characteristics, Kenya Mini DHS 2025–26

	Action taken						
Background characteristic	Mobilise community rescue	Report to police	Report to NGAO (chief, village leader, or other community leader)	Talk to family	Do nothing	Other action	Number of households
Sex							
Male	25.5	52.6	57.9	18.9	5.1	1.0	9,067
Female	23.1	45.0	60.4	19.4	6.8	1.1	17,313
Age							
15–19	23.4	56.4	49.9	15.3	7.0	0.9	1,065
20–24	23.7	53.9	47.7	17.4	6.1	1.1	3,032
25–29	22.7	52.0	50.2	15.4	7.9	1.6	4,047
30–34	22.5	49.3	55.9	18.1	7.4	1.3	3,491
35–39	23.1	46.1	60.9	18.8	6.4	1.5	3,393
40–44	24.7	45.8	62.2	20.2	6.3	1.2	2,466
45–49	24.2	45.4	65.2	19.0	5.4	1.4	2,048
50–54	24.6	46.0	66.5	24.0	5.0	0.4	1,980
55–59	27.3	42.0	67.4	20.6	6.5	0.2	1,224
60–64	29.9	42.8	70.2	22.3	3.2	0.6	1,127
65+	23.4	38.7	74.7	24.8	4.0	0.3	2,499
Education[1]							
No education	22.0	34.0	72.6	30.4	4.0	0.1	2,346
Primary	23.1	42.0	66.1	20.4	5.9	0.6	9,347
Secondary	24.3	50.3	56.3	18.1	6.3	1.4	9,178
More than secondary	25.4	58.5	48.1	14.3	7.5	1.8	5,494
Residence							
Urban	24.0	57.0	44.9	14.5	8.1	2.0	11,975
Rural	23.8	39.8	71.7	23.2	4.7	0.3	14,405
Wealth quintile							
Lowest	22.5	32.1	73.7	26.6	4.3	0.1	4,299
Second	21.8	40.1	71.8	22.0	5.1	0.2	4,719
Middle	24.4	45.9	66.3	20.4	5.2	0.5	5,346
Fourth	23.7	54.4	54.6	16.3	6.8	1.2	6,192
Highest	26.5	59.4	38.1	13.6	8.7	3.0	5,825
Total	23.9	47.6	59.5	19.2	6.2	1.1	26,380

Notes: Because this question was administered in the Household Questionnaire, the sex, age, and education variables refer to those of the household respondent. Total includes eight households that were missing information on the age of the household respondent and 15 households that were missing information on the education level of the household respondent.
NGAO = National Government Administrative Officer
[1] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 14.11C Perceived community response to femicide: County level

Percentage of households by action they would take if they suspected that a woman or a girl was at risk of being killed, according to county, Kenya Mini DHS 2025–26

	Action taken						
County	Mobilise community rescue	Report to police	Report to NGAO (chief, village leader, or other community leader etc.)	Talk to family	Do nothing	Other action	Total number of households
Mombasa	7.0	53.6	53.0	19.8	8.7	0.1	866
Kwale	6.3	49.7	76.5	9.3	1.2	0.1	318
Kilifi	1.1	41.4	67.9	17.5	2.1	0.0	699
Tana River	0.5	56.2	90.9	49.2	0.2	0.0	143
Lamu	3.4	54.0	85.6	16.8	1.6	0.6	62
Taita Taveta	34.0	73.5	75.7	27.8	1.5	0.3	206
Garissa	5.0	51.5	65.9	57.7	2.1	0.5	145
Wajir	1.9	45.9	72.6	56.9	0.2	0.0	126
Mandera	69.1	38.9	67.4	44.4	0.2	0.0	137
Marsabit	14.9	52.5	94.7	42.9	0.1	0.0	138
Isiolo	7.0	60.5	67.8	34.1	2.9	0.2	87
Meru	28.4	32.9	62.8	11.0	12.2	0.2	840
Tharaka-Nithi	19.7	51.4	75.7	32.5	4.2	1.5	209
Embu	25.7	29.1	50.0	31.2	8.5	2.3	385
Kitui	54.4	39.2	82.8	23.7	2.1	0.0	526
Machakos	24.8	38.0	63.7	15.7	11.4	0.7	1,021
Makueni	21.0	53.6	80.0	28.8	3.7	0.2	506
Nyandarua	24.0	52.8	78.7	11.1	3.6	0.4	341
Nyeri	30.3	61.0	49.2	19.2	4.2	0.1	441
Kirinyaga	48.5	34.7	51.1	14.9	9.4	1.0	482
Murang'a	63.6	43.1	62.8	20.1	1.8	0.2	571
Kiambu	25.9	68.2	23.6	6.1	5.5	0.4	2,295
Turkana	32.2	46.5	81.5	19.5	0.4	0.0	232
West Pokot	39.5	34.5	64.6	48.9	0.9	0.0	251
Samburu	25.5	30.5	55.2	24.0	13.7	1.7	109
Trans Nzoia	14.6	13.1	83.0	32.9	3.9	0.0	393
Uasin Gishu	37.1	44.3	69.9	20.2	4.7	0.2	794
Elgeyo-Marakwet	27.6	73.3	81.3	58.1	0.2	0.2	209
Nandi	26.4	63.2	58.6	17.0	2.6	0.0	414
Baringo	19.6	29.1	93.7	11.5	1.2	0.3	242
Laikipia	6.0	50.0	78.5	3.2	2.0	0.0	293
Nakuru	18.4	59.3	67.1	20.0	7.7	0.6	1,442
Narok	13.3	46.2	80.2	14.1	0.8	0.4	576
Kajiado	13.5	55.1	74.9	32.7	1.7	0.8	653
Kericho	20.9	27.3	76.2	26.3	1.7	1.0	464
Bomet	4.4	15.6	76.2	33.3	3.9	0.0	437
Kakamega	30.8	38.5	50.3	29.8	11.0	0.0	850
Vihiga	15.4	44.6	81.0	12.4	2.6	0.2	267
Bungoma	23.3	25.7	79.7	22.1	2.3	0.5	746
Busia	28.6	41.2	46.5	10.9	7.1	0.2	429
Siaya	4.5	50.8	68.1	11.1	2.8	0.0	482
Kisumu	20.3	53.6	63.7	24.3	7.4	2.7	615
Homa Bay	13.7	61.6	79.5	20.5	4.5	0.0	568
Migori	10.5	65.9	64.0	27.3	5.3	0.0	499
Kisii	38.0	29.4	70.4	17.6	6.7	0.0	597
Nyamira	13.7	16.5	58.7	17.3	13.7	0.2	284
Nairobi City	27.8	51.5	35.9	12.0	11.3	4.9	3,992
Total	23.9	47.6	59.5	19.2	6.2	1.1	26,380

Note: Because this question was administered in the Household Questionnaire, the sex, age, and education variables refer to those of the household respondent.
NGAO = National Government Administrative Officer

Table 14.12 Perceived socio-cultural and religious beliefs on justification of violence against women and girls

Percent distribution of households that believe that social, cultural, or religious beliefs justify violence against women or girls, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Justification for violence against women and girls		Total	Number of households
	Yes	No		
Sex				
Male	6.9	93.1	100.0	9,067
Female	8.3	91.7	100.0	17,313
Age				
15–19	7.5	92.5	100.0	1,065
20–24	8.4	91.6	100.0	3,032
25–29	9.1	90.9	100.0	4,047
30–34	7.6	92.4	100.0	3,491
35–39	7.9	92.1	100.0	3,393
40–44	7.7	92.3	100.0	2,466
45–49	7.6	92.4	100.0	2,048
50–54	8.0	92.0	100.0	1,980
55–59	6.6	93.4	100.0	1,224
60–64	5.8	94.2	100.0	1,127
65+	7.0	93.0	100.0	2,499
Education[1]				
No education	8.1	91.9	100.0	2,346
Primary	7.2	92.8	100.0	9,347
Secondary	7.2	92.8	100.0	9,178
More than secondary	9.8	90.2	100.0	5,494
Residence				
Urban	7.5	92.5	100.0	11,975
Rural	8.1	91.9	100.0	14,405
Wealth quintile				
Lowest	9.0	91.0	100.0	4,299
Second	8.0	92.0	100.0	4,719
Middle	6.8	93.2	100.0	5,346
Fourth	7.9	92.1	100.0	6,192
Highest	7.6	92.4	100.0	5,825
Total	7.8	92.2	100.0	26,380

Notes: Because this question was administered in the Household Questionnaire, the sex, age, and education variables refer to those of the household respondent. Total includes eight households that were missing information on the age of the household respondent and 15 households that were missing information on education level of the household respondent.

[1] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 14.12C Perceived socio-cultural and religious beliefs on justification of violence against women and girls: County level

Percent distribution of households that believe that social, cultural, or religious beliefs justify violence against women or girls, according to county, Kenya Mini DHS 2025–26

County	Justification for violence against women and girls		Total	Number of households
	Yes	No		
Mombasa	7.8	92.2	100.0	866
Kwale	6.4	93.6	100.0	318
Kilifi	4.0	96.0	100.0	699
Tana River	2.8	97.2	100.0	143
Lamu	4.5	95.5	100.0	62
Taita Taveta	3.7	96.3	100.0	206
Garissa	3.0	97.0	100.0	145
Wajir	0.0	100.0	100.0	126
Mandera	22.3	77.7	100.0	137
Marsabit	11.2	88.8	100.0	138
Isiolo	12.3	87.7	100.0	87
Meru	5.9	94.1	100.0	840
Tharaka-Nithi	8.6	91.4	100.0	209
Embu	17.5	82.5	100.0	385
Kitui	1.8	98.2	100.0	526
Machakos	2.0	98.0	100.0	1,021
Makueni	2.8	97.2	100.0	506
Nyandarua	4.5	95.5	100.0	341
Nyeri	9.1	90.9	100.0	441
Kirinyaga	5.2	94.8	100.0	482
Murang'a	5.3	94.7	100.0	571
Kiambu	5.5	94.5	100.0	2,295
Turkana	13.1	86.9	100.0	232
West Pokot	19.1	80.9	100.0	251
Samburu	24.6	75.4	100.0	109
Trans Nzoia	6.9	93.1	100.0	393
Uasin Gishu	8.5	91.5	100.0	794
Elgeyo-Marakwet	11.2	88.8	100.0	209
Nandi	2.7	97.3	100.0	414
Baringo	4.7	95.3	100.0	242
Laikipia	7.3	92.7	100.0	293
Nakuru	14.9	85.1	100.0	1,442
Narok	5.1	94.9	100.0	576
Kajiado	14.9	85.1	100.0	653
Kericho	12.3	87.7	100.0	464
Bomet	9.3	90.7	100.0	437
Kakamega	5.8	94.2	100.0	850
Vihiga	3.5	96.5	100.0	267
Bungoma	13.2	86.8	100.0	746
Busia	3.1	96.9	100.0	429
Siaya	5.5	94.5	100.0	482
Kisumu	8.7	91.3	100.0	615
Homa Bay	15.0	85.0	100.0	568
Migori	24.1	75.9	100.0	499
Kisii	6.6	93.4	100.0	597
Nyamira	2.6	97.4	100.0	284
Nairobi City	6.2	93.8	100.0	3,992
Total	7.8	92.2	100.0	26,380

Note: Because this question was administered in the Household Questionnaire, the sex, age, and education variables refer to those of the household respondent.

Table 14.13 Perceptions of victim blaming and privacy of domestic violence

Percent distribution of households that have heard of people who blame women or girls for the violence they experienced and percent distribution of households that think domestic violence is a private matter, according to background characteristics, Kenya Mini DHS 2025–26

Background characteristic	Heard of victim blaming			Perceive domestic violence to be private			Number of households
	Yes	No	Total	Yes	No	Total	
Sex							
Male	33.6	66.4	100.0	41.3	58.7	100.0	9,067
Female	36.7	63.3	100.0	35.4	64.6	100.0	17,313
Age							
15–19	29.1	70.9	100.0	31.6	68.4	100.0	1,065
20–24	35.0	65.0	100.0	37.0	63.0	100.0	3,032
25–29	34.5	65.5	100.0	35.8	64.2	100.0	4,047
30–34	34.9	65.1	100.0	36.9	63.1	100.0	3,491
35–39	38.0	62.0	100.0	38.6	61.4	100.0	3,393
40–44	38.7	61.3	100.0	38.5	61.5	100.0	2,466
45–49	36.2	63.8	100.0	36.3	63.7	100.0	2,048
50–54	40.0	60.0	100.0	37.4	62.6	100.0	1,980
55–59	37.4	62.6	100.0	37.5	62.5	100.0	1,224
60–64	36.1	63.9	100.0	40.5	59.5	100.0	1,127
65+	30.6	69.4	100.0	41.0	59.0	100.0	2,499
Education[1]							
No education	27.8	72.2	100.0	40.0	60.0	100.0	2,346
Primary	34.8	65.2	100.0	41.3	58.7	100.0	9,347
Secondary	35.2	64.8	100.0	38.4	61.6	100.0	9,178
More than secondary	41.1	58.9	100.0	28.2	71.8	100.0	5,494
Residence							
Urban	36.6	63.4	100.0	34.5	65.5	100.0	11,975
Rural	34.8	65.2	100.0	39.9	60.1	100.0	14,405
Wealth quintile							
Lowest	31.6	68.4	100.0	41.3	58.7	100.0	4,299
Second	34.5	65.5	100.0	40.0	60.0	100.0	4,719
Middle	35.1	64.9	100.0	39.4	60.6	100.0	5,346
Fourth	37.8	62.2	100.0	37.4	62.6	100.0	6,192
Highest	37.6	62.4	100.0	30.7	69.3	100.0	5,825
Total	35.6	64.4	100.0	37.5	62.5	100.0	26,380

Notes: Because this question was administered in the Household Questionnaire, the sex, age, and education variables refer to those of the household respondent. Total includes eight households that were missing information on the age of the household respondent and 15 households that were missing information on education level of the household respondent.

[1] No education includes informal education (e.g., madrassa/duksi/adult education). More than secondary includes middle-level colleges and universities. Secondary includes respondents who reported vocational training as their highest level of education attended, as well as respondents who reported junior school as their highest level.

Table 14.13C Perceptions of victim blaming and privacy of domestic violence: County level

Percent distribution of households that have heard of people who blame women or girls for the violence they experienced and percent distribution of households that think domestic violence is a private matter, according to county, Kenya Mini DHS 2025–26

County	Heard of victim blaming			Perceive domestic violence to be private			Number of households
	Yes	No	Total	Yes	No	Total	
Mombasa	27.7	72.3	100.0	32.1	67.9	100.0	866
Kwale	16.4	83.6	100.0	61.3	38.7	100.0	318
Kilifi	21.6	78.4	100.0	35.6	64.4	100.0	699
Tana River	6.6	93.4	100.0	5.8	94.2	100.0	143
Lamu	23.5	76.5	100.0	38.9	61.1	100.0	62
Taita Taveta	26.1	73.9	100.0	51.5	48.5	100.0	206
Garissa	20.6	79.4	100.0	51.4	48.6	100.0	145
Wajir	1.2	98.8	100.0	29.9	70.1	100.0	126
Mandera	31.0	69.0	100.0	74.0	26.0	100.0	137
Marsabit	20.6	79.4	100.0	5.3	94.7	100.0	138
Isiolo	31.1	68.9	100.0	47.0	53.0	100.0	87
Meru	44.1	55.9	100.0	56.3	43.7	100.0	840
Tharaka-Nithi	48.6	51.4	100.0	68.0	32.0	100.0	209
Embu	42.5	57.5	100.0	37.9	62.1	100.0	385
Kitui	16.3	83.7	100.0	12.6	87.4	100.0	526
Machakos	21.2	78.8	100.0	25.8	74.2	100.0	1,021
Makueni	30.9	69.1	100.0	31.6	68.4	100.0	506
Nyandarua	39.7	60.3	100.0	43.7	56.3	100.0	341
Nyeri	28.0	72.0	100.0	39.9	60.1	100.0	441
Kirinyaga	46.1	53.9	100.0	48.8	51.2	100.0	482
Murang'a	43.5	56.5	100.0	39.1	60.9	100.0	571
Kiambu	35.9	64.1	100.0	45.6	54.4	100.0	2,295
Turkana	51.2	48.8	100.0	46.4	53.6	100.0	232
West Pokot	36.6	63.4	100.0	51.0	49.0	100.0	251
Samburu	56.8	43.2	100.0	41.7	58.3	100.0	109
Trans Nzoia	58.8	41.2	100.0	38.3	61.7	100.0	393
Uasin Gishu	34.8	65.2	100.0	28.3	71.7	100.0	794
Elgeyo-Marakwet	19.7	80.3	100.0	15.3	84.7	100.0	209
Nandi	45.0	55.0	100.0	34.6	65.4	100.0	414
Baringo	7.4	92.6	100.0	22.7	77.3	100.0	242
Laikipia	33.4	66.6	100.0	38.0	62.0	100.0	293
Nakuru	42.0	58.0	100.0	36.1	63.9	100.0	1,442
Narok	43.9	56.1	100.0	40.6	59.4	100.0	576
Kajiado	47.3	52.7	100.0	37.2	62.8	100.0	653
Kericho	37.8	62.2	100.0	17.1	82.9	100.0	464
Bomet	24.9	75.1	100.0	48.6	51.4	100.0	437
Kakamega	32.4	67.6	100.0	38.4	61.6	100.0	850
Vihiga	11.7	88.3	100.0	22.4	77.6	100.0	267
Bungoma	47.6	52.4	100.0	21.5	78.5	100.0	746
Busia	13.8	86.2	100.0	29.5	70.5	100.0	429
Siaya	40.0	60.0	100.0	52.8	47.2	100.0	482
Kisumu	39.6	60.4	100.0	38.8	61.2	100.0	615
Homa Bay	43.9	56.1	100.0	56.7	43.3	100.0	568
Migori	46.8	53.2	100.0	24.5	75.5	100.0	499
Kisii	53.7	46.3	100.0	57.1	42.9	100.0	597
Nyamira	28.3	71.7	100.0	62.4	37.6	100.0	284
Nairobi City	37.2	62.8	100.0	31.6	68.4	100.0	3,992
Total	35.6	64.4	100.0	37.5	62.5	100.0	26,380

Note: Because this question was administered in the Household Questionnaire, the sex, age, and education variables refer to those of the household respondent.

ADULT AND MATERNAL MORTALITY 15

Key Findings

- ***Adult mortality:*** Overall, the mortality rate among adults age 15–49 is lower among women (1.73 deaths per 1,000 population) than among men (2.83 deaths per 1,000 population).
- ***Maternal mortality ratio:*** The maternal mortality ratio for the 7-year period before the 2025–26 Kenya Mini Demographic and Health Survey (KMDHS) is estimated at 263 maternal deaths per 100,000 live births; that is, for every 1,000 live births in Kenya, about three women die during pregnancy, during childbirth, or within 42 days of the end of a pregnancy from causes other than accidents or violence.
- ***Pregnancy-related mortality:*** The estimated pregnancy-related mortality ratio for the 7-year period preceding the 2025–26 KMDHS is 287 deaths per 100,000 live births; that is, for every 1,000 live births in Kenya, about three women die during pregnancy or within 2 months of the end of a pregnancy from any cause, including accidents or violence.

Adult and maternal mortality indicators can be used to assess the health status of a population. These indicators provide critical evidence for identifying where health programmes are underperforming, including gaps in disease prevention, maternal care, referral systems, emergency response, and quality of services. They also guide planning, resource allocation, county prioritisation, monitoring of national and Sustainable Development Goal targets, and accountability for reducing preventable adult and maternal deaths. In Kenya, reducing preventable deaths among adults, women of reproductive age, mothers, and newborns is anchored in the broader constitutional commitment of attaining the highest standard of health and is operationalised through national health and reproductive health policy frameworks. The Government of Kenya has positioned mortality reduction as a core priority under universal health coverage, emphasising equitable access to quality services, strengthened primary health care, improved referral systems, health workforce capacity, and the use of data for decision-making.

15.1 DATA

To obtain sibling histories for the 2025–26 KMDHS, each female respondent was asked to list all children born to her biological mother, starting with the firstborn. The respondents were then asked whether each sibling was still alive. For living siblings, the current age of each sibling was recorded in the questionnaire. For deceased siblings, the age at death and the number of years since death were recorded. When precise information on age at death or years since death was not available, approximate quantitative responses were accepted. For sisters who died at age 12 or older, additional questions were asked to determine whether the death occurred during pregnancy, during childbirth, within 42 days after the end of pregnancy, or within 2

months after the end of pregnancy or childbirth. This information was used to classify maternal deaths and pregnancy-related deaths and to estimate adult, maternal, and pregnancy-related mortality indicators.

A total of 103,729 siblings were recorded in the sibling histories in the 2025–26 KMDHS, comprising 50,916 sisters and 52,813 brothers (see **Table C.10** in **Appendix C**). Survival status was reported for all but 23 siblings, indicating near-complete reporting. Overall, 96,669 siblings (93%) were reported to be living, while 7,037 (7%) were reported to have died.

The mean sibship size was 5.7 (including the respondents), and the overall sex ratio of siblings at birth was 102.8 brothers per 100 sisters, which falls within the generally expected range (see **Table C.11** in **Appendix C**). These results indicate a high level of completeness in the sibling history data used to estimate adult and maternal mortality in the 2025–26 KMDHS.

15.2 Direct Estimates of Adult Mortality

Adult mortality rate

The number of adult deaths per 1,000 population age 15–49. Adult mortality rates by 5-year age groups are calculated as the number of deaths among a respondent's siblings in each age group divided by the number of person-years of exposure to the risk of dying in that age group during the 7 years preceding the survey. The number of deaths is the number of siblings (brothers or sisters) reported as having died within the 7 years preceding the survey. The person-years of exposure in each age group are calculated for both surviving and dead siblings based on current age (living siblings) or age at death and years since death (dead siblings).


Sample: Siblings (both living and dead) who were age 15–49 in the 7 years preceding the survey, by sex and 5-year age groups

One way to assess the quality of data used to estimate maternal mortality is to evaluate the plausibility of overall adult mortality estimates. Because maternal deaths are a subset of all female adult deaths, implausible patterns in overall adult mortality would raise concerns about the reliability of maternal mortality estimates. Direct estimates of adult mortality in the 2025–26 KMDHS are based on sibling history data reported by interviewed women, including the survival status of their brothers and sisters and, for deceased siblings, their age at death and years since death.

To ensure a sufficient number of adult deaths for generating a robust estimate of adult mortality, rates are calculated for the 7-year period before the survey (roughly between late 2018/early 2019 and late 2025/early 2026). However, age-specific mortality rates obtained in this manner are subject to considerable sampling variation. Using this 7-year period is a compromise between the desire for the most recent data and the need to minimise the level of sampling error.

Table 15.1 and **Figure 15.1** present age-specific mortality rates for women and men age 15–49 for the 7 years preceding the 2025–26 KMDHS. The results show the expected pattern of increasing mortality with age for both sexes. Among women, mortality rises from 0.36 deaths per 1,000 women age 15–19 to 3.23 deaths per 1,000 women age 45–49. Among men, mortality increases from 1.15 deaths per 1,000 men age 15–19 to 6.32 deaths per 1,000 men age 45–49. Overall, adult mortality is higher among men than among women, with an age-adjusted mortality rate of 2.83 deaths per 1,000 men compared with 1.73 deaths per 1,000 women age 15–49.

Figure 15.1 **Adult mortality rates by age**



The results also show a clear sex differential in adult mortality. Male mortality exceeds female mortality in every 5-year age group, with the gap widening at older ages.

15.3 Trends in Adult Mortality

Table 15.2 shows trends, over the past four surveys, in the probability of dying between exact ages 15 and 50 ($_{35}q_{15}$) in the 7 years before the 2025–26, 2014, 2008–09, and 2003 surveys; $_{35}q_{15}$ is the probability that a woman or man who was age 15 in the 7 years before the survey will have died before reaching age 50 (if the age- and gender-specific mortality rates in the 7 years before the survey hold constant).

The probability of a woman dying between ages 15 and 50 declined from 235 per 1,000 women in the 2003 Kenya Demographic and Health Survey (KDHS) to 214 in the 2008–09 KDHS, 138 in the 2014 KDHS, and 66 in the 2025–26 KMDHS. Among men, the corresponding probability declined from 240 per 1,000 men in the 2003 survey to 231 in the 2008–09 survey, 183 in the 2014 survey, and 108 in the 2025–26 survey.

15.4 Direct Estimates of Maternal Mortality

Maternal mortality rate

The number of maternal deaths per 1,000 women age 15–49. Maternal mortality rates by 5-year age groups are calculated by dividing the number of maternal deaths to female siblings of respondents in each age group by the total person-years of exposure of the sisters to the risk of dying in that age group during the 7 years preceding the survey. The number of deaths is the number of sisters reported as having died in the 7 years preceding the survey during either pregnancy or delivery, or in the 42 days following the delivery or termination of a pregnancy, by their age group at the time of death. Deaths due to accidents or violence are excluded. The person-years of exposure in each age group are calculated for both surviving and dead sisters based on their reported current age (living sisters) or age at death and years since death (dead sisters).

Sample: Sisters (both living and dead) age 15–49 in the 7 years preceding the survey, by 5-year age groups

Maternal mortality ratio

The number of maternal deaths per 100,000 live births. The maternal mortality ratio is calculated by dividing the age-standardised maternal mortality rate for women age 15–49 in the 7 years preceding the survey by the general fertility rate (GFR) for the same period.

Maternal deaths are a subset of all female deaths. They are defined as any deaths that occur during pregnancy or childbirth or within 42 days after the delivery or termination of a pregnancy. Maternal deaths do not include deaths due to accidents or violence. Two methods are generally used to estimate maternal mortality in developing countries: the indirect sisterhood method (Graham et al. 1989) and a direct variant of the sisterhood method (Rutenberg and Sullivan 1991; Stanton et al. 1997).

For the 2025–26 KMDHS, the direct estimates of maternal mortality can be described as follows:

- The maternal mortality rate in Kenya is 0.28 maternal deaths per 1,000 woman-years of exposure for the 7-year period preceding the survey (**Table 15.3**).
- The estimated age-specific maternal mortality rate is highest among women age 30–34, at 0.82 maternal deaths per 1,000 woman-years of exposure, and lowest among women age 15–19, at 0.03.
- Maternal deaths represent 17% of all deaths among women age 15–49 during the 7-year period preceding the survey.
- The maternal mortality ratio for the 7-year period before the survey is estimated at 263 maternal deaths per 100,000 live births. This means that for every 100,000 live births in Kenya, about 263 women die during pregnancy, childbirth, or within 42 days after the end of pregnancy, from causes other than accidents or violence. The confidence interval (CI) surrounding the maternal mortality estimate is 121–405 deaths per 100,000 live births, indicating that the estimate should be interpreted with caution (**Table 15.4**).
- The lifetime risk of maternal death is 0.009. In other words, at current fertility and mortality rates, roughly 1 in every 111 women in Kenya will have a maternal death.

Trends: Trends in the maternal mortality ratio should be interpreted carefully because earlier KDHS surveys largely reported pregnancy-related mortality, which is broader than maternal mortality. Pregnancy-related mortality includes deaths during pregnancy, childbirth, or within 2 months after the end of pregnancy, regardless of cause, while maternal mortality excludes deaths due to accidents or violence and uses the 42-day period after the end of pregnancy. Therefore, the 2025–26 maternal mortality ratio of 263 deaths per 100,000 live births should not be directly compared with earlier pregnancy-related mortality estimates. For trend analysis, the pregnancy-related mortality ratio declined from 743 deaths per 100,000 live births in the 7 years preceding the 1998 KDHS to 287 deaths per 100,000 live births in the 7 years preceding the 2025–26 KMDHS, suggesting a long-term decline in pregnancy-related mortality in Kenya.

15.5 TRENDS IN PREGNANCY-RELATED MORTALITY

Pregnancy-related mortality rate

The number of pregnancy-related deaths per 1,000 women age 15–49. Pregnancy-related mortality rates by 5-year age groups are calculated by dividing the number of pregnancy-related deaths to female siblings of respondents in each age group by the total person-years of exposure of the sisters to the risk of dying in that age group during the 7 years preceding the survey. The number of deaths is the number of sisters reported as having died in the 7 years preceding the survey during either pregnancy or delivery, or in the 2 months following the delivery or termination of a pregnancy, by their age group at the time of death. The person-years of exposure in each age group are calculated for both surviving and dead sisters based on their reported current age (living sisters) or age at death and years since death (dead sisters).

Sample: Sisters (both living and dead) age 15–49 in the 7 years preceding the survey, by 5-year age groups

Pregnancy-related mortality ratio


The number of pregnancy-related deaths per 100,000 live births. The pregnancy-related mortality ratio is calculated by dividing the age-standardized pregnancy-related mortality rate for women age 15–49 in the 7 years preceding the survey by the general fertility rate (GFR) for the same time period.

To allow comparisons with estimates from previous KDHS surveys, the 2025–26 KMDHS defines a pregnancy-related death as the death of a woman during pregnancy or childbirth or within 2 months of delivery or termination of a pregnancy, irrespective of the cause of death. Estimates of pregnancy-related mortality are therefore based solely on the timing of the death in relation to the pregnancy. Note that this definition varies from the World Health Organization definition of a pregnancy-related death, which limits the window to 42 days. What the 2025–26 KMDHS defines as a pregnancy-related death had been labelled a maternal death in the 2014 KDHS and prior surveys.

The estimated pregnancy-related mortality ratio (PRMR) for the 7-year period preceding the 2025–26 KMDHS is 287 deaths per 100,000 live births. This means that for every 1,000 births in Kenya, about three women die during pregnancy, childbirth, or within 2 months of the end of a pregnancy, from any cause including accidents or violence.

Overall, the PRMR has trended downward over time. The PRMR declined from 506 deaths per 100,000 live births in the 7 years preceding the 2003 KDHS to 520 in the 7 years preceding the 2008–09 KDHS, 362 in the 7 years preceding the 2014 KDHS, and 287 in the 7 years preceding the 2025–26 KMDHS (**Figure 15.2**). However, the CIs for the 2025–26 estimate (287; CI: 144–431) and the 2014 estimate (362; CI: 254–471) overlap, meaning that the difference between the last two surveys is not statistically significant.

Figure 15.2 **Trends in the pregnancy-related mortality ratio (PRMR) with confidence intervals**



List of Tables

For more information on adult and maternal mortality, see the following tables:

- **Table 15.1 Adult mortality rates**
- **Table 15.2 Adult mortality probabilities**
- **Table 15.3 Maternal mortality rates**
- **Table 15.4 Maternal mortality ratio**

Table 15.1 Adult mortality rates

Direct estimates of female and male mortality rates for the 7 years preceding the survey, by 5-year age groups, Kenya Mini DHS 2025–26

Age	Number of deaths	Exposure years	Mortality rate[1]
FEMALE			
15–19	15	40,364	0.36
20–24	38	46,422	0.81
25–29	80	43,847	1.82
30–34	91	40,661	2.24
35–39	75	34,167	2.18
40–44	74	24,174	3.04
45–49	50	15,446	3.23
Total 15–49	421	245,081	1.73[a]
MALE			
15–19	47	40,606	1.15
20–24	67	46,211	1.45
25–29	113	45,240	2.50
30–34	109	40,943	2.65
35–39	126	33,790	3.73
40–44	122	24,386	5.01
45–49	101	15,993	6.32
Total 15–49	684	247,169	2.83[a]

[1] Expressed per 1,000 population
[a] Age-adjusted rate

Table 15.2 Adult mortality probabilities

The probability of dying between the ages of 15 and 50 for women and men during the 7 years preceding the survey, Kenya Mini DHS 2025–26

Survey	Female ${}_{35}q_{15}$[1]	Male ${}_{35}q_{15}$[1]
2025–26 Kenya Mini DHS	66	108
2014 KDHS	138	183
2008–09 KDHS	214	231
2003 KDHS	235	240

KDHS = Kenya Demographic and Health Survey
[1] The probability of dying between exact ages 15 and 50, expressed per 1,000 persons at age 15

Table 15.3 Maternal mortality rates

Direct estimates of maternal mortality rates for the 7 years preceding the survey, by 5-year age groups, Kenya Mini DHS 2025–26

Age	Percentage of female deaths that are maternal	Number of maternal deaths[1]	Exposure years	Maternal mortality rate[2]
15–19	9.6	1	40,364	0.03
20–24	12.5	5	46,422	0.10
25–29	17.2	14	43,847	0.31
30–34	36.6	33	40,661	0.82
35–39	12.7	9	34,167	0.28
40–44	10.6	8	24,174	0.32
45–49	4.7	2	15,446	0.15
Total 15–49	17.3	73	245,081	0.28[a]

[1] A maternal death is defined as the death of a woman while pregnant or within 42 days of termination of pregnancy, from any cause except accidents or violence.
[2] Expressed per 1,000 woman-years of exposure
[a] Age-adjusted rate

Table 15.4 Maternal mortality ratio

Total fertility rate, general fertility rate, maternal mortality ratio, and lifetime risk of maternal death for the 7 years preceding the survey, Kenya Mini DHS 2025–26

Total fertility rate (TFR)	3.3
General fertility rate (GFR)[1]	107
Maternal mortality ratio (MMR)[2]	263 (CI: 121, 405)
Lifetime risk of maternal death[3]	0.009

CI = Confidence interval
[1] Age-adjusted rate expressed per 1,000 women age 15–49
[2] Expressed per 100,000 live births; calculated as the age-adjusted maternal mortality rate (shown in Table 15.3) times 100 divided by the age-adjusted general fertility rate
[3] Calculated as $1-(1-MMR)^{TFR}$ where TFR represents the total fertility rate for the 7 years preceding the survey

REFERENCES

Ashraf, S. A. 2025. Food Fortification as a Sustainable Global Strategy to Mitigate Micronutrient Deficiencies and Improve Public Health. *Discover Food* 5(201). https://link.springer.com/article/10.1007/s44187-025-00512-5.

Ayele, D. G., and S. F. Melesse. 2017. Proximate Determinants of Fertility in Eastern Africa: The Case of Kenya, Rwanda and Tanzania. *Scientific Review* 3(4):29–42.

Bradley, S. E. K., T. N. Croft, J. D. Fishel, and C. F. Westoff. 2012. *Revising Unmet Need for Family Planning*. DHS Analytical Studies No. 25. Calverton, Maryland, USA: ICF International. https://dhsprogram.com/pubs/pdf/AS25/AS25%5B12June2012%5D.pdf.

Dewey, K. 2003. *Guiding Principles for Complementary Feeding of the Breastfed Child.* Washington, DC: Pan American Health Organization. https://iris.paho.org/handle/10665.2/752.

Government of Kenya. 2017. *Family Planning 2020 Commitment.* Government of Kenya. https://www.fp2030.org/app/uploads/2023/08/Kenya_FP2020_Commitment_2017_2.pdf.

Graham, W., W. Brass, and R. W. Snow. 1989. Indirect Estimation of Maternal Mortality: The Sisterhood Method. *Studies in Family Planning* 20(3):125–135. https://doi.org/10.2307/1966567.

Haider, B. A., I. Olofin, M. Wang, D. Spiegelman, M. Ezzati, and W. W. Fawzi. Anaemia, Prenatal Iron Use, and Risk of Adverse Pregnancy Outcomes: A Systemic Review and Meta-analysis. *BMJ* 346:f3443. https://doi.org/10.1136/bmj.f3443.

Ministry of Health [Kenya]. 2022a. *Kenya Health Information System and Civil Registration Guidelines.* Ministry of Health.

Ministry of Health [Kenya]. 2022b. *Maternal and Newborn Health Standards for Improving Quality of Care in Health Facilities.* Ministry of Health. https://www.scribd.com/document/880238488/1-Maternal-Newborn-Health-Standards-2-3-22-2.

Ministry of Health [Kenya]. 2024a. *Kenya Malaria Strategy 2023–2027: Towards a Malaria-free Kenya.* Nairobi: National Malaria Control Programme. https://mesamalaria.org/wp-content/uploads/2025/11/KENYA-MALARIA-STRATEGY_2023-2027.pdf.

Ministry of Health [Kenya]. 2024b. *National Guidelines on Quality Obstetrics and Perinatal Care.* Nairobi: Ministry of Health. https://studoke.com/details/1429/.

Moran, A. C., K. Kerber, D. Sitrin, et al. 2013. Measuring Coverage in MNCH: Indicators for Global Tracking of Newborn Care. *PLoS Med* 10(5):e1001415. https://doi.org/10.1371/journal.pmed.1001415.

Republic of Kenya. 1928. "Births and Deaths Registration Act: Cap. 149." *Kenya Gazette,* June 15, 1928. https://kenyalaw.org/akn/ke/act/1928/2/eng@2026-03-10.

Republic of Kenya. 2023. "Social Health Insurance Act: No. 6 of 2023." *Kenya Gazette,* November 24, 2023. https://new.kenyalaw.org/akn/ke/act/2023/16/eng@2023-11-24.

Republic of Kenya. 2025. *Report of the Technical Working Group on Gender-based Violence (GBV) Including Femicide*. Republic of Kenya. https://africa.unwomen.org/sites/default/files/2026-02/technical_working_group_report_on_gbv_including_femicide_0.pdf.

Rutenberg, N., and J. Sullivan. 1991. Direct and Indirect Estimates of Maternal Mortality from the Sisterhood Method. *Proceedings of the Demographic and Health Surveys World Conference* 3:1669–1696. Columbia, Maryland, USA: Institute for Research Development/Macro International Inc.

Shallo, S. A. 2020. Roles of Proximate Determinants of Fertility in Recent Fertility Decline in Ethiopia: Application of the Revised Bongaarts Model. *Open Access Journal of Contraception* 11:33–41. https://doi.org/10.2147/oajc.s251693.

Shonkoff, J. P., and D. A. Phillips, eds. 2000. *From Neurons to Neighborhoods: The Science of Early Childhood Development.* Washington, DC: National Academies Press. https://pubmed.ncbi.nlm.nih.gov/25077268/.

Stanton, C., N. Abderrahim, and K. Hill. 1997. *DHS Maternal Mortality Indicators: An Assessment of Data Quality and Implications for Data Use*. DHS Analytical Reports No. 4. Calverton, Maryland, USA: Macro International Inc.

United Nations. 2015. *Transforming Our World: The 2030 Agenda for Sustainable Development.* United Nations General Assembly. https://docs.un.org/en/A/RES/70/1.

United Nations Children's Fund (UNICEF). 2016. *Advancing Early Childhood Development: From Science to Scale.* Executive Summary. https://www.thelancet.com/pb-assets/Lancet/stories/series/ecd/Lancet_ECD_Executive_Summary.pdf.

United Nations Children's Fund (UNICEF). 2021. "Diarrhoea Remains a Leading Killer of Young Children, Despite the Availability of a Simple Treatment." Accessed 21 August 2026. https://data.unicef.org/topic/child-health/diarrhoeal-disease.

United Nations Children's Fund (UNICEF). 2024. *Large-scale Food Fortification for the Prevention of Micronutrient Deficiencies in Children, Women and Communities: Guidance Note.* New York: UNICEF. https://www.scribd.com/document/868260966/Unicef-Ff-Guidance-V8.

United Nations Population Fund (UNFPA). n.d. "Measuring Technology-facilitated Gender-based Violence: A Growing Threat." Accessed 24 August 2026. https://www.unfpa.org/TFGBV.

United Nations Population Fund (UNFPA). 2021. *Technology-facilitated Gender-based Violence: Making All Spaces Safe.* New York: UNFPA. https://www.unfpa.org/publications/technology-facilitated-gender-based-violence-making-all-spaces-safe.

United Nations Statistics Division. 2013. *Principles and Recommendations for a Vital Statistics System*: Revision 3. New York: United Nations. https://digitallibrary.un.org/record/1474612?v=pdf.

UN Women. 2025. *In Brief: Repository of UN Women's Work on Technology-facilitated Violence against Women and Girls.* New York: UN Women. https://www.unwomen.org/sites/default/files/2025-03/repository-of-un-womens-work-on-technology-facilitated-violence-against-women-and-girls-en.pdf.

World Health Organization (WHO). 2001. *Putting Women First: Ethical and Safety Recommendations for Research on Domestic Violence against Women.* Geneva: WHO. https://www.who.int/publications/i/item/WHO-FCH-GWH-01.1.

World Health Organization (WHO). 2006. *Birth Spacing: Report from a WHO Technical Consultation.* Policy Brief (Department of Reproductive Health and Research). Geneva: WHO. https://iris.who.int/server/api/core/bitstreams/de7cfb95-b1c8-4286-aadf-5833fd3cab81/content.

World Health Organization (WHO). 2011a. *Guideline: Intermittent Iron Supplementation in Preschool and School-age Children.* Geneva: WHO. https://www.who.int/publications/i/item/9789241502009.

World Health Organization (WHO). 2011b. *Guideline: Vitamin A Supplementation in Infants and Children 6–59 Months of Age.* Geneva: WHO. https://www.who.int/publications/i/item/9789241501767.

World Health Organization (WHO). 2013. *Guideline: Updates on the Management of Severe Acute Malnutrition in Infants and Children.* Geneva: WHO. https://www.who.int/publications/i/item/9789241506328.

World Health Organization (WHO). 2015. *WHO Statement on Caesarean Section Rates.* Geneva: WHO. https://iris.who.int/server/api/core/bitstreams/475b39fc-240b-462d-a5e1-429401c98a79/content.

World Health Organization (WHO). 2016a. *Guideline: Daily Iron Supplementation in Infants and Children.* Geneva: WHO. https://www.who.int/publications/i/item/9789241549523.

World Health Organization (WHO). 2016b. *Guideline: Use of Multiple Micronutrient Powders for Point-of-use Fortification of Foods Consumed by Infants and Young Children Aged 6–23 Months and Children Aged 2–12 Years.* Geneva: WHO. https://www.who.int/publications/i/item/9789241549943.

World Health Organization (WHO). 2017a. *Guideline: Assessing and Managing Children at Primary Health-care Facilities to Prevent Overweight and Obesity in the Context of the Double Burden of Malnutrition. Updates for the Integrated Management of Childhood Illness (IMCI).* Geneva: WHO. https://www.who.int/publications/i/item/9789241550123.

World Health Organization (WHO). 2017b. *Guideline: Preventive Chemotherapy to Control Soil-transmitted Helminth Infections in At-risk Population Groups.* Geneva: WHO. https://www.who.int/publications/i/item/9789241550116.

World Health Organization (WHO). 2017c. *Robson Classification: Implementation Manual.* Geneva: WHO. https://www.who.int/publications/i/item/9789241513197.

World Health Organization (WHO). 2018. *Guideline: Counselling of Women to Improve Breastfeeding Practices.* Geneva: WHO. https://www.who.int/publications/i/item/9789241550468.

World Health Organization (WHO). 2021. *Putting Women First: Ethical and Safety Recommendations for Research on Domestic Violence against Women.* Geneva: WHO. https://www.who.int/publications/i/item/WHO-FCH-GWH-01.1.

World Health Organization (WHO). 2022. *WHO Recommendations on Maternal and Newborn Care for a Positive Postnatal Experience.* Geneva: WHO. https://www.who.int/publications/i/item/9789240045989.

World Health Organization (WHO). 2024. *World Health Statistics 2024: Monitoring Health for the SDGs, Sustainable Development Goals.* Geneva: WHO. https://www.who.int/publications/i/item/9789240094703.

World Health Organization (WHO). 2025. *World Malaria Report 2025: Addressing the Threat of Antimalarial Drug Resistance*. Geneva: WHO. https://www.who.int/publications/i/item/9789240117822.

World Health Organization (WHO)/United Nations Children's Fund (UNICEF) Joint Monitoring Programme for Water Supply, Sanitation and Hygiene. 2018. *Core Questions on Water, Sanitation and Hygiene for*

Household Surveys: 2018 Update. New York: UNICEF and WHO. JMP-2018-core-questions-for-household-surveys-1.pdf.

World Health Organization (WHO) and the United Nations Children's Fund. 2014. *Every Newborn: An Action Plan to End Preventable Deaths*. Geneva: WHO. https://apps.who.int/iris/bitstream/handle/10665/127938/9789241507448_eng.pdf;sequence=1.

World Health Organization (WHO) and the United Nations Children's Fund. 2019. *Recommendations for Data Collection, Analysis and Reporting on Anthropometric Indicators in Children under 5 Years Old.* Geneva: WHO. https://www.who.int/publications/i/item/9789241515559.

World Health Organization (WHO) and the United Nations Children's Fund (UNICEF). 2021. *Indicators for Assessing Infant and Young Child Feeding Practices: Definitions and Measurement Methods.* Geneva: WHO. https://www.who.int/publications/i/item/9789240018389.

SAMPLE DESIGN

Appendix A

A.1 INTRODUCTION

This section describes the objectives of the survey and the sample design, including the overall sample size, survey domains, and sampling procedures. The 2025–26 Kenya Mini Demographic and Health Survey (KMDHS) is the first of its kind carried out in Kenya following a successful implementation of the seventh round of the Kenya Demographic and Health Survey (KDHS) in 2022. The primary objective of the 2025–26 KMDHS was to provide timely and accurate data on key demographic and health indicators to inform, design, implement, monitor, and evaluate population, health, and development programmes at both national and county levels and to allow planners and policymakers to adjust programmes and policies based on more current data. A nationally representative sample of 1,000 clusters and 30,000 households has been used for the 2025–26 KMDHS. As with the full KDHS survey, all women age 15–49 who were usual residents of the selected households or who slept in the households the night before the survey were eligible for interviews. An estimated 22,932 eligible women were expected to be interviewed during the survey. The 2025–26 KMDHS was designed to provide reliable estimates at the national level; for rural and urban areas separately; and, for some key indicators, at the county level for all 47 counties.

A.2 SAMPLING FRAME

The sample for the 2025–26 KMDHS was drawn from the Kenya Household Master Sample Frame (K-HMSF), which was developed and maintained by the Kenya National Bureau of Statistics for household-based sample surveys in Kenya. The K-HMSF, based on the enumeration areas (EAs) used in the 2019 Kenya Population and Housing Census (KPHC), comprises 10,000 clusters selected using probability proportional to size. The 10,000 clusters were randomised into four equal subsamples, allowing surveys to utilise one or more subsamples depending on their sample size requirements. The 2025–26 KMDHS sample was drawn from subsample four of the frame. To design the frame, each of the 47 counties in Kenya was stratified into rural and urban, resulting in 92 strata, as Nairobi City and Mombasa Counties are purely urban. The K-HMSF is representative up to the county level.

A total of 12,040,701 households were renumerated in the 2019 KPHC. **Table A.1** shows the distribution of the households and population by county and type of residence based on the 2019 census data. Most (61%) of the households are in rural areas, while 39% are in urban areas, with Nairobi City and Mombasa Counties being purely urban. The share of households in a county relative to the total household population in Kenya varies from 0.3% in Lamu County to 9.2% in Nairobi City County.

Table A.1 Population and household distribution

Distribution of population and households by county and type of residence, Kenya Mini DHS 2025–26

County	Population			Number of households			Share of county households to total number of households	Share of county households that are urban
	Total	Rural	Urban	Total	Rural	Urban		
Mombasa	1,188,639	na	1,188,639	376,212	na	376,212	2.5	100.0
Kwale	858,263	735,530	122,733	172,767	135,863	36,904	1.8	14.3
Kilifi	1,440,289	1,052,712	387,577	297,935	185,363	112,572	3.1	26.9
Tana River	313,384	236,481	76,903	66,964	47,210	19,754	0.7	24.5
Lamu	138,148	100,099	38,049	34,223	23,909	10,314	0.3	27.5
Taita-Taveta	333,643	242,667	90,976	94,453	65,005	29,448	0.7	27.3
Garissa	832,910	625,228	207,682	138,917	101,992	36,925	1.8	24.9
Wajir	774,213	599,554	174,659	126,874	96,880	29,994	1.6	22.6
Mandera	860,132	594,542	265,590	123,935	84,711	39,224	1.8	30.9
Marsabit	446,157	341,625	104,532	76,669	56,445	20,224	0.9	23.4
Isiolo	263,105	138,848	124,257	53,206	25,348	27,858	0.6	47.2
Meru	1,532,938	1,398,935	134,003	423,848	373,233	50,615	3.3	8.7
Tharaka-Nithi	390,825	358,966	31,859	109,425	97,594	11,831	0.8	8.2
Embu	604,407	531,232	73,175	182,413	154,290	28,123	1.3	12.1
Kitui	1,128,899	1,077,561	51,338	261,787	242,581	19,206	2.4	4.5
Machakos	1,410,901	1,002,760	408,141	399,453	256,789	142,664	3.0	28.9
Makueni	976,222	905,404	70,818	243,948	217,026	26,922	2.1	7.3
Nyandarua	634,486	568,540	65,946	178,205	154,989	23,216	1.3	10.4
Nyeri	749,170	604,757	144,413	244,542	189,078	55,464	1.6	19.3
Kirinyaga	604,986	472,185	132,801	203,554	154,238	49,316	1.3	22.0
Murang'a	1,052,386	936,014	116,372	317,459	276,003	41,456	2.2	11.1
Kiambu	2,399,479	708,414	1,691,065	792,147	207,522	584,625	5.1	70.5
Turkana	919,826	781,918	137,908	162,567	132,583	29,984	2.0	15.0
West Pokot	618,341	587,431	30,910	115,738	107,145	8,593	1.3	5.0
Samburu	305,987	260,478	45,509	63,947	51,454	12,493	0.6	14.9
Trans Nzoia	984,462	809,352	175,110	222,967	173,565	49,402	2.1	17.8
Uasin Gishu	1,148,713	647,990	500,723	301,065	139,935	161,130	2.4	43.6
Elgeyo-Marakwet	452,613	432,410	20,203	99,104	93,009	6,095	1.0	4.5
Nandi	883,188	825,109	58,079	199,024	180,495	18,529	1.9	6.6
Baringo	662,087	588,333	73,754	141,867	119,355	22,512	1.4	11.1
Laikipia	510,356	388,618	121,738	145,764	103,013	42,751	1.1	23.9
Nakuru	2,124,581	1,104,869	1,019,712	598,156	269,498	328,658	4.5	48.0
Narok	1,146,240	1,048,539	97,701	238,071	206,255	31,816	2.4	8.5
Kajiado	1,104,068	489,663	614,405	313,130	106,510	206,620	2.3	55.6
Kericho	896,591	806,695	89,896	205,890	175,089	30,801	1.9	10.0
Bomet	872,553	846,721	25,832	187,217	177,874	9,343	1.9	3.0
Kakamega	1,860,113	1,678,502	181,611	432,191	377,522	54,669	3.9	9.8
Vihiga	587,039	529,433	57,606	143,265	127,378	15,887	1.2	9.8
Bungoma	1,662,696	1,476,329	186,367	357,687	307,128	50,559	3.5	11.2
Busia	886,588	777,915	108,673	197,921	167,171	30,750	1.9	12.3
Siaya	988,118	904,706	83,412	249,279	224,358	24,921	2.1	8.4
Kisumu	1,140,672	711,265	429,407	296,766	168,767	127,999	2.4	37.6
Homabay	1,124,025	1,014,692	109,333	260,262	229,154	31,108	2.4	9.7
Migori	1,106,776	943,024	163,752	238,078	195,326	42,752	2.3	14.8
Kisii	1,259,608	1,112,307	147,301	307,224	261,359	45,865	2.7	11.7
Nyamira	602,859	553,576	49,283	150,489	136,583	13,906	1.3	8.2
Nairobi City	4,323,552	na	4,323,552	1,494,096	na	1,494,096	9.2	100.0
Kenya	**47,105,234**	**32,551,929**	**14,553,305**	**12,040,701**	**7,376,595**	**4,664,106**	**100.0**	**na**

Source: 2019 Kenya Population and Housing Census
na = not applicable

Table A.2 indicates the distribution of EAs developed for the 2019 KPHC and the number of EAs in the K-HMSF. In total, 128,239 EAs were developed for the census, of which 79,115 were rural and 49,124 were urban. The K-HMSF, which was also used as the sampling frame for the 2022 KDHS, contains 5,688 rural EAs and 4,312 urban EAs.

Table A.2 Distribution of enumeration areas

Distribution of enumeration areas by county and type of residence, Kenya Mini DHS 2025–26

	Total number of enumeration areas			Number of enumeration areas in the K-HMSF		
County	Total	Rural	Urban	Total	Rural	Urban
Mombasa	4,031	na	4,031	220	na	220
Kwale	1,827	1,462	365	208	124	84
Kilifi	3,112	1,971	1,141	208	108	100
Tana River	833	607	226	220	124	96
Lamu	349	250	99	208	120	88
Taita-Taveta	1,009	710	299	200	112	88
Garissa	1,312	871	441	220	128	92
Wajir	1,297	977	320	220	128	92
Mandera	1,287	901	386	220	124	96
Marsabit	866	651	215	208	120	88
Isiolo	505	232	273	208	104	104
Meru	4,263	3,771	492	228	148	80
Tharaka-Nithi	1,234	1,109	125	200	132	68
Embu	1,818	1,555	263	200	128	72
Kitui	3,731	3,507	224	208	140	68
Machakos	4,133	2,720	1,413	208	112	96
Makueni	2,778	2,481	297	208	136	72
Nyandarua	1,879	1,637	242	200	128	72
Nyeri	2,520	1,960	560	208	124	84
Kirinyaga	1,947	1,490	457	208	120	88
Murang'a	3,199	2,793	406	208	132	76
Kiambu	8,097	2,085	6,012	240	100	140
Turkana	1,694	1,343	351	208	128	80
West Pokot	1,581	1,486	95	208	140	68
Samburu	697	559	138	200	120	80
Trans Nzoia	2,233	1,770	463	208	124	84
Uasin Gishu	3,355	1,492	1,863	232	112	120
Elgeyo-Marakwet	1,163	1,089	74	200	140	60
Nandi	2,094	1,893	201	208	140	68
Baringo	1,897	1,606	291	212	132	80
Laikipia	1,552	1,076	476	208	116	92
Nakuru	6,134	2,703	3,431	236	116	120
Narok	2,620	2,245	375	200	128	72
Kajiado	3,366	1,094	2,272	200	92	108
Kericho	2,165	1,829	336	208	132	76
Bomet	1,933	1,818	115	208	148	60
Kakamega	4,605	3,963	642	212	136	76
Vihiga	1,401	1,241	160	200	132	68
Bungoma	3,732	3,175	557	212	136	76
Busia	2,063	1,727	336	208	128	80
Siaya	2,676	2,396	280	208	136	72
Kisumu	2,933	1,711	1,222	220	116	104
Homabay	2,888	2,526	362	208	136	72
Migori	2,585	2,124	461	208	128	80
Kisii	3,526	2,968	558	224	140	84
Nyamira	1,698	1,541	157	208	140	68
Nairobi City	15,621	na	15,621	300	na	300
Kenya	**128,239**	**79,115**	**49,124**	**10,000**	**5,688**	**4,312**

Source: 2019 Kenya Population and Housing Census
K-HMSF = Kenya Household Master Sample Frame
na = not applicable

A.3 SAMPLE DESIGN AND IMPLEMENTATION

The 2025–26 KMDHS used a two-stage stratified sample design. In the first stage, 1,000 clusters were selected independently in each sampling stratum from the K-HMSF using the equal probability Systematic Sampling Method. Households were relisted in 616 of the 1,000 selected clusters to update household information. The resultant list of households served as the sampling frame for the second stage of selection, in which 30 households were systematically selected from each cluster. The household listing was fully computerised using computer-assisted personal interviewing with fully interactive digital maps; listing data were transmitted to a central server in real time for processing.

Table A.3 shows the distribution of sampled clusters and households by county and type of residence, and **Table A.4** shows the distribution of the expected number of completed interviews with women age 15–49 for each of the 47 counties. To ensure that the precision of the survey would be comparable across counties, a power allocation was used to allocate the sample to each sampling stratum. The survey sample size was computed at 30,000 conventional households and 1,000 clusters, with 354 clusters in urban areas and 646 clusters in rural areas. The sample was expected to result in approximately 22,932 completed interviews with women age 15–49. The expected number of completed interviews was estimated based on the findings of the 2022 KDHS regarding the average number of eligible women per household and the questionnaire completion rates for the household and individual questionnaires.

Table A.3 Sample allocation of clusters and households by county

Sample allocation of clusters and households by county and residence, Kenya Mini DHS 2025–26

	Allocation of clusters			Allocation of households		
County	Urban	Rural	Total	Urban	Rural	Total
Mombasa	23	na	23	690	na	690
Kwale	6	14	20	180	420	600
Kilifi	9	13	22	270	390	660
Tana River	6	12	18	180	360	540
Lamu	5	13	18	150	390	540
Taita-Taveta	7	11	18	210	330	540
Garissa	7	12	19	210	360	570
Wajir	6	13	19	180	390	570
Mandera	7	12	19	210	360	570
Marsabit	6	12	18	180	360	540
Isiolo	8	10	18	240	300	540
Meru	6	20	26	180	600	780
Tharaka-Nithi	4	14	18	120	420	540
Embu	5	15	20	150	450	600
Kitui	4	18	22	120	540	660
Machakos	11	15	26	330	450	780
Makueni	5	17	22	150	510	660
Nyandarua	5	15	20	150	450	600
Nyeri	7	15	22	210	450	660
Kirinyaga	7	14	21	210	420	630
Murang'A	6	17	23	180	510	690
Kiambu	18	12	30	540	360	900
Turkana	6	14	20	180	420	600
West Pokot	4	15	19	120	450	570
Samburu	5	13	18	150	390	540
Trans Nzoia	7	14	21	210	420	630
Uasin Gishu	11	11	22	330	330	660
Elgeyo-Marakwet	4	14	18	120	420	540
Nandi	4	16	20	120	480	600
Baringo	5	14	19	150	420	570
Laikipia	7	12	19	210	360	570
Nakuru	14	12	26	420	360	780
Narok	5	16	21	150	480	630
Kajiado	13	9	22	390	270	660
Kericho	5	15	20	150	450	600
Bomet	4	16	20	120	480	600
Kakamega	6	20	26	180	600	780
Vihiga	4	15	19	120	450	570

Continued…

Table A.3—*Continued*

County	Allocation of clusters			Allocation of households		
	Urban	Rural	Total	Urban	Rural	Total
Bungoma	6	17	23	180	510	690
Busia	5	15	20	150	450	600
Siaya	5	17	22	150	510	660
Kisumu	10	12	22	300	360	660
Homa Bay	5	17	22	150	510	660
Migori	6	15	21	180	450	630
Kisii	6	17	23	180	510	690
Nyamira	4	16	20	120	480	600
Nairobi City	35	na	35	1,050	na	1,050
Kenya	**354**	**646**	**1,000**	**10,620**	**19,380**	**30,000**

na = Not applicable

Table A.4 Expected number of women 15–49 interviews

Sample allocation of expected number of completed interviews for women by county and residence, Kenya Mini DHS 2025–26

County	Women 15–49		
	Urban	Rural	Total
Mombasa	530	na	530
Kwale	151	351	502
Kilifi	217	313	530
Tana River	139	280	419
Lamu	126	329	455
Taita-Taveta	123	193	316
Garissa	180	309	489
Wajir	173	374	547
Mandera	188	323	511
Marsabit	120	240	360
Isiolo	181	227	408
Meru	114	382	496
Tharaka-Nithi	78	275	353
Embu	101	301	402
Kitui	89	403	492
Machakos	243	332	575
Makueni	120	409	529
Nyandarua	102	304	406
Nyeri	123	265	388
Kirinyaga	146	291	437
Murang'A	109	307	416
Kiambu	344	229	573
Turkana	133	311	444
West Pokot	106	400	506
Samburu	112	291	403
Trans Nzoia	169	336	505
Uasin Gishu	261	261	522
Elgeyo-Marakwet	88	305	393
Nandi	99	395	494
Baringo	122	341	463
Laikipia	135	232	367
Nakuru	328	282	610
Narok	124	397	521
Kajiado	271	188	459
Kericho	130	390	520
Bomet	107	427	534
Kakamega	154	512	666
Vihiga	101	381	482
Bungoma	163	464	627
Busia	132	395	527
Siaya	113	383	496
Kisumu	247	296	543
Homa Bay	119	403	522
Migori	156	389	545
Kisii	138	391	529
Nyamira	87	349	436
Nairobi City	684	na	684
Kenya	**7,976**	**14,956**	**22,932**

na = Not applicable

A.4 SAMPLE PROBABILITIES AND SAMPLING WEIGHTS

The 2025–26 KMDHS sample was allocated nonproportionally across sampling strata, plus the probability proportional to size selection of clusters in the K-HMSF and nonresponses resulted in unequal selection probabilities for individuals/households. The sample was therefore not self-weighting, and all data analysis using 2025–26 KMDHS data should be weighted to account for both unequal selection probabilities and nonresponse, ensuring representative survey estimates. Since the 2025–26 KMDHS sample was a two-stage and two-phase stratified cluster sample selected from a master sample, sampling weights were calculated based on sampling probabilities for each sampling stage and phase and for each cluster. The following notations were used:

P_{0hi}: first-phase and first-stage sampling probability of the ith EA in stratum h in the selection of the master sample
P_{1hi}: the second-phase sampling probability of the ith cluster in stratum h from the master sample
P_{2hi}: the second-stage sampling probability within the ith cluster (households)

In the first phase, sample EAs of the master sample were selected with a stratified probability proportional to size procedure. Let a_h be the number of EAs selected in stratum h, M_{hi} the measure of size (number of households) of the ith EA according to the sampling frame, and $\sum M_{hi}$ the total measure of size (total number of households) in stratum h. The probability of selecting the ith EA in the master sample in the first phase and first stage was calculated as follows:

$$P_{0hi} = \frac{a_h M_{hi}}{\sum M_{hi}}$$

Let b_h be the number of clusters selected for the 2025–26 KMDHS in the second phase in stratum h of the master sample. Then the probability of selecting EA i in the sample was:

$$P_{1hi} = \frac{b_h}{a_h}$$

Let L_{hi} be the number of households listed in the household listing operation in cluster i of stratum h, and let g_{hi} be the number of households selected in the cluster. The second stage's selection probability for each household in the cluster was calculated as follows:

$$P_{2hi} = \frac{g_{hi}}{L_{hi}}$$

The overall selection probability of each household in cluster i of stratum h in the 2025–26 KMDHS was therefore the product of the selection probabilities:

$$P_{hi} = P_{0hi} \times P_{1hi} \times P_{2hi} = \frac{b_h M_{hi}}{\sum M_{hi}} \times \frac{g_{hi}}{L_{hi}}$$

The design weight for each household in cluster *i* of stratum *h* was the inverse of its overall selection probability:

$$W_{hi} = 1 / P_{hi}$$

Design weights were adjusted for household nonresponse and individual nonresponse to obtain the sampling weights for households and for women. Nonresponse was adjusted at the sampling stratum level to minimise variability in weights. For the household sampling weight, the household design weight was multiplied by the inverse of the household response rate, by stratum. For the women's individual sampling weight, the household sampling weight was multiplied by the inverse of the women's individual response rate, by stratum. After adjusting for nonresponse, the sampling weights were normalised to obtain the final standard weights that appear in the data files. The normalisation process was aimed at obtaining a total number of unweighted cases equal to the total number of weighted cases using normalised weights at the national level, for the total number of households and for women. Normalisation was achieved by multiplying the sampling weight by the estimated total sampling fraction obtained from the survey, separately for the household weight and for the individual woman's weight. The normalised weights are relative weights that are valid for estimating means, proportions, ratios, and rates but are not valid for estimating population totals or analysing pooled data.

A.4 SURVEY RESULTS

Table A.5 Sample implementation

Percent distribution of households and eligible women age 15–49 by results of the household and individual interviews; and household, eligible women's, and overall women's response rates, according to residence and county (unweighted), Kenya Mini DHS 2025–26

	Selected households												Eligible women										
Result	Completed (C)	Household present but no competent respondent at home (HP)	Postponed (P)	Refused (R)	Dwelling not found (DNF)	Household absent (HA)	Dwelling vacant/ address not a dwelling (DV)	Dwelling destroyed (DD)	Other (O)	Total	Number of sampled households	Household response rate (HRR)[1]	Completed (EWC)	Not at home (EWNH)	Postponed (EWP)	Refused (EWR)	Partly Completed (EWPC)	Incapacitated (EWI)	Other (EWO)	Total	Number of women	Eligible women's response rate (EWRR)[2]	Overall women's response rate (OWRR)[3]
Residence																							
Urban	84.7	5.6	0.0	1.1	0.1	3.8	4.5	0.1	0.1	100.0	10,567	92.6	93.2	5.4	0.0	0.7	0.0	0.6	0.1	100.0	7,771	93.2	86.3
Rural	90.3	2.1	0.0	0.3	0.3	3.4	3.0	0.6	0.1	100.0	19,308	97.1	94.9	3.8	0.0	0.2	0.0	0.9	0.1	100.0	14,736	94.9	92.2
County																							
Mombasa	83.5	5.7	0.0	1.0	0.0	4.3	5.1	0.4	0.0	100.0	690	92.6	87.9	10.1	0.0	0.9	0.2	0.7	0.2	100.0	544	87.9	81.4
Kwale	87.8	4.5	0.0	0.0	0.0	2.2	4.7	0.8	0.0	100.0	600	95.1	88.2	9.6	0.0	0.8	0.0	1.4	0.0	100.0	638	88.2	83.9
Kilifi	86.1	3.0	0.0	0.0	0.0	5.2	4.8	0.9	0.0	100.0	660	96.6	98.1	0.8	0.0	0.0	0.0	1.1	0.0	100.0	526	98.1	94.8
Tana River	77.0	7.2	0.0	0.0	1.3	9.3	4.8	0.4	0.0	100.0	540	90.0	93.0	6.0	0.0	0.0	0.0	0.9	0.0	100.0	430	93.0	83.8
Lamu	90.2	3.5	0.0	0.2	0.0	2.6	2.8	0.4	0.4	100.0	540	96.1	96.8	2.8	0.0	0.0	0.0	0.5	0.0	100.0	431	96.8	92.9
Taita Taveta	81.7	6.1	0.0	0.7	0.2	3.3	7.2	0.7	0.0	100.0	540	92.1	88.7	9.3	0.0	1.2	0.0	0.9	0.0	100.0	335	88.7	81.6
Garissa	86.8	2.1	0.0	1.1	0.2	7.3	1.7	0.8	0.0	100.0	522	96.2	96.3	2.6	0.0	0.4	0.0	0.7	0.0	100.0	536	96.3	92.6
Wajir	87.5	3.9	0.0	0.0	0.0	3.9	4.4	0.4	0.0	100.0	568	95.8	95.2	4.2	0.0	0.0	0.0	0.6	0.0	100.0	544	95.2	91.2
Mandera	91.2	0.4	0.4	0.2	0.2	4.9	2.6	0.2	0.0	100.0	570	98.9	98.4	0.8	0.0	0.0	0.0	0.8	0.0	100.0	492	98.4	97.3
Marsabit	80.9	1.1	0.0	0.2	5.4	8.5	3.3	0.6	0.0	100.0	540	92.4	89.7	8.8	0.0	0.0	0.0	1.5	0.0	100.0	464	89.7	82.8
Isiolo	92.7	0.6	0.0	0.0	0.0	4.2	2.1	0.0	0.4	100.0	518	99.4	97.7	1.6	0.0	0.0	0.0	0.5	0.2	100.0	441	97.7	97.1
Meru	90.1	1.8	0.0	0.5	0.1	2.3	4.6	0.4	0.1	100.0	780	97.4	95.2	3.4	0.0	0.2	0.0	1.3	0.0	100.0	477	95.2	92.7
Tharaka-Nithi	83.7	7.2	0.0	1.3	0.0	4.3	2.8	0.7	0.0	100.0	540	90.8	88.3	6.6	0.0	3.3	0.0	1.8	0.0	100.0	333	88.3	80.1
Embu	94.2	0.8	0.0	1.2	0.0	2.0	1.3	0.3	0.2	100.0	600	97.9	95.0	2.1	0.0	2.1	0.0	0.8	0.0	100.0	380	95.0	93.0
Kitui	90.5	0.6	0.0	0.0	0.0	3.5	5.2	0.3	0.0	100.0	660	99.3	98.3	1.5	0.0	0.0	0.0	0.2	0.0	100.0	476	98.3	97.7
Machakos	88.7	4.2	0.0	0.6	0.0	3.5	2.7	0.1	0.1	100.0	771	94.9	94.7	4.0	0.4	0.2	0.0	0.4	0.2	100.0	472	94.7	89.8
Makueni	88.2	2.3	0.0	0.2	0.0	2.6	6.7	0.2	0.0	100.0	660	97.3	93.7	5.1	0.0	0.0	0.0	1.2	0.0	100.0	415	93.7	91.2
Nyandarua	71.8	12.0	0.0	2.3	0.3	6.0	6.8	0.7	0.0	100.0	600	83.0	87.0	10.4	0.0	0.3	0.7	1.7	0.0	100.0	299	87.0	72.2
Nyeri	86.4	4.2	0.0	1.2	0.0	2.4	5.6	0.0	0.2	100.0	660	94.1	86.4	9.2	0.0	2.2	0.0	0.9	1.3	100.0	316	86.4	81.3
Kirinyaga	92.4	1.7	0.0	1.0	0.3	2.2	2.4	0.0	0.0	100.0	630	96.8	96.1	2.7	0.0	0.0	0.0	0.9	0.3	100.0	332	96.1	93.0
Murang'a	91.4	3.5	0.0	1.2	0.0	1.3	2.5	0.0	0.1	100.0	690	95.2	95.0	3.1	0.0	0.9	0.0	0.9	0.2	100.0	457	95.0	90.4
Kiambu	96.0	0.6	0.0	0.8	0.0	0.4	2.2	0.0	0.0	100.0	900	98.6	96.3	1.7	0.0	1.1	0.0	0.3	0.6	100.0	656	96.3	95.0
Turkana	79.1	3.2	0.0	0.8	0.0	2.5	13.9	0.5	0.0	100.0	599	95.2	94.8	3.6	0.0	0.0	0.0	1.7	0.0	100.0	363	94.8	90.2
West Pokot	88.4	1.4	0.0	0.4	0.0	1.8	6.4	1.6	0.0	100.0	560	98.0	96.3	2.2	0.0	0.7	0.0	0.9	0.0	100.0	461	96.3	94.4
Samburu	85.3	0.6	0.0	0.2	0.2	5.0	6.1	2.4	0.2	100.0	539	98.9	96.3	2.0	0.0	1.0	0.0	0.2	0.5	100.0	410	96.3	95.3
Trans Nzoia	81.3	0.6	0.0	2.1	0.3	7.0	7.8	1.0	0.0	100.0	630	96.4	97.7	2.0	0.0	0.0	0.0	0.4	0.0	100.0	512	97.7	94.2
Uasin Gishu	95.5	1.2	0.0	0.6	0.0	0.9	1.4	0.5	0.0	100.0	660	98.1	95.5	2.9	0.0	0.4	0.0	1.3	0.0	100.0	550	95.5	93.7
Elgeyo-Marakwet	95.9	0.4	0.0	0.0	0.0	2.2	1.5	0.0	0.0	100.0	540	99.6	94.6	4.7	0.0	0.2	0.0	0.4	0.0	100.0	464	94.6	94.2
Nandi	92.3	1.8	0.0	0.0	0.2	3.5	2.0	0.2	0.0	100.0	600	97.9	96.8	3.0	0.0	0.0	0.0	0.2	0.0	100.0	496	96.8	94.7
Baringo	89.3	0.7	0.0	0.2	0.7	4.8	2.5	1.9	0.0	100.0	568	98.3	90.5	8.0	0.0	0.0	0.0	1.5	0.0	100.0	402	90.5	89.0
Laikipia	86.8	4.0	0.0	0.2	0.0	3.5	4.6	0.9	0.0	100.0	570	95.4	93.0	6.4	0.0	0.0	0.0	0.6	0.0	100.0	345	93.0	88.7
Nakuru	88.2	6.3	0.0	0.5	0.3	2.4	1.9	0.3	0.1	100.0	780	92.6	93.9	4.1	0.2	0.6	0.2	0.9	0.2	100.0	538	93.9	86.9
Narok	97.3	0.5	0.0	0.0	0.0	0.5	0.5	0.8	0.5	100.0	630	99.5	99.4	0.6	0.0	0.0	0.0	0.0	0.0	100.0	473	99.4	98.9
Kajiado	80.2	4.8	0.0	1.1	1.2	7.9	4.5	0.3	0.0	100.0	660	91.8	91.3	6.7	0.0	1.2	0.0	0.5	0.2	100.0	404	91.3	83.9
Kericho	93.5	1.0	0.0	0.8	0.0	2.2	2.5	0.0	0.0	100.0	600	98.1	96.3	2.7	0.0	0.6	0.0	0.4	0.0	100.0	481	96.3	94.4
Bomet	94.2	0.5	0.0	0.7	0.0	3.3	1.0	0.3	0.0	100.0	600	98.8	95.0	4.5	0.0	0.0	0.0	0.5	0.0	100.0	625	95.0	93.9
Kakamega	87.7	6.2	0.0	0.4	0.0	2.7	2.7	0.4	0.0	100.0	780	93.1	94.6	4.5	0.0	0.0	0.0	0.9	0.0	100.0	556	94.6	88.0
Vihiga	90.0	1.8	0.0	0.0	0.0	6.1	2.1	0.0	0.0	100.0	570	98.1	95.7	3.9	0.0	0.0	0.0	0.4	0.0	100.0	463	95.7	93.9
Bungoma	96.8	0.1	0.0	0.0	0.0	0.9	1.9	0.3	0.0	100.0	690	99.9	98.6	0.9	0.0	0.0	0.0	0.5	0.0	100.0	582	98.6	98.5
Busia	86.0	4.2	0.0	1.3	0.8	4.5	3.2	0.0	0.0	100.0	600	93.1	91.7	5.0	0.0	1.5	0.0	1.9	0.0	100.0	481	91.7	85.4
Siaya	92.3	0.3	0.0	0.3	0.0	5.0	1.5	0.6	0.0	100.0	660	99.3	95.0	4.0	0.0	0.4	0.0	0.6	0.0	100.0	500	95.0	94.4
Kisumu	91.2	5.5	0.0	0.0	0.0	1.7	1.5	0.2	0.0	100.0	660	94.4	95.7	3.2	0.0	0.2	0.2	0.7	0.0	100.0	562	95.7	90.3
Homa Bay	98.3	0.0	0.0	0.0	0.0	1.7	0.0	0.0	0.0	100.0	660	100.0	98.4	1.0	0.0	0.0	0.0	0.5	0.0	100.0	579	98.4	98.4
Migori	92.1	0.8	0.0	0.2	0.0	4.4	2.1	0.5	0.0	100.0	630	99.0	96.4	3.3	0.0	0.0	0.0	0.3	0.0	100.0	608	96.4	95.4
Kisii	87.4	2.8	0.0	0.0	0.0	5.1	4.2	0.6	0.0	100.0	690	96.9	93.4	5.5	0.0	0.0	0.0	1.1	0.0	100.0	473	93.4	90.6
Nyamira	91.0	0.7	0.0	0.2	0.0	5.0	3.2	0.0	0.0	100.0	600	99.1	95.0	2.5	0.0	0.0	0.0	2.1	0.4	100.0	478	95.0	94.1
Nairobi City	73.1	19.5	0.0	2.0	0.0	1.2	3.4	0.4	0.4	100.0	1,020	77.3	85.1	13.2	0.0	0.8	0.1	0.3	0.4	100.0	707	85.1	65.8
Total	88.3	3.3	0.0	0.6	0.2	3.5	3.6	0.4	0.1	100.0	29,875	95.5	94.3	4.3	0.0	0.4	0.0	0.8	0.1	100.0	22,507	94.3	90.1

[1] Using the number of households falling into specific response categories, the household response rate (HRR) is calculated as:

$$\frac{100 * C}{C + HP + P + R + DNF}$$

[2] The eligible women's response rate (EWRR) is equivalent to the percentage of interviews completed (EWC).
[3] The overall women's response rate (OWRR) is calculated as:

$$OWRR = HRR * EWRR/100$$

ESTIMATES OF SAMPLING ERRORS

Appendix B

The estimates from a sample survey are affected by two types of errors: (1) non-sampling errors, and (2) sampling errors. Non-sampling errors are the results of mistakes made in implementing data collection and data processing, such as failure to locate and interview the correct household, misunderstanding of the questions on the part of either the interviewer or the respondent, and data entry errors. Although numerous efforts were made during the implementation of the 2025–26 Kenya Mini DHS Survey (2025–26 KMDHS) to minimise this type of error, non-sampling errors are impossible to avoid and difficult to evaluate statistically.

Sampling errors, on the other hand, can be evaluated statistically. The sample of respondents selected in the 2025–26 KMDHS is only one of many samples that could have been selected from the same population, using the same design and identical size. Each of these samples would yield results that differ somewhat from the results of the actual sample selected. Sampling errors are a measure of the variability between all possible samples. Although the degree of variability is not known exactly, it can be estimated from the survey results.

A sampling error is usually measured in terms of the standard error for a particular statistic (mean, percentage, etc.), which is the square root of the variance. The standard error can be used to calculate confidence intervals within which the true value for the population can reasonably be assumed to fall. For example, for any given statistic calculated from a sample survey, the value of that statistic will fall within a range of plus or minus two times the standard error of that statistic in 95 percent of all possible samples of identical size and design.

If the sample of respondents had been selected as a simple random sample, it would have been possible to use straightforward formulas for calculating sampling errors. However, the 2025–26 KMDHS sample is the result of a multi-stage stratified design, and, consequently, it was necessary to use more complex formulas. The computer software used to calculate sampling errors for the 2025–26 KMDHS is a SAS program. This program used the Taylor linearisation method for variance estimation for survey estimates that are means, proportions or ratios. The Jackknife repeated replication method is used for variance estimation of more complex statistics such as fertility and mortality rates.

The Taylor linearisation method treats any percentage or average as a ratio estimate, $r = y/x$, where y represents the total sample value for variable y, and x represents the total number of cases in the group or subgroup under consideration. The variance of r is computed using the formula given below, with the standard error being the square root of the variance:

$$SE^2(r) = var(r) = \frac{1}{x^2}\sum_{h=1}^{H}\left[(1-f_h)\frac{m_h}{m_h-1}\left(\sum_{i=1}^{m_h} z_{hi}^2 - \frac{z_h^2}{m_h}\right)\right]$$

in which

$$z_{hi} = y_{hi} - rx_{hi}\text{, and } z_h = y_h - rx_h$$

where h represents the stratum which varies from 1 to H,
m_h is the total number of clusters selected in the hth stratum,
y_{hi} is the sum of the weighted values of variable y in the ith cluster in the hth stratum,
x_{hi} is the sum of the weighted number of cases in the ith cluster in the hth stratum, and
f_h is the sampling fraction of PSU in the hth stratum

The Jackknife repeated replication method derives estimates of complex rates from each of several replications of the parent sample and calculates standard errors for these estimates using simple formulae. Each replication considers all but one cluster in the calculation of the estimates. Pseudo-independent replications are thus created. In the 2025–26 KMDHS, there were 998 non-empty clusters. Hence, 998 replications were created. The variance of a ratio r is calculated as follows:

$$SE^2(r) = var(r) = \frac{1}{k(k-1)} \sum_{i=1}^{k} (r_i - r)^2$$

in which

$$r_i = kr - (k-1)r_{(i)}$$

where r is the estimate computed from the full sample of 998 clusters,
$r_{(i)}$ is the estimate computed from the reduced sample of 997 clusters (i^{th} cluster excluded)
k is the total number of clusters.

In addition to the standard error, the programme computes the design effect (DEFT) for each estimate, which is defined as the ratio between the standard error using the given sample design and the standard error that would result if a simple random sample had been used. A DEFT value of 1.0 indicates that the sample design is as efficient as a simple random sample, while a value greater than 1.0 indicates the increase in the sampling error due to the use of a more complex and less statistically efficient design, such as multistage and cluster sampling. The programme also computes the relative standard error and the confidence limits for the estimates.

Sampling errors for the 2025–26 KMDHS are calculated for selected variables considered to be of primary interest for the survey. The results are presented in this appendix for the country, for the urban and rural areas separately, and for each of the 47 counties. For each variable, the type of statistic (mean, proportion, ratio or rate) and the base population are given in **Table B.1**. **Tables B.2** to **B.53** present the value of the statistic (R), its standard error (SE), the number of unweighted (N-UNWE) and weighted (N-WEIG) cases, the design effect (DEFT), the relative standard error (SE/R), and the 95 percent confidence limits (R±2SE), for each variable. The DEFT is considered undefined when the standard error considering simple random sample is zero (when the estimate is close to 0 or 1).

The confidence interval (e.g., as calculated for *Mean number of children ever born to women aged 15–49*) can be interpreted as follows: the sample estimate from the national sample is 2.112 (R) and its standard error is 0.024 (SE). Therefore, to obtain the 95 percent confidence limits, one adds and subtracts twice the standard error to the sample estimate, i.e., R±2×SE. There is a high probability (95 percent) that the true mean number of children ever born to women aged 15–49 is between 2.063 and 2.160.

For the total sample, the value of the design effect (DEFT), averaged over all variables for the women survey, is 1.775 which means that, due to multistage and clustering of the sample, the average standard error is increased by a factor of 1.775 over that in an equivalent simple random sample.

Table B.1 List of selected variables for sampling errors, Kenya Mini DHS 2025–26

Variable	Estimate	Base population
HOUSEHOLDS AND POPULATION		
Electricity primary source of lighting	Proportion	De jure household population
Primary reliance on clean fuels and technology	Proportion	De jure household population
Births registered with civil authority	Proportion	De jure household population under 5
Ownership of at least one ITN	Proportion	Households
Ownership of at least one ITN for every two persons	Proportion	Households (with at least one person who stayed in the household the night before the survey)
Improved drinking water source	Proportion	De jure household population
At least basic drinking water service	Proportion	De jure household population
Improved sanitation facility	Proportion	De jure household population
At least basic sanitation service	Proportion	De jure household population
Using open defecation	Proportion	De jure household population
Using a handwashing facility with soap and water	Proportion	De jure household population for whom handwashing place was observed or with no on-site place for handwashing
WOMEN		
Urban residence	Proportion	Women 15–49
No education	Proportion	Women 15–49
Secondary education or higher	Proportion	Women 15–49
Literacy	Proportion	Women 15–49
Use of the Internet in last 12 months	Proportion	Women 15–49
Currently married/in union	Proportion	Women 15–49
Had sexual intercourse before age 18	Proportion	Women 20–49
Age specific fertility rate 15–19	Rate	Women-years of exposure to childbearing at age 15–19 in the 3 years preceding the survey
Total fertility rate (3 years)	Rate	Women-years of exposure to childbearing
Currently pregnant	Proportion	Women 15–49
Mean number of children ever born to women age 40–49	Mean	Women 40–49
Mean number of children ever born to women age 15–49	Mean	Women 15–49
Mean number of living children to women age 15–49	Mean	Women 15–49
Median birth interval	Median	Non-first births in the 5 years preceding the survey
First birth before age 18	Proportion	Women 20–49
Want to delay next birth at least 2 years	Proportion	Currently married women 15–49
Want no more children	Proportion	Currently married women 15–49
Ideal number of children	Mean	Women 15–49 with numeric responses
Total wanted fertility rate (3 years)	Rate	Women-years of exposure to childbearing
Currently using any contraceptive method	Proportion	Currently married women 15–49
Currently using any modern method	Proportion	Currently married women 15–49
Currently using pill	Proportion	Currently married women 15–49
Currently using injectables	Proportion	Currently married women 15–49
Currently using implants	Proportion	Currently married women 15–49
Currently using male condom	Proportion	Currently married women 15–49
Currently using any traditional method	Proportion	Currently married women 15–49
12 month discontinuation rate due to method failure	Rate	Women 15–49
12 month discontinuation rate due to any reason	Rate	Women 15–49
12 month discontinuation rate due to switching to another method	Rate	Women 15–49
Unmet need for spacing	Proportion	Currently married women 15–49
Unmet need for limiting	Proportion	Currently married women 15–49
Unmet need total	Proportion	Currently married women 15–49
Demand satisfied by modern methods (married women)	Proportion	Currently married women 15–49
Demand satisfied by modern methods (all women)	Proportion	Women 15–49
Not exposed to any of the ten media sources	Proportion	Women 15–49
Neonatal mortality rate[1]	Rate	Children exposed to the risk of mortality
Post-neonatal mortality rate[1]	Rate	Children exposed to the risk of mortality
Infant mortality rate[1]	Rate	Children exposed to the risk of mortality
Child mortality rate[1]	Rate	Children exposed to the risk of mortality
Under-5 mortality rate[1]	Rate	Children exposed to the risk of mortality
Perinatal mortality rate	Rate	Pregnancies of 28 or more weeks' duration to women 15–49 in the 5 years preceding the survey
Stillbirth rate	Rate	Pregnancies of 28 or more weeks' duration to women 15–49 in the 5 years preceding the survey
Early neonatal rate	Rate	Pregnancies of 28 or more weeks' duration to women 15–49 in the 5 years preceding the survey
In any avoidable high-risk category	Proportion	Children born in the 5 years preceding the survey to women 15–49
Received ANC from a skilled provider	Proportion	Women 15–49 who had a live birth in the 2 years preceding the survey
4+ ANC visits	Proportion	Women 15–49 who had a live birth in the 2 years preceding the survey
8+ ANC visits	Proportion	Women 15–49 who had a live birth in the 2 years preceding the survey
Took any iron-containing supplements	Proportion	Women 15–49 who had a live birth in the 2 years preceding the survey
Mothers protected against tetanus for last birth	Proportion	Women 15–49 with a live birth in the 2 years preceding the survey
Delivered in a health facility (live births)	Proportion	Live births in the 2 years preceding the survey
Delivered by C-section (live births)	Proportion	Live births in the 2 years preceding the survey
Delivered by a skilled provider (live births)	Proportion	Live births in the 2 years preceding the survey
Women with postnatal check during first 2 days	Proportion	Women 15–49 with a live birth in the 2 years preceding the survey

Continued…

Table B.1—*Continued*

Variable	Estimate	Base population
Newborns with postnatal check during first 2 days	Proportion	Most recent live births in the 2 years preceding the survey
Any problem accessing health care	Proportion	Women 15–49
Ever had a vaccination card	Proportion	Children 12–23 months
Received BCG vaccination	Proportion	Children 12–23 months
Received DPT-HepB-Hib vaccination (3 doses)	Proportion	Children 12–23 months
Received pneumococcal vaccination (3 doses)	Proportion	Children 12–23 months
Received MR 1 vaccination	Proportion	Children 12–23 months
Fully vaccinated according to national schedule (12–23 months)	Proportion	Children 12–23 months
Received MR 2 vaccination	Proportion	Children 24–35 months
Fully vaccinated according to national schedule (24–35 months)	Proportion	Children 24–35 months
Received HPV vaccination (2 doses)	Proportion	Women 15–19
Sought treatment for diarrhoea	Proportion	Children under 5 with diarrhoea in last 2 weeks
Treated with ORS	Proportion	Children under 5 with diarrhoea in last 2 weeks
Exclusive breastfeeding	Proportion	Youngest children 0–5 months living with their mother
Minimum dietary diversity (children 6–23 months)	Proportion	Youngest children 6–23 months living with their mother
Child slept under an ITN last night	Proportion	Children under 5 in households
Pregnant women slept under an ITN last night	Proportion	Pregnant women 15–49
Received 3+ doses of SP/Fansidar	Proportion	Women 15–49 with a live birth in the 2 years preceding the survey
Child has fever in last 2 weeks	Proportion	Child under 5
Child had blood taken from finger/heel	Proportion	Child under 5 who had fever in the last 2 weeks
Child took ACT	Proportion	Child under 5 with fever in last 2 weeks who received any antimalarial medications
Condom use at last sex (women 12–24)	Proportion	Women 15–24 with non-marital, non-cohabiting partner in last 12 months
Mobile phone ownership	Proportion	Women 15–49

[1] The mortality rates are calculated for 5 years before the survey for the national sample, urban, and rural samples and for the 10 years before the survey for regional samples.

Table B.2 Sampling errors: Total sample, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases: Unweighted (N)	Number of cases: Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
HOUSEHOLDS AND POPULATION								
Electricity primary source of lighting	0.655	0.009	101,320	92,946	2.445	0.013	0.638	0.673
Primary reliance on clean fuels and technology	0.273	0.011	101,320	92,946	3.655	0.040	0.251	0.296
Births registered with civil authority	0.774	0.009	12,105	10,830	1.913	0.012	0.756	0.792
Ownership of at least one ITN	0.406	0.007	26,380	26,380	2.193	0.016	0.393	0.419
Ownership of at least one ITN for every two persons	0.251	0.005	26,207	26,153	1.856	0.020	0.241	0.261
Improved drinking water source	0.813	0.008	101,320	92,946	2.688	0.010	0.797	0.829
At least basic drinking water service	0.734	0.009	101,320	92,946	2.541	0.012	0.716	0.751
Improved sanitation facility	0.723	0.008	101,320	92,946	2.420	0.011	0.706	0.739
At least basic sanitation service	0.448	0.014	101,320	92,946	3.856	0.031	0.420	0.476
Using open defecation	0.069	0.005	101,320	92,946	2.527	0.076	0.059	0.080
Using a handwashing facility with soap and water	0.532	0.009	99,734	90,983	2.421	0.017	0.514	0.550
WOMEN								
Urban residence	0.463	0.012	21,231	21,231	3.414	0.025	0.440	0.486
No education	0.055	0.005	21,066	21,062	3.333	0.095	0.045	0.066
Secondary education or higher	0.657	0.009	21,066	21,062	2.676	0.013	0.640	0.675
Literacy	0.917	0.006	21,231	21,231	3.093	0.006	0.905	0.928
Use of the internet in last 12 months	0.606	0.009	21,231	21,231	2.659	0.015	0.588	0.624
Currently married/in union	0.550	0.006	21,231	21,231	1.835	0.011	0.537	0.562
Had sexual intercourse before age 18	0.479	0.008	16,831	17,315	1.976	0.016	0.464	0.495
Age specific fertility rate 15–19 (3 years)	63.321	3.299	12,323	11,504	1.475	0.052	56.722	69.920
Total fertility rate (3 years)	3.108	0.068	59,397	60,059	1.727	0.022	2.972	3.245
Currently pregnant	0.047	0.002	21,231	21,231	1.549	0.048	0.043	0.052
Mean number of children ever born to women age 40–49	4.212	0.057	3,956	3,833	1.627	0.013	4.099	4.326
Mean number of children ever born to women age 15–49	2.112	0.024	21,231	21,231	1.695	0.011	2.063	2.160
Mean number of living children to women age 15–49	2.013	0.022	21,231	21,231	1.647	0.011	1.968	2.058
Median birth interval	46.324	0.745	8,322	7,349	1.896	0.016	44.834	47.814
First birth before age 18	0.212	0.005	16,831	17,315	1.560	0.023	0.202	0.221
Want to delay next birth at least 2 years	0.310	0.009	11,686	11,672	1.989	0.027	0.293	0.327
Want no more children	0.441	0.008	11,686	11,672	1.706	0.018	0.425	0.457
Ideal number of children	3.737	0.043	21,135	21,109	3.091	0.011	3.651	3.823
Total wanted fertility rate (3 years)	2.687	0.063	59,397	60,059	1.734	0.024	2.560	2.814
Currently using any contraceptive method	0.607	0.010	11,686	11,672	2.270	0.017	0.586	0.627
Currently using any modern method	0.561	0.010	11,686	11,672	2.099	0.017	0.542	0.581
Currently using pill	0.071	0.006	11,686	11,672	2.482	0.083	0.059	0.082
Currently using injectables	0.187	0.006	11,686	11,672	1.718	0.033	0.175	0.200
Currently using implants	0.228	0.007	11,686	11,672	1.693	0.029	0.215	0.241
Currently using male condoms	0.014	0.002	11,686	11,672	1.610	0.126	0.010	0.017
Currently using any traditional method	0.045	0.003	11,686	11,672	1.621	0.069	0.039	0.051
12 month discontinuation rate due to method failure	2.070	0.699	12,639	14,022	1.453	0.338	0.672	3.467
12 month discontinuation rate due to any reason	27.578	0.176	12,639	14,022	1.832	0.006	27.225	27.930
12 month discontinuation rate due to switching to another method	5.779	0.341	12,639	14,022	1.548	0.059	5.097	6.460
Unmet need for spacing	0.074	0.004	11,686	11,672	1.650	0.054	0.066	0.082
Unmet need for limiting	0.081	0.004	11,686	11,672	1.616	0.050	0.073	0.090
Unmet need total	0.155	0.006	11,686	11,672	1.865	0.040	0.143	0.168
Demand satisfied by modern methods (married women)	0.737	0.008	8,391	8,892	1.811	0.011	0.720	0.754
Demand satisfied by modern methods (all women)	0.732	0.007	10,897	11,581	1.708	0.010	0.718	0.746
Not exposed to any of the ten media sources	0.081	0.005	21,231	21,231	2.692	0.062	0.070	0.091
Neonatal mortality (last 0–4 years)	19.126	2.094	11,371	10,515	1.487	0.109	14.939	23.314
Post-neonatal mortality (last 0–4 years)	10.832	1.399	11,353	10,521	1.394	0.129	8.034	13.630
Infant mortality (last 0–4 years)	29.959	2.722	11,378	10,522	1.594	0.091	24.516	35.402
Child mortality (last 0–4 years)	7.579	1.226	11,475	10,590	1.486	0.162	5.126	10.031
Under-five mortality (last 0–4 years)	37.310	2.926	11,417	10,557	1.548	0.078	31.459	43.162
Stillbirth rate	13.756	2.808	11,451	10,623	2.267	0.204	8.140	19.373
Early neonatal rate	16.616	2.051	11,339	10,519	1.538	0.123	12.515	20.718
Perinatal mortality rate	30.210	3.405	11,451	10,623	1.855	0.113	23.400	37.019
In any avoidable high-risk category	0.395	0.009	11,339	10,519	1.688	0.022	0.377	0.412
Received ANC from a skilled provider	0.979	0.003	4,289	3,983	1.515	0.003	0.973	0.986
4+ ANC visits	0.724	0.010	4,289	3,983	1.413	0.013	0.705	0.743
8+ ANC visits	0.095	0.008	4,289	3,983	1.859	0.088	0.078	0.111
Took any iron-containing supplements	0.889	0.008	4,289	3,983	1.622	0.009	0.873	0.904
Mothers protected against tetanus for last birth	0.785	0.009	4,289	3,983	1.401	0.011	0.767	0.803
Delivered in a health facility (live births)	0.911	0.007	4,446	4,115	1.465	0.007	0.898	0.925
Delivered by a skilled provider (live births)	0.919	0.006	4,446	4,115	1.463	0.007	0.906	0.932
Delivered by C-section (live births)	0.203	0.011	4,446	4,115	1.737	0.056	0.180	0.225
Women with postnatal check during first 2 days	0.846	0.008	4,289	3,983	1.459	0.010	0.830	0.862
Newborns with postnatal check during first 2 days	0.855	0.008	4,289	3,983	1.577	0.010	0.838	0.872
Any problem accessing health care	0.567	0.009	21,231	21,231	2.669	0.016	0.549	0.586
Ever had a vaccination card	0.982	0.004	2,113	1,934	1.480	0.005	0.973	0.991

Continued…

Table B.2—*Continued*

Variable	Value (R)	Standard error (SE)	Number of cases: Unweighted (N)	Number of cases: Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
Received BCG vaccination	0.976	0.004	2,113	1,934	1.227	0.004	0.967	0.985
Received DPT-HepB+Hib vaccination (3 doses)	0.899	0.009	2,113	1,934	1.281	0.010	0.881	0.916
Received pneumococcal vaccination (3 doses)	0.924	0.007	2,113	1,934	1.196	0.008	0.909	0.938
Received measles vaccination	0.887	0.009	2,113	1,934	1.305	0.011	0.868	0.906
Fully vaccinated according to national schedule (12–23 months)	0.456	0.016	2,113	1,934	1.379	0.035	0.425	0.488
Received measles/measles-rubella 2 vaccination (24–35 months)	0.720	0.016	2,100	2,007	1.552	0.022	0.688	0.751
Fully vaccinated according to national schedule (24–35 months)	0.383	0.019	2,100	2,007	1.733	0.050	0.345	0.421
Received HPV vaccination (2 doses) women 15–19	0.394	0.013	4,400	3,916	1.791	0.033	0.368	0.421
Sought treatment for diarrhoea	0.578	0.023	1,033	1,146	1.538	0.040	0.532	0.623
Treated with ORS	0.414	0.023	1,033	1,146	1.548	0.055	0.368	0.460
Exclusive breastfeeding	0.715	0.021	1,072	1,016	1.507	0.029	0.673	0.757
Minimum dietary diversity (children 6–23 months)	0.427	0.013	3,056	2,802	1.443	0.030	0.401	0.453
Child slept under an ITN last night	0.417	0.009	12,151	10,881	1.620	0.021	0.399	0.435
Pregnant women slept under an ITN last night	0.392	0.022	1,126	988	1.413	0.056	0.348	0.436
Received 3+ doses of SP/Fansidar	0.139	0.007	4,289	3,983	1.377	0.052	0.124	0.153
Child has fever in last 2 weeks	0.138	0.006	10,967	10,166	1.759	0.045	0.126	0.151
Child had blood taken from finger/heel	0.295	0.021	1,390	1,403	1.656	0.070	0.253	0.336
Child took ACT	0.741	0.040	258	245	1.405	0.054	0.662	0.821
Condom use at last sex (women 15–24)	0.355	0.017	1,525	1,639	1.394	0.048	0.320	0.389
Mobile phone ownership	0.815	0.005	21,231	21,231	1.938	0.006	0.805	0.825

Table B.3 Sampling errors: Urban sample, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases: Unweighted (N)	Number of cases: Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
HOUSEHOLDS AND POPULATION								
Electricity primary source of lighting	0.935	0.007	29,104	36,354	2.194	0.008	0.920	0.950
Primary reliance on clean fuels and technology	0.611	0.020	29,104	36,354	3.157	0.033	0.570	0.651
Births registered with civil authority	0.826	0.017	3,484	4,431	2.303	0.021	0.792	0.860
Ownership of at least one ITN	0.284	0.010	8,950	11,975	2.141	0.036	0.264	0.304
Ownership of at least one ITN for every two persons	0.178	0.007	8,856	11,813	1.761	0.040	0.164	0.193
Improved drinking water source	0.980	0.003	29,104	36,354	1.751	0.003	0.973	0.987
At least basic drinking water service	0.952	0.005	29,104	36,354	1.759	0.006	0.941	0.963
Improved sanitation facility	0.927	0.009	29,104	36,354	2.616	0.010	0.909	0.945
At least basic sanitation service	0.508	0.032	29,104	36,354	5.006	0.062	0.445	0.572
Using open defecation	0.004	0.001	29,104	36,354	0.755	0.180	0.002	0.005
Using a handwashing facility with soap and water	0.595	0.019	28,460	35,219	2.931	0.031	0.558	0.632
WOMEN								
No education	0.039	0.011	7,213	9,779	4.616	0.269	0.018	0.060
Secondary education or higher	0.747	0.018	7,213	9,779	3.460	0.024	0.712	0.782
Literacy	0.942	0.012	7,241	9,829	4.414	0.013	0.918	0.967
Use of the internet in last 12 months	0.808	0.011	7,241	9,829	2.301	0.013	0.787	0.830
Age specific fertility rate 15–19 (3 years)	53.884	6.378	3,914	4,771	1.717	0.118	41.127	66.640
Total fertility rate (3 years)	2.639	0.106	20,658	28,392	1.899	0.040	2.427	2.851
Currently pregnant	0.047	0.004	7,241	9,829	1.678	0.089	0.039	0.055
Mean number of children ever born to women age 40–49	3.502	0.109	1,123	1,473	1.891	0.031	3.284	3.720
Median birth interval	49.804	1.783	2,342	3,011	2.115	0.036	46.238	53.369
Want no more children	0.361	0.014	3,650	5,174	1.700	0.037	0.334	0.388
Ideal number of children	3.516	0.091	7,204	9,747	3.909	0.026	3.334	3.698
Total wanted fertility rate (3 years)	2.412	0.105	20,658	28,392	1.908	0.043	2.202	2.621
Currently using any contraceptive method	0.604	0.021	3,650	5,174	2.615	0.035	0.561	0.646
Currently using any modern method	0.547	0.020	3,650	5,174	2.381	0.036	0.507	0.586
Currently using pill	0.100	0.012	3,650	5,174	2.492	0.124	0.076	0.125
Currently using injectables	0.156	0.011	3,650	5,174	1.902	0.073	0.133	0.178
Currently using implants	0.206	0.013	3,650	5,174	1.936	0.063	0.181	0.232
Currently using male condoms	0.018	0.004	3,650	5,174	1.596	0.195	0.011	0.025
Currently using any traditional method	0.057	0.006	3,650	5,174	1.635	0.110	0.044	0.070
Unmet need for spacing	0.077	0.008	3,650	5,174	1.787	0.102	0.062	0.093
Unmet need for limiting	0.074	0.008	3,650	5,174	1.839	0.108	0.058	0.089
Unmet need total	0.151	0.013	3,650	5,174	2.165	0.085	0.125	0.177
Demand satisfied by modern methods (married women)	0.724	0.017	2,647	3,904	2.043	0.024	0.690	0.759
Demand satisfied by modern methods (all women)	0.723	0.014	3,608	5,244	1.932	0.019	0.695	0.751
Not exposed to any of the ten media sources	0.079	0.010	7,241	9,829	3.245	0.130	0.058	0.099
Neonatal mortality (last 0–4 years)	17.322	4.150	3,455	4,578	1.691	0.240	9.022	25.621
Post-neonatal mortality (last 0–4 years)	11.153	2.754	3,475	4,604	1.548	0.247	5.645	16.662
Infant mortality (last 0–4 years)	28.475	5.553	3,458	4,581	1.860	0.195	17.370	39.580
Child mortality (last 0–4 years)	8.652	2.484	3,467	4,581	1.593	0.287	3.683	13.621
Under-five mortality (last 0–4 years)	36.880	5.941	3,463	4,590	1.757	0.161	24.999	48.762
Stillbirth rate	17.794	6.076	3,481	4,661	2.364	0.341	5.642	29.946
Early neonatal rate	15.266	4.072	3,445	4,601	1.729	0.267	7.122	23.409
Perinatal mortality rate	32.865	7.122	3,481	4,661	1.996	0.217	18.621	47.110
Received ANC from a skilled provider	0.986	0.006	1,270	1,698	1.667	0.006	0.975	0.997
4+ ANC visits	0.801	0.017	1,270	1,698	1.476	0.021	0.768	0.834
8+ ANC visits	0.127	0.017	1,270	1,698	1.842	0.135	0.093	0.162
Took any iron-containing supplements	0.908	0.015	1,270	1,698	1.864	0.017	0.878	0.938
Mothers protected against tetanus for last birth	0.820	0.015	1,270	1,698	1.417	0.019	0.789	0.850
Delivered in a health facility (live births)	0.987	0.003	1,315	1,755	0.902	0.003	0.981	0.992
Delivered by a skilled provider (live births)	0.989	0.002	1,315	1,755	0.844	0.002	0.984	0.994
Delivered by C-section (live births)	0.291	0.023	1,315	1,755	1.741	0.080	0.244	0.337
Women with postnatal check during first 2 days	0.896	0.013	1,270	1,698	1.476	0.014	0.871	0.922
Newborns with postnatal check during first 2 days	0.886	0.015	1,270	1,698	1.722	0.017	0.855	0.917
Any problem accessing health care	0.469	0.016	7,241	9,829	2.702	0.034	0.437	0.500
Ever had a vaccination card	0.983	0.010	613	784	1.773	0.010	0.963	1.000
Received BCG vaccination	0.985	0.008	613	784	1.514	0.008	0.969	1.000
Received DPT-HepB+Hib vaccination (3 doses)	0.916	0.016	613	784	1.374	0.018	0.884	0.948
Received pneumococcal vaccination (3 doses)	0.942	0.013	613	784	1.306	0.014	0.916	0.968
Received measles vaccination	0.918	0.017	613	784	1.451	0.018	0.885	0.952
Fully vaccinated according to national schedule (12–23 months)	0.466	0.031	613	784	1.467	0.066	0.404	0.527
Received measles/measles-rubella 2 vaccination (24–35 months)	0.754	0.030	649	906	1.733	0.039	0.695	0.813
Fully vaccinated according to national schedule (24–35 months)	0.385	0.037	649	906	1.918	0.096	0.311	0.460
Received HPV vaccination (2 doses) women 15–19	0.360	0.030	1,224	1,398	2.213	0.084	0.299	0.421
Sought treatment for diarrhoea	0.590	0.041	335	554	1.633	0.070	0.507	0.672

Continued…

Table B.3—*Continued*

Variable	Value (R)	Standard error (SE)	Number of cases: Unweighted (N)	Number of cases: Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
Treated with ORS	0.411	0.041	335	554	1.662	0.100	0.328	0.493
Exclusive breastfeeding	0.758	0.040	316	458	1.636	0.052	0.678	0.837
Minimum dietary diversity (children 6–23 months)	0.488	0.025	893	1,156	1.482	0.051	0.439	0.538
Child slept under an ITN last night	0.303	0.015	3,473	4,400	1.616	0.049	0.273	0.333
Pregnant women slept under an ITN last night	0.283	0.037	381	443	1.488	0.131	0.209	0.358
Received 3+ doses of SP/Fansidar	0.104	0.012	1,270	1,698	1.408	0.116	0.079	0.128
Child has fever in last 2 weeks	0.146	0.013	3,326	4,444	1.999	0.087	0.120	0.171
Child had blood taken from finger/heel	0.279	0.039	445	649	1.861	0.140	0.200	0.357
Child took ACT	0.683	0.121	51	65	1.746	0.178	0.440	0.926
Condom use at last sex (women 15–24)	0.331	0.030	540	778	1.471	0.090	0.271	0.391
Mobile phone ownership	0.911	0.006	7,241	9,829	1.877	0.007	0.898	0.923

Table B.4 Sampling errors: Rural sample, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases: Unweighted (N)	Number of cases: Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
HOUSEHOLDS AND POPULATION								
Electricity primary source of lighting	0.475	0.011	72,216	56,592	2.434	0.022	0.454	0.497
Primary reliance on clean fuels and technology	0.057	0.004	72,216	56,592	2.368	0.070	0.049	0.065
Births registered with civil authority	0.738	0.010	8,621	6,400	1.603	0.013	0.719	0.757
Ownership of at least one ITN	0.508	0.007	17,430	14,405	1.735	0.013	0.495	0.521
Ownership of at least one ITN for every two persons	0.310	0.006	17,351	14,340	1.652	0.019	0.299	0.322
Improved drinking water source	0.706	0.012	72,216	56,592	2.965	0.017	0.682	0.730
At least basic drinking water service	0.593	0.012	72,216	56,592	2.768	0.020	0.569	0.617
Improved sanitation facility	0.591	0.010	72,216	56,592	2.382	0.018	0.570	0.612
At least basic sanitation service	0.409	0.009	72,216	56,592	2.068	0.022	0.391	0.427
Using open defecation	0.111	0.008	72,216	56,592	2.773	0.075	0.094	0.128
Using a handwashing facility with soap and water	0.492	0.008	71,274	55,763	1.790	0.016	0.476	0.508
WOMEN								
No education	0.069	0.004	13,853	11,283	1.900	0.059	0.061	0.077
Secondary education or higher	0.580	0.007	13,853	11,283	1.598	0.012	0.566	0.593
Literacy	0.894	0.004	13,990	11,402	1.723	0.005	0.885	0.903
Use of the internet in last 12 months	0.432	0.008	13,990	11,402	1.920	0.019	0.416	0.448
Age specific fertility rate 15–19 (3 years)	70.008	3.230	8,409	6,733	1.164	0.046	63.547	76.469
Total fertility rate (3 years)	3.593	0.065	38,739	31,666	1.303	0.018	3.462	3.724
Currently pregnant	0.048	0.002	13,990	11,402	1.207	0.046	0.044	0.052
Mean number of children ever born to women age 40–49	4.656	0.057	2,833	2,360	1.354	0.012	4.543	4.769
Median birth interval	43.402	0.726	5,980	4,338	1.544	0.017	41.950	44.855
Want no more children	0.504	0.007	8,036	6,498	1.268	0.014	0.490	0.518
Ideal number of children	3.926	0.029	13,931	11,362	1.657	0.007	3.869	3.983
Total wanted fertility rate (3 years)	2.998	0.064	38,739	31,666	1.313	0.021	2.870	3.125
Currently using any contraceptive method	0.609	0.007	8,036	6,498	1.359	0.012	0.594	0.624
Currently using any modern method	0.573	0.007	8,036	6,498	1.344	0.013	0.558	0.588
Currently using pill	0.047	0.003	8,036	6,498	1.209	0.061	0.041	0.053
Currently using injectables	0.213	0.006	8,036	6,498	1.323	0.028	0.201	0.225
Currently using implants	0.245	0.006	8,036	6,498	1.190	0.023	0.234	0.257
Currently using male condoms	0.010	0.001	8,036	6,498	1.214	0.133	0.007	0.013
Currently using any traditional method	0.036	0.003	8,036	6,498	1.222	0.071	0.031	0.041
Unmet need for spacing	0.071	0.003	8,036	6,498	1.180	0.048	0.064	0.078
Unmet need for limiting	0.088	0.004	8,036	6,498	1.160	0.042	0.080	0.095
Unmet need total	0.159	0.005	8,036	6,498	1.140	0.029	0.149	0.168
Demand satisfied by modern methods (married women)	0.746	0.007	5,744	4,988	1.181	0.009	0.733	0.760
Demand satisfied by modern methods (all women)	0.739	0.006	7,289	6,337	1.165	0.008	0.728	0.751
Not exposed to any of the ten media sources	0.082	0.003	13,990	11,402	1.336	0.038	0.076	0.088
Neonatal mortality (last 0–4 years)	20.525	1.880	7,916	5,936	1.086	0.092	16.766	24.284
Post-neonatal mortality (last 0–4 years)	10.597	1.317	7,878	5,916	1.089	0.124	7.962	13.232
Infant mortality (last 0–4 years)	31.121	2.269	7,920	5,941	1.078	0.073	26.583	35.660
Child mortality (last 0–4 years)	6.679	1.050	8,008	6,009	1.090	0.157	4.580	8.778
Under-five mortality (last 0–4 years)	37.592	2.482	7,954	5,966	1.074	0.066	32.629	42.556
Stillbirth rate	10.600	1.322	7,970	5,963	1.068	0.125	7.957	13.243
Early neonatal rate	17.666	1.800	7,894	5,918	1.124	0.102	14.065	21.267
Perinatal mortality rate	28.133	2.321	7,970	5,963	1.155	0.082	23.492	32.775
Received ANC from a skilled provider	0.974	0.004	3,019	2,285	1.370	0.004	0.966	0.982
4+ ANC visits	0.667	0.011	3,019	2,285	1.297	0.017	0.645	0.689
8+ ANC visits	0.071	0.006	3,019	2,285	1.264	0.083	0.059	0.082
Took any iron-containing supplements	0.874	0.008	3,019	2,285	1.258	0.009	0.859	0.889
Mothers protected against tetanus for last birth	0.759	0.010	3,019	2,285	1.265	0.013	0.740	0.779
Delivered in a health facility (live births)	0.855	0.010	3,131	2,360	1.555	0.012	0.834	0.876
Delivered by a skilled provider (live births)	0.866	0.010	3,131	2,360	1.554	0.012	0.846	0.887
Delivered by C-section (live births)	0.137	0.008	3,131	2,360	1.148	0.055	0.122	0.152
Women with postnatal check during first 2 days	0.809	0.010	3,019	2,285	1.432	0.013	0.788	0.829
Newborns with postnatal check during first 2 days	0.833	0.009	3,019	2,285	1.355	0.011	0.814	0.851
Any problem accessing health care	0.653	0.008	13,990	11,402	1.878	0.012	0.637	0.668
Ever had a vaccination card	0.982	0.004	1,500	1,150	0.994	0.004	0.975	0.989
Received BCG vaccination	0.970	0.005	1,500	1,150	1.037	0.005	0.961	0.980
Received DPT-HepB+Hib vaccination (3 doses)	0.887	0.010	1,500	1,150	1.185	0.011	0.867	0.907
Received pneumococcal vaccination (3 doses)	0.911	0.008	1,500	1,150	1.068	0.009	0.895	0.927
Received measles vaccination	0.866	0.011	1,500	1,150	1.164	0.012	0.845	0.887
Fully vaccinated according to national schedule (12–23 months)	0.450	0.016	1,500	1,150	1.207	0.036	0.417	0.482
Received measles/measles-rubella 2 vaccination (24–35 months)	0.692	0.015	1,451	1,100	1.196	0.022	0.661	0.722
Fully vaccinated according to national schedule (24–35 months)	0.381	0.017	1,451	1,100	1.254	0.044	0.348	0.415
Received HPV vaccination (2 doses) women 15–19	0.413	0.011	3,176	2,519	1.271	0.027	0.391	0.435
Sought treatment for diarrhoea	0.566	0.022	698	592	1.158	0.038	0.523	0.610

Continued…

Table B.4—*Continued*

Variable	Value (R)	Standard error (SE)	Number of cases: Unweighted (N)	Number of cases: Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
Treated with ORS	0.418	0.022	698	592	1.171	0.053	0.374	0.462
Exclusive breastfeeding	0.680	0.019	756	558	1.116	0.028	0.642	0.718
Minimum dietary diversity (children 6–23 months)	0.384	0.013	2,163	1,646	1.215	0.033	0.358	0.409
Child slept under an ITN last night	0.494	0.010	8,678	6,481	1.467	0.020	0.475	0.514
Pregnant women slept under an ITN last night	0.481	0.023	745	544	1.187	0.048	0.434	0.527
Received 3+ doses of SP/Fansidar	0.165	0.008	3,019	2,285	1.204	0.049	0.149	0.181
Child has fever in last 2 weeks	0.132	0.005	7,641	5,722	1.212	0.039	0.122	0.142
Child had blood taken from finger/heel	0.309	0.019	945	754	1.189	0.061	0.271	0.346
Child took ACT	0.763	0.030	207	180	1.056	0.040	0.702	0.824
Condom use at last sex (women 15–24)	0.376	0.018	985	861	1.165	0.048	0.340	0.412
Mobile phone ownership	0.733	0.006	13,990	11,402	1.502	0.008	0.721	0.744

Table B.5 Sampling errors: Mombasa sample, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases: Unweighted (N)	Number of cases: Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
HOUSEHOLDS AND POPULATION								
Primary reliance on clean fuels and technology	0.411	0.073	1,913	2,969	3.103	0.177	0.265	0.556
Births registered with civil authority	0.546	0.059	215	345	1.569	0.108	0.428	0.664
Ownership of at least one ITN	0.529	0.044	576	866	2.115	0.083	0.441	0.618
Ownership of at least one ITN for every two persons	0.301	0.026	567	853	1.362	0.087	0.249	0.354
At least basic drinking water service	0.965	0.013	1,913	2,969	1.332	0.014	0.939	0.991
At least basic sanitation service	0.502	0.074	1,913	2,969	3.034	0.148	0.354	0.650
Using open defecation	0.004	0.003	1,913	2,969	0.992	0.745	0.000	0.010
Using a handwashing facility with soap and water	0.291	0.045	1,874	2,926	2.057	0.154	0.202	0.381
WOMEN								
No education	0.055	0.014	476	798	1.378	0.261	0.026	0.084
Secondary education or higher	0.601	0.046	476	798	2.030	0.076	0.509	0.692
Literacy	0.922	0.023	478	803	1.838	0.025	0.877	0.967
Use of the internet in last 12 months	0.762	0.036	478	803	1.853	0.048	0.689	0.834
Total fertility rate (3 years)	2.262	0.217	1,378	2,320	1.016	0.096	1.828	2.697
Currently pregnant	0.061	0.010	478	803	0.900	0.162	0.041	0.081
Mean number of children ever born to women age 40–49	3.398	0.146	71	107	0.806	0.043	3.105	3.691
Median birth interval	50.571	1.778	134	234	0.623	0.035	47.016	54.126
Ideal number of children	3.647	0.151	473	795	1.756	0.041	3.345	3.949
Total wanted fertility rate (3 years)	2.065	0.222	1,378	2,320	1.027	0.108	1.621	2.509
Currently using any contraceptive method	0.475	0.043	255	444	1.359	0.090	0.390	0.560
Currently using any modern method	0.443	0.045	255	444	1.431	0.101	0.353	0.532
Currently using pill	0.040	0.010	255	444	0.812	0.251	0.020	0.060
Currently using injectables	0.174	0.028	255	444	1.163	0.159	0.119	0.229
Currently using implants	0.153	0.035	255	444	1.555	0.230	0.083	0.224
Currently using male condoms	0.014	0.008	255	444	1.039	0.554	0.000	0.029
Currently using any traditional method	0.032	0.015	255	444	1.372	0.473	0.002	0.063
Unmet need for spacing	0.105	0.019	255	444	0.996	0.182	0.067	0.143
Unmet need for limiting	0.128	0.019	255	444	0.901	0.147	0.091	0.166
Unmet need total	0.234	0.033	255	444	1.247	0.142	0.167	0.300
Demand satisfied by modern methods (married women)	0.625	0.049	177	315	1.367	0.079	0.526	0.724
Demand satisfied by modern methods (all women)	0.647	0.045	218	376	1.420	0.070	0.556	0.738
Not exposed to any of the ten media sources	0.101	0.018	478	803	1.276	0.174	0.066	0.136
Neonatal mortality (last 0–9 years)	14.272	9.389	398	703	1.339	0.658	0.000	33.050
Post-neonatal mortality (last 0–9 years)	8.680	4.566	401	711	0.975	0.526	0.000	17.812
Infant mortality (last 0–9 years)	22.952	12.556	398	703	1.500	0.547	0.000	48.064
Child mortality (last 0–9 years)	16.080	8.600	398	704	1.160	0.535	0.000	33.280
Under-five mortality (last 0–9 years)	38.663	19.507	399	705	1.787	0.505	0.000	77.678
Stillbirth rate	22.840	15.419	207	354	1.085	0.675	0.000	53.678
Early neonatal rate	23.510	13.046	201	346	1.021	0.555	0.000	49.602
Perinatal mortality rate	45.814	19.847	207	354	1.070	0.433	6.120	85.508
Received ANC from a skilled provider	0.985	0.015	76	127	1.036	0.015	0.955	1.000
4+ ANC visits	0.856	0.044	76	127	1.093	0.052	0.768	0.945
8+ ANC visits	0.109	0.029	76	127	0.810	0.267	0.051	0.167
Took any iron-containing supplements	0.963	0.021	76	127	0.968	0.022	0.921	1.000
Mothers protected against tetanus for last birth	0.740	0.053	76	127	1.039	0.071	0.635	0.845
Delivered in a health facility (live births)	0.976	0.016	78	131	0.907	0.016	0.944	1.000
Delivered by a skilled provider (live births)	0.989	0.011	78	131	0.934	0.011	0.967	1.000
Delivered by C-section (live births)	0.239	0.052	78	131	1.051	0.217	0.135	0.343
Women with postnatal check during first 2 days	0.921	0.023	76	127	0.753	0.025	0.875	0.968
Newborns with postnatal check during first 2 days	0.907	0.045	76	127	1.332	0.050	0.817	0.997
Any problem accessing health care	0.468	0.063	478	803	2.722	0.134	0.343	0.593
Ever had a vaccination card	1.000	0.000	36	58	na	0.000	1.000	1.000
Received BCG vaccination	1.000	0.000	36	58	na	0.000	1.000	1.000
Received DPT-HepB+Hib vaccination (3 doses)	0.895	0.048	36	58	0.903	0.053	0.799	0.990
Received pneumococcal vaccination (3 doses)	0.954	0.032	36	58	0.887	0.034	0.890	1.000
Received measles vaccination	0.827	0.065	36	58	1.002	0.079	0.696	0.958
Fully vaccinated according to national schedule (12–23 months)	0.443	0.092	36	58	1.074	0.208	0.259	0.628
Received measles/measles-rubella 2 vaccination (24–35 months)	0.632	0.110	30	53	1.228	0.174	0.412	0.853
Fully vaccinated according to national schedule (24–35 months)	0.524	0.130	30	53	1.410	0.247	0.265	0.784
Received HPV vaccination (2 doses) women 15–19	0.252	0.054	61	96	0.958	0.213	0.145	0.359
Sought treatment for diarrhoea	0.618	0.109	18	33	0.976	0.176	0.400	0.835
Treated with ORS	0.556	0.150	18	33	1.314	0.270	0.256	0.856
Exclusive breastfeeding	0.797	0.102	16	25	0.986	0.128	0.593	1.000
Minimum dietary diversity (children 6–23 months)	0.378	0.058	56	95	0.883	0.152	0.262	0.493
Child slept under an ITN last night	0.594	0.056	209	337	1.501	0.095	0.481	0.706
Received 3+ doses of SP/Fansidar	0.328	0.062	76	127	1.144	0.189	0.204	0.453
Child has fever in last 2 weeks	0.119	0.024	190	329	1.006	0.198	0.072	0.166
Child had blood taken from finger/heel	0.366	0.097	24	39	0.961	0.265	0.172	0.560
Child took ACT	na	na	-	-	na	na	0.000	na
Condom use at last sex (women 15–24)	0.193	0.052	41	67	0.829	0.267	0.090	0.296
Mobile phone ownership	0.876	0.023	478	803	1.519	0.026	0.830	0.922

na = not applicable

Table B.6 Sampling errors: Kwale sample, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases: Unweighted (N)	Number of cases: Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
HOUSEHOLDS AND POPULATION								
Primary reliance on clean fuels and technology	0.064	0.011	2,568	1,586	1.168	0.178	0.041	0.087
Births registered with civil authority	0.875	0.031	322	202	1.322	0.036	0.812	0.937
Ownership of at least one ITN	0.705	0.028	527	318	1.419	0.040	0.649	0.762
Ownership of at least one ITN for every two persons	0.397	0.031	526	317	1.447	0.078	0.335	0.459
At least basic drinking water service	0.587	0.079	2,568	1,586	3.061	0.135	0.428	0.746
At least basic sanitation service	0.301	0.048	2,568	1,586	2.038	0.160	0.205	0.398
Using open defecation	0.352	0.071	2,568	1,586	2.788	0.201	0.210	0.494
Using a handwashing facility with soap and water	0.828	0.075	2,562	1,581	3.594	0.091	0.678	0.978
WOMEN								
No education	0.159	0.026	552	368	1.665	0.164	0.107	0.211
Secondary education or higher	0.423	0.029	552	368	1.355	0.067	0.366	0.480
Literacy	0.811	0.026	563	374	1.554	0.032	0.760	0.863
Use of the internet in last 12 months	0.459	0.053	563	374	2.489	0.115	0.354	0.564
Total fertility rate (3 years)	3.374	0.313	1,549	1,027	1.140	0.093	2.748	4.000
Currently pregnant	0.066	0.015	563	374	1.426	0.227	0.036	0.096
Mean number of children ever born to women age 40–49	5.613	0.374	110	72	1.340	0.067	4.865	6.360
Median birth interval	40.082	2.449	229	157	1.299	0.061	35.183	44.981
Ideal number of children	5.085	0.170	559	371	1.712	0.033	4.745	5.424
Total wanted fertility rate (3 years)	3.096	0.310	1,549	1,027	1.075	0.100	2.477	3.716
Currently using any contraceptive method	0.458	0.034	308	205	1.185	0.074	0.390	0.525
Currently using any modern method	0.455	0.034	308	205	1.190	0.074	0.387	0.522
Currently using pill	0.031	0.011	308	205	1.059	0.336	0.010	0.053
Currently using injectables	0.208	0.022	308	205	0.955	0.106	0.164	0.253
Currently using implants	0.202	0.025	308	205	1.080	0.122	0.153	0.252
Currently using male condoms	0.000	0.000	308	205	na	na	0.000	0.000
Currently using any traditional method	0.003	0.003	308	205	0.919	1.010	0.000	0.008
Unmet need for spacing	0.076	0.014	308	205	0.911	0.181	0.049	0.104
Unmet need for limiting	0.110	0.016	308	205	0.888	0.144	0.078	0.141
Unmet need total	0.186	0.017	308	205	0.753	0.090	0.153	0.219
Demand satisfied by modern methods (married women)	0.707	0.027	198	132	0.849	0.039	0.652	0.762
Demand satisfied by modern methods (all women)	0.707	0.025	239	161	0.861	0.036	0.657	0.758
Not exposed to any of the ten media sources	0.048	0.008	563	374	0.884	0.166	0.032	0.064
Neonatal mortality (last 0–9 years)	14.708	6.516	606	406	1.311	0.443	1.675	27.740
Post-neonatal mortality (last 0–9 years)	14.879	5.716	611	410	1.182	0.384	3.446	26.311
Infant mortality (last 0–9 years)	29.586	9.045	607	407	1.387	0.306	11.496	47.677
Child mortality (last 0–9 years)	7.499	3.778	609	409	1.088	0.504	0.000	15.055
Under-five mortality (last 0–9 years)	36.864	11.442	607	407	1.442	0.310	13.980	59.747
Stillbirth rate	8.586	5.644	299	202	1.077	0.657	0.000	19.874
Early neonatal rate	9.893	7.013	297	200	1.245	0.709	0.000	23.920
Perinatal mortality rate	18.395	10.462	299	202	1.389	0.569	0.000	39.319
Received ANC from a skilled provider	1.000	0.000	106	71	na	0.000	1.000	1.000
4+ ANC visits	0.844	0.043	106	71	1.211	0.051	0.758	0.930
8+ ANC visits	0.067	0.021	106	71	0.881	0.321	0.024	0.109
Took any iron-containing supplements	0.961	0.017	106	71	0.885	0.017	0.928	0.995
Mothers protected against tetanus for last birth	0.736	0.058	106	71	1.343	0.079	0.619	0.852
Delivered in a health facility (live births)	0.920	0.033	107	71	1.251	0.036	0.854	0.985
Delivered by a skilled provider (live births)	0.920	0.033	107	71	1.251	0.036	0.854	0.985
Delivered by C-section (live births)	0.093	0.023	107	71	0.810	0.243	0.048	0.139
Women with postnatal check during first 2 days	0.878	0.030	106	71	0.928	0.034	0.819	0.938
Newborns with postnatal check during first 2 days	0.919	0.022	106	71	0.811	0.023	0.876	0.962
Any problem accessing health care	0.675	0.037	563	374	1.858	0.054	0.602	0.749
Ever had a vaccination card	1.000	0.000	51	35	na	0.000	1.000	1.000
Received BCG vaccination	1.000	0.000	51	35	na	0.000	1.000	1.000
Received DPT-HepB+Hib vaccination (3 doses)	0.975	0.025	51	35	1.140	0.025	0.925	1.000
Received pneumococcal vaccination (3 doses)	0.975	0.025	51	35	1.140	0.025	0.925	1.000
Received measles vaccination	0.901	0.047	51	35	1.150	0.053	0.806	0.996
Fully vaccinated according to national schedule (12–23 months)	0.594	0.071	51	35	1.053	0.120	0.452	0.737
Received measles/measles-rubella 2 vaccination (24–35 months)	0.826	0.062	47	32	1.123	0.075	0.703	0.950
Fully vaccinated according to national schedule (24–35 months)	0.783	0.068	47	32	1.132	0.087	0.647	0.918
Received HPV vaccination (2 doses) women 15–19	0.370	0.061	138	94	1.473	0.165	0.248	0.492
Sought treatment for diarrhoea	0.449	0.111	26	18	1.169	0.248	0.226	0.671
Treated with ORS	0.272	0.095	26	18	1.110	0.348	0.083	0.461
Exclusive breastfeeding	0.861	0.069	22	14	0.913	0.080	0.723	0.998
Minimum dietary diversity (children 6–23 months)	0.176	0.054	81	55	1.255	0.305	0.069	0.283
Child slept under an ITN last night	0.634	0.031	321	200	1.043	0.050	0.571	0.697
Received 3+ doses of SP/Fansidar	0.516	0.060	106	71	1.232	0.117	0.395	0.636
Child has fever in last 2 weeks	0.099	0.029	285	192	1.559	0.289	0.042	0.156
Child had blood taken from finger/heel	0.494	0.138	29	19	1.411	0.279	0.218	0.770
Child took ACT	0.145	0.147	9	6	1.262	1.013	0.000	0.438
Condom use at last sex (women 15–24)	0.257	0.074	30	21	0.912	0.287	0.109	0.404
Mobile phone ownership	0.686	0.028	563	374	1.414	0.040	0.630	0.741

na = not applicable

Table B.7 Sampling errors: Kilifi sample, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases: Unweighted (N)	Number of cases: Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
HOUSEHOLDS AND POPULATION								
Primary reliance on clean fuels and technology	0.126	0.029	2,254	2,666	2.106	0.226	0.069	0.183
Births registered with civil authority	0.663	0.041	252	302	1.182	0.062	0.581	0.745
Ownership of at least one ITN	0.739	0.026	568	699	1.385	0.035	0.688	0.790
Ownership of at least one ITN for every two persons	0.531	0.028	563	693	1.352	0.054	0.475	0.588
At least basic drinking water service	0.799	0.061	2,254	2,666	2.850	0.076	0.677	0.921
At least basic sanitation service	0.357	0.039	2,254	2,666	1.613	0.109	0.279	0.435
Using open defecation	0.208	0.050	2,254	2,666	2.326	0.240	0.108	0.308
Using a handwashing facility with soap and water	0.468	0.027	2,254	2,666	1.074	0.057	0.415	0.522
WOMEN								
No education	0.096	0.016	515	608	1.269	0.172	0.063	0.129
Secondary education or higher	0.518	0.034	515	608	1.562	0.067	0.449	0.587
Literacy	0.866	0.017	516	608	1.137	0.020	0.831	0.900
Use of the internet in last 12 months	0.563	0.039	516	608	1.778	0.069	0.485	0.641
Total fertility rate (3 years)	2.976	0.279	1,412	1,677	1.126	0.094	2.418	3.533
Currently pregnant	0.044	0.012	516	608	1.370	0.283	0.019	0.068
Mean number of children ever born to women age 40–49	5.260	0.494	76	85	1.421	0.094	4.272	6.247
Median birth interval	36.374	3.366	176	200	1.162	0.093	29.643	43.105
Ideal number of children	4.657	0.178	516	608	1.671	0.038	4.302	5.013
Total wanted fertility rate (3 years)	2.555	0.266	1,412	1,677	1.177	0.104	2.022	3.088
Currently using any contraceptive method	0.630	0.043	257	302	1.419	0.068	0.544	0.716
Currently using any modern method	0.572	0.040	257	302	1.291	0.070	0.492	0.652
Currently using pill	0.052	0.012	257	302	0.885	0.236	0.027	0.076
Currently using injectables	0.212	0.034	257	302	1.337	0.162	0.143	0.280
Currently using implants	0.248	0.026	257	302	0.949	0.103	0.197	0.300
Currently using male condoms	0.013	0.009	257	302	1.326	0.723	0.000	0.032
Currently using any traditional method	0.058	0.018	257	302	1.249	0.315	0.021	0.094
Unmet need for spacing	0.079	0.023	257	302	1.340	0.286	0.034	0.125
Unmet need for limiting	0.061	0.017	257	302	1.161	0.284	0.027	0.096
Unmet need total	0.141	0.028	257	302	1.301	0.201	0.084	0.197
Demand satisfied by modern methods (married women)	0.742	0.037	199	233	1.174	0.049	0.669	0.815
Demand satisfied by modern methods (all women)	0.730	0.035	257	310	1.289	0.048	0.659	0.801
Not exposed to any of the ten media sources	0.096	0.017	516	608	1.333	0.180	0.061	0.131
Neonatal mortality (last 0–9 years)	10.510	3.935	516	606	0.876	0.374	2.640	18.380
Post-neonatal mortality (last 0–9 years)	22.921	6.628	521	613	1.038	0.289	9.666	36.176
Infant mortality (last 0–9 years)	33.431	7.171	518	610	0.947	0.214	19.090	47.772
Child mortality (last 0–9 years)	7.164	3.379	529	620	0.965	0.472	0.406	13.922
Under-five mortality (last 0–9 years)	40.356	7.615	519	610	0.957	0.189	25.126	55.585
Stillbirth rate	10.346	5.555	244	291	0.866	0.537	0.000	21.457
Early neonatal rate	11.324	6.637	242	289	0.982	0.586	0.000	24.597
Perinatal mortality rate	21.588	9.206	244	291	0.999	0.426	3.176	40.000
Received ANC from a skilled provider	1.000	0.000	85	102	na	0.000	1.000	1.000
4+ ANC visits	0.834	0.048	85	102	1.180	0.058	0.738	0.930
8+ ANC visits	0.091	0.027	85	102	0.848	0.292	0.038	0.144
Took any iron-containing supplements	0.970	0.021	85	102	1.118	0.022	0.928	1.000
Mothers protected against tetanus for last birth	0.947	0.023	85	102	0.928	0.024	0.902	0.992
Delivered in a health facility (live births)	0.888	0.044	88	105	1.213	0.049	0.801	0.975
Delivered by a skilled provider (live births)	0.886	0.042	88	105	1.171	0.048	0.801	0.971
Delivered by C-section (live births)	0.087	0.036	88	105	1.192	0.412	0.015	0.159
Women with postnatal check during first 2 days	0.919	0.030	85	102	1.014	0.033	0.859	0.979
Newborns with postnatal check during first 2 days	0.918	0.028	85	102	0.936	0.030	0.863	0.974
Any problem accessing health care	0.716	0.033	516	608	1.669	0.046	0.650	0.783
Ever had a vaccination card	1.000	0.000	45	54	na	0.000	1.000	1.000
Received BCG vaccination	1.000	0.000	45	54	na	0.000	1.000	1.000
Received DPT-HepB+Hib vaccination (3 doses)	0.950	0.034	45	54	1.051	0.036	0.881	1.000
Received pneumococcal vaccination (3 doses)	0.896	0.042	45	54	0.910	0.046	0.813	0.979
Received measles vaccination	0.916	0.055	45	54	1.318	0.060	0.807	1.000
Fully vaccinated according to national schedule (12–23 months)	0.576	0.061	45	54	0.819	0.106	0.454	0.698
Received measles/measles-rubella 2 vaccination (24–35 months)	0.853	0.050	49	60	0.999	0.059	0.752	0.953
Fully vaccinated according to national schedule (24–35 months)	0.530	0.091	49	60	1.231	0.172	0.348	0.712
Received HPV vaccination (2 doses) women 15–19	0.637	0.048	126	137	1.120	0.076	0.540	0.733
Sought treatment for diarrhoea	0.787	0.086	28	36	1.170	0.109	0.615	0.958
Treated with ORS	0.651	0.080	28	36	0.883	0.123	0.490	0.812
Exclusive breastfeeding	0.609	0.089	22	25	0.845	0.147	0.430	0.788
Minimum dietary diversity (children 6–23 months)	0.271	0.062	59	71	1.068	0.230	0.146	0.395
Child slept under an ITN last night	0.715	0.050	245	290	1.343	0.069	0.616	0.814
Received 3+ doses of SP/Fansidar	0.637	0.053	85	102	1.016	0.084	0.530	0.744
Child has fever in last 2 weeks	0.151	0.043	235	280	1.576	0.284	0.065	0.237
Child had blood taken from finger/heel	0.336	0.097	33	42	1.042	0.289	0.142	0.530
Child took ACT	0.467	0.191	7	9	0.923	0.409	0.085	0.849
Condom use at last sex (women 15–24)	0.189	0.090	35	48	1.326	0.477	0.009	0.369
Mobile phone ownership	0.698	0.031	516	608	1.533	0.045	0.635	0.760

na = not applicable

Table B.8 Sampling errors: Tana River sample, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases: Unweighted (N)	Number of cases: Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
HOUSEHOLDS AND POPULATION								
Primary reliance on clean fuels and technology	0.025	0.011	2,054	721	1.884	0.451	0.002	0.047
Births registered with civil authority	0.650	0.041	311	106	1.246	0.064	0.567	0.732
Ownership of at least one ITN	0.749	0.037	416	143	1.744	0.050	0.674	0.823
Ownership of at least one ITN for every two persons	0.396	0.044	416	143	1.822	0.111	0.308	0.484
At least basic drinking water service	0.553	0.127	2,054	721	4.456	0.229	0.300	0.806
At least basic sanitation service	0.262	0.051	2,054	721	2.105	0.193	0.161	0.364
Using open defecation	0.304	0.065	2,054	721	2.484	0.214	0.174	0.433
Using a handwashing facility with soap and water	0.219	0.050	2,054	721	2.227	0.231	0.118	0.320
WOMEN								
No education	0.371	0.075	398	137	3.068	0.203	0.221	0.522
Secondary education or higher	0.384	0.047	398	137	1.919	0.122	0.290	0.478
Literacy	0.604	0.070	400	137	2.841	0.116	0.464	0.744
Use of the internet in last 12 months	0.462	0.079	400	137	3.123	0.171	0.304	0.619
Total fertility rate (3 years)	5.652	0.554	1,106	377	1.808	0.098	4.545	6.760
Currently pregnant	0.098	0.023	400	137	1.513	0.230	0.053	0.144
Mean number of children ever born to women age 40–49	6.176	0.499	67	24	1.614	0.081	5.177	7.175
Median birth interval	33.969	3.187	240	81	1.965	0.094	27.596	40.343
Ideal number of children	5.836	0.336	400	137	2.686	0.058	5.165	6.508
Total wanted fertility rate (3 years)	5.180	0.506	1,106	377	1.755	0.098	4.168	6.193
Currently using any contraceptive method	0.218	0.057	249	88	2.151	0.260	0.105	0.332
Currently using any modern method	0.218	0.057	249	88	2.151	0.260	0.105	0.332
Currently using pill	0.012	0.005	249	88	0.765	0.442	0.001	0.022
Currently using injectables	0.056	0.021	249	88	1.460	0.381	0.013	0.099
Currently using implants	0.144	0.041	249	88	1.841	0.287	0.061	0.226
Currently using male condoms	0.000	0.000	249	88	na	na	0.000	0.000
Currently using any traditional method	0.000	0.000	249	88	na	na	0.000	0.000
Unmet need for spacing	0.119	0.018	249	88	0.853	0.148	0.084	0.154
Unmet need for limiting	0.174	0.028	249	88	1.156	0.160	0.119	0.230
Unmet need total	0.293	0.032	249	88	1.111	0.110	0.229	0.357
Demand satisfied by modern methods (married women)	0.427	0.073	129	45	1.650	0.170	0.281	0.572
Demand satisfied by modern methods (all women)	0.450	0.069	151	51	1.678	0.152	0.313	0.587
Not exposed to any of the ten media sources	0.037	0.017	400	137	1.821	0.468	0.002	0.071
Neonatal mortality (last 0–9 years)	29.634	5.597	572	198	0.811	0.189	18.440	40.827
Post-neonatal mortality (last 0–9 years)	14.907	4.746	568	197	0.935	0.318	5.416	24.398
Infant mortality (last 0–9 years)	44.541	8.039	572	198	0.938	0.180	28.463	60.619
Child mortality (last 0–9 years)	9.087	5.448	562	195	1.010	0.600	0.000	19.983
Under-five mortality (last 0–9 years)	53.223	11.423	573	198	1.122	0.215	30.378	76.069
Stillbirth rate	12.454	5.692	304	104	0.906	0.457	1.069	23.839
Early neonatal rate	40.233	8.883	300	102	0.792	0.221	22.466	57.999
Perinatal mortality rate	52.185	10.940	304	104	0.856	0.210	30.305	74.066
Received ANC from a skilled provider	0.986	0.014	115	40	1.277	0.014	0.958	1.000
4+ ANC visits	0.751	0.038	115	40	0.939	0.051	0.675	0.827
8+ ANC visits	0.043	0.026	115	40	1.386	0.618	0.000	0.095
Took any iron-containing supplements	0.870	0.032	115	40	1.003	0.036	0.807	0.934
Mothers protected against tetanus for last birth	0.786	0.060	115	40	1.546	0.076	0.667	0.906
Delivered in a health facility (live births)	0.750	0.084	119	42	2.018	0.113	0.581	0.919
Delivered by a skilled provider (live births)	0.756	0.081	119	42	2.013	0.108	0.593	0.919
Delivered by C-section (live births)	0.085	0.028	119	42	1.105	0.334	0.028	0.142
Women with postnatal check during first 2 days	0.906	0.038	115	40	1.366	0.041	0.831	0.981
Newborns with postnatal check during first 2 days	0.897	0.039	115	40	1.374	0.044	0.818	0.976
Any problem accessing health care	0.744	0.039	400	137	1.766	0.052	0.666	0.821
Ever had a vaccination card	0.987	0.013	61	21	0.877	0.013	0.961	1.000
Received BCG vaccination	0.901	0.050	61	21	1.170	0.055	0.801	1.000
Received DPT-HepB+Hib vaccination (3 doses)	0.914	0.032	61	21	0.878	0.035	0.851	0.977
Received pneumococcal vaccination (3 doses)	0.949	0.024	61	21	0.861	0.025	0.901	0.998
Received measles vaccination	0.827	0.075	61	21	1.454	0.090	0.678	0.977
Fully vaccinated according to national schedule (12–23 months)	0.307	0.069	61	21	1.157	0.224	0.169	0.445
Received measles/measles-rubella 2 vaccination (24–35 months)	0.569	0.073	54	18	1.064	0.128	0.424	0.714
Fully vaccinated according to national schedule (24–35 months)	0.224	0.066	54	18	1.153	0.296	0.092	0.357
Received HPV vaccination (2 doses) women 15–19	0.319	0.056	99	35	1.188	0.176	0.207	0.431
Sought treatment for diarrhoea	0.722	0.096	30	10	1.044	0.133	0.531	0.914
Treated with ORS	0.310	0.110	30	10	1.253	0.356	0.090	0.531
Exclusive breastfeeding	0.478	0.117	20	7	1.018	0.244	0.244	0.711
Minimum dietary diversity (children 6–23 months)	0.117	0.036	89	31	1.058	0.309	0.045	0.190
Child slept under an ITN last night	0.718	0.043	305	104	1.355	0.059	0.633	0.804
Received 3+ doses of SP/Fansidar	0.046	0.021	115	40	1.097	0.469	0.003	0.089
Child has fever in last 2 weeks	0.112	0.021	280	96	1.056	0.189	0.070	0.154
Child had blood taken from finger/heel	0.308	0.137	30	11	1.466	0.446	0.034	0.583
Child took ACT	0.453	0.350	2	1	1.187	0.773	0.000	1.000
Condom use at last sex (women 15–24)	0.354	0.176	17	5	1.421	0.496	0.003	0.705
Mobile phone ownership	0.642	0.049	400	137	2.024	0.076	0.545	0.740

na = not applicable

Table B.9 Sampling errors: Lamu sample, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases: Unweighted (N)	Number of cases: Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
HOUSEHOLDS AND POPULATION								
Primary reliance on clean fuels and technology	0.095	0.023	2,038	261	1.645	0.238	0.050	0.140
Births registered with civil authority	0.863	0.030	228	31	1.154	0.034	0.804	0.922
Ownership of at least one ITN	0.680	0.049	487	62	2.294	0.072	0.582	0.777
Ownership of at least one ITN for every two persons	0.479	0.044	486	62	1.946	0.092	0.390	0.567
At least basic drinking water service	0.787	0.064	2,038	261	2.827	0.081	0.659	0.915
At least basic sanitation service	0.676	0.083	2,038	261	3.272	0.123	0.510	0.842
Using open defecation	0.148	0.078	2,038	261	3.701	0.528	0.000	0.305
Using a handwashing facility with soap and water	0.260	0.051	2,012	258	2.208	0.196	0.158	0.362
WOMEN								
No education	0.131	0.066	416	53	3.920	0.504	0.000	0.263
Secondary education or higher	0.513	0.053	416	53	2.145	0.103	0.407	0.618
Literacy	0.830	0.065	417	54	3.501	0.079	0.700	0.961
Use of the internet in last 12 months	0.525	0.068	417	54	2.753	0.130	0.389	0.660
Total fertility rate (3 years)	3.583	0.551	1,153	148	1.581	0.154	2.480	4.685
Currently pregnant	0.066	0.011	417	54	0.903	0.167	0.044	0.088
Mean number of children ever born to women age 40–49	4.623	0.321	85	10	1.320	0.069	3.980	5.266
Median birth interval	42.277	6.217	161	22	1.909	0.147	29.843	54.711
Ideal number of children	4.336	0.281	412	53	2.824	0.065	3.775	4.898
Total wanted fertility rate (3 years)	3.177	0.557	1,153	148	1.749	0.175	2.063	4.291
Currently using any contraceptive method	0.399	0.065	207	27	1.900	0.163	0.269	0.530
Currently using any modern method	0.396	0.065	207	27	1.896	0.164	0.266	0.526
Currently using pill	0.029	0.016	207	27	1.365	0.554	0.000	0.061
Currently using injectables	0.177	0.039	207	27	1.469	0.221	0.099	0.255
Currently using implants	0.131	0.027	207	27	1.161	0.208	0.077	0.186
Currently using male condoms	0.005	0.005	207	27	1.029	1.024	0.000	0.015
Currently using any traditional method	0.004	0.004	207	27	0.906	1.044	0.000	0.011
Unmet need for spacing	0.111	0.021	207	27	0.980	0.194	0.068	0.153
Unmet need for limiting	0.139	0.021	207	27	0.882	0.153	0.096	0.181
Unmet need total	0.249	0.034	207	27	1.137	0.138	0.181	0.318
Demand satisfied by modern methods (married women)	0.610	0.063	136	17	1.485	0.103	0.484	0.736
Demand satisfied by modern methods (all women)	0.617	0.047	170	22	1.271	0.077	0.522	0.711
Not exposed to any of the ten media sources	0.337	0.052	417	54	2.225	0.154	0.233	0.440
Neonatal mortality (last 0–9 years)	13.280	6.946	427	57	1.112	0.523	0.000	27.172
Post-neonatal mortality (last 0–9 years)	12.369	4.341	426	57	0.826	0.351	3.686	21.052
Infant mortality (last 0–9 years)	25.649	7.724	428	57	0.947	0.301	10.200	41.098
Child mortality (last 0–9 years)	2.477	2.482	425	57	0.933	1.002	0.000	7.440
Under-five mortality (last 0–9 years)	28.063	7.398	428	57	0.874	0.264	13.267	42.859
Stillbirth rate	22.542	9.598	220	29	0.994	0.426	3.346	41.738
Early neonatal rate	10.536	7.035	215	29	1.031	0.668	0.000	24.605
Perinatal mortality rate	32.841	9.378	220	29	0.812	0.286	14.085	51.597
Received ANC from a skilled provider	0.979	0.015	78	10	0.949	0.016	0.948	1.000
4+ ANC visits	0.636	0.079	78	10	1.437	0.125	0.478	0.795
8+ ANC visits	0.019	0.014	78	10	0.880	0.718	0.000	0.047
Took any iron-containing supplements	0.976	0.017	78	10	0.957	0.017	0.942	1.000
Mothers protected against tetanus for last birth	0.792	0.067	78	10	1.432	0.084	0.658	0.925
Delivered in a health facility (live births)	0.795	0.110	80	11	2.407	0.138	0.575	1.000
Delivered by a skilled provider (live births)	0.816	0.100	80	11	2.285	0.123	0.616	1.000
Delivered by C-section (live births)	0.271	0.043	80	11	0.848	0.160	0.185	0.358
Women with postnatal check during first 2 days	0.794	0.094	78	10	1.993	0.118	0.606	0.981
Newborns with postnatal check during first 2 days	0.834	0.085	78	10	1.961	0.102	0.664	1.000
Any problem accessing health care	0.849	0.019	417	54	1.063	0.022	0.812	0.886
Ever had a vaccination card	1.000	0.000	34	5	na	0.000	1.000	1.000
Received BCG vaccination	0.971	0.029	34	5	1.009	0.030	0.914	1.000
Received DPT-HepB+Hib vaccination (3 doses)	0.950	0.037	34	5	0.986	0.039	0.877	1.000
Received pneumococcal vaccination (3 doses)	0.912	0.053	34	5	1.108	0.058	0.806	1.000
Received measles vaccination	0.853	0.078	34	5	1.290	0.091	0.697	1.000
Fully vaccinated according to national schedule (12–23 months)	0.336	0.097	34	5	1.200	0.289	0.141	0.531
Received measles/measles-rubella 2 vaccination (24–35 months)	0.739	0.051	43	6	0.773	0.069	0.637	0.842
Fully vaccinated according to national schedule (24–35 months)	0.360	0.085	43	6	1.173	0.237	0.189	0.531
Received HPV vaccination (2 doses) women 15–19	0.412	0.061	106	14	1.261	0.147	0.291	0.534
Sought treatment for diarrhoea	0.267	0.159	4	-	0.697	0.595	0.000	0.586
Treated with ORS	0.535	0.318	4	-	1.235	0.595	0.000	1.000
Exclusive breastfeeding	0.647	0.109	23	3	1.067	0.169	0.429	0.865
Minimum dietary diversity (children 6–23 months)	0.155	0.058	53	7	1.155	0.375	0.039	0.271
Child slept under an ITN last night	0.574	0.076	231	31	1.915	0.133	0.422	0.726
Received 3+ doses of SP/Fansidar	0.254	0.053	78	10	1.063	0.208	0.149	0.360
Child has fever in last 2 weeks	0.035	0.014	209	28	1.091	0.392	0.008	0.062
Child had blood taken from finger/heel	0.136	0.112	7	1	0.885	0.821	0.000	0.359
Child took ACT	na	na	-	-	na	na	0.000	na
Condom use at last sex (women 15–24)	0.326	0.127	11	2	0.866	0.389	0.072	0.579
Mobile phone ownership	0.751	0.032	417	54	1.513	0.043	0.687	0.815

na = not applicable

Table B.10 Sampling errors: Taita Taveta sample, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases Unweighted (N)	Number of cases Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits Lower (R-2SE)	Confidence limits Upper (R+2SE)
HOUSEHOLDS AND POPULATION								
Primary reliance on clean fuels and technology	0.155	0.037	1,474	691	2.070	0.239	0.081	0.229
Births registered with civil authority	0.746	0.042	161	81	1.108	0.056	0.662	0.829
Ownership of at least one ITN	0.761	0.030	441	206	1.465	0.039	0.702	0.821
Ownership of at least one ITN for every two persons	0.584	0.029	436	204	1.246	0.050	0.526	0.643
At least basic drinking water service	0.766	0.089	1,474	691	3.501	0.116	0.588	0.944
At least basic sanitation service	0.505	0.055	1,474	691	1.970	0.109	0.395	0.616
Using open defecation	0.035	0.030	1,474	691	2.433	0.856	0.000	0.096
Using a handwashing facility with soap and water	0.173	0.036	1,467	688	1.724	0.208	0.101	0.245
WOMEN								
No education	0.013	0.007	279	145	1.009	0.522	0.000	0.027
Secondary education or higher	0.613	0.024	279	145	0.811	0.039	0.566	0.661
Literacy	0.968	0.013	297	153	1.257	0.013	0.942	0.994
Use of the internet in last 12 months	0.701	0.031	297	153	1.174	0.045	0.638	0.763
Total fertility rate (3 years)	3.228	0.292	851	439	0.976	0.090	2.645	3.812
Currently pregnant	0.046	0.015	297	153	1.214	0.320	0.017	0.076
Mean number of children ever born to women age 40–49	4.034	0.262	72	35	1.234	0.065	3.509	4.558
Median birth interval	47.154	4.386	124	67	0.709	0.093	38.382	55.926
Ideal number of children	3.772	0.137	297	153	1.336	0.036	3.498	4.047
Total wanted fertility rate (3 years)	2.727	0.296	851	439	0.979	0.109	2.134	3.319
Currently using any contraceptive method	0.722	0.031	163	85	0.870	0.042	0.661	0.783
Currently using any modern method	0.650	0.030	163	85	0.794	0.046	0.591	0.710
Currently using pill	0.086	0.024	163	85	1.108	0.284	0.037	0.135
Currently using injectables	0.205	0.024	163	85	0.770	0.119	0.157	0.254
Currently using implants	0.261	0.043	163	85	1.250	0.166	0.174	0.347
Currently using male condoms	0.029	0.013	163	85	0.959	0.438	0.004	0.054
Currently using any traditional method	0.071	0.027	163	85	1.326	0.377	0.018	0.125
Unmet need for spacing	0.039	0.016	163	85	1.061	0.416	0.006	0.071
Unmet need for limiting	0.072	0.024	163	85	1.168	0.330	0.025	0.120
Unmet need total	0.111	0.027	163	85	1.088	0.242	0.057	0.164
Demand satisfied by modern methods (married women)	0.781	0.037	136	71	1.028	0.047	0.708	0.854
Demand satisfied by modern methods (all women)	0.773	0.037	178	91	1.179	0.048	0.699	0.848
Not exposed to any of the ten media sources	0.042	0.009	297	153	0.756	0.209	0.024	0.060
Neonatal mortality (last 0–9 years)	26.800	10.473	305	159	1.059	0.391	5.853	47.747
Post-neonatal mortality (last 0–9 years)	15.230	7.307	304	159	0.971	0.480	0.616	29.844
Infant mortality (last 0–9 years)	42.030	13.442	305	159	1.113	0.320	15.146	68.914
Child mortality (last 0–9 years)	3.870	3.899	311	163	1.010	1.008	0.000	11.669
Under-five mortality (last 0–9 years)	45.737	12.958	306	160	1.053	0.283	19.821	71.654
Stillbirth rate	12.333	8.319	160	86	0.988	0.675	0.000	28.971
Early neonatal rate	11.931	7.852	158	85	0.947	0.658	0.000	27.634
Perinatal mortality rate	24.117	9.952	160	86	0.870	0.413	4.213	44.021
Received ANC from a skilled provider	0.968	0.029	55	30	1.198	0.030	0.911	1.000
4+ ANC visits	0.735	0.050	55	30	0.830	0.068	0.635	0.834
8+ ANC visits	0.119	0.045	55	30	1.022	0.379	0.029	0.208
Took any iron-containing supplements	0.888	0.037	55	30	0.866	0.042	0.814	0.962
Mothers protected against tetanus for last birth	0.886	0.041	55	30	0.959	0.047	0.803	0.969
Delivered in a health facility (live births)	0.893	0.041	59	31	1.037	0.046	0.812	0.975
Delivered by a skilled provider (live births)	0.913	0.032	59	31	0.897	0.035	0.849	0.977
Delivered by C-section (live births)	0.154	0.073	59	31	1.552	0.474	0.008	0.301
Women with postnatal check during first 2 days	0.819	0.087	55	30	1.641	0.107	0.644	0.994
Newborns with postnatal check during first 2 days	0.870	0.059	55	30	1.276	0.068	0.752	0.987
Any problem accessing health care	0.587	0.036	297	153	1.269	0.062	0.514	0.659
Ever had a vaccination card	1.000	0.000	26	14	na	0.000	1.000	1.000
Received BCG vaccination	1.000	0.000	26	14	na	0.000	1.000	1.000
Received DPT-HepB+Hib vaccination (3 doses)	0.950	0.049	26	14	1.162	0.051	0.853	1.000
Received pneumococcal vaccination (3 doses)	1.000	0.000	26	14	na	0.000	1.000	1.000
Received measles vaccination	0.975	0.026	26	14	0.858	0.026	0.924	1.000
Fully vaccinated according to national schedule (12–23 months)	0.666	0.088	26	14	0.977	0.132	0.489	0.842
Received measles/measles-rubella 2 vaccination (24–35 months)	1.000	0.000	26	14	na	0.000	1.000	1.000
Fully vaccinated according to national schedule (24–35 months)	0.564	0.110	26	14	1.143	0.196	0.343	0.785
Received HPV vaccination (2 doses) women 15–19	0.427	0.116	50	25	1.620	0.273	0.194	0.660
Sought treatment for diarrhoea	0.640	0.150	18	9	1.292	0.235	0.339	0.941
Treated with ORS	0.584	0.153	18	9	1.282	0.263	0.277	0.891
Exclusive breastfeeding	0.780	0.119	14	6	1.031	0.152	0.543	1.000
Minimum dietary diversity (children 6–23 months)	0.591	0.095	37	21	1.149	0.160	0.402	0.780
Child slept under an ITN last night	0.569	0.054	172	85	1.265	0.095	0.461	0.676
Received 3+ doses of SP/Fansidar	0.340	0.061	55	30	0.949	0.180	0.218	0.462
Child has fever in last 2 weeks	0.160	0.035	149	80	1.185	0.220	0.090	0.230
Child had blood taken from finger/heel	0.344	0.091	22	13	0.949	0.264	0.163	0.526
Child took ACT	0.614	0.335	2	2	1.184	0.546	0.000	1.000
Condom use at last sex (women 15–24)	0.425	0.080	27	14	0.827	0.188	0.265	0.584
Mobile phone ownership	0.885	0.022	297	153	1.191	0.025	0.841	0.929

na = not applicable

Table B.11 Sampling errors: Garissa sample, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases: Unweighted (N)	Number of cases: Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
HOUSEHOLDS AND POPULATION								
Primary reliance on clean fuels and technology	0.037	0.008	2,647	852	0.969	0.209	0.021	0.052
Births registered with civil authority	0.483	0.080	377	123	2.443	0.165	0.324	0.642
Ownership of at least one ITN	0.084	0.017	453	145	1.322	0.206	0.049	0.118
Ownership of at least one ITN for every two persons	0.030	0.010	451	145	1.286	0.343	0.009	0.051
At least basic drinking water service	0.601	0.077	2,647	852	2.951	0.128	0.447	0.755
At least basic sanitation service	0.284	0.044	2,647	852	1.738	0.154	0.197	0.371
Using open defecation	0.330	0.086	2,647	852	3.463	0.259	0.159	0.501
Using a handwashing facility with soap and water	0.182	0.029	2,647	852	1.441	0.162	0.123	0.241
WOMEN								
No education	0.557	0.053	516	168	2.392	0.095	0.451	0.662
Secondary education or higher	0.262	0.037	516	168	1.915	0.142	0.188	0.336
Literacy	0.434	0.044	516	168	2.020	0.102	0.345	0.522
Use of the internet in last 12 months	0.510	0.048	516	168	2.157	0.093	0.415	0.606
Total fertility rate (3 years)	5.304	0.804	1,413	461	1.680	0.152	3.696	6.912
Currently pregnant	0.086	0.013	516	168	1.086	0.156	0.059	0.113
Mean number of children ever born to women age 40–49	6.801	0.303	69	22	0.866	0.045	6.195	7.406
Median birth interval	25.285	0.697	328	107	1.160	0.028	23.891	26.679
Ideal number of children	7.732	0.232	516	168	1.771	0.030	7.268	8.196
Total wanted fertility rate (3 years)	5.184	0.809	1,413	461	1.735	0.156	3.566	6.801
Currently using any contraceptive method	0.054	0.020	283	90	1.509	0.377	0.013	0.095
Currently using any modern method	0.050	0.020	283	90	1.577	0.412	0.009	0.090
Currently using pill	0.004	0.004	283	90	1.089	0.996	0.000	0.013
Currently using injectables	0.023	0.010	283	90	1.076	0.420	0.004	0.042
Currently using implants	0.020	0.010	283	90	1.210	0.505	0.000	0.040
Currently using male condoms	0.000	0.000	283	90	na	na	0.000	0.000
Currently using any traditional method	0.004	0.003	283	90	0.837	0.753	0.000	0.011
Unmet need for spacing	0.097	0.015	283	90	0.864	0.157	0.067	0.128
Unmet need for limiting	0.067	0.019	283	90	1.280	0.285	0.029	0.105
Unmet need total	0.164	0.024	283	90	1.072	0.144	0.117	0.212
Demand satisfied by modern methods (married women)	0.227	0.081	61	20	1.505	0.355	0.066	0.388
Demand satisfied by modern methods (all women)	0.284	0.079	73	23	1.453	0.277	0.127	0.442
Not exposed to any of the ten media sources	0.386	0.035	516	168	1.647	0.092	0.316	0.457
Neonatal mortality (last 0–9 years)	34.970	5.125	816	260	0.749	0.147	24.719	45.220
Post-neonatal mortality (last 0–9 years)	14.833	3.888	812	259	0.872	0.262	7.058	22.608
Infant mortality (last 0–9 years)	49.803	7.665	816	260	0.958	0.154	34.472	65.133
Child mortality (last 0–9 years)	10.007	3.322	805	258	0.870	0.332	3.362	16.652
Under-five mortality (last 0–9 years)	59.312	7.385	817	261	0.820	0.125	44.541	74.082
Stillbirth rate	19.073	7.698	396	128	0.905	0.404	3.677	34.469
Early neonatal rate	31.347	6.131	389	126	0.668	0.196	19.085	43.609
Perinatal mortality rate	49.822	8.982	396	128	0.742	0.180	31.858	67.785
Received ANC from a skilled provider	0.768	0.117	129	42	3.043	0.153	0.534	1.000
4+ ANC visits	0.381	0.086	129	42	1.983	0.226	0.209	0.553
8+ ANC visits	0.044	0.033	129	42	1.818	0.752	0.000	0.111
Took any iron-containing supplements	0.487	0.054	129	42	1.213	0.110	0.379	0.594
Mothers protected against tetanus for last birth	0.499	0.082	129	42	1.841	0.165	0.334	0.663
Delivered in a health facility (live births)	0.471	0.109	141	46	2.453	0.232	0.253	0.690
Delivered by a skilled provider (live births)	0.488	0.113	141	46	2.506	0.230	0.263	0.714
Delivered by C-section (live births)	0.081	0.027	141	46	1.168	0.328	0.028	0.134
Women with postnatal check during first 2 days	0.327	0.075	129	42	1.798	0.230	0.177	0.478
Newborns with postnatal check during first 2 days	0.310	0.069	129	42	1.685	0.224	0.171	0.449
Any problem accessing health care	0.587	0.043	516	168	1.966	0.073	0.502	0.673
Ever had a vaccination card	0.820	0.062	60	22	1.327	0.075	0.696	0.943
Received BCG vaccination	0.891	0.068	60	22	1.792	0.076	0.755	1.000
Received DPT-HepB+Hib vaccination (3 doses)	0.587	0.082	60	22	1.366	0.139	0.423	0.750
Received pneumococcal vaccination (3 doses)	0.640	0.078	60	22	1.347	0.123	0.483	0.797
Received measles vaccination	0.551	0.096	60	22	1.582	0.173	0.360	0.742
Fully vaccinated according to national schedule (12–23 months)	0.127	0.054	60	22	1.344	0.428	0.018	0.235
Received measles/measles-rubella 2 vaccination (24–35 months)	0.466	0.102	66	21	1.590	0.220	0.261	0.670
Fully vaccinated according to national schedule (24–35 months)	0.070	0.038	66	21	1.188	0.543	0.000	0.147
Received HPV vaccination (2 doses) women 15–19	0.000	0.000	137	45	na	na	0.000	0.000
Sought treatment for diarrhoea	0.595	0.061	31	11	0.666	0.102	0.474	0.716
Treated with ORS	0.545	0.079	31	11	0.858	0.146	0.386	0.703
Exclusive breastfeeding	0.173	0.063	42	12	1.066	0.365	0.047	0.299
Minimum dietary diversity (children 6–23 months)	0.130	0.042	79	27	1.114	0.327	0.045	0.215
Child slept under an ITN last night	0.051	0.009	385	126	0.673	0.183	0.032	0.070
Received 3+ doses of SP/Fansidar	0.008	0.008	129	42	0.981	0.983	0.000	0.023
Child has fever in last 2 weeks	0.075	0.016	368	119	1.051	0.215	0.043	0.106
Child had blood taken from finger/heel	0.375	0.171	25	9	1.732	0.457	0.032	0.718
Child took ACT	1.000	0.000	1	-	na	0.000	1.000	1.000
Condom use at last sex (women 15–24)	0.000	0.000	5	2	na	na	0.000	0.000
Mobile phone ownership	0.810	0.022	516	168	1.281	0.027	0.766	0.854

na = not applicable

Table B.12 Sampling errors: Wajir sample, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases: Unweighted (N)	Number of cases: Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
HOUSEHOLDS AND POPULATION								
Primary reliance on clean fuels and technology	0.014	0.004	3,404	850	0.978	0.300	0.006	0.022
Births registered with civil authority	0.529	0.074	465	118	2.439	0.140	0.381	0.677
Ownership of at least one ITN	0.255	0.047	497	126	2.374	0.183	0.161	0.348
Ownership of at least one ITN for every two persons	0.067	0.023	495	125	2.045	0.345	0.021	0.113
At least basic drinking water service	0.560	0.076	3,404	850	3.081	0.136	0.408	0.712
At least basic sanitation service	0.343	0.053	3,404	850	2.229	0.155	0.237	0.450
Using open defecation	0.387	0.063	3,404	850	2.621	0.163	0.261	0.513
Using a handwashing facility with soap and water	0.152	0.020	3,404	850	1.089	0.132	0.112	0.192
WOMEN								
No education	0.610	0.038	518	130	1.778	0.063	0.534	0.686
Secondary education or higher	0.336	0.041	518	130	1.957	0.121	0.254	0.417
Literacy	0.376	0.039	518	130	1.833	0.104	0.297	0.454
Use of the internet in last 12 months	0.403	0.046	518	130	2.138	0.115	0.310	0.495
Total fertility rate (3 years)	5.753	0.389	1,421	358	1.250	0.068	4.975	6.532
Currently pregnant	0.065	0.013	518	130	1.176	0.196	0.039	0.090
Mean number of children ever born to women age 40–49	7.460	0.283	94	23	1.147	0.038	6.895	8.026
Median birth interval	26.737	0.763	334	85	1.618	0.029	25.210	28.263
Ideal number of children	8.094	0.092	517	130	0.902	0.011	7.909	8.278
Total wanted fertility rate (3 years)	5.599	0.374	1,421	358	1.139	0.067	4.851	6.346
Currently using any contraceptive method	0.008	0.006	288	72	1.102	0.748	0.000	0.019
Currently using any modern method	0.008	0.006	288	72	1.102	0.748	0.000	0.019
Currently using pill	0.000	0.000	288	72	na	na	0.000	0.000
Currently using injectables	0.005	0.005	288	72	1.194	0.986	0.000	0.015
Currently using implants	0.000	0.000	288	72	na	na	0.000	0.000
Currently using male condoms	0.000	0.000	288	72	na	na	0.000	0.000
Currently using any traditional method	0.000	0.000	288	72	na	na	0.000	0.000
Unmet need for spacing	0.158	0.030	288	72	1.392	0.190	0.098	0.218
Unmet need for limiting	0.119	0.019	288	72	1.003	0.161	0.081	0.157
Unmet need total	0.277	0.025	288	72	0.947	0.090	0.227	0.327
Demand satisfied by modern methods (married women)	0.026	0.020	80	20	1.106	0.743	0.000	0.066
Demand satisfied by modern methods (all women)	0.026	0.020	80	20	1.103	0.743	0.000	0.066
Not exposed to any of the ten media sources	0.153	0.026	518	130	1.608	0.166	0.102	0.204
Neonatal mortality (last 0–9 years)	8.082	2.992	871	221	0.922	0.370	2.097	14.067
Post-neonatal mortality (last 0–9 years)	5.502	2.416	867	220	0.901	0.439	0.671	10.334
Infant mortality (last 0–9 years)	13.585	3.990	871	221	0.935	0.294	5.604	21.565
Child mortality (last 0–9 years)	4.436	2.811	895	225	1.258	0.634	0.000	10.057
Under-five mortality (last 0–9 years)	17.960	5.020	871	221	1.036	0.280	7.920	28.000
Stillbirth rate	49.888	20.797	397	102	1.579	0.417	8.294	91.483
Early neonatal rate	5.786	3.412	380	97	0.889	0.590	0.000	12.610
Perinatal mortality rate	55.406	20.702	397	102	1.529	0.374	14.002	96.811
Received ANC from a skilled provider	0.817	0.066	137	36	1.983	0.081	0.684	0.950
4+ ANC visits	0.596	0.060	137	36	1.431	0.101	0.475	0.717
8+ ANC visits	0.000	0.000	137	36	na	na	0.000	0.000
Took any iron-containing supplements	0.673	0.066	137	36	1.629	0.098	0.541	0.805
Mothers protected against tetanus for last birth	0.688	0.043	137	36	1.079	0.062	0.603	0.774
Delivered in a health facility (live births)	0.522	0.101	149	40	2.325	0.193	0.320	0.723
Delivered by a skilled provider (live births)	0.556	0.102	149	40	2.371	0.183	0.352	0.760
Delivered by C-section (live births)	0.047	0.026	149	40	1.197	0.546	0.000	0.099
Women with postnatal check during first 2 days	0.512	0.070	137	36	1.631	0.137	0.371	0.652
Newborns with postnatal check during first 2 days	0.629	0.078	137	36	1.866	0.124	0.473	0.785
Any problem accessing health care	0.779	0.059	518	130	3.204	0.076	0.661	0.897
Ever had a vaccination card	0.868	0.069	72	19	1.765	0.079	0.731	1.000
Received BCG vaccination	0.927	0.038	72	19	1.267	0.041	0.852	1.000
Received DPT-HepB+Hib vaccination (3 doses)	0.664	0.058	72	19	1.070	0.088	0.548	0.781
Received pneumococcal vaccination (3 doses)	0.766	0.061	72	19	1.246	0.079	0.645	0.887
Received measles vaccination	0.824	0.050	72	19	1.141	0.061	0.724	0.924
Fully vaccinated according to national schedule (12–23 months)	0.330	0.093	72	19	1.666	0.282	0.144	0.517
Received measles/measles-rubella 2 vaccination (24–35 months)	0.393	0.062	74	17	1.040	0.157	0.270	0.516
Fully vaccinated according to national schedule (24–35 months)	0.051	0.039	74	17	1.478	0.777	0.000	0.129
Received HPV vaccination (2 doses) women 15–19	0.019	0.010	157	38	0.962	0.557	0.000	0.040
Sought treatment for diarrhoea	0.458	0.282	8	3	1.899	0.615	0.000	1.000
Treated with ORS	0.458	0.282	8	3	1.899	0.615	0.000	1.000
Exclusive breastfeeding	0.414	0.073	30	7	0.807	0.177	0.267	0.561
Minimum dietary diversity (children 6–23 months)	0.328	0.045	105	28	0.977	0.137	0.238	0.418
Child slept under an ITN last night	0.154	0.032	425	107	1.349	0.207	0.090	0.218
Received 3+ doses of SP/Fansidar	0.000	0.000	137	36	na	na	0.000	0.000
Child has fever in last 2 weeks	0.142	0.028	374	96	1.552	0.201	0.085	0.198
Child had blood taken from finger/heel	0.137	0.053	55	14	1.143	0.389	0.030	0.243
Child took ACT	na	na	-	-	na	na	0.000	na
Condom use at last sex (women 15–24)	na	na	-	na	na	na	0.000	na
Mobile phone ownership	0.816	0.019	518	130	1.099	0.023	0.779	0.854

na = not applicable

Table B.13 Sampling errors: Mandera sample, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases: Unweighted (N)	Number of cases: Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
HOUSEHOLDS AND POPULATION								
Primary reliance on clean fuels and technology	0.019	0.005	3,112	812	1.065	0.260	0.009	0.029
Births registered with civil authority	0.442	0.057	538	135	1.964	0.129	0.329	0.556
Ownership of at least one ITN	0.175	0.029	520	137	1.762	0.168	0.116	0.233
Ownership of at least one ITN for every two persons	0.048	0.017	516	136	1.804	0.356	0.014	0.082
At least basic drinking water service	0.641	0.074	3,112	812	3.127	0.115	0.493	0.788
At least basic sanitation service	0.346	0.052	3,112	812	2.129	0.150	0.242	0.449
Using open defecation	0.265	0.058	3,112	812	2.691	0.218	0.149	0.380
Using a handwashing facility with soap and water	0.544	0.054	3,083	802	2.169	0.099	0.437	0.652
WOMEN								
No education	0.624	0.040	484	123	1.809	0.064	0.545	0.704
Secondary education or higher	0.238	0.037	484	123	1.895	0.155	0.165	0.312
Literacy	0.338	0.036	484	123	1.647	0.105	0.267	0.409
Use of the internet in last 12 months	0.306	0.035	484	123	1.646	0.113	0.236	0.375
Total fertility rate (3 years)	6.787	0.656	1,327	339	1.594	0.097	5.474	8.099
Currently pregnant	0.089	0.012	484	123	0.899	0.131	0.065	0.112
Mean number of children ever born to women age 40–49	7.676	0.363	68	17	1.309	0.047	6.950	8.402
Median birth interval	24.830	0.592	453	109	0.987	0.024	23.647	26.014
Ideal number of children	8.292	0.172	480	122	1.392	0.021	7.947	8.637
Total wanted fertility rate (3 years)	6.676	0.650	1,327	339	1.634	0.097	5.377	7.976
Currently using any contraceptive method	0.003	0.003	307	77	0.767	0.758	0.000	0.008
Currently using any modern method	0.003	0.003	307	77	0.767	0.758	0.000	0.008
Currently using pill	0.000	0.000	307	77	na	na	0.000	0.000
Currently using injectables	0.002	0.002	307	77	0.762	1.036	0.000	0.005
Currently using implants	0.002	0.002	307	77	0.719	1.035	0.000	0.005
Currently using male condoms	0.000	0.000	307	77	na	na	0.000	0.000
Currently using any traditional method	0.000	0.000	307	77	na	na	0.000	0.000
Unmet need for spacing	0.244	0.027	307	77	1.088	0.110	0.190	0.297
Unmet need for limiting	0.004	0.004	307	77	1.077	1.036	0.000	0.011
Unmet need total	0.247	0.026	307	77	1.066	0.106	0.195	0.300
Demand satisfied by modern methods (married women)	0.013	0.010	78	19	0.785	0.772	0.000	0.034
Demand satisfied by modern methods (all women)	0.013	0.010	82	20	0.775	0.767	0.000	0.032
Not exposed to any of the ten media sources	0.559	0.036	484	123	1.577	0.064	0.487	0.630
Neonatal mortality (last 0–9 years)	6.313	2.759	1,067	262	0.959	0.437	0.796	11.831
Post-neonatal mortality (last 0–9 years)	12.704	3.282	1,060	260	0.967	0.258	6.140	19.267
Infant mortality (last 0–9 years)	19.017	4.958	1,067	262	1.120	0.261	9.101	28.933
Child mortality (last 0–9 years)	3.847	1.866	1,075	264	0.928	0.485	0.115	7.579
Under-five mortality (last 0–9 years)	22.791	5.279	1,067	262	1.153	0.232	12.232	33.349
Stillbirth rate	5.864	3.534	504	122	1.017	0.603	0.000	12.932
Early neonatal rate	9.105	5.541	501	121	1.033	0.609	0.000	20.187
Perinatal mortality rate	14.946	5.958	504	122	0.941	0.399	3.030	26.861
Received ANC from a skilled provider	0.788	0.044	160	39	1.349	0.056	0.700	0.875
4+ ANC visits	0.220	0.032	160	39	0.979	0.146	0.156	0.285
8+ ANC visits	0.000	0.000	160	39	na	na	0.000	0.000
Took any iron-containing supplements	0.649	0.042	160	39	1.104	0.064	0.565	0.733
Mothers protected against tetanus for last birth	0.654	0.060	160	39	1.592	0.092	0.533	0.775
Delivered in a health facility (live births)	0.587	0.064	172	42	1.572	0.109	0.459	0.715
Delivered by a skilled provider (live births)	0.593	0.067	172	42	1.668	0.112	0.460	0.727
Delivered by C-section (live births)	0.019	0.010	172	42	0.942	0.531	0.000	0.038
Women with postnatal check during first 2 days	0.406	0.063	160	39	1.600	0.154	0.280	0.531
Newborns with postnatal check during first 2 days	0.421	0.040	160	39	1.029	0.096	0.340	0.501
Any problem accessing health care	0.568	0.038	484	123	1.685	0.067	0.492	0.644
Ever had a vaccination card	0.837	0.043	84	20	1.044	0.051	0.751	0.923
Received BCG vaccination	0.804	0.034	84	20	0.765	0.042	0.736	0.872
Received DPT-HepB+Hib vaccination (3 doses)	0.674	0.039	84	20	0.741	0.058	0.597	0.752
Received pneumococcal vaccination (3 doses)	0.703	0.043	84	20	0.834	0.061	0.618	0.789
Received measles vaccination	0.582	0.057	84	20	1.009	0.097	0.469	0.695
Fully vaccinated according to national schedule (12–23 months)	0.171	0.042	84	20	1.009	0.248	0.086	0.256
Received measles/measles-rubella 2 vaccination (24–35 months)	0.094	0.030	103	26	1.037	0.318	0.034	0.154
Fully vaccinated according to national schedule (24–35 months)	0.015	0.012	103	26	0.974	0.780	0.000	0.038
Received HPV vaccination (2 doses) women 15–19	0.000	0.000	123	31	na	na	0.000	0.000
Sought treatment for diarrhoea	0.691	0.074	20	6	0.773	0.108	0.542	0.840
Treated with ORS	0.356	0.117	20	6	1.173	0.328	0.122	0.591
Exclusive breastfeeding	0.248	0.060	44	11	0.918	0.244	0.127	0.368
Minimum dietary diversity (children 6–23 months)	0.082	0.022	111	27	0.858	0.273	0.038	0.127
Child slept under an ITN last night	0.108	0.012	536	134	0.644	0.107	0.085	0.131
Received 3+ doses of SP/Fansidar	0.004	0.005	160	39	0.856	1.009	0.000	0.014
Child has fever in last 2 weeks	0.145	0.035	487	118	1.808	0.242	0.075	0.215
Child had blood taken from finger/heel	0.250	0.063	69	17	1.072	0.253	0.123	0.376
Child took ACT	0.252	0.169	7	2	0.985	0.671	0.000	0.590
Condom use at last sex (women 15–24)	na	na	-	na	na	na	0.000	na
Mobile phone ownership	0.790	0.025	484	123	1.344	0.032	0.740	0.840

na = not applicable

Table B.14 Sampling errors: Marsabit sample, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases: Unweighted (N)	Number of cases: Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
HOUSEHOLDS AND POPULATION								
Primary reliance on clean fuels and technology	0.064	0.022	2,363	748	1.847	0.347	0.019	0.108
Births registered with civil authority	0.713	0.050	331	105	1.673	0.070	0.613	0.814
Ownership of at least one ITN	0.317	0.045	437	138	2.006	0.141	0.227	0.407
Ownership of at least one ITN for every two persons	0.084	0.020	436	137	1.537	0.244	0.043	0.125
At least basic drinking water service	0.442	0.082	2,363	748	3.166	0.185	0.279	0.605
At least basic sanitation service	0.283	0.050	2,363	748	2.120	0.176	0.183	0.383
Using open defecation	0.182	0.073	2,363	748	3.598	0.400	0.037	0.328
Using a handwashing facility with soap and water	0.411	0.034	2,355	744	1.345	0.083	0.342	0.479
WOMEN								
No education	0.380	0.036	415	141	1.524	0.096	0.308	0.453
Secondary education or higher	0.432	0.037	415	141	1.525	0.086	0.357	0.506
Literacy	0.582	0.039	416	142	1.608	0.067	0.504	0.660
Use of the internet in last 12 months	0.377	0.034	416	142	1.418	0.090	0.309	0.444
Total fertility rate (3 years)	4.923	0.391	1,126	384	1.383	0.079	4.141	5.706
Currently pregnant	0.066	0.012	416	142	0.997	0.183	0.042	0.091
Mean number of children ever born to women age 40–49	5.617	0.288	59	19	1.198	0.051	5.042	6.192
Median birth interval	37.882	2.452	248	83	1.769	0.065	32.979	42.786
Ideal number of children	5.991	0.182	416	142	1.689	0.030	5.627	6.355
Total wanted fertility rate (3 years)	4.746	0.366	1,126	384	1.307	0.077	4.015	5.477
Currently using any contraceptive method	0.350	0.056	226	77	1.741	0.159	0.238	0.461
Currently using any modern method	0.304	0.053	226	77	1.717	0.174	0.198	0.410
Currently using pill	0.020	0.012	226	77	1.282	0.600	0.000	0.044
Currently using injectables	0.103	0.029	226	77	1.434	0.283	0.045	0.161
Currently using implants	0.081	0.029	226	77	1.578	0.355	0.023	0.139
Currently using male condoms	0.004	0.004	226	77	0.967	0.996	0.000	0.012
Currently using any traditional method	0.046	0.015	226	77	1.104	0.337	0.015	0.076
Unmet need for spacing	0.178	0.027	226	77	1.077	0.154	0.123	0.233
Unmet need for limiting	0.057	0.015	226	77	0.948	0.257	0.028	0.086
Unmet need total	0.235	0.034	226	77	1.210	0.146	0.166	0.303
Demand satisfied by modern methods (married women)	0.520	0.068	132	45	1.561	0.131	0.384	0.657
Demand satisfied by modern methods (all women)	0.554	0.056	146	50	1.353	0.100	0.442	0.665
Not exposed to any of the ten media sources	0.205	0.033	416	142	1.658	0.161	0.139	0.271
Neonatal mortality (last 0–9 years)	13.523	6.035	587	201	1.272	0.446	1.454	25.593
Post-neonatal mortality (last 0–9 years)	6.141	4.608	588	201	1.177	0.750	0.000	15.357
Infant mortality (last 0–9 years)	19.664	7.724	587	201	1.171	0.393	4.215	35.113
Child mortality (last 0–9 years)	3.303	2.264	584	199	0.921	0.685	0.000	7.830
Under-five mortality (last 0–9 years)	22.902	7.519	587	201	1.070	0.328	7.865	37.939
Stillbirth rate	8.281	4.694	296	100	0.892	0.567	0.000	17.670
Early neonatal rate	10.385	6.367	293	99	1.073	0.613	0.000	23.119
Perinatal mortality rate	18.580	7.263	296	100	0.930	0.391	4.054	33.107
Received ANC from a skilled provider	0.932	0.044	102	35	1.728	0.047	0.845	1.000
4+ ANC visits	0.772	0.061	102	35	1.457	0.079	0.650	0.894
8+ ANC visits	0.022	0.014	102	35	0.992	0.658	0.000	0.051
Took any iron-containing supplements	0.899	0.042	102	35	1.397	0.047	0.815	0.983
Mothers protected against tetanus for last birth	0.691	0.056	102	35	1.221	0.081	0.579	0.804
Delivered in a health facility (live births)	0.819	0.065	103	35	1.647	0.080	0.688	0.949
Delivered by a skilled provider (live births)	0.859	0.057	103	35	1.649	0.066	0.745	0.973
Delivered by C-section (live births)	0.106	0.031	103	35	1.025	0.294	0.044	0.169
Women with postnatal check during first 2 days	0.834	0.045	102	35	1.221	0.054	0.743	0.925
Newborns with postnatal check during first 2 days	0.844	0.038	102	35	1.050	0.045	0.768	0.920
Any problem accessing health care	0.744	0.044	416	142	2.035	0.059	0.657	0.832
Ever had a vaccination card	1.000	0.000	48	16	na	0.000	1.000	1.000
Received BCG vaccination	1.000	0.000	48	16	na	0.000	1.000	1.000
Received DPT-HepB+Hib vaccination (3 doses)	0.866	0.050	48	16	0.999	0.057	0.767	0.966
Received pneumococcal vaccination (3 doses)	0.916	0.045	48	16	1.121	0.050	0.825	1.000
Received measles vaccination	0.903	0.049	48	16	1.136	0.054	0.805	1.000
Fully vaccinated according to national schedule (12–23 months)	0.287	0.067	48	16	1.012	0.234	0.152	0.421
Received measles/measles-rubella 2 vaccination (24–35 months)	0.698	0.067	60	20	1.107	0.096	0.564	0.831
Fully vaccinated according to national schedule (24–35 months)	0.243	0.061	60	20	1.089	0.251	0.121	0.366
Received HPV vaccination (2 doses) women 15–19	0.251	0.040	114	40	0.972	0.158	0.172	0.330
Sought treatment for diarrhoea	0.467	0.072	26	9	0.747	0.155	0.322	0.611
Treated with ORS	0.493	0.102	26	9	1.048	0.206	0.289	0.696
Exclusive breastfeeding	0.782	0.098	22	8	1.087	0.126	0.585	0.978
Minimum dietary diversity (children 6–23 months)	0.276	0.058	79	27	1.146	0.210	0.160	0.393
Child slept under an ITN last night	0.123	0.029	318	102	1.316	0.235	0.065	0.180
Received 3+ doses of SP/Fansidar	0.030	0.016	102	35	0.930	0.523	0.000	0.062
Child has fever in last 2 weeks	0.089	0.019	288	97	1.118	0.211	0.052	0.127
Child had blood taken from finger/heel	0.328	0.096	23	9	1.028	0.291	0.137	0.519
Child took ACT	0.326	0.160	7	3	0.957	0.492	0.005	0.647
Condom use at last sex (women 15–24)	0.385	0.119	8	3	0.673	0.310	0.147	0.624
Mobile phone ownership	0.762	0.025	416	142	1.181	0.032	0.712	0.811

na = not applicable

Table B.15 Sampling errors: Isiolo sample, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases: Unweighted (N)	Number of cases: Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
HOUSEHOLDS AND POPULATION								
Primary reliance on clean fuels and technology	0.087	0.029	2,255	377	2.537	0.338	0.028	0.146
Births registered with civil authority	0.826	0.032	299	46	1.216	0.039	0.762	0.890
Ownership of at least one ITN	0.298	0.030	480	87	1.412	0.099	0.239	0.358
Ownership of at least one ITN for every two persons	0.128	0.025	480	87	1.623	0.194	0.079	0.178
At least basic drinking water service	0.805	0.050	2,255	377	2.264	0.062	0.705	0.904
At least basic sanitation service	0.256	0.041	2,255	377	1.744	0.161	0.173	0.338
Using open defecation	0.133	0.032	2,255	377	1.708	0.238	0.070	0.196
Using a handwashing facility with soap and water	0.343	0.058	2,255	377	2.299	0.169	0.227	0.459
WOMEN								
No education	0.183	0.039	428	75	2.092	0.215	0.105	0.262
Secondary education or higher	0.431	0.047	428	75	1.969	0.110	0.337	0.526
Literacy	0.798	0.041	431	76	2.128	0.052	0.715	0.881
Use of the internet in last 12 months	0.444	0.035	431	76	1.470	0.079	0.374	0.515
Total fertility rate (3 years)	3.586	0.418	1,195	212	1.238	0.117	2.749	4.422
Currently pregnant	0.050	0.011	431	76	1.072	0.226	0.027	0.072
Mean number of children ever born to women age 40–49	4.884	0.250	74	14	1.004	0.051	4.384	5.384
Median birth interval	41.267	3.091	243	36	1.604	0.075	35.086	47.448
Ideal number of children	5.019	0.244	431	76	1.995	0.049	4.531	5.507
Total wanted fertility rate (3 years)	3.232	0.378	1,195	212	1.097	0.117	2.476	3.988
Currently using any contraceptive method	0.349	0.039	234	36	1.249	0.112	0.271	0.427
Currently using any modern method	0.323	0.036	234	36	1.169	0.111	0.251	0.395
Currently using pill	0.029	0.012	234	36	1.101	0.421	0.005	0.053
Currently using injectables	0.126	0.017	234	36	0.801	0.138	0.091	0.161
Currently using implants	0.149	0.028	234	36	1.185	0.186	0.094	0.204
Currently using male condoms	0.014	0.010	234	36	1.254	0.679	0.000	0.034
Currently using any traditional method	0.026	0.012	234	36	1.194	0.481	0.001	0.051
Unmet need for spacing	0.120	0.030	234	36	1.424	0.253	0.059	0.181
Unmet need for limiting	0.122	0.030	234	36	1.408	0.248	0.062	0.183
Unmet need total	0.242	0.045	234	36	1.586	0.184	0.153	0.332
Demand satisfied by modern methods (married women)	0.547	0.061	123	21	1.430	0.111	0.425	0.668
Demand satisfied by modern methods (all women)	0.612	0.055	149	28	1.413	0.089	0.503	0.721
Not exposed to any of the ten media sources	0.194	0.035	431	76	1.838	0.181	0.124	0.265
Neonatal mortality (last 0–9 years)	13.199	5.615	620	97	1.148	0.425	1.970	24.429
Post-neonatal mortality (last 0–9 years)	0.684	0.692	618	96	0.884	1.011	0.000	2.068
Infant mortality (last 0–9 years)	13.884	5.602	621	97	1.119	0.404	2.679	25.088
Child mortality (last 0–9 years)	7.748	5.323	624	97	1.421	0.687	0.000	18.394
Under-five mortality (last 0–9 years)	21.524	6.537	621	97	1.061	0.304	8.451	34.597
Stillbirth rate	10.564	8.853	293	44	1.374	0.838	0.000	28.271
Early neonatal rate	12.587	7.381	291	43	1.042	0.586	0.000	27.350
Perinatal mortality rate	23.018	10.478	293	44	1.114	0.455	2.061	43.975
Received ANC from a skilled provider	1.000	0.000	109	16	na	0.000	1.000	1.000
4+ ANC visits	0.644	0.068	109	16	1.474	0.106	0.508	0.781
8+ ANC visits	0.055	0.022	109	16	1.013	0.402	0.011	0.100
Took any iron-containing supplements	0.833	0.035	109	16	0.980	0.042	0.763	0.904
Mothers protected against tetanus for last birth	0.589	0.039	109	16	0.827	0.066	0.510	0.667
Delivered in a health facility (live births)	0.892	0.042	113	17	1.303	0.047	0.808	0.976
Delivered by a skilled provider (live births)	0.895	0.040	113	17	1.251	0.045	0.816	0.975
Delivered by C-section (live births)	0.198	0.052	113	17	1.206	0.260	0.095	0.302
Women with postnatal check during first 2 days	0.845	0.044	109	16	1.260	0.052	0.757	0.933
Newborns with postnatal check during first 2 days	0.845	0.044	109	16	1.260	0.052	0.757	0.933
Any problem accessing health care	0.596	0.044	431	76	1.846	0.073	0.508	0.684
Ever had a vaccination card	1.000	0.000	54	8	na	0.000	1.000	1.000
Received BCG vaccination	0.994	0.006	54	8	0.528	0.006	0.983	1.000
Received DPT-HepB+Hib vaccination (3 doses)	0.821	0.060	54	8	1.040	0.073	0.701	0.942
Received pneumococcal vaccination (3 doses)	0.901	0.054	54	8	1.205	0.060	0.793	1.000
Received measles vaccination	0.674	0.075	54	8	1.016	0.111	0.524	0.824
Fully vaccinated according to national schedule (12–23 months)	0.373	0.112	54	8	1.514	0.299	0.150	0.597
Received measles/measles-rubella 2 vaccination (24–35 months)	0.508	0.116	41	6	1.379	0.229	0.275	0.741
Fully vaccinated according to national schedule (24–35 months)	0.234	0.075	41	6	1.054	0.322	0.083	0.384
Received HPV vaccination (2 doses) women 15–19	0.089	0.034	100	18	1.178	0.379	0.022	0.157
Sought treatment for diarrhoea	0.681	0.112	16	3	1.024	0.164	0.458	0.904
Treated with ORS	0.734	0.096	16	3	0.931	0.130	0.543	0.925
Exclusive breastfeeding	0.914	0.048	24	4	0.821	0.052	0.819	1.000
Minimum dietary diversity (children 6–23 months)	0.492	0.069	82	12	1.232	0.140	0.354	0.629
Child slept under an ITN last night	0.309	0.035	300	47	1.087	0.112	0.240	0.378
Received 3+ doses of SP/Fansidar	0.017	0.013	109	16	1.041	0.757	0.000	0.043
Child has fever in last 2 weeks	0.051	0.016	284	42	1.149	0.319	0.018	0.083
Child had blood taken from finger/heel	0.448	0.144	13	2	1.020	0.322	0.159	0.736
Child took ACT	0.000	0.000	2	-	na	na	0.000	0.000
Condom use at last sex (women 15–24)	0.649	0.139	19	5	1.223	0.215	0.370	0.928
Mobile phone ownership	0.794	0.026	431	76	1.310	0.032	0.743	0.845

na = not applicable

Table B.16 Sampling errors: Meru sample, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases: Unweighted (N)	Number of cases: Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
HOUSEHOLDS AND POPULATION								
Primary reliance on clean fuels and technology	0.120	0.028	2,195	2,693	2.194	0.230	0.065	0.175
Births registered with civil authority	0.607	0.067	220	270	1.884	0.110	0.473	0.741
Ownership of at least one ITN	0.241	0.025	703	840	1.561	0.105	0.190	0.291
Ownership of at least one ITN for every two persons	0.131	0.021	695	832	1.610	0.157	0.090	0.172
At least basic drinking water service	0.822	0.043	2,195	2,693	2.424	0.052	0.736	0.908
At least basic sanitation service	0.542	0.052	2,195	2,693	2.392	0.097	0.437	0.647
Using open defecation	0.052	0.044	2,195	2,693	4.348	0.846	0.000	0.140
Using a handwashing facility with soap and water	0.409	0.060	2,085	2,560	2.691	0.146	0.289	0.528
WOMEN								
No education	0.044	0.024	451	526	2.474	0.549	0.000	0.092
Secondary education or higher	0.535	0.033	451	526	1.402	0.062	0.469	0.601
Literacy	0.904	0.028	454	530	2.001	0.031	0.849	0.960
Use of the internet in last 12 months	0.473	0.041	454	530	1.736	0.086	0.391	0.554
Total fertility rate (3 years)	2.788	0.329	1,270	1,485	1.428	0.118	2.130	3.445
Currently pregnant	0.041	0.008	454	530	0.852	0.192	0.026	0.057
Mean number of children ever born to women age 40–49	3.613	0.251	112	137	1.487	0.070	3.110	4.115
Median birth interval	56.262	4.492	130	163	1.778	0.080	47.278	65.245
Ideal number of children	3.148	0.140	453	529	2.141	0.044	2.868	3.428
Total wanted fertility rate (3 years)	2.638	0.321	1,270	1,485	1.437	0.122	1.996	3.279
Currently using any contraceptive method	0.764	0.034	256	308	1.279	0.045	0.696	0.833
Currently using any modern method	0.715	0.035	256	308	1.227	0.049	0.645	0.784
Currently using pill	0.095	0.019	256	308	1.059	0.204	0.056	0.134
Currently using injectables	0.292	0.029	256	308	1.031	0.100	0.234	0.351
Currently using implants	0.246	0.033	256	308	1.222	0.134	0.180	0.312
Currently using male condoms	0.000	0.000	256	308	na	na	0.000	0.000
Currently using any traditional method	0.050	0.012	256	308	0.873	0.239	0.026	0.074
Unmet need for spacing	0.028	0.011	256	308	1.019	0.374	0.007	0.050
Unmet need for limiting	0.052	0.012	256	308	0.858	0.229	0.028	0.076
Unmet need total	0.080	0.018	256	308	1.084	0.230	0.043	0.117
Demand satisfied by modern methods (married women)	0.846	0.026	216	261	1.053	0.031	0.794	0.898
Demand satisfied by modern methods (all women)	0.860	0.023	265	312	1.073	0.027	0.814	0.905
Not exposed to any of the ten media sources	0.102	0.020	454	530	1.433	0.200	0.061	0.143
Neonatal mortality (last 0–9 years)	7.390	4.967	370	450	1.115	0.672	0.000	17.325
Post-neonatal mortality (last 0–9 years)	2.877	2.897	366	445	1.072	1.007	0.000	8.672
Infant mortality (last 0–9 years)	10.267	5.497	370	450	1.059	0.535	0.000	21.261
Child mortality (last 0–9 years)	9.830	5.024	383	469	1.023	0.511	0.000	19.877
Under-five mortality (last 0–9 years)	19.996	6.666	371	451	0.935	0.333	6.665	33.327
Stillbirth rate	8.142	6.319	189	227	0.986	0.776	0.000	20.779
Early neonatal rate	14.600	9.075	188	226	1.082	0.622	0.000	32.751
Perinatal mortality rate	22.707	10.115	189	227	0.979	0.445	2.477	42.936
Received ANC from a skilled provider	0.975	0.023	62	72	1.151	0.024	0.929	1.000
4+ ANC visits	0.475	0.075	62	72	1.167	0.158	0.325	0.625
8+ ANC visits	0.043	0.027	62	72	1.041	0.630	0.000	0.097
Took any iron-containing supplements	0.846	0.048	62	72	1.029	0.056	0.750	0.941
Mothers protected against tetanus for last birth	0.779	0.090	62	72	1.660	0.115	0.599	0.958
Delivered in a health facility (live births)	0.887	0.090	62	72	2.201	0.102	0.706	1.000
Delivered by a skilled provider (live births)	0.911	0.068	62	72	1.847	0.074	0.776	1.000
Delivered by C-section (live births)	0.248	0.061	62	72	1.090	0.244	0.127	0.370
Women with postnatal check during first 2 days	0.808	0.049	62	72	0.965	0.060	0.711	0.906
Newborns with postnatal check during first 2 days	0.849	0.043	62	72	0.941	0.051	0.763	0.935
Any problem accessing health care	0.747	0.034	454	530	1.660	0.045	0.679	0.815
Ever had a vaccination card	1.000	0.000	28	34	na	0.000	1.000	1.000
Received BCG vaccination	0.964	0.037	28	34	1.069	0.039	0.889	1.000
Received DPT-HepB+Hib vaccination (3 doses)	0.830	0.068	28	34	0.967	0.082	0.694	0.966
Received pneumococcal vaccination (3 doses)	0.804	0.070	28	34	0.948	0.087	0.664	0.945
Received measles vaccination	0.826	0.106	28	34	1.495	0.129	0.613	1.000
Fully vaccinated according to national schedule (12–23 months)	0.467	0.107	28	34	1.150	0.230	0.253	0.682
Received measles/measles-rubella 2 vaccination (24–35 months)	0.701	0.134	37	48	1.838	0.192	0.432	0.970
Fully vaccinated according to national schedule (24–35 months)	0.408	0.110	37	48	1.384	0.270	0.188	0.629
Received HPV vaccination (2 doses) women 15–19	0.158	0.037	86	103	0.941	0.235	0.084	0.233
Sought treatment for diarrhoea	0.529	0.088	37	47	1.081	0.166	0.353	0.705
Treated with ORS	0.352	0.087	37	47	1.088	0.247	0.178	0.526
Exclusive breastfeeding	0.862	0.095	16	20	1.065	0.110	0.672	1.000
Minimum dietary diversity (children 6–23 months)	0.604	0.108	46	52	1.458	0.178	0.388	0.819
Child slept under an ITN last night	0.341	0.052	217	269	1.498	0.153	0.236	0.445
Received 3+ doses of SP/Fansidar	0.000	0.000	62	72	na	na	0.000	0.000
Child has fever in last 2 weeks	0.158	0.023	184	221	0.809	0.147	0.111	0.204
Child had blood taken from finger/heel	0.293	0.099	29	35	1.046	0.339	0.094	0.491
Child took ACT	0.400	0.205	6	7	1.004	0.512	0.000	0.810
Condom use at last sex (women 15–24)	0.431	0.122	20	19	1.069	0.283	0.187	0.675
Mobile phone ownership	0.762	0.041	454	530	2.053	0.054	0.680	0.845

na = not applicable

Table B.17 Sampling errors: Tharaka-Nithi sample, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases: Unweighted (N)	Number of cases: Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
HOUSEHOLDS AND POPULATION								
Primary reliance on clean fuels and technology	0.069	0.018	1,468	704	1.582	0.270	0.032	0.106
Births registered with civil authority	0.873	0.030	142	69	1.162	0.035	0.813	0.934
Ownership of at least one ITN	0.316	0.045	452	209	2.027	0.141	0.227	0.405
Ownership of at least one ITN for every two persons	0.186	0.034	449	208	1.821	0.181	0.119	0.253
At least basic drinking water service	0.548	0.097	1,468	704	3.452	0.177	0.354	0.741
At least basic sanitation service	0.556	0.044	1,468	704	1.595	0.079	0.469	0.644
Using open defecation	0.006	0.006	1,468	704	1.101	0.949	0.000	0.018
Using a handwashing facility with soap and water	0.828	0.026	1,468	704	1.250	0.032	0.775	0.880
WOMEN								
No education	0.012	0.007	282	140	1.037	0.553	0.000	0.026
Secondary education or higher	0.675	0.040	282	140	1.428	0.059	0.595	0.755
Literacy	0.920	0.028	294	147	1.782	0.031	0.864	0.977
Use of the internet in last 12 months	0.588	0.047	294	147	1.623	0.080	0.495	0.682
Total fertility rate (3 years)	2.671	0.323	820	407	1.089	0.121	2.026	3.317
Currently pregnant	0.039	0.009	294	147	0.840	0.244	0.020	0.058
Mean number of children ever born to women age 40–49	3.773	0.332	57	29	1.328	0.088	3.109	4.436
Median birth interval	48.738	3.518	76	41	0.693	0.072	41.703	55.774
Ideal number of children	3.064	0.112	294	147	1.506	0.037	2.840	3.289
Total wanted fertility rate (3 years)	2.240	0.222	820	407	0.805	0.099	1.795	2.684
Currently using any contraceptive method	0.691	0.029	154	80	0.771	0.042	0.634	0.749
Currently using any modern method	0.649	0.031	154	80	0.801	0.048	0.587	0.711
Currently using pill	0.040	0.016	154	80	0.983	0.389	0.009	0.071
Currently using injectables	0.282	0.029	154	80	0.800	0.103	0.224	0.340
Currently using implants	0.180	0.034	154	80	1.098	0.190	0.112	0.248
Currently using male condoms	0.019	0.010	154	80	0.874	0.504	0.000	0.039
Currently using any traditional method	0.043	0.013	154	80	0.807	0.309	0.016	0.069
Unmet need for spacing	0.055	0.021	154	80	1.134	0.379	0.013	0.097
Unmet need for limiting	0.049	0.021	154	80	1.211	0.430	0.007	0.092
Unmet need total	0.105	0.028	154	80	1.143	0.270	0.048	0.161
Demand satisfied by modern methods (married women)	0.815	0.035	122	64	0.993	0.043	0.745	0.885
Demand satisfied by modern methods (all women)	0.810	0.033	167	83	1.066	0.040	0.744	0.875
Not exposed to any of the ten media sources	0.165	0.029	294	147	1.346	0.177	0.107	0.224
Neonatal mortality (last 0–9 years)	30.407	7.446	251	131	0.725	0.245	15.515	45.299
Post-neonatal mortality (last 0–9 years)	5.118	5.141	255	133	1.028	1.004	0.000	15.401
Infant mortality (last 0–9 years)	35.525	8.694	251	131	0.762	0.245	18.138	52.913
Child mortality (last 0–9 years)	10.293	6.092	254	132	0.936	0.592	0.000	22.478
Under-five mortality (last 0–9 years)	45.453	12.786	251	131	0.979	0.281	19.882	71.024
Stillbirth rate	0.000	0.000	124	65	na	na	0.000	0.000
Early neonatal rate	0.000	0.000	124	65	na	na	0.000	0.000
Perinatal mortality rate	0.000	0.000	124	65	na	na	0.000	0.000
Received ANC from a skilled provider	0.969	0.028	42	22	1.047	0.029	0.913	1.000
4+ ANC visits	0.780	0.069	42	22	1.058	0.088	0.643	0.917
8+ ANC visits	0.046	0.037	42	22	1.138	0.814	0.000	0.120
Took any iron-containing supplements	0.894	0.045	42	22	0.935	0.050	0.805	0.984
Mothers protected against tetanus for last birth	0.780	0.058	42	22	0.899	0.074	0.664	0.896
Delivered in a health facility (live births)	0.878	0.058	45	24	1.189	0.066	0.762	0.993
Delivered by a skilled provider (live births)	0.897	0.056	45	24	1.250	0.063	0.785	1.000
Delivered by C-section (live births)	0.337	0.096	45	24	1.202	0.283	0.146	0.528
Women with postnatal check during first 2 days	0.727	0.092	42	22	1.307	0.126	0.544	0.911
Newborns with postnatal check during first 2 days	0.888	0.044	42	22	0.886	0.049	0.800	0.975
Any problem accessing health care	0.747	0.037	294	147	1.443	0.049	0.674	0.821
Ever had a vaccination card	0.967	0.033	19	9	0.778	0.035	0.900	1.000
Received BCG vaccination	0.967	0.033	19	9	0.778	0.035	0.900	1.000
Received DPT-HepB+Hib vaccination (3 doses)	0.778	0.101	19	9	1.012	0.129	0.577	0.979
Received pneumococcal vaccination (3 doses)	0.842	0.099	19	9	1.131	0.117	0.644	1.000
Received measles vaccination	0.896	0.081	19	9	1.104	0.090	0.734	1.000
Fully vaccinated according to national schedule (12–23 months)	0.757	0.102	19	9	0.991	0.134	0.554	0.960
Received measles/measles-rubella 2 vaccination (24–35 months)	0.523	0.154	22	12	1.515	0.295	0.215	0.832
Fully vaccinated according to national schedule (24–35 months)	0.259	0.119	22	12	1.335	0.459	0.021	0.498
Received HPV vaccination (2 doses) women 15–19	0.446	0.085	52	27	1.213	0.190	0.276	0.616
Sought treatment for diarrhoea	0.521	0.123	24	13	1.236	0.235	0.276	0.766
Treated with ORS	0.328	0.096	24	13	0.994	0.293	0.136	0.520
Exclusive breastfeeding	0.545	0.085	14	8	0.628	0.155	0.376	0.714
Minimum dietary diversity (children 6–23 months)	0.750	0.064	28	14	0.770	0.085	0.623	0.877
Child slept under an ITN last night	0.381	0.055	141	69	1.200	0.145	0.270	0.491
Received 3+ doses of SP/Fansidar	0.000	0.000	42	22	na	na	0.000	0.000
Child has fever in last 2 weeks	0.296	0.040	123	65	0.945	0.135	0.216	0.375
Child had blood taken from finger/heel	0.210	0.096	37	19	1.396	0.458	0.018	0.403
Child took ACT	0.500	0.000	2	1	0.000	0.000	0.500	0.500
Condom use at last sex (women 15–24)	0.284	0.065	21	9	0.652	0.229	0.154	0.413
Mobile phone ownership	0.846	0.032	294	147	1.514	0.038	0.783	0.910

na = not applicable

Table B.18 Sampling errors: Embu sample, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases: Unweighted (N)	Number of cases: Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
HOUSEHOLDS AND POPULATION								
Primary reliance on clean fuels and technology	0.260	0.057	1,529	1,035	2.771	0.220	0.145	0.374
Births registered with civil authority	0.958	0.021	128	81	1.155	0.022	0.916	1.000
Ownership of at least one ITN	0.242	0.025	565	385	1.397	0.104	0.192	0.293
Ownership of at least one ITN for every two persons	0.155	0.020	555	378	1.326	0.131	0.114	0.196
At least basic drinking water service	0.722	0.068	1,529	1,035	2.991	0.094	0.586	0.858
At least basic sanitation service	0.490	0.048	1,529	1,035	1.934	0.098	0.394	0.586
Using open defecation	0.000	0.000	1,529	1,035	na	na	0.000	0.000
Using a handwashing facility with soap and water	0.387	0.044	1,502	1,020	1.840	0.115	0.298	0.475
WOMEN								
No education	0.002	0.002	357	236	0.903	0.997	0.000	0.007
Secondary education or higher	0.704	0.035	357	236	1.442	0.050	0.634	0.774
Literacy	0.955	0.012	361	239	1.061	0.012	0.932	0.978
Use of the internet in last 12 months	0.586	0.041	361	239	1.571	0.070	0.505	0.668
Total fertility rate (3 years)	2.229	0.293	1,026	679	1.055	0.132	1.642	2.816
Currently pregnant	0.048	0.010	361	239	0.916	0.216	0.027	0.068
Mean number of children ever born to women age 40–49	3.141	0.224	84	58	1.229	0.071	2.692	3.589
Median birth interval	68.514	1.860	81	52	0.809	0.027	64.795	72.234
Ideal number of children	2.888	0.084	361	239	1.241	0.029	2.720	3.056
Total wanted fertility rate (3 years)	1.939	0.292	1,026	679	1.166	0.151	1.355	2.522
Currently using any contraceptive method	0.734	0.034	193	129	1.054	0.046	0.666	0.801
Currently using any modern method	0.697	0.033	193	129	0.983	0.047	0.632	0.762
Currently using pill	0.153	0.036	193	129	1.367	0.233	0.082	0.224
Currently using injectables	0.264	0.033	193	129	1.025	0.124	0.199	0.329
Currently using implants	0.209	0.034	193	129	1.141	0.160	0.142	0.276
Currently using male condoms	0.010	0.007	193	129	0.981	0.693	0.000	0.025
Currently using any traditional method	0.037	0.017	193	129	1.218	0.452	0.004	0.070
Unmet need for spacing	0.038	0.013	193	129	0.973	0.355	0.011	0.064
Unmet need for limiting	0.028	0.009	193	129	0.786	0.334	0.009	0.047
Unmet need total	0.066	0.016	193	129	0.911	0.248	0.033	0.098
Demand satisfied by modern methods (married women)	0.872	0.027	155	103	1.002	0.031	0.818	0.926
Demand satisfied by modern methods (all women)	0.837	0.024	210	139	0.939	0.029	0.789	0.885
Not exposed to any of the ten media sources	0.056	0.007	361	239	0.591	0.128	0.042	0.071
Neonatal mortality (last 0–9 years)	28.530	12.417	276	178	1.101	0.435	3.696	53.364
Post-neonatal mortality (last 0–9 years)	4.229	4.262	274	176	1.097	1.008	0.000	12.753
Infant mortality (last 0–9 years)	32.759	12.666	276	178	1.073	0.387	7.428	58.091
Child mortality (last 0–9 years)	19.204	19.291	278	181	1.125	1.005	0.000	57.786
Under-five mortality (last 0–9 years)	51.334	19.853	278	180	1.013	0.387	11.627	91.040
Stillbirth rate	0.000	0.000	122	77	na	na	0.000	0.000
Early neonatal rate	16.322	15.512	122	77	1.324	0.950	0.000	47.347
Perinatal mortality rate	16.322	15.512	122	77	1.324	0.950	0.000	47.347
Received ANC from a skilled provider	1.000	0.000	44	27	na	0.000	1.000	1.000
4+ ANC visits	0.659	0.059	44	27	0.814	0.089	0.542	0.777
8+ ANC visits	0.153	0.046	44	27	0.838	0.300	0.061	0.245
Took any iron-containing supplements	0.977	0.022	44	27	0.983	0.023	0.932	1.000
Mothers protected against tetanus for last birth	0.957	0.027	44	27	0.870	0.028	0.904	1.000
Delivered in a health facility (live births)	0.980	0.020	46	29	0.956	0.021	0.939	1.000
Delivered by a skilled provider (live births)	1.000	0.000	46	29	na	0.000	1.000	1.000
Delivered by C-section (live births)	0.430	0.112	46	29	1.472	0.260	0.206	0.654
Women with postnatal check during first 2 days	0.908	0.047	44	27	1.066	0.052	0.814	1.000
Newborns with postnatal check during first 2 days	0.906	0.034	44	27	0.780	0.038	0.837	0.975
Any problem accessing health care	0.552	0.036	361	239	1.358	0.065	0.480	0.623
Ever had a vaccination card	0.915	0.059	23	15	1.005	0.065	0.797	1.000
Received BCG vaccination	0.961	0.039	23	15	0.958	0.041	0.883	1.000
Received DPT-HepB+Hib vaccination (3 doses)	0.961	0.039	23	15	0.958	0.041	0.883	1.000
Received pneumococcal vaccination (3 doses)	0.961	0.039	23	15	0.958	0.041	0.883	1.000
Received measles vaccination	0.919	0.055	23	15	0.958	0.060	0.809	1.000
Fully vaccinated according to national schedule (12–23 months)	0.741	0.102	23	15	1.108	0.138	0.536	0.945
Received measles/measles-rubella 2 vaccination (24–35 months)	0.926	0.051	24	16	0.955	0.055	0.825	1.000
Fully vaccinated according to national schedule (24–35 months)	0.656	0.089	24	16	0.882	0.135	0.478	0.833
Received HPV vaccination (2 doses) women 15–19	0.324	0.068	62	42	1.139	0.211	0.187	0.460
Sought treatment for diarrhoea	0.533	0.140	11	8	0.981	0.263	0.252	0.813
Treated with ORS	0.634	0.132	11	8	0.953	0.208	0.371	0.897
Exclusive breastfeeding	0.794	0.132	12	7	1.073	0.166	0.530	1.000
Minimum dietary diversity (children 6–23 months)	0.753	0.079	28	18	0.947	0.104	0.596	0.910
Child slept under an ITN last night	0.455	0.057	130	83	1.129	0.126	0.340	0.569
Received 3+ doses of SP/Fansidar	0.040	0.038	44	27	1.254	0.949	0.000	0.115
Child has fever in last 2 weeks	0.161	0.047	119	75	1.345	0.288	0.068	0.254
Child had blood taken from finger/heel	0.223	0.084	19	12	0.859	0.375	0.056	0.390
Child took ACT	0.670	0.268	3	2	1.013	0.400	0.135	1.000
Condom use at last sex (women 15–24)	0.365	0.075	27	17	0.798	0.205	0.215	0.515
Mobile phone ownership	0.855	0.023	361	239	1.212	0.026	0.810	0.900

na = not applicable

Table B.19 Sampling errors: Kitui sample, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases: Unweighted (N)	Number of cases: Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
HOUSEHOLDS AND POPULATION								
Primary reliance on clean fuels and technology	0.067	0.017	2,254	2,122	1.738	0.249	0.033	0.100
Births registered with civil authority	0.791	0.046	213	208	1.621	0.059	0.698	0.884
Ownership of at least one ITN	0.305	0.038	597	526	2.028	0.126	0.228	0.382
Ownership of at least one ITN for every two persons	0.176	0.029	589	520	1.831	0.164	0.119	0.234
At least basic drinking water service	0.313	0.068	2,254	2,122	3.175	0.216	0.178	0.448
At least basic sanitation service	0.500	0.063	2,254	2,122	2.641	0.125	0.374	0.625
Using open defecation	0.156	0.062	2,254	2,122	3.303	0.394	0.033	0.279
Using a handwashing facility with soap and water	0.229	0.046	2,254	2,122	2.425	0.200	0.137	0.321
WOMEN								
No education	0.022	0.010	437	380	1.436	0.456	0.002	0.043
Secondary education or higher	0.483	0.042	437	380	1.736	0.086	0.400	0.567
Literacy	0.930	0.020	468	412	1.727	0.022	0.890	0.971
Use of the internet in last 12 months	0.400	0.042	468	412	1.862	0.106	0.315	0.485
Total fertility rate (3 years)	2.767	0.264	1,324	1,165	1.125	0.095	2.240	3.295
Currently pregnant	0.049	0.008	468	412	0.782	0.160	0.033	0.064
Mean number of children ever born to women age 40–49	4.782	0.299	111	99	1.707	0.063	4.183	5.380
Median birth interval	48.404	3.687	128	121	1.042	0.076	41.031	55.777
Ideal number of children	3.333	0.127	468	412	1.793	0.038	3.080	3.587
Total wanted fertility rate (3 years)	2.147	0.289	1,324	1,165	1.226	0.135	1.568	2.726
Currently using any contraceptive method	0.643	0.027	273	251	0.939	0.042	0.589	0.698
Currently using any modern method	0.626	0.026	273	251	0.876	0.041	0.574	0.677
Currently using pill	0.023	0.011	273	251	1.159	0.458	0.002	0.044
Currently using injectables	0.242	0.040	273	251	1.525	0.164	0.162	0.321
Currently using implants	0.289	0.030	273	251	1.080	0.103	0.229	0.348
Currently using male condoms	0.000	0.000	273	251	na	na	0.000	0.000
Currently using any traditional method	0.018	0.007	273	251	0.912	0.412	0.003	0.032
Unmet need for spacing	0.056	0.015	273	251	1.067	0.265	0.026	0.086
Unmet need for limiting	0.103	0.022	273	251	1.184	0.212	0.059	0.146
Unmet need total	0.159	0.025	273	251	1.137	0.159	0.108	0.209
Demand satisfied by modern methods (married women)	0.780	0.026	219	201	0.932	0.033	0.728	0.832
Demand satisfied by modern methods (all women)	0.755	0.028	283	252	1.118	0.038	0.698	0.812
Not exposed to any of the ten media sources	0.056	0.015	468	412	1.407	0.267	0.026	0.086
Neonatal mortality (last 0–9 years)	32.242	8.365	410	377	0.844	0.259	15.512	48.972
Post-neonatal mortality (last 0–9 years)	12.257	4.998	408	375	0.918	0.408	2.260	22.254
Infant mortality (last 0–9 years)	44.499	8.352	410	377	0.761	0.188	27.795	61.204
Child mortality (last 0–9 years)	0.000	0.000	405	373	na	na	0.000	0.000
Under-five mortality (last 0–9 years)	44.499	8.352	410	377	0.761	0.188	27.795	61.204
Stillbirth rate	0.000	0.000	191	181	na	na	0.000	0.000
Early neonatal rate	30.871	11.428	193	182	0.950	0.370	8.016	53.727
Perinatal mortality rate	31.160	11.482	191	181	0.945	0.369	8.195	54.124
Received ANC from a skilled provider	0.986	0.014	73	64	0.988	0.014	0.958	1.000
4+ ANC visits	0.824	0.057	73	64	1.273	0.070	0.710	0.939
8+ ANC visits	0.073	0.035	73	64	1.146	0.482	0.003	0.144
Took any iron-containing supplements	0.925	0.031	73	64	0.995	0.033	0.863	0.987
Mothers protected against tetanus for last birth	0.858	0.050	73	64	1.211	0.058	0.758	0.958
Delivered in a health facility (live births)	0.922	0.041	75	66	1.336	0.045	0.840	1.000
Delivered by a skilled provider (live births)	0.922	0.041	75	66	1.336	0.045	0.840	1.000
Delivered by C-section (live births)	0.193	0.053	75	66	1.143	0.276	0.087	0.300
Women with postnatal check during first 2 days	0.853	0.055	73	64	1.307	0.064	0.743	0.962
Newborns with postnatal check during first 2 days	0.841	0.070	73	64	1.616	0.084	0.700	0.982
Any problem accessing health care	0.754	0.041	468	412	2.072	0.055	0.671	0.837
Ever had a vaccination card	1.000	0.000	44	39	na	0.000	1.000	1.000
Received BCG vaccination	1.000	0.000	44	39	na	0.000	1.000	1.000
Received DPT-HepB+Hib vaccination (3 doses)	0.817	0.079	44	39	1.339	0.097	0.659	0.974
Received pneumococcal vaccination (3 doses)	0.922	0.046	44	39	1.138	0.050	0.830	1.000
Received measles vaccination	0.824	0.082	44	39	1.413	0.100	0.660	0.988
Fully vaccinated according to national schedule (12–23 months)	0.363	0.102	44	39	1.395	0.282	0.159	0.568
Received measles/measles-rubella 2 vaccination (24–35 months)	0.886	0.051	35	33	0.973	0.058	0.784	0.988
Fully vaccinated according to national schedule (24–35 months)	0.421	0.122	35	33	1.453	0.290	0.176	0.665
Received HPV vaccination (2 doses) women 15–19	0.808	0.045	92	82	1.083	0.055	0.718	0.897
Sought treatment for diarrhoea	0.566	0.116	22	19	1.072	0.206	0.333	0.799
Treated with ORS	0.417	0.122	22	19	1.133	0.293	0.173	0.662
Exclusive breastfeeding	0.776	0.082	19	17	0.843	0.106	0.612	0.941
Minimum dietary diversity (children 6–23 months)	0.565	0.094	53	45	1.351	0.166	0.378	0.753
Child slept under an ITN last night	0.308	0.067	212	205	1.989	0.216	0.175	0.441
Received 3+ doses of SP/Fansidar	0.056	0.030	73	64	1.109	0.540	0.000	0.116
Child has fever in last 2 weeks	0.109	0.030	186	175	1.358	0.279	0.048	0.170
Child had blood taken from finger/heel	0.130	0.090	22	19	1.232	0.695	0.000	0.310
Child took ACT	0.546	0.271	4	4	1.112	0.496	0.004	1.000
Condom use at last sex (women 15–24)	0.541	0.123	43	40	1.566	0.226	0.296	0.787
Mobile phone ownership	0.781	0.017	468	412	0.899	0.022	0.746	0.815

na = not applicable

Table B.20 Sampling errors: Machakos sample, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases: Unweighted (N)	Number of cases: Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
HOUSEHOLDS AND POPULATION								
Primary reliance on clean fuels and technology	0.367	0.093	2,075	2,993	4.404	0.252	0.182	0.553
Births registered with civil authority	0.936	0.019	219	354	1.024	0.021	0.897	0.975
Ownership of at least one ITN	0.242	0.020	684	1,021	1.207	0.082	0.202	0.281
Ownership of at least one ITN for every two persons	0.107	0.013	680	1,016	1.085	0.120	0.081	0.133
At least basic drinking water service	0.692	0.065	2,075	2,993	2.927	0.094	0.561	0.822
At least basic sanitation service	0.490	0.036	2,075	2,993	1.604	0.074	0.417	0.563
Using open defecation	0.001	0.001	2,075	2,993	1.001	0.741	0.000	0.002
Using a handwashing facility with soap and water	0.404	0.032	2,035	2,941	1.431	0.079	0.340	0.468
WOMEN								
No education	0.002	0.002	428	652	0.842	1.016	0.000	0.005
Secondary education or higher	0.773	0.045	428	652	2.216	0.058	0.683	0.863
Literacy	1.000	0.000	447	677	na	0.000	1.000	1.000
Use of the internet in last 12 months	0.692	0.043	447	677	1.968	0.062	0.606	0.778
Total fertility rate (3 years)	3.209	0.329	1,277	1,951	1.807	0.103	2.551	3.867
Currently pregnant	0.056	0.011	447	677	1.006	0.195	0.034	0.078
Mean number of children ever born to women age 40–49	3.708	0.331	90	113	1.826	0.089	3.047	4.370
Median birth interval	57.596	3.705	134	212	0.941	0.064	50.186	65.005
Ideal number of children	2.973	0.061	446	670	1.149	0.020	2.852	3.094
Total wanted fertility rate (3 years)	2.759	0.401	1,277	1,951	2.107	0.145	1.956	3.561
Currently using any contraceptive method	0.751	0.035	267	420	1.299	0.046	0.682	0.820
Currently using any modern method	0.681	0.038	267	420	1.330	0.056	0.605	0.757
Currently using pill	0.083	0.028	267	420	1.646	0.338	0.027	0.138
Currently using injectables	0.216	0.026	267	420	1.043	0.122	0.163	0.268
Currently using implants	0.322	0.051	267	420	1.765	0.158	0.221	0.424
Currently using male condoms	0.011	0.006	267	420	0.964	0.572	0.000	0.023
Currently using any traditional method	0.070	0.024	267	420	1.537	0.345	0.022	0.118
Unmet need for spacing	0.061	0.021	267	420	1.411	0.339	0.020	0.103
Unmet need for limiting	0.022	0.009	267	420	1.031	0.420	0.004	0.041
Unmet need total	0.084	0.018	267	420	1.054	0.214	0.048	0.119
Demand satisfied by modern methods (married women)	0.816	0.024	223	350	0.915	0.029	0.769	0.864
Demand satisfied by modern methods (all women)	0.777	0.022	300	472	0.950	0.029	0.733	0.822
Not exposed to any of the ten media sources	0.037	0.011	447	677	1.292	0.314	0.014	0.059
Neonatal mortality (last 0–9 years)	12.568	6.662	385	634	0.977	0.530	0.000	25.892
Post-neonatal mortality (last 0–9 years)	18.405	9.000	386	635	1.103	0.489	0.406	36.404
Infant mortality (last 0–9 years)	30.972	6.617	385	634	0.774	0.214	17.738	44.207
Child mortality (last 0–9 years)	1.896	1.998	369	600	0.886	1.054	0.000	5.892
Under-five mortality (last 0–9 years)	32.810	6.540	385	634	0.747	0.199	19.730	45.890
Stillbirth rate	10.000	6.360	211	349	0.982	0.636	0.000	22.720
Early neonatal rate	10.597	6.407	210	348	0.953	0.605	0.000	23.410
Perinatal mortality rate	20.547	10.671	211	349	1.162	0.519	0.000	41.889
Received ANC from a skilled provider	1.000	0.000	93	156	na	0.000	1.000	1.000
4+ ANC visits	0.726	0.048	93	156	1.026	0.066	0.630	0.821
8+ ANC visits	0.079	0.026	93	156	0.908	0.322	0.028	0.131
Took any iron-containing supplements	0.908	0.021	93	156	0.710	0.023	0.865	0.951
Mothers protected against tetanus for last birth	0.738	0.027	93	156	0.599	0.037	0.683	0.792
Delivered in a health facility (live births)	0.974	0.015	96	159	0.976	0.016	0.943	1.000
Delivered by a skilled provider (live births)	0.989	0.009	96	159	0.858	0.009	0.972	1.000
Delivered by C-section (live births)	0.221	0.023	96	159	0.562	0.104	0.175	0.267
Women with postnatal check during first 2 days	0.890	0.023	93	156	0.722	0.026	0.843	0.937
Newborns with postnatal check during first 2 days	0.927	0.036	93	156	1.327	0.039	0.854	0.999
Any problem accessing health care	0.380	0.033	447	677	1.453	0.088	0.313	0.447
Ever had a vaccination card	1.000	0.000	48	92	na	0.000	1.000	1.000
Received BCG vaccination	0.978	0.019	48	92	1.006	0.020	0.939	1.000
Received DPT-HepB+Hib vaccination (3 doses)	0.863	0.030	48	92	0.666	0.034	0.804	0.923
Received pneumococcal vaccination (3 doses)	0.978	0.019	48	92	1.006	0.020	0.939	1.000
Received measles vaccination	0.879	0.026	48	92	0.623	0.030	0.826	0.931
Fully vaccinated according to national schedule (12–23 months)	0.492	0.089	48	92	1.364	0.180	0.315	0.669
Received measles/measles-rubella 2 vaccination (24–35 months)	0.897	0.062	34	59	1.266	0.069	0.772	1.000
Fully vaccinated according to national schedule (24–35 months)	0.668	0.138	34	59	1.814	0.207	0.391	0.945
Received HPV vaccination (2 doses) women 15–19	0.366	0.064	77	98	1.160	0.175	0.238	0.495
Sought treatment for diarrhoea	0.650	0.041	30	73	0.592	0.063	0.569	0.732
Treated with ORS	0.482	0.041	30	73	0.564	0.085	0.400	0.563
Exclusive breastfeeding	0.781	0.097	26	34	1.169	0.125	0.586	0.976
Minimum dietary diversity (children 6–23 months)	0.476	0.041	63	117	0.656	0.087	0.393	0.559
Child slept under an ITN last night	0.244	0.029	219	355	0.967	0.118	0.186	0.301
Received 3+ doses of SP/Fansidar	0.022	0.015	93	156	0.959	0.673	0.000	0.051
Child has fever in last 2 weeks	0.199	0.026	202	339	0.974	0.131	0.147	0.252
Child had blood taken from finger/heel	0.291	0.096	35	67	1.400	0.331	0.099	0.484
Child took ACT	0.729	0.227	4	12	1.394	0.311	0.276	1.000
Condom use at last sex (women 15–24)	0.242	0.050	37	66	0.707	0.207	0.141	0.342
Mobile phone ownership	0.880	0.021	447	677	1.358	0.024	0.838	0.922

na = not applicable

Table B.21 Sampling errors: Makueni sample, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases: Unweighted (N)	Number of cases: Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
HOUSEHOLDS AND POPULATION								
Primary reliance on clean fuels and technology	0.100	0.030	1,875	1,665	2.672	0.302	0.040	0.161
Births registered with civil authority	0.660	0.040	185	167	1.056	0.061	0.579	0.741
Ownership of at least one ITN	0.353	0.022	582	506	1.100	0.062	0.309	0.397
Ownership of at least one ITN for every two persons	0.212	0.020	581	506	1.168	0.093	0.173	0.252
At least basic drinking water service	0.559	0.066	1,875	1,665	2.675	0.118	0.427	0.691
At least basic sanitation service	0.707	0.049	1,875	1,665	2.352	0.069	0.610	0.805
Using open defecation	0.000	0.000	1,875	1,665	0.946	1.000	0.000	0.001
Using a handwashing facility with soap and water	0.294	0.037	1,875	1,665	1.630	0.125	0.221	0.368
WOMEN								
No education	0.003	0.003	375	336	1.121	1.018	0.000	0.010
Secondary education or higher	0.718	0.045	375	336	1.946	0.063	0.627	0.809
Literacy	0.991	0.005	389	350	0.998	0.005	0.981	1.000
Use of the internet in last 12 months	0.618	0.056	389	350	2.238	0.090	0.507	0.729
Total fertility rate (3 years)	3.274	0.320	1,093	985	1.046	0.098	2.634	3.913
Currently pregnant	0.038	0.009	389	350	0.971	0.250	0.019	0.056
Mean number of children ever born to women age 40–49	3.735	0.185	88	79	1.046	0.049	3.366	4.105
Median birth interval	53.316	3.595	112	104	1.245	0.067	46.125	60.506
Ideal number of children	2.935	0.054	389	350	0.997	0.018	2.827	3.043
Total wanted fertility rate (3 years)	2.916	0.247	1,093	985	0.917	0.085	2.422	3.409
Currently using any contraceptive method	0.796	0.024	210	192	0.866	0.030	0.747	0.844
Currently using any modern method	0.717	0.019	210	192	0.601	0.026	0.679	0.754
Currently using pill	0.055	0.015	210	192	0.934	0.268	0.026	0.085
Currently using injectables	0.272	0.029	210	192	0.930	0.105	0.214	0.329
Currently using implants	0.274	0.023	210	192	0.737	0.083	0.229	0.319
Currently using male condoms	0.010	0.007	210	192	0.969	0.668	0.000	0.023
Currently using any traditional method	0.079	0.020	210	192	1.061	0.251	0.039	0.118
Unmet need for spacing	0.022	0.009	210	192	0.877	0.402	0.004	0.040
Unmet need for limiting	0.016	0.008	210	192	0.940	0.507	0.000	0.033
Unmet need total	0.038	0.010	210	192	0.764	0.264	0.018	0.059
Demand satisfied by modern methods (married women)	0.859	0.018	175	160	0.686	0.021	0.823	0.895
Demand satisfied by modern methods (all women)	0.801	0.022	253	229	0.880	0.027	0.757	0.845
Not exposed to any of the ten media sources	0.024	0.006	389	350	0.775	0.251	0.012	0.036
Neonatal mortality (last 0–9 years)	40.003	12.846	331	307	1.234	0.321	14.311	65.694
Post-neonatal mortality (last 0–9 years)	2.814	2.840	331	308	1.015	1.009	0.000	8.494
Infant mortality (last 0–9 years)	42.816	13.626	331	307	1.191	0.318	15.565	70.068
Child mortality (last 0–9 years)	1.630	1.647	309	286	na	1.010	0.000	4.923
Under-five mortality (last 0–9 years)	44.376	13.560	331	307	1.191	0.306	17.256	71.497
Stillbirth rate	15.638	8.898	175	161	0.978	0.569	0.000	33.435
Early neonatal rate	20.093	12.976	172	159	1.254	0.646	0.000	46.046
Perinatal mortality rate	35.417	18.473	175	161	1.209	0.522	0.000	72.364
Received ANC from a skilled provider	1.000	0.000	63	59	na	0.000	1.000	1.000
4+ ANC visits	0.891	0.050	63	59	1.246	0.056	0.792	0.990
8+ ANC visits	0.238	0.074	63	59	1.361	0.311	0.090	0.387
Took any iron-containing supplements	0.961	0.022	63	59	0.874	0.022	0.918	1.000
Mothers protected against tetanus for last birth	0.945	0.026	63	59	0.909	0.028	0.892	0.997
Delivered in a health facility (live births)	0.971	0.019	67	62	0.927	0.019	0.933	1.000
Delivered by a skilled provider (live births)	0.985	0.014	67	62	0.964	0.014	0.957	1.000
Delivered by C-section (live births)	0.302	0.070	67	62	1.167	0.231	0.163	0.442
Women with postnatal check during first 2 days	0.956	0.024	63	59	0.932	0.025	0.908	1.000
Newborns with postnatal check during first 2 days	0.981	0.018	63	59	1.028	0.018	0.946	1.000
Any problem accessing health care	0.541	0.051	389	350	2.026	0.095	0.438	0.644
Ever had a vaccination card	1.000	0.000	24	21	na	0.000	1.000	1.000
Received BCG vaccination	1.000	0.000	24	21	na	0.000	1.000	1.000
Received DPT-HepB+Hib vaccination (3 doses)	1.000	0.000	24	21	na	0.000	1.000	1.000
Received pneumococcal vaccination (3 doses)	1.000	0.000	24	21	na	0.000	1.000	1.000
Received measles vaccination	1.000	0.000	24	21	na	0.000	1.000	1.000
Fully vaccinated according to national schedule (12–23 months)	0.688	0.089	24	21	0.923	0.130	0.509	0.867
Received measles/measles-rubella 2 vaccination (24–35 months)	0.716	0.072	40	37	1.015	0.100	0.572	0.860
Fully vaccinated according to national schedule (24–35 months)	0.517	0.062	40	37	0.785	0.119	0.394	0.641
Received HPV vaccination (2 doses) women 15–19	0.341	0.087	72	64	1.530	0.255	0.167	0.515
Sought treatment for diarrhoea	0.454	0.135	18	17	1.124	0.296	0.185	0.724
Treated with ORS	0.324	0.119	18	17	1.079	0.368	0.086	0.563
Exclusive breastfeeding	0.712	0.077	23	23	0.808	0.109	0.557	0.867
Minimum dietary diversity (children 6–23 months)	0.785	0.082	37	33	1.191	0.105	0.620	0.949
Child slept under an ITN last night	0.383	0.025	188	170	0.676	0.065	0.333	0.432
Received 3+ doses of SP/Fansidar	0.016	0.015	63	59	0.942	0.953	0.000	0.045
Child has fever in last 2 weeks	0.129	0.025	164	152	0.942	0.193	0.079	0.179
Child had blood taken from finger/heel	0.181	0.089	22	20	1.075	0.492	0.003	0.359
Child took ACT	0.471	0.254	4	3	0.995	0.538	0.000	0.979
Condom use at last sex (women 15–24)	0.483	0.094	27	24	0.962	0.195	0.295	0.671
Mobile phone ownership	0.847	0.023	389	350	1.241	0.027	0.801	0.892

na = not applicable

Table B.22 Sampling errors: Nyandarua sample, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases Unweighted (N)	Number of cases Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits Lower (R-2SE)	Confidence limits Upper (R+2SE)
HOUSEHOLDS AND POPULATION								
Primary reliance on clean fuels and technology	0.090	0.019	1,306	1,060	1.291	0.211	0.052	0.128
Births registered with civil authority	0.919	0.025	130	99	0.999	0.027	0.870	0.969
Ownership of at least one ITN	0.089	0.012	431	341	0.906	0.140	0.064	0.114
Ownership of at least one ITN for every two persons	0.040	0.007	428	339	0.761	0.180	0.026	0.055
At least basic drinking water service	0.891	0.036	1,306	1,060	2.063	0.040	0.820	0.962
At least basic sanitation service	0.455	0.043	1,306	1,060	1.525	0.095	0.368	0.541
Using open defecation	0.001	0.001	1,306	1,060	1.076	0.999	0.000	0.003
Using a handwashing facility with soap and water	0.316	0.032	1,303	1,058	1.208	0.101	0.253	0.380
WOMEN								
No education	0.000	0.000	259	222	na	na	0.000	0.000
Secondary education or higher	0.693	0.027	259	222	0.954	0.040	0.638	0.747
Literacy	0.987	0.007	260	223	1.034	0.007	0.972	1.000
Use of the internet in last 12 months	0.590	0.035	260	223	1.133	0.059	0.520	0.659
Total fertility rate (3 years)	3.780	0.207	740	636	0.723	0.055	3.366	4.195
Currently pregnant	0.027	0.009	260	223	0.890	0.331	0.009	0.045
Mean number of children ever born to women age 40–49	4.025	0.273	65	60	1.207	0.068	3.479	4.572
Median birth interval	57.306	3.690	80	71	0.905	0.064	49.925	64.686
Ideal number of children	3.531	0.151	258	221	1.430	0.043	3.229	3.833
Total wanted fertility rate (3 years)	3.359	0.255	740	636	0.771	0.076	2.850	3.868
Currently using any contraceptive method	0.686	0.031	139	122	0.796	0.046	0.623	0.749
Currently using any modern method	0.656	0.033	139	122	0.820	0.050	0.590	0.722
Currently using pill	0.113	0.030	139	122	1.114	0.265	0.053	0.174
Currently using injectables	0.190	0.038	139	122	1.140	0.201	0.114	0.266
Currently using implants	0.222	0.035	139	122	1.001	0.160	0.151	0.292
Currently using male condoms	0.033	0.019	139	122	1.225	0.561	0.000	0.071
Currently using any traditional method	0.029	0.017	139	122	1.149	0.562	0.000	0.063
Unmet need for spacing	0.070	0.022	139	122	1.033	0.320	0.025	0.115
Unmet need for limiting	0.082	0.025	139	122	1.060	0.302	0.032	0.131
Unmet need total	0.152	0.027	139	122	0.870	0.175	0.099	0.205
Demand satisfied by modern methods (married women)	0.783	0.035	116	103	0.913	0.045	0.713	0.853
Demand satisfied by modern methods (all women)	0.787	0.030	146	126	0.874	0.038	0.728	0.846
Not exposed to any of the ten media sources	0.094	0.023	260	223	1.273	0.246	0.048	0.140
Neonatal mortality (last 0–9 years)	31.237	15.140	241	205	1.309	0.485	0.958	61.517
Post-neonatal mortality (last 0–9 years)	16.188	7.590	239	204	0.941	0.469	1.008	31.368
Infant mortality (last 0–9 years)	47.425	16.387	241	205	1.091	0.346	14.650	80.200
Child mortality (last 0–9 years)	0.000	0.000	233	201	na	na	0.000	0.000
Under-five mortality (last 0–9 years)	47.425	16.387	241	205	1.091	0.346	14.650	80.200
Stillbirth rate	0.000	0.000	125	104	na	na	0.000	0.000
Early neonatal rate	21.257	14.041	126	105	1.095	0.661	0.000	49.338
Perinatal mortality rate	21.469	14.298	125	104	1.093	0.666	0.000	50.066
Received ANC from a skilled provider	0.980	0.020	58	48	1.085	0.020	0.940	1.000
4+ ANC visits	0.582	0.074	58	48	1.136	0.128	0.434	0.731
8+ ANC visits	0.103	0.051	58	48	1.268	0.498	0.000	0.206
Took any iron-containing supplements	0.938	0.033	58	48	1.044	0.036	0.871	1.000
Mothers protected against tetanus for last birth	0.811	0.056	58	48	1.081	0.069	0.699	0.923
Delivered in a health facility (live births)	1.000	0.000	58	48	na	0.000	1.000	1.000
Delivered by a skilled provider (live births)	1.000	0.000	58	48	na	0.000	1.000	1.000
Delivered by C-section (live births)	0.230	0.053	58	48	0.935	0.229	0.124	0.335
Women with postnatal check during first 2 days	0.777	0.071	58	48	1.286	0.092	0.634	0.919
Newborns with postnatal check during first 2 days	0.959	0.027	58	48	1.042	0.029	0.904	1.000
Any problem accessing health care	0.693	0.043	260	223	1.503	0.062	0.607	0.779
Ever had a vaccination card	0.966	0.032	30	25	0.947	0.033	0.902	1.000
Received BCG vaccination	1.000	0.000	30	25	na	0.000	1.000	1.000
Received DPT-HepB+Hib vaccination (3 doses)	0.884	0.076	30	25	1.272	0.085	0.733	1.000
Received pneumococcal vaccination (3 doses)	1.000	0.000	30	25	na	0.000	1.000	1.000
Received measles vaccination	0.933	0.041	30	25	0.878	0.044	0.852	1.000
Fully vaccinated according to national schedule (12–23 months)	0.546	0.095	30	25	1.032	0.174	0.356	0.736
Received measles/measles-rubella 2 vaccination (24–35 months)	0.757	0.090	23	19	0.978	0.119	0.576	0.937
Fully vaccinated according to national schedule (24–35 months)	0.430	0.114	23	19	1.047	0.265	0.202	0.657
Received HPV vaccination (2 doses) women 15–19	0.724	0.064	41	36	0.912	0.089	0.595	0.853
Sought treatment for diarrhoea	0.490	0.186	10	7	1.025	0.379	0.119	0.862
Treated with ORS	0.490	0.186	10	7	1.025	0.379	0.119	0.862
Exclusive breastfeeding	0.616	0.157	14	11	1.150	0.255	0.302	0.930
Minimum dietary diversity (children 6–23 months)	0.571	0.081	44	37	1.066	0.141	0.410	0.732
Child slept under an ITN last night	0.107	0.039	139	107	1.343	0.366	0.029	0.185
Received 3+ doses of SP/Fansidar	0.000	0.000	58	48	na	na	0.000	0.000
Child has fever in last 2 weeks	0.063	0.029	123	102	1.271	0.454	0.006	0.121
Child had blood taken from finger/heel	0.000	0.000	7	6	na	na	0.000	0.000
Child took ACT	na	na	-	-	na	na	0.000	na
Condom use at last sex (women 15–24)	0.373	0.113	18	16	0.967	0.303	0.147	0.600
Mobile phone ownership	0.880	0.024	260	223	1.208	0.028	0.831	0.929

na = not applicable

Table B.23 Sampling errors: Nyeri sample, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases: Unweighted (N)	Number of cases: Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
HOUSEHOLDS AND POPULATION								
Primary reliance on clean fuels and technology	0.188	0.031	1,480	1,148	1.743	0.166	0.125	0.250
Births registered with civil authority	0.942	0.026	108	82	1.004	0.028	0.889	0.995
Ownership of at least one ITN	0.034	0.009	570	441	1.231	0.275	0.015	0.053
Ownership of at least one ITN for every two persons	0.014	0.004	565	437	0.868	0.308	0.005	0.022
At least basic drinking water service	0.887	0.034	1,480	1,148	2.025	0.039	0.819	0.955
At least basic sanitation service	0.416	0.041	1,480	1,148	1.678	0.100	0.333	0.499
Using open defecation	0.000	0.000	1,480	1,148	na	na	0.000	0.000
Using a handwashing facility with soap and water	0.790	0.034	1,457	1,134	1.662	0.043	0.722	0.858
WOMEN								
No education	0.010	0.007	270	220	1.214	0.735	0.000	0.025
Secondary education or higher	0.778	0.032	270	220	1.277	0.042	0.713	0.843
Literacy	0.984	0.008	273	223	1.046	0.008	0.968	1.000
Use of the internet in last 12 months	0.639	0.028	273	223	0.968	0.044	0.583	0.695
Total fertility rate (3 years)	2.818	0.273	788	642	0.924	0.097	2.271	3.365
Currently pregnant	0.029	0.011	273	223	1.118	0.394	0.006	0.051
Mean number of children ever born to women age 40–49	3.041	0.140	70	59	1.012	0.046	2.761	3.320
Median birth interval	55.597	5.484	62	53	0.858	0.099	44.629	66.565
Ideal number of children	2.838	0.061	273	223	0.816	0.022	2.715	2.961
Total wanted fertility rate (3 years)	2.220	0.254	788	642	1.037	0.114	1.712	2.728
Currently using any contraceptive method	0.615	0.039	125	104	0.890	0.063	0.537	0.692
Currently using any modern method	0.607	0.042	125	104	0.951	0.069	0.524	0.690
Currently using pill	0.182	0.045	125	104	1.300	0.248	0.091	0.272
Currently using injectables	0.112	0.026	125	104	0.926	0.234	0.060	0.165
Currently using implants	0.210	0.042	125	104	1.149	0.200	0.126	0.294
Currently using male condoms	0.000	0.000	125	104	na	na	0.000	0.000
Currently using any traditional method	0.007	0.007	125	104	0.951	0.986	0.000	0.022
Unmet need for spacing	0.049	0.018	125	104	0.915	0.362	0.013	0.084
Unmet need for limiting	0.144	0.028	125	104	0.891	0.195	0.088	0.200
Unmet need total	0.192	0.029	125	104	0.813	0.149	0.135	0.250
Demand satisfied by modern methods (married women)	0.752	0.035	100	84	0.812	0.047	0.682	0.822
Demand satisfied by modern methods (all women)	0.768	0.030	139	115	0.843	0.039	0.708	0.828
Not exposed to any of the ten media sources	0.098	0.015	273	223	0.844	0.155	0.068	0.129
Neonatal mortality (last 0–9 years)	20.261	10.083	222	182	1.078	0.498	0.094	40.428
Post-neonatal mortality (last 0–9 years)	4.256	4.266	220	180	0.942	1.002	0.000	12.789
Infant mortality (last 0–9 years)	24.517	10.212	222	182	0.985	0.417	4.093	44.942
Child mortality (last 0–9 years)	13.177	7.456	213	175	0.935	0.566	0.000	28.089
Under-five mortality (last 0–9 years)	37.371	12.339	223	183	0.855	0.330	12.692	62.050
Stillbirth rate	11.128	10.840	104	86	1.072	0.974	0.000	32.808
Early neonatal rate	18.033	17.314	103	85	1.337	0.960	0.000	52.660
Perinatal mortality rate	28.960	18.925	104	86	1.178	0.653	0.000	66.810
Received ANC from a skilled provider	1.000	0.000	50	42	na	0.000	1.000	1.000
4+ ANC visits	0.801	0.064	50	42	1.124	0.080	0.673	0.930
8+ ANC visits	0.094	0.043	50	42	1.031	0.458	0.008	0.180
Took any iron-containing supplements	0.906	0.041	50	42	0.985	0.045	0.824	0.988
Mothers protected against tetanus for last birth	0.891	0.037	50	42	0.827	0.041	0.817	0.964
Delivered in a health facility (live births)	1.000	0.000	50	42	na	0.000	1.000	1.000
Delivered by a skilled provider (live births)	1.000	0.000	50	42	na	0.000	1.000	1.000
Delivered by C-section (live births)	0.369	0.076	50	42	1.121	0.206	0.217	0.521
Women with postnatal check during first 2 days	0.965	0.022	50	42	0.850	0.023	0.921	1.000
Newborns with postnatal check during first 2 days	0.965	0.022	50	42	0.850	0.023	0.921	1.000
Any problem accessing health care	0.501	0.031	273	223	1.036	0.063	0.438	0.564
Ever had a vaccination card	1.000	0.000	23	20	na	0.000	1.000	1.000
Received BCG vaccination	1.000	0.000	23	20	na	0.000	1.000	1.000
Received DPT-HepB+Hib vaccination (3 doses)	0.862	0.071	23	20	1.009	0.082	0.720	1.000
Received pneumococcal vaccination (3 doses)	0.911	0.064	23	20	1.102	0.070	0.782	1.000
Received measles vaccination	0.966	0.030	23	20	0.801	0.031	0.907	1.000
Fully vaccinated according to national schedule (12–23 months)	0.551	0.109	23	20	1.071	0.198	0.333	0.769
Received measles/measles-rubella 2 vaccination (24–35 months)	0.852	0.090	16	13	1.007	0.105	0.672	1.000
Fully vaccinated according to national schedule (24–35 months)	0.567	0.100	16	13	0.805	0.176	0.367	0.768
Received HPV vaccination (2 doses) women 15–19	0.383	0.080	41	33	1.035	0.208	0.224	0.543
Sought treatment for diarrhoea	0.167	0.132	5	4	0.783	0.791	0.000	0.431
Treated with ORS	0.167	0.132	5	4	0.783	0.791	0.000	0.431
Exclusive breastfeeding	0.921	0.076	12	10	0.943	0.083	0.769	1.000
Minimum dietary diversity (children 6–23 months)	0.678	0.080	35	29	1.004	0.119	0.518	0.839
Child slept under an ITN last night	0.070	0.028	114	87	1.178	0.407	0.013	0.126
Received 3+ doses of SP/Fansidar	0.000	0.000	50	42	na	na	0.000	0.000
Child has fever in last 2 weeks	0.146	0.026	101	84	0.750	0.178	0.094	0.199
Child had blood taken from finger/heel	0.090	0.087	15	12	1.175	0.964	0.000	0.265
Child took ACT	na	na	-	-	na	na	0.000	na
Condom use at last sex (women 15–24)	0.403	0.092	17	15	0.758	0.228	0.219	0.586
Mobile phone ownership	0.896	0.015	273	223	0.824	0.017	0.866	0.927

na = not applicable

Table B.24 Sampling errors: Kirinyaga sample, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases: Unweighted (N)	Number of cases: Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
HOUSEHOLDS AND POPULATION								
Primary reliance on clean fuels and technology	0.277	0.037	1,458	1,210	1.771	0.132	0.204	0.350
Births registered with civil authority	0.872	0.026	123	100	0.853	0.030	0.820	0.925
Ownership of at least one ITN	0.446	0.068	582	482	3.265	0.152	0.310	0.582
Ownership of at least one ITN for every two persons	0.395	0.063	568	471	3.024	0.158	0.270	0.520
At least basic drinking water service	0.847	0.058	1,458	1,210	3.369	0.069	0.730	0.964
At least basic sanitation service	0.582	0.034	1,458	1,210	1.471	0.058	0.514	0.650
Using open defecation	0.001	0.001	1,458	1,210	0.976	0.993	0.000	0.002
Using a handwashing facility with soap and water	0.340	0.050	1,338	1,115	2.043	0.146	0.241	0.439
WOMEN								
No education	0.009	0.003	319	258	0.613	0.361	0.002	0.015
Secondary education or higher	0.675	0.032	319	258	1.233	0.048	0.610	0.740
Literacy	0.953	0.012	319	258	1.055	0.013	0.929	0.978
Use of the internet in last 12 months	0.674	0.031	319	258	1.191	0.047	0.611	0.736
Total fertility rate (3 years)	2.684	0.313	926	752	1.125	0.117	2.058	3.310
Currently pregnant	0.045	0.011	319	258	0.925	0.238	0.024	0.067
Mean number of children ever born to women age 40–49	2.682	0.110	94	77	1.039	0.041	2.463	2.902
Median birth interval	63.756	4.178	75	60	1.394	0.066	55.400	72.112
Ideal number of children	2.753	0.066	317	255	1.141	0.024	2.622	2.885
Total wanted fertility rate (3 years)	2.454	0.323	926	752	1.168	0.132	1.808	3.100
Currently using any contraceptive method	0.767	0.034	176	141	1.073	0.045	0.699	0.836
Currently using any modern method	0.712	0.034	176	141	1.003	0.048	0.644	0.781
Currently using pill	0.209	0.041	176	141	1.330	0.196	0.127	0.291
Currently using injectables	0.196	0.026	176	141	0.864	0.132	0.144	0.248
Currently using implants	0.196	0.034	176	141	1.147	0.176	0.127	0.265
Currently using male condoms	0.005	0.005	176	141	0.966	1.004	0.000	0.016
Currently using any traditional method	0.055	0.016	176	141	0.927	0.290	0.023	0.087
Unmet need for spacing	0.030	0.014	176	141	1.080	0.467	0.002	0.057
Unmet need for limiting	0.040	0.014	176	141	0.983	0.366	0.011	0.069
Unmet need total	0.069	0.016	176	141	0.847	0.234	0.037	0.102
Demand satisfied by modern methods (married women)	0.851	0.017	148	118	0.580	0.020	0.817	0.885
Demand satisfied by modern methods (all women)	0.832	0.020	203	160	0.760	0.024	0.792	0.872
Not exposed to any of the ten media sources	0.034	0.009	319	258	0.884	0.266	0.016	0.052
Neonatal mortality (last 0–9 years)	35.959	12.703	246	201	1.064	0.353	10.553	61.365
Post-neonatal mortality (last 0–9 years)	5.311	3.793	246	200	0.761	0.714	0.000	12.897
Infant mortality (last 0–9 years)	41.271	12.502	246	201	0.990	0.303	16.266	66.275
Child mortality (last 0–9 years)	8.005	5.694	245	201	1.010	0.711	0.000	19.394
Under-five mortality (last 0–9 years)	48.945	12.121	246	201	0.891	0.248	24.703	73.188
Stillbirth rate	0.000	0.000	122	98	na	na	0.000	0.000
Early neonatal rate	34.685	24.194	122	98	1.482	0.698	0.000	83.072
Perinatal mortality rate	34.685	24.194	122	98	1.482	0.698	0.000	83.072
Received ANC from a skilled provider	0.981	0.020	46	37	0.975	0.020	0.942	1.000
4+ ANC visits	0.815	0.059	46	37	1.017	0.072	0.697	0.933
8+ ANC visits	0.115	0.048	46	37	1.017	0.421	0.018	0.212
Took any iron-containing supplements	0.967	0.024	46	37	0.915	0.025	0.918	1.000
Mothers protected against tetanus for last birth	0.735	0.057	46	37	0.872	0.078	0.621	0.850
Delivered in a health facility (live births)	1.000	0.000	47	38	na	0.000	1.000	1.000
Delivered by a skilled provider (live births)	1.000	0.000	47	38	na	0.000	1.000	1.000
Delivered by C-section (live births)	0.375	0.048	47	38	0.668	0.128	0.279	0.471
Women with postnatal check during first 2 days	0.871	0.057	46	37	1.129	0.065	0.758	0.984
Newborns with postnatal check during first 2 days	0.944	0.034	46	37	0.999	0.036	0.876	1.000
Any problem accessing health care	0.546	0.036	319	258	1.281	0.066	0.474	0.617
Ever had a vaccination card	1.000	0.000	25	20	na	0.000	1.000	1.000
Received BCG vaccination	1.000	0.000	25	20	na	0.000	1.000	1.000
Received DPT-HepB+Hib vaccination (3 doses)	0.875	0.077	25	20	1.170	0.089	0.720	1.000
Received pneumococcal vaccination (3 doses)	0.928	0.055	25	20	1.061	0.059	0.818	1.000
Received measles vaccination	0.954	0.045	25	20	1.076	0.047	0.864	1.000
Fully vaccinated according to national schedule (12–23 months)	0.505	0.132	25	20	1.323	0.262	0.240	0.770
Received measles/measles-rubella 2 vaccination (24–35 months)	0.865	0.096	19	17	1.266	0.111	0.673	1.000
Fully vaccinated according to national schedule (24–35 months)	0.480	0.115	19	17	1.035	0.240	0.250	0.711
Received HPV vaccination (2 doses) women 15–19	0.591	0.067	37	30	0.818	0.113	0.457	0.724
Sought treatment for diarrhoea	0.622	0.148	10	8	0.961	0.239	0.325	0.919
Treated with ORS	0.544	0.110	10	8	0.696	0.203	0.323	0.765
Exclusive breastfeeding	0.717	0.183	9	8	1.127	0.255	0.352	1.000
Minimum dietary diversity (children 6–23 months)	0.668	0.098	35	28	1.206	0.147	0.472	0.864
Child slept under an ITN last night	0.548	0.087	126	102	1.761	0.159	0.374	0.722
Received 3+ doses of SP/Fansidar	0.057	0.032	46	37	0.942	0.573	0.000	0.121
Child has fever in last 2 weeks	0.135	0.040	114	91	1.261	0.298	0.054	0.216
Child had blood taken from finger/heel	0.000	0.000	14	12	na	na	0.000	0.000
Child took ACT	0.000	0.000	2	2	na	na	0.000	0.000
Condom use at last sex (women 15–24)	0.184	0.071	15	11	0.703	0.389	0.041	0.326
Mobile phone ownership	0.905	0.019	319	258	1.136	0.021	0.868	0.943

na = not applicable

Table B.25 Sampling errors: Murang'a sample, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases: Unweighted (N)	Number of cases: Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
HOUSEHOLDS AND POPULATION								
Primary reliance on clean fuels and technology	0.175	0.027	1,886	1,708	1.625	0.155	0.120	0.229
Births registered with civil authority	0.941	0.023	167	150	1.246	0.024	0.896	0.986
Ownership of at least one ITN	0.188	0.026	631	571	1.657	0.137	0.137	0.240
Ownership of at least one ITN for every two persons	0.111	0.018	626	568	1.412	0.160	0.076	0.147
At least basic drinking water service	0.825	0.036	1,886	1,708	1.990	0.043	0.753	0.896
At least basic sanitation service	0.564	0.033	1,886	1,708	1.461	0.059	0.497	0.631
Using open defecation	0.000	0.000	1,886	1,708	na	na	0.000	0.000
Using a handwashing facility with soap and water	0.701	0.044	1,824	1,669	2.053	0.063	0.612	0.789
WOMEN								
No education	0.003	0.003	434	377	1.069	1.001	0.000	0.008
Secondary education or higher	0.660	0.033	434	377	1.447	0.050	0.594	0.726
Literacy	0.964	0.011	434	377	1.213	0.011	0.942	0.986
Use of the internet in last 12 months	0.597	0.042	434	377	1.784	0.071	0.513	0.681
Total fertility rate (3 years)	3.388	0.307	1,204	1,044	1.166	0.091	2.775	4.002
Currently pregnant	0.031	0.007	434	377	0.888	0.239	0.016	0.046
Mean number of children ever born to women age 40–49	3.434	0.119	124	114	0.836	0.035	3.196	3.672
Median birth interval	57.074	3.954	118	102	1.060	0.069	49.167	64.982
Ideal number of children	3.312	0.063	434	377	0.894	0.019	3.187	3.438
Total wanted fertility rate (3 years)	3.071	0.249	1,204	1,044	1.017	0.081	2.574	3.569
Currently using any contraceptive method	0.784	0.030	206	182	1.060	0.039	0.723	0.845
Currently using any modern method	0.756	0.033	206	182	1.091	0.043	0.690	0.821
Currently using pill	0.138	0.020	206	182	0.829	0.145	0.098	0.178
Currently using injectables	0.268	0.039	206	182	1.263	0.146	0.190	0.347
Currently using implants	0.204	0.029	206	182	1.037	0.143	0.146	0.263
Currently using male condoms	0.012	0.007	206	182	0.940	0.589	0.000	0.027
Currently using any traditional method	0.028	0.011	206	182	0.926	0.381	0.007	0.049
Unmet need for spacing	0.018	0.009	206	182	1.014	0.528	0.000	0.036
Unmet need for limiting	0.050	0.016	206	182	1.066	0.324	0.018	0.083
Unmet need total	0.068	0.016	206	182	0.914	0.236	0.036	0.100
Demand satisfied by modern methods (married women)	0.887	0.022	176	155	0.905	0.024	0.844	0.931
Demand satisfied by modern methods (all women)	0.866	0.020	228	196	0.859	0.023	0.827	0.905
Not exposed to any of the ten media sources	0.012	0.006	434	377	1.161	0.511	0.000	0.024
Neonatal mortality (last 0–9 years)	28.674	9.641	354	308	1.097	0.336	9.392	47.955
Post-neonatal mortality (last 0–9 years)	4.433	3.328	354	307	0.977	0.751	0.000	11.088
Infant mortality (last 0–9 years)	33.107	9.727	354	308	1.035	0.294	13.653	52.561
Child mortality (last 0–9 years)	0.000	0.000	370	321	na	na	0.000	0.000
Under-five mortality (last 0–9 years)	33.107	9.727	354	308	1.035	0.294	13.653	52.561
Stillbirth rate	4.983	4.885	167	146	0.903	0.980	0.000	14.753
Early neonatal rate	34.605	16.149	166	145	1.154	0.467	2.307	66.902
Perinatal mortality rate	39.415	17.738	167	146	1.199	0.450	3.939	74.891
Received ANC from a skilled provider	0.985	0.015	66	59	1.016	0.015	0.955	1.000
4+ ANC visits	0.678	0.064	66	59	1.094	0.094	0.551	0.805
8+ ANC visits	0.093	0.037	66	59	1.026	0.399	0.019	0.166
Took any iron-containing supplements	0.944	0.028	66	59	0.982	0.030	0.888	1.000
Mothers protected against tetanus for last birth	0.947	0.029	66	59	1.028	0.030	0.890	1.000
Delivered in a health facility (live births)	0.967	0.023	68	60	1.087	0.024	0.921	1.000
Delivered by a skilled provider (live births)	0.967	0.023	68	60	1.087	0.024	0.921	1.000
Delivered by C-section (live births)	0.356	0.086	68	60	1.500	0.241	0.184	0.528
Women with postnatal check during first 2 days	0.958	0.024	66	59	0.962	0.025	0.911	1.000
Newborns with postnatal check during first 2 days	0.959	0.020	66	59	0.827	0.021	0.918	0.999
Any problem accessing health care	0.366	0.041	434	377	1.755	0.111	0.285	0.448
Ever had a vaccination card	1.000	0.000	38	32	na	0.000	1.000	1.000
Received BCG vaccination	0.969	0.032	38	32	1.107	0.033	0.904	1.000
Received DPT-HepB+Hib vaccination (3 doses)	0.933	0.047	38	32	1.116	0.050	0.839	1.000
Received pneumococcal vaccination (3 doses)	0.969	0.032	38	32	1.107	0.033	0.904	1.000
Received measles vaccination	0.964	0.029	38	32	0.857	0.030	0.906	1.000
Fully vaccinated according to national schedule (12–23 months)	0.706	0.081	38	32	1.060	0.115	0.544	0.869
Received measles/measles-rubella 2 vaccination (24–35 months)	0.822	0.053	39	34	0.859	0.064	0.717	0.927
Fully vaccinated according to national schedule (24–35 months)	0.468	0.073	39	34	0.920	0.157	0.321	0.615
Received HPV vaccination (2 doses) women 15–19	0.565	0.059	89	77	1.112	0.104	0.447	0.683
Sought treatment for diarrhoea	0.615	0.204	6	5	1.023	0.331	0.207	1.000
Treated with ORS	0.382	0.215	6	5	1.083	0.563	0.000	0.812
Exclusive breastfeeding	1.000	0.000	12	12	na	0.000	1.000	1.000
Minimum dietary diversity (children 6–23 months)	0.596	0.088	51	45	1.264	0.148	0.419	0.772
Child slept under an ITN last night	0.305	0.050	184	163	1.262	0.164	0.205	0.406
Received 3+ doses of SP/Fansidar	0.000	0.000	66	59	na	na	0.000	0.000
Child has fever in last 2 weeks	0.045	0.015	159	138	0.910	0.334	0.015	0.075
Child had blood taken from finger/heel	0.477	0.202	8	6	1.077	0.424	0.072	0.881
Child took ACT	0.000	0.000	2	2	na	na	0.000	0.000
Condom use at last sex (women 15–24)	0.405	0.079	27	23	0.830	0.196	0.246	0.563
Mobile phone ownership	0.840	0.027	434	377	1.550	0.033	0.785	0.895

na = not applicable

Table B.26 Sampling errors: Kiambu sample, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases: Unweighted (N)	Number of cases: Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
HOUSEHOLDS AND POPULATION								
Primary reliance on clean fuels and technology	0.752	0.027	2,410	6,076	1.503	0.036	0.698	0.807
Births registered with civil authority	0.921	0.020	239	652	1.107	0.022	0.882	0.961
Ownership of at least one ITN	0.168	0.022	864	2,295	1.707	0.130	0.124	0.211
Ownership of at least one ITN for every two persons	0.100	0.016	864	2,295	1.593	0.163	0.067	0.132
At least basic drinking water service	0.969	0.011	2,410	6,076	1.428	0.011	0.948	0.990
At least basic sanitation service	0.551	0.066	2,410	6,076	3.284	0.120	0.419	0.684
Using open defecation	0.000	0.000	2,410	6,076	na	na	0.000	0.000
Using a handwashing facility with soap and water	0.526	0.065	2,170	5,596	3.027	0.123	0.396	0.656
WOMEN								
No education	0.009	0.004	632	1,689	1.049	0.434	0.001	0.017
Secondary education or higher	0.839	0.025	632	1,689	1.710	0.030	0.789	0.889
Literacy	0.978	0.006	632	1,689	0.954	0.006	0.967	0.989
Use of the internet in last 12 months	0.795	0.031	632	1,689	1.916	0.039	0.734	0.857
Total fertility rate (3 years)	2.328	0.248	1,820	4,885	1.443	0.107	1.832	2.825
Currently pregnant	0.040	0.008	632	1,689	0.965	0.188	0.025	0.055
Mean number of children ever born to women age 40–49	3.385	0.238	106	234	1.321	0.070	2.909	3.862
Median birth interval	55.413	3.186	187	494	0.881	0.057	49.041	61.785
Ideal number of children	3.106	0.070	627	1,677	1.167	0.022	2.966	3.245
Total wanted fertility rate (3 years)	2.063	0.247	1,820	4,885	1.415	0.120	1.568	2.557
Currently using any contraceptive method	0.616	0.029	297	819	1.030	0.047	0.558	0.674
Currently using any modern method	0.573	0.029	297	819	0.995	0.050	0.516	0.630
Currently using pill	0.153	0.027	297	819	1.279	0.175	0.099	0.206
Currently using injectables	0.130	0.019	297	819	0.983	0.148	0.092	0.169
Currently using implants	0.197	0.023	297	819	0.991	0.116	0.152	0.243
Currently using male condoms	0.005	0.004	297	819	0.940	0.752	0.000	0.013
Currently using any traditional method	0.043	0.011	297	819	0.972	0.266	0.020	0.066
Unmet need for spacing	0.099	0.014	297	819	0.816	0.143	0.071	0.128
Unmet need for limiting	0.080	0.015	297	819	0.969	0.191	0.050	0.111
Unmet need total	0.179	0.019	297	819	0.849	0.105	0.142	0.217
Demand satisfied by modern methods (married women)	0.720	0.024	237	652	0.834	0.034	0.672	0.769
Demand satisfied by modern methods (all women)	0.713	0.024	315	849	0.956	0.034	0.664	0.762
Not exposed to any of the ten media sources	0.040	0.011	632	1,689	1.468	0.287	0.017	0.063
Neonatal mortality (last 0–9 years)	12.148	5.514	532	1,402	0.965	0.454	1.120	23.175
Post-neonatal mortality (last 0–9 years)	9.915	5.036	529	1,391	1.094	0.508	0.000	19.987
Infant mortality (last 0–9 years)	22.063	6.613	532	1,402	0.939	0.300	8.837	35.288
Child mortality (last 0–9 years)	1.770	1.357	511	1,321	0.670	0.767	0.000	4.485
Under-five mortality (last 0–9 years)	23.794	7.135	532	1,402	0.901	0.300	9.523	38.064
Stillbirth rate	2.115	2.092	270	735	0.756	0.989	0.000	6.298
Early neonatal rate	18.390	9.616	270	740	1.084	0.523	0.000	37.621
Perinatal mortality rate	20.643	10.703	270	735	1.139	0.518	0.000	42.049
Received ANC from a skilled provider	0.982	0.011	100	269	0.820	0.011	0.960	1.000
4+ ANC visits	0.843	0.036	100	269	0.972	0.042	0.771	0.914
8+ ANC visits	0.158	0.040	100	269	1.083	0.251	0.079	0.238
Took any iron-containing supplements	0.950	0.024	100	269	1.086	0.025	0.902	0.998
Mothers protected against tetanus for last birth	0.817	0.045	100	269	1.143	0.055	0.728	0.906
Delivered in a health facility (live births)	0.983	0.011	102	272	0.882	0.012	0.960	1.000
Delivered by a skilled provider (live births)	0.986	0.011	102	272	0.961	0.011	0.963	1.000
Delivered by C-section (live births)	0.386	0.057	102	272	1.159	0.147	0.273	0.500
Women with postnatal check during first 2 days	0.811	0.049	100	269	1.230	0.060	0.714	0.908
Newborns with postnatal check during first 2 days	0.849	0.042	100	269	1.166	0.050	0.764	0.933
Any problem accessing health care	0.366	0.037	632	1,689	1.917	0.101	0.292	0.439
Ever had a vaccination card	0.992	0.008	43	104	0.566	0.008	0.976	1.000
Received BCG vaccination	0.938	0.055	43	104	1.429	0.059	0.827	1.000
Received DPT-HepB+Hib vaccination (3 doses)	0.918	0.058	43	104	1.317	0.063	0.803	1.000
Received pneumococcal vaccination (3 doses)	0.901	0.044	43	104	0.917	0.049	0.814	0.989
Received measles vaccination	0.875	0.062	43	104	1.175	0.071	0.751	1.000
Fully vaccinated according to national schedule (12–23 months)	0.344	0.084	43	104	1.110	0.245	0.175	0.513
Received measles/measles-rubella 2 vaccination (24–35 months)	0.679	0.097	47	133	1.447	0.142	0.486	0.872
Fully vaccinated according to national schedule (24–35 months)	0.425	0.092	47	133	1.302	0.217	0.240	0.610
Received HPV vaccination (2 doses) women 15–19	0.435	0.046	100	245	0.914	0.105	0.344	0.526
Sought treatment for diarrhoea	0.410	0.150	22	62	1.402	0.365	0.110	0.709
Treated with ORS	0.474	0.163	22	62	1.482	0.343	0.148	0.799
Exclusive breastfeeding	0.839	0.069	33	96	1.062	0.082	0.701	0.977
Minimum dietary diversity (children 6–23 months)	0.332	0.061	61	157	0.997	0.183	0.210	0.453
Child slept under an ITN last night	0.232	0.040	250	685	1.310	0.171	0.152	0.311
Received 3+ doses of SP/Fansidar	0.020	0.013	100	269	0.940	0.664	0.000	0.046
Child has fever in last 2 weeks	0.057	0.020	261	714	1.325	0.355	0.016	0.097
Child had blood taken from finger/heel	0.144	0.090	18	41	0.982	0.624	0.000	0.323
Child took ACT	na	na	-	-	na	na	0.000	na
Condom use at last sex (women 15–24)	0.302	0.066	47	139	0.972	0.218	0.170	0.433
Mobile phone ownership	0.925	0.014	632	1,689	1.371	0.016	0.897	0.954

na = not applicable

Table B.27 Sampling errors: Turkana sample, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases: Unweighted (N)	Number of cases: Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
HOUSEHOLDS AND POPULATION								
Primary reliance on clean fuels and technology	0.018	0.008	2,351	1,183	1.642	0.462	0.001	0.034
Births registered with civil authority	0.691	0.051	358	187	1.785	0.074	0.589	0.793
Ownership of at least one ITN	0.844	0.032	474	232	1.882	0.037	0.781	0.907
Ownership of at least one ITN for every two persons	0.515	0.038	472	231	1.662	0.074	0.439	0.592
At least basic drinking water service	0.443	0.086	2,351	1,183	3.298	0.193	0.272	0.614
At least basic sanitation service	0.148	0.043	2,351	1,183	2.144	0.291	0.062	0.234
Using open defecation	0.712	0.063	2,351	1,183	2.632	0.089	0.586	0.839
Using a handwashing facility with soap and water	0.556	0.033	2,347	1,181	1.319	0.060	0.489	0.623
WOMEN								
No education	0.518	0.061	344	178	2.264	0.119	0.395	0.640
Secondary education or higher	0.312	0.053	344	178	2.099	0.169	0.207	0.418
Literacy	0.462	0.065	344	178	2.409	0.141	0.332	0.593
Use of the internet in last 12 months	0.178	0.050	344	178	2.416	0.282	0.078	0.279
Total fertility rate (3 years)	5.629	0.588	971	502	1.660	0.105	4.452	6.805
Currently pregnant	0.061	0.015	344	178	1.142	0.243	0.031	0.090
Mean number of children ever born to women age 40–49	5.894	0.315	61	30	1.133	0.054	5.263	6.525
Median birth interval	32.315	1.069	253	136	1.419	0.033	30.177	34.453
Ideal number of children	5.625	0.165	332	171	1.341	0.029	5.294	5.956
Total wanted fertility rate (3 years)	5.094	0.512	971	502	1.580	0.100	4.071	6.117
Currently using any contraceptive method	0.214	0.027	220	116	0.989	0.128	0.159	0.268
Currently using any modern method	0.170	0.030	220	116	1.200	0.179	0.109	0.231
Currently using pill	0.008	0.008	220	116	1.328	0.991	0.000	0.024
Currently using injectables	0.068	0.017	220	116	0.989	0.248	0.034	0.101
Currently using implants	0.081	0.020	220	116	1.094	0.249	0.041	0.121
Currently using male condoms	0.000	0.000	220	116	na	na	0.000	0.000
Currently using any traditional method	0.043	0.012	220	116	0.887	0.281	0.019	0.068
Unmet need for spacing	0.256	0.029	220	116	0.987	0.114	0.198	0.314
Unmet need for limiting	0.083	0.018	220	116	0.985	0.222	0.046	0.119
Unmet need total	0.339	0.036	220	116	1.139	0.108	0.266	0.412
Demand satisfied by modern methods (married women)	0.308	0.043	121	64	1.035	0.141	0.221	0.395
Demand satisfied by modern methods (all women)	0.348	0.035	140	73	0.875	0.101	0.278	0.418
Not exposed to any of the ten media sources	0.148	0.022	344	178	1.148	0.149	0.104	0.193
Neonatal mortality (last 0–9 years)	6.876	3.910	633	334	0.869	0.569	0.000	14.697
Post-neonatal mortality (last 0–9 years)	19.820	6.666	630	333	0.996	0.336	6.489	33.151
Infant mortality (last 0–9 years)	26.696	7.744	633	334	0.965	0.290	11.207	42.184
Child mortality (last 0–9 years)	10.064	5.228	626	335	0.931	0.519	0.000	20.520
Under-five mortality (last 0–9 years)	36.491	9.106	636	336	0.927	0.250	18.280	54.703
Stillbirth rate	16.917	9.941	315	170	1.404	0.588	0.000	36.800
Early neonatal rate	4.127	3.255	311	167	0.912	0.789	0.000	10.638
Perinatal mortality rate	20.975	10.105	315	170	1.286	0.482	0.765	41.185
Received ANC from a skilled provider	0.990	0.010	118	62	1.038	0.010	0.971	1.000
4+ ANC visits	0.509	0.083	118	62	1.782	0.163	0.343	0.675
8+ ANC visits	0.000	0.000	118	62	na	na	0.000	0.000
Took any iron-containing supplements	0.823	0.042	118	62	1.186	0.051	0.739	0.907
Mothers protected against tetanus for last birth	0.799	0.031	118	62	0.841	0.039	0.736	0.861
Delivered in a health facility (live births)	0.518	0.056	121	63	1.203	0.108	0.406	0.630
Delivered by a skilled provider (live births)	0.559	0.054	121	63	1.176	0.097	0.450	0.667
Delivered by C-section (live births)	0.100	0.033	121	63	1.196	0.328	0.035	0.166
Women with postnatal check during first 2 days	0.620	0.063	118	62	1.402	0.102	0.494	0.747
Newborns with postnatal check during first 2 days	0.632	0.051	118	62	1.144	0.081	0.530	0.734
Any problem accessing health care	0.735	0.041	344	178	1.702	0.055	0.654	0.816
Ever had a vaccination card	1.000	0.000	61	33	na	0.000	1.000	1.000
Received BCG vaccination	0.965	0.028	61	33	1.219	0.029	0.908	1.000
Received DPT-HepB+Hib vaccination (3 doses)	0.899	0.036	61	33	0.953	0.040	0.827	0.971
Received pneumococcal vaccination (3 doses)	0.909	0.045	61	33	1.236	0.049	0.820	0.998
Received measles vaccination	0.815	0.061	61	33	1.258	0.075	0.692	0.938
Fully vaccinated according to national schedule (12–23 months)	0.104	0.037	61	33	0.971	0.358	0.029	0.178
Received measles/measles-rubella 2 vaccination (24–35 months)	0.578	0.061	55	30	0.920	0.106	0.455	0.701
Fully vaccinated according to national schedule (24–35 months)	0.099	0.035	55	30	0.888	0.354	0.029	0.170
Received HPV vaccination (2 doses) women 15–19	0.083	0.036	74	37	1.126	0.440	0.010	0.155
Sought treatment for diarrhoea	0.689	0.080	43	23	1.137	0.116	0.529	0.849
Treated with ORS	0.634	0.061	43	23	0.829	0.096	0.512	0.756
Exclusive breastfeeding	0.734	0.113	32	16	1.401	0.154	0.508	0.960
Minimum dietary diversity (children 6–23 months)	0.157	0.060	81	43	1.464	0.382	0.037	0.277
Child slept under an ITN last night	0.813	0.042	354	184	1.708	0.052	0.729	0.896
Received 3+ doses of SP/Fansidar	0.020	0.012	118	62	0.903	0.586	0.000	0.043
Child has fever in last 2 weeks	0.244	0.026	302	162	1.032	0.105	0.193	0.295
Child had blood taken from finger/heel	0.695	0.057	72	40	1.108	0.083	0.580	0.810
Child took ACT	0.833	0.078	31	18	1.237	0.093	0.677	0.988
Condom use at last sex (women 15–24)	0.453	0.156	11	4	0.989	0.343	0.142	0.764
Mobile phone ownership	0.624	0.035	344	178	1.337	0.056	0.554	0.694

na = not applicable

Table B.28 Sampling errors: West Pokot sample, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases: Unweighted (N)	Number of cases: Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
HOUSEHOLDS AND POPULATION								
Primary reliance on clean fuels and technology	0.017	0.008	2,377	1,259	1.759	0.476	0.001	0.033
Births registered with civil authority	0.469	0.072	384	201	2.270	0.153	0.326	0.613
Ownership of at least one ITN	0.654	0.057	495	251	2.660	0.088	0.539	0.769
Ownership of at least one ITN for every two persons	0.288	0.031	491	249	1.532	0.109	0.225	0.350
At least basic drinking water service	0.405	0.106	2,377	1,259	4.071	0.261	0.194	0.617
At least basic sanitation service	0.475	0.088	2,377	1,259	3.343	0.185	0.299	0.650
Using open defecation	0.326	0.087	2,377	1,259	3.585	0.268	0.151	0.501
Using a handwashing facility with soap and water	0.298	0.044	2,338	1,236	1.835	0.148	0.210	0.386
WOMEN								
No education	0.182	0.053	441	223	2.850	0.291	0.076	0.287
Secondary education or higher	0.490	0.065	441	223	2.705	0.132	0.360	0.620
Literacy	0.776	0.053	444	225	2.675	0.069	0.670	0.883
Use of the internet in last 12 months	0.211	0.052	444	225	2.679	0.248	0.107	0.316
Total fertility rate (3 years)	5.128	0.455	1,235	627	1.324	0.089	4.219	6.038
Currently pregnant	0.075	0.016	444	225	1.289	0.215	0.043	0.108
Mean number of children ever born to women age 40–49	6.093	0.392	57	31	1.156	0.064	5.309	6.876
Median birth interval	30.458	1.808	271	140	1.600	0.059	26.841	34.075
Ideal number of children	6.257	0.396	444	225	3.079	0.063	5.466	7.049
Total wanted fertility rate (3 years)	4.815	0.476	1,235	627	1.346	0.099	3.864	5.767
Currently using any contraceptive method	0.183	0.048	278	145	2.073	0.265	0.086	0.280
Currently using any modern method	0.176	0.048	278	145	2.088	0.273	0.080	0.272
Currently using pill	0.012	0.007	278	145	1.004	0.541	0.000	0.026
Currently using injectables	0.057	0.018	278	145	1.316	0.322	0.020	0.094
Currently using implants	0.088	0.025	278	145	1.448	0.280	0.039	0.138
Currently using male condoms	0.009	0.005	278	145	0.962	0.610	0.000	0.020
Currently using any traditional method	0.007	0.005	278	145	1.047	0.744	0.000	0.018
Unmet need for spacing	0.216	0.037	278	145	1.496	0.172	0.142	0.290
Unmet need for limiting	0.199	0.025	278	145	1.033	0.125	0.149	0.248
Unmet need total	0.415	0.039	278	145	1.317	0.094	0.337	0.493
Demand satisfied by modern methods (married women)	0.294	0.068	166	87	1.913	0.231	0.158	0.431
Demand satisfied by modern methods (all women)	0.320	0.068	218	111	2.140	0.212	0.184	0.455
Not exposed to any of the ten media sources	0.148	0.028	444	225	1.637	0.187	0.093	0.203
Neonatal mortality (last 0–9 years)	10.994	3.354	700	367	0.844	0.305	4.287	17.702
Post-neonatal mortality (last 0–9 years)	5.060	2.410	699	367	0.911	0.476	0.239	9.881
Infant mortality (last 0–9 years)	16.054	3.763	701	368	0.801	0.234	8.529	23.580
Child mortality (last 0–9 years)	1.576	1.344	700	365	0.886	0.852	0.000	4.264
Under-five mortality (last 0–9 years)	17.606	4.484	701	368	0.847	0.255	8.638	26.573
Stillbirth rate	10.698	8.137	361	186	1.522	0.761	0.000	26.971
Early neonatal rate	12.724	5.804	358	184	0.924	0.456	1.116	24.331
Perinatal mortality rate	23.285	8.627	361	186	1.068	0.371	6.031	40.539
Received ANC from a skilled provider	0.919	0.029	142	73	1.244	0.031	0.862	0.977
4+ ANC visits	0.374	0.070	142	73	1.703	0.187	0.234	0.513
8+ ANC visits	0.013	0.012	142	73	1.314	0.977	0.000	0.038
Took any iron-containing supplements	0.755	0.054	142	73	1.487	0.072	0.646	0.863
Mothers protected against tetanus for last birth	0.643	0.061	142	73	1.512	0.095	0.521	0.766
Delivered in a health facility (live births)	0.668	0.071	148	76	1.774	0.106	0.527	0.809
Delivered by a skilled provider (live births)	0.730	0.065	148	76	1.739	0.089	0.600	0.860
Delivered by C-section (live births)	0.112	0.034	148	76	1.204	0.305	0.044	0.181
Women with postnatal check during first 2 days	0.630	0.059	142	73	1.456	0.094	0.511	0.749
Newborns with postnatal check during first 2 days	0.579	0.065	142	73	1.565	0.113	0.449	0.710
Any problem accessing health care	0.806	0.052	444	225	2.735	0.064	0.702	0.909
Ever had a vaccination card	0.934	0.040	67	34	1.325	0.043	0.854	1.000
Received BCG vaccination	0.913	0.030	67	34	0.884	0.033	0.853	0.973
Received DPT-HepB+Hib vaccination (3 doses)	0.754	0.076	67	34	1.430	0.101	0.601	0.906
Received pneumococcal vaccination (3 doses)	0.722	0.081	67	34	1.463	0.113	0.559	0.885
Received measles vaccination	0.606	0.069	67	34	1.152	0.114	0.468	0.744
Fully vaccinated according to national schedule (12–23 months)	0.116	0.032	67	34	0.834	0.279	0.051	0.181
Received measles/measles-rubella 2 vaccination (24–35 months)	0.572	0.060	61	32	0.955	0.104	0.453	0.691
Fully vaccinated according to national schedule (24–35 months)	0.070	0.042	61	32	1.313	0.604	0.000	0.154
Received HPV vaccination (2 doses) women 15–19	0.146	0.036	97	52	1.008	0.249	0.073	0.218
Sought treatment for diarrhoea	0.818	0.111	18	10	1.251	0.136	0.596	1.000
Treated with ORS	0.838	0.059	18	10	0.751	0.071	0.719	0.956
Exclusive breastfeeding	0.609	0.108	33	16	1.245	0.178	0.393	0.826
Minimum dietary diversity (children 6–23 months)	0.307	0.059	105	56	1.295	0.191	0.190	0.425
Child slept under an ITN last night	0.583	0.064	383	201	2.097	0.110	0.455	0.712
Received 3+ doses of SP/Fansidar	0.021	0.012	142	73	1.023	0.585	0.000	0.046
Child has fever in last 2 weeks	0.061	0.016	350	181	1.162	0.262	0.029	0.094
Child had blood taken from finger/heel	0.459	0.110	23	11	1.059	0.239	0.240	0.679
Child took ACT	0.678	0.128	9	4	0.838	0.189	0.422	0.933
Condom use at last sex (women 15–24)	0.087	0.042	51	27	1.047	0.481	0.003	0.170
Mobile phone ownership	0.570	0.035	444	225	1.474	0.061	0.501	0.639

Table B.29 Sampling errors: Samburu sample, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases: Unweighted (N)	Number of cases: Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
HOUSEHOLDS AND POPULATION								
Primary reliance on clean fuels and technology	0.022	0.007	2,265	554	1.288	0.327	0.007	0.036
Births registered with civil authority	0.620	0.061	346	85	1.975	0.099	0.497	0.742
Ownership of at least one ITN	0.129	0.026	460	109	1.655	0.201	0.077	0.181
Ownership of at least one ITN for every two persons	0.023	0.008	458	109	1.084	0.328	0.008	0.039
At least basic drinking water service	0.326	0.065	2,265	554	2.715	0.200	0.196	0.456
At least basic sanitation service	0.138	0.054	2,265	554	2.928	0.392	0.030	0.246
Using open defecation	0.579	0.063	2,265	554	2.477	0.109	0.453	0.705
Using a handwashing facility with soap and water	0.469	0.032	2,237	547	1.244	0.069	0.405	0.534
WOMEN								
No education	0.410	0.062	395	98	2.496	0.152	0.285	0.534
Secondary education or higher	0.409	0.059	395	98	2.352	0.143	0.292	0.526
Literacy	0.569	0.063	395	98	2.490	0.110	0.444	0.694
Use of the internet in last 12 months	0.266	0.054	395	98	2.411	0.203	0.158	0.374
Total fertility rate (3 years)	4.997	0.322	1,085	268	1.151	0.064	4.353	5.640
Currently pregnant	0.050	0.014	395	98	1.269	0.280	0.022	0.078
Mean number of children ever born to women age 40–49	5.821	0.366	59	13	1.232	0.063	5.090	6.553
Median birth interval	32.396	1.302	240	60	1.847	0.040	29.792	35.001
Ideal number of children	5.401	0.331	385	96	2.463	0.061	4.739	6.062
Total wanted fertility rate (3 years)	4.271	0.314	1,085	268	1.123	0.074	3.642	4.899
Currently using any contraceptive method	0.446	0.057	242	58	1.781	0.128	0.331	0.560
Currently using any modern method	0.374	0.062	242	58	1.969	0.165	0.250	0.497
Currently using pill	0.008	0.004	242	58	0.633	0.466	0.001	0.015
Currently using injectables	0.210	0.053	242	58	2.022	0.255	0.103	0.316
Currently using implants	0.113	0.026	242	58	1.275	0.230	0.061	0.165
Currently using male condoms	0.020	0.007	242	58	0.778	0.352	0.006	0.034
Currently using any traditional method	0.072	0.018	242	58	1.086	0.251	0.036	0.108
Unmet need for spacing	0.147	0.027	242	58	1.169	0.182	0.094	0.200
Unmet need for limiting	0.067	0.018	242	58	1.127	0.270	0.031	0.104
Unmet need total	0.214	0.031	242	58	1.185	0.146	0.152	0.277
Demand satisfied by modern methods (married women)	0.566	0.066	152	38	1.672	0.116	0.434	0.698
Demand satisfied by modern methods (all women)	0.573	0.050	204	52	1.452	0.087	0.473	0.672
Not exposed to any of the ten media sources	0.050	0.012	395	98	1.096	0.239	0.026	0.075
Neonatal mortality (last 0–9 years)	21.998	4.252	594	147	0.715	0.193	13.494	30.503
Post-neonatal mortality (last 0–9 years)	5.970	3.107	590	146	0.912	0.520	0.000	12.184
Infant mortality (last 0–9 years)	27.969	6.539	594	147	0.970	0.234	14.890	41.047
Child mortality (last 0–9 years)	2.601	1.846	572	139	0.823	0.710	0.000	6.293
Under-five mortality (last 0–9 years)	30.497	6.130	594	147	0.874	0.201	18.237	42.757
Stillbirth rate	6.041	4.333	310	78	0.986	0.717	0.000	14.708
Early neonatal rate	19.786	6.785	309	78	0.858	0.343	6.216	33.356
Perinatal mortality rate	25.796	8.938	310	78	0.991	0.346	7.920	43.672
Received ANC from a skilled provider	0.912	0.030	123	31	1.170	0.033	0.852	0.972
4+ ANC visits	0.653	0.048	123	31	1.114	0.074	0.557	0.749
8+ ANC visits	0.056	0.029	123	31	1.415	0.530	0.000	0.115
Took any iron-containing supplements	0.832	0.050	123	31	1.458	0.060	0.733	0.931
Mothers protected against tetanus for last birth	0.631	0.062	123	31	1.418	0.099	0.506	0.755
Delivered in a health facility (live births)	0.682	0.078	130	33	1.911	0.115	0.526	0.839
Delivered by a skilled provider (live births)	0.682	0.078	130	33	1.869	0.115	0.526	0.839
Delivered by C-section (live births)	0.098	0.039	130	33	1.536	0.399	0.020	0.177
Women with postnatal check during first 2 days	0.631	0.068	123	31	1.553	0.108	0.495	0.768
Newborns with postnatal check during first 2 days	0.654	0.071	123	31	1.644	0.109	0.512	0.797
Any problem accessing health care	0.649	0.046	395	98	1.904	0.071	0.557	0.740
Ever had a vaccination card	0.965	0.022	56	14	0.880	0.022	0.922	1.000
Received BCG vaccination	0.955	0.027	56	14	0.968	0.028	0.901	1.000
Received DPT-HepB+Hib vaccination (3 doses)	0.806	0.055	56	14	1.033	0.068	0.695	0.916
Received pneumococcal vaccination (3 doses)	0.798	0.062	56	14	1.152	0.078	0.673	0.923
Received measles vaccination	0.732	0.065	56	14	1.088	0.089	0.602	0.862
Fully vaccinated according to national schedule (12–23 months)	0.260	0.054	56	14	0.856	0.208	0.152	0.369
Received measles/measles-rubella 2 vaccination (24–35 months)	0.441	0.063	56	14	0.967	0.143	0.314	0.567
Fully vaccinated according to national schedule (24–35 months)	0.169	0.042	56	14	0.843	0.247	0.085	0.252
Received HPV vaccination (2 doses) women 15–19	0.310	0.065	82	22	1.253	0.208	0.181	0.440
Sought treatment for diarrhoea	0.742	0.073	25	6	0.809	0.098	0.595	0.888
Treated with ORS	0.491	0.133	25	6	1.289	0.271	0.225	0.757
Exclusive breastfeeding	1.000	0.000	29	7	na	0.000	1.000	1.000
Minimum dietary diversity (children 6–23 months)	0.249	0.062	89	22	1.338	0.249	0.125	0.372
Child slept under an ITN last night	0.100	0.034	350	87	1.725	0.340	0.032	0.168
Received 3+ doses of SP/Fansidar	0.010	0.009	123	31	1.053	0.958	0.000	0.029
Child has fever in last 2 weeks	0.100	0.020	301	75	1.165	0.198	0.060	0.139
Child had blood taken from finger/heel	0.209	0.076	30	8	1.032	0.363	0.057	0.361
Child took ACT	0.000	0.000	2	-	na	na	0.000	0.000
Condom use at last sex (women 15–24)	0.390	0.094	41	11	1.206	0.240	0.203	0.577
Mobile phone ownership	0.652	0.027	395	98	1.134	0.042	0.598	0.707

na = not applicable

Table B.30 Sampling errors: Trans Nzoia sample, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases: Unweighted (N)	Number of cases: Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
HOUSEHOLDS AND POPULATION								
Primary reliance on clean fuels and technology	0.063	0.012	2,013	1,568	1.178	0.188	0.039	0.087
Births registered with civil authority	0.740	0.046	226	173	1.379	0.063	0.647	0.833
Ownership of at least one ITN	0.804	0.022	512	393	1.257	0.027	0.760	0.848
Ownership of at least one ITN for every two persons	0.568	0.025	512	393	1.132	0.044	0.519	0.618
At least basic drinking water service	0.609	0.083	2,013	1,568	3.341	0.136	0.443	0.775
At least basic sanitation service	0.521	0.048	2,013	1,568	1.911	0.092	0.425	0.617
Using open defecation	0.001	0.001	2,013	1,568	1.083	0.990	0.000	0.002
Using a handwashing facility with soap and water	0.910	0.019	2,012	1,567	1.400	0.021	0.871	0.949
WOMEN								
No education	0.013	0.007	498	369	1.329	0.513	0.000	0.027
Secondary education or higher	0.618	0.028	498	369	1.299	0.046	0.561	0.675
Literacy	0.933	0.014	500	370	1.270	0.015	0.904	0.961
Use of the internet in last 12 months	0.541	0.041	500	370	1.837	0.076	0.459	0.623
Total fertility rate (3 years)	3.000	0.274	1,384	1,021	1.011	0.091	2.451	3.548
Currently pregnant	0.041	0.012	500	370	1.310	0.284	0.018	0.064
Mean number of children ever born to women age 40–49	4.304	0.204	107	84	0.975	0.047	3.896	4.713
Median birth interval	54.765	2.732	146	110	0.851	0.050	49.302	60.228
Ideal number of children	3.577	0.075	496	368	0.956	0.021	3.427	3.727
Total wanted fertility rate (3 years)	2.410	0.229	1,384	1,021	0.912	0.095	1.952	2.869
Currently using any contraceptive method	0.567	0.039	264	197	1.278	0.069	0.489	0.645
Currently using any modern method	0.543	0.037	264	197	1.205	0.068	0.469	0.617
Currently using pill	0.022	0.008	264	197	0.889	0.368	0.006	0.038
Currently using injectables	0.251	0.030	264	197	1.115	0.119	0.192	0.311
Currently using implants	0.185	0.023	264	197	0.978	0.126	0.139	0.232
Currently using male condoms	0.006	0.005	264	197	0.989	0.781	0.000	0.016
Currently using any traditional method	0.024	0.009	264	197	0.965	0.378	0.006	0.042
Unmet need for spacing	0.086	0.018	264	197	1.053	0.211	0.050	0.123
Unmet need for limiting	0.110	0.034	264	197	1.764	0.310	0.042	0.179
Unmet need total	0.197	0.040	264	197	1.608	0.201	0.118	0.276
Demand satisfied by modern methods (married women)	0.711	0.045	206	150	1.399	0.063	0.621	0.801
Demand satisfied by modern methods (all women)	0.704	0.040	257	188	1.382	0.056	0.624	0.783
Not exposed to any of the ten media sources	0.110	0.018	500	370	1.297	0.165	0.074	0.146
Neonatal mortality (last 0–9 years)	9.758	4.396	438	330	0.808	0.450	0.966	18.550
Post-neonatal mortality (last 0–9 years)	7.237	4.191	437	330	1.050	0.579	0.000	15.618
Infant mortality (last 0–9 years)	16.995	5.913	438	330	0.896	0.348	5.170	28.820
Child mortality (last 0–9 years)	4.984	4.250	437	331	1.197	0.853	0.000	13.484
Under-five mortality (last 0–9 years)	21.895	8.382	439	331	1.073	0.383	5.130	38.659
Stillbirth rate	16.063	11.804	214	156	1.367	0.735	0.000	39.670
Early neonatal rate	5.887	5.774	210	153	1.083	0.981	0.000	17.434
Perinatal mortality rate	21.855	12.683	214	156	1.262	0.580	0.000	47.220
Received ANC from a skilled provider	0.984	0.010	78	57	0.720	0.010	0.964	1.000
4+ ANC visits	0.603	0.072	78	57	1.280	0.119	0.460	0.747
8+ ANC visits	0.114	0.047	78	57	1.303	0.417	0.019	0.208
Took any iron-containing supplements	0.772	0.045	78	57	0.932	0.058	0.682	0.861
Mothers protected against tetanus for last birth	0.680	0.058	78	57	1.099	0.086	0.563	0.797
Delivered in a health facility (live births)	0.858	0.032	81	59	0.812	0.037	0.794	0.922
Delivered by a skilled provider (live births)	0.939	0.026	81	59	0.984	0.028	0.886	0.992
Delivered by C-section (live births)	0.058	0.030	81	59	1.150	0.522	0.000	0.118
Women with postnatal check during first 2 days	0.694	0.069	78	57	1.310	0.100	0.556	0.832
Newborns with postnatal check during first 2 days	0.815	0.064	78	57	1.442	0.079	0.687	0.944
Any problem accessing health care	0.767	0.025	500	370	1.309	0.032	0.717	0.816
Ever had a vaccination card	0.902	0.052	31	22	0.960	0.058	0.797	1.000
Received BCG vaccination	0.922	0.050	31	22	1.017	0.054	0.822	1.000
Received DPT-HepB+Hib vaccination (3 doses)	0.883	0.078	31	22	1.337	0.089	0.726	1.000
Received pneumococcal vaccination (3 doses)	0.883	0.058	31	22	0.981	0.065	0.767	0.998
Received measles vaccination	0.869	0.055	31	22	0.890	0.063	0.760	0.979
Fully vaccinated according to national schedule (12–23 months)	0.417	0.104	31	22	1.153	0.249	0.210	0.625
Received measles/measles-rubella 2 vaccination (24–35 months)	0.678	0.078	51	37	1.156	0.116	0.521	0.835
Fully vaccinated according to national schedule (24–35 months)	0.376	0.089	51	37	1.261	0.237	0.198	0.554
Received HPV vaccination (2 doses) women 15–19	0.356	0.062	113	84	1.369	0.175	0.232	0.481
Sought treatment for diarrhoea	0.626	0.201	7	5	1.032	0.322	0.223	1.000
Treated with ORS	0.502	0.206	7	5	1.022	0.411	0.090	0.915
Exclusive breastfeeding	0.703	0.097	25	19	1.040	0.138	0.508	0.897
Minimum dietary diversity (children 6–23 months)	0.280	0.073	52	37	1.150	0.259	0.135	0.426
Child slept under an ITN last night	0.734	0.053	229	176	1.590	0.072	0.628	0.840
Received 3+ doses of SP/Fansidar	0.000	0.000	78	57	na	na	0.000	0.000
Child has fever in last 2 weeks	0.096	0.023	205	150	1.110	0.235	0.051	0.141
Child had blood taken from finger/heel	0.292	0.112	17	14	1.080	0.382	0.069	0.516
Child took ACT	0.456	0.364	2	2	1.190	0.798	0.000	1.000
Condom use at last sex (women 15–24)	0.113	0.061	32	22	1.065	0.538	0.000	0.234
Mobile phone ownership	0.748	0.027	500	370	1.376	0.036	0.694	0.801

na = not applicable

Table B.31 Sampling errors: Uasin Gishu sample, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases: Unweighted (N)	Number of cases: Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
HOUSEHOLDS AND POPULATION								
Primary reliance on clean fuels and technology	0.230	0.029	2,226	2,677	1.611	0.124	0.173	0.288
Births registered with civil authority	0.919	0.019	254	307	1.005	0.021	0.880	0.958
Ownership of at least one ITN	0.420	0.043	630	794	2.185	0.103	0.334	0.507
Ownership of at least one ITN for every two persons	0.255	0.028	623	780	1.611	0.111	0.199	0.311
At least basic drinking water service	0.897	0.032	2,226	2,677	2.073	0.035	0.834	0.960
At least basic sanitation service	0.510	0.040	2,226	2,677	1.685	0.078	0.431	0.590
Using open defecation	0.005	0.003	2,226	2,677	0.845	0.647	0.000	0.011
Using a handwashing facility with soap and water	0.439	0.035	2,225	2,676	1.484	0.079	0.369	0.509
WOMEN								
No education	0.005	0.003	525	667	1.069	0.665	0.000	0.011
Secondary education or higher	0.797	0.015	525	667	0.872	0.019	0.766	0.828
Literacy	0.957	0.014	525	667	1.595	0.015	0.928	0.985
Use of the internet in last 12 months	0.717	0.026	525	667	1.299	0.036	0.666	0.768
Total fertility rate (3 years)	2.961	0.326	1,490	1,901	1.156	0.110	2.309	3.612
Currently pregnant	0.044	0.008	525	667	0.871	0.178	0.028	0.059
Mean number of children ever born to women age 40–49	3.949	0.207	92	118	1.038	0.052	3.535	4.364
Median birth interval	53.748	3.581	159	199	0.712	0.067	46.586	60.910
Ideal number of children	3.101	0.073	524	666	1.234	0.023	2.956	3.247
Total wanted fertility rate (3 years)	2.695	0.269	1,490	1,901	0.992	0.100	2.156	3.234
Currently using any contraceptive method	0.631	0.026	261	335	0.870	0.041	0.579	0.683
Currently using any modern method	0.565	0.031	261	335	1.004	0.055	0.503	0.627
Currently using pill	0.049	0.012	261	335	0.885	0.242	0.025	0.073
Currently using injectables	0.166	0.035	261	335	1.517	0.211	0.096	0.236
Currently using implants	0.276	0.024	261	335	0.880	0.088	0.227	0.325
Currently using male condoms	0.019	0.009	261	335	1.029	0.462	0.001	0.036
Currently using any traditional method	0.066	0.016	261	335	1.004	0.233	0.035	0.097
Unmet need for spacing	0.094	0.026	261	335	1.446	0.280	0.041	0.146
Unmet need for limiting	0.100	0.017	261	335	0.930	0.173	0.065	0.135
Unmet need total	0.194	0.030	261	335	1.210	0.153	0.134	0.253
Demand satisfied by modern methods (married women)	0.685	0.041	213	277	1.296	0.060	0.602	0.767
Demand satisfied by modern methods (all women)	0.688	0.027	302	390	1.017	0.039	0.634	0.742
Not exposed to any of the ten media sources	0.091	0.015	525	667	1.183	0.164	0.061	0.121
Neonatal mortality (last 0–9 years)	16.255	5.976	467	583	1.003	0.368	4.303	28.208
Post-neonatal mortality (last 0–9 years)	13.375	6.167	470	588	1.122	0.461	1.042	25.709
Infant mortality (last 0–9 years)	29.631	6.515	468	584	0.844	0.220	16.602	42.660
Child mortality (last 0–9 years)	11.859	6.792	447	571	1.280	0.573	0.000	25.443
Under-five mortality (last 0–9 years)	41.139	10.663	468	584	1.167	0.259	19.813	62.464
Stillbirth rate	18.809	9.137	261	321	1.076	0.486	0.535	37.083
Early neonatal rate	20.257	9.674	256	315	1.093	0.478	0.909	39.606
Perinatal mortality rate	38.685	11.055	261	321	0.929	0.286	16.575	60.796
Received ANC from a skilled provider	1.000	0.000	105	123	na	0.000	1.000	1.000
4+ ANC visits	0.820	0.036	105	123	0.946	0.043	0.749	0.891
8+ ANC visits	0.145	0.050	105	123	1.433	0.343	0.046	0.245
Took any iron-containing supplements	0.913	0.032	105	123	1.151	0.035	0.849	0.977
Mothers protected against tetanus for last birth	0.755	0.031	105	123	0.746	0.042	0.692	0.818
Delivered in a health facility (live births)	0.990	0.010	106	124	1.002	0.010	0.969	1.000
Delivered by a skilled provider (live births)	0.990	0.010	106	124	1.002	0.010	0.969	1.000
Delivered by C-section (live births)	0.104	0.022	106	124	0.701	0.209	0.061	0.148
Women with postnatal check during first 2 days	0.867	0.042	105	123	1.252	0.048	0.783	0.951
Newborns with postnatal check during first 2 days	0.907	0.027	105	123	0.958	0.030	0.852	0.961
Any problem accessing health care	0.419	0.028	525	667	1.298	0.067	0.363	0.475
Ever had a vaccination card	1.000	0.000	50	61	na	0.000	1.000	1.000
Received BCG vaccination	0.984	0.016	50	61	0.894	0.016	0.952	1.000
Received DPT-HepB+Hib vaccination (3 doses)	0.806	0.088	50	61	1.530	0.109	0.630	0.982
Received pneumococcal vaccination (3 doses)	0.887	0.054	50	61	1.166	0.060	0.780	0.994
Received measles vaccination	0.900	0.056	50	61	1.278	0.062	0.789	1.000
Fully vaccinated according to national schedule (12–23 months)	0.432	0.073	50	61	1.012	0.169	0.286	0.578
Received measles/measles-rubella 2 vaccination (24–35 months)	0.773	0.058	45	57	0.930	0.075	0.657	0.890
Fully vaccinated according to national schedule (24–35 months)	0.254	0.080	45	57	1.227	0.316	0.093	0.414
Received HPV vaccination (2 doses) women 15–19	0.465	0.064	95	112	1.237	0.137	0.337	0.592
Sought treatment for diarrhoea	0.612	0.118	22	24	1.045	0.193	0.375	0.848
Treated with ORS	0.399	0.102	22	24	0.900	0.256	0.195	0.603
Exclusive breastfeeding	0.615	0.095	19	22	0.836	0.155	0.425	0.806
Minimum dietary diversity (children 6–23 months)	0.547	0.073	82	96	1.313	0.133	0.401	0.693
Child slept under an ITN last night	0.446	0.042	251	302	1.159	0.094	0.362	0.529
Received 3+ doses of SP/Fansidar	0.030	0.016	105	123	0.989	0.555	0.000	0.063
Child has fever in last 2 weeks	0.137	0.025	249	304	1.077	0.183	0.087	0.187
Child had blood taken from finger/heel	0.056	0.042	35	42	1.036	0.746	0.000	0.140
Child took ACT	1.000	0.000	1	1	na	0.000	1.000	1.000
Condom use at last sex (women 15–24)	0.461	0.079	57	74	1.188	0.172	0.302	0.619
Mobile phone ownership	0.842	0.019	525	667	1.214	0.023	0.804	0.881

na = not applicable

Table B.32 Sampling errors: Elgeyo-Marakwet sample, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases: Unweighted (N)	Number of cases: Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
HOUSEHOLDS AND POPULATION								
Primary reliance on clean fuels and technology	0.060	0.024	2,054	848	2.738	0.405	0.011	0.109
Births registered with civil authority	0.877	0.036	235	102	1.449	0.041	0.805	0.948
Ownership of at least one ITN	0.350	0.046	518	209	2.166	0.130	0.259	0.442
Ownership of at least one ITN for every two persons	0.171	0.039	514	207	2.340	0.229	0.093	0.249
At least basic drinking water service	0.718	0.066	2,054	848	2.770	0.091	0.587	0.849
At least basic sanitation service	0.475	0.052	2,054	848	1.983	0.109	0.372	0.578
Using open defecation	0.020	0.011	2,054	848	1.681	0.562	0.000	0.043
Using a handwashing facility with soap and water	0.611	0.064	2,054	848	2.528	0.105	0.483	0.739
WOMEN								
No education	0.002	0.002	436	176	0.998	1.015	0.000	0.007
Secondary education or higher	0.727	0.030	436	176	1.418	0.042	0.666	0.788
Literacy	0.970	0.013	439	178	1.535	0.013	0.945	0.995
Use of the internet in last 12 months	0.430	0.055	439	178	2.303	0.127	0.321	0.540
Total fertility rate (3 years)	2.887	0.306	1,228	497	1.071	0.106	2.275	3.499
Currently pregnant	0.035	0.010	439	178	1.092	0.276	0.016	0.054
Mean number of children ever born to women age 40–49	5.029	0.269	78	32	1.168	0.054	4.491	5.568
Median birth interval	36.467	1.574	141	63	1.119	0.043	33.318	39.616
Ideal number of children	3.816	0.096	439	178	1.205	0.025	3.624	4.009
Total wanted fertility rate (3 years)	2.585	0.285	1,228	497	1.096	0.110	2.015	3.155
Currently using any contraceptive method	0.693	0.030	229	92	0.975	0.043	0.633	0.752
Currently using any modern method	0.633	0.019	229	92	0.603	0.030	0.594	0.671
Currently using pill	0.035	0.012	229	92	1.019	0.356	0.010	0.059
Currently using injectables	0.194	0.031	229	92	1.173	0.158	0.133	0.256
Currently using implants	0.306	0.032	229	92	1.047	0.104	0.242	0.370
Currently using male condoms	0.018	0.008	229	92	0.892	0.437	0.002	0.034
Currently using any traditional method	0.060	0.015	229	92	0.977	0.256	0.029	0.091
Unmet need for spacing	0.028	0.010	229	92	0.915	0.355	0.008	0.048
Unmet need for limiting	0.105	0.020	229	92	0.972	0.188	0.066	0.145
Unmet need total	0.134	0.020	229	92	0.887	0.149	0.094	0.174
Demand satisfied by modern methods (married women)	0.766	0.022	189	76	0.715	0.029	0.722	0.810
Demand satisfied by modern methods (all women)	0.723	0.020	261	107	0.730	0.028	0.683	0.763
Not exposed to any of the ten media sources	0.072	0.019	439	178	1.535	0.264	0.034	0.110
Neonatal mortality (last 0–9 years)	8.082	4.071	418	178	0.937	0.504	0.000	16.225
Post-neonatal mortality (last 0–9 years)	15.382	5.624	418	178	0.905	0.366	4.134	26.630
Infant mortality (last 0–9 years)	23.465	8.056	418	178	1.027	0.343	7.353	39.576
Child mortality (last 0–9 years)	0.000	0.000	434	184	na	na	0.000	0.000
Under-five mortality (last 0–9 years)	23.465	8.056	418	178	1.027	0.343	7.353	39.576
Stillbirth rate	5.242	5.206	205	87	1.057	0.993	0.000	15.654
Early neonatal rate	0.000	0.000	204	87	na	na	0.000	0.000
Perinatal mortality rate	5.242	5.206	205	87	1.057	0.993	0.000	15.654
Received ANC from a skilled provider	0.965	0.024	75	32	1.101	0.024	0.918	1.000
4+ ANC visits	0.536	0.060	75	32	1.040	0.112	0.416	0.657
8+ ANC visits	0.078	0.028	75	32	0.914	0.365	0.021	0.135
Took any iron-containing supplements	0.913	0.026	75	32	0.801	0.029	0.861	0.965
Mothers protected against tetanus for last birth	0.646	0.063	75	32	1.134	0.098	0.520	0.773
Delivered in a health facility (live births)	0.951	0.025	76	33	1.037	0.026	0.901	1.000
Delivered by a skilled provider (live births)	0.951	0.025	76	33	1.037	0.026	0.901	1.000
Delivered by C-section (live births)	0.122	0.031	76	33	0.854	0.256	0.059	0.184
Women with postnatal check during first 2 days	0.905	0.040	75	32	1.155	0.044	0.826	0.984
Newborns with postnatal check during first 2 days	0.919	0.028	75	32	0.884	0.030	0.863	0.975
Any problem accessing health care	0.250	0.027	439	178	1.313	0.109	0.196	0.304
Ever had a vaccination card	1.000	0.000	39	16	na	0.000	1.000	1.000
Received BCG vaccination	0.969	0.032	39	16	1.145	0.033	0.905	1.000
Received DPT-HepB+Hib vaccination (3 doses)	0.936	0.046	39	16	1.181	0.049	0.844	1.000
Received pneumococcal vaccination (3 doses)	0.936	0.046	39	16	1.181	0.049	0.844	1.000
Received measles vaccination	0.940	0.033	39	16	0.877	0.035	0.874	1.000
Fully vaccinated according to national schedule (12–23 months)	0.520	0.064	39	16	0.798	0.122	0.393	0.647
Received measles/measles-rubella 2 vaccination (24–35 months)	0.718	0.057	35	14	0.738	0.079	0.605	0.831
Fully vaccinated according to national schedule (24–35 months)	0.429	0.081	35	14	0.959	0.189	0.267	0.591
Received HPV vaccination (2 doses) women 15–19	0.454	0.059	88	35	1.096	0.129	0.337	0.572
Sought treatment for diarrhoea	0.592	0.151	19	8	1.262	0.256	0.289	0.894
Treated with ORS	0.412	0.149	19	8	1.288	0.360	0.115	0.709
Exclusive breastfeeding	1.000	0.000	18	8	na	0.000	1.000	1.000
Minimum dietary diversity (children 6–23 months)	0.613	0.060	57	25	0.927	0.098	0.493	0.734
Child slept under an ITN last night	0.353	0.046	235	102	1.299	0.129	0.262	0.444
Received 3+ doses of SP/Fansidar	0.061	0.026	75	32	0.938	0.426	0.009	0.114
Child has fever in last 2 weeks	0.068	0.021	203	86	1.238	0.311	0.026	0.110
Child had blood taken from finger/heel	0.411	0.128	14	6	0.991	0.313	0.154	0.667
Child took ACT	0.000	0.000	1	-	na	na	0.000	0.000
Condom use at last sex (women 15–24)	0.622	0.075	48	22	1.063	0.121	0.471	0.772
Mobile phone ownership	0.748	0.026	439	178	1.266	0.035	0.695	0.800

na = not applicable

Table B.33 Sampling errors: Nandi sample, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases: Unweighted (N)	Number of cases: Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
HOUSEHOLDS AND POPULATION								
Primary reliance on clean fuels and technology	0.052	0.010	2,024	1,550	1.187	0.200	0.031	0.073
Births registered with civil authority	0.714	0.039	230	176	1.203	0.055	0.636	0.793
Ownership of at least one ITN	0.538	0.037	554	414	1.740	0.069	0.464	0.612
Ownership of at least one ITN for every two persons	0.323	0.032	554	414	1.608	0.099	0.259	0.387
At least basic drinking water service	0.639	0.056	2,024	1,550	2.364	0.087	0.528	0.750
At least basic sanitation service	0.577	0.041	2,024	1,550	1.729	0.072	0.494	0.660
Using open defecation	0.024	0.022	2,024	1,550	2.995	0.928	0.000	0.068
Using a handwashing facility with soap and water	0.566	0.058	1,909	1,468	2.298	0.103	0.450	0.682
WOMEN								
No education	0.003	0.002	480	361	0.924	0.791	0.000	0.007
Secondary education or higher	0.645	0.025	480	361	1.128	0.038	0.596	0.694
Literacy	0.959	0.015	480	361	1.662	0.016	0.929	0.989
Use of the internet in last 12 months	0.416	0.040	480	361	1.756	0.095	0.337	0.496
Total fertility rate (3 years)	3.371	0.250	1,340	1,002	0.956	0.074	2.872	3.870
Currently pregnant	0.024	0.005	480	361	0.748	0.219	0.013	0.034
Mean number of children ever born to women age 40–49	4.104	0.221	99	73	1.244	0.054	3.661	4.547
Median birth interval	49.281	3.254	152	114	0.999	0.066	42.773	55.789
Ideal number of children	2.888	0.061	477	359	0.863	0.021	2.766	3.011
Total wanted fertility rate (3 years)	2.286	0.176	1,340	1,002	0.866	0.077	1.933	2.639
Currently using any contraceptive method	0.559	0.045	250	188	1.416	0.080	0.470	0.649
Currently using any modern method	0.555	0.045	250	188	1.440	0.082	0.464	0.646
Currently using pill	0.036	0.010	250	188	0.818	0.267	0.017	0.056
Currently using injectables	0.190	0.034	250	188	1.380	0.181	0.121	0.259
Currently using implants	0.255	0.042	250	188	1.514	0.164	0.171	0.339
Currently using male condoms	0.009	0.006	250	188	1.055	0.686	0.000	0.022
Currently using any traditional method	0.005	0.005	250	188	1.061	0.990	0.000	0.014
Unmet need for spacing	0.079	0.015	250	188	0.870	0.188	0.049	0.109
Unmet need for limiting	0.121	0.020	250	188	0.946	0.162	0.082	0.160
Unmet need total	0.200	0.019	250	188	0.737	0.093	0.162	0.237
Demand satisfied by modern methods (married women)	0.731	0.026	192	143	0.793	0.035	0.680	0.782
Demand satisfied by modern methods (all women)	0.683	0.025	259	192	0.864	0.037	0.633	0.733
Not exposed to any of the ten media sources	0.240	0.033	480	361	1.701	0.138	0.174	0.307
Neonatal mortality (last 0–9 years)	11.009	6.020	438	328	1.193	0.547	0.000	23.050
Post-neonatal mortality (last 0–9 years)	20.050	10.396	435	325	0.979	0.519	0.000	40.843
Infant mortality (last 0–9 years)	31.059	10.617	438	328	0.918	0.342	9.825	52.293
Child mortality (last 0–9 years)	0.000	0.000	442	329	na	na	0.000	0.000
Under-five mortality (last 0–9 years)	31.059	10.617	438	328	0.918	0.342	9.825	52.293
Stillbirth rate	20.529	11.891	226	170	1.151	0.579	0.000	44.311
Early neonatal rate	12.097	6.919	221	167	0.945	0.572	0.000	25.934
Perinatal mortality rate	32.379	16.072	226	170	1.296	0.496	0.235	64.522
Received ANC from a skilled provider	1.000	0.000	98	75	na	0.000	1.000	1.000
4+ ANC visits	0.619	0.042	98	75	0.858	0.068	0.534	0.703
8+ ANC visits	0.044	0.025	98	75	1.205	0.571	0.000	0.094
Took any iron-containing supplements	0.697	0.058	98	75	1.233	0.083	0.582	0.813
Mothers protected against tetanus for last birth	0.697	0.052	98	75	1.115	0.075	0.593	0.802
Delivered in a health facility (live births)	0.891	0.033	101	77	0.990	0.037	0.826	0.956
Delivered by a skilled provider (live births)	0.902	0.032	101	77	1.000	0.035	0.838	0.965
Delivered by C-section (live births)	0.201	0.036	101	77	0.903	0.178	0.130	0.273
Women with postnatal check during first 2 days	0.722	0.043	98	75	0.947	0.060	0.636	0.808
Newborns with postnatal check during first 2 days	0.596	0.056	98	75	1.115	0.093	0.485	0.707
Any problem accessing health care	0.405	0.039	480	361	1.727	0.096	0.327	0.483
Ever had a vaccination card	1.000	0.000	57	43	na	0.000	1.000	1.000
Received BCG vaccination	0.954	0.036	57	43	1.284	0.037	0.883	1.000
Received DPT-HepB+Hib vaccination (3 doses)	0.854	0.045	57	43	0.956	0.052	0.764	0.943
Received pneumococcal vaccination (3 doses)	0.915	0.038	57	43	1.038	0.042	0.838	0.991
Received measles vaccination	0.896	0.043	57	43	1.068	0.048	0.809	0.982
Fully vaccinated according to national schedule (12–23 months)	0.415	0.071	57	43	1.084	0.170	0.274	0.556
Received measles/measles-rubella 2 vaccination (24–35 months)	0.608	0.079	37	27	0.966	0.129	0.451	0.765
Fully vaccinated according to national schedule (24–35 months)	0.263	0.072	37	27	0.983	0.274	0.119	0.407
Received HPV vaccination (2 doses) women 15–19	0.366	0.040	109	86	0.871	0.110	0.285	0.446
Sought treatment for diarrhoea	0.638	0.066	11	8	0.447	0.103	0.507	0.770
Treated with ORS	0.397	0.144	11	8	0.956	0.362	0.109	0.684
Exclusive breastfeeding	0.577	0.115	24	19	1.113	0.200	0.347	0.808
Minimum dietary diversity (children 6–23 months)	0.504	0.060	72	55	1.011	0.119	0.384	0.625
Child slept under an ITN last night	0.493	0.048	229	175	1.353	0.096	0.398	0.588
Received 3+ doses of SP/Fansidar	0.094	0.034	98	75	1.148	0.362	0.026	0.163
Child has fever in last 2 weeks	0.107	0.015	213	161	0.707	0.140	0.077	0.137
Child had blood taken from finger/heel	0.398	0.100	23	17	0.979	0.250	0.199	0.598
Child took ACT	0.745	0.160	4	3	0.701	0.215	0.425	1.000
Condom use at last sex (women 15–24)	0.446	0.071	43	34	0.921	0.158	0.305	0.587
Mobile phone ownership	0.726	0.037	480	361	1.798	0.051	0.652	0.799

na = not applicable

Table B.34 Sampling errors: Baringo sample, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases: Unweighted (N)	Number of cases: Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
HOUSEHOLDS AND POPULATION								
Primary reliance on clean fuels and technology	0.035	0.018	2,096	1,012	2.640	0.517	0.000	0.071
Births registered with civil authority	0.783	0.063	270	128	2.157	0.081	0.657	0.910
Ownership of at least one ITN	0.505	0.051	507	242	2.298	0.102	0.402	0.607
Ownership of at least one ITN for every two persons	0.302	0.048	506	242	2.338	0.159	0.206	0.398
At least basic drinking water service	0.383	0.060	2,096	1,012	2.417	0.158	0.262	0.504
At least basic sanitation service	0.401	0.055	2,096	1,012	2.104	0.137	0.291	0.510
Using open defecation	0.267	0.082	2,096	1,012	3.365	0.309	0.102	0.431
Using a handwashing facility with soap and water	0.402	0.057	2,079	1,004	2.202	0.141	0.289	0.516
WOMEN								
No education	0.085	0.056	360	185	3.709	0.654	0.000	0.196
Secondary education or higher	0.663	0.049	360	185	1.949	0.074	0.565	0.761
Literacy	0.907	0.056	364	187	3.582	0.061	0.796	1.000
Use of the internet in last 12 months	0.428	0.052	364	187	2.010	0.122	0.324	0.533
Total fertility rate (3 years)	4.303	0.452	1,004	516	1.536	0.105	3.399	5.207
Currently pregnant	0.062	0.013	364	187	1.057	0.216	0.035	0.089
Mean number of children ever born to women age 40–49	5.748	0.343	54	28	1.083	0.060	5.061	6.434
Median birth interval	38.270	2.928	179	89	1.474	0.077	32.414	44.127
Ideal number of children	4.572	0.338	360	186	2.764	0.074	3.895	5.249
Total wanted fertility rate (3 years)	3.671	0.448	1,004	516	1.591	0.122	2.774	4.567
Currently using any contraceptive method	0.666	0.065	212	109	1.993	0.098	0.535	0.796
Currently using any modern method	0.477	0.055	212	109	1.587	0.115	0.368	0.587
Currently using pill	0.029	0.013	212	109	1.095	0.436	0.004	0.054
Currently using injectables	0.143	0.030	212	109	1.240	0.209	0.083	0.203
Currently using implants	0.244	0.050	212	109	1.692	0.206	0.143	0.344
Currently using male condoms	0.027	0.010	212	109	0.899	0.369	0.007	0.048
Currently using any traditional method	0.188	0.036	212	109	1.326	0.190	0.117	0.260
Unmet need for spacing	0.093	0.019	212	109	0.957	0.206	0.055	0.131
Unmet need for limiting	0.043	0.010	212	109	0.708	0.230	0.023	0.063
Unmet need total	0.136	0.022	212	109	0.931	0.162	0.092	0.180
Demand satisfied by modern methods (married women)	0.595	0.039	166	87	1.034	0.066	0.517	0.674
Demand satisfied by modern methods (all women)	0.593	0.044	219	116	1.325	0.073	0.506	0.680
Not exposed to any of the ten media sources	0.089	0.036	364	187	2.390	0.405	0.017	0.160
Neonatal mortality (last 0–9 years)	23.005	12.330	470	240	1.642	0.536	0.000	47.666
Post-neonatal mortality (last 0–9 years)	3.355	2.402	472	242	0.888	0.716	0.000	8.158
Infant mortality (last 0–9 years)	26.360	11.987	470	240	1.519	0.455	2.386	50.334
Child mortality (last 0–9 years)	3.722	2.641	465	239	0.936	0.710	0.000	9.005
Under-five mortality (last 0–9 years)	29.984	11.774	470	240	1.418	0.393	6.436	53.533
Stillbirth rate	10.495	6.939	243	123	1.066	0.661	0.000	24.372
Early neonatal rate	27.325	14.524	242	122	1.387	0.532	0.000	56.373
Perinatal mortality rate	37.639	16.102	243	123	1.335	0.428	5.436	69.843
Received ANC from a skilled provider	0.967	0.021	103	53	1.196	0.022	0.925	1.000
4+ ANC visits	0.615	0.077	103	53	1.594	0.126	0.460	0.770
8+ ANC visits	0.032	0.022	103	53	1.259	0.688	0.000	0.076
Took any iron-containing supplements	0.918	0.036	103	53	1.302	0.039	0.846	0.989
Mothers protected against tetanus for last birth	0.575	0.047	103	53	0.965	0.082	0.481	0.670
Delivered in a health facility (live births)	0.819	0.090	107	55	2.295	0.109	0.639	0.998
Delivered by a skilled provider (live births)	0.819	0.090	107	55	2.295	0.109	0.639	0.998
Delivered by C-section (live births)	0.144	0.039	107	55	1.106	0.270	0.066	0.222
Women with postnatal check during first 2 days	0.832	0.063	103	53	1.697	0.076	0.706	0.959
Newborns with postnatal check during first 2 days	0.841	0.077	103	53	2.090	0.091	0.688	0.995
Any problem accessing health care	0.635	0.060	364	187	2.364	0.095	0.515	0.756
Ever had a vaccination card	0.980	0.017	46	25	0.834	0.017	0.946	1.000
Received BCG vaccination	0.980	0.020	46	25	1.004	0.021	0.939	1.000
Received DPT-HepB+Hib vaccination (3 doses)	0.955	0.027	46	25	0.895	0.028	0.901	1.000
Received pneumococcal vaccination (3 doses)	0.920	0.031	46	25	0.792	0.033	0.859	0.982
Received measles vaccination	0.973	0.022	46	25	0.928	0.022	0.930	1.000
Fully vaccinated according to national schedule (12–23 months)	0.633	0.101	46	25	1.457	0.159	0.431	0.835
Received measles/measles-rubella 2 vaccination (24–35 months)	0.663	0.092	46	22	1.269	0.138	0.480	0.847
Fully vaccinated according to national schedule (24–35 months)	0.278	0.085	46	22	1.240	0.306	0.108	0.448
Received HPV vaccination (2 doses) women 15–19	0.622	0.066	77	39	1.176	0.105	0.491	0.754
Sought treatment for diarrhoea	0.617	0.052	22	11	0.498	0.084	0.513	0.721
Treated with ORS	0.571	0.063	22	11	0.592	0.110	0.445	0.697
Exclusive breastfeeding	0.555	0.060	30	14	0.660	0.109	0.435	0.676
Minimum dietary diversity (children 6–23 months)	0.544	0.060	71	38	1.003	0.110	0.424	0.663
Child slept under an ITN last night	0.403	0.059	269	127	1.612	0.146	0.285	0.520
Received 3+ doses of SP/Fansidar	0.000	0.000	103	53	na	na	0.000	0.000
Child has fever in last 2 weeks	0.107	0.026	234	118	1.239	0.241	0.055	0.158
Child had blood taken from finger/heel	0.488	0.084	23	13	0.833	0.172	0.320	0.656
Child took ACT	1.000	0.000	4	3	na	0.000	1.000	1.000
Condom use at last sex (women 15–24)	0.396	0.060	38	21	0.748	0.151	0.276	0.515
Mobile phone ownership	0.724	0.048	364	187	2.019	0.066	0.629	0.819

na = not applicable

Table B.35 Sampling errors: Laikipia sample, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases: Unweighted (N)	Number of cases: Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
HOUSEHOLDS AND POPULATION								
Primary reliance on clean fuels and technology	0.175	0.032	1,539	931	1.745	0.180	0.112	0.238
Births registered with civil authority	0.655	0.034	175	105	0.830	0.051	0.588	0.723
Ownership of at least one ITN	0.154	0.025	495	293	1.546	0.163	0.103	0.204
Ownership of at least one ITN for every two persons	0.099	0.020	493	292	1.466	0.200	0.059	0.138
At least basic drinking water service	0.683	0.073	1,539	931	2.860	0.107	0.537	0.829
At least basic sanitation service	0.391	0.068	1,539	931	2.657	0.174	0.255	0.528
Using open defecation	0.090	0.065	1,539	931	4.072	0.727	0.000	0.220
Using a handwashing facility with soap and water	0.306	0.040	1,539	931	1.696	0.131	0.226	0.386
WOMEN								
No education	0.084	0.054	319	194	3.433	0.645	0.000	0.193
Secondary education or higher	0.664	0.047	319	194	1.780	0.071	0.570	0.759
Literacy	0.936	0.046	321	195	3.317	0.049	0.844	1.000
Use of the internet in last 12 months	0.534	0.051	321	195	1.819	0.095	0.432	0.636
Total fertility rate (3 years)	3.719	0.433	918	559	1.414	0.116	2.854	4.584
Currently pregnant	0.028	0.009	321	195	0.970	0.318	0.010	0.046
Mean number of children ever born to women age 40–49	3.787	0.227	83	49	1.229	0.060	3.334	4.241
Median birth interval	46.925	4.708	125	76	1.298	0.100	37.510	56.340
Ideal number of children	3.758	0.248	321	195	2.331	0.066	3.261	4.255
Total wanted fertility rate (3 years)	3.035	0.440	918	559	1.579	0.145	2.156	3.915
Currently using any contraceptive method	0.625	0.071	167	105	1.872	0.113	0.483	0.767
Currently using any modern method	0.558	0.060	167	105	1.553	0.108	0.438	0.678
Currently using pill	0.137	0.037	167	105	1.375	0.269	0.063	0.211
Currently using injectables	0.266	0.036	167	105	1.055	0.136	0.193	0.338
Currently using implants	0.048	0.016	167	105	0.950	0.328	0.017	0.080
Currently using male condoms	0.010	0.007	167	105	0.891	0.691	0.000	0.024
Currently using any traditional method	0.067	0.021	167	105	1.087	0.316	0.025	0.109
Unmet need for spacing	0.028	0.016	167	105	1.236	0.571	0.000	0.059
Unmet need for limiting	0.130	0.031	167	105	1.170	0.235	0.069	0.191
Unmet need total	0.158	0.037	167	105	1.295	0.233	0.084	0.231
Demand satisfied by modern methods (married women)	0.713	0.048	131	82	1.211	0.067	0.617	0.809
Demand satisfied by modern methods (all women)	0.701	0.043	169	103	1.216	0.061	0.615	0.787
Not exposed to any of the ten media sources	0.108	0.041	321	195	2.325	0.376	0.027	0.189
Neonatal mortality (last 0–9 years)	29.521	11.277	322	200	1.072	0.382	6.967	52.076
Post-neonatal mortality (last 0–9 years)	19.673	8.812	317	197	0.982	0.448	2.049	37.298
Infant mortality (last 0–9 years)	49.195	16.278	322	200	1.234	0.331	16.640	81.750
Child mortality (last 0–9 years)	1.568	1.588	304	190	0.973	1.013	0.000	4.745
Under-five mortality (last 0–9 years)	50.686	16.630	323	201	1.266	0.328	17.426	83.946
Stillbirth rate	9.578	6.824	177	108	0.932	0.712	0.000	23.226
Early neonatal rate	36.447	13.274	175	107	0.963	0.364	9.898	62.995
Perinatal mortality rate	45.676	14.394	177	108	0.940	0.315	16.888	74.463
Received ANC from a skilled provider	0.942	0.026	82	50	0.982	0.027	0.890	0.993
4+ ANC visits	0.650	0.051	82	50	0.966	0.079	0.547	0.752
8+ ANC visits	0.057	0.030	82	50	1.148	0.521	0.000	0.116
Took any iron-containing supplements	0.893	0.035	82	50	1.008	0.039	0.823	0.962
Mothers protected against tetanus for last birth	0.818	0.048	82	50	1.118	0.059	0.723	0.914
Delivered in a health facility (live births)	0.826	0.099	84	51	2.230	0.120	0.627	1.000
Delivered by a skilled provider (live births)	0.841	0.090	84	51	2.078	0.107	0.662	1.000
Delivered by C-section (live births)	0.169	0.061	84	51	1.479	0.362	0.047	0.291
Women with postnatal check during first 2 days	0.822	0.102	82	50	2.327	0.124	0.619	1.000
Newborns with postnatal check during first 2 days	0.826	0.089	82	50	2.079	0.108	0.647	1.000
Any problem accessing health care	0.385	0.055	321	195	2.001	0.142	0.275	0.494
Ever had a vaccination card	0.977	0.022	42	26	0.972	0.023	0.932	1.000
Received BCG vaccination	0.977	0.022	42	26	0.972	0.023	0.932	1.000
Received DPT-HepB+Hib vaccination (3 doses)	0.871	0.072	42	26	1.407	0.083	0.727	1.000
Received pneumococcal vaccination (3 doses)	0.847	0.074	42	26	1.345	0.087	0.699	0.994
Received measles vaccination	0.892	0.052	42	26	1.094	0.058	0.789	0.995
Fully vaccinated according to national schedule (12–23 months)	0.473	0.045	42	26	0.597	0.096	0.382	0.564
Received measles/measles-rubella 2 vaccination (24–35 months)	0.906	0.064	21	13	1.028	0.071	0.778	1.000
Fully vaccinated according to national schedule (24–35 months)	0.224	0.111	21	13	1.247	0.495	0.002	0.446
Received HPV vaccination (2 doses) women 15–19	0.546	0.100	51	32	1.403	0.183	0.346	0.745
Sought treatment for diarrhoea	0.408	0.143	16	10	1.110	0.351	0.122	0.695
Treated with ORS	0.191	0.113	16	10	1.132	0.594	0.000	0.418
Exclusive breastfeeding	0.907	0.064	19	11	0.939	0.071	0.778	1.000
Minimum dietary diversity (children 6–23 months)	0.278	0.072	59	37	1.217	0.259	0.134	0.421
Child slept under an ITN last night	0.135	0.038	174	104	1.339	0.282	0.059	0.211
Received 3+ doses of SP/Fansidar	0.000	0.000	82	50	na	na	0.000	0.000
Child has fever in last 2 weeks	0.121	0.028	163	100	1.061	0.228	0.066	0.176
Child had blood taken from finger/heel	0.121	0.057	19	12	0.776	0.469	0.007	0.234
Child took ACT	0.500	0.000	2	1	0.000	0.000	0.500	0.500
Condom use at last sex (women 15–24)	0.246	0.213	9	5	1.330	0.864	0.000	0.673
Mobile phone ownership	0.822	0.033	321	195	1.535	0.040	0.756	0.888

na = not applicable

Table B.36 Sampling errors: Nakuru sample, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases: Unweighted (N)	Number of cases: Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
HOUSEHOLDS AND POPULATION								
Primary reliance on clean fuels and technology	0.231	0.036	2,262	4,659	2.092	0.157	0.159	0.304
Births registered with civil authority	0.744	0.054	278	572	1.753	0.072	0.636	0.852
Ownership of at least one ITN	0.166	0.020	688	1,442	1.411	0.121	0.126	0.207
Ownership of at least one ITN for every two persons	0.099	0.012	685	1,435	1.094	0.127	0.074	0.124
At least basic drinking water service	0.809	0.060	2,262	4,659	3.082	0.074	0.689	0.929
At least basic sanitation service	0.355	0.057	2,262	4,659	2.610	0.161	0.241	0.469
Using open defecation	0.042	0.041	2,262	4,659	3.797	0.974	0.000	0.124
Using a handwashing facility with soap and water	0.659	0.032	2,239	4,614	1.479	0.049	0.595	0.723
WOMEN								
No education	0.020	0.013	504	1,077	1.984	0.617	0.000	0.045
Secondary education or higher	0.724	0.020	504	1,077	1.006	0.028	0.683	0.764
Literacy	0.946	0.014	505	1,078	1.408	0.015	0.918	0.974
Use of the internet in last 12 months	0.604	0.040	505	1,078	1.834	0.066	0.524	0.684
Total fertility rate (3 years)	3.455	0.242	1,427	3,052	0.879	0.070	2.970	3.939
Currently pregnant	0.047	0.010	505	1,078	1.036	0.208	0.027	0.066
Mean number of children ever born to women age 40–49	4.016	0.321	88	187	1.441	0.080	3.374	4.659
Median birth interval	50.183	3.702	186	386	1.446	0.074	42.779	57.586
Ideal number of children	3.837	0.140	503	1,074	1.664	0.036	3.558	4.117
Total wanted fertility rate (3 years)	3.050	0.217	1,427	3,052	0.810	0.071	2.616	3.484
Currently using any contraceptive method	0.654	0.038	291	614	1.345	0.058	0.579	0.729
Currently using any modern method	0.622	0.033	291	614	1.150	0.053	0.557	0.688
Currently using pill	0.074	0.017	291	614	1.127	0.234	0.040	0.109
Currently using injectables	0.256	0.037	291	614	1.452	0.145	0.182	0.331
Currently using implants	0.230	0.023	291	614	0.933	0.100	0.184	0.276
Currently using male condoms	0.011	0.006	291	614	0.972	0.536	0.000	0.023
Currently using any traditional method	0.032	0.014	291	614	1.382	0.448	0.003	0.060
Unmet need for spacing	0.058	0.014	291	614	0.990	0.235	0.031	0.085
Unmet need for limiting	0.067	0.015	291	614	0.988	0.216	0.038	0.096
Unmet need total	0.125	0.020	291	614	1.032	0.160	0.085	0.165
Demand satisfied by modern methods (married women)	0.799	0.028	226	479	1.058	0.035	0.742	0.855
Demand satisfied by modern methods (all women)	0.785	0.027	283	605	1.114	0.035	0.731	0.839
Not exposed to any of the ten media sources	0.068	0.011	505	1,078	1.017	0.168	0.045	0.090
Neonatal mortality (last 0–9 years)	18.547	5.104	511	1,074	0.878	0.275	8.339	28.754
Post-neonatal mortality (last 0–9 years)	9.745	4.248	512	1,077	1.000	0.436	1.250	18.240
Infant mortality (last 0–9 years)	28.292	6.018	511	1,074	0.846	0.213	16.256	40.327
Child mortality (last 0–9 years)	12.644	6.012	501	1,059	1.196	0.475	0.620	24.667
Under-five mortality (last 0–9 years)	40.578	7.147	511	1,074	0.833	0.176	26.283	54.872
Stillbirth rate	11.385	6.372	271	569	0.982	0.560	0.000	24.129
Early neonatal rate	15.366	6.879	269	565	0.913	0.448	1.609	29.123
Perinatal mortality rate	26.647	9.795	271	569	0.998	0.368	7.058	46.237
Received ANC from a skilled provider	1.000	0.000	108	222	na	0.000	1.000	1.000
4+ ANC visits	0.743	0.038	108	222	0.900	0.051	0.667	0.819
8+ ANC visits	0.060	0.027	108	222	1.156	0.442	0.007	0.113
Took any iron-containing supplements	0.954	0.024	108	222	1.161	0.025	0.907	1.000
Mothers protected against tetanus for last birth	0.781	0.038	108	222	0.954	0.049	0.705	0.858
Delivered in a health facility (live births)	0.963	0.022	109	224	1.175	0.023	0.920	1.000
Delivered by a skilled provider (live births)	0.969	0.019	109	224	1.121	0.020	0.931	1.000
Delivered by C-section (live births)	0.085	0.025	109	224	0.905	0.291	0.035	0.134
Women with postnatal check during first 2 days	0.889	0.028	108	222	0.931	0.032	0.833	0.946
Newborns with postnatal check during first 2 days	0.917	0.025	108	222	0.920	0.027	0.867	0.966
Any problem accessing health care	0.600	0.043	505	1,078	1.949	0.071	0.515	0.685
Ever had a vaccination card	1.000	0.000	53	109	na	0.000	1.000	1.000
Received BCG vaccination	1.000	0.000	53	109	na	0.000	1.000	1.000
Received DPT-HepB+Hib vaccination (3 doses)	0.924	0.043	53	109	1.162	0.047	0.838	1.000
Received pneumococcal vaccination (3 doses)	0.966	0.024	53	109	0.932	0.025	0.918	1.000
Received measles vaccination	0.904	0.036	53	109	0.879	0.040	0.831	0.976
Fully vaccinated according to national schedule (12–23 months)	0.635	0.074	53	109	1.093	0.116	0.487	0.782
Received measles/measles-rubella 2 vaccination (24–35 months)	0.776	0.063	51	111	1.083	0.081	0.650	0.902
Fully vaccinated according to national schedule (24–35 months)	0.503	0.078	51	111	1.120	0.155	0.347	0.659
Received HPV vaccination (2 doses) women 15–19	0.469	0.058	91	188	1.093	0.123	0.353	0.584
Sought treatment for diarrhoea	0.615	0.105	35	74	1.221	0.171	0.405	0.825
Treated with ORS	0.517	0.090	35	74	1.141	0.174	0.337	0.698
Exclusive breastfeeding	0.690	0.099	25	51	1.048	0.144	0.492	0.888
Minimum dietary diversity (children 6–23 months)	0.523	0.066	74	151	1.120	0.125	0.392	0.654
Child slept under an ITN last night	0.146	0.025	281	577	1.027	0.171	0.096	0.197
Received 3+ doses of SP/Fansidar	0.000	0.000	108	222	na	na	0.000	0.000
Child has fever in last 2 weeks	0.086	0.021	257	542	1.161	0.246	0.044	0.129
Child had blood taken from finger/heel	0.197	0.087	22	47	1.011	0.441	0.023	0.372
Child took ACT	na	na	-	-	na	na	0.000	na
Condom use at last sex (women 15–24)	0.366	0.093	34	76	1.111	0.255	0.179	0.553
Mobile phone ownership	0.829	0.020	505	1,078	1.192	0.024	0.789	0.869

na = not applicable

Table B.37 Sampling errors: Narok sample, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases: Unweighted (N)	Number of cases: Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
HOUSEHOLDS AND POPULATION								
Primary reliance on clean fuels and technology	0.079	0.016	2,399	2,265	1.525	0.205	0.047	0.112
Births registered with civil authority	0.501	0.080	279	254	2.260	0.159	0.342	0.660
Ownership of at least one ITN	0.513	0.060	613	576	2.932	0.116	0.394	0.632
Ownership of at least one ITN for every two persons	0.310	0.042	611	574	2.245	0.136	0.226	0.395
At least basic drinking water service	0.261	0.038	2,399	2,265	1.983	0.147	0.184	0.338
At least basic sanitation service	0.224	0.058	2,399	2,265	2.888	0.259	0.108	0.340
Using open defecation	0.161	0.072	2,399	2,265	3.977	0.449	0.016	0.305
Using a handwashing facility with soap and water	0.729	0.048	2,399	2,265	2.266	0.066	0.633	0.824
WOMEN								
No education	0.131	0.051	470	428	3.213	0.387	0.030	0.232
Secondary education or higher	0.503	0.048	470	428	2.090	0.096	0.406	0.600
Literacy	0.804	0.050	470	428	2.710	0.062	0.704	0.904
Use of the internet in last 12 months	0.349	0.036	470	428	1.634	0.103	0.277	0.421
Total fertility rate (3 years)	3.248	0.483	1,315	1,194	1.846	0.149	2.283	4.214
Currently pregnant	0.077	0.014	470	428	1.146	0.183	0.049	0.106
Mean number of children ever born to women age 40–49	4.788	0.332	68	66	1.166	0.069	4.124	5.452
Median birth interval	35.891	1.994	208	181	1.106	0.056	31.903	39.880
Ideal number of children	4.362	0.256	467	425	2.820	0.059	3.850	4.874
Total wanted fertility rate (3 years)	2.691	0.379	1,315	1,194	1.679	0.141	1.934	3.448
Currently using any contraceptive method	0.598	0.031	312	283	1.118	0.052	0.536	0.661
Currently using any modern method	0.553	0.036	312	283	1.269	0.065	0.481	0.625
Currently using pill	0.011	0.005	312	283	0.921	0.497	0.000	0.022
Currently using injectables	0.170	0.025	312	283	1.154	0.145	0.120	0.219
Currently using implants	0.303	0.032	312	283	1.216	0.105	0.240	0.367
Currently using male condoms	0.000	0.000	312	283	na	na	0.000	0.000
Currently using any traditional method	0.045	0.015	312	283	1.309	0.341	0.014	0.076
Unmet need for spacing	0.081	0.024	312	283	1.557	0.298	0.033	0.130
Unmet need for limiting	0.102	0.023	312	283	1.364	0.230	0.055	0.148
Unmet need total	0.183	0.026	312	283	1.196	0.144	0.130	0.235
Demand satisfied by modern methods (married women)	0.708	0.040	242	221	1.369	0.056	0.628	0.788
Demand satisfied by modern methods (all women)	0.690	0.034	282	256	1.217	0.049	0.623	0.757
Not exposed to any of the ten media sources	0.050	0.010	470	428	0.995	0.200	0.030	0.070
Neonatal mortality (last 0–9 years)	8.482	3.602	624	549	0.876	0.425	1.279	15.685
Post-neonatal mortality (last 0–9 years)	3.992	2.750	622	548	1.036	0.689	0.000	9.491
Infant mortality (last 0–9 years)	12.474	3.718	624	549	0.757	0.298	5.038	19.910
Child mortality (last 0–9 years)	1.482	1.499	645	571	0.967	1.011	0.000	4.481
Under-five mortality (last 0–9 years)	13.938	4.360	624	549	0.766	0.313	5.219	22.657
Stillbirth rate	7.927	4.666	280	245	0.866	0.589	0.000	17.259
Early neonatal rate	14.242	8.175	277	243	0.973	0.574	0.000	30.592
Perinatal mortality rate	22.056	8.646	280	245	0.879	0.392	4.764	39.348
Received ANC from a skilled provider	0.948	0.024	100	87	1.063	0.025	0.901	0.996
4+ ANC visits	0.567	0.081	100	87	1.616	0.143	0.405	0.730
8+ ANC visits	0.030	0.017	100	87	0.997	0.572	0.000	0.064
Took any iron-containing supplements	0.891	0.035	100	87	1.117	0.039	0.821	0.961
Mothers protected against tetanus for last birth	0.775	0.059	100	87	1.406	0.077	0.656	0.893
Delivered in a health facility (live births)	0.728	0.114	105	91	2.505	0.156	0.500	0.955
Delivered by a skilled provider (live births)	0.733	0.113	105	91	2.512	0.155	0.506	0.959
Delivered by C-section (live births)	0.136	0.047	105	91	1.292	0.342	0.043	0.230
Women with postnatal check during first 2 days	0.705	0.078	100	87	1.691	0.111	0.548	0.861
Newborns with postnatal check during first 2 days	0.771	0.066	100	87	1.551	0.086	0.639	0.903
Any problem accessing health care	0.670	0.042	470	428	1.921	0.062	0.587	0.754
Ever had a vaccination card	0.961	0.037	51	44	1.345	0.039	0.887	1.000
Received BCG vaccination	0.971	0.021	51	44	0.867	0.021	0.929	1.000
Received DPT-HepB+Hib vaccination (3 doses)	0.983	0.018	51	44	0.974	0.019	0.946	1.000
Received pneumococcal vaccination (3 doses)	0.893	0.046	51	44	1.042	0.052	0.801	0.985
Received measles vaccination	0.861	0.036	51	44	0.727	0.042	0.789	0.933
Fully vaccinated according to national schedule (12–23 months)	0.125	0.060	51	44	1.272	0.483	0.004	0.245
Received measles/measles-rubella 2 vaccination (24–35 months)	0.505	0.080	41	39	1.044	0.158	0.346	0.664
Fully vaccinated according to national schedule (24–35 months)	0.114	0.052	41	39	1.066	0.454	0.011	0.217
Received HPV vaccination (2 doses) women 15–19	0.251	0.042	80	78	0.862	0.167	0.167	0.335
Sought treatment for diarrhoea	0.729	0.126	16	14	1.126	0.173	0.477	0.982
Treated with ORS	0.686	0.144	16	14	1.227	0.210	0.398	0.973
Exclusive breastfeeding	0.721	0.120	21	18	1.186	0.167	0.481	0.962
Minimum dietary diversity (children 6–23 months)	0.352	0.059	76	66	1.076	0.169	0.234	0.471
Child slept under an ITN last night	0.468	0.062	278	253	1.758	0.132	0.344	0.591
Received 3+ doses of SP/Fansidar	0.021	0.014	100	87	0.982	0.677	0.000	0.049
Child has fever in last 2 weeks	0.086	0.018	269	236	1.020	0.211	0.050	0.122
Child had blood taken from finger/heel	0.372	0.062	23	20	0.635	0.168	0.247	0.497
Child took ACT	0.336	0.277	3	3	1.077	0.825	0.000	0.889
Condom use at last sex (women 15–24)	0.292	0.084	23	22	0.872	0.288	0.124	0.460
Mobile phone ownership	0.759	0.018	470	428	0.935	0.024	0.722	0.796

na = not applicable

Table B.38 Sampling errors: Kajiado sample, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases: Unweighted (N)	Number of cases: Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
HOUSEHOLDS AND POPULATION								
Primary reliance on clean fuels and technology	0.517	0.056	1,793	1,991	2.122	0.108	0.406	0.629
Births registered with civil authority	0.936	0.026	255	289	1.379	0.027	0.884	0.987
Ownership of at least one ITN	0.345	0.029	529	653	1.401	0.084	0.287	0.403
Ownership of at least one ITN for every two persons	0.192	0.026	527	651	1.532	0.137	0.139	0.244
At least basic drinking water service	0.779	0.070	1,793	1,991	2.898	0.090	0.639	0.920
At least basic sanitation service	0.396	0.069	1,793	1,991	2.735	0.174	0.259	0.534
Using open defecation	0.153	0.072	1,793	1,991	3.309	0.472	0.009	0.298
Using a handwashing facility with soap and water	0.813	0.027	1,793	1,991	1.256	0.033	0.759	0.867
WOMEN								
No education	0.091	0.028	366	464	1.846	0.306	0.036	0.147
Secondary education or higher	0.684	0.047	366	464	1.934	0.069	0.590	0.779
Literacy	0.872	0.030	369	467	1.709	0.034	0.812	0.932
Use of the internet in last 12 months	0.641	0.046	369	467	1.822	0.071	0.549	0.732
Total fertility rate (3 years)	3.629	0.497	1,057	1,346	1.753	0.137	2.634	4.623
Currently pregnant	0.070	0.016	369	467	1.174	0.224	0.038	0.101
Mean number of children ever born to women age 40–49	3.623	0.442	54	69	1.409	0.122	2.739	4.507
Median birth interval	38.364	4.317	172	188	1.534	0.113	29.730	46.997
Ideal number of children	3.589	0.193	367	466	1.722	0.054	3.202	3.976
Total wanted fertility rate (3 years)	3.243	0.459	1,057	1,346	1.613	0.142	2.325	4.161
Currently using any contraceptive method	0.633	0.042	217	261	1.288	0.067	0.549	0.718
Currently using any modern method	0.538	0.046	217	261	1.340	0.085	0.447	0.629
Currently using pill	0.100	0.020	217	261	0.999	0.204	0.059	0.141
Currently using injectables	0.179	0.027	217	261	1.039	0.151	0.125	0.234
Currently using implants	0.177	0.033	217	261	1.261	0.185	0.112	0.243
Currently using male condoms	0.011	0.009	217	261	1.222	0.790	0.000	0.028
Currently using any traditional method	0.095	0.017	217	261	0.867	0.182	0.061	0.130
Unmet need for spacing	0.047	0.014	217	261	0.982	0.300	0.019	0.076
Unmet need for limiting	0.021	0.010	217	261	0.997	0.459	0.002	0.041
Unmet need total	0.069	0.018	217	261	1.025	0.257	0.033	0.104
Demand satisfied by modern methods (married women)	0.767	0.043	147	183	1.247	0.056	0.681	0.852
Demand satisfied by modern methods (all women)	0.723	0.038	195	250	1.184	0.052	0.647	0.799
Not exposed to any of the ten media sources	0.033	0.010	369	467	1.066	0.299	0.013	0.053
Neonatal mortality (last 0–9 years)	24.048	7.311	465	525	1.014	0.304	9.426	38.671
Post-neonatal mortality (last 0–9 years)	4.187	2.956	465	525	0.944	0.706	0.000	10.100
Infant mortality (last 0–9 years)	28.236	8.223	465	525	1.069	0.291	11.789	44.682
Child mortality (last 0–9 years)	8.936	5.400	451	496	1.088	0.604	0.000	19.736
Under-five mortality (last 0–9 years)	36.919	8.739	466	526	0.984	0.237	19.442	54.396
Stillbirth rate	5.499	4.181	240	280	0.841	0.760	0.000	13.860
Early neonatal rate	24.959	10.083	238	278	0.975	0.404	4.793	45.125
Perinatal mortality rate	30.321	11.964	240	280	1.057	0.395	6.393	54.248
Received ANC from a skilled provider	0.992	0.008	96	123	0.896	0.008	0.976	1.000
4+ ANC visits	0.756	0.042	96	123	0.955	0.056	0.671	0.840
8+ ANC visits	0.094	0.038	96	123	1.246	0.397	0.019	0.169
Took any iron-containing supplements	0.941	0.028	96	123	1.172	0.030	0.885	0.998
Mothers protected against tetanus for last birth	0.928	0.028	96	123	1.038	0.030	0.873	0.983
Delivered in a health facility (live births)	0.853	0.075	102	130	2.147	0.088	0.702	1.000
Delivered by a skilled provider (live births)	0.858	0.076	102	130	2.203	0.089	0.706	1.000
Delivered by C-section (live births)	0.263	0.059	102	130	1.267	0.226	0.144	0.382
Women with postnatal check during first 2 days	0.741	0.077	96	123	1.693	0.104	0.587	0.895
Newborns with postnatal check during first 2 days	0.829	0.066	96	123	1.703	0.080	0.696	0.962
Any problem accessing health care	0.395	0.054	369	467	2.091	0.136	0.288	0.502
Ever had a vaccination card	1.000	0.000	45	58	na	0.000	1.000	1.000
Received BCG vaccination	1.000	0.000	45	58	na	0.000	1.000	1.000
Received DPT-HepB+Hib vaccination (3 doses)	0.969	0.021	45	58	0.807	0.021	0.927	1.000
Received pneumococcal vaccination (3 doses)	0.966	0.022	45	58	0.829	0.023	0.922	1.000
Received measles vaccination	0.935	0.052	45	58	1.431	0.055	0.832	1.000
Fully vaccinated according to national schedule (12–23 months)	0.483	0.085	45	58	1.157	0.175	0.314	0.653
Received measles/measles-rubella 2 vaccination (24–35 months)	0.665	0.082	40	42	1.013	0.124	0.500	0.830
Fully vaccinated according to national schedule (24–35 months)	0.391	0.101	40	42	1.197	0.258	0.190	0.593
Received HPV vaccination (2 doses) women 15–19	0.365	0.084	56	65	1.284	0.230	0.197	0.532
Sought treatment for diarrhoea	0.563	0.130	17	20	1.097	0.231	0.303	0.823
Treated with ORS	0.492	0.134	17	20	1.029	0.273	0.224	0.760
Exclusive breastfeeding	0.914	0.061	30	39	1.161	0.067	0.792	1.000
Minimum dietary diversity (children 6–23 months)	0.585	0.109	62	81	1.706	0.187	0.366	0.803
Child slept under an ITN last night	0.404	0.052	254	286	1.340	0.129	0.300	0.507
Received 3+ doses of SP/Fansidar	0.009	0.009	96	123	0.941	1.022	0.000	0.027
Child has fever in last 2 weeks	0.115	0.027	227	268	1.225	0.239	0.060	0.170
Child had blood taken from finger/heel	0.229	0.095	26	31	1.113	0.414	0.040	0.419
Child took ACT	0.794	0.231	2	3	0.962	0.292	0.331	1.000
Condom use at last sex (women 15–24)	0.380	0.092	33	39	1.073	0.243	0.195	0.565
Mobile phone ownership	0.861	0.029	369	467	1.634	0.034	0.802	0.920

na = not applicable

Table B.39 Sampling errors: Kericho sample, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases: Unweighted (N)	Number of cases: Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
HOUSEHOLDS AND POPULATION								
Primary reliance on clean fuels and technology	0.054	0.014	2,057	1,777	1.584	0.262	0.025	0.082
Births registered with civil authority	0.565	0.047	201	173	1.222	0.082	0.472	0.658
Ownership of at least one ITN	0.696	0.036	561	464	1.831	0.051	0.625	0.767
Ownership of at least one ITN for every two persons	0.454	0.038	548	455	1.790	0.084	0.377	0.530
At least basic drinking water service	0.571	0.061	2,057	1,777	2.475	0.106	0.449	0.692
At least basic sanitation service	0.618	0.039	2,057	1,777	1.709	0.063	0.539	0.696
Using open defecation	0.004	0.002	2,057	1,777	1.013	0.463	0.000	0.007
Using a handwashing facility with soap and water	0.705	0.027	2,050	1,771	1.171	0.038	0.651	0.759
WOMEN								
No education	0.002	0.001	460	384	0.591	0.620	0.000	0.004
Secondary education or higher	0.684	0.034	460	384	1.558	0.050	0.616	0.752
Literacy	0.954	0.011	463	387	1.156	0.012	0.931	0.976
Use of the internet in last 12 months	0.486	0.034	463	387	1.464	0.070	0.418	0.554
Total fertility rate (3 years)	2.608	0.261	1,306	1,091	1.159	0.100	2.086	3.131
Currently pregnant	0.040	0.007	463	387	0.770	0.176	0.026	0.054
Mean number of children ever born to women age 40–49	4.345	0.320	80	68	1.404	0.074	3.705	4.984
Median birth interval	50.349	4.376	123	106	1.013	0.087	41.598	59.101
Ideal number of children	3.226	0.070	460	384	1.218	0.022	3.087	3.366
Total wanted fertility rate (3 years)	2.062	0.237	1,306	1,091	1.078	0.115	1.588	2.536
Currently using any contraceptive method	0.628	0.033	234	198	1.053	0.053	0.561	0.694
Currently using any modern method	0.587	0.031	234	198	0.951	0.052	0.525	0.648
Currently using pill	0.006	0.004	234	198	0.883	0.754	0.000	0.015
Currently using injectables	0.221	0.031	234	198	1.135	0.140	0.159	0.282
Currently using implants	0.284	0.039	234	198	1.311	0.137	0.207	0.362
Currently using male condoms	0.014	0.008	234	198	1.002	0.552	0.000	0.029
Currently using any traditional method	0.041	0.009	234	198	0.725	0.229	0.022	0.060
Unmet need for spacing	0.063	0.013	234	198	0.797	0.202	0.037	0.088
Unmet need for limiting	0.068	0.015	234	198	0.937	0.227	0.037	0.099
Unmet need total	0.131	0.022	234	198	0.974	0.164	0.088	0.174
Demand satisfied by modern methods (married women)	0.773	0.027	179	150	0.851	0.035	0.720	0.827
Demand satisfied by modern methods (all women)	0.789	0.028	259	217	1.100	0.035	0.733	0.845
Not exposed to any of the ten media sources	0.021	0.007	463	387	1.020	0.322	0.008	0.035
Neonatal mortality (last 0–9 years)	7.044	4.096	401	343	0.979	0.581	0.000	15.237
Post-neonatal mortality (last 0–9 years)	11.948	6.917	396	337	1.276	0.579	0.000	25.783
Infant mortality (last 0–9 years)	18.993	8.137	401	343	1.205	0.428	2.719	35.266
Child mortality (last 0–9 years)	7.848	4.357	408	346	0.985	0.555	0.000	16.563
Under-five mortality (last 0–9 years)	26.692	9.109	401	343	1.143	0.341	8.474	44.910
Stillbirth rate	17.747	9.407	194	166	1.007	0.530	0.000	36.560
Early neonatal rate	14.456	8.227	194	166	0.975	0.569	0.000	30.910
Perinatal mortality rate	32.132	13.425	194	166	0.925	0.418	5.282	58.981
Received ANC from a skilled provider	1.000	0.000	77	68	na	0.000	1.000	1.000
4+ ANC visits	0.551	0.049	77	68	0.853	0.088	0.454	0.649
8+ ANC visits	0.018	0.018	77	68	1.183	0.999	0.000	0.055
Took any iron-containing supplements	0.879	0.043	77	68	1.137	0.048	0.794	0.964
Mothers protected against tetanus for last birth	0.863	0.036	77	68	0.912	0.042	0.791	0.935
Delivered in a health facility (live births)	0.908	0.030	79	70	0.976	0.034	0.847	0.969
Delivered by a skilled provider (live births)	0.917	0.030	79	70	1.000	0.032	0.858	0.977
Delivered by C-section (live births)	0.215	0.035	79	70	0.798	0.165	0.144	0.286
Women with postnatal check during first 2 days	0.786	0.043	77	68	0.924	0.055	0.699	0.873
Newborns with postnatal check during first 2 days	0.944	0.018	77	68	0.699	0.019	0.908	0.981
Any problem accessing health care	0.300	0.027	463	387	1.270	0.090	0.246	0.355
Ever had a vaccination card	1.000	0.000	42	36	na	0.000	1.000	1.000
Received BCG vaccination	1.000	0.000	42	36	na	0.000	1.000	1.000
Received DPT-HepB+Hib vaccination (3 doses)	0.979	0.019	42	36	0.883	0.020	0.940	1.000
Received pneumococcal vaccination (3 doses)	0.960	0.024	42	36	0.818	0.025	0.912	1.000
Received measles vaccination	0.957	0.039	42	36	1.261	0.041	0.880	1.000
Fully vaccinated according to national schedule (12–23 months)	0.582	0.061	42	36	0.817	0.105	0.459	0.704
Received measles/measles-rubella 2 vaccination (24–35 months)	0.942	0.041	33	28	1.023	0.044	0.859	1.000
Fully vaccinated according to national schedule (24–35 months)	0.524	0.075	33	28	0.911	0.143	0.374	0.674
Received HPV vaccination (2 doses) women 15–19	0.576	0.069	84	70	1.261	0.119	0.439	0.713
Sought treatment for diarrhoea	0.622	0.136	14	12	1.058	0.218	0.351	0.893
Treated with ORS	0.369	0.170	14	12	1.331	0.461	0.029	0.708
Exclusive breastfeeding	0.720	0.080	22	20	0.825	0.111	0.560	0.881
Minimum dietary diversity (children 6–23 months)	0.603	0.079	53	46	1.160	0.131	0.445	0.761
Child slept under an ITN last night	0.615	0.048	201	175	1.207	0.079	0.518	0.712
Received 3+ doses of SP/Fansidar	0.044	0.025	77	68	1.072	0.574	0.000	0.094
Child has fever in last 2 weeks	0.087	0.020	187	159	0.964	0.235	0.046	0.128
Child had blood taken from finger/heel	0.089	0.031	19	14	0.444	0.350	0.027	0.152
Child took ACT	na	na	-	-	na	na	0.000	na
Condom use at last sex (women 15–24)	0.450	0.083	45	39	1.103	0.184	0.284	0.615
Mobile phone ownership	0.808	0.023	463	387	1.257	0.028	0.762	0.854

na = not applicable

Table B.40 Sampling errors: Bomet sample, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases: Unweighted (N)	Number of cases: Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
HOUSEHOLDS AND POPULATION								
Primary reliance on clean fuels and technology	0.048	0.020	2,385	1,963	2.841	0.425	0.007	0.088
Births registered with civil authority	0.772	0.032	273	224	1.115	0.042	0.708	0.837
Ownership of at least one ITN	0.648	0.033	565	437	1.636	0.051	0.582	0.714
Ownership of at least one ITN for every two persons	0.353	0.041	564	437	2.040	0.117	0.271	0.436
At least basic drinking water service	0.389	0.058	2,385	1,963	2.553	0.150	0.272	0.506
At least basic sanitation service	0.148	0.026	2,385	1,963	1.476	0.173	0.097	0.199
Using open defecation	0.001	0.001	2,385	1,963	1.045	0.983	0.000	0.004
Using a handwashing facility with soap and water	0.384	0.033	2,370	1,951	1.408	0.086	0.318	0.450
WOMEN								
No education	0.002	0.002	591	478	1.053	1.005	0.000	0.006
Secondary education or higher	0.648	0.035	591	478	1.777	0.054	0.578	0.718
Literacy	0.989	0.005	594	480	1.139	0.005	0.980	0.999
Use of the internet in last 12 months	0.350	0.044	594	480	2.216	0.124	0.263	0.437
Total fertility rate (3 years)	3.354	0.336	1,650	1,327	1.485	0.100	2.681	4.027
Currently pregnant	0.040	0.011	594	480	1.385	0.280	0.018	0.062
Mean number of children ever born to women age 40–49	5.207	0.272	103	86	1.182	0.052	4.664	5.751
Median birth interval	41.669	3.046	189	156	1.465	0.073	35.577	47.760
Ideal number of children	3.593	0.073	594	480	1.162	0.020	3.447	3.738
Total wanted fertility rate (3 years)	2.345	0.293	1,650	1,327	1.458	0.125	1.759	2.931
Currently using any contraceptive method	0.550	0.029	295	242	1.015	0.054	0.491	0.609
Currently using any modern method	0.513	0.032	295	242	1.086	0.062	0.450	0.576
Currently using pill	0.018	0.007	295	242	0.866	0.369	0.005	0.032
Currently using injectables	0.191	0.023	295	242	1.022	0.123	0.144	0.238
Currently using implants	0.245	0.027	295	242	1.065	0.109	0.191	0.298
Currently using male condoms	0.000	0.000	295	242	0.307	1.007	0.000	0.001
Currently using any traditional method	0.037	0.015	295	242	1.338	0.398	0.008	0.067
Unmet need for spacing	0.090	0.017	295	242	1.027	0.190	0.056	0.125
Unmet need for limiting	0.114	0.017	295	242	0.929	0.151	0.080	0.149
Unmet need total	0.205	0.022	295	242	0.931	0.107	0.161	0.249
Demand satisfied by modern methods (married women)	0.680	0.033	227	183	1.041	0.048	0.614	0.745
Demand satisfied by modern methods (all women)	0.639	0.030	296	236	1.055	0.046	0.580	0.699
Not exposed to any of the ten media sources	0.038	0.010	594	480	1.315	0.273	0.017	0.058
Neonatal mortality (last 0–9 years)	7.865	3.345	575	479	0.915	0.425	1.176	14.555
Post-neonatal mortality (last 0–9 years)	5.382	2.913	570	475	0.961	0.541	0.000	11.209
Infant mortality (last 0–9 years)	13.247	4.492	575	479	0.957	0.339	4.262	22.232
Child mortality (last 0–9 years)	2.811	1.955	571	477	0.990	0.696	0.000	6.721
Under-five mortality (last 0–9 years)	16.021	4.295	575	479	0.917	0.268	7.431	24.610
Stillbirth rate	13.203	7.385	280	231	1.101	0.559	0.000	27.974
Early neonatal rate	1.631	1.625	277	228	0.678	0.997	0.000	4.882
Perinatal mortality rate	14.813	7.554	280	231	1.066	0.510	0.000	29.922
Received ANC from a skilled provider	1.000	0.000	106	89	na	0.000	1.000	1.000
4+ ANC visits	0.692	0.045	106	89	1.006	0.065	0.602	0.783
8+ ANC visits	0.106	0.029	106	89	0.961	0.272	0.048	0.164
Took any iron-containing supplements	0.973	0.019	106	89	1.192	0.020	0.935	1.000
Mothers protected against tetanus for last birth	0.653	0.062	106	89	1.331	0.095	0.529	0.777
Delivered in a health facility (live births)	0.960	0.018	106	89	0.966	0.019	0.924	0.996
Delivered by a skilled provider (live births)	0.960	0.018	106	89	0.966	0.019	0.924	0.996
Delivered by C-section (live births)	0.144	0.042	106	89	1.242	0.288	0.061	0.228
Women with postnatal check during first 2 days	0.920	0.033	106	89	1.229	0.035	0.855	0.985
Newborns with postnatal check during first 2 days	0.918	0.032	106	89	1.195	0.035	0.854	0.982
Any problem accessing health care	0.601	0.040	594	480	1.995	0.067	0.520	0.681
Ever had a vaccination card	0.985	0.015	61	52	0.962	0.015	0.955	1.000
Received BCG vaccination	0.944	0.039	61	52	1.380	0.042	0.866	1.000
Received DPT-HepB+Hib vaccination (3 doses)	0.878	0.050	61	52	1.231	0.057	0.777	0.978
Received pneumococcal vaccination (3 doses)	0.915	0.046	61	52	1.332	0.051	0.823	1.000
Received measles vaccination	0.867	0.081	61	52	1.916	0.094	0.704	1.000
Fully vaccinated according to national schedule (12–23 months)	0.374	0.054	61	52	0.890	0.143	0.267	0.481
Received measles/measles-rubella 2 vaccination (24–35 months)	0.795	0.061	61	49	1.177	0.077	0.673	0.918
Fully vaccinated according to national schedule (24–35 months)	0.416	0.072	61	49	1.138	0.174	0.272	0.561
Received HPV vaccination (2 doses) women 15–19	0.427	0.039	127	106	0.885	0.091	0.349	0.505
Sought treatment for diarrhoea	0.346	0.214	11	8	1.435	0.620	0.000	0.774
Treated with ORS	0.327	0.196	11	8	1.334	0.601	0.000	0.720
Exclusive breastfeeding	0.893	0.056	21	17	0.818	0.063	0.781	1.000
Minimum dietary diversity (children 6–23 months)	0.349	0.053	84	71	1.021	0.153	0.242	0.456
Child slept under an ITN last night	0.497	0.039	282	233	1.186	0.078	0.419	0.575
Received 3+ doses of SP/Fansidar	0.001	0.001	106	89	0.304	1.010	0.000	0.003
Child has fever in last 2 weeks	0.052	0.019	275	227	1.443	0.369	0.014	0.091
Child had blood taken from finger/heel	0.009	0.010	19	12	0.398	1.071	0.000	0.029
Child took ACT	0.000	0.000	1	1	na	na	0.000	0.000
Condom use at last sex (women 15–24)	0.388	0.077	63	50	1.239	0.199	0.234	0.542
Mobile phone ownership	0.698	0.034	594	480	1.792	0.048	0.630	0.766

na = not applicable

Table B.41 Sampling errors: Kakamega sample, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases: Unweighted (N)	Number of cases: Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
HOUSEHOLDS AND POPULATION								
Primary reliance on clean fuels and technology	0.068	0.011	2,493	3,162	1.181	0.169	0.045	0.091
Births registered with civil authority	0.905	0.022	283	360	0.961	0.024	0.862	0.948
Ownership of at least one ITN	0.740	0.019	684	850	1.156	0.026	0.702	0.779
Ownership of at least one ITN for every two persons	0.493	0.020	680	847	1.049	0.041	0.453	0.533
At least basic drinking water service	0.747	0.045	2,493	3,162	2.241	0.060	0.658	0.836
At least basic sanitation service	0.501	0.032	2,493	3,162	1.451	0.064	0.437	0.565
Using open defecation	0.000	0.000	2,493	3,162	na	na	0.000	0.000
Using a handwashing facility with soap and water	0.230	0.044	2,493	3,162	2.337	0.191	0.142	0.318
WOMEN								
No education	0.021	0.006	523	660	0.996	0.299	0.008	0.033
Secondary education or higher	0.616	0.028	523	660	1.293	0.045	0.561	0.671
Literacy	0.945	0.012	526	665	1.187	0.013	0.921	0.968
Use of the internet in last 12 months	0.569	0.034	526	665	1.585	0.060	0.500	0.637
Total fertility rate (3 years)	3.476	0.239	1,472	1,862	1.058	0.069	2.999	3.953
Currently pregnant	0.052	0.010	526	665	1.076	0.202	0.031	0.072
Mean number of children ever born to women age 40–49	4.792	0.195	111	140	1.001	0.041	4.403	5.181
Median birth interval	45.938	3.506	182	234	1.427	0.076	38.926	52.951
Ideal number of children	3.602	0.071	526	665	1.028	0.020	3.459	3.744
Total wanted fertility rate (3 years)	2.984	0.239	1,472	1,862	1.182	0.080	2.507	3.462
Currently using any contraceptive method	0.670	0.030	314	406	1.125	0.045	0.610	0.730
Currently using any modern method	0.661	0.029	314	406	1.082	0.044	0.603	0.719
Currently using pill	0.048	0.013	314	406	1.087	0.273	0.022	0.074
Currently using injectables	0.184	0.025	314	406	1.141	0.136	0.134	0.234
Currently using implants	0.359	0.026	314	406	0.950	0.072	0.308	0.411
Currently using male condoms	0.001	0.001	314	406	0.627	1.016	0.000	0.004
Currently using any traditional method	0.009	0.007	314	406	1.246	0.734	0.000	0.023
Unmet need for spacing	0.054	0.011	314	406	0.894	0.210	0.032	0.077
Unmet need for limiting	0.091	0.011	314	406	0.682	0.122	0.069	0.113
Unmet need total	0.146	0.016	314	406	0.807	0.110	0.114	0.178
Demand satisfied by modern methods (married women)	0.810	0.020	256	331	0.801	0.024	0.771	0.850
Demand satisfied by modern methods (all women)	0.813	0.019	322	410	0.874	0.023	0.775	0.851
Not exposed to any of the ten media sources	0.098	0.017	526	665	1.323	0.175	0.064	0.133
Neonatal mortality (last 0–9 years)	23.954	7.247	483	624	1.045	0.303	9.460	38.448
Post-neonatal mortality (last 0–9 years)	23.704	6.096	484	624	0.903	0.257	11.512	35.896
Infant mortality (last 0–9 years)	47.658	10.716	484	624	1.073	0.225	26.225	69.091
Child mortality (last 0–9 years)	5.446	3.182	473	614	1.026	0.584	0.000	11.809
Under-five mortality (last 0–9 years)	52.844	11.168	484	624	1.093	0.211	30.508	75.180
Stillbirth rate	3.377	3.398	260	329	0.947	1.006	0.000	10.174
Early neonatal rate	3.529	3.622	261	330	0.986	1.026	0.000	10.773
Perinatal mortality rate	6.922	4.948	260	329	0.960	0.715	0.000	16.817
Received ANC from a skilled provider	0.982	0.018	101	126	1.340	0.018	0.947	1.000
4+ ANC visits	0.755	0.047	101	126	1.081	0.062	0.662	0.848
8+ ANC visits	0.055	0.021	101	126	0.907	0.375	0.014	0.096
Took any iron-containing supplements	0.882	0.029	101	126	0.909	0.033	0.823	0.940
Mothers protected against tetanus for last birth	0.873	0.038	101	126	1.128	0.043	0.798	0.948
Delivered in a health facility (live births)	0.926	0.031	104	130	1.181	0.033	0.864	0.988
Delivered by a skilled provider (live births)	0.922	0.031	104	130	1.163	0.034	0.860	0.984
Delivered by C-section (live births)	0.135	0.037	104	130	1.093	0.276	0.061	0.210
Women with postnatal check during first 2 days	0.784	0.044	101	126	1.079	0.057	0.695	0.873
Newborns with postnatal check during first 2 days	0.856	0.033	101	126	0.935	0.038	0.790	0.921
Any problem accessing health care	0.727	0.039	526	665	2.015	0.054	0.648	0.805
Ever had a vaccination card	0.981	0.019	49	62	0.973	0.019	0.943	1.000
Received BCG vaccination	0.981	0.019	49	62	0.973	0.019	0.943	1.000
Received DPT-HepB+Hib vaccination (3 doses)	0.853	0.049	49	62	0.952	0.057	0.755	0.950
Received pneumococcal vaccination (3 doses)	0.849	0.049	49	62	0.954	0.058	0.750	0.947
Received measles vaccination	0.942	0.032	49	62	0.949	0.034	0.878	1.000
Fully vaccinated according to national schedule (12–23 months)	0.477	0.070	49	62	0.949	0.146	0.337	0.616
Received measles/measles-rubella 2 vaccination (24–35 months)	0.835	0.064	51	65	1.219	0.077	0.707	0.963
Fully vaccinated according to national schedule (24–35 months)	0.533	0.076	51	65	1.058	0.142	0.382	0.684
Received HPV vaccination (2 doses) women 15–19	0.390	0.039	109	139	0.836	0.101	0.312	0.468
Sought treatment for diarrhoea	0.564	0.095	44	55	1.249	0.168	0.374	0.753
Treated with ORS	0.313	0.063	44	55	0.886	0.201	0.187	0.439
Exclusive breastfeeding	0.554	0.085	26	31	0.864	0.154	0.384	0.725
Minimum dietary diversity (children 6–23 months)	0.174	0.049	68	86	1.056	0.282	0.076	0.272
Child slept under an ITN last night	0.632	0.044	293	370	1.391	0.069	0.545	0.720
Received 3+ doses of SP/Fansidar	0.419	0.047	101	126	0.948	0.112	0.325	0.512
Child has fever in last 2 weeks	0.194	0.033	252	318	1.304	0.172	0.127	0.260
Child had blood taken from finger/heel	0.137	0.046	48	62	0.947	0.336	0.045	0.230
Child took ACT	0.936	0.064	11	14	0.859	0.068	0.808	1.000
Condom use at last sex (women 15–24)	0.254	0.067	46	56	1.036	0.265	0.119	0.389
Mobile phone ownership	0.748	0.030	526	665	1.563	0.040	0.688	0.807

Table B.42 Sampling errors: Vihiga sample, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases Unweighted (N)	Number of cases Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits Lower (R-2SE)	Confidence limits Upper (R+2SE)
HOUSEHOLDS AND POPULATION								
Primary reliance on clean fuels and technology	0.038	0.011	2,002	1,055	1.401	0.295	0.016	0.061
Births registered with civil authority	0.880	0.027	228	120	1.082	0.030	0.827	0.934
Ownership of at least one ITN	0.817	0.021	513	267	1.228	0.026	0.775	0.859
Ownership of at least one ITN for every two persons	0.602	0.027	513	267	1.264	0.045	0.547	0.657
At least basic drinking water service	0.838	0.032	2,002	1,055	1.651	0.038	0.774	0.901
At least basic sanitation service	0.507	0.031	2,002	1,055	1.209	0.060	0.446	0.568
Using open defecation	0.062	0.013	2,002	1,055	1.142	0.211	0.036	0.088
Using a handwashing facility with soap and water	0.372	0.019	1,941	1,022	0.788	0.052	0.333	0.411
WOMEN								
No education	0.003	0.003	443	233	0.930	0.760	0.000	0.009
Secondary education or higher	0.698	0.020	443	233	0.935	0.029	0.657	0.739
Literacy	0.973	0.008	443	233	1.054	0.008	0.956	0.989
Use of the internet in last 12 months	0.525	0.030	443	233	1.245	0.056	0.466	0.584
Total fertility rate (3 years)	3.542	0.386	1,195	625	1.409	0.109	2.769	4.314
Currently pregnant	0.051	0.010	443	233	0.972	0.199	0.031	0.072
Mean number of children ever born to women age 40–49	4.616	0.222	97	51	1.100	0.048	4.172	5.060
Median birth interval	49.452	3.833	130	69	1.009	0.078	41.786	57.118
Ideal number of children	3.595	0.053	439	230	0.814	0.015	3.489	3.700
Total wanted fertility rate (3 years)	2.831	0.416	1,195	625	1.493	0.147	1.999	3.662
Currently using any contraceptive method	0.628	0.044	225	118	1.357	0.070	0.540	0.716
Currently using any modern method	0.613	0.047	225	118	1.439	0.077	0.519	0.707
Currently using pill	0.028	0.011	225	118	1.011	0.399	0.006	0.050
Currently using injectables	0.152	0.015	225	118	0.615	0.097	0.123	0.182
Currently using implants	0.353	0.041	225	118	1.279	0.116	0.271	0.435
Currently using male condoms	0.010	0.006	225	118	0.852	0.573	0.000	0.021
Currently using any traditional method	0.015	0.008	225	118	0.937	0.509	0.000	0.030
Unmet need for spacing	0.062	0.012	225	118	0.766	0.200	0.037	0.086
Unmet need for limiting	0.129	0.025	225	118	1.108	0.193	0.079	0.179
Unmet need total	0.191	0.028	225	118	1.084	0.149	0.134	0.248
Demand satisfied by modern methods (married women)	0.749	0.042	181	97	1.316	0.056	0.664	0.833
Demand satisfied by modern methods (all women)	0.736	0.036	237	126	1.267	0.049	0.664	0.809
Not exposed to any of the ten media sources	0.071	0.012	443	233	1.014	0.174	0.047	0.096
Neonatal mortality (last 0–9 years)	29.884	9.015	370	196	0.912	0.302	11.854	47.913
Post-neonatal mortality (last 0–9 years)	6.809	4.729	364	193	1.107	0.695	0.000	16.266
Infant mortality (last 0–9 years)	36.692	10.002	370	196	0.937	0.273	16.688	56.697
Child mortality (last 0–9 years)	7.172	4.128	354	188	0.996	0.576	0.000	15.428
Under-five mortality (last 0–9 years)	43.601	10.746	371	196	1.007	0.246	22.109	65.093
Stillbirth rate	28.046	10.307	196	102	0.880	0.368	7.431	48.660
Early neonatal rate	17.623	10.349	192	99	0.955	0.587	0.000	38.320
Perinatal mortality rate	45.174	11.685	196	102	0.758	0.259	21.805	68.544
Received ANC from a skilled provider	1.000	0.000	71	38	na	0.000	1.000	1.000
4+ ANC visits	0.763	0.053	71	38	1.038	0.069	0.657	0.869
8+ ANC visits	0.158	0.044	71	38	1.004	0.277	0.070	0.245
Took any iron-containing supplements	0.948	0.033	71	38	1.261	0.035	0.881	1.000
Mothers protected against tetanus for last birth	0.936	0.026	71	38	0.881	0.028	0.884	0.987
Delivered in a health facility (live births)	0.932	0.024	71	38	0.801	0.025	0.885	0.979
Delivered by a skilled provider (live births)	0.914	0.033	71	38	1.003	0.036	0.848	0.979
Delivered by C-section (live births)	0.135	0.039	71	38	0.979	0.289	0.057	0.213
Women with postnatal check during first 2 days	0.889	0.054	71	38	1.438	0.061	0.781	0.998
Newborns with postnatal check during first 2 days	0.860	0.047	71	38	1.124	0.054	0.767	0.954
Any problem accessing health care	0.462	0.026	443	233	1.112	0.057	0.409	0.515
Ever had a vaccination card	1.000	0.000	27	15	na	0.000	1.000	1.000
Received BCG vaccination	1.000	0.000	27	15	na	0.000	1.000	1.000
Received DPT-HepB+Hib vaccination (3 doses)	1.000	0.000	27	15	na	0.000	1.000	1.000
Received pneumococcal vaccination (3 doses)	1.000	0.000	27	15	na	0.000	1.000	1.000
Received measles vaccination	1.000	0.000	27	15	na	0.000	1.000	1.000
Fully vaccinated according to national schedule (12–23 months)	0.650	0.074	27	15	0.833	0.114	0.502	0.799
Received measles/measles-rubella 2 vaccination (24–35 months)	0.923	0.047	39	20	1.097	0.051	0.830	1.000
Fully vaccinated according to national schedule (24–35 months)	0.600	0.072	39	20	0.913	0.119	0.457	0.743
Received HPV vaccination (2 doses) women 15–19	0.600	0.055	123	66	1.228	0.091	0.491	0.709
Sought treatment for diarrhoea	0.702	0.159	16	8	1.284	0.226	0.384	1.000
Treated with ORS	0.517	0.115	16	8	0.826	0.222	0.288	0.746
Exclusive breastfeeding	0.565	0.115	23	12	1.087	0.204	0.334	0.796
Minimum dietary diversity (children 6–23 months)	0.526	0.117	47	25	1.562	0.222	0.292	0.760
Child slept under an ITN last night	0.687	0.044	235	124	1.283	0.064	0.599	0.774
Received 3+ doses of SP/Fansidar	0.623	0.061	71	38	1.060	0.099	0.500	0.746
Child has fever in last 2 weeks	0.119	0.030	184	95	1.197	0.252	0.059	0.178
Child had blood taken from finger/heel	0.412	0.101	23	11	0.998	0.245	0.210	0.615
Child took ACT	1.000	0.000	4	2	na	0.000	1.000	1.000
Condom use at last sex (women 15–24)	0.401	0.088	30	15	0.964	0.219	0.226	0.576
Mobile phone ownership	0.706	0.025	443	233	1.139	0.035	0.656	0.755

na = not applicable

Table B.43 Sampling errors: Bungoma sample, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases Unweighted (N)	Number of cases Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits Lower (R-2SE)	Confidence limits Upper (R+2SE)
HOUSEHOLDS AND POPULATION								
Primary reliance on clean fuels and technology	0.054	0.012	2,678	3,052	1.497	0.218	0.031	0.078
Births registered with civil authority	0.759	0.045	319	368	1.682	0.059	0.669	0.849
Ownership of at least one ITN	0.766	0.021	668	746	1.285	0.028	0.724	0.808
Ownership of at least one ITN for every two persons	0.500	0.035	666	744	1.780	0.069	0.431	0.569
At least basic drinking water service	0.670	0.034	2,678	3,052	1.583	0.051	0.602	0.738
At least basic sanitation service	0.433	0.034	2,678	3,052	1.517	0.078	0.365	0.500
Using open defecation	0.000	0.000	2,678	3,052	na	na	0.000	0.000
Using a handwashing facility with soap and water	0.741	0.036	2,665	3,038	1.805	0.049	0.668	0.814
WOMEN								
No education	0.009	0.004	573	624	0.930	0.409	0.002	0.016
Secondary education or higher	0.649	0.019	573	624	0.969	0.030	0.610	0.688
Literacy	0.959	0.008	574	625	0.952	0.008	0.944	0.975
Use of the internet in last 12 months	0.342	0.033	574	625	1.679	0.098	0.275	0.408
Total fertility rate (3 years)	3.555	0.354	1,578	1,707	1.483	0.100	2.846	4.263
Currently pregnant	0.075	0.012	574	625	1.097	0.161	0.051	0.099
Mean number of children ever born to women age 40–49	5.472	0.214	108	116	0.949	0.039	5.043	5.901
Median birth interval	50.999	4.280	196	217	1.773	0.084	42.439	59.559
Ideal number of children	3.917	0.119	572	623	1.614	0.030	3.678	4.156
Total wanted fertility rate (3 years)	2.980	0.278	1,578	1,707	1.380	0.093	2.425	3.536
Currently using any contraceptive method	0.579	0.031	317	349	1.128	0.054	0.516	0.641
Currently using any modern method	0.558	0.033	317	349	1.168	0.058	0.493	0.623
Currently using pill	0.017	0.009	317	349	1.268	0.548	0.000	0.035
Currently using injectables	0.206	0.020	317	349	0.859	0.095	0.167	0.245
Currently using implants	0.256	0.025	317	349	1.009	0.097	0.206	0.305
Currently using male condoms	0.011	0.006	317	349	1.051	0.560	0.000	0.023
Currently using any traditional method	0.021	0.009	317	349	1.083	0.419	0.003	0.038
Unmet need for spacing	0.105	0.021	317	349	1.207	0.199	0.063	0.146
Unmet need for limiting	0.088	0.015	317	349	0.910	0.165	0.059	0.117
Unmet need total	0.193	0.028	317	349	1.249	0.144	0.137	0.248
Demand satisfied by modern methods (married women)	0.723	0.033	243	269	1.162	0.046	0.657	0.790
Demand satisfied by modern methods (all women)	0.718	0.030	303	329	1.150	0.041	0.659	0.778
Not exposed to any of the ten media sources	0.033	0.006	574	625	0.867	0.197	0.020	0.046
Neonatal mortality (last 0–9 years)	21.871	4.057	576	627	0.678	0.185	13.757	29.984
Post-neonatal mortality (last 0–9 years)	10.421	3.802	574	623	0.904	0.365	2.817	18.026
Infant mortality (last 0–9 years)	32.292	5.553	576	627	0.765	0.172	21.186	43.398
Child mortality (last 0–9 years)	7.877	3.680	558	609	0.977	0.467	0.517	15.237
Under-five mortality (last 0–9 years)	39.914	6.572	577	628	0.807	0.165	26.771	53.058
Stillbirth rate	4.181	4.170	286	315	1.098	0.997	0.000	12.521
Early neonatal rate	19.798	6.743	285	314	0.828	0.341	6.312	33.284
Perinatal mortality rate	23.896	7.124	286	315	0.799	0.298	9.648	38.145
Received ANC from a skilled provider	0.992	0.009	115	128	1.019	0.009	0.974	1.000
4+ ANC visits	0.731	0.049	115	128	1.185	0.068	0.632	0.829
8+ ANC visits	0.104	0.036	115	128	1.253	0.344	0.033	0.176
Took any iron-containing supplements	0.794	0.054	115	128	1.414	0.068	0.687	0.902
Mothers protected against tetanus for last birth	0.793	0.052	115	128	1.363	0.065	0.689	0.897
Delivered in a health facility (live births)	0.896	0.035	118	131	1.265	0.039	0.827	0.966
Delivered by a skilled provider (live births)	0.896	0.035	118	131	1.265	0.039	0.827	0.966
Delivered by C-section (live births)	0.132	0.034	118	131	1.080	0.259	0.063	0.200
Women with postnatal check during first 2 days	0.839	0.038	115	128	1.112	0.046	0.762	0.916
Newborns with postnatal check during first 2 days	0.860	0.042	115	128	1.297	0.049	0.775	0.945
Any problem accessing health care	0.826	0.021	574	625	1.302	0.025	0.785	0.868
Ever had a vaccination card	1.000	0.000	54	63	na	0.000	1.000	1.000
Received BCG vaccination	0.959	0.026	54	63	0.982	0.027	0.908	1.000
Received DPT-HepB+Hib vaccination (3 doses)	0.904	0.044	54	63	1.132	0.048	0.816	0.992
Received pneumococcal vaccination (3 doses)	0.953	0.033	54	63	1.186	0.035	0.887	1.000
Received measles vaccination	0.839	0.035	54	63	0.731	0.042	0.769	0.910
Fully vaccinated according to national schedule (12–23 months)	0.585	0.071	54	63	1.103	0.122	0.443	0.728
Received measles/measles-rubella 2 vaccination (24–35 months)	0.634	0.068	51	56	0.966	0.107	0.498	0.769
Fully vaccinated according to national schedule (24–35 months)	0.496	0.084	51	56	1.163	0.170	0.327	0.664
Received HPV vaccination (2 doses) women 15–19	0.393	0.042	136	158	1.004	0.107	0.309	0.478
Sought treatment for diarrhoea	0.608	0.112	23	22	1.002	0.183	0.385	0.831
Treated with ORS	0.401	0.120	23	22	1.039	0.300	0.160	0.641
Exclusive breastfeeding	0.544	0.082	28	31	0.857	0.150	0.380	0.708
Minimum dietary diversity (children 6–23 months)	0.343	0.060	82	92	1.144	0.176	0.222	0.464
Child slept under an ITN last night	0.729	0.041	329	379	1.504	0.057	0.647	0.812
Received 3+ doses of SP/Fansidar	0.390	0.056	115	128	1.219	0.143	0.278	0.501
Child has fever in last 2 weeks	0.129	0.025	276	305	1.256	0.194	0.079	0.179
Child had blood taken from finger/heel	0.485	0.088	35	39	1.046	0.181	0.310	0.660
Child took ACT	1.000	0.000	12	13	na	0.000	1.000	1.000
Condom use at last sex (women 15–24)	0.346	0.085	50	54	1.247	0.246	0.175	0.516
Mobile phone ownership	0.663	0.025	574	625	1.241	0.037	0.614	0.712

na = not applicable

Table B.44 Sampling errors: Busia sample, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases: Unweighted (N)	Number of cases: Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
HOUSEHOLDS AND POPULATION								
Primary reliance on clean fuels and technology	0.081	0.021	2,129	1,758	1.724	0.254	0.040	0.122
Births registered with civil authority	0.676	0.036	243	204	1.033	0.053	0.605	0.747
Ownership of at least one ITN	0.786	0.026	516	429	1.411	0.032	0.735	0.837
Ownership of at least one ITN for every two persons	0.550	0.029	512	426	1.320	0.053	0.492	0.608
At least basic drinking water service	0.680	0.074	2,129	1,758	3.024	0.109	0.532	0.828
At least basic sanitation service	0.428	0.036	2,129	1,758	1.405	0.083	0.356	0.499
Using open defecation	0.014	0.007	2,129	1,758	1.183	0.525	0.000	0.028
Using a handwashing facility with soap and water	0.942	0.022	1,885	1,567	1.668	0.023	0.898	0.985
WOMEN								
No education	0.019	0.006	441	373	0.981	0.340	0.006	0.031
Secondary education or higher	0.658	0.027	441	373	1.212	0.042	0.603	0.713
Literacy	0.941	0.012	441	373	1.072	0.013	0.917	0.965
Use of the internet in last 12 months	0.481	0.031	441	373	1.292	0.064	0.419	0.542
Total fertility rate (3 years)	3.567	0.226	1,222	1,037	0.837	0.063	3.114	4.020
Currently pregnant	0.051	0.011	441	373	1.058	0.217	0.029	0.073
Mean number of children ever born to women age 40–49	4.916	0.286	75	65	1.099	0.058	4.345	5.488
Median birth interval	48.226	3.050	131	111	0.973	0.063	42.126	54.327
Ideal number of children	3.691	0.075	434	367	0.899	0.020	3.540	3.841
Total wanted fertility rate (3 years)	2.879	0.180	1,222	1,037	0.933	0.063	2.519	3.239
Currently using any contraceptive method	0.591	0.042	246	212	1.320	0.070	0.508	0.674
Currently using any modern method	0.564	0.042	246	212	1.337	0.075	0.479	0.649
Currently using pill	0.028	0.012	246	212	1.134	0.426	0.004	0.052
Currently using injectables	0.145	0.021	246	212	0.920	0.143	0.104	0.187
Currently using implants	0.317	0.031	246	212	1.045	0.098	0.255	0.379
Currently using male condoms	0.007	0.005	246	212	0.989	0.738	0.000	0.018
Currently using any traditional method	0.026	0.012	246	212	1.179	0.458	0.002	0.051
Unmet need for spacing	0.080	0.016	246	212	0.931	0.202	0.047	0.112
Unmet need for limiting	0.128	0.024	246	212	1.137	0.189	0.080	0.177
Unmet need total	0.208	0.029	246	212	1.130	0.141	0.149	0.266
Demand satisfied by modern methods (married women)	0.707	0.034	196	169	1.038	0.048	0.639	0.774
Demand satisfied by modern methods (all women)	0.709	0.031	238	205	1.073	0.044	0.646	0.771
Not exposed to any of the ten media sources	0.052	0.013	441	373	1.270	0.258	0.025	0.079
Neonatal mortality (last 0–9 years)	24.541	10.571	394	338	1.231	0.431	3.399	45.683
Post-neonatal mortality (last 0–9 years)	5.443	3.842	393	337	1.027	0.706	0.000	13.128
Infant mortality (last 0–9 years)	29.984	11.243	394	338	1.128	0.375	7.497	52.471
Child mortality (last 0–9 years)	15.374	7.695	377	322	0.989	0.501	0.000	30.765
Under-five mortality (last 0–9 years)	44.898	16.820	395	339	1.335	0.375	11.258	78.538
Stillbirth rate	5.178	5.270	202	173	1.047	1.018	0.000	15.717
Early neonatal rate	25.007	12.329	204	174	0.961	0.493	0.350	49.665
Perinatal mortality rate	30.466	13.392	202	173	0.953	0.440	3.683	57.250
Received ANC from a skilled provider	0.988	0.012	89	78	0.994	0.012	0.964	1.000
4+ ANC visits	0.785	0.049	89	78	1.126	0.063	0.686	0.884
8+ ANC visits	0.099	0.030	89	78	0.944	0.303	0.039	0.160
Took any iron-containing supplements	0.903	0.035	89	78	1.092	0.038	0.833	0.972
Mothers protected against tetanus for last birth	0.694	0.042	89	78	0.864	0.061	0.609	0.779
Delivered in a health facility (live births)	0.916	0.036	89	78	1.220	0.039	0.845	0.987
Delivered by a skilled provider (live births)	0.928	0.031	89	78	1.143	0.034	0.865	0.990
Delivered by C-section (live births)	0.128	0.030	89	78	0.863	0.236	0.068	0.189
Women with postnatal check during first 2 days	0.913	0.027	89	78	0.889	0.029	0.860	0.966
Newborns with postnatal check during first 2 days	0.955	0.021	89	78	0.946	0.022	0.914	0.997
Any problem accessing health care	0.691	0.025	441	373	1.144	0.036	0.640	0.741
Ever had a vaccination card	1.000	0.000	45	40	na	0.000	1.000	1.000
Received BCG vaccination	0.981	0.020	45	40	1.001	0.020	0.942	1.000
Received DPT-HepB+Hib vaccination (3 doses)	0.871	0.050	45	40	1.012	0.057	0.772	0.970
Received pneumococcal vaccination (3 doses)	0.891	0.042	45	40	0.920	0.047	0.807	0.974
Received measles vaccination	0.839	0.062	45	40	1.156	0.074	0.715	0.963
Fully vaccinated according to national schedule (12–23 months)	0.527	0.052	45	40	0.713	0.099	0.423	0.631
Received measles/measles-rubella 2 vaccination (24–35 months)	0.797	0.055	46	39	0.917	0.069	0.686	0.907
Fully vaccinated according to national schedule (24–35 months)	0.425	0.061	46	39	0.798	0.142	0.304	0.546
Received HPV vaccination (2 doses) women 15–19	0.419	0.048	108	87	1.001	0.114	0.323	0.514
Sought treatment for diarrhoea	0.486	0.073	31	27	0.816	0.149	0.341	0.631
Treated with ORS	0.129	0.057	31	27	0.957	0.443	0.015	0.243
Exclusive breastfeeding	0.626	0.106	24	21	1.047	0.169	0.415	0.838
Minimum dietary diversity (children 6–23 months)	0.389	0.057	60	52	0.901	0.147	0.275	0.503
Child slept under an ITN last night	0.731	0.047	235	196	1.356	0.064	0.638	0.824
Received 3+ doses of SP/Fansidar	0.469	0.042	89	78	0.790	0.089	0.386	0.553
Child has fever in last 2 weeks	0.198	0.017	196	168	0.586	0.087	0.164	0.232
Child had blood taken from finger/heel	0.363	0.069	38	33	0.914	0.190	0.225	0.501
Child took ACT	1.000	0.000	7	6	na	0.000	1.000	1.000
Condom use at last sex (women 15–24)	0.284	0.120	25	21	1.285	0.423	0.044	0.523
Mobile phone ownership	0.727	0.027	441	373	1.265	0.037	0.673	0.780

na = not applicable

Table B.45 Sampling errors: Siaya sample, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases: Un-weighted (N)	Number of cases: Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
HOUSEHOLDS AND POPULATION								
Primary reliance on clean fuels and technology	0.060	0.011	2,262	1,804	1.161	0.183	0.038	0.081
Births registered with civil authority	0.496	0.032	252	195	0.863	0.065	0.431	0.560
Ownership of at least one ITN	0.687	0.025	609	482	1.336	0.037	0.637	0.737
Ownership of at least one ITN for every two persons	0.417	0.027	606	479	1.357	0.065	0.362	0.471
At least basic drinking water service	0.589	0.061	2,262	1,804	2.620	0.104	0.466	0.712
At least basic sanitation service	0.350	0.032	2,262	1,804	1.406	0.091	0.286	0.413
Using open defecation	0.067	0.027	2,262	1,804	2.301	0.398	0.014	0.120
Using a handwashing facility with soap and water	0.249	0.025	2,224	1,769	1.197	0.100	0.200	0.299
WOMEN								
No education	0.020	0.007	475	370	1.052	0.343	0.006	0.033
Secondary education or higher	0.553	0.022	475	370	0.942	0.039	0.510	0.596
Literacy	0.948	0.009	475	370	0.883	0.009	0.930	0.966
Use of the internet in last 12 months	0.451	0.038	475	370	1.653	0.084	0.376	0.527
Total fertility rate (3 years)	2.881	0.230	1,330	1,034	1.045	0.080	2.421	3.340
Currently pregnant	0.027	0.006	475	370	0.780	0.217	0.015	0.038
Mean number of children ever born to women age 40–49	4.766	0.371	89	69	1.427	0.078	4.025	5.507
Median birth interval	45.638	3.660	165	129	1.291	0.080	38.318	52.957
Ideal number of children	3.603	0.079	475	370	0.986	0.022	3.446	3.761
Total wanted fertility rate (3 years)	2.155	0.137	1,330	1,034	0.658	0.064	1.881	2.430
Currently using any contraceptive method	0.516	0.034	261	206	1.090	0.065	0.449	0.584
Currently using any modern method	0.513	0.034	261	206	1.085	0.066	0.446	0.580
Currently using pill	0.012	0.007	261	206	1.079	0.606	0.000	0.027
Currently using injectables	0.221	0.029	261	206	1.121	0.131	0.163	0.278
Currently using implants	0.206	0.024	261	206	0.967	0.118	0.157	0.254
Currently using male condoms	0.013	0.007	261	206	0.961	0.522	0.000	0.026
Currently using any traditional method	0.003	0.003	261	206	0.928	0.999	0.000	0.010
Unmet need for spacing	0.080	0.016	261	206	0.950	0.200	0.048	0.112
Unmet need for limiting	0.147	0.027	261	206	1.227	0.184	0.093	0.200
Unmet need total	0.226	0.026	261	206	1.002	0.115	0.174	0.278
Demand satisfied by modern methods (married women)	0.691	0.030	195	153	0.908	0.044	0.630	0.751
Demand satisfied by modern methods (all women)	0.694	0.031	254	201	1.082	0.045	0.632	0.756
Not exposed to any of the ten media sources	0.034	0.014	475	370	1.659	0.406	0.006	0.062
Neonatal mortality (last 0–9 years)	12.894	4.897	462	362	0.834	0.380	3.100	22.687
Post-neonatal mortality (last 0–9 years)	12.136	5.848	455	356	0.938	0.482	0.440	23.831
Infant mortality (last 0–9 years)	25.029	8.185	462	362	0.967	0.327	8.659	41.399
Child mortality (last 0–9 years)	13.040	5.009	454	359	1.018	0.384	3.022	23.059
Under-five mortality (last 0–9 years)	37.743	8.187	464	363	0.878	0.217	21.369	54.117
Stillbirth rate	0.000	0.000	208	161	na	na	0.000	0.000
Early neonatal rate	13.634	7.981	211	164	1.002	0.585	0.000	29.597
Perinatal mortality rate	13.849	8.083	208	161	0.999	0.584	0.000	30.015
Received ANC from a skilled provider	0.992	0.008	71	53	0.751	0.008	0.976	1.000
4+ ANC visits	0.702	0.071	71	53	1.290	0.101	0.561	0.844
8+ ANC visits	0.059	0.029	71	53	1.018	0.485	0.002	0.117
Took any iron-containing supplements	0.840	0.048	71	53	1.082	0.057	0.745	0.935
Mothers protected against tetanus for last birth	0.531	0.078	71	53	1.300	0.147	0.375	0.687
Delivered in a health facility (live births)	0.924	0.039	75	55	1.252	0.042	0.847	1.000
Delivered by a skilled provider (live births)	0.924	0.039	75	55	1.252	0.042	0.847	1.000
Delivered by C-section (live births)	0.132	0.047	75	55	1.197	0.360	0.037	0.227
Women with postnatal check during first 2 days	0.861	0.054	71	53	1.299	0.063	0.753	0.969
Newborns with postnatal check during first 2 days	0.869	0.052	71	53	1.281	0.060	0.765	0.973
Any problem accessing health care	0.686	0.038	475	370	1.790	0.056	0.609	0.762
Ever had a vaccination card	0.935	0.052	31	22	1.116	0.056	0.831	1.000
Received BCG vaccination	0.966	0.024	31	22	0.685	0.024	0.919	1.000
Received DPT-HepB+Hib vaccination (3 doses)	0.966	0.024	31	22	0.685	0.024	0.919	1.000
Received pneumococcal vaccination (3 doses)	0.921	0.040	31	22	0.783	0.043	0.841	1.000
Received measles vaccination	0.966	0.024	31	22	0.685	0.024	0.919	1.000
Fully vaccinated according to national schedule (12–23 months)	0.312	0.071	31	22	0.781	0.227	0.170	0.453
Received measles/measles-rubella 2 vaccination (24–35 months)	0.616	0.088	46	38	1.256	0.142	0.441	0.792
Fully vaccinated according to national schedule (24–35 months)	0.285	0.084	46	38	1.300	0.296	0.116	0.454
Received HPV vaccination (2 doses) women 15–19	0.428	0.067	96	76	1.323	0.157	0.293	0.563
Sought treatment for diarrhoea	0.707	0.148	12	9	1.093	0.209	0.411	1.000
Treated with ORS	0.646	0.158	12	9	1.111	0.245	0.330	0.961
Exclusive breastfeeding	0.865	0.098	22	16	1.290	0.113	0.670	1.000
Minimum dietary diversity (children 6–23 months)	0.289	0.060	48	36	0.912	0.208	0.168	0.409
Child slept under an ITN last night	0.561	0.039	250	194	1.076	0.069	0.483	0.638
Received 3+ doses of SP/Fansidar	0.370	0.073	71	53	1.257	0.197	0.224	0.515
Child has fever in last 2 weeks	0.048	0.019	207	161	1.262	0.389	0.011	0.086
Child had blood taken from finger/heel	0.053	0.055	9	8	0.772	1.036	0.000	0.162
Child took ACT	0.697	0.266	3	3	1.147	0.382	0.164	1.000
Condom use at last sex (women 15–24)	0.339	0.092	30	23	1.041	0.271	0.155	0.522
Mobile phone ownership	0.753	0.027	475	370	1.345	0.035	0.699	0.806

na = not applicable

Table B.46 Sampling errors: Kisumu sample, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases: Unweighted (N)	Number of cases: Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
HOUSEHOLDS AND POPULATION								
Primary reliance on clean fuels and technology	0.185	0.027	2,209	2,293	1.646	0.144	0.132	0.238
Births registered with civil authority	0.828	0.023	245	258	0.862	0.028	0.783	0.874
Ownership of at least one ITN	0.766	0.019	602	615	1.081	0.024	0.729	0.803
Ownership of at least one ITN for every two persons	0.479	0.027	595	609	1.332	0.057	0.424	0.534
At least basic drinking water service	0.814	0.026	2,209	2,293	1.365	0.032	0.762	0.865
At least basic sanitation service	0.363	0.043	2,209	2,293	1.857	0.118	0.278	0.449
Using open defecation	0.040	0.023	2,209	2,293	2.554	0.567	0.000	0.086
Using a handwashing facility with soap and water	0.491	0.029	2,095	2,189	1.209	0.060	0.432	0.550
WOMEN								
No education	0.011	0.004	535	547	0.928	0.388	0.002	0.019
Secondary education or higher	0.686	0.028	535	547	1.397	0.041	0.629	0.742
Literacy	0.968	0.008	538	551	1.026	0.008	0.953	0.984
Use of the internet in last 12 months	0.569	0.034	538	551	1.588	0.060	0.501	0.637
Total fertility rate (3 years)	3.167	0.276	1,518	1,558	1.478	0.087	2.616	3.719
Currently pregnant	0.036	0.009	538	551	1.098	0.245	0.018	0.054
Mean number of children ever born to women age 40–49	4.384	0.210	95	99	1.016	0.048	3.965	4.803
Median birth interval	50.896	3.212	180	186	1.132	0.063	44.473	57.319
Ideal number of children	3.608	0.104	538	551	1.457	0.029	3.401	3.815
Total wanted fertility rate (3 years)	2.441	0.241	1,518	1,558	1.559	0.099	1.959	2.924
Currently using any contraceptive method	0.740	0.022	281	296	0.842	0.030	0.696	0.784
Currently using any modern method	0.661	0.020	281	296	0.692	0.030	0.622	0.700
Currently using pill	0.042	0.012	281	296	0.970	0.278	0.018	0.065
Currently using injectables	0.233	0.031	281	296	1.237	0.134	0.170	0.295
Currently using implants	0.278	0.028	281	296	1.041	0.100	0.222	0.334
Currently using male condoms	0.043	0.013	281	296	1.092	0.306	0.017	0.070
Currently using any traditional method	0.079	0.013	281	296	0.778	0.159	0.054	0.104
Unmet need for spacing	0.068	0.016	281	296	1.066	0.235	0.036	0.101
Unmet need for limiting	0.048	0.014	281	296	1.133	0.301	0.019	0.077
Unmet need total	0.116	0.019	281	296	0.990	0.163	0.078	0.154
Demand satisfied by modern methods (married women)	0.772	0.019	242	254	0.686	0.024	0.734	0.809
Demand satisfied by modern methods (all women)	0.803	0.019	353	366	0.884	0.023	0.766	0.841
Not exposed to any of the ten media sources	0.057	0.007	538	551	0.706	0.124	0.043	0.071
Neonatal mortality (last 0–9 years)	8.958	3.801	509	536	0.938	0.424	1.356	16.559
Post-neonatal mortality (last 0–9 years)	10.215	4.556	513	540	0.993	0.446	1.104	19.327
Infant mortality (last 0–9 years)	19.173	4.625	509	536	0.727	0.241	9.924	28.422
Child mortality (last 0–9 years)	12.592	4.398	517	544	0.969	0.349	3.795	21.389
Under-five mortality (last 0–9 years)	31.523	6.147	510	537	0.744	0.195	19.230	43.817
Stillbirth rate	7.999	5.540	249	262	1.000	0.693	0.000	19.079
Early neonatal rate	4.236	4.257	247	260	1.042	1.005	0.000	12.750
Perinatal mortality rate	12.201	7.048	249	262	1.032	0.578	0.000	26.298
Received ANC from a skilled provider	0.983	0.012	92	101	0.907	0.013	0.958	1.000
4+ ANC visits	0.676	0.045	92	101	0.921	0.067	0.586	0.766
8+ ANC visits	0.107	0.019	92	101	0.604	0.182	0.068	0.146
Took any iron-containing supplements	0.980	0.014	92	101	0.960	0.014	0.952	1.000
Mothers protected against tetanus for last birth	0.842	0.031	92	101	0.803	0.036	0.781	0.904
Delivered in a health facility (live births)	0.957	0.029	96	106	1.418	0.030	0.900	1.000
Delivered by a skilled provider (live births)	0.967	0.021	96	106	1.189	0.022	0.924	1.000
Delivered by C-section (live births)	0.222	0.048	96	106	1.084	0.216	0.126	0.317
Women with postnatal check during first 2 days	0.891	0.033	92	101	1.022	0.037	0.825	0.958
Newborns with postnatal check during first 2 days	0.907	0.036	92	101	1.184	0.040	0.835	0.979
Any problem accessing health care	0.707	0.028	538	551	1.430	0.040	0.651	0.763
Ever had a vaccination card	1.000	0.000	47	52	na	0.000	1.000	1.000
Received BCG vaccination	0.979	0.021	47	52	1.007	0.021	0.937	1.000
Received DPT-HepB+Hib vaccination (3 doses)	0.901	0.039	47	52	0.924	0.044	0.822	0.980
Received pneumococcal vaccination (3 doses)	0.958	0.030	47	52	1.046	0.031	0.899	1.000
Received measles vaccination	1.000	0.000	47	52	na	0.000	1.000	1.000
Fully vaccinated according to national schedule (12–23 months)	0.701	0.059	47	52	0.910	0.085	0.582	0.820
Received measles/measles-rubella 2 vaccination (24–35 months)	0.798	0.041	56	60	0.771	0.051	0.716	0.880
Fully vaccinated according to national schedule (24–35 months)	0.426	0.081	56	60	1.214	0.189	0.265	0.587
Received HPV vaccination (2 doses) women 15–19	0.457	0.052	106	106	1.067	0.113	0.354	0.561
Sought treatment for diarrhoea	0.679	0.100	38	42	1.354	0.147	0.479	0.879
Treated with ORS	0.471	0.092	38	42	1.172	0.196	0.286	0.656
Exclusive breastfeeding	0.669	0.099	26	28	1.053	0.148	0.471	0.868
Minimum dietary diversity (children 6–23 months)	0.636	0.052	62	68	0.844	0.082	0.532	0.740
Child slept under an ITN last night	0.689	0.035	251	264	1.076	0.051	0.619	0.758
Received 3+ doses of SP/Fansidar	0.500	0.073	92	101	1.378	0.145	0.355	0.646
Child has fever in last 2 weeks	0.222	0.029	243	256	1.003	0.129	0.165	0.279
Child had blood taken from finger/heel	0.448	0.064	55	57	0.873	0.142	0.321	0.575
Child took ACT	0.856	0.089	19	21	1.126	0.103	0.679	1.000
Condom use at last sex (women 15–24)	0.539	0.065	65	62	1.039	0.120	0.409	0.668
Mobile phone ownership	0.807	0.024	538	551	1.423	0.030	0.759	0.856

na = not applicable

Table B.47 Sampling errors: Homa Bay sample, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases Unweighted (N)	Number of cases Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits Lower (R-2SE)	Confidence limits Upper (R+2SE)
HOUSEHOLDS AND POPULATION								
Primary reliance on clean fuels and technology	0.061	0.020	2,519	2,246	2.170	0.324	0.021	0.100
Births registered with civil authority	0.900	0.025	299	265	1.196	0.027	0.851	0.949
Ownership of at least one ITN	0.713	0.040	649	568	2.217	0.055	0.634	0.792
Ownership of at least one ITN for every two persons	0.375	0.035	648	567	1.852	0.094	0.305	0.446
At least basic drinking water service	0.627	0.061	2,519	2,246	2.787	0.098	0.504	0.750
At least basic sanitation service	0.195	0.030	2,519	2,246	1.750	0.155	0.134	0.256
Using open defecation	0.112	0.037	2,519	2,246	2.467	0.325	0.039	0.186
Using a handwashing facility with soap and water	0.736	0.023	2,519	2,246	1.142	0.031	0.690	0.781
WOMEN								
No education	0.003	0.002	569	484	0.950	0.712	0.000	0.008
Secondary education or higher	0.617	0.031	569	484	1.496	0.049	0.556	0.679
Literacy	0.937	0.013	570	485	1.240	0.013	0.912	0.962
Use of the internet in last 12 months	0.503	0.034	570	485	1.615	0.067	0.435	0.571
Total fertility rate (3 years)	3.349	0.233	1,561	1,322	0.960	0.070	2.882	3.815
Currently pregnant	0.051	0.009	570	485	0.959	0.173	0.034	0.069
Mean number of children ever born to women age 40–49	5.023	0.289	90	82	1.186	0.058	4.444	5.602
Median birth interval	45.282	3.330	211	180	1.291	0.074	38.622	51.942
Ideal number of children	4.103	0.082	570	485	1.070	0.020	3.940	4.266
Total wanted fertility rate (3 years)	2.595	0.195	1,561	1,322	0.992	0.075	2.205	2.984
Currently using any contraceptive method	0.689	0.037	325	279	1.425	0.053	0.615	0.762
Currently using any modern method	0.653	0.035	325	279	1.319	0.053	0.583	0.723
Currently using pill	0.033	0.012	325	279	1.239	0.372	0.009	0.058
Currently using injectables	0.213	0.029	325	279	1.267	0.135	0.155	0.271
Currently using implants	0.324	0.023	325	279	0.878	0.070	0.279	0.370
Currently using male condoms	0.038	0.014	325	279	1.283	0.360	0.011	0.065
Currently using any traditional method	0.036	0.014	325	279	1.363	0.395	0.007	0.064
Unmet need for spacing	0.083	0.019	325	279	1.261	0.233	0.044	0.122
Unmet need for limiting	0.047	0.014	325	279	1.213	0.302	0.019	0.076
Unmet need total	0.131	0.025	325	279	1.351	0.194	0.080	0.181
Demand satisfied by modern methods (married women)	0.797	0.034	266	228	1.387	0.043	0.729	0.866
Demand satisfied by modern methods (all women)	0.785	0.029	363	313	1.328	0.036	0.728	0.842
Not exposed to any of the ten media sources	0.010	0.005	570	485	1.133	0.484	0.000	0.019
Neonatal mortality (last 0–9 years)	25.385	7.674	586	503	1.020	0.302	10.037	40.733
Post-neonatal mortality (last 0–9 years)	15.403	5.099	590	507	1.012	0.331	5.204	25.602
Infant mortality (last 0–9 years)	40.788	10.229	588	505	1.147	0.251	20.331	61.245
Child mortality (last 0–9 years)	12.228	6.228	612	524	1.270	0.509	0.000	24.685
Under-five mortality (last 0–9 years)	52.518	12.030	590	507	1.242	0.229	28.457	76.578
Stillbirth rate	4.586	4.498	290	250	1.139	0.981	0.000	13.582
Early neonatal rate	31.243	12.077	289	249	1.069	0.387	7.089	55.397
Perinatal mortality rate	35.686	15.165	290	250	1.280	0.425	5.357	66.015
Received ANC from a skilled provider	1.000	0.000	93	81	na	0.000	1.000	1.000
4+ ANC visits	0.736	0.055	93	81	1.198	0.075	0.626	0.846
8+ ANC visits	0.115	0.027	93	81	0.817	0.236	0.061	0.169
Took any iron-containing supplements	0.941	0.024	93	81	0.994	0.026	0.892	0.990
Mothers protected against tetanus for last birth	0.801	0.044	93	81	1.051	0.055	0.713	0.888
Delivered in a health facility (live births)	0.907	0.031	98	86	0.946	0.034	0.845	0.969
Delivered by a skilled provider (live births)	0.907	0.031	98	86	0.946	0.034	0.845	0.969
Delivered by C-section (live births)	0.190	0.049	98	86	1.133	0.258	0.092	0.288
Women with postnatal check during first 2 days	0.922	0.033	93	81	1.179	0.036	0.855	0.988
Newborns with postnatal check during first 2 days	0.891	0.030	93	81	0.912	0.033	0.832	0.950
Any problem accessing health care	0.789	0.029	570	485	1.684	0.037	0.732	0.847
Ever had a vaccination card	1.000	0.000	44	38	na	0.000	1.000	1.000
Received BCG vaccination	0.960	0.039	44	38	1.318	0.041	0.881	1.000
Received DPT-HepB+Hib vaccination (3 doses)	0.761	0.071	44	38	1.018	0.094	0.618	0.903
Received pneumococcal vaccination (3 doses)	0.915	0.041	44	38	0.966	0.045	0.833	0.997
Received measles vaccination	0.765	0.045	44	38	0.689	0.058	0.676	0.854
Fully vaccinated according to national schedule (12–23 months)	0.322	0.070	44	38	0.974	0.217	0.182	0.462
Received measles/measles-rubella 2 vaccination (24–35 months)	0.689	0.070	63	53	1.166	0.101	0.550	0.829
Fully vaccinated according to national schedule (24–35 months)	0.233	0.062	63	53	1.145	0.267	0.109	0.358
Received HPV vaccination (2 doses) women 15–19	0.439	0.057	153	131	1.410	0.130	0.325	0.553
Sought treatment for diarrhoea	0.504	0.102	38	34	1.286	0.202	0.300	0.708
Treated with ORS	0.440	0.112	38	34	1.418	0.254	0.217	0.664
Exclusive breastfeeding	0.454	0.130	18	15	1.074	0.287	0.194	0.715
Minimum dietary diversity (children 6–23 months)	0.514	0.065	66	58	1.043	0.126	0.385	0.643
Child slept under an ITN last night	0.692	0.037	299	263	1.213	0.053	0.618	0.765
Received 3+ doses of SP/Fansidar	0.549	0.062	93	81	1.186	0.112	0.425	0.672
Child has fever in last 2 weeks	0.331	0.029	275	237	0.994	0.088	0.272	0.389
Child had blood taken from finger/heel	0.502	0.056	89	78	1.019	0.112	0.389	0.615
Child took ACT	0.898	0.059	26	25	1.034	0.065	0.780	1.000
Condom use at last sex (women 15–24)	0.474	0.065	66	57	1.049	0.137	0.344	0.604
Mobile phone ownership	0.720	0.033	570	485	1.750	0.046	0.654	0.786

na = not applicable

Table B.48 Sampling errors: Migori sample, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases: Un-weighted (N)	Number of cases: Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
HOUSEHOLDS AND POPULATION								
Primary reliance on clean fuels and technology	0.129	0.024	2,542	2,240	1.708	0.183	0.082	0.177
Births registered with civil authority	0.900	0.023	374	335	1.280	0.025	0.855	0.946
Ownership of at least one ITN	0.802	0.022	580	499	1.333	0.028	0.758	0.846
Ownership of at least one ITN for every two persons	0.484	0.034	580	499	1.654	0.071	0.415	0.553
At least basic drinking water service	0.662	0.039	2,542	2,240	1.757	0.060	0.584	0.741
At least basic sanitation service	0.346	0.043	2,542	2,240	1.879	0.123	0.261	0.432
Using open defecation	0.047	0.023	2,542	2,240	2.271	0.501	0.000	0.093
Using a handwashing facility with soap and water	0.517	0.020	2,523	2,224	0.873	0.040	0.476	0.558
WOMEN								
No education	0.011	0.007	586	509	1.701	0.657	0.000	0.026
Secondary education or higher	0.607	0.036	586	509	1.789	0.060	0.535	0.679
Literacy	0.937	0.009	586	509	0.945	0.010	0.919	0.956
Use of the internet in last 12 months	0.575	0.033	586	509	1.615	0.057	0.509	0.641
Total fertility rate (3 years)	3.814	0.251	1,646	1,433	1.280	0.066	3.311	4.317
Currently pregnant	0.072	0.017	586	509	1.631	0.243	0.037	0.106
Mean number of children ever born to women age 40–49	5.670	0.260	85	78	1.147	0.046	5.149	6.191
Median birth interval	50.042	2.083	256	225	0.916	0.042	45.876	54.209
Ideal number of children	4.012	0.131	585	508	1.911	0.033	3.751	4.274
Total wanted fertility rate (3 years)	3.180	0.326	1,646	1,433	1.654	0.103	2.527	3.833
Currently using any contraceptive method	0.605	0.038	361	314	1.472	0.063	0.529	0.681
Currently using any modern method	0.570	0.041	361	314	1.560	0.071	0.489	0.652
Currently using pill	0.025	0.007	361	314	0.918	0.305	0.010	0.040
Currently using injectables	0.137	0.021	361	314	1.130	0.149	0.096	0.178
Currently using implants	0.320	0.027	361	314	1.108	0.085	0.266	0.375
Currently using male condoms	0.021	0.008	361	314	1.129	0.409	0.004	0.038
Currently using any traditional method	0.034	0.008	361	314	0.819	0.229	0.019	0.050
Unmet need for spacing	0.100	0.023	361	314	1.429	0.227	0.055	0.145
Unmet need for limiting	0.067	0.013	361	314	1.012	0.198	0.041	0.094
Unmet need total	0.167	0.026	361	314	1.319	0.155	0.115	0.219
Demand satisfied by modern methods (married women)	0.739	0.040	280	242	1.501	0.054	0.660	0.818
Demand satisfied by modern methods (all women)	0.735	0.035	375	325	1.528	0.047	0.665	0.805
Not exposed to any of the ten media sources	0.018	0.005	586	509	0.809	0.244	0.009	0.027
Neonatal mortality (last 0–9 years)	17.042	6.547	689	608	1.149	0.384	3.948	30.136
Post-neonatal mortality (last 0–9 years)	10.467	4.030	684	604	1.031	0.385	2.408	18.526
Infant mortality (last 0–9 years)	27.509	7.597	689	608	1.117	0.276	12.314	42.704
Child mortality (last 0–9 years)	11.083	3.692	661	583	0.887	0.333	3.698	18.467
Under-five mortality (last 0–9 years)	38.287	9.112	692	611	1.163	0.238	20.063	56.511
Stillbirth rate	11.044	5.272	370	328	0.989	0.477	0.499	21.589
Early neonatal rate	17.374	8.841	368	327	1.162	0.509	0.000	35.057
Perinatal mortality rate	28.325	11.224	370	328	1.229	0.396	5.877	50.773
Received ANC from a skilled provider	1.000	0.000	146	131	na	0.000	1.000	1.000
4+ ANC visits	0.704	0.035	146	131	0.914	0.049	0.635	0.774
8+ ANC visits	0.093	0.022	146	131	0.899	0.233	0.049	0.136
Took any iron-containing supplements	0.901	0.029	146	131	1.173	0.032	0.842	0.959
Mothers protected against tetanus for last birth	0.619	0.046	146	131	1.147	0.075	0.527	0.712
Delivered in a health facility (live births)	0.932	0.020	155	139	1.004	0.021	0.892	0.972
Delivered by a skilled provider (live births)	0.935	0.020	155	139	1.036	0.022	0.895	0.975
Delivered by C-section (live births)	0.143	0.037	155	139	1.277	0.259	0.069	0.216
Women with postnatal check during first 2 days	0.973	0.013	146	131	0.967	0.013	0.947	0.999
Newborns with postnatal check during first 2 days	0.945	0.019	146	131	1.017	0.020	0.906	0.983
Any problem accessing health care	0.528	0.035	586	509	1.699	0.067	0.458	0.598
Ever had a vaccination card	1.000	0.000	78	71	na	0.000	1.000	1.000
Received BCG vaccination	1.000	0.000	78	71	na	0.000	1.000	1.000
Received DPT-HepB+Hib vaccination (3 doses)	0.979	0.016	78	71	1.055	0.017	0.946	1.000
Received pneumococcal vaccination (3 doses)	0.954	0.019	78	71	0.830	0.020	0.915	0.992
Received measles vaccination	0.931	0.029	78	71	1.046	0.031	0.873	0.990
Fully vaccinated according to national schedule (12–23 months)	0.557	0.049	78	71	0.888	0.087	0.460	0.654
Received measles/measles-rubella 2 vaccination (24–35 months)	0.630	0.051	54	50	0.800	0.081	0.528	0.732
Fully vaccinated according to national schedule (24–35 months)	0.484	0.054	54	50	0.816	0.112	0.376	0.593
Received HPV vaccination (2 doses) women 15–19	0.363	0.056	109	96	1.199	0.153	0.252	0.474
Sought treatment for diarrhoea	0.477	0.051	67	62	0.857	0.106	0.375	0.578
Treated with ORS	0.258	0.042	67	62	0.810	0.163	0.174	0.342
Exclusive breastfeeding	0.776	0.067	31	29	0.877	0.086	0.643	0.909
Minimum dietary diversity (children 6–23 months)	0.343	0.059	111	99	1.303	0.172	0.225	0.461
Child slept under an ITN last night	0.702	0.027	389	349	1.054	0.039	0.647	0.756
Received 3+ doses of SP/Fansidar	0.393	0.036	146	131	0.878	0.091	0.322	0.464
Child has fever in last 2 weeks	0.257	0.023	360	320	0.984	0.091	0.211	0.304
Child had blood taken from finger/heel	0.432	0.054	94	82	1.032	0.125	0.323	0.540
Child took ACT	0.884	0.052	43	42	1.117	0.059	0.780	0.988
Condom use at last sex (women 15–24)	0.371	0.074	51	46	1.086	0.200	0.222	0.519
Mobile phone ownership	0.745	0.021	586	509	1.168	0.028	0.702	0.787

na = not applicable

Table B.49 Sampling errors: Kisii sample, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases: Unweighted (N)	Number of cases: Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
HOUSEHOLDS AND POPULATION								
Primary reliance on clean fuels and technology	0.109	0.016	2,171	2,210	1.249	0.146	0.077	0.141
Births registered with civil authority	0.873	0.035	235	237	1.353	0.040	0.802	0.944
Ownership of at least one ITN	0.546	0.040	603	597	1.953	0.073	0.466	0.625
Ownership of at least one ITN for every two persons	0.374	0.025	600	593	1.247	0.066	0.325	0.424
At least basic drinking water service	0.660	0.055	2,171	2,210	2.454	0.083	0.550	0.770
At least basic sanitation service	0.376	0.039	2,171	2,210	1.789	0.105	0.297	0.455
Using open defecation	0.000	0.000	2,171	2,210	na	na	0.000	0.000
Using a handwashing facility with soap and water	0.601	0.030	2,148	2,188	1.328	0.050	0.541	0.661
WOMEN								
No education	0.008	0.004	442	439	0.908	0.495	0.000	0.015
Secondary education or higher	0.707	0.035	442	439	1.594	0.049	0.638	0.776
Literacy	0.965	0.009	442	439	1.080	0.010	0.946	0.984
Use of the internet in last 12 months	0.403	0.035	442	439	1.510	0.088	0.332	0.473
Total fertility rate (3 years)	2.896	0.260	1,257	1,238	1.111	0.090	2.376	3.416
Currently pregnant	0.019	0.007	442	439	1.032	0.353	0.006	0.032
Mean number of children ever born to women age 40–49	4.394	0.225	82	87	1.054	0.051	3.945	4.843
Median birth interval	43.641	3.431	154	157	1.089	0.079	36.780	50.502
Ideal number of children	2.948	0.074	442	439	1.301	0.025	2.801	3.095
Total wanted fertility rate (3 years)	1.883	0.191	1,257	1,238	0.900	0.101	1.501	2.265
Currently using any contraceptive method	0.713	0.038	254	253	1.337	0.053	0.637	0.789
Currently using any modern method	0.687	0.037	254	253	1.268	0.054	0.613	0.761
Currently using pill	0.062	0.012	254	253	0.810	0.198	0.037	0.086
Currently using injectables	0.310	0.029	254	253	0.986	0.093	0.252	0.367
Currently using implants	0.240	0.023	254	253	0.864	0.097	0.194	0.286
Currently using male condoms	0.016	0.008	254	253	1.068	0.534	0.000	0.032
Currently using any traditional method	0.025	0.011	254	253	1.116	0.434	0.003	0.048
Unmet need for spacing	0.023	0.009	254	253	0.939	0.389	0.005	0.040
Unmet need for limiting	0.100	0.022	254	253	1.172	0.221	0.056	0.145
Unmet need total	0.123	0.021	254	253	1.041	0.175	0.080	0.166
Demand satisfied by modern methods (married women)	0.822	0.025	210	212	0.969	0.031	0.772	0.873
Demand satisfied by modern methods (all women)	0.815	0.026	261	263	1.089	0.032	0.763	0.867
Not exposed to any of the ten media sources	0.030	0.009	442	439	1.050	0.283	0.013	0.047
Neonatal mortality (last 0–9 years)	26.671	10.578	397	404	1.403	0.397	5.515	47.827
Post-neonatal mortality (last 0–9 years)	12.171	5.521	399	407	1.002	0.454	1.129	23.213
Infant mortality (last 0–9 years)	38.842	10.391	397	404	1.188	0.268	18.061	59.623
Child mortality (last 0–9 years)	2.833	2.848	402	407	1.083	1.005	0.000	8.529
Under-five mortality (last 0–9 years)	41.565	10.179	398	405	1.137	0.245	21.207	61.922
Stillbirth rate	13.200	10.323	218	218	1.337	0.782	0.000	33.845
Early neonatal rate	13.023	7.562	214	215	0.991	0.581	0.000	28.146
Perinatal mortality rate	26.051	11.980	218	218	1.124	0.460	2.091	50.011
Received ANC from a skilled provider	0.984	0.017	81	84	1.169	0.017	0.950	1.000
4+ ANC visits	0.578	0.058	81	84	1.057	0.101	0.462	0.695
8+ ANC visits	0.034	0.020	81	84	0.984	0.588	0.000	0.074
Took any iron-containing supplements	0.922	0.034	81	84	1.123	0.037	0.854	0.989
Mothers protected against tetanus for last birth	0.714	0.059	81	84	1.168	0.083	0.596	0.833
Delivered in a health facility (live births)	0.945	0.024	81	84	0.955	0.025	0.897	0.993
Delivered by a skilled provider (live births)	0.945	0.024	81	84	0.955	0.025	0.897	0.993
Delivered by C-section (live births)	0.215	0.040	81	84	0.877	0.185	0.136	0.294
Women with postnatal check during first 2 days	0.923	0.031	81	84	1.048	0.034	0.861	0.986
Newborns with postnatal check during first 2 days	0.869	0.033	81	84	0.876	0.038	0.803	0.935
Any problem accessing health care	0.757	0.031	442	439	1.528	0.041	0.694	0.819
Ever had a vaccination card	1.000	0.000	36	36	na	0.000	1.000	1.000
Received BCG vaccination	0.967	0.034	36	36	1.123	0.035	0.899	1.000
Received DPT-HepB+Hib vaccination (3 doses)	0.938	0.042	36	36	1.023	0.045	0.854	1.000
Received pneumococcal vaccination (3 doses)	0.939	0.043	36	36	1.053	0.046	0.853	1.000
Received measles vaccination	0.851	0.066	36	36	1.093	0.077	0.720	0.983
Fully vaccinated according to national schedule (12–23 months)	0.312	0.101	36	36	1.289	0.324	0.110	0.515
Received measles/measles-rubella 2 vaccination (24–35 months)	0.727	0.075	35	36	1.012	0.104	0.577	0.878
Fully vaccinated according to national schedule (24–35 months)	0.410	0.086	35	36	1.045	0.210	0.238	0.582
Received HPV vaccination (2 doses) women 15–19	0.273	0.052	82	85	1.050	0.190	0.169	0.378
Sought treatment for diarrhoea	0.576	0.125	10	10	0.812	0.217	0.326	0.826
Treated with ORS	0.513	0.164	10	10	1.048	0.319	0.186	0.840
Exclusive breastfeeding	0.736	0.126	15	14	1.067	0.172	0.483	0.989
Minimum dietary diversity (children 6–23 months)	0.440	0.055	62	64	0.871	0.126	0.329	0.550
Child slept under an ITN last night	0.539	0.040	247	249	1.067	0.073	0.460	0.618
Received 3+ doses of SP/Fansidar	0.012	0.012	81	84	1.000	0.999	0.000	0.037
Child has fever in last 2 weeks	0.075	0.023	207	208	1.253	0.309	0.029	0.121
Child had blood taken from finger/heel	0.411	0.164	14	15	1.299	0.400	0.082	0.739
Child took ACT	0.475	0.364	2	2	1.064	0.765	0.000	1.000
Condom use at last sex (women 15–24)	0.367	0.073	46	43	1.014	0.199	0.221	0.513
Mobile phone ownership	0.721	0.029	442	439	1.376	0.041	0.662	0.779

na = not applicable

Table B.50 Sampling errors: Nyamira sample, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases: Unweighted (N)	Number of cases: Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
HOUSEHOLDS AND POPULATION								
Primary reliance on clean fuels and technology	0.075	0.023	2,017	1,050	1.992	0.303	0.030	0.121
Births registered with civil authority	0.921	0.024	200	106	1.258	0.026	0.874	0.969
Ownership of at least one ITN	0.782	0.033	546	284	1.861	0.042	0.716	0.848
Ownership of at least one ITN for every two persons	0.578	0.033	546	284	1.547	0.057	0.512	0.643
At least basic drinking water service	0.464	0.034	2,017	1,050	1.425	0.073	0.396	0.533
At least basic sanitation service	0.418	0.036	2,017	1,050	1.521	0.086	0.346	0.490
Using open defecation	0.000	0.000	2,017	1,050	na	na	0.000	0.000
Using a handwashing facility with soap and water	0.663	0.011	2,017	1,050	0.497	0.017	0.640	0.685
WOMEN								
No education	0.002	0.002	453	237	0.976	1.014	0.000	0.006
Secondary education or higher	0.772	0.023	453	237	1.162	0.030	0.726	0.817
Literacy	0.949	0.016	454	238	1.591	0.017	0.916	0.982
Use of the internet in last 12 months	0.421	0.025	454	238	1.094	0.060	0.370	0.472
Total fertility rate (3 years)	3.161	0.308	1,251	652	1.140	0.097	2.546	3.776
Currently pregnant	0.042	0.010	454	238	1.069	0.239	0.022	0.062
Mean number of children ever born to women age 40–49	4.253	0.220	98	51	1.156	0.052	3.814	4.692
Median birth interval	50.169	2.954	135	71	0.985	0.059	44.261	56.076
Ideal number of children	2.864	0.056	454	238	0.839	0.019	2.753	2.976
Total wanted fertility rate (3 years)	2.124	0.253	1,251	652	1.225	0.119	1.618	2.630
Currently using any contraceptive method	0.690	0.035	239	123	1.151	0.050	0.621	0.759
Currently using any modern method	0.590	0.035	239	123	1.090	0.059	0.520	0.659
Currently using pill	0.032	0.014	239	123	1.220	0.435	0.004	0.060
Currently using injectables	0.305	0.032	239	123	1.065	0.104	0.241	0.369
Currently using implants	0.209	0.028	239	123	1.065	0.134	0.153	0.265
Currently using male condoms	0.005	0.005	239	123	1.066	0.994	0.000	0.014
Currently using any traditional method	0.100	0.024	239	123	1.208	0.235	0.053	0.147
Unmet need for spacing	0.052	0.014	239	123	0.974	0.271	0.024	0.080
Unmet need for limiting	0.067	0.018	239	123	1.088	0.262	0.032	0.103
Unmet need total	0.119	0.018	239	123	0.847	0.149	0.084	0.155
Demand satisfied by modern methods (married women)	0.729	0.031	194	100	0.974	0.043	0.667	0.791
Demand satisfied by modern methods (all women)	0.717	0.028	271	140	1.025	0.039	0.660	0.774
Not exposed to any of the ten media sources	0.085	0.016	454	238	1.245	0.192	0.053	0.118
Neonatal mortality (last 0–9 years)	7.074	4.081	370	194	0.941	0.577	0.000	15.236
Post-neonatal mortality (last 0–9 years)	20.902	7.876	369	193	1.050	0.377	5.150	36.653
Infant mortality (last 0–9 years)	27.976	8.810	370	194	1.032	0.315	10.355	45.597
Child mortality (last 0–9 years)	8.803	4.767	370	192	0.948	0.542	0.000	18.337
Under-five mortality (last 0–9 years)	36.532	9.069	370	194	0.929	0.248	18.394	54.671
Stillbirth rate	33.440	15.686	194	103	0.971	0.469	2.067	64.813
Early neonatal rate	5.356	5.290	188	99	0.995	0.988	0.000	15.936
Perinatal mortality rate	38.617	15.713	194	103	0.933	0.407	7.191	70.043
Received ANC from a skilled provider	0.989	0.011	74	40	0.941	0.012	0.966	1.000
4+ ANC visits	0.674	0.064	74	40	1.172	0.096	0.545	0.803
8+ ANC visits	0.087	0.034	74	40	1.031	0.391	0.019	0.155
Took any iron-containing supplements	0.806	0.043	74	40	0.924	0.053	0.720	0.891
Mothers protected against tetanus for last birth	0.710	0.066	74	40	1.233	0.092	0.579	0.842
Delivered in a health facility (live births)	0.977	0.017	78	42	0.984	0.017	0.943	1.000
Delivered by a skilled provider (live births)	0.977	0.017	78	42	0.984	0.017	0.943	1.000
Delivered by C-section (live births)	0.101	0.027	78	42	0.797	0.267	0.047	0.155
Women with postnatal check during first 2 days	0.748	0.063	74	40	1.230	0.084	0.622	0.873
Newborns with postnatal check during first 2 days	0.805	0.054	74	40	1.155	0.067	0.697	0.912
Any problem accessing health care	0.712	0.032	454	238	1.479	0.044	0.648	0.775
Ever had a vaccination card	0.989	0.011	43	23	0.699	0.011	0.968	1.000
Received BCG vaccination	0.989	0.011	43	23	0.699	0.011	0.968	1.000
Received DPT-HepB+Hib vaccination (3 doses)	0.880	0.053	43	23	1.079	0.061	0.773	0.986
Received pneumococcal vaccination (3 doses)	0.989	0.011	43	23	0.699	0.011	0.968	1.000
Received measles vaccination	0.953	0.033	43	23	1.018	0.034	0.887	1.000
Fully vaccinated according to national schedule (12–23 months)	0.599	0.087	43	23	1.120	0.145	0.426	0.772
Received measles/measles-rubella 2 vaccination (24–35 months)	0.768	0.066	38	20	0.967	0.085	0.637	0.900
Fully vaccinated according to national schedule (24–35 months)	0.523	0.105	38	20	1.310	0.202	0.312	0.733
Received HPV vaccination (2 doses) women 15–19	0.453	0.044	109	61	0.921	0.097	0.365	0.541
Sought treatment for diarrhoea	0.393	0.101	19	10	0.878	0.257	0.191	0.595
Treated with ORS	0.341	0.074	19	10	0.669	0.218	0.192	0.489
Exclusive breastfeeding	0.603	0.140	16	9	1.101	0.232	0.323	0.883
Minimum dietary diversity (children 6–23 months)	0.276	0.065	55	29	1.075	0.237	0.145	0.406
Child slept under an ITN last night	0.757	0.041	201	106	1.231	0.054	0.675	0.839
Received 3+ doses of SP/Fansidar	0.059	0.033	74	40	1.191	0.557	0.000	0.125
Child has fever in last 2 weeks	0.086	0.030	181	96	1.424	0.348	0.026	0.146
Child had blood taken from finger/heel	0.082	0.077	16	8	1.117	0.940	0.000	0.236
Child took ACT	1.000	0.000	1	1	na	0.000	1.000	1.000
Condom use at last sex (women 15–24)	0.364	0.067	51	27	0.982	0.184	0.230	0.497
Mobile phone ownership	0.686	0.030	454	238	1.383	0.044	0.625	0.746

na = not applicable

Table B.51 Sampling errors: Nairobi City sample, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases: Unweighted (N)	Number of cases: Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
HOUSEHOLDS AND POPULATION								
Primary reliance on clean fuels and technology	0.813	0.038	2,439	12,891	2.189	0.046	0.738	0.888
Births registered with civil authority	0.817	0.039	290	1,582	1.540	0.048	0.739	0.894
Ownership of at least one ITN	0.152	0.015	746	3,992	1.134	0.098	0.123	0.182
Ownership of at least one ITN for every two persons	0.086	0.010	731	3,896	0.976	0.118	0.066	0.106
At least basic drinking water service	0.973	0.010	2,439	12,891	1.096	0.010	0.953	0.993
At least basic sanitation service	0.561	0.077	2,439	12,891	3.613	0.137	0.407	0.715
Using open defecation	0.000	0.000	2,439	12,891	na	na	0.000	0.000
Using a handwashing facility with soap and water	0.658	0.037	2,355	12,430	1.724	0.055	0.585	0.731
WOMEN								
No education	0.051	0.028	597	3,521	3.085	0.550	0.000	0.107
Secondary education or higher	0.716	0.042	597	3,521	2.274	0.059	0.631	0.800
Literacy	0.930	0.032	602	3,550	3.023	0.034	0.867	0.994
Use of the internet in last 12 months	0.874	0.021	602	3,550	1.523	0.024	0.833	0.915
Total fertility rate (3 years)	2.594	0.246	1,741	0,313	1.135	0.095	2.103	3.085
Currently pregnant	0.040	0.010	602	3,550	1.197	0.240	0.021	0.059
Mean number of children ever born to women age 40–49	3.423	0.240	97	571	1.267	0.070	2.943	3.904
Median birth interval	49.220	4.978	185	1,109	1.479	0.101	39.265	59.176
Ideal number of children	3.641	0.235	594	3,505	2.632	0.064	3.172	4.110
Total wanted fertility rate (3 years)	2.470	0.245	1,741	0,313	1.148	0.099	1.981	2.960
Currently using any contraceptive method	0.615	0.053	318	1,920	1.918	0.086	0.510	0.721
Currently using any modern method	0.546	0.048	318	1,920	1.706	0.088	0.451	0.642
Currently using pill	0.124	0.029	318	1,920	1.550	0.232	0.067	0.182
Currently using injectables	0.152	0.026	318	1,920	1.310	0.174	0.099	0.205
Currently using implants	0.194	0.027	318	1,920	1.221	0.140	0.139	0.248
Currently using male condoms	0.027	0.009	318	1,920	0.945	0.318	0.010	0.044
Currently using any traditional method	0.069	0.014	318	1,920	0.985	0.203	0.041	0.097
Unmet need for spacing	0.051	0.019	318	1,920	1.496	0.361	0.014	0.089
Unmet need for limiting	0.077	0.019	318	1,920	1.256	0.245	0.039	0.114
Unmet need total	0.128	0.031	318	1,920	1.663	0.244	0.066	0.191
Demand satisfied by modern methods (married women)	0.735	0.042	235	1,428	1.468	0.058	0.650	0.819
Demand satisfied by modern methods (all women)	0.735	0.033	324	1,921	1.351	0.045	0.669	0.801
Not exposed to any of the ten media sources	0.113	0.025	602	3,550	1.942	0.222	0.063	0.163
Neonatal mortality (last 0–9 years)	19.055	7.073	535	3,186	0.966	0.371	4.909	33.201
Post-neonatal mortality (last 0–9 years)	12.683	6.444	537	3,189	1.315	0.508	0.000	25.571
Infant mortality (last 0–9 years)	31.738	10.450	535	3,186	1.171	0.329	10.838	52.638
Child mortality (last 0–9 years)	18.456	7.134	537	3,193	1.176	0.387	4.187	32.724
Under-five mortality (last 0–9 years)	49.608	13.383	538	3,201	1.285	0.270	22.841	76.375
Stillbirth rate	31.872	14.994	281	1,699	1.289	0.470	1.884	61.860
Early neonatal rate	17.681	9.808	276	1,657	1.050	0.555	0.000	37.297
Perinatal mortality rate	49.116	17.280	281	1,699	1.112	0.352	14.556	83.675
Received ANC from a skilled provider	0.979	0.015	96	577	1.016	0.015	0.949	1.000
4+ ANC visits	0.837	0.037	96	577	0.984	0.045	0.762	0.911
8+ ANC visits	0.154	0.043	96	577	1.165	0.280	0.068	0.241
Took any iron-containing supplements	0.849	0.041	96	577	1.109	0.048	0.767	0.930
Mothers protected against tetanus for last birth	0.863	0.032	96	577	0.899	0.037	0.799	0.926
Delivered in a health facility (live births)	1.000	0.000	101	611	na	0.000	1.000	1.000
Delivered by a skilled provider (live births)	1.000	0.000	101	611	na	0.000	1.000	1.000
Delivered by C-section (live births)	0.307	0.056	101	611	1.146	0.183	0.195	0.419
Women with postnatal check during first 2 days	0.945	0.023	96	577	0.974	0.024	0.899	0.990
Newborns with postnatal check during first 2 days	0.865	0.035	96	577	1.009	0.041	0.794	0.936
Any problem accessing health care	0.496	0.033	602	3,550	1.612	0.066	0.430	0.562
Ever had a vaccination card	0.961	0.030	42	256	1.007	0.031	0.902	1.000
Received BCG vaccination	1.000	0.000	42	256	na	0.000	1.000	1.000
Received DPT-HepB+Hib vaccination (3 doses)	0.958	0.028	42	256	0.920	0.029	0.902	1.000
Received pneumococcal vaccination (3 doses)	0.973	0.027	42	256	1.066	0.027	0.920	1.000
Received measles vaccination	0.942	0.035	42	256	0.992	0.038	0.871	1.000
Fully vaccinated according to national schedule (12–23 months)	0.412	0.073	42	256	0.979	0.178	0.265	0.559
Received measles/measles-rubella 2 vaccination (24–35 months)	0.731	0.058	59	360	1.007	0.079	0.615	0.847
Fully vaccinated according to national schedule (24–35 months)	0.279	0.065	59	360	1.110	0.232	0.149	0.408
Received HPV vaccination (2 doses) women 15–19	0.329	0.077	85	498	1.482	0.233	0.176	0.482
Sought treatment for diarrhoea	0.584	0.072	39	232	0.919	0.124	0.439	0.729
Treated with ORS	0.313	0.073	39	232	0.980	0.232	0.167	0.458
Exclusive breastfeeding	0.710	0.094	26	164	1.039	0.133	0.521	0.899
Minimum dietary diversity (children 6–23 months)	0.466	0.054	66	390	0.872	0.115	0.359	0.574
Child slept under an ITN last night	0.132	0.022	285	1,546	1.001	0.169	0.087	0.177
Received 3+ doses of SP/Fansidar	0.039	0.022	96	577	1.129	0.578	0.000	0.084
Child has fever in last 2 weeks	0.186	0.032	266	1,600	1.316	0.170	0.123	0.249
Child had blood taken from finger/heel	0.226	0.077	48	297	1.309	0.339	0.072	0.379
Child took ACT	0.581	0.303	4	22	1.062	0.521	0.000	1.000
Condom use at last sex (women 15–24)	0.348	0.075	45	245	1.037	0.214	0.199	0.498
Mobile phone ownership	0.933	0.013	602	3,550	1.247	0.014	0.908	0.959

na = not applicable

Table B.52 Sampling errors for adult and maternal mortality rates, Kenya Mini DHS 2025–26

Variable	Value (R)	Standard error (SE)	Number of cases: Unweighted (N)	Number of cases: Weighted (NW)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
			WOMEN					
Adult mortality rates								
15–19	0.360	0.092	42,086	40,364	0.978	0.257	0.175	0.545
20–24	0.811	0.160	47,230	46,422	1.195	0.197	0.492	1.130
25–29	1.822	0.380	45,058	43,847	1.866	0.209	1.062	2.582
30–34	2.236	0.485	42,011	40,661	2.039	0.217	1.266	3.207
35–39	2.181	0.326	35,010	34,167	1.292	0.149	1.530	2.833
40–44	3.042	0.526	24,347	24,174	1.482	0.173	1.991	4.093
45–49	3.233	0.686	15,113	15,446	1.506	0.212	1.860	4.605
15–49 (age-adjusted)	1.726	0.147	250,855	245,081	1.603	0.085	1.433	2.019
Adult mortality probabilities								
$_{35}q_{15}$ 2025–26	66	5	250,855	245,081	2.073	0.083	55	77
$_{35}q_{15}$ 2014	138	7	201,204	199,731	1.835	0.052	124	153
Maternal mortality rates								
15–19	0.034	0.025	42,086	40,364	0.872	0.739	0.000	0.085
20–24	0.101	0.044	47,230	46,422	0.938	0.432	0.014	0.189
25–29	0.313	0.201	45,058	43,847	2.372	0.641	0.000	0.714
30–34	0.819	0.427	42,011	40,661	3.013	0.521	0.000	1.673
35–39	0.278	0.084	35,010	34,167	0.935	0.303	0.109	0.446
40–44	0.322	0.251	24,347	24,174	2.180	0.781	0.000	0.824
45–49	0.151	0.126	15,113	15,446	1.275	0.835	0.000	0.403
15–49 (age-adjusted)	0.282	0.077	250,855	245,081	2.350	0.273	0.128	0.435
Maternal mortality ratio (MMR) 2025–26	263	71	250,855	245,081	2.350	0.270	121	405
Pregnancy-related mortality ratio (PRMR) 2025–26	287	72	250,855	245,081	2.258	0.249	144	431
Pregnancy-related mortality ratio (PRMR) 2014	362	54	201,204	199,731	1.443	0.150	254	471
			MEN					
Adult mortality rates								
15–19	1.151	0.248	43,014	40,606	1.435	0.216	0.654	1.648
20–24	1.448	0.279	48,267	46,211	1.545	0.192	0.891	2.006
25–29	2.497	0.389	46,582	45,240	1.592	0.156	1.720	3.274
30–34	2.652	0.384	42,387	40,943	1.499	0.145	1.885	3.419
35–39	3.727	0.435	34,911	33,790	1.297	0.117	2.857	4.598
40–44	5.010	0.647	25,105	24,386	1.429	0.129	3.717	6.303
45–49	6.319	0.946	16,142	15,993	1.485	0.150	4.426	8.211
15–49 (age-adjusted)	2.828	0.180	256,408	247,169	1.463	0.064	2.467	3.189
Adult mortality probabilities								
$_{35}q_{15}$ 2025–26	108	7	256,408	247,169	2.150	0.060	95	121
$_{35}q_{15}$ 2014	183	9	199,611	196,802	1.648	0.050	165	202

Table B.53 Sampling errors for Early Childhood Development Index 2030, Kenya Mini DHS 2025–26

Background characteristic	Value (R)	Standard error (SE)	Number of cases: Unweighted (N)	Number of cases: Weighted (WN)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
Sex								
Male	0.803	0.011	3,179	2,862	1.527	0.013	0.782	0.825
Female	0.852	0.009	3,084	2,882	1.405	0.011	0.834	0.870
Residence								
Urban	0.867	0.011	1,856	2,482	1.401	0.013	0.845	0.889
Rural	0.798	0.008	4,407	3,262	1.356	0.010	0.782	0.814
Age (in months)								
24–35	0.871	0.010	2,001	1,863	1.359	0.012	0.851	0.891
36–47	0.824	0.011	2,162	2,001	1.332	0.013	0.802	0.846
48–59	0.788	0.012	2,100	1,881	1.361	0.015	0.764	0.813
Mother's education								
No education	0.638	0.029	1,362	582	2.190	0.045	0.581	0.695
Primary	0.788	0.013	2,007	1,922	1.457	0.017	0.761	0.814
Secondary	0.865	0.011	1,970	2,105	1.402	0.013	0.843	0.886
More than secondary	0.924	0.012	924	1,135	1.342	0.013	0.900	0.947
Wealth quintile								
Lowest	0.698	0.014	2,253	1,291	1.458	0.020	0.670	0.726
Second	0.824	0.013	1,254	1,076	1.205	0.016	0.798	0.850
Middle	0.855	0.014	1,133	1,030	1.359	0.017	0.826	0.883
Fourth	0.875	0.015	1,022	1,226	1.478	0.018	0.844	0.905
Highest	0.905	0.018	601	1,120	1.479	0.020	0.869	0.940
Total	**0.828**	**0.007**	**6,263**	**5,744**	**1.419**	**0.008**	**0.814**	**0.841**
County								
Mombasa	0.858	0.050	104	186	1.449	0.058	0.758	0.958
Kwale	0.744	0.043	168	114	1.279	0.058	0.657	0.831
Kilifi	0.877	0.027	135	159	0.938	0.030	0.824	0.930
Tana River	0.692	0.057	164	55	1.580	0.083	0.578	0.807
Lamu	0.817	0.048	123	16	1.363	0.059	0.721	0.913
Taita Taveta	0.766	0.036	87	47	0.784	0.047	0.695	0.837
Garissa	0.622	0.064	223	72	1.946	0.102	0.495	0.750
Wajir	0.746	0.032	224	56	1.095	0.043	0.682	0.810
Mandera	0.629	0.036	312	75	1.311	0.057	0.557	0.700
Marsabit	0.582	0.049	183	62	1.333	0.084	0.485	0.680
Isiolo	0.591	0.058	167	25	1.505	0.097	0.476	0.707
Meru	0.847	0.035	115	143	1.032	0.041	0.777	0.916
Tharaka-Nithi	0.828	0.035	75	39	0.804	0.043	0.757	0.898
Embu	0.737	0.040	67	43	0.740	0.054	0.657	0.817
Kitui	0.589	0.057	110	108	1.198	0.096	0.476	0.702
Machakos	0.779	0.037	102	167	0.906	0.048	0.705	0.854
Makueni	0.848	0.034	89	82	0.891	0.040	0.780	0.916
Nyandarua	0.819	0.059	60	51	1.171	0.072	0.701	0.937
Nyeri	0.965	0.020	49	40	0.744	0.020	0.926	1.004
Kirinyaga	0.895	0.060	65	51	1.556	0.067	0.774	1.016
Murang'a	0.893	0.037	89	78	1.109	0.041	0.819	0.966
Kiambu	0.857	0.029	146	401	1.008	0.034	0.799	0.916
Turkana	0.509	0.042	177	96	1.103	0.082	0.426	0.592
West Pokot	0.552	0.069	193	100	1.916	0.126	0.413	0.690
Samburu	0.694	0.057	172	43	1.609	0.082	0.580	0.808
Trans Nzoia	0.965	0.023	114	85	1.342	0.024	0.918	1.012
Uasin Gishu	0.880	0.036	131	165	1.250	0.041	0.808	0.951
Elgeyo-Marakwet	0.855	0.040	120	52	1.238	0.047	0.775	0.935
Nandi	0.871	0.032	106	79	0.981	0.037	0.807	0.935
Baringo	0.783	0.059	127	64	1.600	0.075	0.665	0.901
Laikipia	0.834	0.053	75	47	1.224	0.064	0.727	0.940
Nakuru	0.879	0.031	140	299	1.134	0.036	0.816	0.942
Narok	0.708	0.057	160	142	1.583	0.081	0.594	0.823
Kajiado	0.884	0.026	122	132	0.892	0.029	0.833	0.936
Kericho	0.932	0.020	103	86	0.802	0.021	0.892	0.972
Bomet	0.851	0.029	161	134	1.041	0.034	0.792	0.910
Kakamega	0.825	0.041	140	179	1.269	0.050	0.743	0.907
Vihiga	0.859	0.034	108	55	1.020	0.040	0.790	0.927
Bungoma	0.865	0.035	146	159	1.245	0.041	0.794	0.936
Busia	0.946	0.032	98	83	1.387	0.034	0.882	1.010
Siaya	0.801	0.042	129	103	1.183	0.052	0.717	0.884

Continued…

Table B.53—*Continued*

Background characteristic	Value (R)	Standard error (SE)	Number of cases: Unweighted (N)	Number of cases: Weighted (WN)	Design effect (DEFT)	Relative error (SE/R)	Confidence limits: Lower (R-2SE)	Confidence limits: Upper (R+2SE)
Kisumu	0.785	0.044	133	136	1.241	0.057	0.696	0.874
Homa Bay	0.840	0.028	178	152	1.016	0.033	0.784	0.896
Migori	0.905	0.020	198	174	0.936	0.022	0.865	0.944
Kisii	0.923	0.018	123	124	0.733	0.019	0.888	0.959
Nyamira	0.801	0.042	99	53	1.040	0.052	0.717	0.885
Nairobi City	0.891	0.021	153	934	0.814	0.023	0.850	0.932

DATA QUALITY TABLES

Appendix C

Table C.1 Household age distribution

Single-year age distribution of the de facto household population by sex (weighted), Kenya Mini DHS 2025–26

	Male		Female			Male		Female	
Age	Number	Percent	Number	Percent	Age	Number	Percent	Number	Percent
0	1,017	2.2	957	2.1	42	453	1.0	542	1.2
1	948	2.1	1,039	2.3	43	405	0.9	428	1.0
2	1,043	2.3	1,098	2.5	44	250	0.6	221	0.5
3	1,213	2.7	1,130	2.5	45	486	1.1	533	1.2
4	1,183	2.6	1,208	2.7	46	339	0.7	321	0.7
5	1,135	2.5	1,208	2.7	47	386	0.9	357	0.8
6	1,070	2.4	1,205	2.7	48	285	0.6	339	0.8
7	1,109	2.4	1,236	2.8	49	277	0.6	295	0.7
8	1,169	2.6	1,178	2.6	50	433	1.0	474	1.1
9	1,067	2.4	1,153	2.6	51	440	1.0	234	0.5
10	1,145	2.5	1,218	2.7	52	476	1.0	345	0.8
11	1,035	2.3	1,040	2.3	53	379	0.8	275	0.6
12	1,210	2.7	1,089	2.4	54	260	0.6	203	0.5
13	1,177	2.6	1,161	2.6	55	307	0.7	302	0.7
14	1,010	2.2	1,059	2.4	56	274	0.6	263	0.6
15	875	1.9	1,068	2.4	57	214	0.5	224	0.5
16	842	1.9	972	2.2	58	182	0.4	183	0.4
17	941	2.1	1,035	2.3	59	149	0.3	142	0.3
18	853	1.9	1,029	2.3	60	260	0.6	304	0.7
19	704	1.6	805	1.8	61	154	0.3	120	0.3
20	791	1.7	816	1.8	62	198	0.4	210	0.5
21	687	1.5	735	1.6	63	238	0.5	169	0.4
22	766	1.7	757	1.7	64	200	0.4	140	0.3
23	807	1.8	719	1.6	65	260	0.6	217	0.5
24	739	1.6	624	1.4	66	145	0.3	104	0.2
25	1,052	2.3	923	2.1	67	167	0.4	122	0.3
26	786	1.7	640	1.4	68	153	0.3	148	0.3
27	724	1.6	744	1.7	69	114	0.3	122	0.3
28	752	1.7	656	1.5	70	247	0.5	209	0.5
29	651	1.4	654	1.5	71	103	0.2	84	0.2
30	884	1.9	810	1.8	72	133	0.3	112	0.3
31	554	1.2	537	1.2	73	121	0.3	88	0.2
32	737	1.6	743	1.7	74	61	0.1	75	0.2
33	575	1.3	493	1.1	75	125	0.3	113	0.3
34	464	1.0	474	1.1	76	75	0.2	70	0.2
35	722	1.6	724	1.6	77	56	0.1	47	0.1
36	604	1.3	464	1.0	78	75	0.2	54	0.1
37	513	1.1	443	1.0	79	49	0.1	40	0.1
38	673	1.5	633	1.4	80+	545	1.2	335	0.7
39	597	1.3	477	1.1	Don't know	31	0.1	44	0.1
40	698	1.5	793	1.8					
41	353	0.8	379	0.8	Total	45,380	100	44,730	100.5

Note: The de facto population includes all residents and nonresidents who stayed in the household the night before the interview.

Figure C1 Population pyramid

Percent distribution of the household population by single-year




Table C.2 Age distribution of eligible and interviewed women

De facto household population of women age 10–54, number and percent distribution of interviewed women age 15–49, and percentage of eligible women who were interviewed (weighted), by 5-year age groups, Kenya Mini DHS 2025–26

	Household population of women age 10–54	Interviewed women age 15–49		Percentage of eligible women interviewed
Age group	Number	Number	Percentage	
10–14	5,577	na	na	na
15–19	4,215	3,871	18.6	91.8
20–24	3,791	3,558	17.1	93.9
25–29	3,964	3,701	17.8	93.4
30–34	3,215	3,052	14.7	94.9
35–39	3,110	2,928	14.1	94.1
40–44	2,159	1,993	9.6	92.4
45–49	1,773	1,659	8.0	93.6
50–54	1,988	na	na	na
15–49	22,227	20,761	100.0	93.4
Ratio				
10–14 to 15–19	132.3	na	na	na
50–54 to 45–49	112.1	na	na	na

Notes: The de facto population includes all residents and nonresidents who stayed in the household the night before the interview. Weights for both the household population of women and interviewed women are household weights. Age is based on the Household Questionnaire.
na = not applicable

Table C.3 Age displacement at ages 14/15

Number of women age 12–18 listed in the household schedule by single-year age and age ratio 15/14, according to region (weighted), Kenya Mini DHS 2025–26

	Age							Total age 12–18	Age ratio (age 15/ age 14)
	12	13	14	15	16	17	18		
County									
Mombasa	57	18	45	30	21	29	31	231	67.0
Kwale	29	30	20	30	15	26	23	172	151.9
Kilifi	48	34	42	44	27	35	19	249	103.4
Tana River	11	11	10	10	11	10	11	73	97.8
Lamu	3	3	3	3	3	4	3	22	117.3
Taita Taveta	10	8	5	8	6	7	10	53	166.0
Garissa	19	16	13	9	13	11	8	89	68.2
Wajir	15	15	13	8	11	11	12	84	61.9
Mandera	13	11	14	10	6	4	7	67	72.1
Marsabit	13	15	10	15	9	13	10	84	146.7
Isiolo	6	6	4	4	5	4	3	31	104.7
Meru	29	20	34	24	30	36	20	193	71.0
Tharaka-Nithi	7	10	11	9	9	10	6	63	80.2
Embu	15	11	11	7	11	10	10	75	61.0
Kitui	24	46	27	22	20	24	16	179	80.6
Machakos	23	23	30	25	20	21	28	170	83.7
Makueni	13	19	18	24	22	22	18	136	132.1
Nyandarua	14	14	18	8	13	9	9	86	43.9
Nyeri	9	11	7	7	6	10	11	62	96.3
Kirinyaga	14	10	7	6	13	13	8	70	80.6
Murang'a	19	19	13	21	16	23	13	124	157.6
Kiambu	55	56	43	48	39	41	63	346	112.7
Turkana	27	28	17	10	8	9	10	109	54.6
West Pokot	24	23	23	11	13	9	11	114	48.9
Samburu	10	9	10	7	8	6	4	55	72.4
Trans Nzoia	22	26	18	23	21	25	11	147	130.2
Uasin Gishu	44	33	26	29	34	31	27	224	107.9
Elgeyo-Marakwet	13	18	13	8	14	8	10	85	62.7
Nandi	19	18	19	21	20	27	22	145	108.0
Baringo	16	30	12	11	12	11	9	101	94.8
Laikipia	9	15	9	7	12	12	6	70	72.8
Nakuru	83	54	52	57	58	40	37	381	110.6
Narok	20	37	44	27	23	17	21	188	62.1
Kajiado	29	22	23	16	15	15	23	142	67.8
Kericho	20	23	32	19	24	19	20	156	58.2
Bomet	32	34	17	35	40	28	28	214	206.6
Kakamega	49	51	46	38	47	61	24	316	82.5
Vihiga	16	19	19	21	17	21	12	125	114.6
Bungoma	40	55	33	40	33	46	40	288	118.7
Busia	27	39	30	20	19	25	22	181	65.1
Siaya	35	41	25	20	17	21	21	179	77.5
Kisumu	37	51	18	27	28	26	36	224	149.8
Homa Bay	35	39	32	34	26	33	25	223	106.1
Migori	43	39	28	32	20	38	18	218	115.4
Kisii	41	41	29	25	29	37	23	225	85.9
Nyamira	22	15	15	15	17	18	14	116	98.2
Nairobi City	134	89	120	89	101	160	124	817	74.3
Total	1,293	1,252	1,109	1,011	980	1,115	938	7,699	91.2

Table C.4 Age displacement at ages 49/50

Number of women age 47–53 listed in the household schedule by single-year age and age ratio 50/49, according to region (weighted), Kenya Mini DHS 2025–26

Region	Age							Total age 47–53	Age ratio (age 50/ age 49)
	47	48	49	50	51	52	53		
County									
Mombasa	9	4	5	19	8	15	10	71	417.9
Kwale	9	6	9	8	6	5	4	48	94.8
Kilifi	15	9	7	5	13	17	8	74	70.2
Tana River	3	4	1	1	3	2	3	17	211.2
Lamu	1	1	0	1	2	1	2	8	262.7
Taita Taveta	2	3	2	5	5	4	4	25	280.0
Garissa	4	1	1	5	1	2	4	18	323.2
Wajir	2	2	0	3	2	3	1	15	773.4
Mandera	2	1	1	4	3	1	3	14	491.2
Marsabit	2	1	3	4	4	3	2	17	132.3
Isiolo	1	1	0	1	1	2	1	8	186.4
Meru	13	12	7	10	12	24	10	88	141.4
Tharaka-Nithi	5	2	2	6	6	4	4	28	290.0
Embu	9	5	4	8	9	4	6	45	231.3
Kitui	8	9	5	13	22	9	19	84	297.8
Machakos	12	9	13	19	15	28	11	108	151.4
Makueni	10	7	7	7	14	11	16	72	89.8
Nyandarua	6	8	4	2	6	3	4	34	50.4
Nyeri	6	5	4	13	13	15	10	65	341.1
Kirinyaga	10	1	5	12	11	14	13	65	232.8
Murang'a	10	8	15	11	4	14	7	69	69.3
Kiambu	23	14	13	18	18	29	17	132	136.9
Turkana	4	3	1	8	8	4	2	30	750.9
West Pokot	4	3	3	5	5	3	4	28	155.8
Samburu	1	3	1	1	2	4	2	14	92.8
Trans Nzoia	14	12	7	6	4	7	6	56	84.6
Uasin Gishu	13	10	5	11	11	9	6	64	248.4
Elgeyo-Marakwet	3	3	2	6	2	2	2	20	316.0
Nandi	6	4	6	9	11	9	4	50	149.9
Baringo	3	2	2	2	5	3	7	24	138.2
Laikipia	9	5	2	8	9	5	5	42	418.8
Nakuru	8	10	7	18	19	31	25	119	266.9
Narok	8	6	5	30	9	9	7	74	631.6
Kajiado	3	9	1	8	13	11	6	50	1,064.1
Kericho	8	5	7	6	12	11	10	58	88.8
Bomet	7	7	9	9	10	4	9	56	93.2
Kakamega	19	12	8	15	17	16	14	103	177.0
Vihiga	7	4	5	4	3	6	4	33	72.8
Bungoma	12	18	15	11	12	17	12	98	73.2
Busia	5	5	8	8	6	7	10	50	105.4
Siaya	5	6	4	10	9	12	7	53	261.9
Kisumu	8	9	9	5	7	10	2	49	56.8
Homa Bay	4	4	8	6	10	9	11	52	72.9
Migori	4	6	8	7	6	5	8	43	94.2
Kisii	9	12	5	8	12	19	20	86	153.7
Nyamira	8	5	4	7	6	7	7	44	203.9
Nairobi City	60	14	45	62	71	53	37	343	139.1
Total	392	289	284	445	459	484	388	2,741	156.9

Table C.5 Pregnancy outcomes by years preceding the survey

Number of pregnancy outcomes, percentage with year and month of birth given or end of pregnancy given, sex ratio at birth of live births, and ratio by years preceding the survey, according to living children, dead children, stillbirths, miscarriages/abortions, and total pregnancy outcomes (weighted), Kenya Mini DHS 2025–26

	Number of pregnancy outcomes					Percentage with year and month of birth given or end of pregnancy given					Sex ratio at birth of live births[1]			Ratio by years preceding survey[2]				
Years preceding survey	Living children	Dead children	Stillbirths	Miscar-riages/ abortions	Total	Living children	Dead children	Stillbirths	Miscar-riages/ abortions	Total	Living children	Dead children	Total	Living children	Dead children	Stillbirths	Miscar-riages/ abortions	Total
0	2,041	58	41	243	2,383	100.0	100.0	100.0	100.0	100.0	93.6	160.9	95.0	na	na	na	na	na
1	1,934	81	28	217	2,261	100.0	100.0	100.0	100.0	100.0	108.3	152.5	109.7	95.6	117.2	98.5	93.9	96.1
2	2,007	80	16	220	2,322	100.0	100.0	100.0	100.0	100.0	103.0	169.8	104.9	98.6	101.2	55.0	113.8	99.4
3	2,136	77	30	169	2,412	100.0	100.0	100.0	100.0	100.0	95.7	150.9	97.3	105.4	112.6	127.0	92.7	104.8
4	2,048	57	31	145	2,281	100.0	99.0	100.0	100.0	100.0	100.2	191.7	101.9	97.6	83.0	110.2	80.8	96.0
5	2,063	60	26	190	2,339	100.0	100.0	100.0	100.0	100.0	107.3	92.9	106.9	100.2	79.3	93.2	150.0	102.2
6	2,070	94	26	108	2,297	99.4	96.8	91.1	96.7	99.1	124.4	129.5	124.6	103.6	158.9	78.9	65.3	101.9
7	1,933	59	39	141	2,171	99.0	96.5	98.1	80.7	97.7	101.9	60.2	100.4	94.7	67.6	175.8	139.7	96.5
8	2,010	79	18	94	2,201	99.1	88.4	88.0	96.1	98.5	109.9	84.7	108.8	106.9	103.3	50.3	65.7	103.1
9	1,826	95	34	144	2,099	99.2	95.9	72.1	93.0	98.2	111.5	84.2	109.9	96.0	112.1	182.3	147.9	99.8
0–4	10,166	353	146	994	11,659	100.0	99.8	100.0	100.0	100.0	99.9	163.0	101.5	na	na	na	na	na
5–9	9,901	386	144	676	11,107	99.3	95.3	89.7	93.4	98.7	110.9	90.5	110.0	na	na	na	na	na
10–14	8,485	379	112	411	9,387	98.6	91.9	87.5	69.1	96.9	99.9	113.2	100.4	na	na	na	na	na
15–10	7,246	371	108	258	7,983	98.0	92.0	87.1	80.3	97.0	105.0	119.1	105.7	na	na	na	na	na
20+	6,938	610	84	226	7,858	96.6	89.4	77.5	74.2	95.2	95.5	130.8	97.9	na	na	na	na	na
All	42,737	2,099	594	2,565	47,995	98.7	93.1	89.6	89.1	97.8	102.4	121.4	103.3	na	na	na	na	na

[1] (Bm/Bf)x100, where Bm and Bf are the numbers of male and female births, respectively
[2] [2Px/(Px-1+Px+1)]x100, where Px is the number of pregnancy outcomes in year x preceding the survey
na = not applicable

Table C.6 Completeness of reporting

Percentage of observations missing information for selected demographic and health questions (weighted), Kenya Mini DHS 2025–26

Subject	Percentage with information missing	Number of cases
Date of live birth or stillbirth (last 15 years)		
Missing day only	1.12	30,072
Missing month, but year reported	0.82	30,072
Missing year	0.15	30,072
Date of live birth or stillbirth (last 5 years)		
Missing day only	0.00	10,665
Missing year	0.01	10,665
Date of birth of women		
Missing month, but year reported	1.72	21,231
Missing year	0.36	21,231
Diarrhoea in last 2 weeks	0.69	10,166

Table C.7 Observation of mosquito nets

Percentage of all mosquito nets observed by the interviewers, according to background characteristics (weighted), Kenya Mini DHS 2025–26

Background characteristic	Percentage of mosquito nets observed by interviewers	Number of mosquito nets
Residence		
Urban	83.3	9,660
Rural	89.9	17,035
County		
Mombasa	70.8	1,040
Kwale	98.1	602
Kilifi	100.0	1,181
Tana River	98.7	280
Lamu	92.1	91
Taita Taveta	88.8	420
Garissa	99.3	320
Wajir	97.0	218
Mandera	97.4	219
Marsabit	95.8	117
Isiolo	78.0	80
Meru	78.7	415
Tharaka-Nithi	86.6	144
Embu	80.0	198
Kitui	99.6	382
Machakos	69.0	602
Makueni	96.1	432
Nyandarua	79.7	75
Nyeri	70.3	87
Kirinyaga	68.4	476
Murang'a	66.9	265
Kiambu	77.6	1,035
Turkana	99.8	556
West Pokot	67.2	385
Samburu	58.0	35
Trans Nzoia	91.7	796
Uasin Gishu	97.0	756
Elgeyo-Marakwet	92.2	112
Nandi	91.2	466
Baringo	89.7	267
Laikipia	97.6	90
Nakuru	87.2	607
Narok	99.7	671
Kajiado	59.3	444
Kericho	89.6	684
Bomet	63.7	633
Kakamega	95.2	1,440
Vihiga	93.9	538
Bungoma	99.9	1,351
Busia	80.4	819
Siaya	94.9	602
Kisumu	91.4	1,098
Homa Bay	99.5	810
Migori	79.5	1,071
Kisii	84.6	693
Nyamira	92.7	549
Nairobi City	83.5	2,541
Wealth quintile		
Lowest	92.9	4,300
Second	91.7	5,545
Middle	88.1	5,805
Fourth	84.7	5,699
Highest	81.3	5,346
Total	87.6	26,695

Table C.8 Observation of handwashing facility

Percent distribution of handwashing facilities in all households by whether or not they were observed by the interviewers, according to background characteristics (weighted), Kenya Mini DHS 2025–26

Background characteristic	Handwashing facility observed		Handwashing facility not observed			Total	Number of households
	Fixed place	Mobile	Not in dwelling, yard, or plot	No permission to see	Other reason		
Residence							
Urban	45.0	43.6	8.3	1.3	1.8	100.0	11,975
Rural	18.5	67.4	12.3	0.3	1.5	100.0	14,405
County							
Mombasa	26.3	32.8	39.0	0.6	1.3	100.0	866
Kwale	9.4	88.3	2.0	0.3	0.0	100.0	318
Kilifi	13.9	86.1	0.0	0.0	0.0	100.0	699
Tana River	5.9	94.1	0.0	0.0	0.0	100.0	143
Lamu	16.7	47.6	34.0	1.2	0.4	100.0	62
Taita Taveta	35.8	42.3	21.4	0.6	0.0	100.0	206
Garissa	6.9	41.9	51.2	0.0	0.0	100.0	145
Wajir	7.5	64.6	27.9	0.0	0.0	100.0	126
Mandera	8.3	89.4	0.7	0.0	1.6	100.0	137
Marsabit	6.7	68.4	24.2	0.0	0.7	100.0	138
Isiolo	27.3	30.5	42.1	0.0	0.0	100.0	87
Meru	31.6	62.7	0.3	1.1	4.2	100.0	840
Tharaka-Nithi	31.0	68.5	0.5	0.0	0.0	100.0	209
Embu	46.0	49.9	2.7	1.0	0.4	100.0	385
Kitui	16.0	84.0	0.0	0.0	0.0	100.0	526
Machakos	20.5	44.8	32.2	1.8	0.7	100.0	1,021
Makueni	23.6	72.3	4.2	0.0	0.0	100.0	506
Nyandarua	34.7	65.0	0.1	0.0	0.2	100.0	341
Nyeri	49.2	37.3	11.9	0.1	1.5	100.0	441
Kirinyaga	49.6	39.5	2.8	0.6	7.5	100.0	482
Murang'a	46.9	50.4	0.7	0.8	1.1	100.0	571
Kiambu	73.6	17.5	0.8	1.1	7.0	100.0	2,295
Turkana	10.2	86.6	3.1	0.2	0.0	100.0	232
West Pokot	10.0	64.8	23.5	1.7	0.0	100.0	251
Samburu	10.5	56.2	31.6	0.0	1.6	100.0	109
Trans Nzoia	14.4	80.7	4.6	0.2	0.0	100.0	393
Uasin Gishu	36.2	62.3	1.4	0.2	0.0	100.0	794
Elgeyo-Marakwet	29.3	68.7	1.9	0.0	0.0	100.0	209
Nandi	28.4	60.1	4.1	0.3	7.0	100.0	414
Baringo	10.3	64.5	24.4	0.5	0.3	100.0	242
Laikipia	41.2	36.6	22.1	0.0	0.0	100.0	293
Nakuru	27.1	65.0	6.9	0.7	0.4	100.0	1,442
Narok	5.2	80.6	14.2	0.0	0.0	100.0	576
Kajiado	34.6	64.8	0.5	0.0	0.0	100.0	653
Kericho	26.1	71.4	1.5	0.1	0.9	100.0	464
Bomet	10.6	38.9	50.2	0.0	0.4	100.0	437
Kakamega	16.4	74.7	8.9	0.0	0.0	100.0	850
Vihiga	6.6	33.9	56.2	2.9	0.4	100.0	267
Bungoma	8.5	86.2	4.9	0.1	0.4	100.0	746
Busia	11.8	76.5	0.0	0.4	11.4	100.0	429
Siaya	7.4	45.7	45.0	0.0	1.9	100.0	482
Kisumu	18.0	77.8	0.0	0.1	4.1	100.0	615
Homa Bay	11.6	88.4	0.0	0.0	0.0	100.0	568
Migori	15.8	39.1	44.5	0.5	0.0	100.0	499
Kisii	6.0	60.8	31.8	1.2	0.2	100.0	597
Nyamira	5.8	94.2	0.0	0.0	0.0	100.0	284
Nairobi City	48.3	44.9	3.7	2.2	0.9	100.0	3,992
Wealth quintile							
Lowest	2.6	79.5	15.7	0.5	1.7	100.0	4,299
Second	8.8	74.3	14.9	0.2	1.8	100.0	4,719
Middle	21.1	64.5	12.0	0.7	1.6	100.0	5,346
Fourth	32.8	55.5	8.9	0.6	2.2	100.0	6,192
Highest	74.8	19.2	3.4	1.6	1.0	100.0	5,825
Total	30.5	56.6	10.5	0.8	1.6	100.0	26,380

Table C.9 Vaccination cards photographed

Percentage of children under age 3 reported to have a vaccination card, percentage whose vaccination card was seen by the interviewer, and percentages whose vaccination card was photographed and was not photographed by reason; and among children with a vaccination card seen, percentage whose card was photographed, according to background characteristics (weighted), Kenya Mini DHS 2025–26

Background characteristic	Percentage of children reported to have a vaccination card	Percentage of children whose vaccination card was seen by interviewer	Percentage of children whose vaccination card was photographed	Percentage of children whose vaccination card was not photographed as permission was not received	Percentage of children whose vaccination card was not photographed for other reasons	Number of children	Among children with a vaccination card seen: Percentage whose vaccination card was photographed	Among children with a vaccination card seen: Number of children
Age in months								
0–11	90.4	82.9	81.3	1.4	0.2	2,041	98.0	1,693
12–23	80.4	69.1	68.1	0.8	0.1	1,934	98.6	1,337
24–35	72.6	55.7	54.1	1.5	0.0	2,007	97.2	1,117
Residence								
Urban	77.5	62.4	60.7	1.6	0.1	2,592	97.3	1,617
Rural	84.0	74.6	73.4	1.0	0.2	3,390	98.4	2,529
County								
Mombasa	73.9	71.1	70.3	0.0	0.8	176	98.8	125
Kwale	77.1	77.1	77.1	0.0	0.0	102	100.0	78
Kilifi	82.1	74.4	74.4	0.0	0.0	163	100.0	121
Tana River	81.3	80.9	78.1	1.7	1.0	58	96.6	47
Lamu	85.8	80.0	79.0	0.0	1.1	16	98.7	13
Taita Taveta	89.8	79.0	77.1	0.0	1.9	42	97.6	33
Garissa	53.5	53.2	52.0	0.0	1.1	64	97.8	34
Wajir	45.7	35.7	35.7	0.0	0.0	56	100.0	20
Mandera	48.5	29.1	27.4	0.6	1.1	67	94.2	19
Marsabit	91.9	68.0	68.0	0.0	0.0	55	100.0	37
Isiolo	84.2	78.0	77.2	0.0	0.8	23	98.9	18
Meru	95.1	81.2	81.2	0.0	0.0	119	100.0	97
Tharaka-Nithi	84.8	78.7	78.7	0.0	0.0	36	100.0	28
Embu	80.4	71.7	71.7	0.0	0.0	44	100.0	31
Kitui	80.7	73.5	73.5	0.0	0.0	98	100.0	72
Machakos	73.8	58.5	53.9	4.6	0.0	212	92.1	124
Makueni	95.1	78.5	78.5	0.0	0.0	95	100.0	75
Nyandarua	90.8	71.1	71.1	0.0	0.0	67	100.0	48
Nyeri	89.3	77.6	77.6	0.0	0.0	54	100.0	42
Kirinyaga	79.7	71.6	71.6	0.0	0.0	54	100.0	38
Murang'a	84.6	78.1	77.0	1.2	0.0	93	98.5	73
Kiambu	73.0	60.8	60.1	0.8	0.0	396	98.7	241
Turkana	86.1	80.6	80.1	0.5	0.0	92	99.4	74
West Pokot	71.9	52.5	23.1	29.1	0.3	107	44.0	56
Samburu	78.2	72.1	72.1	0.0	0.0	46	100.0	33
Trans Nzoia	75.4	72.2	70.5	1.6	0.0	95	97.7	68
Uasin Gishu	84.7	57.9	57.9	0.0	0.0	179	100.0	104
Elgeyo-Marakwet	87.0	85.1	83.1	0.0	2.0	47	97.6	40
Nandi	82.9	80.0	79.5	0.4	0.0	104	99.5	83
Baringo	84.1	78.5	78.5	0.0	0.0	76	100.0	59
Laikipia	82.0	68.7	68.7	0.0	0.0	62	100.0	42
Nakuru	86.8	71.9	71.2	0.7	0.0	322	99.0	232
Narok	81.0	80.3	80.3	0.0	0.0	127	100.0	102
Kajiado	80.0	70.0	68.6	1.0	0.5	169	97.9	118
Kericho	90.8	77.3	76.3	0.0	1.0	96	98.7	74
Bomet	85.6	81.7	81.7	0.0	0.0	138	100.0	112
Kakamega	87.8	75.8	75.8	0.0	0.0	190	100.0	144
Vihiga	92.2	79.1	79.1	0.0	0.0	59	100.0	46
Bungoma	84.0	77.8	77.8	0.0	0.0	183	100.0	143
Busia	93.5	70.9	70.9	0.0	0.0	116	100.0	82
Siaya	71.7	56.9	56.9	0.0	0.0	93	100.0	53
Kisumu	82.8	67.8	67.8	0.0	0.0	165	100.0	112
Homa Bay	81.0	73.7	73.7	0.0	0.0	132	100.0	97
Migori	88.3	78.9	78.9	0.0	0.0	188	100.0	148
Kisii	89.7	73.1	73.1	0.0	0.0	115	100.0	84
Nyamira	86.1	69.8	68.8	0.0	1.0	60	98.6	42
Nairobi City	78.9	62.2	59.8	2.4	0.0	936	96.2	582
Wealth quintile								
Lowest	79.6	72.9	71.2	1.5	0.2	1,244	97.7	907
Second	85.3	75.1	74.1	0.7	0.3	1,113	98.7	836
Middle	84.3	73.4	72.8	0.4	0.1	1,081	99.3	793
Fourth	80.0	65.7	63.7	1.8	0.1	1,340	97.1	880
Highest	77.7	60.7	59.0	1.6	0.0	1,204	97.3	730
Total	81.2	69.3	67.9	1.3	0.1	5,982	98.0	4,146

Note: Vaccination cards include cards, booklets, or other home-based records.

Table C.10 Completeness of information on siblings

Completeness of data on survival status of sisters and brothers reported by interviewed women, age of living siblings, and age at death (AD) and years since death (YSD) of dead siblings (unweighted), Kenya Mini DHS 2025–26

	Sisters		Brothers		All siblings	
	Number	Percent	Number	Percent	Number	Percent
All siblings	50,916	100.0	52,813	100.0	103,729	100.0
Living	47,747	93.8	48,922	92.6	96,669	93.2
Dead	3,155	6.2	3,882	7.4	7,037	6.8
Survival status unknown	14	0.0	9	0.0	23	0.0
Living siblings	47,747	100.0	48,922	100.0	96,669	100.0
Age reported	47,747	100.0	48,922	100.0	96,669	100.0
Dead siblings	3,155	100.0	3,882	100.0	7,037	100.0
AD and YSD reported	3,155	100.0	3,882	100.0	7,037	100.0

Table C.11 Sibship size and sex ratio of siblings

Mean sibship size and sex ratio of siblings at birth, Kenya Mini DHS 2025–26

Age of respondents	Mean sibship size[1]	Sex ratio of siblings at birth[2]
15–19	5.0	100.5
20–24	5.3	98.0
25–29	5.5	105.5
30–34	5.7	104.6
35–39	6.1	101.7
40–44	6.2	102.2
45–49	6.6	110.2
Total	5.7	102.8

[1] Includes the respondent
[2] Excludes the respondent

Table C.12 Pregnancy-related mortality trends

Direct estimates of pregnancy-related mortality rates for the 7 years preceding each survey, by 5-year age groups, Kenya Mini DHS 2025–26

	Pregnancy-related mortality rates[1,2]				
Age	2019–2026	2007–2014	2002–2009	1996–2003	1991–1998
15–19	0.03	0.11	0.38	0.45	0.68
20–24	0.11	0.46	0.67	1.01	1.58
25–29	0.40	0.73	0.92	1.06	1.53
30–34	0.86	0.65	0.86	1.23	1.41
35–39	0.30	0.87	1.68	1.17	1.23
40–44	0.32	0.50	0.68	0.40	1.41
45–49	0.15	0.22	1.09	0.30	0.16
Total 15–49 [a]	0.31	0.51	0.82	0.83	1.19
Total fertility rate (TFR)	3.3	3.9	4.6	4.9	4.7
General fertility rate (GFR)[3]	107	141	161	171	166
Pregnancy-related mortality ratio (PRMR)[4]	287	362	520	506	743
Confidence interval	(144, 431)	(254, 471)	(343, 696)	(398, 614)	(591, 895)
Lifetime risk of pregnancy-related death[5]	0.010	0.015	0.025	0.025	0.036

[1] Pregnancy-related mortality is defined as the death of a woman while pregnant or within 2 months of termination of pregnancy, from any cause including accidents or violence.
[2] Expressed per 1,000 woman-years of exposure
[3] Age-adjusted rate expressed per 1,000 women age 15–49
[4] Expressed per 100,000 live births; calculated as the age-adjusted pregnancy-related mortality rate times 100 divided by the age-adjusted general fertility rate
[5] Calculated as $1-(1-PRMR)^{TFR}$ where TFR represents the total fertility rate for the 7 years preceding the survey
[a] Age-adjusted rate

PERSONS INVOLVED IN THE 2025–26 KENYA MINI DEMOGRAPHIC AND HEALTH SURVEY

Appendix D

TECHNICAL AND ADMINISTRATIVE STAFF

National Coordinator

Dr Macdonald Obudho

National Coordinator

Rosemary Chepkoech Bowen

Regional Coordinators

Dr Jane Weru
Robert Nderitu
Benjamin Muchiri
Benjamin Avusevwa

Project Management

Francis Kundu
Scholastica Kingi

Senior Technical Staff

Name	Role
James Ng'ang'a	Technical Coordinator/Sampler/Trainer/Author
Elias Nyaga	Technical Coordinator/Trainer/Author
Vivianne Nyarunda	Technical Coordinator/Trainer/Author
Andrew Imbwaga	Technical Coordinator/Trainer/Author
Zachary Ochola	Trainer/Sampler/Author
Jim Kirimi	Trainer/ICT/Data Team
Kevin Nyawade	Trainer/ICT/Data Team/Author
Paul Waweru	Trainer/ICT/Data Team
Peter Kamau	Trainer/ICT/Data Team/Author
Maurice Kamau	Trainer/ICT/Data Team
Mirriam Mwaura	Trainer/Data Editor/Author
Kevin Dede	Trainer/Data Editor/Author
Jackline Chepkorir	Trainer/Data Editor
Caneble Oganga	Trainer/Data Editor/Author
Renice Bunde	Trainer/Data Editor/Author
Winnie Kemunto	Trainer/Data Editor/Author
Hiram Mbatia	Trainer/Author
Paul Kuria	Trainer/Author
Peter Wanjohi	Trainer/Author
Isaac Ndegwa	Trainer/Author
Christine Mbuli	Trainer/Author
Dr Martin Mwangi	Trainer/Author
Pius Kibet	Trainer/Data Editor/Author
Bernard Kiprotich	Trainer/Author
Dr Hellen Kiarie	Trainer/Author
Bueter Obura	Trainer/Author
Prof. George Otieno Orwa	Trainer/Author
Scolastica Wabwire	Trainer/Author

Prof. Alfred Agwanda	Trainer/Author
Dr Samwel Wakibi	Trainer/Author
Samwel Murage	Trainer/Author
Oliver Watima	Trainer/Author
Richard Kimenye	Trainer/Author
Benson Karugu	Trainer/Author
Benard Wambu	Trainer/Author
Stanley Wambua	Trainer/Author
John Anampiu	Trainer/Author
Loise Mathenge	Trainer/Author
Feisal Hassan	Author
Alinoor Hapicha	Author
Prof Lawrence Ikamari	Author
Paul Samoei	Author
Dekar Muktar	Sampler/Author
Prisca Mwangi	Trainer/Author/Logistics Coordinator
Franklin Mwamba	Logistics Coordinator/Author
Geoffrey Kariuki	Trainer/Logistics Coordinator
Ronald Kirui	Trainer/Logistics Coordinator
Samwel Maosa	Logistics Coordinator
Baldwin Anyiga	Trainer
John Bore	Trainer
Medina Ibrahim	Trainer
Rose Awino	Trainer
Katunge Kiilu	Trainer
Dr Camlus Odhus	Author
Lucy Maina	Author
Vivian Nyakangi	Author
Prof Amina Abubakar	Author
Dr Makeba Shiroya	Author

Cluster Mobilisation Officers

Daniel Ochieng Okoo
Simeon Owino Onyisi
Vincent Kipkemboi Kemei
Vincent Obiri
Raphael Namatsi
Sophia Nafula Nyongesa
Janet Nyanchama Nchobera
Jane Adungo
David Kipkorir Kigen
Shadrack Kimeli Kitur
Monicah Mugure Karanja
Bernard Muhoro Karenye
Hassan Sanbur Adan
Zeinab Ahmed Abdullahi
Jared Orwa
Julius Ochieng Ochanda
Shinda Abdullahi Ware
Hamdi Bishar
Samwel Leteipa Ngoibuni
Douglas Nyakango Sagwe
Morice Okech Motegi
Calectus Billy Onyango
Rhoda Nyakundi
Joseph Kimani
David Mutinda Mueke
David Kyengo Amos
Reuben Saro
Harriet Wawira Githinji
Paul Juma Wekesa
Mohamed Ahmed Hassan
Wilson Galana Malibe
Redempta Katungwa Muyuma
Samson Musila Kalungu
Veronica Kalii Ndeto
Christopher Muasya Kyangu
Abdikadir Abdirahman Adan
Ahmed Abdullahi Abdirahman
Guyo Sora Galgallo
Philemon Kiprono Kipkirwok
Jonah Kipkoech Chesir
Simion Chelimo Bowen
Fred Leyian Kamwaro
Martin Ogoti Mogambi
Joash Nyasani Ondieki
John Muriithi Murekio
Phoebe Kageni Kariuki
Joel Ndichu Ng'Ang'A
Esther Wanjiru Wamithi
James Maina Kangara
Jacob Kaaka Tumpeine
Irine Odegi
Charles Omondi Obudho
Abdiwahab Mohamed
Kennedy Mugeni
Mohamed Adan Ibrahim
Mohamed Ahmed Kerow
Robert Kioko

Mourice Peter Sila
Edwin Muthune Agola
Daniel Asiko Olukaka
Judy Chepkirui Rono
Jackson Wafula Wamalwa
Johnstone Matheka Kitema
Winnie Wambui Kiraka
Joseph Chengo Thoya
Sawa Mwatela Kamanza
Catherine Wairimu Wanyoro
Mary Njoki Maina
Stephen Kapterema
Stephen Kapterema
Peter Ltamalan Ngolei
Gabriel Ntabo
Hellen Kathao Kobia
Moses Nyanjong Osano
Maurice Opiyo Otieno
Stanslaus Nzuki
Washington Saitoti Madadi Dula
Maxwell Kirima Macharia
Benson Chege
Philomena Munyoro
Aisha Razik Said
Korir Evans Kipyegon
Annah Sereya Purkei
Boaz Mweu
Emmanuel Kirwa Kurgat
David Williams Ochieng
Dickson Osolo Warega
Eris Dalmas Ewet
Dominic Kiptanui Kemei
Dorcas Aono Angolo
Frankline Wesonga Apwokha
Annalice Lumumba
Abdinoor Salat Mohamed
Hassan Mohamed Abdullahi

Field Supervisors

Abdifatah Mohamed Kosar
Amos Otieno Nyadera
Arani Job Mumbo
Azaneth Jebet
Beulah Jepngetich
Christine A Onyango
Cindy Wanjiru Mahinda
Elvis Kibet Changwony
Enoch Obuolo
Eunice Kwamboka Mokua
Everlyne Wangui Mwangi
Felix Maroa Chacha
Frank Jara
Gao Ken Lwaki
George Odhiambo
Hesbon Nyamoko Nyakweba
Hilbert Omamo
Hope Lily Wanjiru Mucheru
Irene Kittur
Isaac Karuna Karia
Jaafar Adan
Janet Asenwa Lunayo
Janeth Nyarinda Mamboreo
Jesca Nanjala Wabwile
Josephine Marani
Joyce Kavemba Mwangangi
Kigen Kipchirchir Benard
Lekakeny Karoi
Leonard Kipruto Koech
Mageto Jeremiah
Margaret Mwaila
Maureen Achieng Shikuku
Mohamed Noor
Moses Ouma
Nicodemus Barasa Okwara
Richard Ngunjiri Mwinga
Roseline Rehema Kiragu
Salome Mutie
Samuel Njoroge Kanyi
Scholasticah Chemtai Ngere
Simbe Teresa
Simeon Kamau Wanjiru
Stephen Mulinge Ndambuki
Stephen Mureithi Theuri
Tabitha Adoyo Adongo
Victoria Mutiso
Wato Galgallo Elema

Interviewers

Abdirissack Abdullahi Abdi
Abdirizak Dahir
Abdullahi Elmi Mohamed
Abigael Losike Longiti
Alicia Wabuti Wambani
Aliya Mohamed Noor
Amina Abdi Dida
Andrew Mutwii
Anne Silole Sorimpan
Beatrice Jeluget Kosgei
Beatrice Wanja Githendu
Bernice Aketch Mulongo
Beryl Cherotich
Beth Wanjiku Kamau
Bevelyne Jepkurui Koech
Caroline Jemutai Koima
Catherine Mukethe Matuku
Catherine Wagaki Kigombe
Getrude Jeruto Kosgei
Gilbert Majaliwa Abel
Glory Chepng'Eno Kilel
Grace Mbeyu James
Grace Moende Kivuti
Grace Njeri Kibathi
Hamdi Mohamed Hassan
Hanan Ahmed
Harriet Amina Masitsa
Husna Mwalimu Shee
Irene Nairesiae Konah
Isabellah Ratemo
Itai Farah Issa
Ivy Asimani Amang'Alia
Jeanne Wanjiku Gitau
Jonathan Saitoti Tumaina
Joy Kendi Mutwiri
Joyce Karaye Lodungokiok
Molly Akinyi Ojwando
Monica Wambua
Mulki Elmi
Mwanarusi Kigoto Mwazani
Naima Ali Golicha
Naliaka Valarie Otieno
Mkala Clemence Saru
Naomi Chenangat
Napunyu Silantoi Lengaina
Newton Mwirigi Gichuru
Ngandu Jane Wambui
Nimriyan Hamud Sheikh
Norah Achaira Muliro
Pamela Gakii
Pamela Nyakio Kivuti
Patrick Kahiga Wanjohi
Peris Chelagat Metto
Geldine Mbaabu

Celestine Kerubo Maake
Charity Ekure
Charity Ikote Mercy
Chebet Mercy Sigei
Chepkemoi Juliet
Chepkemoi Winnie Tonui
Christabel Shisia Juma
Christine Ann Mutete Wambua
Christine Mwende Gitari
Collins Kipleting Chepsiror
Damaris Basweti Anari
David Kiarie
Diana Jeruto
Diana Sang
Dorcas Jeruto
Dorcas Ndanu Mutuku
Dorry Atieno Omondi
Edith Joseph
Elizabeth Mukhaya
Emmaculate Cheptoo
Ephy Achieng Ogembo
Faith Malia Vaati
Mohamud Sharif
Faith Muthoni
Faith Vaati Kasyula
Fanis Chemtai
Farhan Abudi
Fatuma Abdalla Matamu
Fauzia Monica Achieng
Feizah Mohamed
Judy Chemutai
Judy Njeri Ngure
Julie Sharon Khaoma
Juliet Wandia Muthoni
Kelsie Rogito
Kennedy Rotich Loling'Anyang
Kevin Kipyegon Korir
Keziah Obukheri Washiali
Lavender Anyango
Leila Haret Sugow
Lilian Wanjiku Mugwanja
Loice Mawia Mumo
Lydia Zawadi Karisa
Lydiah Ogembo
Margaret Akeng'O Omusale
Margaret Senewa Mesenka
Martha Arbe Barile
Martha Njambi Chege
Mary Joy Akinyi Otieno
Mary Wairimu Nganga
Mary Waithira Mwangi
Mary Wangui Gakure
Maryann Wanja Kariuki
Maureen Njambi Njuguna
Mercy Mbinya Musau
Mercy Mwangangi
Mercy Wanjiru Kang'Ethe
Michelle Mburu
Millicent Akinyi
Millicent Kanini Lengaur
Peter Kampois Lemarkele
Phoebe Nasimiyu Wafula
Quinter Auma Odhiambo
Risper Kemunto Ongaki
Rose Adongo Okoth
Rukia Ali Mbwana
Ruth Nthambi Maingi
Sahra Mohamed Abdi
Salma Abdalla Ahmed
Salma Nyangule Wangara
Sarah Shali
Sarah Siwa
Shago Marysellah
Shanice Wami Gatumo
Sharon Chepkirui
George Thige Muchemi
Siteyia Tiampati
Solet Amondi
Stella Muthusi
Timothy Kiprotich Cheruiyot
Vanessa Mwende Mwema
Victoria Chemtai Koech
Vivian Mmbone
Wanjohi Mary Wanjiru
Wilkista Akinyi
Winjoy Mwende Mwenda
Winnie Awuor Awino
Zaynab Maende Olum
Zeinab Gababa Issack

ICF Staff

Name	Role
Sara Scates	Survey Manager
Keith Purvis	Data Processing Specialist
Ruilin Ren	Sampling Specialist
Natalie Shattuck	Production Specialist
Chris Gramer	Production Specialist
Kerry Aradhya	Editor
Paulo Medina	GIS Specialist
Sabina Behague	Dissemination Specialist
Olsen Hanner	Data Archivist
Yodit Bekele	Technical Reviewer
Annie Allnutt	Technical Reviewer
Michelle Winner	Technical Reviewer
Monica Kothari	Technical Reviewer
Joy Fishel	Technical Reviewer
Trevor Croft	Technical Reviewer
Cameron Taylor	Technical Reviewer
Suzanne Arrington	Technical Reviewer

QUESTIONNAIRES

Appendix **E**

Date: 20 July 2025

2025 KENYA MINI DEMOGRAPHIC AND HEALTH SURVEY (KMDHS)
HOUSEHOLD QUESTIONNAIRE

KENYA NATIONAL BUREAU OF STATISTICS

IDENTIFICATION

COUNTY _____

SUB COUNTY _____

LOCATION _____

SUB LOCATION _____

NAME OF HOUSEHOLD HEAD _____

KDHS CLUSTER NUMBER . . . . . [ ][ ][ ][ ]

HOUSEHOLD NUMBER . . . . . [ ][ ][ ][ ]

INTERVIEWER VISITS

	1	2	3	FINAL VISIT
DATE				DAY [ ][ ]
				MONTH [ ][ ]
				YEAR 2025
INTERVIEWER'S NAME				INT. NO. [ ][ ][ ][ ]
RESULT*				RESULT* [ ]
NEXT VISIT: DATE				TOTAL NUMBER OF VISITS [ ]
TIME				

*RESULT CODES:

1 COMPLETED
2 NO HOUSEHOLD MEMBER AT HOME OR NO COMPETENT RESPONDENT AT HOME AT TIME OF VISIT
3 ENTIRE HOUSEHOLD ABSENT FOR EXTENDED PERIOD OF TIME
4 POSTPONED
5 REFUSED
6 DWELLING VACANT OR ADDRESS NOT A DWELLING
7 DWELLING DESTROYED
8 DWELLING NOT FOUND
9 OTHER _____ (SPECIFY)

TOTAL PERSONS IN HOUSEHOLD [ ][ ]

TOTAL ELIGIBLE WOMEN [ ][ ]

LINE NO. OF RESPONDENT TO HOUSEHOLD QUESTIONNAIRE [ ][ ]

LANGUAGE OF QUESTIONNAIRE** 01 | LANGUAGE OF INTERVIEW** [ ][ ] | NATIVE LANGUAGE OF RESPONDENT** [ ][ ] | TRANSLATOR USED (YES = 1, NO = 2) [ ]

LANGUAGE OF QUESTIONNAIRE** **ENGLISH**

**LANGUAGE CODES:

01 ENGLISH	06 KAMBA	11 LUO	16 SOMALI
02 KISWAHILI	07 KIKUYU	12 MAASAI	17 TURKANA
03 BORANA	08 KISII	13 MERU	96 OTHER _____ (SPECIFY)
04 EMBU	09 LUHYA	14 MIJIKENDA	
05 KALENJIN	10 MARAGOLI	15 POKOT	

TEAM NUMBER [ ][ ]

TEAM SUPERVISOR NAME _____ NUMBER [ ][ ][ ][ ]

THIS PAGE IS INTENTIONALLY BLANK

INTRODUCTION AND CONSENT

Hello. My name is _____. I am working with Kenya National Bureau of Statistics. We are conducting a survey about health and other topics all over Kenya. The information we collect will help the government to plan health services. Your household was selected for the survey. I would like to ask you some questions about your household. The questions usually take about 30 to 60 minutes. All of the answers you give will be confidential and will not be shared with anyone other than members of our survey team. You don't have to be in the survey, but we hope you will agree to answer the questions since your views are important. If I ask you any question you don't want to answer, just let me know and I will go on to the next question or you can stop the interview at any time. In case you need more information about the survey, you may contact the person listed on this card.

GIVE CARD WITH CONTACT INFORMATION

Do you have any questions?
May I begin the interview now?

SIGNATURE OF INTERVIEWER _____ DATE _____

RESPONDENT AGREES TO BE INTERVIEWED . . 1 ↓

RESPONDENT DOES NOT AGREE TO BE INTERVIEWED . . 2 → END

##	RECORD THE TIME.	HOURS . . . . . [ ][ ] MINUTES . . . . . [ ][ ]

HOUSEHOLD SCHEDULE

LINE NO.	USUAL RESIDENTS AND VISITORS	RELATIONSHIP TO HEAD OF HOUSEHOLD	SEX	RESIDENCE		AGE	MORE PEOPLE	IF AGE 15 OR OLDER: MARITAL STATUS	ELIGIBILITY
1	2	3	4	5	6	7	7-1	8	9
	Please give me the names of the persons who usually live in your household and guests of the household who stayed here last night, starting with the head of the household. RECORD THE FIRST NAME OF THE HEAD OF THE HOUSEHOLD AFTER LISTING THE NAMES AND RECORDING THE RELATIONSHIP, SEX, RESIDENCE, AND AGE FOR EACH PERSON, ASK QUESTIONS 7A-7C TO BE SURE THAT THE LISTING IS COMPLETE. THEN ASK APPROPRIATE QUESTIONS IN COLUMNS 8-20 FOR EACH PERSON.	What is the relationship of {FULL NAME} to the head of the household? SEE CODES BELOW.	Is {FULL NAME} male or female?	Does {FULL NAME} usually live here?	Did {FULL NAME} stay here last night?	How old is {FULL NAME}? IF 95 OR MORE, RECORD '95'	Are there any other persons living in this household?	What is {FIRST NAME}'s current marital status? 1 = MARRIED OR LIVING TOGETHER 2 = DIVORCED 3=SEPARATED 4 = WIDOWED 5 = NEVER-MARRIED AND NEVER LIVED TOGETHER	CIRCLE LINE NUMBER OF ALL WOMEN AGE 15-49
01			M F 1 2	Y N 1 2	Y N 1 2	IN YEARS	Y N 1 → GO TO NEXT LINE 2 GO TO 7A ←		01
02			1 2	1 2	1 2		1 → GO TO NEXT LINE 2 GO TO 7A ←		02
03			1 2	1 2	1 2		1 → GO TO NEXT LINE 2 GO TO 7A ←		03
04			1 2	1 2	1 2		1 → GO TO NEXT LINE 2 GO TO 7A ←		04
05			1 2	1 2	1 2		1 → GO TO NEXT LINE 2 GO TO 7A ←		05

HOUSEHOLD SCHEDULE

LINE NO.	IF AGE 3 YEARS OR OLDER: EVER ATTENDED SCHOOL			IF AGE 0-4 YEARS: BIRTH REGISTRATION	IF USUAL HOUSEHOLD MEMBER: HEALTH INSURANCE			
	16	17A	17B	20	27	28	29	30
	Has {FIRST NAME} ever attended school or any early childhood education program?	What is the highest level of school {FIRST NAME} has attended?	What is the highest grade {FIRST NAME} completed at that level?	Does {FIRST NAME} have a birth certificate? IF NO, PROBE: Has {FIRST NAME}'s birth ever been registered with the civil authority? 1=YES REGISTERED WITH BIRTH CERTIFICATE 2 = YES REGISTERED WITHOUT BIRTH CERTIFICATE 3=NOT REGISTERED 8 = DON'T KNOW	CHECK 7: IF 18+: Is {FIRST NAME} registered under the Social Health Authority (SHA)? CHECK 7: IF LESS THAN 18: Is {FIRST NAME} registered under the Social Health Authority (SHA) or a dependent of a registered adult? IF YES, PROBE: Are they registered in their own right or as a dependent? 1 = YES, REGISTERED IN THEIR OWN RIGHT 2 = DEPENDENT 3 = NOT REGISTERED 8 = DON'T KNOW	Has {FIRST NAME's} premiums been paid for the current month for Social Health Insurance Fund (SHIF)? 1 = YES 2 = NO 8 = DON'T KNOW	Is {FIRST NAME} covered by any other health insurance? 1 = YES 2 = NO 8 = DON'T KNOW	What type of health insurance is {FIRST NAME} covered by? A=PRIVATE MEDICAL INSURANCE B = COMMUNITY BASED HEALTH INSURANCE C = COUNTY INSURANCE SCHEME D=EMPLOYER MANAGED INSURANCE SCHEME X = OTHER Z = DON'T KNOW
01	Y N 1 2 GO TO 20	LEVEL SEE CODES BELOW.	GRADE SEE CODES BELOW.		Y D N DK 1 2 3 8 GO TO 29	Y N DK 1 2 8	Y N DK 1 2 8 GO TO 101	A B C D X Z
02	1 2 GO TO 20				1 2 3 8 GO TO 29	1 2 8	1 2 8 GO TO 101	A B C D X Z
03	1 2 GO TO 20				1 2 3 8 GO TO 29	1 2 8	1 2 8 GO TO 101	A B C D X Z
04	1 2 GO TO 20				1 2 3 8 GO TO 29	1 2 8	1 2 8 GO TO 101	A B C D X Z
05	1 2 GO TO 20				1 2 3 8 GO TO 29	1 2 8	1 2 8 GO TO 101	A B C D X Z

7A) Just to make sure that I have a complete listing: are there any other people such as small children or infants that we have not listed? YES → ADD TO TABLE NO

7B) Are there any other people who may not be members of your family, such as domestic servants, lodgers, or friends who usually live here? YES → ADD TO TABLE NO

7C) Are there any guests or temporary visitors staying here, or anyone else who stayed here last night, who have not been listed? YES → ADD TO TABLE NO

CODES FOR Q. 3: RELATIONSHIP TO HEAD OF HOUSEHOLD

01 = HEAD
02 = WIFE OR HUSBAND
03 = SON OR DAUGHTER
04 = SON-IN-LAW OR DAUGHTER-IN-LAW
05 = GRANDCHILD
06 = PARENT
07 = PARENT-IN-LAW
08 = BROTHER OR SISTER
09 = OTHER RELATIVE
10 = ADOPTED/FOSTER/
11 = NOT RELATED
98 = DON'T KNOW

CODES FOR Q17 : EDUCATION

LEVEL

0 = PRE-PRIMARY
1 = PRIMARY
2 = JUNIOR
3 = POST-PRIMARY VOCATIONAL
4 = SECONDARY/"A" LEVEL/SENIOR SCHOOL
5 = MIDDLE LEVEL COLLEGE/TVET (CERTIFICATE/DIPLOMA)
6 = UNIVERSITY
7 = SNE (STAGE BASED)
8 = INFORMAL EDUCATION (MADRASA/ADULT BASIC)
9 = DON'T KNOW

GRADE

00 = LESS THAN 1 YEAR COMPLETED (USE '00' FOR Q. 17 ONLY. THIS CODE IS NOT ALLOWED FOR Q. 19.)
98 = DON'T KNOW

HOUSEHOLD CHARACTERISTICS

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
101	What is the main source of drinking water for members of your household?	**PIPED WATER** PIPED INTO DWELLING . . . . . 11 PIPED TO YARD/PLOT . . . . . 12 PIPED TO NEIGHBOR . . . . . 13 PUBLIC TAP/STANDPIPE . . . . . 14 TUBE WELL OR BOREHOLE . . . . . 21 **DUG WELL** PROTECTED WELL . . . . . 31 UNPROTECTED WELL . . . . . 32 **WATER FROM SPRING** PROTECTED SPRING . . . . . 41 UNPROTECTED SPRING . . . . . 42 RAINWATER . . . . . 51 TANKER TRUCK . . . . . 61 CART WITH SMALL TANK . . . . . 71 SURFACE WATER (RIVER/DAM/ LAKE/POND/STREAM/CANAL/ IRRIGATION CHANNEL) . . . . . 81 BOTTLED WATER . . . . . 91 OTHER _____ 96 (SPECIFY)	→ 109 → 103 → 103
102	What is the main source of water used by your household for other purposes such as cooking and handwashing?	**PIPED WATER** PIPED INTO DWELLING . . . . . 11 PIPED TO YARD/PLOT . . . . . 12 PIPED TO NEIGHBOR . . . . . 13 PUBLIC TAP/STANDPIPE . . . . . 14 TUBE WELL OR BOREHOLE . . . . . 21 **DUG WELL** PROTECTED WELL . . . . . 31 UNPROTECTED WELL . . . . . 32 **WATER FROM SPRING** PROTECTED SPRING . . . . . 41 UNPROTECTED SPRING . . . . . 42 RAINWATER . . . . . 51 TANKER TRUCK . . . . . 61 CART WITH SMALL TANK . . . . . 71 SURFACE WATER (RIVER/DAM/ LAKE/POND/STREAM/CANAL/ IRRIGATION CHANNEL) . . . . . 81 OTHER _____ 96 (SPECIFY)	→ 109
103	Where is that water source located?	IN OWN DWELLING . . . . . 1 IN OWN YARD/PLOT . . . . . 2 ELSEWHERE 3	→ 109
104	How long does it take to go there, get water, and come back?	MINUTES . . . . . ☐☐☐ DON'T KNOW 998	

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
109	What kind of toilet facility do members of your household usually use? IF NOT POSSIBLE TO DETERMINE, ASK PERMISSION TO OBSERVE THE FACILITY.	**FLUSH OR POUR FLUSH TOILET** FLUSH TO PIPED SEWER SYSTEM . . . . . 11 FLUSH TO SEPTIC TANK . . . . . 12 FLUSH TO PIT LATRINE . . . . . 13 FLUSH TO SOMEWHERE ELSE . . . . . 14 FLUSH, DON'T KNOW WHERE . . . . . 15 **PIT LATRINE** VENTILATED IMPROVED PIT LATRINE . . . . 21 PIT LATRINE WITH SLAB . . . . . 22 PIT LATRINE WITHOUT SLAB/OPEN PIT . . 23 COMPOSTING TOILET . . . . . 31 BIODIGESTER . . . . . 32 BUCKET TOILET . . . . . 41 HANGING TOILET/HANGING LATRINE . . . . . 51 NO FACILITY/BUSH/FIELD . . . . . 61 OTHER _____ 96 (SPECIFY)	→ 117
110	Do you share this toilet facility with other households?	YES . . . . . 1 NO 2	→ 112
111	Including your own household, how many households use this toilet facility?	NO. OF HOUSEHOLDS IF LESS THAN 10 . . . . . 0 _ 10 OR MORE HOUSEHOLDS . . . . . 95 DON'T KNOW 98	
112	Where is this toilet facility located?	IN OWN DWELLING . . . . . 1 IN OWN YARD/PLOT . . . . . 2 ELSEWHERE 3	
113	CHECK 109: CODES 12, 13, 21, 22, 23, OR 31 CIRCLED ☐ ↓ OTHER ☐		→ 117
114	CHECK 109: CODE 12 ☐ ↓ a) Has your septic tank ever been emptied? CODE 13, 21, 22, OR 23 ☐ ↓ b) Has your pit latrine ever been emptied? CODE 31 ☐ ↓ c) Has your composting toilet ever been emptied?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	☐ → 117
115	CHECK 109: CODE 12 ☐ ↓ a) The last time the septic tank was emptied, was it emptied by a service CODE 13, 21, 22, OR 23 ☐ ↓ b) The last time the pit latrine was emptied, was it emptied by a service provider? CODE 31 ☐ ↓ c) The last time the composting toilet was emptied, was it emptied by a service provider?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
116	Where were the contents emptied to?	A TREATMENT PLANT . . . . . 1 BURIED IN A COVERED PIT . . . . . 2 UNCOVERED PIT/BUSH/FIELD/OPEN GROUND 3 SURFACE WATER (RIVER/DAM/ LAKE/POND/STREAM/CANAL/ IRRIGATION CHANNEL) . . . . . 4 OTHER _____ 6 (SPECIFY) DON'T KNOW 8	
117	In your household, what type of cooking device or cookstove is mainly used for cooking?	ELECTRIC STOVE . . . . . 01 SOLAR COOKER . . . . . 02 LIQUEFIED PETROLEUM GAS (LPG)/ COOKING GAS STOVE . . . . . 03 PIPED NATURAL GAS STOVE . . . . . 04 BIOGAS STOVE . . . . . 05 LIQUID FUEL STOVE . . . . . 06 MANUFACTURED SOLID FUEL STOVE (JIKO . . 07 TRADITIONAL SOLID FUEL STOVE . . . . . 08 THREE STONE STOVE/OPEN FIRE . . . . . 09 NO FOOD COOKED IN HOUSEHOLD . . . . . 95 OTHER _____ 96 (SPECIFY)	→ 121 → 120 → 120 → 123 → 120
118	Does the stove have a chimney?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
120	What type of fuel or energy source is used in this cookstove?	ALCOHOL/ETHANOL . . . . . 01 GASOLINE/DIESEL . . . . . 02 KEROSENE/PARAFFIN . . . . . 03 COAL/LIGNITE . . . . . 04 CHARCOAL . . . . . 05 WOOD . . . . . 06 STRAW/SHRUBS/GRASS . . . . . 07 AGRICULTURAL CROP . . . . . 08 ANIMAL DUNG/WASTE . . . . . 09 PROCESSED BIOMASS (PELLETS) OR WOODCHIPS . . . . . 10 GARBAGE/PLASTIC . . . . . 11 SAWDUST . . . . . 12 OTHER _____ 96 (SPECIFY)	
121	Is the cooking usually done in the house, in a separate building, or outdoors?	IN THE HOUSE . . . . . 1 IN A SEPARATE BUILDING . . . . . 2 OUTDOORS . . . . . 3 OTHER _____ 6 (SPECIFY)	→ 123
122	Do you have a separate room which is used as a kitchen?	YES . . . . . 1 NO 2	

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
123	What does this household use to heat the home when needed? IF THE RESPONDENT SAYS ELECTRICITY OR GAS, ASK: What type of heater is the (electricity/gas) used in?	CENTRAL HEATING . . . . . 01 MANUFACTURED SPACE HEATER . . . . . 02 TRADITIONAL SPACE HEATER . . . . . 03 MANUFACTURED COOKSTOVE . . . . . 04 TRADITIONAL COOKSTOVE . . . . . 05 THREE STONE STOVE/OPEN FIRE . . . . . 06 FIRE PLACE . . . . . 07 NO HEATING IN HOUSEHOLD/NO NEED . . . . . 95 OTHER _____ 96 (SPECIFY)	→ 125 → 125 → 126 → 125
124	Does it have a chimney?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
125	What type of fuel or energy source is used in this heater?	ELECTRICITY . . . . . 01 PIPED NATURAL GAS . . . . . 02 SOLAR AIR HEATER . . . . . 03 LIQUEFIED PETROLEUM GAS (LPG) / COOKING GAS . . . . . 04 BIOGAS . . . . . 05 ALCOHOL/ETHANOL . . . . . 06 GASOLINE/DIESEL . . . . . 07 KEROSENE/PARAFFIN . . . . . 08 COAL/LIGNITE . . . . . 09 CHARCOAL . . . . . 10 WOOD . . . . . 11 STRAW/SHRUBS/GRASS . . . . . 12 AGRICULTURAL CRO . . . . . 13 ANIMAL DUNG/WASTI . . . . . 14 PROCESSED BIOMASS (PELLETS) OR WOODCHIPS . . . . . 15 GARBAGE/PLASTIC . . . . . 16 SAWDUST . . . . . 17 OTHER _____ 96 (SPECIFY)	

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
126	At night, what does your household mainly use to light the home?	ELECTRICITY FROM NATIONAL GRID .... 01 ELECTRICITY FROM GENERATOR ..... 02 ELECRICITY FROM SOLAR PANEL ..... 03 SOLAR LANTERN ..... 04 RE RE RE RECHARGEABLE FLASHLIGHT, TORCH OR LANTERN ..... 05 BATTERY POWERED FLASHLIGHT, TORCH OR LANTERN ..... 06 BIOGAS LAMP ..... 07 GASOLINE LAMP ..... 08 KEROSENE OR PARAFFIN LAMP ..... 09 CHARCOAL ..... 10 WOOD ..... 11 STRAW/SHRUBS/GRASS ..... 12 AGRICULTURAL CRO ..... 13 ANIMAL DUNG/WAST ..... 14 OIL LAMP ..... # CANDLE ..... # NO LIGHTING IN HOUSEHOLD ..... 95 OTHER _____ 96 (SPECIFY)	
127	How many rooms in this household are used for sleeping?	ROOMS	
128	Does this household own any livestock, herds, other farm animals, or poultry?	YES ..... 1 NO 2	→ 130

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
129	How many of the following animals does this household own? IF NONE, RECORD '00'. IF MORE THAN 95, RECORD '95'. IF UNKNOWN, RECORD '98'. a) Local cattle (indigenous)? b) Exotic/grade cattle? c) Horses, donkeys, or camels? d) Goats? e) Sheep? f) Chickens or other poultry? g) Pigs? h) Rabbits	a) LOCAL CATTLE (INDIGENOUS) . . [ ][ ] b) EXOTIC/GRADE CATTLE/ CROSSBREED . . . . . [ ][ ] c) HORSES/DONKEYS/MULES . . . . [ ][ ] d) GOATS . . . . . [ ][ ] e) SHEEP . . . . . [ ][ ] f) CHICKENS/POULTRY . . . . . [ ][ ] g) PIGS . . . . . [ ][ ] h) RABBITS . . . . . [ ][ ]	
130	Does any member of this household own any agricultural land?	YES . . . . . 1 NO 2	→ 131A
131	How many acres or hectares of agricultural land do members of this household own? ACRES/HECTARES: IF 995 OR MORE, RECORD '995.0' IN APPROPRIATE BOX. PLOT SIZE (SQ FT): IF 999995 OR MORE, RECORD '999995.0' IN APPROPRIATE BOX	ACRES . . . . . 1 [ ][ ][ ].[ ] HECTARES . . . . 2 [ ][ ][ ].[ ] SQ FT 3 [ ][ ][ ][ ][ ].[ ] DON'T KNOW 999998	
131A	Does any member of this household own any non-agricultural land?	YES . . . . . 1 NO 2	→ 132
131B	How many acres or hectares of non-agricultural land do members of this household own? ACRES/HECTARES: IF 995 OR MORE, RECORD '995.0' IN APPROPRIATE BOX. PLOT SIZE (SQ FT): IF 999995 OR MORE, RECORD '999995.0' IN APPROPRIATE BOX	ACRES . . . . . 1 [ ][ ][ ].[ ] HECTARES . . . . 2 [ ][ ][ ].[ ] SQ FT 3 [ ][ ][ ][ ][ ].[ ] DON'T KNOW 999998	

HOUSEHOLD CHARACTERISTICS

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES			SKIP
132	Does your household have:		YES	NO	
	a) Electricity?	a) ELECTRICITY	1	2	
	b) A radio?	b) RADIO	1	2	
	c) A television?	c) TELEVISION	1	2	
	d) A non-mobile telephone?	d) NON-MOBILE TELEPHONE	1	2	
	e) A computer?	e) COMPUTER	1	2	
	f) A refrigerator?	f) REFRIGERATOR	1	2	
	g) A solar panel?	g) SOLAR PANEL	1	2	
	h) A table?	h) TABLE	1	2	
	i) A chair?	i) CHAIR	1	2	
	j) A sofa?	j) SOFA	1	2	
	k) A bed?	k) BED	1	2	
	l) A cupboard?	l) CUPBOARD	1	2	
	m) A clock?	m) CLOCK	1	2	
	n) A microwave oven?	n) MICROWAVE OVEN	1	2	
	o) A DVD player?	o) DVD PLAYER	1	2	
	p) A cassette or CD player?	p) CASSETTE/CD PLAYER	1	2	
133	Does any member of this household own:		YES	NO	
	a) A watch?	a) WATCH	1	2	
	b) A mobile phone?	b) MOBILE PHONE	1	2	
	c) A bicycle?	c) BICYCLE	1	2	
	d) A motorcycle or motor scooter?	d) MOTORCYCLE/SCOOTER	1	2	
	e) An animal-drawn cart?	e) ANIMAL-DRAWN CART	1	2	
	f) A car or truck?	f) CAR/TRUCK	1	2	
	g) A boat with a motor?	g) BOAT WITH MOTOR	1	2	
134	Does any member of this household have an account in a bank or other financial institution?	YES NO		1 2	
135	Does any member of this household use a mobile phone to make financial transactions such as sending or receiving money, paying bills, purchasing goods or services, or receiving wages?	YES NO		1 2	
136	How often does anyone smoke inside your house? Would you say daily, weekly, monthly, less often than once a month, or never?	DAILY WEEKLY MONTHLY LESS OFTEN THAN ONCE A MONTH NEVER		1 2 3 4 5	
137	Does your household have any mosquito nets?	YES NO		1 2	→ 149
138	How many mosquito nets does your household have?	NUMBER OF NETS		☐	

MOSQUITO NETS

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
	ASK THE RESPONDENT TO SHOW YOU ALL THE NETS IN THE HOUSEHOLD. OBSERVE AND ANSWER THE QUESTIONS FOR EACH NET, ONE BY ONE.		
139	ASSIGN EACH NET A SEQUENTIAL NUMBER AND RECORD THE NUMBER HERE.	NET NUMBER	
140	I would like to take a look at the nets. Can you show me the first net? RECORD IF THE NET WAS OBSERVED OR NOT OBSERVED.	OBSERVED . . . . . 1 NOT OBSERVED 2	
141	How many months ago did your household get the mosquito net? IF LESS THAN ONE MONTH AGO, RECORD '00'.	MONTHS AGO . . . . . MORE THAN 36 MONTHS AGO . . . . . 95 NOT SURE 98	
142	OBSERVE OR ASK THE BRAND/TYPE OF MOSQUITO NET. IF BRAND IS UNKNOWN AND YOU CANNOT OBSERVE THE NET, SHOW PICTURES OF TYPICAL NET TYPES/BRANDS TO RESPONDENT.	**LONG-LASTING INSECTICIDE-TREATED NET (LLIN)** OLYSET (SUPANET EXTRA) . . . . . 11 PERMANET (SUPANET EXTRA) . . . . . 12 NETPROTECT . . . . . 13 YORKOOL . . . . . 14 DAWA PLUS . . . . . 15 DURANET . . . . . 16 MAGNET . . . . . 17 VEERALIN . . . . . 18 MIRANET . . . . . 19 OTHER/DON'T KNOW BRAND (LLI . . . . . 21 OTHER TYPE (NOT LLIN) . . . . . 96 DON'T KNOW TYPE 98	
143	Did you get the net through a mass distribution campaign, during an antenatal care visit, or during an immunization visit?	YES, MASS DISTRIBUTION CAMPAIGN 1 YES, ANC . . . . . 2 YES, CHILD WELFARE VISIT / IMMUNIZATION VISIT . . . . . 3 NO 4	→ 145

MOSQUITO NETS

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
144	Where did you get the net?	GOVERNMENT HEALTH FACILITY . . . . . 01 PRIVATE HEALTH FACILITY . . . . . 02 PHARMACY . . . . . 03 SHOP/MARKET . . . . . 04 CHW . . . . . 05 RELIGIOUS INSTITUTION . . . . . 06 SCHOOL . . . . . 07 OTHER . . . . . 96 DON'T KNOW 98	
145	Did anyone sleep under this mosquito net last night?	YES . . . . . 1 NO . . . . . 2 NOT SURE 8	 → 147 → 148

MOSQUITO NETS

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
146	Who slept under this mosquito net last night? RECORD THE PERSON'S NAME AND LINE NUMBER FROM HOUSEHOLD SCHEDULE.	NAME _____ LINE NUMBER . . . . . ☐☐ NAME _____ LINE NUMBER . . . . . ☐☐ NAME _____ LINE NUMBER . . . . . ☐☐ NAME _____ LINE NUMBER . . . . . ☐☐	→ 148
147	What was the main reason this net was not used last night?	TOO HOT . . . . . 01 DON'T LIKE NET SHAPE/COLOR/SIZE . . . . . 02 DON'T LIKE SMELL . . . . . 03 UNABLE TO HANG NET . . . . . 04 SLEPT OUTDOORS . . . . . 05 USUAL USER DIDN'T SLEEP HERE LAST NIGH 06 NO MOSQUITOES/NO MALARIA . . . . . 07 EXTRA NET/SAVING FOR LATE . . . . . 08 NET TOO SMALL/SHORT . . . . . 09 NET BROUGHT BEDBUGS . . . . . 10 NET TOO OLD/ TORN/DAMAGE . . . . . 11 NET IS TOO DIRTY . . . . . 12 NO PLACE TO HANG NET . . . . . 13 OTHER _____ 96 (SPECIFY)	
148	GO BACK TO 139 FOR NEXT NET; OR, IF NO MORE NETS, GO TO 149.		

HOUSEHOLD ENTERPRISES

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
154A	Now I would like to ask about any business or income-generating activity owned or operated by members of this household, excluding paid employment or unpaid work in another family member's business.		
154B	In the past month, excluding paid employment or unpaid work in another family member's business, has anyone in this household owned or operated any income-generating enterprise? Examples include a shop, tailoring or carpentry business, posho mill, jua kali work, water vending, or waste collection.	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	→ 160
154C	Please tell me the income-generating activities that members of this household have operated in the past month and the county where each activity is located. LIST UPTO 4 ESTABLISHMENTS/ENTERPRISES AND COUNTY OF OPERATION BEFORE COLLECTING INFORMATION ON EACH.	NAME OF ENTERPRISE _____ COUNTY CODE ☐☐	
154D	Now I would like to ask you some additional details about the businesses that members of your household operate or own. Please remember that your answers are completely confidential and will not be linked to your name or identity.		
154E	Who in the household operates or owns [ENTERPRISE]?	NAME _____ LINE NUMBER ☐☐ NAME _____ LINE NUMBER ☐☐	
154F	What is the main economic activity of [ENTERPRISE]?	MANUFACTURING . . . . . 1 SERVICES . . . . . 2 AGRI-BUSINESS 3	
154G	Is/was [ENTERPRISE] registered or licensed with any of the following?		

	YES	NO	DK
a) BUSINESS REGISTRATION SERVICE (BRS) FORMERLY	1	2	8
b) COUNTY GOVERNMENT	1	2	8
c) MICRO AND SMALL	1	2	8
d) PROFESSIONAL BODIES	1	2	8
e) SOCIAL DEVELOPMENT DEPARTMENT	1	2	8
f) OTHER (SPECIFY) .	1	2	8

a) Business Registration Service (BRS) formerly Registrar of Companies
b) County Government
c) Micro and Small Enterprise Authority (MSEA)
d) Professional Bodies
e) Social Development Department
f) Others (SPECIFY)

HOUSEHOLD ENTERPRISES

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
154H	How many persons are/were engaged in this income generating activity [ENTERPRISE]? IF THE ENTERPRISE IS OPERATED OR OWNED BY THE RESPONDENT, HE/SHE SHOULD BE INCLUDED IN THE COUNT	NUMBER OF PERSONS ENGAGED ☐☐	
154I	What is the total revenue from sales generated by [ENTERPRISE] during the [last month/last period of operation]? IF THE RESPONDENT IS ONLY ABLE TO PROVIDE THE DAILY SALES, PROBE TO ESTABLISH THE AVERAGE NUMBER OF DAYS WORKED IN A MONTH AND MULTIPLY WITH THE DAILY SALES TO OBTAIN THE MONTHLY VALUES	TOTAL REVENUE FROM SALES FOR A MONTH (KSh) ☐☐☐☐☐☐	
154J	GO BACK TO 154E FOR ADDITIONAL HOUSEHOLD ENTERPRISES OWNED OR OPERATED; OR, IF NO MORE ENTERPRISES, GO TO 160.		

HOUSEHOLD MEMBER DEATH REGISTRATION

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
160	Now, I would like to ask about recent deaths of your hourshold members. Since January 2020, has this household experienced the loss of any usual member, whether due to natural causes or other circumstances? Please do not include any pregnancy losses (including stillbirths or miscarriages).	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	→ 177A
161	How many usual members of this household have died between 2020-2025?	NUMBER OF HOUSEHOLD MEMBERS WHO DIED IN 2020-2025	
162	Now I would like to record the names of members of this households who died between 2020-2025, starting with the member who died most recently. LIST ALL USUAL HOUSEHOLD MEMBERS WHO DIED SINCE JANUARY 2020 STARTING FROM THE MOST RECENT. ASK Q163 TO QH175 FOR EACH DEATH ONE BY ONE.		
163	What is the name of the member of this household who died (most recently/previously)?	NAME _____	
164	Was {NAME IN QH163} male or female?	MALE . . . . . 1 FEMALE 2	
164A	What is the relationship of {NAME IN QH163} to the head of the household? SEE CODES ON PAGE 1		
165	In what month and year did {NAME IN QH163} die?	MONTH . . . . . YEAR	
166	IF MALE ↓ a) How old was {NAME IN QH163} when he died? IF DON'T KNOW, PROBE AND ASK ADDITIONAL QUESTIONS IF FEMALE ↓ b) How old was {NAME IN QH163} when she died? IF DON'T KNOW, PROBE AND ASK ADDITIONAL QUESTIONS	AGE AT DEATH IN COMPLETED YEARS	
167	Where did {NAME IN QH163} die?	IN A HEALTH FACILITY . . . . . 1 IN THE COMMUNITY . . . . . 2 DON'T KNOW 98	
167A	Was a death certificate issued for {NAME IN QH163} death? SHOW PICTURE	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 98	→ 177 → 168

HOUSEHOLD MEMBER DEATH REGISTRATION

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
167B	What was the main reason why the death certificate was not obtained for {NAME IN QH163}?	CULTURAL AND SOCIAL REASON . . . . . 1 ACCESS AND GEOGRAPHICAL ISSUE . . . . . 2 COST . . . . . 3 PERCEIVED LACK OF BENEFIT . . . . . 4 OTHER _____ 96 (SPECIFY) DON'T KNOW 98	
168	Was a burial permit issued when {NAME IN QH163} died?	YES . . . . . 1 → 177 NO . . . . . 2 DON'T KNOW . . . . . 98 → 177	
169	What was the main reason why a burial permit was not issued for {NAME IN QH163}?	CULTURAL AND SOCIAL REASON . . . . . 1 ACCESS AND GEOGRAPHICAL ISSUE . . . . . 2 COST . . . . . 3 PERCEIVED LACK OF BENEFIT . . . . . 4 OTHER _____ 96 (SPECIFY) DON'T KNOW 98	
177	GO BACK TO 162 FOR PREVIOUS DEATH THAT OCCURRED IN 2020-2025; OR, IF NO MORE DEATHS, GO TO 200.		

PERCEPTIONS OF HARM AGAINST WOMEN

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
177A	Now I'd like to ask you a few questions about how people in your community think and feel, so we can better understand the experiences of women and girls and whether they may face harm or violence.		
177AA	Are there specific groups of women or girls in your community who are at higher risk of violence? (e.g., indigenous women , migrants, etc.)	YES . . . . . 1 NO 2	
177AB	How safe do you think women or girls are in your community? Would you say very safe, safe, somewhat safe not very safe?	VERY SAFE . . . . . 1 SAFE . . . . . 2 SOMEWHAT SAFE . . . . . 3 NOT VERY SAFE 4	
177AC	How do you think people in your community would react if they suspected a woman or a girl was at risk of being killed by her husband, partner or other family member or anyone else ?	MOBILIZE COMMUNITY RESCU . . . . . A REPORT TO POLICE . . . . . B REPORT TO NGAO (CHIEF, VILLAGE LEADER OR OTHER COMMUNITY . . . . . C TALK TO FAMILY . . . . . D DO NOTHING . . . . . E OTHER _____ X (SPECIFY)	
177AD	In your community, are there social, cultural or religious beliefs that justify violence against women or girls ?	YES . . . . . 1 NO 2	
177AE	Have you heard people blame women or girls for the violence they experience?	YES . . . . . 1 NO 2	
177AF	Do you think domestic violence is a private matter that should be handled within the family?	YES . . . . . 1 NO 2	
205	RECORD THE TIME.	HOURS . . . . . [ ][ ] MINUTES . . . . . [ ][ ]	

ADDITIONAL HOUSEHOLD CHARACTERISTICS

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
149	We would like to learn about the places that households use to wash their hands. Can you please show me where members of your household most often wash their hands?	OBSERVED, FIXED PLACE . . . . . 1 OBSERVED, MOBILE . . . . . 2 NOT OBSERVED, NOT IN DWELLING/YARD/PL 3 NOT OBSERVED, NO PERMISSION TO S . . . . 4 NOT OBSERVED, OTHER REASOI 5	→ 152
150	OBSERVE PRESENCE OF WATER AT THE PLACE FOR HANDWASHING. RECORD OBSERVATION.	WATER IS AVAILABLE . . . . . 1 WATER IS NOT AVAILABLE 2	
151	OBSERVE PRESENCE OF SOAP, DETERGENT, OR OTHER CLEANSING AGENT AT THE PLACE OF HANDWASHING. RECORD OBSERVATION.	SOAP OR DETERGENT (BAR, LIQUID, POWDER, PASTE) . . . . . A ASH, MUD, SAND . . . . . B NONE . . . . . Y	
152	OBSERVE MAIN MATERIAL OF THE FLOOR OF THE DWELLING. RECORD OBSERVATION.	**NATURAL FLOOR** EARTH/SAND . . . . . 11 DUNG . . . . . 12 **RUDIMENTARY FLOOR** WOOD PLANKS . . . . . 21 PALM/BAMBOO . . . . . 22 **FINISHED FLOOR** PARQUET OR POLISHED WOO . . . . . 31 VINYL OR ASPHALT STRIPS . . . . . 32 CERAMIC TILES . . . . . 33 CEMENT . . . . . 34 CARPET . . . . . 35 OTHER _____ 96 (SPECIFY)	
153	OBSERVE MAIN MATERIAL OF THE ROOF OF THE DWELLING. RECORD OBSERVATION.	**NATURAL ROOFING** NO ROOF . . . . . 11 THATCH/GRASS/MAKUTI . . . . . 12 SOD/MUD/DUNG . . . . . 13 **RUDIMENTARY ROOFING** RUSTIC MAT . . . . . 21 PALM/BAMBOO . . . . . 22 WOOD PLANKS . . . . . 23 CARDBOARD . . . . . 24 TIN CANS . . . . . 25 **FINISHED ROOFING** IRON SHEETS/METAL . . . . . 31 WOOD . . . . . 32 CALAMINE/CEMENT FIBER . . . . . 33 CERAMIC TILES . . . . . 34 CEMENT . . . . . 35 ROOFING SHINGLES . . . . . 36 ASBESTOS SHEET . . . . . 37 OTHER _____ 96 (SPECIFY)	
154	OBSERVE MAIN MATERIAL OF THE EXTERIOR WALLS OF THE DWELLING. RECORD OBSERVATION.	**NATURAL WALLS** NO WALLS . . . . . 11 CANE/PALM/TRUNKS . . . . . 12 DIRT/ MUD 13 **RUDIMENTARY WALLS** BAMBOO WITH MUD . . . . . 21 STONE WITH MUD . . . . . 22 UNCOVERED ADOBE . . . . . 23 PLYWOOD . . . . . 24 CARDBOARD . . . . . 25 REUSED WOOD . . . . . 26 IRON SHEETS . . . . . 27 **FINISHED WALLS** CEMENT . . . . . 31 STONE WITH LIME/CEMENT . . . . . 32 BRICKS 33	

ADDITIONAL HOUSEHOLD CHARACTERISTICS

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
		CEMENT BLOCKS . . . . . 34 COVERED ADOBE . . . . . 35 WOOD PLANKS/SHINGLES . . . . . 36 OTHER _____ 96 (SPECIFY)	

INTERVIEWER'S OBSERVATIONS

TO BE FILLED IN AFTER COMPLETING INTERVIEW

COMMENTS ABOUT INTERVIEW:

COMMENTS ON SPECIFIC QUESTIONS:

ANY OTHER COMMENTS:

SUPERVISOR'S OBSERVATIONS

Date 20 July 2025

2025 KENYA MINI DEMOGRAPHIC AND HEALTH SURVEY (KMDHS)
WOMAN'S QUESTIONNAIRE

KENYA NATIONAL BUREAU OF STATISTICS

IDENTIFICATION

COUNTY _____

SUB COUNTY _____

LOCATION _____

SUB LOCATION _____

NAME OF HOUSEHOLD HEAD _____

KDHS CLUSTER NUMBER . . . . . [ ][ ][ ][ ]

HOUSEHOLD NUMBER . . . . . [ ][ ][ ][ ]

NAME AND LINE NUMBER OF WOMAN _____ [ ][ ]

WOMAN SELECTED FOR TFGBV MODULE? (1=YES, 2=NO) . . . . . [ ]

INTERVIEWER VISITS

	1	2	3	FINAL VISIT
DATE				DAY [ ][ ]
				MONTH [ ][ ]
				YEAR 2025
INTERVIEWER'S NAME				INT. NO. [ ][ ][ ][ ]
RESULT*				RESULT* [ ]
NEXT VISIT: DATE				TOTAL NUMBER OF VISITS [ ]
TIME				

*RESULT CODES: 1 COMPLETED; 2 NOT AT HOME; 3 POSTPONED; 4 REFUSED; 5 PARTLY COMPLETED; 6 INCAPACITATED; 7 OTHER _____ (SPECIFY)

LANGUAGE OF QUESTIONNAIRE** [0 1] LANGUAGE OF INTERVIEW** [ ][ ] NATIVE LANGUAGE OF RESPONDENT** [ ][ ] TRANSLATOR USED (YES = 1, NO = 2) [ ]

LANGUAGE OF QUESTIONNAIRE** **ENGLISH**

**LANGUAGE CODES:

01 ENGLISH	07 KIKUYU	12 MAASAI	17 TURKANA
02 KISWAHILI	08 KISII	13 MERU	96 OTHER _____ (SPECIFY)
03 BORANA	09 LUHYA	14 MIJIKENDA	
04 KALENJIN	10 MARAGOLI	15 POKOT	
06 KAMBA	11 LUO	16 SOMALI	

TEAM NUMBER [ ][ ]

TEAM SUPERVISOR NAME _____ NUMBER [ ][ ][ ][ ]

INTRODUCTION AND CONSENT

Hello. My name is _____. I am working with Kenya National Bureau of Statistics. We are conducting a survey about health and other topics all over Kenya. The information we collect will help the government to plan health services. Your household was selected for the survey. The questions usually take about 30 to 90 minutes. All of the answers you give will be confidential and will not be shared with anyone other than members of our survey team. You don't have to be in the survey, but we hope you will agree to answer the questions since your views are important. If I ask you any question you don't want to answer, just let me know and I will go on to the next question or you can stop the interview at any time.

In case you need more information about the survey, you may contact the person listed on the card that has already been given to your household.

Do you have any questions?
May I begin the interview now?

SIGNATURE OF INTERVIEWER _____ DATE _____

RESPONDENT AGREES TO BE INTERVIEWED . . 1 ↓

RESPONDENT DOES NOT AGREE TO BE INTERVIEWED . . 2 → END

SECTION 1. RESPONDENT'S BACKGROUND

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
101	RECORD THE TIME.	HOURS . . . . . ☐☐ MINUTES . . . . . ☐☐	
102	What county were you born in? REFER TO COUNTY CODE LIST IN INTERVIEWER'S MANUAL	COUNTY CODE . . . . . ☐☐ OUTSIDE OF KENYA 96	→ 104
103	What country were you born in?	COUNTRY _____ ☐☐	
104	How long have you been living continuously in (NAME OF CURRENT CITY, TOWN OR VILLAGE OF RESIDENCE)? IF LESS THAN ONE YEAR, RECORD '00' YEARS.	YEARS . . . . . ☐☐ ALWAYS/NEVER MOVED . . . . . 95 VISITOR 96	→ 110
105	CHECK 104: 00 - 04 YEARS ☐ ↓ 05 YEARS OR MORE ☐ →		107
106	In what month and year did you move here?	MONTH . . . . . ☐☐ DON'T KNOW MONTH . . . . . 98 YEAR . . . . . ☐☐☐☐ DON'T KNOW YEAR 9998	
107	Just before you moved here, which county did you live in? REFER TO COUNTY CODE LIST IN	COUNTY CODE . . . . . ☐☐ OUTSIDE OF KENYA 96	
108	Just before you moved here, did you live in a city, in a town, or in a rural area?	CITY (NAIROBI, KISUMU, MOMBASA, NAKURU) 1 TOWN . . . . . 2 RURAL AREA 3	

SECTION 1. RESPONDENT'S BACKGROUND

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
109	Why did you move to this place?	ECONOMIC REASONS/BUSINESS .... 01 EDUCATION/TRAINING .... 02 MARRIAGE FORMATION .... 03 FAMILY REUNIFICATION/OTHER FAMILY-RELATED REASON .... 04 FORCED DISPLACEMEN .... 05 OTHER _____ 96 (SPECIFY)	
110	In what month and year were you born?	MONTH .... ☐☐ DON'T KNOW MONTH .... 98 YEAR .... ☐☐☐☐ DON'T KNOW YEAR 9998	
111	How old were you at your last birthday? COMPARE AND CORRECT 110 AND/OR 111 IF INCONSISTENT.	AGE IN COMPLETED YEARS .... ☐☐	
112	In general, would you say your health is very good, good, moderate, bad, or very bad?	VERY GOOD .... 1 GOOD .... 2 MODERATE .... 3 BAD .... 4 VERY BAD 5	
113	Have you ever attended school?	YES .... 1 NO 2	→ 117
114	What is the highest level of school you attended: primary, post-primary vocational, secondary/'A' level, college, or university?	PRIMARY .... 1 JUNIOR 2 POST-PRIMARY VOCATIONAL TRAINING .. 3 SECONDARY/ 'A' LEVEL/SENIOR SCHOOL 4 MIDDLE LEVEL COLLEGE (CERTIFICATE/ DIPLOMA) .... 5 UNIVERSITY .... 6 SNE (STAGE BASED) 7 INFORMAL EDUCATION (MADRASA/ ADULT BASIC) 8	
115	What is the highest (standard/grade/form/year) you completed at that level? IF COMPLETED LESS THAN ONE YEAR AT THAT LEVEL, RECORD '00'.	STANDARD/GRADE/FORM/YEAR .. ☐☐	
116	CHECK 114: PRIMARY, POST-PRIMARY VOCAT., VOCAT.SECONDARY, INFORMAL ☐ ↓ UNIVERSITY ☐ →		119
117	Now I would like you to read this sentence to me. SHOW CARD TO RESPONDENT. IF RESPONDENT CANNOT READ WHOLE SENTENCE, PROBE: Can you read any part of the sentence to me?	CANNOT READ AT ALL .... 1 ABLE TO READ ONLY PART OF THE SENTENC .... 2 ABLE TO READ WHOLE SENTENCE .... 3 NO CARD WITH REQUIRED LANGUAGE _____ 4 (SPECIFY LANGUAGE) BLIND/VISUALLY IMPAIRED 5	

SECTION 1. RESPONDENT'S BACKGROUND

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
118	CHECK 117: CODE '2', '3' OR '4' CIRCLED ☐↓ CODE '1' OR '5' CIRCLED ☐→ 120		
119	Do you read a newspaper or magazine at least once a week, less than once a week or not at all?	AT LEAST ONCE A WEEI . . . . . 1 LESS THAN ONCE A WEEK . . . . . 2 NOT AT ALL 3	
120	Do you listen to the radio at least once a week, less than once a week or not at all?	AT LEAST ONCE A WEEI . . . . . 1 LESS THAN ONCE A WEEK . . . . . 2 NOT AT ALL 3	
121	Do you watch television at least once a week, less than once a week or not at all?	AT LEAST ONCE A WEEI . . . . . 1 LESS THAN ONCE A WEEK . . . . . 2 NOT AT ALL 3	
122	Do you own a mobile phone?	YES . . . . . 1 NO 2	→ 127
123	Is your mobile phone a smart phone?	YES . . . . . 1 NO 2	
127	Have you ever used the Internet from any location on any device?	YES . . . . . 1 NO 2	→ 130
128	In the last 12 months, have you used the Internet? IF NECESSARY, PROBE FOR USE FROM ANY LOCATION, WITH ANY DEVICE.	YES . . . . . 1 NO 2	→ 130
129	During the last one month, how often did you use the Internet: almost every day, at least once a week, less than once a week, or not at all?	ALMOST EVERY DAY . . . . . 1 AT LEAST ONCE A WEEI . . . . . 2 LESS THAN ONCE A WEEK . . . . . 3 NOT AT ALL 4	
130	What is your religion?	CATHOLIC . . . . . 01 PROTESTANT . . . . . 02 EVANGELICAL CHURCHES . . . . . 03 AFRICAN INSTITUTED CHURCHES . . . . . 04 ORTHODOX . . . . . 05 OTHER CHRISTIAN . . . . . 06 ISLAM . . . . . 07 HINDU . . . . . 08 TRADITIONISTS . . . . . 09 NO RELIGION/ ATHEISTS . . . . . 10 OTHER _____ 96 (SPECIFY)	
131	What is your ethnic group?	EMBU . . . . . 01 KALENJIN . . . . . 02 KAMBA . . . . . 03 KIKUYU . . . . . 04 KISII . . . . . 05 LUHYA . . . . . 06 LUO . . . . . 07 MAASAI 08	

SECTION 1. RESPONDENT'S BACKGROUND

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
		MERU . . . . . 09 MIJIKENDA/SWAHILI . . . . . 10 SOMALI . . . . . 11 TAITA/TAVETA . . . . . 12 OTHER _____ 96 (SPECIFY)	

SECTION 2. REPRODUCTION

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
201	Now I would like to ask about all the births you have had during your life. Have you ever given birth?	YES . . . . . 1 NO 2	→ 206
202	Do you have any sons or daughters to whom you have given birth who are now living with you?	YES . . . . . 1 NO 2	→ 204
203	a) How many sons live with you? IF NONE, RECORD '00'. b) And how many daughters live with you? IF NONE, RECORD '00'.	a) SONS AT HOME . . . . . ☐☐ b) DAUGHTERS AT HOME . . . . . ☐☐	
204	Do you have any sons or daughters to whom you have given birth who are alive but do not live with	YES . . . . . 1 NO 2	→ 206
205	a) How many sons are alive but do not live with you? b) And how many daughters are alive but do not live with you? IF NONE, RECORD '00'.	a) SONS ELSEWHERE . . . . . ☐☐ b) DAUGHTERS ELSEWHERE . . . . ☐☐	
206	Have you ever given birth to a boy or girl who was born alive but later died? IF NO, PROBE: Any baby who cried, who made any movement, sound, or effort to breathe, or who showed any other signs of life even if for a very short	YES . . . . . 1 NO 2	→ 208
207	a) How many boys have died? IF NONE, RECORD '00'. b) And how many girls have died? IF NONE, RECORD '00'.	a) BOYS DEAD . . . . . ☐☐ b) GIRLS DEAD . . . . . ☐☐	
208	SUM ANSWERS TO 203, 205, AND 207, AND ENTER TOTAL. IF NONE, RECORD '00'.	TOTAL LIVE BIRTHS . . . . . ☐☐	
209	Just to make sure that I have this right: you have had in total {NUMBER OF BIRTHS} births during your life. Is that correct? YES ☐ ↓ NO ☐ → PROBE AND CORRECT 201-208 AS NECESSARY		
210	Women sometimes have a pregnancy that does not result in a live birth. For example, a pregnancy can end in a miscarriage, an abortion, or the child can be born dead. Have you ever had a pregnancy that did not end in a live birth?	YES . . . . . 1 NO 2	→ 212
211	How many miscarriages, abortions, and stillbirths have you had?	PREGNANCY LOSSES . . . . . ☐☐	
212	SUM ANSWERS TO 208 AND 211 AND ENTER TOTAL. IF NONE, RECORD '00'.	TOTAL PREGNANCY OUTCOMES . . ☐☐	
213	CHECK 212: ONE OR MORE PAST PREGNANCIES ☐ ↓ NO PAST PREGNANCIES ☐ →		→ 232

214 Now I would like to record all your pregnancies including live births, stillbirths, miscarriages, and abortions, starting with your first pregnancy.

RECORD ALL PREGNANCIES. RECORD TWINS AND TRIPLETS ON SEPARATE LINES.

PREGNANCY HISTORY LINE NUMBER	215	216	217	218	219	220	221	222
	IF ROW=01: Think back to your first pregnancy. Was that a single pregnancy, twins, or triplets? IF ROW>01: Think back to your next pregnancy. Was that a single pregnancy, twins, or triplets?	IF 215=SING: Was the baby born alive, born dead, or did you have a miscarriage or abortion? IF 215>1: FIRST OF MULT. Was the first baby in this pregnancy born alive or born dead? NEXT MULT. Was the next baby in this pregnancy born alive or born dead?	Did the baby cry, move, or breathe?	What name was given to the baby? RECORD NAME.	Is {NAME IN 218} a boy or a girl?	CHECK 216 AND 217: TYPE OF PREGNANCY OUTCOME. NOTE: IF 217=1, THEN PREGNANCY OUTCOME = BORN ALIVE. IF BORN On what day, month, and year was {NAME IN 218} born? IF BORN DEAD, MISCARRIAGE OR On what day, month, and year did this pregnancy end?	How long did this pregnancy last in weeks or months? RECORD IN COMPLETED WEEKS OR MONTHS.	IF ROW=01: Were there any other pregnancies before this pregnancy? IF ROW>01: Were there any other pregnancies between the previous pregnancy and this pregnancy? IF 215>1 AND THIS IS NOT THE FIRST BIRTH OF THE PREGNANCY, SKIP TO 216 IN NEXT ROW.
01	SING ..... 1 TWINS ..... 2 TRIP ..... 3 QUAD ..... 4 QUIN 5	BORN ALIVE ..... 1 (SKIP TO 218) BORN DEAD ..... 2 MISCARRIAGE 3 (SKIP TO 220) ABORTION 4	YES ... 1 NO ..... 2 (SKIP TO 220)	NAME	BOY ... 1 GIRL .. 2	DAY MONTH YEAR	WEEKS 1 MONTHS 2	YES ..... 1 (ADD PREGNANCY) NO ..... 2 (NEXT ROW)
02	SING ..... 1 TWINS ..... 2 TRIP ..... 3 QUAD ..... 4 QUIN 5	BORN ALIVE ..... 1 (SKIP TO 218) BORN DEAD ..... 2 MISCARRIAGE 3 (SKIP TO 220) ABORTION 4	YES ... 1 NO ..... 2 (SKIP TO 220)	NAME	BOY ... 1 GIRL .. 2	DAY MONTH YEAR	WEEKS 1 MONTHS 2	YES ..... 1 (ADD PREGNANCY) NO ..... 2 (NEXT ROW)
03	SING ..... 1 TWINS ..... 2 TRIP ..... 3 QUAD ..... 4 QUIN 5	BORN ALIVE ..... 1 (SKIP TO 218) BORN DEAD ..... 2 MISCARRIAGE 3 (SKIP TO 220) ABORTION 4	YES ... 1 NO ..... 2 (SKIP TO 220)	NAME	BOY ... 1 GIRL .. 2	DAY MONTH YEAR	WEEKS 1 MONTHS 2	YES ..... 1 (ADD PREGNANCY) NO ..... 2 (NEXT ROW)

222A	Have you had any pregnancies that ended since the last pregnancy	YES ..... 1→ADD TO TABLE NO 2
222B	READ THE LIST OF PREGNANCY OUTCOMES IN ORDER TO THE RESPONDENT AND ASK IF THEY ARE ALL THAT SHE HAS EVER HAD, AND IF THEY ARE LISTED IN ORDER STARTING FROM THE FIRST ONE. DOES THE RESPONDENT AGREE? IF NOT, PROBE FOR THE CORRECT INFORMATION AND REVISE THE PREGNANCY HISTORY ACCORDINGLY.	

PREGNANCY HISTORY LINE NUMBER	223	224	225	226	227	228
			IF BORN ALIVE AND STILL LIVING:			IF BORN ALIVE AND NOW DEAD:
	CHECK 216, 217 AND 221: IF 216=1 OR 217=1, THEN PREGNANCY OUTCOME = BORN ALIVE. IF 216=2 OR 3, THEN CHECK 221. IF 221 ≥ 7 MONTHS OR 28 WEEKS, THEN PREGNANCY OUTCOME = BORN DEAD. IF 221 < 7 MONTHS OR 28 WEEKS, FINAL PREGNANCY OUTCOME = MISCARRIAGE. IF 216=4, THEN PREGNANCY OUTCOME = ABORTION.	Is {NAME IN 218} still alive?	IF 219=BOY How old was {NAME IN 218} at his last birthday? RECORD AGE IN COMPLETED IF 219=GIR How old was {NAME IN 218} at her last birthday? RECORD AGE IN COMPLETED YEARS.	Is {NAME IN 218} living with you?	RECORD HOUSEHOLD LINE NUMBER OF CHILD. RECORD '00' IF CHILD NOT LISTED IN HOUSEHOLD.	IF 219=BOY: How old was {NAME IN 218} when he died? IF '12 MONTHS' OR '1 YR', ASK: Did {NAME IN 218} have his first birthday? THEN ASK: Exactly how many months old was {NAME IN 218} when he died? RECORD DAYS IF LESS THAN 1 MONTH; MONTHS IF LESS THAN TWO YEARS; OR YEARS. IF 219=GIRL: How old was {NAME IN 218} when she died? IF '12 MONTHS' OR '1 YR', ASK: Did {NAME IN 218} have her first birthday? THEN ASK: Exactly how many months old was {NAME IN 218} when she died? RECORD DAYS IF LESS THAN 1 MONTH; MONTHS IF LESS THAN TWO YEARS; OR YEARS.
01	BORN ALIVE ..... 1 1 BORN DEAD ..... 2 MISCARRIAGE .... 3 ABORTION 4	YES ..... 1 NO ..... 2 (SKIP TO 228)	AGE IN YEARS ☐☐	YES ..... 1 NO 2	HOUSEHOLD LINE NUMBER ☐☐ (SKIP TO 223 IN NEXT ROW)	DAYS 1 ☐☐ MONTHS 2 ☐☐ YEARS 3 ☐☐ (SKIP TO 223 IN NEXT ROW)
02	BORN ALIVE ..... 1 BORN DEAD ..... 2 MISCARRIAGE .... 3 ABORTION 4	YES ..... 1 NO ..... 2 (SKIP TO 228)	AGE IN YEARS ☐☐	YES ..... 1 NO 2	HOUSEHOLD LINE NUMBER ☐☐ (SKIP TO 223 IN NEXT ROW)	DAYS 1 ☐☐ MONTHS 2 ☐☐ YEARS 3 ☐☐ (SKIP TO 223 IN NEXT ROW)
03	BORN ALIVE ..... 1 BORN DEAD ..... 2 MISCARRIAGE .... 3 ABORTION 4	YES ..... 1 NO ..... 2 (SKIP TO 228)	AGE IN YEARS ☐☐	YES ..... 1 NO 2	HOUSEHOLD LINE NUMBER ☐☐ (SKIP TO 223 IN NEXT ROW)	DAYS 1 ☐☐ MONTHS 2 ☐☐ YEARS 3 ☐☐ (SKIP TO 223 IN NEXT ROW)

SECTION 2. REPRODUCTION

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
230	COMPARE 212 WITH NUMBER OF PREGNANCY OUTCOMES IN PREGNANCY HISTORY NUMBER IN PREGNANCY HISTORY IS GREATER THAN OR EQUAL TO 212 ☐ NUMBER IN PREGNANCY HISTORY IS LESS THAN 212 ☐ (PROBE AND RECONCILE)		
231	**C** FOR EACH LIVE BIRTH IN 2020-2025, ENTER 'B' IN THE MONTH OF BIRTH IN THE CALENDAR. WRITE THE NAME OF THE CHILD TO THE LEFT OF THE 'B' CODE. FOR EACH LIVE BIRTH, RECORD 'P' IN EACH OF THE PRECEDING MONTHS ACCORDING TO THE DURATION OF PREGNANCY. (NOTE: THE NUMBER OF 'P's MUST BE ONE LESS THAN THE NUMBER OF MONTHS THAT THE PREGNANCY LASTED.) FOR EACH PREGNANCY THAT DID NOT END IN A LIVE BIRTH IN 2020-2025, ENTER 'T' IN THE CALENDAR IN THE MONTH THAT THE PREGNANCY TERMINATED AND 'P' FOR THE REMAINING NUMBER OF COMPLETED MONTHS OF PREGNANCY. IF DURATION OF PREGNANCY WAS REPORTED IN WEEKS, MULTIPLY THE NUMBER OF WEEKS BY 0.23 TO CONVERT TO THE NUMBER OF MONTHS. ROUND DOWN TO THE NEAREST WHOLE NUMBER TO GET THE NUMBER OF COMPLETED MONTHS.		
232	Are you pregnant now?	YES . . . . . 1 NO . . . . . 2 UNSURE 8	2 → 236
233	How many weeks or months pregnant are you? RECORD NUMBER OF COMPLETED WEEKS OR MONTHS. **C** ENTER 'P's IN THE CALENDAR, BEGINNING WITH THE MONTH OF INTERVIEW AND FOR THE TOTAL NUMBER OF COMPLETED MONTHS. IF DURATION OF PREGNANCY WAS REPORTED IN WEEKS, MULTIPLY THE NUMBER OF WEEKS BY 0.23 TO CONVERT TO THE NUMBER OF MONTHS. ROUND DOWN TO THE NEAREST WHOLE NUMBER TO GET THE NUMBER OF COMPLETED	WEEKS . . . . . 1 ☐☐ MONTHS . . . . . 2 ☐☐	
234	When you got pregnant, did you want to get pregnant at that time?	YES . . . . . 1 NO 2	2 → 236
235	CHECK 208: TOTAL NUMBER OF LIVE BIRTHS ONE OR MORE ☐ / NONE ☐ a) Did you want to have a baby later on or did you not want any more children? b) Did you want to have a baby later on or did you not want any children?	LATER . . . . . 1 NO MORE/NONE 2	
236	When did your last menstrual period start? _____ (DATE, IF GIVEN)	DAYS AGO . . . . . 1 ☐☐ WEEKS AGO . . . . . 2 ☐☐ MONTHS AGO . . . . . 3 ☐☐ YEARS AGO . . . . . 4 ☐☐ IN MENOPAUSE/HAS HAD HYSTERECT(. . 994 BEFORE LAST PREGNANC` . . . . . 995 NEVER MENSTRUATED 996	994, 995 → 240 996 → 241

SECTION 2. REPRODUCTION

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
237	CHECK 236: WAS THE LAST MENSTRUAL PERIOD WITHIN THE LAST YEAR? YES, WITHIN LAST YEAR ☐ ↓ NO, ONE YEAR OR MORE ☐ → 240		
238	During your last menstrual period, what did you use to collect or absorb your menstrual blood? Anything else?	REUSABLE SANITARY PAD . . . . . A DISPOSABLE SANITARY PADS . . . . . B TAMPONS . . . . . C MENSTRUAL CUP . . . . . D CLOTH . . . . . E TOILET PAPER . . . . . F COTTON WOOL . . . . . G UNDERWEAR ONLY . . . . . H OTHER _____ X (SPECIFY) NOTHING . . . . . Y	
239	During your last menstrual period, were you able to wash and change in privacy while at home?	YES . . . . . 1 NO . . . . . 2 AWAY FROM HOME DURING LAST MENSTRUAL PERIOD 3	
240	How old were you when you had your first menstrual period?	AGE . . . . . ☐☐ DON'T KNOW 98	
241	From one menstrual period to the next, are there certain days when a woman is more likely to become pregnant?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	→ 243
242	Is this time just before her period begins, during her period, right after her period has ended, or halfway between two periods?	JUST BEFORE HER PERIOD BEGINS . . . . . 1 DURING HER PERIOD . . . . . 2 RIGHT AFTER HER PERIOD HAS ENDED . . . . 3 HALFWAY BETWEEN TWO PERIODS . . . . . 4 OTHER _____ 6 (SPECIFY) DON'T KNOW 8	
243	After the birth of a child, can a woman become pregnant before her menstrual period has returned?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	

SECTION 3. CONTRACEPTION

301	Now I would like to talk about family planning - the various ways or methods that a couple can use to delay or avoid a pregnancy.	
01	Have you heard of Female Sterilization? PROBE: Women can have an operation to avoid having any more children.	YES . . . . . 1 NO 2
02	Have you heard of Male Sterilization? PROBE: Men can have an operation to avoid having any more children. May also be referred to as vasectomy.	YES . . . . . 1 NO 2
03	Have you heard of IUD? PROBE: Women can have a loop or coil placed inside them by a doctor or a nurse which can prevent pregnancy for one or more years.	YES . . . . . 1 NO 2
04	Have you heard of Injectables? PROBE: Women can have an injection by a health provider that stops them from becoming pregnant for one or more months.	YES . . . . . 1 NO 2
05	Have you heard of Implants? PROBE: Women can have one or more small rods placed in their upper arm by a doctor, nurse, or other health provider which can prevent pregnancy for one or more years.	YES . . . . . 1 NO 2
06	Have you heard of Pill? PROBE: Women can take a pill every day to avoid becoming pregnant.	YES . . . . . 1 NO 2
07	Have you heard of Male Condom? PROBE: Men can put a rubber sheath on their penis before sexual intercourse.	YES . . . . . 1 NO 2
08	Have you heard of Female Condom? PROBE: Women can place a sheath in their vagina before sexual intercourse.	YES . . . . . 1 NO 2
09	Have you heard of Emergency Contraception? PROBE: As an emergency measure, within 3 days after they have unprotected sexual intercourse, women can take special pills to prevent pregnancy.	YES . . . . . 1 NO 2
10	Have you heard of Standard Days Method? PROBE: A woman uses a string of colored beads to know the days she can get pregnant. On the days she can get pregnant, she uses a condom or does not have sexual intercourse.	YES . . . . . 1 NO 2
11	Have you heard of Lactational Amenorrhea Method (LAM)? PROBE: Up to 6 months after childbirth, before the menstrual period has returned, women use a method requiring frequent breastfeeding day and night.	YES . . . . . 1 NO 2
12	Have you heard of Rhythm Method? PROBE: To avoid pregnancy, women do not have sexual intercourse on the days of the month they think they can get	YES . . . . . 1 NO 2
13	Have you heard of Withdrawal? PROBE: Men can be careful and pull out before climax.	YES . . . . . 1 NO 2
14	Have you heard of any other ways or methods that women or men can use to avoid pregnancy?	YES, MODERN METHOD A _____ (SPECIFY) YES, TRADITIONAL METHOD B _____ (SPECIFY) NO . . . . . Y

SECTION 3. CONTRACEPTION

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
302	CHECK 232: NOT PREGNANT OR UNSURE ☐ PREGNANT ☐ → 317		
303	Are you or your partner currently doing something or using any method to delay or avoid getting	YES . . . . . 1 NO 2	→ 307
304	Are you or your partner sterilized? IF YES: Who is sterilized, you or your partner?	YES, RESPONDENT STERILIZED ONLY . . . . 1 YES, PARTNER STERILIZED ONLY . . . . . 2 YES, BOTH STERILIZED . . . . . 3 NO, NEITHER STERILIZED 4	→ 306
305	CHECK 304: RESPONDENT STERILIZED ONLY ☐ → PROCEED TO 307. CIRCLE CODE 'A' AND FOLLOW THE SKIP INSTRUCTION. PARTNER STERILIZED ONLY ☐ → PROCEED TO 307. CIRCLE CODE 'B' AND FOLLOW THE SKIP INSTRUCTION. BOTH STERILIZED ☐ → PROCEED TO 307. CIRCLE CODE 'A' AND CODE 'B' AND FOLLOW		
306	Just to check, are you or your partner doing any of the following to avoid pregnancy: deliberately avoiding sex on certain days, using a condom, using withdrawal or using emergency	YES . . . . . 1 NO 2	→ 317
307	Which method are you using? RECORD ALL MENTIONED. IF MORE THAN ONE METHOD MENTIONED, FOLLOW SKIP INSTRUCTION FOR HIGHEST METHOD IN LIST.	FEMALE STERILIZATION . . . . . A MALE STERILIZATION . . . . . B IUD . . . . . C INJECTABLES . . . . . D IMPLANTS . . . . . E PILL . . . . . F MALE CONDOM . . . . . G FEMALE CONDOM . . . . . H EMERGENCY CONTRACEPTIO . . . . . I STANDARD DAYS METHOD . . . . . J LACTATIONAL AMENORRHEA METHOD . . . . K RHYTHM METHOD . . . . . L WITHDRAWAL . . . . . M OTHER MODERN METHOD . . . . . X OTHER TRADITIONAL METHOD . . . . . Y	A → 312 B C → 314 D E → 314 F → 314 G → 314 H I J K L → 314 M X Y
308	Now I'm going to show you two pictures. Please point to the picture that best matches what was used the last time you received your injectable. SHOW IMAGES OF SAYANA PRESS AND REGULAR SYRINGE.	DMPA-SC/SAYANA PRESS . . . . . 1 NEEDLE AND SYRINGE . . . . . 2 DON'T KNOW 8	→ 314
309	The last time you received your injectable, did you inject DMPA-SC/Sayana Press yourself or did a health care provider do it for you?	SELF-INJECTION . . . . . 1 INJECTION GIVEN BY HEALTH CARE PROVIDER . . . . . 2 OTHER (FRIEND/RELATIVE ETC) . . . . . 3 DON'T KNOW 8	→ 314

SECTION 3. CONTRACEPTION

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
312	In what facility did the sterilization take place? PROBE TO IDENTIFY THE TYPE OF SOURCE. IF UNABLE TO DETERMINE IF PUBLIC, PRIVATE, OR NGO SECTOR, RECORD '96' AND WRITE THE NAME OF THE PLACE.	**PUBLIC SECTOR** GOVERNMENT HOSPITAL . . . . . 11 GOVERNMENT HEALTH CENTER . . . . 12 GOVERNMENT DISPENSARY . . . . . 13 MOBILE CLINIC . . . . . 14 OTHER PUBLIC SECTOR _____ 16 (SPECIFY) **PRIVATE MEDICAL SECTOR** PRIVATE HOSPITAL . . . . . 21 PRIVATE CLINIC . . . . . 22 PRIVATE DOCTOR'S OFFIC . . . . . 23 MOBILE CLINIC . . . . . 24 OTHER PRIVATE MEDICAL SECTOR _____ 26 (SPECIFY) **NGO MEDICAL SECTOR** NGO HOSPITAI . . . . . 31 NGO CLINIC . . . . . 32 OTHER NGO MEDICAL SECTOR _____ 36 (SPECIFY) **FBO MEDICAL SECTOR** FBO/MISSION HOSPITAL . . . . . 41 FBO/MISSION CLINIC . . . . . 42 FBO MOBILE CLINIC . . . . . 43 OTHER FBO SECTOR _____ 46 (SPECIFY) OTHER _____ 96 (SPECIFY) DON'T KNOW 98	
313	In what month and year was the sterilization performed?	MONTH . . . . . ☐☐ YEAR . . . . . ☐☐☐☐	→ 315
314	Since what month and year have you been using {METHOD} without stopping? PROBE: For how long have you been using {METHOD} now without stopping?	MONTH . . . . . ☐☐ YEAR . . . . . ☐☐☐☐	
315	CHECK 313 AND 314, AND 220: ANY LIVE BIRTH, STILLBIRTH, MISSCARRIAGE OR ABORTION AFTER MONTH AND YEAR OF START OF USE OF CONTRACEPTION IN 313 OR 314? NO ☐ ↓ YES ☐ → GO BACK TO 313 OR 314, PROBE AND RECORD MONTH AND YEAR AT START OF CONTINUOUS USE OF CURRENT METHOD (MUST BE AFTER LAST BIRTH OR PREGNANCY		

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
316	CHECK 313 AND 314: YEAR IS 2020-2025 ☐ **C** ENTER CODE FOR METHOD USED IN MONTH OF INTERVIEW IN THE CALENDAR AND IN EACH MONTH BACK TO THE DATE STARTED USING. THEN CONTINUE ↓ YEAR IS 2019 OR EARLIER ☐ **C** ENTER CODE FOR METHOD USED IN MONTH OF INTERVIEW IN THE CALENDAR AND EACH MONTH BACK TO JANUARY 2020 . THEN → (SKIP TO 329)		
317	I would like to ask you some questions about the times you or your partner may have used a method to avoid getting pregnant during the last few years. USE CALENDAR TO PROBE FOR EARLIER PERIODS OF USE AND NONUSE, STARTING WITH MOST RECENT USE, BACK TO JANUARY 2020. USE NAMES OF CHILDREN, DATES OF BIRTH, AND PERIODS	**C**	
317A	MONTH AND YEAR OF START OF INTERVAL OF USE OR NON-USE.	MONTH . . . . . ☐☐ YEAR . . . . . ☐☐☐☐	
317B	Between {EVENT ONE} in {MONTH/YEAR ONE} and {EVENT TWO} in {MONTH/YEAR TWO}, did you or your partner use any method of	YES . . . . . 1 NO 2	→ 317I
317C	Which method was that?	METHOD CODE . . . . . ☐	
317D	How many months after {EVENT ONE} in {MONTH/YEAR ONE} did you start to use the {METHOD}? RECORD '95' IF THE RESPONDENT SAYS THE DATE OF STARTING TO USE THE METHOD.	IMMEDIATELY . . . . . 00 MONTHS . . . . . ☐☐ DATE GIVEN 95	→ 317F
317E	RECORD MONTH AND YEAR RESPONDENT STARTED USING METHOD.	MONTH . . . . . ☐☐ YEAR . . . . . ☐☐☐☐	
317F	For how many months did you use the {METHOD} continuously? RECORD '95' IF RESPONDENT GAVE THE DATE OF TERMINATION OF USE	MONTHS . . . . . ☐☐ DATE GIVEN 95	→ 317H
317G	RECORD MONTH AND YEAR RESPONDENT STOPPED USING METHOD.	MONTH . . . . . ☐☐ YEAR . . . . . ☐☐☐☐	
317H	Why did you stop using {METHOD}?	REASON STOPPED . . . . . ☐	
317I	GO BACK TO 317A FOR NEXT GAP; OR, IF NO MORE GAPS, GO TO 318.		

SECTION 3. CONTRACEPTION

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
318	Have you used emergency contraception in the last 12 months? That is, have you taken special pills within 3 days after having unprotected sexual intercourse to prevent pregnancy?	YES . . . . . 1 NO 2	
319	CHECK THE CALENDAR FOR USE OF ANY CONTRACEPTIVE METHOD IN ANY MONTH NO METHOD USED ☐ ↓ ANY METHOD USED ☐ →		321
320	Have you ever used anything or tried in any way to delay or avoid getting pregnant?	YES . . . . . 1 NO 2	331
321	CHECK 307: CIRCLE METHOD CODE: IF MORE THAN ONE METHOD CODE CIRCLED IN 307, CIRCLE CODE FOR HIGHEST METHOD IN LIST.	NO CODE CIRCLED . . . . . 00 FEMALE STERILIZATION . . . . . 01 MALE STERILIZATION . . . . . 02 IUD . . . . . 03 INJECTABLES . . . . . 04 IMPLANTS . . . . . 05 PILL . . . . . 06 MALE CONDOM . . . . . 07 FEMALE CONDOM . . . . . 08 EMERGENCY CONTRACEPTION . . . . . 09 STANDARD DAYS METHOD . . . . . 10 LACTATIONAL AMENORRHEA METHOD . . . . 11 RHYTHM METHOD . . . . . 12 WITHDRAWAL . . . . . 13 OTHER MODERN METHOD . . . . . 95 OTHER TRADITIONAL METHOD 96	331 324 332 332 332
322	You first started using {METHOD} in {DATE FROM 314}. Where did you get it at that time? PROBE TO IDENTIFY THE TYPE OF SOURCE. IF UNABLE TO DETERMINE IF PUBLIC, PRIVATE, OR NGO SECTOR, RECORD '96' AND WRITE THE NAME OF THE PLACE.	**PUBLIC SECTOR** GOVERNMENT HOSPITAL . . . . . 11 GOVERNMENT HEALTH CENTER . . . . . 12 GOVERNMENT DISPENSARY . . . . . 13 MOBILE CLINIC . . . . . 14 COMMUNITY HEALTH WORKER/ FIELDWORKER . . . . . 15 OTHER PUBLIC SECTOR _____ 16 (SPECIFY) **PRIVATE MEDICAL SECTOR** PRIVATE HOSPITAL . . . . . 21 PRIVATE CLINI . . . . . 22 PHARMACY . . . . . 23 PRIVATE DOCTOR . . . . . 24 MOBILE CLINIC . . . . . 25 COMMUNITY HEALTH WORKER / FIELDWORKER . . . . . 26 OTHER PRIVATE MEDICAL SECTOR _____ 27 (SPECIFY) **NGO MEDICAL SECTOR** NGO HOSPITAL . . . . . 31 NGO CLINIC . . . . . 32 OTHER NGO MEDICAL SECTOR _____ 36 (SPECIFY) **FBO MEDICAL SECTOR** FBO/MISSION HOSPITAL . . . . . 41 FBO/MISSION CLINIC . . . . . 42 FBO MOBILE CLINIC . . . . . 43 OTHER FBO SECTOR _____ 46 (SPECIFY)	

SECTION 3. CONTRACEPTION

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
		OTHER SOURCE SHOP . . . . . 51 CHURCH . . . . . 52 FRIEND/RELATIVE . . . . . 53 OTHER _____ 96 (SPECIFY)	

SECTION 3. CONTRACEPTION

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
323	At that time, were you told about side effects or problems you might have with the method?	YES . . . . . 1 NO 2	→ 325
324	When you got sterilized, were you told about side effects or problems you might have with the	YES . . . . . 1 NO 2	
325	Were you told what to do if you experienced side effects or problems?	YES . . . . . 1 NO 2	
326	At that time, were you told about other methods of family planning that you could use?	YES . . . . . 1 NO 2	
327	CHECK 307: CIRCLE METHOD CODE: IF MORE THAN ONE METHOD CODE CIRCLED IN 307, CIRCLE CODE FOR HIGHEST METHOD IN LIST.	FEMALE STERILIZATION . . . . . 01 IUD . . . . . 03 INJECTABLES . . . . . 04 IMPLANTS . . . . . 05 PILL . . . . . 06 MALE CONDOM . . . . . 07 FEMALE CONDOM . . . . . 08 EMERGENCY CONTRACEPTION. . . . . 09 STANDARD DAYS METHOD . . . . . 10 OTHER MODERN METHOD 95	→ 332
328	At that time, were you told that you could switch to another method if you wanted to or needed to?	YES . . . . . 1 NO 2	→ 330
329	CHECK 307: CIRCLE METHOD CODE: IF MORE THAN ONE METHOD CODE CIRCLED IN 307, CIRCLE CODE FOR HIGHEST METHOD IN LIST.	FEMALE STERILIZATION . . . . . 01 MALE STERILIZATION . . . . . 02 IUD . . . . . 03 INJECTABLES . . . . . 04 IMPLANTS . . . . . 05 PILL . . . . . 06 MALE CONDOM . . . . . 07 FEMALE CONDOM . . . . . 08 EMERGENCY CONTRACEPTION . . . . . 09 STANDARD DAYS METHOD . . . . . 10 LACTATIONAL AMENORRHEA METHOD. . . . . 11 RHYTHM METHOD. . . . . 12 WITHDRAWAL . . . . . 13 OTHER MODERN METHOD . . . . . 95 OTHER TRADITIONAL METHOD. 96	→ 332 → 332 → 332

SECTION 3. CONTRACEPTION

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
330	Where did you obtain {METHOD} the last time? PROBE TO IDENTIFY THE TYPE OF SOURCE. IF UNABLE TO DETERMINE IF PUBLIC, PRIVATE, OR NGO SECTOR, RECORD '96' AND WRITE THE NAME OF THE PLACE.	**PUBLIC SECTOR** GOVERNMENT HOSPITAL . . . . . 11 GOVERNMENT HEALTH CENTER . . . . . 12 GOVERNMENT DISPENSARY . . . . . 13 MOBILE CLINIC . . . . . 14 COMMUNITY HEALTH WORKER/ FIELDWORKER . . . . . 15 OTHER PUBLIC SECTOR _____ 16 (SPECIFY) **PRIVATE MEDICAL SECTOR** PRIVATE HOSPITAL . . . . . 21 PRIVATE CLINIC . . . . . 22 PHARMACY . . . . . 23 PRIVATE DOCTOR . . . . . 24 MOBILE CLINIC . . . . . 25 COMMUNITY HEALTH WORKER FIELDWORKER . . . . . 26 OTHER PRIVATE MEDICAL SECTOR _____ 27 (SPECIFY) **NGO MEDICAL SECTOR** NGO HOSPITAI . . . . . 31 NGO CLINIC . . . . . 32 OTHER NGO MEDICAL SECTOR _____ 36 (SPECIFY) **FBO MEDICAL SECTOR** FBO/MISSION HOSPITAL . . . . . 41 FBO/MISSION CLINIC . . . . . 42 FBO MOBILE CLINIC . . . . . 43 OTHER FBO SECTOR _____ 46 (SPECIFY) **OTHER SOURCE** SHOP . . . . . 51 CHURCH . . . . . 52 FRIEND/RELATIVE . . . . . 53 OTHER _____ 96 (SPECIFY)	→ 332
331	Do you know of a place where you can obtain a method of family planning?	YES . . . . . 1 NO 2	
332	In the last 12 months, were you visited by a fieldworker?	YES . . . . . 1 NO 2	→ 334
333	Did the fieldworker talk to you about family planning?	YES . . . . . 1 NO 2	
334	CHECK 202: CHILDREN LIVING WITH YES ↓ NO ↓ a) In the last 12 months, | b) In the last 12 months,	YES 1	

SECTION 3. CONTRACEPTION

NO.	QUESTIONS AND FILTERS		CODING CATEGORIES	SKIP
	have you visited a health facility for care for yourself or your children?	have you visited a health facility for care for yourself?	NO 2	→ 401
335	Did any staff member at the health facility speak to you about family planning methods?		YES . . . . . 1 NO 2	

SECTION 4. PREGNANCY AND POSTNATAL CARE

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
401	CHECK 220 AND 225: ONE OR MORE PREGNANCY OUTCOMES 0-35 MONTHS BEFORE THE SURVEY ☐ NO PREGNANCY OUTCOMES 0-35 MONTHS BEFORE ☐ → 601		
402	CHECK 220. LIST THE PREGNANCY HISTORY NUMBER IN 215 FOR EACH PREGNANCY OUTCOME 0-35 MONTHS BEFORE THE SURVEY, STARTING FROM THE LAST ONE. CLASSIFY EACH PREGNANCY OUTCOME BY TYPE USING 223 AND THE ORDER OF OUTCOMES IN THE PREGNANCY HISTORY. **PREGNANCY OUTCOME TYPE** MOST RECENT LIVE BIRTH 1 PRIOR LIVE BIRTH 2 MOST RECENT STILLBIRTH 3 PRIOR STILLBIRTH 4 ABORTION OR MISCARRIAGE 5 PREGNANCY HISTORY NUMBE . . . . ☐☐ PREGNANCY OUTCOME TYPE . . . . . ☐ PREGNANCY HISTORY NUMBE . . . . ☐☐ PREGNANCY OUTCOME TYPE . . . . . ☐ PREGNANCY HISTORY NUMBE . . . . ☐☐ PREGNANCY OUTCOME TYPE . . . . . ☐ PREGNANCY HISTORY NUMBE . . . . ☐☐ PREGNANCY OUTCOME TYPE . . . . . ☐ PREGNANCY HISTORY NUMBE . . . . ☐☐ PREGNANCY OUTCOME TYPE . . . . . ☐ PREGNANCY HISTORY NUMBE . . . . ☐☐ PREGNANCY OUTCOME TYPE . . . . . ☐		
403	Now I would like to ask some questions about your pregnancies in the last 3 years. We will talk about each separately, starting with the last one you had.		
404	PREGNANCY HISTORY NUMBER FROM 402.	PREGNANCY HISTORY NUMBER . . . . . ☐☐	
405	PREGNANCY OUTCOME TYPE FROM 402.	MOST RECENT LIVE BIRTH . . . . . 1 PRIOR LIVE BIRTH . . . . . 2 MOST RECENT STILLBIRTH . . . . . 3 PRIOR STILLBIRTH . . . . . 4 MISCARRIAGE/ABORTION 5	→ 407
406	RECORD DATE PREGNANCY ENDED FROM 220.	DAY . . . . . ☐☐ MONTH . . . . . ☐☐ YEAR . . . . . ☐☐☐☐	→ 408
407	RECORD NAME FROM 218. NAME_____		
408	CHECK 405: PREGNANCY TYPE 1 OR 2 ☐ → a) When you got pregnant with {NAME IN 407}, did you want to get pregnant at that time? PREGNANCY TYPE 3, 4, OR 5 ☐ → b) When you got pregnant with the pregnancy that ended in {DATE FROM 406}, did you want to get pregnant at that time?	YES . . . . . 1 NO 2	→ 411

SECTION 4. PREGNANCY AND POSTNATAL CARE

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
409	Did you want to have a baby later on, or not at all?	LATER . . . . . 1 NOT AT ALL 2	→ 411
410	How much longer did you want to wait?	MONTHS . . . . . 1 [ ][ ] YEARS . . . . . 2 [ ][ ] DON'T KNOW 998	
411	CHECK 405: PREGNANCY OUTCOME TYPE	MOST RECENT LIVE BIRTH . . . . . 1 PRIOR LIVE BIRTH . . . . . 2 MOST RECENT STILLBIRTH . . . . . 3 PRIOR STILLBIRTH . . . . . 4 ABORTION/MISCARRIAGE 5	→ 434 → 434 → 475
412	Did you see anyone for antenatal care for this pregnancy?	YES . . . . . 1 NO 2	→ 414
413	CHECK 405: PREGNANCY OUTCOME TYPE MOST RECENT LIVE BIRTH (SKIP TO 420) ← ☐ MOST RECENT STILLBIRTH ☐ →		→ 426
414	Whom did you see? Anyone else? PROBE TO IDENTIFY EACH TYPE OF PERSON AND RECORD ALL MENTIONED.	**HEALTH PERSONNEL** DOCTOR . . . . . A NURSE/MIDWIFE/CLINICAL OFFICER . . . . B **OTHER PERSON** TRADITIONAL BIRTH ATTENDANT . . . . . D COMMUNITY HEALTH WORKER/ FIELD WORKER . . . . . E OTHER _____ X (SPECIFY)	
415	Where did you receive antenatal care for this pregnancy? Anywhere else? PROBE TO IDENTIFY TYPE OF SOURCE. IF UNABLE TO DETERMINE IF PUBLIC, PRIVATE, OR NGO SECTOR, RECORD 'X' AND WRITE THE NAME OF THE PLACE(S).	**HOME** HER HOME . . . . . A OTHER HOME . . . . . B **PUBLIC SECTOR** GOVERNMENT HOSPITAL . . . . . C GOVERNMENT HEALTH CENTER . . . . . D GOVERNMENT DISPENSARY . . . . . E OTHER PUBLIC SECTOR F _____ (SPECIFY) **PRIVATE MEDICAL SECTOR** PRIVATE HOSPITAL . . . . . G PRIVATE CLINIC . . . . . H OTHER PRIVATE MEDICAL SECTOR I _____ (SPECIFY) **NGO MEDICAL SECTOR** NGO HOSPITAL . . . . . J NGO CLINIC . . . . . K OTHER NGO MEDICAL SECTOR L _____ (SPECIFY) **FBO MEDICAL SECTOR** FBO/MISSION HOSPITAL . . . . . M FBO/MISSION CLINIC . . . . . N OTHER FBO MEDICAL SECTOR O _____ (SPECIFY)	

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
		OTHER _____ (SPECIFY) X	

SECTION 4. PREGNANCY AND POSTNATAL CARE

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
416	How many weeks or months pregnant were you when you first received antenatal care for this pregnancy?	WEEKS . . . . . 1 [ ][ ] MONTHS . . . . . 2 [ ][ ] DON'T KNOW 998	
417	How many times did you receive antenatal care during this pregnancy?	NUMBER OF TIMES . . . . . [ ][ ] DON'T KNOW 98	
418	As part of your antenatal care during this pregnancy, did a healthcare provider do any of the following: a) Measure your blood pressure? b) Take a urine sample? c) Take a blood sample? d) Listen to the baby's heartbeat? e) Talk with you about which foods or how much food you should eat? f) Talk with you about breastfeeding? g) Ask you if you had vaginal bleeding?	YES NO DK a) BP . . . . . 1 2 8 b) URINE . . . . . 1 2 8 c) BLOOD . . . . . 1 2 8 d) HEARTBEAT . . . . . 1 2 8 e) FOODS . . . . . 1 2 8 f) BREASTFEED . . . . . 1 2 8 g) BLEEDING 1 2 8	
419	CHECK 405: PREGNANCY OUTCOME TYPE MOST RECENT LIVE BIRTH ☐ ↓ MOST RECENT STILLBIRTH ☐ →		426
420	During this pregnancy, were you given an injection in the arm to prevent the baby from getting tetanus after birth?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	423

SECTION 4. PREGNANCY AND POSTNATAL CARE

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
421	During this pregnancy, how many times did you get a tetanus injection?	TIMES . . . . . ☐☐ DON'T KNOW 8	
422	CHECK 421: ONE TIME OR DK ☐ ↓ TWO OR MORE TIMES ☐ →		426
423	At any time before this pregnancy, did you receive any tetanus injections?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	→ 426
424	Before this pregnancy, how many times did you receive a tetanus injection? IF 7 OR MORE TIMES, RECORD '7'.	TIMES . . . . . ☐☐ DON'T KNOW 8	
425	CHECK 424: ONLY ONE ☐ ↓ a) How many years ago did you receive that tetanus injection? MORE THAN ONE ☐ ↓ b) How many years ago did you receive the last tetanus injection prior to this	YEARS AGC . . . . . ☐☐	
426	During this pregnancy, were you given or did you buy any iron tablets, IFAS tablets or iron syrup? SHOW IRON TABLETS/IFAS TABLETS/SYRUP/IRON SYRUPS.	YES, IRON TABLETS . . . . . 1 YES, IFAS TABLETS . . . . . 2 NO . . . . . 3 DON'T KNOW 8	→ 429

SECTION 4. PREGNANCY AND POSTNATAL CARE

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
427	Where did you get the iron tablets/IFAS or syrup? Anywhere else? PROBE TO IDENTIFY THE TYPE OF SOURCE. IF UNABLE TO DETERMINE IF PUBLIC, PRIVATE, OR NGO SECTOR, RECORD 'X' AND WRITE THE NAME OF THE PLACE(S).	**PUBLIC SECTOR** GOVERNMENT HOSPITAL . . . . . A GOVERNMENT HEALTH CENTER . . . . . B GOVERNMENT DISPENSARY . . . . . C MOBILE CLINIC . . . . . D COMMUNITY HEALTH WORKER/ FIELDWORKER . . . . . E OTHER PUBLIC SECTOR F _____ (SPECIFY) **PRIVATE MEDICAL SECTOR** PRIVATE HOSPITAL . . . . . G PRIVATE CLINIC . . . . . H PHARMACY . . . . . I PRIVATE DOCTOR . . . . . J MOBILE CLINIC . . . . . K COMMUNITY HEALTH WORKER/ FIELDWORKER . . . . . L OTHER PRIVATE MEDICAL SECTOR M _____ (SPECIFY) **NGO MEDICAL SECTOR** NGO HOSPITAL . . . . . N NGO CLINIC . . . . . O OTHER NGO MEDICAL SECTOR P _____ (SPECIFY) **FBO MEDICAL SECTOR** FBO/MISSION HOSPITAL . . . . . Q FBO/MISSION CLINIC . . . . . R FBO MOBILE CLINIC . . . . . S OTHER FBO MEDICAL SECTOR T _____ (SPECIFY) **OTHER SOURCE** SHOP . . . . . U MARKET . . . . . V MASS DISTRIBUTION CAMPAIGN . . . . W OTHER _____ X (SPECIFY)	
428	During the whole pregnancy, for how many days did you take the iron tablets/IFAS or syrup? IF ANSWER IS NOT NUMERIC, PROBE FOR APPROXIMATE NUMBER OF DAYS.	DAYS . . . . . ☐☐☐ DON'T KNOW 998	
429	During this pregnancy, did you take any medicine for intestinal worms?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
431	During this pregnancy, did you take SP/Fansidar to keep you from getting malaria?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	→ 434
432	How many times did you take SP/Fansidar during this pregnancy?	TIMES . . . . . ☐☐	

SECTION 4. PREGNANCY AND POSTNATAL CARE

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
433	Did you get the SP/Fansidar during any antenatal care visit, during another visit to a health facility or from another source? IF MORE THAN ONE SOURCE, RECORD THE HIGHEST SOURCE ON THE LIST.	ANTENATAL VISIT . . . . . 1 ANOTHER FACILITY VISIT . . . . . 2 OTHER SOURCE 6	
434	CHECK 405: PREGNANCY TYPE 1 OR 2 → a) Who assisted with the delivery of {NAME IN 407}? Anyone else? PROBE FOR THE TYPE(S) OF PERSON(S) AND RECORD ALL MENTIONED. IF RESPONDENT SAYS NO ONE ASSISTED, PROBE TO DETERMINE WHETHER ANY ADULTS WERE PRESENT AT THE DELIVERY. PREGNANCY TYPE 3 OR 4 → b) Who assisted with the delivery of the stillbirth you had in {DATE FROM 406}? Anyone else? PROBE FOR THE TYPE(S) OF PERSON(S) AND RECORD ALL MENTIONED. IF RESPONDENT SAYS NO ONE ASSISTED, PROBE TO DETERMINE WHETHER ANY ADULTS WERE PRESENT AT THE DELIVERY.	**HEALTH PERSONNEL** DOCTOR . . . . . A NURSE/MIDWIFE/CLINICAL OFFICER . . . . B **OTHER PERSON** TRADITIONAL BIRTH ATTENDANT . . . . . C RELATIVE/FRIEND . . . . . D OTHER _____ X (SPECIFY) NO ONE ASSISTED . . . . . Y	
435	CHECK 405: PREGNANCY TYPE 1 OR 2 → a) Where did you give birth to {NAME IN 407}? PROBE TO IDENTIFY THE TYPE OF SOURCE. IF UNABLE TO DETERMINE IF PUBLIC, PRIVATE, OR NGO SECTOR, RECORD '96' AND WRITE THE NAME OF THE PLACE. PREGNANCY TYPE 3 OR 4 → b) Where did you deliver this stillbirth? PROBE TO IDENTIFY THE TYPE OF SOURCE. IF UNABLE TO DETERMINE IF PUBLIC, PRIVATE, OR NGO SECTOR, RECORD '96' AND WRITE THE NAME OF THE PLACE.	**HOME** HER HOME . . . . . 11 OTHER HOME . . . . . 12 → 437 **PUBLIC SECTOR** GOVERNMENT HOSPITAL . . . . . 21 GOVERNMENT HEALTH CENTER . . . . . 22 GOVERNMENT DISPENSARY . . . . . 23 OTHER PUBLIC SECTOR 26 _____ (SPECIFY) **PRIVATE MEDICAL SECTOR** PRIVATE HOSPITAL . . . . . 31 PRIVATE CLINIC . . . . . 32 OTHER PRIVATE MEDICAL SECTOR 36 _____ (SPECIFY) **NGO MEDICAL SECTOR** NGO HOSPITAL . . . . . 41 NGO CLINIC . . . . . 42 OTHER NGO MEDICAL SECTOR 46 _____ (SPECIFY) **FBO MEDICAL SECTOR** FBO/MISSION HOSPITAL . . . . . 51 FBO/MISSION CLINIC . . . . . 52 OTHER FBO MEDICAL SECTOR 56 _____ (SPECIFY) OTHER _____ 96 → 437 (SPECIFY)	

SECTION 4. PREGNANCY AND POSTNATAL CARE

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
436	CHECK 405: PREGNANCY TYPE 1 OR 2 → a) Was {NAME IN 407} delivered by caesarean, that is, did they cut your belly open to take the baby out? PREGNANCY TYPE 3 OR 4 → b) Was this stillbirth delivered by caesarean, that is, did they cut your belly open to take the baby out?	YES . . . . . 1 NO 2	
437	CHECK 405: PREGNANCY OUTCOME TYPE	MOST RECENT LIVE BIRTH . . . . . 1 PRIOR LIVE BIRTH . . . . . 2 MOST RECENT STILLBIRTH . . . . . 3 PRIOR STILLBIRTH 4	→ 441 → 445 → 487
438	After the birth, was {NAME IN 407} put on your chest?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	→ 441
439	Was {NAME IN 407}'s bare skin touching your bare skin?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	→ 441
440	How long after birth was {NAME IN 407} put on the bare skin of your chest? PROBE FOR A NUMERIC RESPONSE. IF LESS THAN 1 HOUR, RECORD '00' HOURS; IF 24 HOURS OR MORE, RECORD 24.	IMMEDIATELY . . . . . 00 HOURS . . . . . ☐☐	
441	When {NAME IN 407} was born, was {NAME IN 407} very large, larger than average, average, smaller than average, or very small?	VERY LARGE . . . . . 1 LARGER THAN AVERAG . . . . . 2 AVERAGE . . . . . 3 SMALLER THAN AVERAGE . . . . . 4 VERY SMALL . . . . . 5 DON'T KNOW 8	
442	Was {NAME IN 407} weighed at birth?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	→ 444
443	How much did {NAME IN 407} weigh? RECORD WEIGHT IN KILOGRAMS FROM HEALTH CARD, IF AVAILABLE.	KG FROM CARD 1 ☐.☐☐☐ KG FROM RECALL 2 ☐.☐☐☐ DON'T KNOW 99998	
444	CHECK 405: PREGNANCY OUTCOME TYPE MOST RECENT LIVE BIRTH →	PRIOR LIVE BIRTH ☐	→ 480
445	CHECK 435: PLACE OF DELIVERY FACILITY BIRTH: ANY CODE 21 THROUGH 56 CIRCLED →	CODE 11, 12, OR 96 CIRCLED ☐	→ 464

SECTION 4. PREGNANCY AND POSTNATAL CARE

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
447	CHECK 405: PREGNANCY TYPE 1 → a) How long after {NAME IN 407} was delivered did you stay in the {FACILITY IN 435}? PREGNANCY TYPE 3 → b) For the stillbirth you had in {DATE FROM 406}, how long after the baby was born did you stay in the {FACILITY IN 435}? IF LESS THAN ONE DAY, RECORD HOURS; IF LESS THAN ONE WEEK, RECORD DAYS.	HOURS . . . . . 1 ☐☐ DAYS . . . . . 2 ☐☐ WEEKS . . . . . 3 ☐☐ DON'T KNOW 998	
448	I would like to talk to you about checks on your health after delivery, for example, someone asking you questions about your health or examining you. Before you left the facility, did anyone check on your health?	YES . . . . . 1 NO 2	→ 451
449	How long after delivery did the first check take place? IF LESS THAN ONE DAY, RECORD HOURS; IF LESS THAN ONE WEEK, RECORD DAYS.	HOURS . . . . . 1 ☐☐ DAYS . . . . . 2 ☐☐ WEEKS . . . . . 3 ☐☐ DON'T KNOW 998	
450	Who checked on your health at that time? PROBE FOR MOST QUALIFIED PERSON.	**HEALTH PERSONNEL** DOCTOR . . . . . 11 NURSE/MIDWIFE/CLINICAL OFFICER . . . . 12 **OTHER PERSON** TRADITIONAL BIRTH ATTENDANT . . . . . 21 COMMUNITY HEALTH WORKER/ FIELD WORKER . . . . . 22 OTHER _____ 96 (SPECIFY)	
451	CHECK 405: PREGNANCY OUTCOME TYPE MOST RECENT LIVE BIRTH MOST RECENT STILLBIRTH		→ 455
452	Now I would like to talk to you about checks on {NAME IN 407}'s health -- for example, someone examining {NAME IN 407}, checking the cord, or talking to you about how to care for {NAME IN 407}. Before {NAME IN 407} left the facility, did anyone check on {NAME IN 407}'s health?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	→ 455
453	How long after delivery was {NAME IN 407}'s health first checked? IF LESS THAN ONE DAY, RECORD HOURS; IF LESS THAN ONE WEEK, RECORD DAYS.	HOURS . . . . . 1 ☐☐ DAYS . . . . . 2 ☐☐ WEEKS . . . . . 3 ☐☐ DON'T KNOW 998	

SECTION 4. PREGNANCY AND POSTNATAL CARE

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
454	Who checked on {NAME IN 407}'s health at that time? PROBE FOR MOST QUALIFIED PERSON.	**HEALTH PERSONNEL** DOCTOR . . . . . 11 NURSE/MIDWIFE/CLINICAL OFFICER . . . . 12 **OTHER PERSON** TRADITIONAL BIRTH ATTENDANT . . . . . 21 COMMUNITY HEALTH WORKER/ FIELD WORKER . . . . . 22 OTHER _____ 96 (SPECIFY)	
455	Now I would like to talk to you about what happened after you left the facility. Did anyone check on your health after you left the facility?	YES . . . . . 1 NO 2	→ 459
456	How long after delivery did that check take place? IF LESS THAN ONE DAY, RECORD HOURS; IF LESS THAN ONE WEEK, RECORD DAYS.	HOURS . . . . . 1 ☐☐ DAYS . . . . . 2 ☐☐ WEEKS . . . . . 3 ☐☐ DON'T KNOW 998	
457	Who checked on your health at that time? PROBE FOR MOST QUALIFIED PERSON.	**HEALTH PERSONNEL** DOCTOR . . . . . 11 NURSE/MIDWIFE/CLINICAL OFFICER . . . . 12 **OTHER PERSON** TRADITIONAL BIRTH ATTENDANT . . . . . 21 COMMUNITY HEALTH WORKER/ FIELD WORKER . . . . . 22 OTHER _____ 96 (SPECIFY)	
458	Where did the check take place? PROBE TO IDENTIFY THE TYPE OF SOURCE. IF UNABLE TO DETERMINE IF PUBLIC, PRIVATE, OR NGO SECTOR, RECORD '96' AND WRITE THE NAME OF THE PLACE.	**HOME** HER HOME . . . . . 11 OTHER HOME . . . . . 12 **PUBLIC SECTOR** GOVERNMENT HOSPITAL . . . . . 21 GOVERNMENT HEALTH CENTER . . . . . 22 GOVERNMENT DISPENSARY . . . . . 23 OTHER PUBLIC SECTOR 26 _____ (SPECIFY) **PRIVATE MEDICAL SECTOR** PRIVATE HOSPITAL . . . . . 31 PRIVATE CLINIC . . . . . 32 OTHER PRIVATE MEDICAL SECTOR 36 _____ (SPECIFY) **NGO MEDICAL SECTOR** NGO HOSPITAl . . . . . 41 NGO CLINIC . . . . . 42 OTHER NGO MEDICAL SECTOR 46 _____ (SPECIFY) **FBO MEDICAL SECTOR** FBO/MISSION HOSPITAL . . . . . 51 FBO/MISSION CLINIC . . . . . 52 OTHER FBO MEDICAL SECTOR 56 _____ (SPECIFY) OTHER _____ 96 (SPECIFY)	

SECTION 4. PREGNANCY AND POSTNATAL CARE

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
459	CHECK 405: PREGNANCY OUTCOME TYPE MOST RECENT LIVE BIRTH ↓ MOST RECENT STILLBIRTH → 474		
460	After {NAME IN 407} left the {FACILITY IN 435} did any health care provider or a traditional birth attendant check on {NAME IN 407}'s health?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	→ 473
461	How long after the birth of {NAME IN 407} did that check take place? IF LESS THAN ONE DAY, RECORD HOURS; IF LESS THAN ONE WEEK, RECORD DAYS.	HOURS . . . . . 1 DAYS . . . . . 2 WEEKS . . . . . 3 DON'T KNOW 998	
462	Who checked on {NAME IN 407}'s health at that time? PROBE FOR MOST QUALIFIED PERSON.	**HEALTH PERSONNEL** DOCTOR . . . . . 11 NURSE/MIDWIFE/CLINICAL OFFICER . . . . 12 **OTHER PERSON** TRADITIONAL BIRTH ATTENDANT . . . . . 21 COMMUNITY HEALTH WORKER/ FIELD WORKER . . . . . 22 OTHER _____ (SPECIFY) 96	
463	Where did this check of {NAME IN 407} take place? PROBE TO IDENTIFY THE TYPE OF SOURCE. IF UNABLE TO DETERMINE IF PUBLIC, PRIVATE, OR NGO SECTOR, RECORD '96' AND WRITE THE NAME OF THE PLACE.	**HOME** HER HOME . . . . . 11 OTHER HOME . . . . . 12 **PUBLIC SECTOR** GOVERNMENT HOSPITAL . . . . . 21 GOVERNMENT HEALTH CENTER . . . . . 22 GOVERNMENT DISPENSARY . . . . . 23 OTHER PUBLIC SECTOR 26 _____ (SPECIFY) **PRIVATE MEDICAL SECTOR** PRIVATE HOSPITAL . . . . . 31 PRIVATE CLINIC . . . . . 32 OTHER PRIVATE MEDICAL SECTOR 36 _____ (SPECIFY) **NGO MEDICAL SECTOR** NGO HOSPITAl . . . . . 41 NGO CLINIC . . . . . 42 OTHER NGO MEDICAL SECTOR 46 _____ (SPECIFY) **FBO MEDICAL SECTOR** FBO/MISSION HOSPITAL . . . . . 51 FBO/MISSION CLINIC . . . . . 52 OTHER FBO MEDICAL SECTOR 56 _____ (SPECIFY) OTHER _____ (SPECIFY) 96	→ 473

SECTION 4. PREGNANCY AND POSTNATAL CARE

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
464	CHECK 405: PREGNANCY TYPE 1 → a) I would like to talk to you about checks on your health after delivery, for example, someone asking you questions about your health or examining you. Did anyone check on your health after you gave birth to {NAME IN 407}? PREGNANCY TYPE 3 → b) I would like to talk to you about checks on your health after delivery, for example, someone asking you questions about your health or examining you. Did anyone check on your health after you delivered the stillbirth you had in {DATE FROM 406}?	YES . . . . . 1 NO 2	→ 468
465	How long after delivery did the first check take place? IF LESS THAN ONE DAY, RECORD HOURS; IF LESS THAN ONE WEEK, RECORD DAYS.	HOURS . . . . . 1 ☐☐ DAYS . . . . . 2 ☐☐ WEEKS . . . . . 3 ☐☐ DON'T KNOW 998	
466	Who checked on your health at that time? PROBE FOR MOST QUALIFIED PERSON.	**HEALTH PERSONNEL** DOCTOR . . . . . 11 NURSE/MIDWIFE/CLINICAL OFFICER . . . . 12 **OTHER PERSON** TRADITIONAL BIRTH ATTENDANT . . . . . 21 COMMUNITY HEALTH WORKER/ FIELD WORKER . . . . . 22 OTHER _____ 96 (SPECIFY)	

SECTION 4. PREGNANCY AND POSTNATAL CARE

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
467	Where did this first check take place? PROBE TO IDENTIFY THE TYPE OF SOURCE. IF UNABLE TO DETERMINE IF PUBLIC, PRIVATE, OR NGO SECTOR, RECORD '96' AND WRITE THE NAME OF THE PLACE.	**HOME** HER HOME . . . . . 11 OTHER HOME . . . . . 12 **PUBLIC SECTOR** GOVERNMENT HOSPITAL . . . . . 21 GOVERNMENT HEALTH CENTER . . . . . 22 GOVERNMENT DISPENSARY . . . . . 23 OTHER PUBLIC SECTOR 26 _____ (SPECIFY) **PRIVATE MEDICAL SECTOR** PRIVATE HOSPITAL . . . . . 31 PRIVATE CLINIC . . . . . 32 OTHER PRIVATE MEDICAL SECTOR 36 _____ (SPECIFY) **NGO MEDICAL SECTOR** NGO HOSPITAI . . . . . 41 NGO CLINIC . . . . . 42 OTHER NGO MEDICAL SECTOR 46 _____ (SPECIFY) **FBO MEDICAL SECTOR** FBO/MISSION HOSPITAL . . . . . 51 FBO/MISSION CLINIC . . . . . 52 OTHER FBO MEDICAL SECTOR 56 _____ (SPECIFY) OTHER _____ 96 (SPECIFY)	

SECTION 4. PREGNANCY AND POSTNATAL CARE

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
468	CHECK 405: PREGNANCY OUTCOME TYPE MOST RECENT LIVE BIRTH ☐ ↓ MOST RECENT STILLBIRTH ☐ → 474		
469	I would like to talk to you about checks on {NAME IN 407}'s health -- for example, someone examining {NAME IN 407}, checking the cord, or talking to you about how to care for {NAME IN 407}. After {NAME IN 407} was born, did any health care provider or a traditional birth attendant check on {NAME IN 407}'s health?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	→ 473
470	How long after the birth of {NAME IN 407} did that check take place? IF LESS THAN ONE DAY, RECORD HOURS; IF LESS THAN ONE WEEK, RECORD DAYS.	HOURS . . . . . 1 ☐☐ DAYS . . . . . 2 ☐☐ WEEKS . . . . . 3 ☐☐ DON'T KNOW 998	
471	Who checked on {NAME IN 407}'s health at that time? PROBE FOR MOST QUALIFIED PERSON.	**HEALTH PERSONNEL** DOCTOR . . . . . 11 NURSE/MIDWIFE/CLINICAL OFFICER . . . . 12 **OTHER PERSON** TRADITIONAL BIRTH ATTENDANT . . . . . 21 COMMUNITY HEALTH WORKER/ FIELD WORKER . . . . . 22 OTHER _____ 96 (SPECIFY)	
472	Where did this first check of {NAME IN 407} take place? PROBE TO IDENTIFY THE TYPE OF SOURCE. IF UNABLE TO DETERMINE IF PUBLIC, PRIVATE, OR NGO SECTOR, RECORD '96' AND WRITE THE NAME OF THE PLACE.	**HOME** HER HOME . . . . . 11 OTHER HOME . . . . . 12 **PUBLIC SECTOR** GOVERNMENT HOSPITAL . . . . . 21 GOVERNMENT HEALTH CENTER . . . . . 22 GOVERNMENT DISPENSARY . . . . . 23 OTHER PUBLIC SECTOR 26 _____ (SPECIFY) **PRIVATE MEDICAL SECTOR** PRIVATE HOSPITAL . . . . . 31 PRIVATE CLINIC . . . . . 32 OTHER PRIVATE MEDICAL SECTOR 36 _____ (SPECIFY) **NGO MEDICAL SECTOR** NGO HOSPITAI . . . . . 41 NGO CLINIC . . . . . 42 OTHER NGO MEDICAL SECTOR 46 _____ (SPECIFY) **FBO MEDICAL SECTOR** FBO/MISSION HOSPITAL . . . . . 51 FBO/MISSION CLINIC . . . . . 52 OTHER FBO MEDICAL SECTOR 56 _____ (SPECIFY) OTHER _____ 96 (SPECIFY)	

SECTION 4. PREGNANCY AND POSTNATAL CARE

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
473	During the first 2 days after {NAME IN 407}'s birth, did any health care provider do the following: a) Examine the cord? b) Measure {NAME IN 407}'s temperature? c) Tell you how to recognize if your baby needs immediate medical attention? d) Talk with you about breastfeeding? e) Observe {NAME IN 407} breastfeeding to see if you are doing it correctly? f) Show how you can clean the cord?	YES NO DK a) CORD . . . . . 1 2 8 b) TEMPERATURE . . . . . 1 2 8 c) MEDICAL ATTENTION . . . . . 1 2 8 d) TALK ABOUT BREASTFEEDING 1 2 8 e) OBSERVE BREASTFEEDING . . 1 2 8 f) CORD CARE 1 2 8	
474	During the first 2 days after the birth, did any healthcare provider do the following to you: a) Measure your blood pressure? b) Discuss your vaginal bleeding with you? c) Discuss family planning with you?	YES NO DK a) BLOOD PRESSURE . . . . . 1 2 8 b) BLEEDING . . . . . 1 2 8 c) FAMILY PLANNING 1 2 8	
475	CHECK 215: IS THIS PREGNANCY THE WOMAN'S LAST PREGNANCY? YES ☐ NO ☐ → 479		
476	CHECK 405: PREGNANCY TYPE 1 ☐ / PREGNANCY TYPE 3 OR 5 ☐ a) Has your menstrual period returned since the birth of {NAME IN 407}? b) Has your menstrual period returned since the pregnancy that ended in {DATE FROM 406}?	YES . . . . . 1 NO 2	
477	CHECK 232: IS RESPONDENT PREGNANT? NOT PREGNANT ☐ PREGNANT OR UNSURE ☐ → 479		
478	CHECK 405: PREGNANCY TYPE 1 ☐ / PREGNANCY TYPE 3 OR 5 ☐ a) Have you had sexual intercourse since the birth of {NAME IN 407}? b) Have you had sexual intercourse since the pregnancy that ended in {DATE FROM 406}?	YES . . . . . 1 NO 2	
479	CHECK 405: PREGNANCY OUTCOME TYPE	MOST RECENT LIVE BIRTH . . . . . 1 MOST RECENT STILLBIRTH . . . . . 3 MISCARRIAGE/ABORTION . . . . . 5 → 487	
480	Did you ever breastfeed {NAME IN 407}?	YES . . . . . 1 NO . . . . . 2 → 482	
481	CHECK 224 FOR CHILD:	LIVING ☐ → 486 DEAD ☐ → 487	
482	How long after birth did you first put {NAME IN 407} to the breast? IF LESS THAN 1 HOUR, RECORD '00' HOURS; IF LESS THAN 24 HOURS, RECORD HOURS; OTHERWISE, RECORD DAYS.	IMMEDIATELY . . . . . 000 HOURS . . . . . 1 ☐☐ DAYS . . . . . 2 ☐☐	

SECTION 4. PREGNANCY AND POSTNATAL CARE

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
483	In the first 2 days after delivery, was {NAME IN 407} given anything other than breast milk to eat or drink – anything at all like water, infant formula, or herbs?	YES . . . . . 1 NO 2	
484	CHECK 224 FOR CHILD: LIVING ☐ (↓) DEAD ☐ (→ 487)		
485	Are you still breastfeeding {NAME IN 407}?	YES . . . . . 1 NO 2	
486	Did {NAME IN 407} drink anything from a bottle with a nipple yesterday during the day or at night?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
487	CHECK 402: ANY MORE PREGNANCY OUTCOMES 0-35 MONTHS BEFORE THE SURVEY? MORE PREGNANCY OUTCOMES 0-35 MONTHS BEFORE THE SURVEY ☐ (↓) (GO TO 404 FOR THE NEXT PREGNANCY NO MORE PREGNANCY OUTCOMES 0-35 MONTHS BEFORE ☐ (→ 501)		

SECTION 5. CHILD IMMUNIZATION

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
501	CHECK 220, 224 AND 225 IN THE PREGNANCY HISTORY: ANY SURVIVING CHILDREN BORN 0-35 MONTHS BEFORE THE SURVEY? ONE OR MORE SURVIVING CHILDREN BORN 0-35 MONTHS BEFORE THE SURVEY ☐ NO SURVIVING CHILDREN BORN 0-35 MONTHS BEFORE THE ☐ → 601		
502	Now I would like to ask some questions about vaccinations received by your children born in the last 3 years. We will talk about each separately, starting with the youngest.		
503	RECORD THE NAME AND PREGNANCY HISTORY NUMBER FROM 215 AND 218 OF THE SURVIVING CHILDREN BORN 0-35 MONTHS BEFORE THE SURVEY, STARTING WITH THE LAST ONE. NAME OF CHILD _____	PREGNANCY HISTORY NUMBI . . . . ☐☐	
504	Do you have a card or other document where {NAME IN 503}'s vaccinations are written down?	YES, HAS ONLY A CARD . . . . . 1 YES, HAS ONLY ANOTHER DOCUMENT . . . . 2 YES, HAS CARD AND OTHER DOCUME . . . . 3 NO, NO CARD AND NO OTHER DOCUMEN . . 4	→ 507 → 507
505	Did you ever have a vaccination card for {NAME IN 503}?	YES . . . . . 1 NO 2	
506	CHECK 504: CODE '2' CIRCLED ☐ CODE '4' CIRCLED ☐ → 513		
507	May I see the card or other document where {NAME IN 503}'s vaccinations are written down?	YES, ONLY CARD SEEN . . . . . 1 YES, ONLY OTHER DOCUMENT SEE . . . . . 2 YES, CARD AND OTHER DOCUMENT SEEN . . 3 NO CARD AND NO OTHER DOCUMENT SE . . 4	 → 513
508	RECORD (NAME'S) DATE OF BIRTH FROM THE VACCINATION CARD OR OTHER DOCUMENT.	DAY . . . . . ☐☐ MONTH . . . . . ☐☐ YEAR . . . . . ☐☐☐☐ DATE OF BIRTH NOT ON CARI 95	

SECTION 5. CHILD IMMUNIZATION

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
	NAME OF LIVE BIRTH _____	PREGNANCY HISTORY NUMBER	
509	COPY VACCINATION DATES FROM THE CARD FOR (NAME). RECORD '44' IN 'DAY' COLUMN IF CARD SHOWS THAT A DOSE WAS GIVEN, BUT NO DATE IS RECORDED. RECORD '00' IN 'DAY' COLUMN IF CARD IS BLANK FOR THE DOSE. EXTRA INSTRUCTION: IF HEXAXIM IS RECORDED AS THE BRAND FOR ANY OPV DOSE, RECORD '00' FOR THAT DOSE AND RECORD THE DATE IN THE IPV/.HEXAXIM FIELD INSTEAD		

	DAY		MONTH		YEAR			
BCG								
ORAL POLIO VACCINE (OPV) 0 (BIRTH DOSE)								
ORAL POLIO VACCINE (OPV) 1								
ORAL POLIO VACCINE (OPV) 2								
ORAL POLIO VACCINE (OPV) 3								
INACTIVATED POLIO VACCINE (IPV)								

511 CHECK: ANY OTHER IPV/ HEXAXIM RECORDED ON CARD?

☐ YES: RECORD INACTIVATED POLIO/ HEXAXIM DOSES

☐ NO: CONTINUE TO DPT-HEPB-HIB (PENTAVALENT)

INACTIVATED POLIO VACCINE (IPV)/ HEXAXIM								
INACTIVATED POLIO VACCINE (IPV)/ HEXAXIM								
INACTIVATED POLIO VACCINE (IPV)/ HEXAXIM								
DPT-HEP.B-HIB (PENTAVALENT) 1 / HEXAXIM								
DPT-HEP.B-HIB (PENTAVALENT) 2 / HEXAXIM								
DPT-HEP.B-HIB (PENTAVALENT) 3 / HEXAXIM								
PNEUMOCOCCAL 1								
PNEUMOCOCCAL 2								
PNEUMOCOCCAL 3								
ROTAVIRUS 1								
ROTAVIRUS 2								
ROTAVIRUS 3								
MEASLES RUBELLA (MR1) (9 MONTHS)								
MEASLES RUBELLA (MR2) (18 MONTHS)								
YELLOW FEVER								
VITAMIN A (MOST RECENT)								
DEWORMER (ALBENDAZOLE) (MOST RECENT)								

510	ASK THE RESPONDENT FOR PERMISSION TO PHOTOGRAPH VACCINATION CARD OR OTHER DOCUMENT WHERE VACCINATIONS ARE WRITTEN. IF PERMISSION IS GRANTED, PHOTOGRAPH CARD.	PHOTOGRAPH TAKE . . . . . 1 PHOTOGRAPH NOT TAKEN, PERMISSION NOT RECEIVED . . . . . 2 PHOTOGRAPH NOT TAKEN, OTHER REAS . . 6 _____ (SPECIFY)	
511	CHECK 509: 'BCG' TO 'YELLOW FEVER' ALL HAVE A DATE RECORDED OR '44' RECORDED IN THE 'DAY' COLUMN? NO ☐ YES ☐ → 529		

SECTION 5. CHILD IMMUNIZATION

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
	NAME OF LIVE BIRTH _____	PREGNANCY HISTORY NUMBER . . . . [ ][ ]	
512	In addition to what is recorded on (this document/these documents), did {NAME IN 503} receive any other vaccinations, including vaccinations received in campaigns or immunization days or child health days? RECORD 'YES' ONLY IF THE RESPONDENT MENTIONS AT LEAST ONE OF THE VACCINATIONS IN 509 THAT ARE NOT RECORDED AS HAVING BEEN GIVEN.	YES . . . . . 1 (USE THE LIST SHOWN IN CAPI TO SELECT THE OTHER VACCINATIONS GIVEN. NOTE THAT CAPI WILL CHANGE THE ANSWER IN 509 IN THE 'DAY' COLUMN FROM '00' TO '66' FOR THE SELECTED (THEN SKIP TO 529) NO . . . . . 2 DON'T KNOW 8	
512A	CHECK 509: ANY VACCINATIONS RECORDED ON THE CARD? YES [ ] → SKIP TO 529 NO [ ]		→ 530
513	Did {NAME IN 503} ever receive any vaccinations to prevent {NAME IN 503} from getting diseases, including vaccinations received in campaigns or immunization days or child health days?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	→ 530
514	Has {NAME IN 503} ever received a BCG vaccination against tuberculosis, that is, an injection in the arm or shoulder that usually causes a scar?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
517	Has {NAME IN 503} ever received oral polio vaccine, that is, about two drops in the mouth to prevent polio?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	→ 521
518	Did {NAME IN 503} receive the first oral polio vaccine in the first 2 weeks after birth or later?	FIRST TWO WEEKS . . . . . 1 LATER 2	
519	How many times did {NAME IN 503} receive the oral polio vaccine?	NUMBER OF TIMES . . . . . [ ]	
520	The last time {NAME IN 503} received the polio drops, did {NAME IN 503} also get an IPV injection in the right thigh to protect against polio?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
521	Has {NAME IN 503} ever received a pentavalent vaccination, that is, an injection given in the left thigh sometimes at the same time as polio drops?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	→ 523
522	How many times did {NAME IN 503} receive the pentavalent vaccine?	NUMBER OF TIMES . . . . . [ ]	
523	Has {NAME IN 503} ever received a pneumococcal vaccination, that is, an injection in the right thigh to prevent pneumonia?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	→ 525

SECTION 5. CHILD IMMUNIZATION

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
	NAME OF LIVE BIRTH _____	PREGNANCY HISTORY NUMBER	
524	How many times did {NAME IN 503} receive the pneumococcal vaccine?	NUMBER OF TIMES	
525	Has {NAME IN 503} ever received a rotavirus vaccination, that is, liquid in the mouth to prevent diarrhea?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	→ 527
526	How many times did {NAME IN 503} receive the rotavirus vaccine?	NUMBER OF TIMES	
527	Has {NAME IN 503} ever received a measles vaccination, that is, an injection in the arm to prevent measles and rubella?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	→ 528A
528	How many times did {NAME IN 503} receive the measles and rubella vaccine?	NUMBER OF TIMES	
528A	Has {NAME IN 503} ever received a yellow fever vaccination, that is, an injection in the left arm to prevent yellow fever?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
529	Where did {NAME IN 503} receive most of his/her vaccinations? PROBE TO IDENTIFY THE TYPE OF SOURCE. IF UNABLE TO DETERMINE IF PUBLIC, PRIVATE, OR NGO SECTOR, RECORD '96' AND WRITE THE NAME OF THE PLACE.	**PUBLIC SECTOR** GOVERNMENT HOSPITAL . . . . . 11 GOVERNMENT HEALTH CENTER . . . . . 12 GOVERNMENT DISPENSARY . . . . . 13 MOBILE CLINIC . . . . . 14 COMMUNITY HEALTH WORKER/ FIELDWORKER . . . . . 15 OTHER PUBLIC SECTOR _____ 16 (SPECIFY) **PRIVATE MEDICAL SECTOR** PRIVATE HOSPITAL . . . . . 21 PRIVATE CLINIC . . . . . 22 PHARMACY . . . . . 23 PRIVATE DOCTOR . . . . . 24 MOBILE CLINIC . . . . . 25 COMMUNITY HEALTH WORKER/ FIELDWORKER . . . . . 26 OTHER PRIVATE MEDICAL SECTOR _____ 27 (SPECIFY) **NGO MEDICAL SECTOR** NGO HOSPITAL . . . . . 31 NGO CLINIC . . . . . 32 OTHER NGO MEDICAL SECTOR _____ 36 (SPECIFY) **NGO MEDICAL SECTOR** FBO/MISSION HOSPITAL . . . . . 41 FBO/MISSION CLINIC . . . . . 42 FBO MOBILE CLINIC . . . . . 43 _____ 46 (SPECIFY) **OTHER SOURCE** VACCINATION CAMPAIGN . . . . . 51 OTHER _____ 96 (SPECIFY)	
530	CHECK 220 AND 224 IN PREGNANCY HISTORY: ANY MORE SURVIVING CHILDREN BORN 0-35 MONTHS BEFORE THE SURVEY? MORE SURVIVING CHILDREN BORN 0-35 MONTHS BEFORE THE SURVEY	NO MORE SURVIVING CHILDREN BORN 0-35 MONTHS BEFORE THE	→ 601

SECTION 5. CHILD IMMUNIZATION

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
	NAME OF LIVE BIRTHS _____	PREGNANCY HISTORY NUMBER. . . ☐☐	
	(GO TO 503 FOR THE NEXT SURVIVING CHILD) ←	SURVEY	

SECTION 6. CHILD HEALTH AND NUTRITION

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
601	CHECK 220, 224, AND 225 IN THE PREGNANCY HISTORY: ANY SURVIVING CHILDREN BORN 0-59 MONTHS BEFORE THE SURVEY? ONE OR MORE SURVIVING CHILDREN BORN 0-59 MONTHS BEFORE THE ☐ NO SURVIVING CHILDREN BORN 0-59 MONTHS BEFORE THE SURVEY ☐ → 643		
602	Now I would like to ask some questions about the health of your children born in the last 5 years. We will talk about each separately, starting with the youngest.		
603	RECORD THE NAME FROM 218 AND PREGNANCY HISTORY NUMBER FROM 215 OF THE SURVIVING CHILDREN BORN 0-59 MONTHS BEFORE THE SURVEY, STARTING WITH THE LAST ONE. NAME OF CHILD _____ PREGNANCY HISTORY NUMBER . . ☐☐		
604	In the last 12 months, was {NAME IN 603} given any of the following: a) Iron tablets or syrup? SHOW COMMON TYPES OF TABLETS/SYRUPS. b) Micronutrient powder or syrup? SHOW COMMON TYPES OF MULTIPLE MICRONUTRIENT POWDERS.	YES NO DK a) TABLETS/SYRUP . . . . . 1 2 8 b) MULTIPLE MICRONUTRIENT POWDERS 1 2 8	
605	In the last 6 months, was {NAME IN 603} given a vitamin A dose like [this/any of these]? SHOW COMMON TYPES OF AMPULES/CAPSULES/SYRUPS.	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
606	In the last 6 months, was {NAME IN 603} given any medicine for intestinal worms?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
607	In the last 3 months, has any healthcare provider or community health worker measured: a) {NAME IN 603}'s weight? b) {NAME IN 603}'s length or height? c) Around {NAME IN 603}'s upper arm? SHOW IMAGE OF MUAC TAPE.	YES NO DK a) WEIGHT . . . . . 1 2 8 b) LENGTH/HEIGHT . . . . . 1 2 8 c) UPPER ARM 1 2 8	
608	Has {NAME IN 603} had diarrhea in the last 2 weeks?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	→ 618

SECTION 6. CHILD HEALTH AND NUTRITION

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
NO.	NAME OF LIVE BIRTH _____	PREGNANCY HISTORY NUMBE . . . . ☐☐	
609	CHECK 485: CURRENTLY BREASTFEEDING? YES ☐↓ — NO/NOT ☐↓ a) Now I would like to know how much {NAME IN 603} was given to drink during the diarrhea, including breast milk. Was {NAME IN 603} given less than usual to drink, about the same amount, or more than usual to drink? IF LESS, PROBE: Was {NAME IN 603} given much less than usual to drink or b) Now I would like to know how much {NAME IN 603} was given to drink during the diarrhea. Was {NAME IN 603} given less than usual to drink, about the same amount, or more than usual to drink? IF LESS, PROBE: Was {NAME IN 603} given much less than usual to drink or somewhat less?	MUCH LESS . . . . . 1 SOMEWHAT LESS . . . . . 2 ABOUT THE SAME . . . . . 3 MORE . . . . . 4 NOTHING TO DRINK . . . . . 5 DON'T KNOW 8	
610	When {NAME IN 603} had diarrhea, was {NAME IN 603} given less than usual to eat, about the same amount, more than usual, or nothing to eat? IF LESS, PROBE: Was {NAME IN 603} given much less than usual to eat or somewhat less?	MUCH LESS . . . . . 1 SOMEWHAT LESS . . . . . 2 ABOUT THE SAME . . . . . 3 MORE . . . . . 4 STOPPED FOOD . . . . . 5 NEVER GAVE FOOD . . . . . 6 DON'T KNOW 8	
611	Did you seek advice or treatment for the diarrhea from any source?	YES . . . . . 1 NO 2	→ 615

SECTION 6. CHILD HEALTH AND NUTRITION

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
NO.	NAME OF LIVE BIRTH _____	PREGNANCY HISTORY NUMBE . . . . [ ][ ]	
612	Where did you seek advice or treatment? Anywhere else? PROBE TO IDENTIFY EACH TYPE OF SOURCE. IF UNABLE TO DETERMINE IF PUBLIC, PRIVATE, OR NGO SECTOR, RECORD 'X' AND WRITE THE NAME OF THE PLACE(S).	**PUBLIC SECTOR** GOVERNMENT HOSPITAL . . . . . A GOVERNMENT HEALTH CENTER . . . . . B GOVERNMENT DISPENSARY . . . . . C MOBILE CLINIC . . . . . D COMMUNITY HEALTH WORKER/ FIELDWORKER . . . . . E OTHER PUBLIC SECTOR _____ (SPECIFY) F **PRIVATE MEDICAL SECTOR** PRIVATE HOSPITAL . . . . . G PRIVATE CLINIC . . . . . H PHARMACY . . . . . I PRIVATE DOCTOR . . . . . J MOBILE CLINIC . . . . . K COMMUNITY HEALTH WORKER/ FIELDWORKER . . . . . L OTHER PRIVATE MEDICAL SECTOR _____ (SPECIFY) M **NGO MEDICAL SECTOR** NGO HOSPITAL . . . . . N NGO CLINIC . . . . . O OTHER NGO MEDICAL SECTOR _____ (SPECIFY) P **FBO MEDICAL SECTOR** FBO/MISSION HOSPITAL . . . . . Q FBO/MISSION CLINIC . . . . . R FBO MOBILE CLINIC . . . . . S OTHER FBO MEDICAL SECTOR _____ (SPECIFY) T **OTHER SOURCE** SHOP . . . . . U TRADITIONAL PRACTITIONER . . . . . V MARKET . . . . . W ITINERANT DRUG SELLER . . . . . X OTHER _____ (SPECIFY) Y	
613	CHECK 612: TWO OR MORE CODES CIRCLED ONLY ONE CODE CIRCLED → 615		
614	Where did you first seek advice or treatment? USE LETTER CODE FROM 612.	FIRST PLACE . . . . . [ ]	
615	Was {NAME IN 603} given any of the following at any time since {NAME IN 603} started having the	YES NO DK	
	a) A fluid made from a special packet called ORS?	a) FLUID FROM ORS PACKET . . 1 2 8	
	b) Zinc tablets or syrup?	b) ZINC 1 2 8	
	c) A homemade sugar-salt solution?	c) HOMEMADE	

SECTION 6. CHILD HEALTH AND NUTRITION

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES				SKIP
NO.	NAME OF LIVE BIRTH _____	PREGNANCY HISTORY NUMBE				
		SUGAR-SALT SOLUTION	1	2	8	
	d) Other homemade fluids such as porridge, soup, yoghurt, coconut water, fresh fruit juice, tea, milk, or rice water?	d) OTHER HOMEMADE FLUID	1	2	8	

SECTION 6. CHILD HEALTH AND NUTRITION

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
NO.	NAME OF LIVE BIRTH _____	PREGNANCY HISTORY NUMBE . . . . ☐☐	
616	CHECK 615: ANY 'YES' ☐ → a) Was anything else given to treat the diarrhea? ALL 'NO' OR 'DK' ☐ → b) Was anything given to treat the diarrhea?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	→ 618
617	CHECK 615: ANY 'YES' ☐ → a) What else was given to treat the diarrhea? Anything else? RECORD ALL TREATMENTS GIVEN. ALL 'NO' OR 'DK' ☐ → b) What was given to treat the diarrhea? Anything else? RECORD ALL TREATMENTS GIVEN.	**PILL OR SYRUP** ANTIBIOTIC . . . . . A ANTIMOTILITY . . . . . B OTHER (NOT ANTIBIOTIC OR ANTIMOTILIT\ C UNKNOWN PILL OR SYRUP . . . . . D THERAPEUTIC VITAMIN A . . . . . E **INJECTION** ANTIBIOTIC . . . . . F NON-ANTIBIOTIC . . . . . G UNKNOWN INJECTION . . . . . H (IV) INTRAVENOUS . . . . . I HOME REMEDY/HERBAL MEDICIN . . . . . J OTHER _____ X (SPECIFY)	
618	Has {NAME IN 603} been ill with a fever at any time in the last 2 weeks?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	→ 621
619	At any time during the illness, did {NAME IN 603} have blood taken from {NAME IN 603}'s finger or heel for testing?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
620	Were you told by a healthcare provider that {NAME IN 603} had malaria?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
621	Has {NAME IN 603} had an illness with a cough at any time in the last 2 weeks?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
622	Has {NAME IN 603} had fast, short, rapid breaths or difficulty breathing at any time in the last 2 weeks?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	→ 624
623	Was the fast or difficult breathing due to a problem in the chest or to a blocked or runny nose?	CHEST ONLY . . . . . 1 NOSE ONLY . . . . . 2 BOTH . . . . . 3 OTHER _____ 6 (SPECIFY) DON'T KNOW 8	→ 625
624	CHECK 618: HAD FEVER? YES ☐	NO OR DON'T KNOW ☐	→ 634
625	Did you seek advice or treatment for the illness from any source?	YES . . . . . 1 NO 2	→ 630

SECTION 6. CHILD HEALTH AND NUTRITION

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
NO.	NAME OF LIVE BIRTH _____	PREGNANCY HISTORY NUMBE . . . . ☐☐	
626	Where did you seek advice or treatment? Anywhere else? PROBE TO IDENTIFY EACH TYPE OF SOURCE. IF UNABLE TO DETERMINE IF PUBLIC, PRIVATE, OR NGO SECTOR, RECORD 'X' AND WRITE THE NAME OF THE PLACE(S).	**PUBLIC SECTOR** GOVERNMENT HOSPITAL . . . . . A GOVERNMENT HEALTH CENTER . . . . . B GOVERNMENT DISPENSARY . . . . . C MOBILE CLINIC . . . . . D COMMUNITY HEALTH WORKER/ FIELDWORKER . . . . . E OTHER PUBLIC SECTOR F SECTOR _____ (SPECIFY) **PRIVATE MEDICAL SECTOR** PRIVATE HOSPITAL . . . . . G PRIVATE CLINIC . . . . . H PHARMACY . . . . . I PRIVATE DOCTOR . . . . . J MOBILE CLINIC . . . . . K COMMUNITY HEALTH WORKER/ FIELDWORKER . . . . . L OTHER PRIVATE MEDICAL SECTOR M _____ (SPECIFY) **NGO MEDICAL SECTOR** NGO HOSPITAL . . . . . N NGO CLINIC . . . . . O OTHER NGO MEDICAL SECTOR P _____ (SPECIFY) **FBO MEDICAL SECTOR** FBO/MISSION HOSPITAL . . . . . Q FBO/MISSION CLINIC . . . . . R FBO MOBILE CLINIC . . . . . S OTHER FBO MEDICAL SECTOR _____ T (SPECIFY) **OTHER SOURCE** SHOP . . . . . U TRADITIONAL PRACTITIONER . . . . . V MARKET . . . . . W ITINERANT DRUG SELLER . . . . . X OTHER _____ Y (SPECIFY)	
627	CHECK 626: TWO OR MORE CODES CIRCLED ☐↓ ONLY ONE CODE CIRCLED ☐ → 629		
628	Where did you first seek advice or treatment? USE LETTER CODE FROM 626.	FIRST PLACE . . . . . ☐	
629	How many days after the illness began did you first seek advice or treatment for {NAME IN 603}? IF SAME DAY, RECORD '00'.	DAYS . . . . . ☐☐	
630	At any time during the illness, did {NAME IN 603} take any medicine for the illness?	YES . . . . . 1 NO . . . . . 2 → 634 DON'T KNOW 8	

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
NO.	NAME OF LIVE BIRTH _____	PREGNANCY HISTORY NUMBE . . . . ☐☐	
631	What medicine did {NAME IN 603} take? Any other medicine? RECORD ALL MENTIONED. IF MEDICINE NOT KNOWN, ASK TO SEE THE PACKAGE OR PRESCRIPTION	**ANTIMALARIAL MEDICINE** ACT - AL . . . . . A ACT- DHAP . . . . . B OTHER ACT (NOT AL OR DHAP) . . . . . C **NON-ACT ANTIMALARIAL** SP/FANSIDAR . . . . . D CHLOROQUINE . . . . . E AMODIAQUINE . . . . . F QUININE PILLS . . . . . G INJECTION/IV . . . . . H ARTESUNATE RECTAL . . . . . I INJECTION/IV . . . . . J OTHER ANTIMALARIAL K _____ (SPECIFY) **ANTIBIOTIC MEDICINE** AMOXICILLIN DT TABLETS . . . . . L AMOXICILLIN SYRUP . . . . . M COTRIMOXAZOLE . . . . . N OTHER PILL/SYRUP . . . . . O OTHER INJECTION/IV . . . . . P **OTHER MEDICINE** ASPIRIN . . . . . Q PARACETAMOL/PANADOL/ACETAMINOPHE R IBUPROFEN . . . . . S OXYGEN . . . . . T OTHER _____ X (SPECIFY) DON'T KNOW . . . . . Z	
632	CHECK 631: ARTEMISININ COMBINATION THERAPY ('A') GIVEN CODE 'A', 'B', OR 'C' CIRCLED ☐ ↓ CODE 'A', 'B', OR 'C' NOT CIRCLED ☐ → 634		
633	How long after the fever started did {NAME IN 603} first take an artemisinin combination therapy?	SAME DAY . . . . . 0 NEXT DAY . . . . . 1 TWO DAYS AFTER FEVER . . . . . 2 THREE OR MORE DAYS AFTER FEVE . . . . . 3 DON'T KNOW 8	
634	CHECK 220, 224, AND 225 IN PREGNANCY HISTORY: ANY MORE SURVIVING CHILDREN BORN 0-59 MONTHS BEFORE THE SURVEY? MORE SURVIVING CHILDREN BORN 0-59 MONTHS BEFORE THE SURVEY ☐ → (GO TO 603 FOR THE NEXT SURVIVING CHILD) NO MORE SURVIVING CHILDREN BORN 0-59 MONTHS BEFORE ☐ → 635		

SECTION 6. CHILD HEALTH AND NUTRITION

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
635	CHECK 220, 225 AND 226, ALL ROWS: NUMBER OF CHILDREN BORN 0-23 MONTHS BEFORE THE SURVEY LIVING WITH THE RESPONDENT ONE OR MORE ☐ ↓ (NAME OF YOUNGEST CHILD LIVING WITH HER) ↓	NONE ☐ →	→ 643
636	Now I would like to ask you about liquids that {NAME IN 635} had yesterday during the day or at night. Please tell me about all drinks, whether {NAME IN 635} had them at home, or somewhere else. Yesterday during the day or at night, did {NAME IN 635} drink:	YES NO DK	
	a) Plain water?	a) 1 2 8	
	b) Infant formula such as NAN (Nestle), SMA, or S26? IF YES: b1) How many times did {NAME IN 635} drink infant formula? IF 7 OR MORE TIMES, RECORD '7'.	b) . . . . . 1 2 8 NUMBER OF TIMES DRANK ☐ 8	
	c) Milk from animals, such as fresh milk, powdered milk, or milk tea? IF YES: c1) How many times did {NAME IN 635} drink milk? IF 7 OR MORE TIMES, RECORD '7'. c2) Was the milk a sweet or flavored type of milk?	c) . . . . . 1 2 8 NUMBER OF TIMES DRANK ☐ 8 SWEET/ FLAVORED . . 1 2 8	
	f) Milo or cocoa?	f) 1 2 8	
	g) Fruit juice or fruit drinks?	g) 1 2 8	
	h) Soft drinks such as Coca-Cola, Fanta, Sprite, or energy drinks such as Red Bull?	h) 1 2 8	
	i) Tea, coffee, or herbal drinks? IF YES: i1) Was the drink sweetened?	i) . . . . . 1 2 8 SWEETENED 1 2 8	
	j) Clear broth or clear soup?	j) 1 2 8	
		YES NO DK	
	k) Any other liquids? IF YES: k1) What was the drink? MARK THE APPROPRIATE GROUP FOR EACH ADDITIONAL DRINK, IF THE GROUP IS NOT YET CODED 'YES'. IF UNABLE TO DETERMINE WHICH GROUP THE ADDITIONAL DRINK BELONGS TO, SELECT OPTION "Z" AND A SCREEN WILL BE DISPLAYED TO REGISTER THE NAME OF k2) Was the drink sweetened?	k) . . . . . 1 2 8 OTHER DRINK(S) _____ (SPECIFY) SWEETENED 1 2 8	

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
637	Now I would like to ask you about foods that {NAME IN 635} had yesterday during the day or at night. I am interested in foods your child ate whether at home or somewhere else. Please think about snacks and small meals as well as main meals. I will ask you about different foods, and I would like to know whether your child ate the food even if it was combined with other foods. Please do not answer 'yes' for any food or ingredient only used in a small amount to add flavor to a dish. Yesterday during the day or at night, did {NAME IN 635} have:	YES NO DK	
	a) Yogurt or mala?	a) 1 2 8	
	IF YES: a1) How many times did {NAME IN 635} eat yogurt or mala? IF 7 OR MORE TIMES, RECORD '7'.	NUMBER OF TIMES ATE ☐ 8	
	a2) Did {NAME IN 635} have any yogurt or mala as a drink?	HAD YOGURT AS A DRINK . . 1 2 8	
	IF YES: a3) Was the yogurt drink a sweet or flavored type of	SWEETENED 1 2 8	
	b) Ugali, porridge, rice, bread, chapati, pasta, or green maize?	b) 1 2 8	
	c) Carrots, pumpkin, butternut, or sweet potato that are yellow or orange inside?	c) 1 2 8	
	d) Irish potato, white sweet potato, green banana, nduma (arrowroot), yam, or cassava?	d) 1 2 8	
		YES NO DK	
	e) Sukuma wiki, spinach, managu (nightshade), terere (amaranth), saget, or kunde (cowpea leaves)?	e) 1 2 8	
	e1) Khandira (Ethiopian kale), mrenda (jute mallow), pumpkin leaves, nderema (Malabar spinach), mitoo, broccoli, or mchunga?	h) 1 2 8	
	f) Any other vegetables, such as tomatoes, cabbage, green capsicum, mushrooms, cucumber or other vegetables?	f) 1 2 8	
	g) Ripe pawpaw, ripe mango, passionfruit, or matunda ya damu?	g) 1 2 8	
	h) Any other fruits, such as banana, pineapple, avocado, watermelon, orange or other fruits?	h) 1 2 8	
	i) Fish, dagaa, canned tuna, or seafood?	i) 1 2 8	
	j) Liver, blood, kidney, lung, gizzard, or heart?	j) 1 2 8	

SECTION 6. CHILD HEALTH AND NUTRITION

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
	k) Sausages, Smokies, hot dogs, salami, or ham?	k) 1 2 8	
	l) Any other meat, such as goat, beef, minced beef, mutton, pork, wild game, or chicken?	l) 1 2 8	

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
	m) Eggs?	m) 1 2 8	
	n) Beans, githeri, ndengu (green gram), njahi (black gram), kamande (lentils), pigeon peas, or chickpeas?	n) 1 2 8	
	o) Groundnuts, cashews, pumpkin seeds, simsim (sesame seeds), or peanut butter?	o) 1 2 8	
	p) Cheese?	p) 1 2 8	
	q) Termites, locusts, or grasshoppers?	q) 1 2 8	
	r) Any sweet foods such as cakes, sweet biscuits, candy, chocolates, ice cream, or ice lollies?	r) 1 2 8	
		YES NO DK	
	t) Crisps, chips, ngumu, mandaazi, samosa, bhajias, or Indomie?	t) 1 2 8	
	v) Any other solid, semi-solid, or soft food? IF YES: v1) What was the food? MARK THE APPROPRIATE FOOD GROUP FOR EACH ADDITIONAL FOOD, IF THE GROUP IS NOT YET CODED 'YES'. IF UNABLE TO DETERMINE WHICH GROUP THE ADDITIONAL FOOD BELONGS TO, SELECT OPTION "Z" AND A SCREEN WILL BE DISPLAYED TO REGISTER THE NAME OF	v) . . . . . 1 2 8 OTHER FOOD(S) _____ (SPECIFY)	
638	CHECK 637 (CATEGORIES 'a' THROUGH 'v'): NOT A SINGLE 'YES' ☐ ↓ AT LEAST ONE 'YES' ☐ → 640		
639	Did {NAME IN 635} eat any solid, semi-solid, or soft foods yesterday during the day or at night? IF 'YES' PROBE: What kind of solid, semi-solid or soft foods did {NAME IN 635} eat?	YES . . . . . 1 ☐ (GO BACK TO 637 TO RECORD FOOD EATEN YESTERDAY) ← (THEN CONTINUE TO 640) ← NO . . . . . 2 → 641	
640	How many times did {NAME IN 635} eat solid, semi-solid, or soft foods yesterday during the day or at night?	NUMBER OF TIMES . . . . . ☐☐ DON'T KNOW 8	
641	In the last 6 months, did any healthcare provider or community health worker talk with you about how or what to feed {NAME IN 635}?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
642	The last time {NAME IN 635} passed stools, what was done to dispose of the stools?	CHILD USED TOILET OR LATRINE . . . . . 01 PUT/RINSED INTO TOILET OR LATRINE . . . . 02 PUT/RINSED INTO DRAIN OR DITCH . . . . . 03 THROWN INTO GARBAG 04	

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
		BURIED . . . . . 05 LEFT IN THE OPEI . . . . . 06 OTHER _____ 96 (SPECIFY)	

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
650A	CHECK 220, 224, 225, AND 226 IN THE PREGNANCY HISTORY: CHILDREN BORN 3-59 MONTHS BEFORE THE SURVEY LIVING WITH THE RESPONDENT? ONE OR MORE ☐ NONE ☐ → 701		
650B	I would like to ask you some questions about your children age 3-59 months who live with you. These questions are about certain things that they are currently able to do. Please keep in mind that children can develop and learn at a different pace. For example, some start talking earlier than others, or they might already say some words but not yet form sentences. So, it is fine if your child is not able to do all the things I am going to ask you about. You can let me know if you have any doubts about what answer to give.		
650C	RECORD THE NAME AND PREGNANCY HISTORY NUMBER FROM 215 AND 218 OF CHILDREN BORN 3-59 MONTHS BEFORE THE SURVEY LIVING WITH THE RESPONDENT, STARTING WITH THE LAST NAME OF CHILD _____	PREGNANCY HISTORY NUMBER . . ☐☐ CHILD'S AGE . . . . . ☐☐ M F CHILD'S SEX 1 2	
650C	CHECK THE CHILD'S AGE FROM 650C: 3-5 MONTHS ☐	6-11 MONTHS ☐ → 661 12-17 MONTHS ☐ → 673 18-23 MONTHS ☐ → 679 24-59 MONTHS ☐ → ECD01	
651	CHECK 652: MALE ☐ / FEMALE ☐		
	a) When you talk to {ECD CHILD NAME}, does he smile, make noises or move arms, legs or trunk in response?	b) When you talk to {ECD CHILD NAME}, does she smile, make noises or move arms, legs or trunk in response?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8
652	a) When you are about to pick up {ECD CHILD NAME}, does he act happy or	b) When you are about to pick up {ECD CHILD NAME}, does she act happy or	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8
653	a) Does {ECD CHILD NAME} turn his/her head towards your voice or some other noise?	b) Does {ECD CHILD NAME} turn his/her head towards your voice or some other noise?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8

NO.	QUESTIONS AND FILTERS		CODING CATEGORIES	SKIP
654	a) Does {ECD CHILD NAME} sometimes suck his thumb or fingers? SHOW PICTURE.	b) Does {ECD CHILD NAME} sometimes suck her thumb or fingers? SHOW PICTURE.	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
655	a) While {ECD CHILD NAME} is on his back, can he bring his hands together such that the hands touch each other?	b) While {ECD CHILD NAME} is on her back, can she bring her hands together such that the hands touch each other?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
656	a) Does {ECD CHILD NAME} move excitedly, kick legs, move arms or trunk or make cooing noises when a known person enters the room or speaks to him?	b) Does {ECD CHILD NAME} move excitedly, kick legs, move arms or trunk or make cooing noises when a known person enters the room or speaks to her?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
657	Does {ECD CHILD NAME} make noise or gestures to get your attention?		YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
658	CHECK 652: MALE ☐ ↓	FEMALE ☐ ↓		
	a) Does {ECD CHILD NAME} recognize you or other family members, for example smilng when someone enters a room or moves	b) Does {ECD CHILD NAME} recognize you or other family members, for example smilng when someone enters a room or moves	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
659	Does {ECD CHILD NAME} laugh?		YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
660	CHECK 652: MALE ☐ ↓	FEMALE ☐ ↓		
	a) When {ECD CHILD NAME} is on his stomach, can he turn his head to the side? SHOW PICTURE.	b) When {ECD CHILD NAME} is on her stomach, can she turn hee head to the side? SHOW PICTURE.	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
661	a) Is {ECD CHILD NAME} interested when he sees other children playing? Does he watch, smile or look excited?	b) Is {ECD CHILD NAME} interested when she sees other children playing? Does she watch, smile or look excited?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
662	a) Can {ECD CHILD NAME} hold his head steady for at least a few seconds, without it flopping to the side? b) Can {ECD CHILD NAME} hold her head steady for at least a few seconds, without it flopping to the side?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
663	a) When {ECD CHILD NAME} is on his stomach, can she hold his head up off the ground? SHOW PICTURE. b) When {ECD CHILD NAME} is on her stomach, can she hold her head up off the ground? SHOW PICTURE.	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
664	a) Does {ECD CHILD NAME} show interest in new objects that are put in front of him by reaching out for them? SHOW PICTURE. b) Does {ECD CHILD NAME} show interest in new objects that are put in front of her by reaching out for them? SHOW PICTURE.	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
665	a) When {ECD CHILD NAME} is on his tummy, can he hold his head straight up, looking around for more than a few seconds? He can rest on his arms while doing this. SHOW PICTURE. b) When {ECD CHILD NAME} is on her tummy, can she hold her head straight up, looking around for more than a few seconds? She can rest on her arms while doing this. SHOW PICTURE.	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
666	Can {ECD CHILD NAME} pick up a small object, such as a small toy or small stone, using just one hand? SHOW PICTURE.	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
667	CHECK 652: MALE ☐ ↓ FEMALE ☐ ↓ a) Can {ECD CHILD NAME} sit with support, either leaning against something such as furniture or person, or by leaning forward on his hands? SHOW PICTURE. b) Can {ECD CHILD NAME} sit with support, either leaning against something such as furniture or person, or by leaning forward on her hands? SHOW PICTURE.	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
668	If an object falls to the ground out of view, does {ECD CHILD NAME} look for it? SHOW PICTURE.	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	

NO.	QUESTIONS AND FILTERS		CODING CATEGORIES	SKIP
669	CHECK 652: MALE ☐ ↓	FEMALE ☐ ↓		
	a) When lying on his stomach, can {ECD CHILD NAME} hold his head and chest off the ground using only his hands and arms for support? SHOW PICTURE.	b) When lying on her stomach, can {ECD CHILD NAME} hold her head and chest off the ground using only her hands and arms for support? SHOW PICTURE.	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
670	a) When lying on his back, does {ECD CHILD NAME} grab his feet? SHOW PICTURE.	b) When lying on her back, does {ECD CHILD NAME} grab her feet? SHOW PICTURE.	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
671C	CHECK THE CHILD'S AGE FROM 650C: 6-11 MONTHS ☐ ↓	3-5 MONTHS ☐ →		ECD21
671	a) Does {ECD CHILD NAME} look for an object of interest when it is removed from sight or hidden from her, for example when it is placed under a cover, behind	b) Does {ECD CHILD NAME} look for an object of interest when it is removed from sight or hidden from her, for example when it is placed under a cover, behind	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
672	a) Can {ECD CHILD NAME} hold himself in a sitting position without help or support for longer than a few seconds? SHOW PICTURE.	b) Can {ECD CHILD NAME} hold herself in a sitting position without help or support for longer than a few seconds? SHOW PICTURE.	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
673	a) Can {ECD CHILD NAME} pick up small bits of food and feed himself using his hand? SHOW PICTURE.	b) Can {ECD CHILD NAME} pick up small bits of food and feed herself using her hand? SHOW PICTURE.	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
674	Can {ECD CHILD NAME} pick up and drop a small object such as a piece of food, small toy or small stone, into a bucket or bowl while sitting? SHOW PICTURE.		YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
675	Can {ECD CHILD NAME} maintain a standing position while holding onto a person or object such as a wall or furniture? SHOW PICTURE.	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
676	CHECK 652: MALE ☐ ↓ FEMALE ☐ ↓ a) Can {ECD CHILD NAME} pull himself up from the floor while holding onto something? For example, can {ECD CHILD NAME} pull himself up using a chair, a person or some other object? b) Can {ECD CHILD NAME} pull herself up from the floor while holding onto something? For example, can {ECD CHILD NAME} pull herself up using a chair, a person or some other object?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
677	a) Does {ECD CHILD NAME} stop what he is doing when you say "Stop!", even if just for a second? b) Does {ECD CHILD NAME} stop what she is doing when you say "Stop!", even if just for a second?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
678	Can {ECD CHILD NAME} walk several steps while holding onto a person or object such as a wall or furniture? SHOW PICTURE.	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
679	CHECK 652: MALE ☐ ↓ FEMALE ☐ ↓ a) Does {ECD CHILD NAME} make a gesture to indicate "No", such as shaking his head? b) Does {ECD CHILD NAME} make a gesture to indicate "No", such as shaking her head?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
680	Can {ECD CHILD NAME} stand up without holding onto anything, even if just for a few seconds? SHOW PICTURE.	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
671C	CHECK THE CHILD'S AGE FROM 650C: 12-23 MONTHS ☐ ↓ 6-11 MONTHS ☐ →		ECD21
681	CHECK 652: MALE ☐ ↓ FEMALE ☐ ↓ a) Does {ECD CHILD NAME} put his hands out to have them washed? b) Does {ECD CHILD NAME} put her hands out to have them washed?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
682	a) Can {ECD CHILD NAME} maintain a standing position on his own, without holding on or receiving support? SHOW PICTURE. b) Can {ECD CHILD NAME} maintain a standing position on her own, without holding on or receiving support? SHOW PICTURE.	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
683	Can {ECD CHILD NAME} climb onto an object, such as a rock, porch, step, chair, bed, or low table? SHOW PICTURE.	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
684	Can {ECD CHILD NAME} make any light marks on paper or in dirt with a crayon or a stick? SHOW PICTURE.	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
685	Can {ECD CHILD NAME} follow a simple spoken command or direction without you making a gesture?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
686	Does {ECD CHILD NAME} share with others, such as by sharing food?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
687	CHECK 652: MALE ☐ → a) Can {ECD CHILD NAME} make a scribble on paper, or in dirt, in a back-and-forth manner? For example, can he move a pen, pencil or stick back-and-forth? FEMALE ☐ → b) Can {ECD CHILD NAME} make a scribble on paper, or in dirt, in a back-and-forth manner? For example, can she move a pen, pencil or stick back-and-forth?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
688	a) Can {ECD CHILD NAME} move around by walking, rather than by crawling on his hands and knees? b) Can {ECD CHILD NAME} move around by walking, rather than by crawling on her hands and knees?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
689	Can {ECD CHILD NAME} stack at least 2 objects on top of each other, such as bottle tops, blocks, or stones.? SHOW PICTURE.	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
690	Can {ECD CHILD NAME} say 5 or more distinct words, such as names like "Mama" or objects like "ball"?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
691	CHECK 652: MALE ☐ → FEMALE ☐ → a) Can {ECD CHILD NAME} ask for something, such as food or water, by name when he wants it? b) Can {ECD CHILD NAME} ask for something, such as food or water, by name when she wants it?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
692	Can {ECD CHILD NAME} run well, without falling or bumping into objects? SHOW PICTURE.	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
692C	CHECK THE CHILD'S AGE FROM 650C: 18-23 MONTHS ☐ → 12-17 MONTHS ☐ →		ECD21
693	CHECK 652: MALE ☐ → FEMALE ☐ → a) Can {ECD CHILD NAME} wash hands by himself? b) Can {ECD CHILD NAME} wash hands by herself?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
694	a) If you show {ECD CHILD NAME} an object he knows well, such as a cup or animal, can he consistently name it? By consistently, we mean that he uses the same word to refer to the same object, even if the word used is not fully correct. b) If you show {ECD CHILD NAME} an object she knows well, such as a cup or animal, can she consistently name it? By consistently, we mean that she uses the same word to refer to the same object, even if the word used is not fully	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
695	Can {ECD CHILD NAME} say 10 or more words in addition to "Mama" and "Dada"?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
696	Can {ECD CHILD NAME} speak using sentences of 3 or more words that go together, such as "I want water", or "The house is big"?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
697	Can {ECD CHILD NAME} jump with both feet leaving the ground? SHOW PICTURE.	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
698	Can {ECD CHILD NAME} correctly use any of the words "I", "you", "she" or "he", for example "I go to store", or "He eats rice"?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	→ ECD21

EARLY CHILDHOOD DEVELOPMENT MODULE (ECDI)

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
ECD01	Can {ECD CHILD NAME} walk on an uneven surface, for example, a bumpy or steep road, without falling?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
ECD02	Can {ECD CHILD NAME} jump up with both feet leaving the ground?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
ECD03	CHECK ECDC: MALE ☐ ↓ a) Can {ECD CHILD NAME} dress himself, that is, put on pants and a shirt, without help? FEMALE ☐ ↓ b) Can {ECD CHILD NAME} dress herself, that is, put on pants and a shirt, without help?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
ECD04	Can {ECD CHILD NAME} fasten and unfasten buttons without help?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
ECD05	Can {ECD CHILD NAME} say 10 or more words, like 'mama' or 'ball'?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
ECD06	Can {ECD CHILD NAME} speak using sentences of 3 or more words that go together, for example, "I want water" or "The house is big"?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	☐ → ECD08
ECD07	Can {ECD CHILD NAME} speak using sentences of 5 or more words that go together, for example, "The house is very big"?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
ECD08	Can {ECD CHILD NAME} correctly use any of the words 'I', 'you', 'she', or 'he', for example, "I want water" or "He eats rice"?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
ECD09	CHECK ECDC: MALE ☐ ↓ a) If you show {ECD CHILD NAME} an object he knows well, such as a cup or animal, can he consistently name it? By consistently, we mean that he uses the same word to refer to the same object, even if the word used is not fully correct. FEMALE ☐ ↓ b) If you show {ECD CHILD NAME} an object she knows well, such as a cup or animal, can she consistently name it? By consistently, we mean that she uses the same word to refer to the same object, even if the word used is not fully correct.	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
ECD10	Can {ECD CHILD NAME} recognize at least 5 letters of the alphabet?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
ECD11	CHECK ECDC: MALE ☐ → a) Can {ECD CHILD NAME} write his name? FEMALE ☐ → b) Can {ECD CHILD NAME} write her name?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
ECD12	Can {ECD CHILD NAME} recognize all numbers from 1 to 5?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
ECD13	CHECK ECDC: MALE ☐ → a) If you ask {ECD CHILD NAME} to give you 3 objects, such as 3 stones or 3 beans, does he give you the correct amount? FEMALE ☐ → b) If you ask {ECD CHILD NAME} to give you 3 objects, such as 3 stones or 3 beans, does she give you the correct amount?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
ECD14	Can {ECD CHILD NAME} count 10 objects, for example, 10 fingers or 10 blocks, without mistakes?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
ECD15	Can {ECD CHILD NAME} do an activity, such as coloring or playing with building blocks, without repeatedly asking for help or giving up too quickly?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
ECD16	Does {ECD CHILD NAME} ask about familiar people other than parents when they are not there, for example, "Where is Grandma?"?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
ECD17	Does {ECD CHILD NAME} offer to help someone who seems to need help?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
ECD18	Does {ECD CHILD NAME} get along well with other children?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
ECD19	How often does {ECD CHILD NAME} seem to be very sad or depressed? Would you say: daily, weekly, monthly, a few times a year, or never?	DAILY . . . . . 1 WEEKLY . . . . . 2 MONTHLY . . . . . 3 A FEW TIMES A YEAR . . . . . 4 NEVER . . . . . 5 DON'T KNOW 8	
ECD20	Compared with other children of the same age, how much does {ECD CHILD NAME} kick, bite, or hit other children or adults? Would you say: not at all, the same or less, more, or a lot more?	NOT AT ALL . . . . . 1 THE SAME OR LESS . . . . . 2 MORE . . . . . 3 A LOT MORE . . . . . 4 DON'T KNOW 8	
ECD21	CHECK 220, 224, 225 AND 226 IN PREGNANCY HISTORY: ANY MORE CHILDREN BORN 3-59 MONTHS BEFORE THE SURVEY LIVING WITH THE RESPONDENT? MORE CHILDREN BORN 3-59 MONTHS BEFORE THE SURVEY LIVING WITH THE RESPONDENT ☐ → (GO TO 650C FOR THE NEXT CHILD) NO MORE ☐ → 701		

SECTION 7. MARRIAGE AND SEXUAL ACTIVITY

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
701	Are you currently married or living together with a man as if married?	YES, CURRENTLY MARRIED . . . . . 1 YES, LIVING WITH A MA . . . . . 2 NO, NOT IN UNION 3	→706A
702	Have you ever been married or lived together with a man as if married?	YES, FORMERLY MARRIED . . . . . 1 YES, LIVED WITH A MAN . . . . . 2 NO 3	→721
703	What is your marital status now: are you widowed, divorced, or separated?	WIDOWED . . . . . 1 DIVORCED . . . . . 2 SEPARATED 3	→721
706A	Do you have a marriage certificate or other document recognizing this (marriage/union)?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	→707
706B	What document or documents do you have? Any other document? RECORD ALL MENTIONED.	MARRIAGE CERTIFICATE FROM A CHURCH, MOSQUE OR OTHER RELIGIOUS INSTITUTION . . . . . A MARRIAGE CERTIFICATE FROM A CIVIL AUTHORITY . . . . . B OTHER DOCUMENT FROM A RELIGIOUS INSTITUTION . . . . . C OTHER DOCUMENT FROM A CIVIL AUTHORITY . . . . . D OTHER _____ X (SPECIFY)	→721
707	Was this marriage ever registered with the civil authority?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
721	**CHECK FOR PRESENCE OF OTHERS. BEFORE CONTINUING, MAKE EVERY EFFORT TO ENSURE**		
722	Now I would like to ask some questions about sexual activity in order to gain a better understanding of some important life issues. Let me assure you again that your answers are completely confidential and will not be told to anyone. If we should come to any question that you don't want to answer, just let me know and we will go to the next question. How old were you when you had sexual intercourse for the very first time?	NEVER HAD SEXUAL INTERCOURSE . . . . . 00 AGE IN YEARS	→738
723	I would like to ask you about your recent sexual activity. When was the last time you had sexual intercourse? IF LESS THAN 12 MONTHS, ANSWER MUST BE RECORDED IN DAYS, WEEKS OR MONTHS. IF 12 MONTHS (ONE YEAR) OR MORE, ANSWER MUST BE RECORDED IN YEARS.	DAYS AGO . . . . . 1 WEEKS AGO . . . . . 2 MONTHS AGO . . . . . 3 YEARS AGO 4	→737

SECTION 7. MARRIAGE AND SEXUAL ACTIVITY

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
724	CHECK 232: NOT PREGNANT OR UNSURE ☐ PREGNANT ☐		→ 727
725	The last time you had sexual intercourse, did you or your partner do something or use any method to delay or avoid getting pregnant?	YES . . . . . 1 NO 2	→ 727
726	Which method did you use? RECORD ALL MENTIONED. IF CODES 'G' OR 'H' ARE CIRCLED, SKIP TO 728 EVEN IF ANOTHER METHOD WAS ALSO USED.	FEMALE STERILIZATION . . . . . A MALE STERILIZATION . . . . . B IUD . . . . . C INJECTABLES . . . . . D IMPLANTS . . . . . E PILL . . . . . F CONDOM . . . . . G FEMALE CONDOM . . . . . H EMERGENCY CONTRACEPTIO . . . . . I STANDARD DAYS METHOD . . . . . J LACTATIONAL AMENORRHEA METHOD . . . . K RHYTHM METHOD . . . . . L WITHDRAWAL . . . . . M OTHER MODERN METHOD . . . . . X OTHER TRADITIONAL METHOD . . . . . Y	→ 730
727	The last time you had sexual intercourse, was a condom used?	YES . . . . . 1 NO 2	→ 730
730	What was your relationship to this person with whom you had sexual intercourse? IF BOYFRIEND: Were you living together as if married? IF YES, RECORD '2'. IF NO, RECORD '3'.	HUSBAND . . . . . 1 LIVE-IN PARTNER . . . . . 2 BOYFRIEND NOT LIVING WITH RESPONDE . . 3 CASUAL ACQUAINTANCE . . . . . 4 CLIENT/SEX WORKER . . . . . 5 OTHER _____ 6 (SPECIFY)	
731	Apart from this person, have you had sexual intercourse with any other person in the last 12 months?	YES . . . . . 1 NO 2	→ 737
732 (2)	The last time you had sexual intercourse with this second person, was a condom used?	YES . . . . . 1 NO 2	
733	What was your relationship to this second person with whom you had sexual intercourse? IF BOYFRIEND: Were you living together as if married? IF YES, RECORD '2'. IF NO, RECORD '3'.	HUSBAND . . . . . 1 LIVE-IN PARTNER . . . . . 2 BOYFRIEND NOT LIVING WITH RESPONDENT 3 CASUAL ACQUAINTANCE . . . . . 4 CLIENT/SEX WORKER . . . . . 5 OTHER _____ 6 (SPECIFY)	

SECTION 7. MARRIAGE AND SEXUAL ACTIVITY

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
734	Apart from these two people, have you had sexual intercourse with any other person in the last 12 months?	YES . . . . . 1 NO 2	→ 737
735 (2)	The last time you had sexual intercourse with this third person, was a condom used?	YES . . . . . 1 NO 2	
736	What was your relationship to this third person with whom you had sexual intercourse? IF BOYFRIEND: Were you living together as if married? IF YES, RECORD '2'. IF NO, RECORD '3'.	HUSBAND . . . . . 1 LIVE-IN PARTNER . . . . . 2 BOYFRIEND NOT LIVING WITH RESPONDE . . 3 CASUAL ACQUAINTANCE . . . . . 4 CLIENT/SEX WORKER . . . . . 5 OTHER _____ 6 (SPECIFY)	
737	In total, with how many different people have you had sexual intercourse in your lifetime? IF NON-NUMERIC ANSWER, PROBE TO GET AN ESTIMATE. IF NUMBER OF PARTNERS IS 95 OR MORE, RECORD '95'.	NUMBER OF PARTNERS IN LIFETIME ☐☐ DON'T KNOW 98	
738	PRESENCE OF OTHERS DURING THIS SECTION.	YES NO CHILDREN <10 . . . . . 1 2 MALE ADULTS . . . . . 1 2 FEMALE ADULTS 1 2	

SECTION 8. FERTILITY PREFERENCES

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
801	CHECK 307: NOT ASKED ☐ ↓ NEITHER ARE STERILIZED ☐ ↓	HE OR SHE STERILIZED ☐	→ 813
802	CHECK 232: PREGNANT ☐ ↓	NOT PREGNANT OR UNSURE ☐	→ 804
803	Now I have some questions about the future. After the child you are expecting now, would you like to have another child, or would you prefer not to have any more children?	HAVE ANOTHER CHILD . . . . . 1 NO MORE . . . . . 2 UNDECIDED/DON'T KNOW 8	→ 805 → 812
804	CHECK 208: HAS HAD A CHILD ☐ ↓ a) Now I have some questions about the future. Would you like to have another child, or would you prefer not to have any more children? HAS NOT HAD A CHILD ☐ ↓ b) Now I have some questions about the future. Would you like to have a child, or would you prefer not to have any children?	HAVE (A/ANOTHER) CHILD . . . . . 1 NO MORE/NONE . . . . . 2 SAYS SHE CAN'T GET PREGNANT . . . . . 3 UNDECIDED/DON'T KNOW 8	→ 807 → 813 → 811
805	CHECK 208 AND 232: NOT PREG. OR UNSURE AND HAS HAD A CHILD ☐ ↓ a) How long would you like to wait from now before the birth of another child? NOT PREG. OR UNSURE AND HAS NOT HAD A CHILD ☐ ↓ b) How long would you like to wait from now before the birth of a child? PREGNANT ☐ ↓ c) After the birth of the child you are expecting now, how long would you like to wait before the birth of another	MONTHS . . . . . 1 ☐☐ YEARS . . . . . 2 ☐☐ SOON/NOW . . . . . 993 SAYS SHE CAN'T GET PREGNANT . . . . . 994 AFTER MARRIAGE . . . . . 995 OTHER _____ (SPECIFY) 996 DON'T KNOW 998	→ 811 → 813 → 811
806	CHECK 232: NOT PREGNANT OR UNSURE ☐ ↓	PREGNANT ☐	→ 812
807	CHECK 307: USING A CONTRACEPTIVE METHOD? NOT ASKED ☐ ↓	CURRENTLY USING ☐	→ 813
808	CHECK 805: '24' OR MORE MONTHS OR '02' OR MORE YEARS ☐ ↓ NOT ASKED ☐ ↓	'00-23' MONTHS OR '00-01' YEAR ☐	→ 812
809	CHECK 723: DAYS, WEEKS OR MONTHS AGO ☐ ↓	YEARS AGO ☐ NOT ASKED ☐	→ 811 → 811

SECTION 8. FERTILITY PREFERENCES

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
810	CHECK 208 AND 804: HAS HAD A CHILD AND WANTS TO HAVE ANOTHER CHILD ☐↓ a) You have said that you do not want another child soon. Can you tell me why you are not using a method to prevent pregnancy? Any other reason? RECORD ALL REASONS MENTIONED. HAS HAD A CHILD AND WANTS NO MORE ☐↓ b) You have said that you do not want any more children. Can you tell me why you are not using a method to prevent pregnancy? Any other reason? RECORD ALL REASONS MENTIONED. HAS NOT HAD A CHILD AND WANTS TO HAVE A CHILD ☐↓ c) You have said that you do not want a child soon. Can you tell me why you are not using a method to prevent pregnancy? Any other reason? RECORD ALL REASONS MENTIONED. HAS NOT HAD A CHILD AND WANTS NO CHILDREN ☐↓ d) You have said that you do not want any children. Can you tell me why you are not using a method to prevent pregnancy? Any other reason? RECORD ALL REASONS MENTIONED.	NOT MARRIED . . . . . A **FERTILITY-RELATED REASONS** NOT HAVING SEX . . . . . B INFREQUENT SEX . . . . . C MENOPAUSAL/HYSTERECTOMY . . . . . D CAN'T GET PREGNANT . . . . . E NOT MENSTRUATED SINCE LAST BIRTH F BREASTFEEDING . . . . . G UP TO GOD/FATALISTIC . . . . . H **OPPOSITION TO USE** RESPONDENT OPPOSEI . . . . . I HUSBAND/PARTNER OPPOSEI . . . . . J OTHERS OPPOSED . . . . . K RELIGIOUS PROHIBITIO . . . . . L **LACK OF KNOWLEDGE** KNOWS NO METHOD . . . . . M KNOWS NO SOURCE . . . . . N **METHOD-RELATED REASONS** INCONVENIENT TO USE . . . . . O CHANGES IN MENSTRUAL BLEEDIN(. . . . P METHODS COULD CAUSE INFERTILITY . . Q INTERFERES WITH BODY'S NORMAL PROCESSES . . . . . R OTHER SIDE EFFECTS . . . . . S **COST/ACCESS/AVAILABILITY** LACK OF ACCESS/TOO FAI . . . . . T COSTS TOO MUCH . . . . . U PREFERRED METHOD NOT AVAILABLE . . V NO METHOD AVAILABLE . . . . . W OTHER _____ X (SPECIFY) DON'T KNOW . . . . . Z	
811	CHECK 307: USING A CONTRACEPTIVE METHOD? NOT ASKED ☐↓ YES, CURRENTLY USING ☐ →		813
812	Do you think you will use a contraceptive method to delay or avoid pregnancy at any time in the future?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	

SECTION 8. FERTILITY PREFERENCES

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
813	CHECK 224: HAS LIVING CHILDREN ↓ / NO LIVING CHILDREN ↓ a) If you could go back to the time you did not have any children and could choose exactly the number of children to have in your whole life, how many would that be? PROBE FOR A NUMERIC RESPONSE. b) If you could choose exactly the number of children to have in your whole life, how many would that be? PROBE FOR A NUMERIC RESPONSE.	NONE . . . . . 00 NUMBER . . . . . [ ][ ] OTHER _____ 96 (SPECIFY)	→ 815 → 815
814	How many of these children would you like to be boys, how many would you like to be girls and for how many would it not matter if it's a boy or a girl?	BOYS GIRLS EITHER NUMBER . . [ ][ ] [ ][ ] [ ][ ] OTHER _____ 96 (SPECIFY)	
815	In the last 12 months have you:	YES NO	
	a) Heard about family planning on the radio?	a) RADIO 1 2	
	b) Seen anything about family planning on the television?	b) TELEVISION 1 2	
	c) Read about family planning in a newspaper or magazine?	c) NEWSPAPER OR MAGAZIN 1 2	
	d) Received a voice or text message about family planning on a mobile phone?	d) MOBILE PHONI 1 2	
	e) Seen anything about family planning on social media such as Facebook, Twitter, Instagram, Tiktok, or WhatsApp?	e) FACEBOOK/TWITTER/ INSTAGRAM/TIKTOK/WHATSAPP 1 2	
	f) Seen anything about family planning on a poster, leaflet or brochure?	f) POSTER/LEAFLET/BROCHURE 1 2	
	g) Seen anything about family planning on an outdoor sign or billboard?	g) OUTDOOR SIGN/BILLBOAR 1 2	
	h) Heard anything about family planning at community meetings or events?	h) COMMUNITY MEETINGS/EVEN 1 2	
	i) Heard anything about family planning from friends/peers?	i) FRIENDS/PEERS 1 2	
	j) Seen anything on family planning on the internet?	j) INTERNET 1 2	

SECTION 11. OTHER HEALTH ISSUES

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
1101	How long does it take in minutes to go from your home to the nearest healthcare facility, which could be a hospital, a medical clinic, a health center, a dispensary, or a medical doctor?	MINUTES . . . . . [ ][ ][ ]	
1102	How do you travel to this healthcare facility from your home? IF MORE THAN ONE WAY OF TRAVEL IS MENTIONED, RECORD THE ONE HIGHEST ON THE LIST.	**MOTORIZED** CAR/TRUCK . . . . . 01 PUBLIC BUS . . . . . 02 MOTORCYCLE/SCOOTER/TUK TUK . . . . 03 BOAT WITH MOTOR . . . . . 04 **NOT MOTORIZED** ANIMAL-DRAWN CART . . . . . 05 BICYCLE . . . . . 06 BOAT WITHOUT MOTOR . . . . . 07 WALKING . . . . . 08 OTHER _____ 96 (SPECIFY)	
1103	Has a doctor or other healthcare provider examined your breasts to check for breast cancer?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
1104	Now I'm going to ask you about tests a healthcare worker can do to check for cervical cancer, which is cancer in the cervix. The cervix connects the womb to the vagina. To be checked for cervical cancer, a woman is asked to lie on her back with her legs apart. Then the healthcare worker will use a brush or swab to collect a sample from inside her. The sample is sent to a laboratory for testing. This test is called a Pap smear or HPV test. Another method is called a VIA or Visual Inspection with Acetic Acid. In this test, the healthcare worker puts vinegar on the cervix to see if there is a reaction.		
1105	Has a doctor or other healthcare worker ever tested you for cervical cancer?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	
1113	Many different factors can prevent women from getting medical advice or treatment for themselves. When you are sick and want to get medical advice or treatment, is each of the following a big problem or not a big problem:	BIG PROBLEM / NOT A BIG PROBLEM a) PERMISSION TO GO . . . . 1 2 b) GETTING MONEY . . . . . 1 2 c) DISTANCE . . . . . 1 2 d) GO ALONE . . . . . 1 2 e) RELIGION . . . . . 1 2 f) CULTURE 1 2	
1114	Are you registered / covered by the Social Health Insurance (SHA) scheme?	YES . . . . . 1 NO 2	→ 1115A
1127	How easy was it to register and enroll in the SHA insurance scheme? Would you say it was very easy, easy, difficult, or very difficult?	VERY EASY . . . . . 1 EASY . . . . . 2 DIFFICULT . . . . . 3 VERY DIFFICULT . . . . . 4 DON'T KNOW 8	→ 1116

SECTION 11. OTHER HEALTH ISSUES

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
1115A	Why are you not registered under the Social Health Authority (SHA) scheme? RECORD ALL MENTIONED	INCONVENIENT/ BURDENSOM . . . . . A HAVE NOT HEARD OF SHA/ NOT AWARE . . . . B AWARE, BUT DO NOT KNOW HOW TO REGIST C DISTRUST IN SYSTEM/ GOVERNMENT . . . . E FINANCIAL BARRIERS/ TOO EXPENSIVE . . . . F PROBLEMS WITH REQUIRED DOCUMENTATIC G INADEQUATE COVERAGE (SERVICES NEEDED ARE NOT COVERED) . . . . . H NOT INTERESTED / SATISFIED WITH OTHER COVERAGE J OTHER _____ X (SPECIFY)	→HPV01
1116	In the last 12 months, did you receive care from a health provider, or a pharmacy?	YES . . . . . 1 NO 2	→ HPV01
1117	Where did you get care most recently? Anywhere else? PROBE TO IDENTIFY EACH TYPE OF SOURCE. IF UNABLE TO DETERMINE IF PUBLIC, PRIVATE, OR NGO SECTOR, RECORD 'Y' AND WRITE THE NAME OF THE PLACE(S).	**PUBLIC SECTOR** GOVERNMENT HOSPITAL . . . . . A GOVERNMENT HEALTH CENTER . . . . . B GOVERNMENT DISPENSARY . . . . . C MOBILE CLINIC . . . . . D COMMUNITY HEALTH WORKER/ FIELDWORKER . . . . . E OTHER PUBLIC SECTOR F SECTOR _____ (SPECIFY) **PRIVATE MEDICAL SECTOR** PRIVATE HOSPITAL . . . . . G PRIVATE CLINIC . . . . . H PHARMACY . . . . . I PRIVATE DOCTOR . . . . . J MOBILE CLINIC . . . . . K COMMUNITY HEALTH WORKER/ FIELDWORKER . . . . . L OTHER PRIVATE MEDICAL SECTOR M _____ (SPECIFY) **NGO MEDICAL SECTOR** NGO HOSPITAL . . . . . N NGO CLINIC . . . . . O OTHER NGO MEDICAL SECTOR P _____ (SPECIFY) **FBO MEDICAL SECTOR** FBO/MISSION HOSPITAL . . . . . Q FBO/MISSION CLINIC . . . . . R FBO MOBILE CLINIC . . . . . S OTHER FBO MEDICAL SECTOR _____ T (SPECIFY) **OTHER SOURCE** SHOP . . . . . U TRADITIONAL PRACTITIONER . . . . . V MARKET . . . . . W ITINERANT DRUG SELLER . . . . . X OTHER _____ Y (SPECIFY)	

SECTION 11. OTHER HEALTH ISSUES

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
1118	What was the main reason for seeking care this most recent time?	OUTPATIENT SERVICES/CONSULTATIO .... 11 INPATIENT CARE/HOSPITALIZATION ..... 12 ACCIDENT/INJURY ..... 13 ANTENATAL CARE SERVICES ..... 14 DELIVERY AND MATERNITY SERVICES .... 15 POSTNATAL CARE SERVICI ..... 16 CHRONIC ILLNESS MANAGEMENT ..... 17 ROUTINE MEDICAL CHECK UP / EXAMINATION 18 LABORATORY/DIAGNOSTIC SERVICE ..... 19 MEDICINES / PHARMACY ..... 20 OTHER _____ 96 (SPECIFY)	
1120	Were any of the health care services you received most recently covered under the SHA scheme?	YES ..... 1 NO 2	
1121	At the time you most recently received health care, did you experience any difficulties accessing SHA services?	NO CHALLENGES FACED ..... A LONG WAITING TIMES ..... B SOME OF THE SERVICES NOT AVAILABLE AT PREFERRED HEALTH FACILITY ..... C LACK OF MEDICINES OR SUPPLIES ..... D POOR PROVIDER ATTITUDE ..... E CONFUSION ABOUT BENEFITS ..... F INADEQUACY OF BENEFITS COVERED .... G NOT ACCEPTED AT PREFERRED HEALTH FACILITY ..... H PREMIUMS NOT PAID .... I SHA SYSTEM DOWN J OTHER _____ X (SPECIFY)	
1121A	CHECK 1120: YES ☐ NO ☐ → HPV01		
1122	When you most recently received health care services, what benefits did you receive through SHA coverage?	REDUCED OUT-OF-POCKET SPENDING ON HEALTHCARE ..... A INCREASED ACCESS TO QUALITY HEALTH SERVICES ..... B TIMELY MEDICAL ATTENTION ..... C IMPROVED PERSONAL HEALTH ..... D REDUCED STRESS OVER HEALTH EXPENSES E INCLUSION OF DEPENDENT ..... F OTHER _____ X (SPECIFY)	
1126	Now, I would like to ask you some questions about your experiences and satisfaction with the services provided under the Social Health Authority (SHA) insurance scheme. Your responses will help us assess the quality, accessibility, and effectiveness of SHA services and identify areas for improvement.		
1128	How satisfied are you with the availability of SHA services in contracted facilities? Would you say you are not satisfied at all, somewhat satisfied, satisfied, or very satisfied?	NOT SATISFIED AT ALL ..... 1 SOMEWHAT SATISFIED ..... 2 SATISFIED ..... 3 VERY SATISFIED ..... 4 NOT APPLICABLE ..... 5 DC 8	

SECTION 11. OTHER HEALTH ISSUES

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
1129	How satisfied are you with the quality of care you received? Would you say you are not satisfied at all, somewhat satisfied, satisfied, or very satisfied?	NOT SATISFIED AT ALL . . . . . 1 SOMEWHAT SATISFIED . . . . . 2 SATISFIED . . . . . 3 VERY SATISFIED . . . . . 4 NOT APPLICABLE . . . . . 5 DON'T KNOW 8	
1130	How satisfied are you with the respect shown by healthcare providers? Would you say you are not satisfied at all, somewhat satisfied, satisfied, or very satisfied?	NOT SATISFIED AT ALL . . . . . 1 SOMEWHAT SATISFIED . . . . . 2 SATISFIED . . . . . 3 VERY SATISFIED . . . . . 4 NOT APPLICABLE . . . . . 5 DON'T KNOW 8	
1131	How satisfied are you with the timeliness of service delivery? Would you say you are not satisfied at all, somewhat satisfied, satisfied, or very satisfied?	NOT SATISFIED AT ALL . . . . . 1 SOMEWHAT SATISFIED . . . . . 2 SATISFIED . . . . . 3 VERY SATISFIED . . . . . 4 NOT APPLICABLE . . . . . 5 DON'T KNOW 8	
1132	How satisfied are you with the coverage of essential medicines and diagnostics? Would you say you are not satisfied at all, somewhat satisfied, satisfied, or very satisfied?	NOT SATISFIED AT ALL . . . . . 1 SOMEWHAT SATISFIED . . . . . 2 SATISFIED . . . . . 3 VERY SATISFIED . . . . . 4 NOT APPLICABLE . . . . . 5 DON'T KNOW 8	
1133	How satisfied are you with the communication and public information provided? Would you say you are not satisfied at all, somewhat satisfied, satisfied, or very satisfied?	NOT SATISFIED AT ALL . . . . . 1 SOMEWHAT SATISFIED . . . . . 2 SATISFIED . . . . . 3 VERY SATISFIED . . . . . 4 NOT APPLICABLE . . . . . 5 DON'T KNOW 8	
1134	Overall, how satisfied are you with SHA? Would you say you are not satisfied at all, somewhat satisfied, satisfied, or very satisfied?	NOT SATISFIED AT ALL . . . . . 1 SOMEWHAT SATISFIED . . . . . 2 SATISFIED . . . . . 3 VERY SATISFIED . . . . . 4 NOT APPLICABLE . . . . . 5 DON'T KNOW 8	
1135	How likely are you to recommend SHA services to your friends, family, and others? Would you say you are not at all likely, unlikely, likely, or very likely?	NOT ALL LIKELY . . . . . 1 UNLIKELY . . . . . 2 LIKELY . . . . . 3 VERY LIKELY . . . . . 4 NOT APPLICABLE . . . . . 5 DON'T KNOW 8	

HPV VACCINATION MODULE

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
HPV02	Now I would like to ask some questions about human papillomavirus or HPV vaccinations that you have received. An HPV vaccine is an injection given in the left or right upper arm to girls between the ages of 10-14 years, as a protection against cervical cancer. In Kenya, the HPV vaccine is also commonly referred to as Cervarix/Gardasil and is commonly given at school/at a medical facility.		
HPV01	CHECK 111: 15-19 YEARS OLD ☐ ↓ 20-49 YEARS OLD ☐		→ HPV07
HPV03	Do you have a card or other document where HPV vaccination is written down?	YES, HAS CARD OR OTHER DOCUMENT . . . . . 1 NO, NO CARD AND NO OTHER DOCUMENT . . 2	→ HPV05
HPV04	Did you ever receive an HPV vaccination card ?	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	→ HPV07
HPV05	May I see the card or other document where HPV vaccination is written down?	YES, CARD OR OTHER DOCUMENT SEEN . . . . . 1 NO CARD AND NO OTHER DOCUMENT SEEN . . 4	→ HPV07
HPV06	COPY VACCINATION DATES FROM THE CARD. RECORD '44' IN 'DAY' COLUMN IF CARD SHOWS THAT A DOSE WAS GIVEN, BUT NO DATE IS RECORDED. RECORD '00' IN 'DAY' COLUMN IF CARD IS BLANK FOR THE DOSE. DAY MONTH YEAR HUMAN PAPILLOMAVIRUS (HPV) 1 ☐☐ ☐☐ ☐☐☐☐ HUMAN PAPILLOMAVIRUS (HPV) 2 ☐☐ ☐☐ ☐☐☐☐		→ HPV09
HPV07	Have you ever received a vaccination against HPV, that is, an injection in the left or right upper arm to protect against cervical cancer? IF NO OR DON'T KNOW: In Kenya, the HPV vaccine is also referred to as Cervarix/Gardasil and is commonly given at school/at a medical facility to girls between the ages of 10-14.	YES . . . . . 1 NO . . . . . 2 DON'T KNOW 8	→ MM01
HPV08	Did you receive one or two doses of the HPV vaccine?	ONE DOSE . . . . . 1 TWO DOSES . . . . . 2 DON'T KNOW 8	
HPV09	Where did you receive your most recent HPV vaccination? PROBE TO IDENTIFY THE TYPE OF SOURCE. IF UNABLE TO CLASSIFY THE SOURCE, RECORD '96' AND WRITE THE NAME OF THE PLACE.	**HEALTH FACILITY** PUBLIC HEALTH FACILITY . . . . . 11 PRIVATE HEALTH FACILITY . . . . . 12 NGO HEALTH FACILITY . . . . . 13 SCHOOL . . . . . 21 OTHER _____ 96 (SPECIFY) DON'T KNOW 98	

SECTION MM. ADULT AND MATERNAL MORTALITY MODULE

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
MM01	Please tell me, which brother or sister was born first? And which was born next? RECORD '01' FOR THE ORDER NUMBER FOR THE FIRST BROTHER OR SISTER, '02' FOR THE SECOND, AND SO ON UNTIL YOU HAVE RECORDED THE ORDER NUMBER FOR ALL BROTHERS AND SISTERS. NAME / ORDER NUMBER a _____ ☐☐ b _____ ☐☐ c _____ ☐☐ d _____ ☐☐ e _____ ☐☐ f _____ ☐☐ g _____ ☐☐ h _____ ☐☐ i _____ ☐☐ j _____ ☐☐ NAME / ORDER NUMBER k _____ ☐☐ l _____ ☐☐ m _____ ☐☐ n _____ ☐☐ o _____ ☐☐ p _____ ☐☐ q _____ ☐☐ r _____ ☐☐ s _____ ☐☐ t _____ ☐☐		
MM02	CHECK MM01:	ONE OR MORE BROTHERS OR SISTERS LISTED ☐ NO BROTHERS OR SISTERS LISTED ☐ →	MM04
MM03	READ THE NAMES OF THE BROTHERS AND SISTERS TO THE RESPONDENT AND AFTER THE LAST ONE ASK: Are there any other brothers and sisters from the same mother that you have not mentioned?	NO ☐ YES ☐ → LIST ADDITIONAL BROTHERS AND SISTERS IN MM01.	
MM04	Sometimes people forget to mention children born to their biological mother because they do not live with them or they do not see them very often. Are there any brothers or sisters who do not live with you that you have not mentioned?	NO ☐ YES ☐ → LIST ADDITIONAL BROTHERS AND SISTERS IN MM01.	
MM05	Sometimes people forget to mention children born to their biological mother because they have died. Are there any brothers or sisters who died that you have not mentioned?	NO ☐ YES ☐ → LIST ADDITIONAL BROTHERS AND SISTERS IN MM01.	
MM06	Some people have brothers or sisters from the same mother but a different father. Are there any brothers or sisters born to your biological mother, but who have a different natural father, that you have not mentioned?	NO ☐ YES ☐ → LIST ADDITIONAL BROTHERS AND SISTERS IN MM01.	
MM07	COUNT THE NUMBER OF BROTHERS AND SISTERS RECORDED IN MM01.	TOTAL BROTHERS AND SISTERS . . ☐☐	

SECTION MM. ADULT AND MATERNAL MORTALITY MODULE

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
MM08	CHECK MM07: Just to make sure that I have this right: Your mother had in total {NUMBER OF BIRTHS TO MOTHER} births, excluding you, during her lifetime. Is that correct? YES ☐ NO ☐ → PROBE AND CORRECT MM01 AND/OR MM07.		
MM09	CHECK MM07: ONE OR MORE BROTHERS/SISTERS ☐ NO BROTHER OR SISTER ☐ →		DV00
MM10	Please tell me, which brother or sister was born first? And which was born next? RECORD '01' FOR THE ORDER NUMBER IN MM01 FOR THE FIRST BROTHER OR SISTER, '02' FOR THE SECOND, AND SO ON UNTIL YOU HAVE RECORDED THE ORDER NUMBER FOR ALL BROTHERS		
MM11	How many births did your mother have before you were born?	NUMBER OF PRECEDING BIRTHS . . ☐☐	

SECTION MM. ADULT AND MATERNAL MORTALITY MODULE

MM12	LIST THE BROTHERS AND SISTERS ACCORDING TO THE ORDER NUMBER IN MM01. ASK MM13 TO MM24 FOR ONE BROTHER OR SISTER BEFORE ASKING ABOUT THE NEXT BROTHER OR SISTER.			
MM13	NAME OF BROTHER OR SISTER.	(01) _____	(02) _____	(03) _____
MM14	Is {NAME IN MM13} male or female?	MALE . . . . . 1 FEMALE 2	MALE . . . . . 1 FEMALE 2	MALE . . . . . 1 FEMALE 2
MM15	Is {NAME IN MM13} still alive?	YES . . . . . 1 NO . . . . . 2 → GO TO MM17 DK . . . . . 8 → GO TO (02)	YES . . . . . 1 NO . . . . . 2 → GO TO MM17 DK . . . . . 8 → GO TO (03)	YES . . . . . 1 NO . . . . . 2 → GO TO MM17 DK . . . . . 8 → GO TO (04)
MM16	How old is {NAME IN MM13}?	AGE . . . . ☐☐ GO TO (02)	AGE . . . . ☐☐ GO TO (03)	AGE . . . . ☐☐ GO TO (04)
MM17	How many years ago did {NAME IN MM13} die?	YEARS AGO . . ☐☐	YEARS AGO . . ☐☐	YEARS AGO . . ☐☐
MM18	IF MALE ↓ a) How old was {NAME IN MM13} when he died? IF DON'T KNOW, PROBE AND ASK ADDITIONAL QUESTIONS TO GET AN ESTIMATE. IF FEMALE ↓ b) How old was {NAME IN MM13} when she died? IF DON'T KNOW, PROBE AND ASK ADDITIONAL QUESTIONS TO GET AN ESTIMATE.	AGE . . . . ☐☐ IF MALE OR DIED BEFORE 12 YEARS OF AGE, GO TO MM23	AGE . . . . ☐☐ IF MALE OR DIED BEFORE 12 YEARS OF AGE, GO TO MM23	AGE . . . . ☐☐ IF MALE OR DIED BEFORE 12 YEARS OF AGE, GO TO MM23

MM19	Was {NAME IN MM13} pregnant when she died?	YES . . . . . 1 → GO TO MM23 NO 2	YES . . . . . 1 → GO TO MM23 NO 2	YES . . . . . 1 → GO TO MM23 NO 2
MM20	Did {NAME IN MM13} die during childbirth?	YES . . . . . 1 → GO TO (02) NO 2	YES . . . . . 1 → GO TO (03) NO 2	YES . . . . . 1 → GO TO (04) NO 2
MM21	Did {NAME IN MM13} die within two months after the end of a pregnancy or childbirth?	YES . . . . . 1 NO . . . . . 2 → GO TO MM23	YES . . . . . 1 NO . . . . . 2 → GO TO MM23	YES . . . . . 1 NO . . . . . 2 → GO TO MM23
MM22	How many days after the end of the pregnancy or childbirth did {NAME IN MM13} die?	DAYS . . ☐☐	DAYS . . ☐☐	DAYS . . ☐☐
MM23	Was {NAME IN MM13}'s death due to an act of violence?	YES . . . . . 1 → GO TO MM24A NO 2	YES . . . . . 1 → GO TO MM24A NO 2	YES . . . . . 1 → GO TO MM24A NO 2
MM24	Was {NAME IN MM13}'s death due to an accident?	YES . . . . . 1 NO . . . . . 2 GO TO (02)	YES . . . . . 1 NO . . . . . 2 GO TO (03)	YES . . . . . 1 NO . . . . . 2 GO TO (04)

MM24A	CHECK MM14	IF MALE GO TO (02) IF FEMALE ☐	IF MALE GO TO (03) IF FEMALE ☐	IF MALE GO TO (04) IF FEMALE ☐
MM24B	To the best of your knowledge, who killed {NAME IN MM13}?	**INTIMATE PARTNER** HUSBAND/PARTNER 11 BOYFRIEND 12 FORMER HUSBAND/ PARTNER 13 FORMER BOYFRIEND 14 **OTHER FAMILY MEMBERS** MOTHER/STEP MOTHER 21 FATHER/STEP FATHER 22 BROTHER 23 SISTER 24 COUSIN 25 OTHER RELATIVE 26 **PERSONS IN AUTHORITY** POLICE/SOLDIER 31 CHIEF/ADMINISTRATION OFFICER 32 TEACHER/LECTURER 33 **OTHER** NEIGHBOUR 41 BUSINESS PERSON 42 PRIEST/RELIGIOUS LEADER 43 FRIEND/FAMILY FRIEND 44 SCHOOLMATE/CLASSMATE 45 EMPLOYER/SOMEONE AT WORK 46 COACH/MENTOR 47 **SUICIDE** 51 ☐← (GO TO (02)) OTHER _____ 96 (SPECIFY) DON'T KNOW KILLER 98	SAME RESPONSE OPTIONS	SAME RESPONSE OPTION
MM24C	To the best of your knowledge, what was the main reason {NAME IN MM13} was killed?	SUCCESSION/FAMILY PROPERTY-RELATED CONFLICTS 01 INFIDELITY/CHEATING 02 CRIME OF PASSION 03 SPOUSAL INSECURITY 04 MARRITAL CONFLICT 05 CRIMINAL DEAL GONE WRONG/CONC 06 BUSINESS RIVALRY 07 RETALIATION/REVENGE 08 HONOUR KILLINGS 09 WITCHCRAFT ALLEGATION 10 OTHER _____ 96 SPECIFY DON'T KNOW 98	SAME RESPONSE OPTIONS	SAME RESPONSE OPTION
MM24D	Where was {NAME IN MM13} killed?	VICTIM'S RESIDENCE 01 PERPETRATOR'S RESIDENCE 02 BUSH/RIVER 03 VACATION 04 HOTEL/BAR 05 AMUSEMENT PARK 06 OTHER _____ 96 SPECIFY DON'T KNOW 98 GO TO (02)	SAME RESPONSE OPTIONS GO TO (03)	SAME RESPONSE OPTION GO TO (04)
IF NO MORE BROTHERS OR SISTERS, GO TO NEXT SECTION.				

TECHNOLOGY-FACILITATED GENDER-BASED VIOLENCE

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
DV00	CHECK COVER PAGE: WOMAN SELECTED FOR TFGBV MODULE? WOMAN SELECTED FOR THIS SECTION ↓ WOMAN NOT SELECTED →		NEXT SECT.
DV01	CHECK FOR PRESENCE OF OTHERS: DO NOT CONTINUE UNTIL PRIVACY IS ENSURED. PRIVACY OBTAINED . . . . . 1 ↓ PRIVACY NOT POSSIBLE . . . . . 2 →		TV13
DV02	Now I would like to ask you questions about some other important aspects of a woman's life. You may find some of these questions very personal. However, your answers are crucial for helping to understand the condition of women in Kenya. Let me assure you that your answers are completely confidential and will not be told to anyone and no one else in your household will know that you were asked these questions. If I ask you any question you don't want to answer, just let me know and I will go on to the next question.		
DV03	CHECK 701 AND 702: CURRENTLY MARRIED/ LIVING WITH A MAN → NEVER MARRIED/ NEVER LIVED WITH A MAN ↓ FORMERLY MARRIED/ LIVED WITH A MAN (READ IN PAST TENSE AND USE 'LAST' WITH 'HUSBAND/ MALE PARTNER') →		DV06 DV06
DV04	You have said that you are not married and are not living with a man as if married. Are you currently in an intimate relationship with a man even though you are not living with him?	YES . . . . . 1 NO 2	→DV06
DV05	Have you ever been in an intimate relationship with a man even though you did not ever live with him?	YES . . . . . 1 NO 2	→TV09
DV06	Now, I am going to ask you about some situations that can happen between some women and their (husband/male partner).		

	A. Please tell me if these descriptions apply to your relationship with your (last) (husband/male partner).	EVER	B. How often did this happen during the last 12 months: often, only sometimes, or not at all? OFTEN	SOME-TIMES	NOT IN LAST 12 MONTHS
a)	He (is/was) jealous or angry if you (talk/talked) to other men?	YES 1 NO 2 ↓	→ 1	2	3
b)	He wrongly (accuses/accused) you of being unfaithful?	YES 1 NO 2 ↓	→ 1	2	3
c)	He (does/did) not permit you to meet your female friends?	YES 1 NO 2 ↓	→ 1	2	3
d)	He (tries/tried) to limit your contact with your family?	YES 1 NO 2 ↓	→ 1	2	3
e)	He (insists/insisted) on knowing where you (are/were) at all times?	YES 1 NO 2 ↓	→ 1	2	3
f)	He (insists/insisted) on reading call log/messages/emails from your phone/tablet/laptop or other digital tools without your consent?	YES 1 NO 2 ↓	→ 1	2	3
g)	He (insists/insisted) on you giving him passwords, pin, or patterns for your phone, tablet, laptop, email, facebook, Instagram, whatsapp, snapchat or other digital platforms?	YES 1 NO 2 ↓	→ 1	2	3
h)	He (uses/used) mobile, internet, or other technology including phone calls, facetime, video calling, or tracking software, to check on you or track where you (are/were) in ways that (makes/made) you feel controlled?	YES 1 NO 2 ↓	→ 1	2	3

TECHNOLOGY-FACILITATED GENDER-BASED VIOLENCE

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
TV07	A. Did your (last) (husband/male partner) ever do any of the following things to you:	B. How often did this happen during the last 12 months: often, only sometimes, or not at all?	

		EVER		OFTEN	SOME-TIMES	NOT IN LAST 12 MONTHS
a)	REVENGE PORNOGRAPHY: share with other people your sexually explicit messages, images or videos?	YES NO	1 2 ↓	1 →	2	3
b)	DOXXING:share your private or personal information (e.g. address, phone number, email) with malicious intent?	YES NO	1 2 ↓	1 →	2	3
c)	BLACKMAIL/SEXTORTION: threaten to share sensitive information , images, audio or video about you unless certain demands, financial, sexual, emotional are	YES NO	1 2 ↓	1 →	2	3
d)	HACKING: gain unauthorized access to your personal devices, social media, or emails to track,control or harm you?	YES NO	1 2 ↓	1 →	2	3
e)	CYBERBULLYNG: use digital tools to spread false, distorted or exaggerated information about you?	YES NO	1 2 ↓	1 →	2	3
f)	ONLINE HARASSMENT: Send you threatening messages (via the internet, text messages, instant messages, social media or through other digital tools) or call/called you repeatedly to harass you ?	YES NO	1 2 ↓	1 →	2	3
g)	CYBERSTALKING: use technology or digital devices to monitor and stalk you in real-time or historically with malicious intent?	YES NO	1 2 ↓	1 →	2	3

TV08	A. So far we have been talking about the behavior of your (current/last) (husband/male partner). Now I want to ask you about the behavior of any previous husband or any other current or previous male partner that you may have ever had.			B. How long ago did this last happen?			
		EVER		0 - 11 MONTHS AGO	12+ MONTHS AGO	DON'T REMEMBER	
		HAS NEVER HAD ANOTHER HUSBAND/ MALE PARTNER		6 →			TV09
a)	Did any previous husband or any other current or previous male partner use digital tools to share with other people your sexually explicit messages,images, audio, videos, or your personal information, or spread false information about you or even gain unauthorized access to your personal devices, social media, or emails to track, control,or harm you?	YES NO	1 2 ↓	1 →	2	3	

TECHNOLOGY-FACILITATED GENDER-BASED VIOLENCE

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
TV09	CHECK 701 AND 702 AND DV04 AND DV05: EVER MARRIED/EVER LIVED WITH A MAN/ EVER HAD A MALE PARTNER ☐ → a) NEVER MARRIED/ NEVER LIVED WITH A MAN/ NEVER HAD A MALE PARTNER ☐ → b)		

	a) At any time in your life,as a child or an adult, has anyone other than any of your previous husband, or any other current, or previous male partner that you may have ever had,done any of the following things to you.Remember I do not want you to include any husband or male partner	b) At any time in your life,as a child or an adult, has anyone ever done any of the following things to you:	EVER		B. How often did this happen during the last 12 months: often, only sometimes, or not at all? OFTEN	SOME-TIMES	NOT IN LAST 12 MONTHS	
Aa)		REVENGE PORNOGRAPHY:share with other people your sexually explicit messages, images or videos?	YES NO	1 2 →	1	2	3	
Ab)		ONLINE HARASSMENT: Sent you threatening messages via the internet, text messages, instant messages, social media or through other digital tools/digital platforms or call/called you repeatedly to harass you ?	YES NO	1 2 →	1	2	3	
Ac)		DOXXING: shared your private or personal information (e.g. address, phone number, email) with malicious intent?	YES NO	1 2 →	1	2	3	
Ad)		BLACKMAIL/SEXTORTION: threatened to share sensitive information, images, audio or video unless certain demands, financial, sexual, emotional are met?	YES NO	1 2 →	1	2	3	
Ae)		HACKING: gained unauthorized access to your personal devices, social media, or emails to track,control or harm you?	YES NO	1 2 →	1	2	3	
Af)		CYBERBULLYING: used digital tools (internet, text messages, instant messages, phone calls, social media or any other digital tools) to spread false, distorted or exaggerated information about you or publicly humiliate you?	YES NO	1 2 →	1	2	3	
Ag)		CYBERSTALKING: used technology or digital devices to monitor and stalk you in real-time or historically with malicious intent ?	YES NO	1 2 →	1	2	3	
Ah)		ONLINE SEXUAL HARASSMENT OR CYBER SEXUAL ABUSE: Sent you unwanted sexual explicit messages, images, audio or videos	YES NO	1 2	1	2	3	

TV10	CHECK TV09A (a-g) : AT LEAST ONE 'YES' ☐ →	NOT A SINGLE YES ☐ →	TV12

TECHNOLOGY-FACILITATED GENDER-BASED VIOLENCE

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
TV11	Who has done or did these things to you? Anyone else? RECORD ALL MENTIONED	FATHER/STEP-FATHER ..... A MOTHER/STEP-MOTHER ..... B BROTHER/STEP-BROTHER ..... C SISTER/STEP-SISTER ..... D OTHER RELATIVE ..... E CURRENT BOYFRIEND ..... F FORMER BOYFRIEND ..... G IN-LAW ..... H OWN FRIEND/ACQUAINTANCE ..... I FAMILY FRIEND ..... J TEACHER ..... K SCHOOLMATE/CLASSMATE ..... L EMPLOYER/SOMEONE AT WORK .. M POLICE/SOLDIER ..... N PRIEST/RELIGIOUS LEADER O STRANGER ..... P ONLINE ATTACKER ..... Q COACH/MENTOR ..... R SCHOOLMATE/CLASSMATE ..... S OTHER _____ X (SPECIFY)	
TV11A	CHECK TV07(a-g) ==1 OR TV08Aa ==1 OR TV09A (Aa-Ai) AT LEAST ONE 'YES' ☐ ↓	NOT A SINGLE YES ☐ →	TV12
TV11B	On which platforms did you experience these acts/behaviors? RECORD ALL MENTIONED	**SOCIAL MEDIA** FACEBOOK ..... A X (FORMERLY TWITTER ..... B INSTAGRAM ..... C THREADS ..... D TIKTOK ..... E YOUTUBE ..... F REDDIT ..... G OTHER SOCIAL MEDIA ..... H **MESSAGING APPS** TEXT MESSAGE/ IMESSAGE ..... I WHATSAPP ..... J SIGNAL ..... K TELEGRAM ..... L OTHER MESSAGING API ..... M **DATING APPS** .. TINDER ..... N BUMBLE ..... O KENYA SINGLES DATING APP ..... P OTHER DATING APP ..... Q **OTHER** EMAIL ..... R VIRTUAL MEETING TOOLS ..... S ONLINE GAMING ..... T ONLINE NEWS/ BLOGS ..... U LINKED IN ..... V **OTHER** _____ X (SPECIFY)	
	THANK THE RESPONDENT FOR HER COOPERATION AND REASSURE HER ABOUT THE CONFIDENTIALITY OF HER ANSWERS. FILL OUT THE QUESTIONS BELOW WITH REFERENCE TO THE TECHNOLOGY-FACILITATED GENDER-BASED VIOLENCE MODULE ONLY.		

TECHNOLOGY-FACILITATED GENDER-BASED VIOLENCE

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
TV12	DID YOU HAVE TO INTERRUPT THE INTERVIEW BECAUSE SOME ADULT WAS TRYING TO LISTEN, OR CAME INTO THE ROOM, OR INTERFERED IN ANY OTHER WAY?	YES, ONCE / YES, MORE THAN ONCE / NO HUSBAND . . . . . 1 / 2 / 3 OTHER MALE ADULT . . 1 / 2 / 3 FEMALE ADULT . . . . . 1 / 2 / 3	
TV13	INTERVIEWER'S COMMENTS/EXPLANATION FOR NOT COMPLETING THE TECHNOLOGY-FACILITATED GENDER-BASED VIOLENCE MODULE.		
1500	RECORD THE TIME.	HOURS . . . . . [ ][ ] MINUTE . . . . . [ ][ ]	

INSTRUCTIONS:

ONLY ONE CODE SHOULD APPEAR IN ANY BOX.
COLUMN 1 REQUIRES A CODE IN EVERY MONTH.

CODES FOR EACH COLUMN:

COLUMN 1: BIRTHS, PREGNANCIES, CONTRACEPTIVE USE (2)

- B BIRTHS
- P PREGNANCIES
- T TERMINATIONS

- 0 NO METHOD
- 1 FEMALE STERILIZATION
- 2 MALE STERILIZATION
- 3 IUD
- 4 INJECTABLES
- 5 IMPLANTS
- 6 PILL
- 7 CONDOM
- 8 FEMALE CONDOM
- 9 EMERGENCY CONTRACEPTION
- J STANDARD DAYS METHOD
- K LACTATIONAL AMENORRHEA METHOD
- L RHYTHM METHOD
- M WITHDRAWAL
- X OTHER MODERN METHOD
- Y OTHER TRADITIONAL METHOD

COLUMN 2: DISCONTINUATION OF CONTRACEPTIVE USE

- 0 INFREQUENT SEX/HUSBAND AWAY
- 1 BECAME PREGNANT WHILE USING
- 2 WANTED TO BECOME PREGNANT
- 3 HUSBAND/PARTNER DISAPPROVED
- 4 WANTED MORE EFFECTIVE METHOD
- 5 CHANGES IN MENSTRUAL BLEEDING
- 6 OTHER SIDE EFFECTS/HEALTH CONCERNS
- 7 LACK OF ACCESS/TOO FAR
- 8 COSTS TOO MUCH
- N INCONVENIENT TO USE
- F UP TO GOD/FATALISTIC
- A DIFFICULT TO GET PREGNANT/MENOPAUSAL
- D MARITAL DISSOLUTION/SEPARATION
- E METHOD EXPIRED
- X OTHER _____ (SPECIFY)
- Z DON'T KNOW

				COL. 1	COL. 2	
	12	DEC	01			
	11	NOV	02			
	10	OCT	03			
2	09	SEP	04			2
	08	AUG	05			
0	07	JUL	06			0
2	06	JUN	07			2
	05	MAY	08			
5	04	APR	09			5
	03	MAR	10			
	02	FEB	11			
	01	JAN	12			
	12	DEC	13			
	11	NOV	14			
	10	OCT	15			
2	09	SEP	16			2
	08	AUG	17			
0	07	JUL	18			0
2	06	JUN	19			2
	05	MAY	20			
4	04	APR	21			4
	03	MAR	22			
	02	FEB	23			
	01	JAN	24			
	12	DEC	25			
	11	NOV	26			
	10	OCT	27			
2	09	SEP	28			2
	08	AUG	29			
0	07	JUL	30			0
2	06	JUN	31			2
	05	MAY	32			
3	04	APR	33			3
	03	MAR	34			
	02	FEB	35			
	01	JAN	36			
	12	DEC	37			
	11	NOV	38			
	10	OCT	39			
2	09	SEP	40			2
	08	AUG	41			
0	07	JUL	42			0
2	06	JUN	43			2
	05	MAY	44			
2	04	APR	45			2
	03	MAR	46			
	02	FEB	47			
	01	JAN	48			
	12	DEC	49			
	11	NOV	50			
	10	OCT	51			
2	09	SEP	52			2
	08	AUG	53			
0	07	JUL	54			0
2	06	JUN	55			2
	05	MAY	56			
1	04	APR	57			1
	03	MAR	58			
	02	FEB	59			
	01	JAN	60			
	12	DEC	61			
	11	NOV	62			
	10	OCT	63			
2	09	SEP	64			2
	08	AUG	65			
0	07	JUL	66			0
2	06	JUN	67			2
	05	MAY	68			
0	04	APR	69			0
	03	MAR	70			
	02	FEB	71			
	01	JAN	72			

INTERVIEWER'S OBSERVATIONS

TO BE FILLED IN AFTER COMPLETING INTERVIEW

COMMENTS ABOUT INTERVIEW:

COMMENTS ON SPECIFIC QUESTIONS:

ANY OTHER COMMENTS:

SUPERVISOR'S OBSERVATIONS

2025 KENYA MINI DEMOGRAPHIC AND HEALTH SURVEY (KMDHS)
FIELDWORKER QUESTIONNAIRE

KENYA
KENYA NATIONAL BUREAU OF STATISTICS

LANGUAGE OF QUESTIONNAIRE ENGLISH

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
100	What is your name?	NAME _____	
101	RECORD FIELDWORKER NUMBER	NUMBER . . . . . [ ][ ][ ][ ]	

INSTRUCTIONS

Information on all KMDHS field workers is collected as part of the KMDHS survey. Please fill out the questions below. The information you provide will be part of the survey data file; however, your name will be removed and will not be part of the data file. Thank you for providing the information needed.

102	In what county do you live? REFER TO COUNTY CODE LIST IN INTERVIEWER'S MANUAL	COUNTY CODE . . . . . [ ][ ]	
103	Do you live in a city, town, or rural area?	CITY (NAIROBI, KISUMU, MOMBASA, NAKURU) . 1 TOWN . . . . . 2 RURAL AREA 3	
104	How old are you? RECORD AGE IN COMPLETED YEARS.	AGE . . . . . [ ][ ]	
105	Are you male or female?	MALE . . . . . 1 FEMALE 2	
106	What is your current marital status?	CURRENTLY MARRIED . . . . . 1 LIVING WITH A MAN/WOMAN . . . . . 2 WIDOWED . . . . . 3 DIVORCED . . . . . 4 SEPARATED . . . . . 5 NEVER MARRIED OR LIVED WITH A MAN/WOMAN 6	
107	How many living children do you have? INCLUDE ONLY CHILDREN WHO ARE YOUR BIOLOGICAL CHILDREN.	LIVING CHILDREN . . . . . [ ][ ]	
108	Have you ever had a child who died?	YES . . . . . 1 NO 2	
109	What is the highest level of school you attended: primary, secondary/'A' level, college, university, or vocational?	PRIMARY . . . . . 1 SECONDARY/ 'A' LEVEL . . . . . 2 MIDDLE LEVEL COLLEGE (CERTIFICATE/ DIPLOMA) . . . . . 3 UNIVERSITY . . . . . 4 VOCATIONAL TRAINING . . . . . 5 INFORMAL EDUCATION (MADRASA/ ADULT BASIC) 6	
110	What is the highest (standard/grade/form/year) you completed at that level? IF COMPLETED LESS THAN ONE YEAR AT THAT LEVEL, RECORD '00'.	STANDARD/GRADE/FORM/YEAR . . [ ][ ]	

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
111	What is your religion?	CATHOLIC . . . . . 01 PROTESTANT . . . . . 02 EVANGELICAL CHURCHES . . . . . 03 AFRICAN INSTITUTED CHURCHES . . . . . 04 ORTHODOX . . . . . 05 OTHER CHRISTIAN . . . . . 06 ISLAM . . . . . 07 HINDU . . . . . 08 TRADITIONISTS . . . . . 09 NO RELIGION . . . . . 95 OTHER _____ 96 (SPECIFY)	
112	What is your ethnic group?	EMBU . . . . . 01 KALENJIN . . . . . 02 KAMBA . . . . . 03 KIKUYU . . . . . 04 KISII . . . . . 05 LUHYA . . . . . 06 LUO . . . . . 07 MAASAI . . . . . 08 MERU . . . . . 09 MIJIKENDA/SWAHILI . . . . . 10 SOMALI . . . . . 11 TAITA/TAVETA . . . . . 12 OTHER _____ 96 (SPECIFY)	
113	What languages can you speak? RECORD ALL LANGUAGES YOU CAN SPEAK.	ENGLISH . . . . . A KISWAHILI . . . . . B BORANA . . . . . C EMBU . . . . . D KALENJIN . . . . . E KAMBA . . . . . F KIKUYU . . . . . G KISII . . . . . H LUHYA . . . . . I MARAGOLI . . . . . J LUO . . . . . K MAASAI . . . . . L MERU . . . . . M MIJIKENDA . . . . . N POKOT . . . . . O SOMALI . . . . . P TURKANA . . . . . Q OTHER _____ X (SPECIFY)	

NO.	QUESTIONS AND FILTERS	CODING CATEGORIES	SKIP
114	What is your mother tongue/native language (language spoken at home growing up)?	ENGLISH . . . . . 01 KISWAHILI . . . . . 02 BORANA . . . . . 03 EMBU . . . . . 04 KALENJIN . . . . . 05 KAMBA . . . . . 06 KIKUYU . . . . . 07 KISII . . . . . 08 LUHYA . . . . . 09 MARAGOLI . . . . . 10 LUO . . . . . 11 MAASAI . . . . . 12 MERU . . . . . 13 MIJIKENDA . . . . . 14 POKOT . . . . . 15 SOMALI . . . . . 16 TURKANA . . . . . 17 OTHER _____ 96 (SPECIFY)	
115	Have you ever worked on: a) a DHS prior to this survey? b) an MIS prior to this survey? c) any other survey prior to this survey?	YES NO a) DHS . . . . . 1 2 b) MIS . . . . . 1 2 c) OTHER SURVEY 1 2	
116	Were you already working for KNBS at the time you were employed to work on this KMDHS?	YES, KNBS . . . . . 1 NO 3	→ 118
117	Are you a permanent or temporary employee of KNBS?	PERMANENT . . . . . 1 TEMPORARY 2	
118	If you have comments, please write them here.		

ADDITIONAL DHS PROGRAM RESOURCES

The DHS Program Website – Download free DHS reports, standard documentation, key indicator data, and training tools, and view announcements.	DHSprogram.com	
STATcompiler – Build custom tables, graphs, and maps with data from 90 countries and thousands of indicators.	Statcompiler.com	
DHS Program Mobile App – Access key DHS indicators for 90 countries on your mobile device (Apple, Android, or Windows).	Search DHS Program in your iTunes or Google Play store	
DHS Program User Forum – Post questions about DHS data and search our archive of FAQs.	userforum.DHSprogram.com	
Tutorial Videos – Watch interviews with experts and learn DHS basics, such as sampling and weighting, downloading datasets, and How to Read DHS Tables.	www.youtube.com/DHSProgram	
Datasets – Download DHS datasets for analysis.	DHSprogram.com/Data	

Spatial Data Repository – Download geographically linked health and demographic data for mapping in a geographic information system (GIS).	spatialdata.DHSprogram.com	
Learning Hub – Access online courses for independent learning and workshop participation, communities of practice, and other training resources.	Learning.DHSprogram.com	
GitHub – Open access to Stata, SPSS and R code for DHS indicators for public use.	Github.com/DHSprogram	

YouTube
www.youtube.com/DHSprogram

Blog
Blog.DHSprogram.com



REPUBLIC OF KENYA

KNBS
KENYA NATIONAL
BUREAU OF STATISTICS
Keeping you informed

Kenya National Bureau of Statistics
P.O. Box 30266 - 00100 Nairobi GPO, Kenya

Telephone: +254-20-3317583, +254-20-2911000/1,
+254-20-3317612/22/23/51

Physical Address:
Real Towers, Hospital Road, Upper Hill

Email: info@knbs.or.ke
Website: www.knbs.or.ke
Facebook: Kenya Stats
X-handle: @KNBStats

ISBN 978-9914-9428-7-3
9 789914 942873