

New *Phytologist* Supporting Information

Article title: Tree height and leaf drought tolerance traits shape growth responses across droughts in a temperate broadleaf forest

Authors: Ian R. McGregor, Ryan Helcoski, Norbert Kunert, Alan J. Tepley, Erika B. Gonzalez-Akre, Valentine Herrmann, Joseph Zailaa, Atticus E.L. Stovall, Norman A. Bourg, William J. McShea, Neil Pederson, Lawren Sack, Kristina J. Anderson-Teixeira

Article acceptance date: 02 October 2020

The following Supporting Information is available for this article:

Table S1 . Monthly Palmer Drought Severity Index (PDSI), and its rank among all years between 1950 and 2009, for focal droughts.

Table S2 Species-specific bark thickness regression equations.

Table S3 Species-specific height regression equations.

Table S4 Individual tests of species traits as drivers of drought resistance, where Rt is used as the response variable.

Table S5 Individual tests of species traits as drivers of drought resistance where Rt_{ARIMA} is used as the response variable.

Table S6 Individual tests of species traits as drivers of drought recovery (Rc).

Table S7 Individual tests of species traits as drivers of drought resilience (Rs).

Table S8 Summary of top full models for each drought instance, where Rt is used as the response variable.

Table S9 Summary of top full models for each drought instance, where Rt_{ARIMA} is used as the response variable.

Table S10 Summary of top full models for each drought instance, where Rc is used as the response variable.

Table S11 Summary of top full models for each drought instance, where Rs is used as the response variable.

Fig. S1 Time series of Palmer Drought Severity Index (PDSI) for the 2 years prior and after each focal drought.

Fig. S2 Map of ForestGEO plot showing topographic wetness index and location of cored trees.

Fig. S3 Distribution of reconstructed tree heights across drought years.

Fig. S4 Distribution of independent variables by species.

Fig. S5 Comparison of R_t and R_{tARIMA} results, with residuals, for each drought scenario.

Fig. S6 Density plot of drought recovery (R_c) values for each focal drought year.

Fig. S7 Drought recovery (R_c) across species for the three focal droughts.

Methods S1 Further Package Citations

Table S1 Monthly Palmer Drought Severity Index (PDSI), and its rank among all years between 1950 and 2009 (driest=1), for focal droughts.

year	month	PDSI	rank
1966	May	-2.98	2
	June	-3.40	2
	July	-4.08	2
	August	-4.82	1
1977	May	-2.96	3
	June	-3.28	3
	July	-3.61	3
	August	-3.68	3
1999	May	-3.63	1
	June	-4.21	1
	July	-4.53	1
	August	-4.64	2

Table S2 Species-specific regression equations for bark thickness (mm) as a function of diameter at breast height without bark (mm).

Species	Equations	R^2
<i>Carya cordiformis</i>	$\ln[r_{\text{bark}}] = -1.56 + 0.416 \cdot \ln[\text{DBH}]$	0.226
<i>Carya glabra</i>	$\ln[r_{\text{bark}}] = -0.393 + 0.268 \cdot \ln[\text{DBH}]$	0.04
<i>Carya ovalis</i>	$\ln[r_{\text{bark}}] = -2.18 + 0.651 \cdot \ln[\text{DBH}]$	0.389
<i>Carya tomentosa</i>	$\ln[r_{\text{bark}}] = -0.477 + 0.301 \cdot \ln[\text{DBH}]$	0.297
<i>Fagus grandifolia</i>	-	-
<i>Fraxinus americana</i>	$\ln[r_{\text{bark}}] = 0.418 + 0.26 \cdot \ln[\text{DBH}]$	0.256
<i>Juglans nigra</i>	$\ln[r_{\text{bark}}] = 0.346 + 0.279 \cdot \ln[\text{DBH}]$	0.246
<i>Liriodendron tulipifera</i>	$\ln[r_{\text{bark}}] = -1.14 + 0.463 \cdot \ln[\text{DBH}]$	0.545
<i>Quercus alba</i>	$\ln[r_{\text{bark}}] = -2.09 + 0.637 \cdot \ln[\text{DBH}]$	0.603
<i>Quercus prinus</i>	$\ln[r_{\text{bark}}] = -1.31 + 0.528 \cdot \ln[\text{DBH}]$	0.577
<i>Quercus rubra</i>	$\ln[r_{\text{bark}}] = -0.593 + 0.292 \cdot \ln[\text{DBH}]$	0.101
all	$\ln[r_{\text{bark}}] = 0.245 + 0.219 \cdot \ln[\text{DBH}]$	0.087

We used linear regression on log-transformed data to relate r_{bark} to the diameter inside bark from 2008 data. These were then used to determine r_{bark} in the DBH_Y reconstruction (DBH in year Y). No bark correction was applied for *Fagus grandifolia*, which has thin bark.

Table S3 Species-specific regression equations for height (m) as a function of DBH (cm).

Species	Equations	R^2
Carya cordiformis	$\ln[H] = 0.332 + 0.808 \cdot \ln[DBH]$	0.874
Carya glabra	$\ln[H] = 0.685 + 0.691 \cdot \ln[DBH]$	0.841
Carya ovalis	$\ln[H] = 0.533 + 0.741 \cdot \ln[DBH]$	0.924
Carya tomentosa	$\ln[H] = 0.726 + 0.713 \cdot \ln[DBH]$	0.897
Fagus grandifolia	$\ln[H] = 0.708 + 0.662 \cdot \ln[DBH]$	0.857
Liriodendron tulipifera	$\ln[H] = 1.33 + 0.52 \cdot \ln[DBH]$	0.771
Quercus alba	$\ln[H] = 0.74 + 0.645 \cdot \ln[DBH]$	0.719
Quercus prinus	$\ln[H] = 0.41 + 0.757 \cdot \ln[DBH]$	0.886
Quercus rubra	$\ln[H] = 1.00 + 0.574 \cdot \ln[DBH]$	0.755
all	$\ln[H] = 0.839 + 0.642 \cdot \ln[DBH]$	0.857

Table S4 Individual tests of species traits as drivers of drought resistance, where *Rt* is used as the response variable.

variable	category	all droughts		1966		1977		1999	
		ΔAICc	coefficients	ΔAICc	coefficients	ΔAICc	coefficients	ΔAICc	coefficients
xylem porosity	R	-0.8	0.0630	2.29**	0.190	1.92	-0.152	3.36**	0.1500
	D/SR		0.0000		0.000		0.000		0.0000
<i>PLA</i>		6.7**	-0.0140	9.13**	-0.025	-0.32	-0.010	-0.95	-0.0070
<i>LMA</i>		-2.01	0.0002	-1.9	0.001	-1.68	-0.002	-2.03	0.0003
π_{tlp}		1.33	-0.1740	-1.65	-0.107	1.23	-0.245	-0.1	-0.1690
<i>WD</i>		-1.97	-0.0310	-1.26	-0.206	-1.44	-0.154	0.66	0.2720

Variable abbreviations are as in Table 2. ΔAICc is the AICc of a model excluding the trait minus that of the model including it.

** $\Delta\text{AICc} > 2$: variable considered significant as an individual predictor

Table S5 Individual tests of species traits as drivers of drought resistance, where Rt_{ARIMA} is used as the response variable.

variable	category	all droughts		1966		1977		1999	
		$\Delta AICc$	coefficients	$\Delta AICc$	coefficients	$\Delta AICc$	coefficients	$\Delta AICc$	coefficients
xylem porosity	R	-1.47	0.0420	0.95	0.1520	2.84**	-0.171	2.27**	0.155
	D/SR		0.0000		0.0000		0.000		0.000
<i>PLA</i>		4.48**	-0.0120	10.15**	-0.0240	-0.9	-0.008	-1.67	-0.005
<i>LMA</i>		-1.99	-0.0003	-2.02	0.0005	-0.42	-0.003	-1.9	0.001
π_{tlp}		0.42	-0.1510	-1.94	-0.0530	-0.53	-0.179	0.04	-0.200
<i>WD</i>		-1.94	-0.0390	-0.08	-0.3040	-1.57	-0.142	0.83	0.316

Variable abbreviations are as in Table 2. $\Delta AICc$ is the $AICc$ of a model excluding the trait minus that of the model including it.

** $\Delta AICc > 2$: variable considered significant as an individual predictor

Table S6 Individual test of species traits as drivers of drought recovery (R_c).

variable	category	all droughts		1966		1977		1999	
		$\Delta AICc$	coefficients	$\Delta AICc$	coefficients	$\Delta AICc$	coefficients	$\Delta AICc$	coefficients
xylem porosity	R	15.25**	-0.280	9.9**	-0.474	-1.67	-0.0370	17.06**	-0.3380
	D/SR		0.000		0.000		0.0000		0.0000
<i>PLA</i>		-1.98	0.002	-1.33	0.014	1.10	-0.0090	-2.03	0.0010
<i>LMA</i>		-1.35	-0.002	0.32	-0.008	-2.04	-0.0001	-2.03	-0.0005
π_{tip}		-1.13	-0.149	-1.94	-0.101	1.08	-0.1630	-1.14	-0.2020
<i>WD</i>		-1.86	-0.088	-1.6	0.278	-1.68	-0.0980	-1.03	-0.2950

Variable abbreviations are as in Table 2. $\Delta AICc$ is the AICc of a model excluding the trait minus that of the model including it.

** $\Delta AICc > 2$: variable considered significant as an individual predictor

Table S7 Individual test of species traits as drivers of drought resilience (R_s).

variable	category	all droughts		1966		1977		1999	
		$\Delta AICc$	coefficients	$\Delta AICc$	coefficients	$\Delta AICc$	coefficients	$\Delta AICc$	coefficients
xylem porosity	R	0.24	-0.147	-1.29	-0.110	1.42	-0.263	-1.11	-0.0840
	D/SR		0.000		0.000		0.000		0.0000
<i>PLA</i>		1.09	-0.016	1.09	-0.020	-0.51	-0.017	0.67	-0.0130
<i>LMA</i>		-1.9	-0.001	-1.00	-0.004	-1.95	-0.001	-2.02	-0.0004
π_{tlp}		2.5**	-0.347	-1.11	-0.212	1.57	-0.468	6.11**	-0.3730
<i>WD</i>		-1.83	-0.109	-2.05	-0.020	-1.37	-0.298	-2.02	0.0360

Variable abbreviations are as in Table 2. $\Delta AICc$ is the AICc of a model excluding the trait minus that of the model including it.

** $\Delta AICc > 2$: variable considered significant as an individual predictor

Table S8 Summary of top full models for each drought instance, where Rt is used as the response variable.

drought	$\Delta AICc$	<i>Marginal R^2</i>	<i>Conditional R^2</i>	Intercept	$\ln[H]$	$\ln[TWI]$	$\ln[H] * \ln[TWI]$	PLA	π_{tlp}
all	0.000	0.08	0.12	1.131	-0.057	-0.086	-	-0.012	-0.113
	0.583	0.06	0.11	1.423	-0.055	-0.086	-	-0.013	-
	0.726	0.08	0.12	1.537	-0.202	-0.326	0.082	-0.012	-0.114
	1.352	0.06	0.11	1.826	-0.198	-0.324	0.081	-0.013	-
1966	0.000	0.16	0.25	1.622	-0.135	-	-	-0.025	-
1977	0.000	0.06	0.22	0.503	-	-0.144	-	-	-0.24
	0.908	0.01	0.21	1.069	-	-0.144	-	-	-
	0.988	0.06	0.22	0.568	-0.03	-0.139	-	-	-0.246
	1.144	0.08	0.24	0.684	-	-0.142	-	-0.007	-0.204
	1.267	0.04	0.22	1.211	-	-0.141	-	-0.01	-
1999	0.000	0.01	0.18	1.061	-	-0.102	-	-	-
	0.023	0.04	0.19	0.659	-	-0.101	-	-	-0.169
	0.954	0.02	0.19	1.157	-	-0.1	-	-0.007	-
	1.513	0.05	0.21	0.783	-	-0.1	-	-0.005	-0.145
	1.803	0.01	0.18	1.024	0.013	-0.103	-	-	-
	1.901	0.04	0.19	0.635	0.011	-0.102	-	-	-0.166

Models are ranked by AICc. Shown are all models whose AICc value falls within 2.0 ($\Delta AICc < 1$) of the best model (bold). R^2 refers to conditional R^2 . Year was included in the model for all drought years, but its effect was not included in any top models, and coefficients were small (1966: 0, 1977: -0.019, 1999: -0.005; same values in all top models).

Table S9 Summary of top full models for each drought instance, where Rt_{ARIMA} is used as the response variable.

drought	$\Delta AICc$	Marginal R^2	Conditional R^2	Intercept	$\ln[H]$	$\ln[TWI]$	$\ln[H] * \ln[TWI]$	PLA	π_{tlp}
all	0.000	0.05	0.09	2.113	-0.307	-0.506	0.14	-0.012	-
	0.419	0.06	0.10	1.872	-0.31	-0.508	0.141	-0.011	-0.096
	1.217	0.05	0.09	1.395	-0.06	-0.1	-	-0.012	-
	1.698	0.06	0.10	1.153	-0.062	-0.1	-	-0.011	-0.095
1966	0.000	0.17	0.23	1.660	-0.154	-	-	-0.024	-
	1.393	0.17	0.23	1.735	-0.152	-0.047	-	-0.024	-
	1.457	0.16	0.23	1.859	-0.152	-	-	-0.025	0.078
1977	0.000	0.01	0.16	1.130	-	-0.18	-	-	-
	0.424	0.02	0.16	2.453	-0.461	-0.896	0.25	-	-
	0.688	0.03	0.17	0.720	-	-0.179	-	-	-0.173
	0.922	0.04	0.17	2.040	-0.466	-0.898	0.251	-	-0.18
	0.927	0.03	0.17	1.248	-	-0.177	-	-0.008	-
	1.322	0.03	0.17	2.569	-0.461	-0.893	0.25	-0.008	-
	1.709	0.01	0.15	1.183	-0.02	-0.177	-	-	-
1999	0.000	0.04	0.20	0.563	-	-0.076	-	-	-0.2
	0.064	0.03	0.19	0.421	-	-	-	-	-0.202
	0.127	0.00	0.18	1.036	-	-0.077	-	-	-
	0.256	0.00	0.18	0.899	-	-	-	-	-
	1.777	0.04	0.20	0.529	0.016	-0.078	-	-	-0.195
	1.797	0.01	0.20	1.101	-	-0.076	-	-0.004	-
	1.815	0.00	0.18	0.986	0.018	-0.079	-	-	-
	1.838	0.01	0.20	0.972	-	-	-	-0.005	-
	1.933	0.03	0.19	0.391	0.012	-	-	-	-0.199
	1.979	0.04	0.21	0.612	-	-0.075	-	-0.002	-0.19
	1.999	0.04	0.21	0.482	-	-	-	-0.002	-0.19

Models are ranked by AICc. Shown are all models whose AICc value falls within 2.0 ($\Delta AICc < 1$) of the best model (bold). R^2 refers to conditional R^2 . Year was included in the model for all drought years and appeared in all its top models, but coefficients were small (1966: 0, 1977: -0.03, 1999: 0.008; same values in all top models).

Table S10 Summary of top full models for each drought instance, where R_c is used as the response variable.

drought	$\Delta AICc$	Marginal R^2	Conditional R^2	Intercept	$\ln[H]$	$\ln[TWT]$	$\ln[H] * \ln[TWT]$	PLA	π_{tlp}
all	0.000	0.05	0.17	0.434	0.345	0.844	-0.269	-	-
	0.995	0.05	0.17	1.913	-0.126	-	-	-	-
	1.135	0.06	0.17	0.077	0.344	0.845	-0.269	-	-0.152
	1.991	0.05	0.18	0.410	0.346	0.843	-0.269	0.002	-
1966	0.000	0.01	0.28	-0.797	0.89	1.263	-0.475	-	-
	1.040	0.00	0.25	1.577	-	-	-	-	-
	1.367	0.02	0.30	-0.984	0.888	1.257	-0.474	0.013	-
	1.785	0.00	0.26	1.781	-	-0.114	-	-	-
	1.956	0.01	0.30	-1.025	0.89	1.261	-0.475	-	-0.097
1977	0.000	0.17	0.17	2.485	-0.482	-	-	-	-0.157
	0.299	0.17	0.17	2.943	-0.47	-	-	-0.008	-
	0.716	0.17	0.18	2.657	-0.477	-	-	-0.006	-0.114
	0.807	0.17	0.18	1.152	0.071	1.026	-0.308	-0.009	-
	0.875	0.17	0.18	2.729	-0.47	0.124	-	-0.009	-
	0.891	0.17	0.18	2.271	-0.479	0.115	-	-	-0.158
	0.910	0.17	0.18	0.712	0.054	1.004	-0.304	-	-0.159
	1.315	0.17	0.18	0.871	0.065	1.023	-0.308	-0.006	-0.112
	1.331	0.16	0.17	2.805	-0.464	-	-	-	-
	1.372	0.17	0.18	2.445	-0.475	0.122	-	-0.006	-0.112
	1.974	0.16	0.17	2.597	-0.466	0.118	-	-	-
1999	0.000	0.00	0.16	1.281	-	-	-	-	-
	0.532	0.00	0.17	1.093	-	0.105	-	-	-
	1.091	0.02	0.19	0.779	-	-	-	-	-0.212
	1.609	0.02	0.19	0.578	-	0.106	-	-	-0.217
	1.755	0.00	0.17	1.200	0.027	-	-	-	-
	1.996	0.00	0.18	1.251	-	-	-	0.002	-

Models are ranked by AICc. Shown are all models whose AICc value falls within 2.0 ($\Delta AICc < 1$) of the best model (bold). R^2 refers to conditional R^2 . Year was included in the model for all drought years and appeared in all its top models (1966: 0, 1977: -0.14, 1999: -0.217; same values in all top models).

Table S11 Summary of top full models for each drought instance, where R_s is used as the response variable.

drought	$\Delta AICc$	<i>Marginal R^2</i>	<i>Conditional R^2</i>	Intercept	$\ln[H]$	$\ln[TWT]$	$\ln[H] * \ln[TWT]$	PLA	π_{tlp}
all	0.000	0.10	0.17	-0.265	0.348	0.864	-0.291	-0.012	-0.287
	0.176	0.08	0.16	-0.572	0.347	0.859	-0.291	-	-0.347
	1.518	0.07	0.16	0.458	0.354	0.866	-0.292	-0.016	-
	1.552	0.09	0.17	1.253	-0.166	-	-	-0.011	-0.288
	1.698	0.08	0.16	0.940	-0.166	-	-	-	-0.348
1966	0.000	0.04	0.15	1.834	-0.085	-	-	-0.02	-
	0.402	0.03	0.16	1.589	-	-	-	-0.02	-
	1.189	0.00	0.14	1.534	-0.082	-	-	-	-
	1.313	0.00	0.15	1.293	-	-	-	-	-
	1.692	0.04	0.16	1.534	-0.085	-	-	-0.018	-0.116
1977	0.000	0.14	0.28	-0.932	0.294	1.207	-0.384	-	-0.467
	0.497	0.13	0.28	1.194	-0.383	-	-	-	-0.469
	1.304	0.15	0.30	-0.648	0.294	1.208	-0.383	-0.011	-0.411
	1.542	0.13	0.28	1.026	-0.387	0.095	-	-	-0.472
	1.555	0.09	0.28	0.138	0.304	1.211	-0.385	-	-
	1.852	0.14	0.29	1.467	-0.381	-	-	-0.01	-0.416
1999	0.000	0.07	0.13	0.237	-	-	-	-	-0.366
	0.313	0.08	0.14	0.472	-	-	-	-0.008	-0.317
	0.503	0.07	0.13	0.358	-0.048	-	-	-	-0.376
	0.532	0.07	0.13	0.394	-	-0.086	-	-	-0.364
	0.726	0.09	0.14	0.588	-0.047	-	-	-0.008	-0.328
	1.079	0.09	0.15	0.602	-	-0.081	-	-0.008	-0.319
	1.249	0.07	0.13	0.495	-0.044	-0.08	-	-	-0.374
	1.706	0.09	0.14	0.699	-0.044	-0.075	-	-0.007	-0.329

Models are ranked by AICc. Shown are all models whose AICc value falls within 2.0 ($\Delta AICc < 1$) of the best model (bold). R^2 refers to conditional R^2 . Year was included in the model for all drought years and appeared in all its top models (1966: 0, 1977: -0.099, -0.099, -0.099, -0.097, -0.097; 1999: -0.174, -0.174, -0.174, -0.173, -0.172).

Fig. S1 Time series of Palmer Drought Severity Index (PDSI) for each focal drought year.




Fig. S2 Map of ForestGEO plot showing topographic wetness index and location of cored trees. Scale units are in meters.




Fig. S3 Distribution of reconstructed tree heights across drought years.




Fig. S4 Distribution of independent variable values by species. Species codes are given in Table 2. Boxes in plots (a) – (f) represent the interquartile range, with the horizontal line at the median, whiskers representing the range within 2.7 SD, and dots representing outliers. For plots (a) – (f), species that are assigned the same letter are not significantly different from each other with regard to the tested variable. Similarly, letter groupings do not transfer between variables. Meanwhile, plot (g) shows the number of trees in each crown position per species. Descriptions of variables (e.g. $\ln[H]$) can be found in Table 3.




Fig. S5 Comparisons of R_t and R_{tARIMA} results, with residuals, for each drought scenario.




Fig. S6 Density plot of drought recovery (R_c) for each focal drought year.





Fig. S7 Drought recovery (R_c) across species for the three focal droughts. Species codes are given in Table 2. Shaded boxes represent the interquartile range, with horizontal line at median, whiskers represent the range within 2.7 SD, and dots represent outliers. The dotted line at $y=1$ represents no change in R_c from the five years prior to drought. Letters illustrate significance groupings per year (colored and ordered, top to bottom, 1966, 1977, 1999). That is, a group of species with the same letter above their boxplot (e.g. "b") are statistically different from species in another group (e.g. "a"). See Fig. 4 for parallel plot for resistance (R_t) and resilience (R_s). Analysis conducted using *agricolae* package in R.



Methods S1 Further Package Citations

While there were several R-packages we used for a specific purpose in our methods, numerous packages were immensely helpful for this research behind the scenes. As in all of science, this study is a representation of the work done by both the authors of this paper as well as countless others. While acknowledging everyone is impossible, we want to at least give thanks to those who made this work possible.

R-packages not already cited in the main manuscript include the following, listed alphabetically by corresponding package name:

R base (R Core Team, 2019); broom (Robinson & Hayes, 2020); car (Fox *et al.*, 2019); cowplot (Wilke, 2019); data.table (Dowle & Srinivasan, 2019); devtools (Wickham *et al.*, 2020b); dplR (Bunn *et al.*, 2019); dplyr (Wickham *et al.*, 2020a); extrafont (Winston Chang, 2014); ggplot2 (Wickham *et al.*, 2019); ggpubr (Kassambara, 2020); ggthemes (Arnold, 2019); gridExtra (Auguie, 2017); knitr (Xie, 2020); lubridate (Spinu *et al.*, 2018); MuMIn (Barton, 2019); piecewiseSEM (Lefcheck *et al.*, 2019); png (Urbanek, 2013); purrr (Henry & Wickham, 2019); raster (Hijmans, 2020); rasterVis (Perpinan Lamigueiro & Hijmans, 2019); RCurl (Temple Lang, 2020); readxl (Wickham & Bryan, 2019); reshape2 (Wickham, 2017); rgdal (Bivand *et al.*, 2019); rgeos (Bivand & Rundel, 2019); rmarkdown (Allaire *et al.*, 2020); sf (Pebesma, 2020); stringi (Gagolewski *et al.*, 2020); stringr (Wickham, 2019); tidyr (Wickham & Henry, 2020)

Allaire, JJ, Yihui Xie, Jonathan McPherson, Javier Luraschi, Kevin Ushey, Aron Atkins, Hadley Wickham, Joe Cheng, Winston Chang, and Richard Iannone. 2020. *Rmarkdown: Dynamic Documents for R*. <https://CRAN.R-project.org/package=rmarkdown>.

Arnold, Jeffrey B. 2019. *Ggthemes: Extra Themes, Scales and Geoms for 'Ggplot2'*. <https://CRAN.R-project.org/package=ggthemes>.

Auguie, Baptiste. 2017. *GridExtra: Miscellaneous Functions for "Grid" Graphics*. <https://CRAN.R-project.org/package=gridExtra>.

Barton, Kamil. 2019. *MuMIn: Multi-Model Inference*. <https://CRAN.R-project.org/package=MuMIn>.

Bivand, Roger, Tim Keitt, and Barry Rowlingson. 2019. *Rgdal: Bindings for the 'Geospatial' Data Abstraction Library*. <https://CRAN.R-project.org/package=rgdal>.

Bivand, Roger, and Colin Rundel. 2019. *Rgeos: Interface to Geometry Engine - Open Source ('Geos')*. <https://CRAN.R-project.org/package=rgeos>.

Bunn, Andy, Mikko Korpela, Franco Biondi, Filipe Campelo, Pierre Mérian, Fares Qeadan, and Christian Zang. 2019. *DplR: Dendrochronology Program Library in R*. <https://CRAN.R-project.org/package=dplR>.

- Dowle, Matt, and Arun Srinivasan. 2019. *Data.table: Extension of 'Data.frame'*. <https://CRAN.R-project.org/package=data.table>.
- Fox, John, Sanford Weisberg, and Brad Price. 2019. *Car: Companion to Applied Regression*. <https://CRAN.R-project.org/package=car>.
- Gagolewski, Marek, Bartek Tartanus, other contributors; IBM, Unicode, Inc., other contributors; Unicode, and Inc. 2020. *Stringi: Character String Processing Facilities*. <https://CRAN.R-project.org/package=stringi>.
- Henry, Lionel, and Hadley Wickham. 2019. *Purrr: Functional Programming Tools*. <https://CRAN.R-project.org/package=purrr>.
- Hijmans, Robert J. 2020. *Raster: Geographic Data Analysis and Modeling*. <https://CRAN.R-project.org/package=raster>.
- Kassambara, Alboukadel. 2020. *Ggpubr: 'Ggplot2' Based Publication Ready Plots*. <https://CRAN.R-project.org/package=ggpubr>.
- Lefcheck, Jon, Jarrett Byrnes, and James Grace. 2019. *PiecewiseSEM: Piecewise Structural Equation Modeling*. <https://CRAN.R-project.org/package=piecewiseSEM>.
- Pebesma, Edzer. 2020. *Sf: Simple Features for R*. <https://CRAN.R-project.org/package=sf>.
- Perpinan Lamigueiro, Oscar, and Robert Hijmans. 2019. *RasterVis: Visualization Methods for Raster Data*. <https://CRAN.R-project.org/package=rasterVis>.
- R Core Team. 2019. *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing. <https://www.R-project.org/>.
- Robinson, David, and Alex Hayes. 2020. *Broom: Convert Statistical Analysis Objects into Tidy Tibbles*. <https://CRAN.R-project.org/package=broom>.
- Spinu, Vitalie, Garrett Grolemond, and Hadley Wickham. 2018. *Lubridate: Make Dealing with Dates a Little Easier*. <https://CRAN.R-project.org/package=lubridate>.
- Temple Lang, Duncan. 2020. *RCurl: General Network (Http/Ftp/...) Client Interface for R*. <https://CRAN.R-project.org/package=RCurl>.
- Urbanek, Simon. 2013. *Png: Read and Write Png Images*. <https://CRAN.R-project.org/package=png>.
- Wickham, Hadley. 2017. *Reshape2: Flexibly Reshape Data: A Reboot of the Reshape Package*. <https://CRAN.R-project.org/package=reshape2>.
- . 2019. *Stringr: Simple, Consistent Wrappers for Common String Operations*. <https://CRAN.R-project.org/package=stringr>.
- Wickham, Hadley, and Jennifer Bryan. 2019. *Readxl: Read Excel Files*. <https://CRAN.R-project.org/package=readxl>.

Wickham, Hadley, Winston Chang, Lionel Henry, Thomas Lin Pedersen, Kohske Takahashi, Claus Wilke, Kara Woo, and Hiroaki Yutani. 2019. *Ggplot2: Create Elegant Data Visualisations Using the Grammar of Graphics*. <https://CRAN.R-project.org/package=ggplot2>.

Wickham, Hadley, Romain François, Lionel Henry, and Kirill Müller. 2020. *Dplyr: A Grammar of Data Manipulation*. <https://CRAN.R-project.org/package=dplyr>.

Wickham, Hadley, and Lionel Henry. 2020. *Tidyr: Tidy Messy Data*. <https://CRAN.R-project.org/package=tidyr>.

Wickham, Hadley, Jim Hester, and Winston Chang. 2020. *Devtools: Tools to Make Developing R Packages Easier*. <https://CRAN.R-project.org/package=devtools>.

Wilke, Claus O. 2019. *Cowplot: Streamlined Plot Theme and Plot Annotations for 'Ggplot2'*. <https://CRAN.R-project.org/package=cowplot>.

Winston Chang. 2014. *Extrafont: Tools for Using Fonts*. <https://CRAN.R-project.org/package=extrafont>.

Xie, Yihui. 2020. *Knitr: A General-Purpose Package for Dynamic Report Generation in R*. <https://CRAN.R-project.org/package=knitr>.