

**Title:** Tree height and leaf drought tolerance traits shape growth responses across droughts in a temperate broadleaf forest

**Authors:** Ian R. McGregor<sup>1,2</sup>, Ryan Helcoski<sup>1</sup>, Norbert Kunert<sup>1,3</sup>, Alan J. Tepley<sup>1,4</sup>, Erika B. Gonzalez-Akre<sup>1</sup>, Valentine Herrmann<sup>1</sup>, Joseph Zailaa<sup>1,5</sup>, Atticus E.L. Stovall<sup>1,6,7</sup>, Norman A. Bourg<sup>1</sup>, William J. McShea<sup>1</sup>, Neil Pederson<sup>8</sup>, Lawren Sack<sup>9,10</sup>, Kristina J. Anderson-Teixeira<sup>1,3\*</sup>

**Author Affiliations:**

1. Conservation Ecology Center; Smithsonian Conservation Biology Institute; National Zoological Park, Front Royal, VA 22630, USA
2. Center for Geospatial Analytics; North Carolina State University; Raleigh, NC 27607, USA
3. Center for Tropical Forest Science-Forest Global Earth Observatory; Smithsonian Tropical Research Institute; Panama, Republic of Panama
4. Canadian Forest Service, Northern Forestry Centre, Edmonton, Alberta, Canada
5. Biological Sciences Department; California State University; Los Angeles, CA 90032, USA
6. Department of Environmental Sciences, University of Virginia, Charlottesville, VA 22903, USA
7. NASA Goddard Space Flight Center; Greenbelt, MD 20771, USA
8. Harvard Forest, Petersham, MA 01366, USA
9. Department of Ecology and Evolutionary Biology; University of California, Los Angeles; Los Angeles, CA 90095, USA
10. Institute of the Environment and Sustainability; University of California, Los Angeles; Los Angeles, CA 90095, USA

\*corresponding author: teixeirak@si.edu; +1 540 635 6546

Text	word count	other	n
Total word count (excluding summary, references and legends)	5,365	No. of figures	4 (all colour)
Summary	198	No. of Tables	3
Introduction	1,034	No of Supporting Information files	#
Materials and Methods	1,945		
Results	697		
Discussion	1467		
Acknowledgements	125		

## Summary

- As climate change is driving increased drought frequency and severity in many forested regions around the world, mechanistic understanding of the factors conferring drought resistance in trees is increasingly important. The dendrochronological record provides a window through which we can understand how tree size and species' traits shape tree growth responses during droughts.
- We analyzed tree-ring records for twelve species that comprise 97% of the woody productivity of the 25.6-ha ForestGEO plot in a broadleaf deciduous forest of northern Virginia (USA) to test hypotheses on how tree height, microenvironment characteristics, and species' traits shaped drought responses across the three strongest regional droughts over a 60-year period (1950 - 2009).
- Individual-level drought resistance decreased with tree height, which was strongly correlated with exposure to higher evaporative demand and solar radiation. The potentially greater rooting volume of larger trees did not confer an advantage in sites with low topographic wetness index. Resistance was greater among species whose leaves experienced less shrinkage upon desiccation and lost turgor (wilted) at more negative water potentials.
- We conclude that tree height and leaf drought tolerance traits influence growth responses during drought, as recorded in the tree-ring record spanning historical droughts. Thus, these factors can be useful for predicting future drought responses under climate change.

*Key words:* annual growth; crown exposure; drought; Forest Global Earth Observatory (ForestGEO); leaf drought tolerance traits; temperate broadleaf deciduous forest; tree height; tree-ring

## 41 Introduction

42 Forests play a critical global role in climate regulation (Bonan, 2008), yet there remains enormous  
43 uncertainty as to how the forest-dominated terrestrial carbon sink will respond to climate change  
44 (Friedlingstein et al., 2006). An important aspect of this uncertainty lies with physiological responses of  
45 trees to drought (Kennedy et al., 2019). In many forested regions around the world, the risk of severe  
46 drought is increasing (Trenberth et al., 2014; Dai et al., 2018), often despite increasing precipitation  
47 (Intergovernmental Panel on Climate Change, 2015; Cook et al., 2015). Droughts, intensified by climate  
48 change, have been affecting forests worldwide and are expected to continue as one of the most important  
49 drivers of forest change in the future (Allen et al., 2010, 2015). Understanding forest responses to drought  
50 requires elucidation of how tree size, microenvironment, and species’ traits jointly influence individual-level  
51 drought resistance, defined here as a tree’s ability to maintain growth during drought, and the extent to  
52 which their influence is consistent across droughts. Because the resistance and resilience (i.e., post-drought  
53 recovery) of growth to drought are linked to trees’ probability of surviving drought (DeSoto et al., 2020;  
54 Liu et al., 2019), understanding growth responses can also help elucidate which trees are most vulnerable  
55 to drought-induced mortality. However, it has proven difficult to resolve the many factors affecting tree  
56 growth during drought with available forest census data, which only rarely captures extreme drought, and  
57 with tree-ring records, which capture multiple droughts but usually only sample a subset of a forest  
58 community, typically focusing on a single species or the largest individuals.

59 Many studies have shown that within and across species, large trees tend to be more affected by drought.  
60 Greater growth reductions for larger trees were first shown on a global scale by Bennett et al. (2015), and  
61 subsequent studies have reinforced this finding (*e.g.*, Hacket-Pain et al., 2016; Gillerot et al., 2020). It has  
62 yet to be resolved which of several potential underlying mechanisms most strongly shape these trends in  
63 drought response. First, tree height itself may be a primary driver. Taller trees face the biophysical  
64 challenge of lifting water greater distances against the effects of gravity and friction (McDowell et al., 2011;  
65 McDowell and Allen, 2015; Ryan et al., 2006; Couvreur et al., 2018). Vertical gradients in stem and leaf  
66 traits—including smaller and thicker leaves (higher leaf mass per area, LMA), greater resistance to  
67 hydraulic dysfunction (*i.e.*, more negative water potential at 50% loss of hydraulic conductivity, more  
68 negative P50), and lower hydraulic conductivity at greater heights (Couvreur et al., 2018; Koike et al.,  
69 2001; McDowell et al., 2011)—enable trees to become tall (Couvreur et al., 2018). Greater stem capacitance  
70 (*i.e.*, water storage capacity) of larger trees may also confer resistance to transient droughts [Phillips et al.  
71 (2003)\*; Scholz et al. (2011)]. Indeed, tall trees require xylem of greater hydraulic efficiency, such that  
72 xylem conduit diameters are wider in the basal portions of taller trees, both within and across species  
73 (Olson et al., 2018; Liu et al., 2019), and throughout the conductive systems of angiosperms (Zach et al.,  
74 2010; Olson et al., 2014, 2018). Wider xylem conduits plausibly make large trees more vulnerable to  
75 embolism during drought (Olson et al., 2018), and traits conducive to efficient water transport may also  
76 lead to poor ability to recover from or re-route water around embolisms (Roskilly et al., 2019).

77 Larger trees may also have lower drought resistance because of microenvironmental and ecological factors.  
78 Their crowns tend to occupy more exposed canopy positions, which are associated with higher evaporative  
79 demand (Kunert et al., 2017). Subcanopy trees tend to fare better specifically due to the benefits of a  
80 buffered environment (Pretzsch et al., 2018). Counteracting the liabilities associated with tall height, large  
81 trees tend to have larger root systems (Enquist and Niklas, 2002), potentially mitigating some of the  
82 biophysical challenges they face by allowing greater access to water. Larger root systems—if they grant

access to deeper water sources—would be particularly advantageous in drier microenvironments (e.g., hilltops, as compared to valleys and streambeds) during drought. Finally, tree size-related responses to drought can be modified by species’ traits and their distribution across size classes (Meakem et al., 2018; Liu et al., 2019). Understanding the mechanisms driving the greater relative growth reductions of larger trees during drought requires sorting out the interactive effects of height and associated exposure, root water access, and species’ traits.

Debates have also arisen regarding the traits influencing tree growth responses to drought. Studies within temperate broadleaf forests have observed ring-porous species showing higher drought tolerance than diffuse-porous species (Friedrichs et al., 2009; Elliott et al., 2015; Kannenberg et al., 2019), but this distinction would not hold in the global context (Wheeler et al., 2007; Olson et al., 2020) and does not resolve differences among the many species within each category. Commonly-measured traits including wood density and leaf mass per area (*LMA*) have been linked to drought responses within some temperate deciduous forests (Abrams, 1990; Guerfel et al., 2009; Hoffmann et al., 2011; Martin-Benito and Pederson, 2015) and across forests worldwide (Greenwood et al., 2017). However, in other cases these traits could not explain drought tolerance (e.g., in a tropical rainforest; Maréchaux et al., 2019), or the direction of response was not always consistent. For instance, higher wood density has been associated with greater drought resistance at a global scale (Greenwood et al., 2017), but correlated negatively with tree performance during drought in a broadleaf deciduous forest in the southeastern United States (Hoffmann et al., 2011). Thus, the perceived influence of these traits on drought resistance may actually reflect indirect correlations with other traits that more directly drive drought responses (Hoffmann et al., 2011).

In contrast, hydraulic traits have direct physiological linkages to tree growth and mortality responses to drought. For instance, water potentials at which percent the loss of conductivity surpasses a certain threshold (e.g., P50 and P88, representing 50 and 88% loss of conductivity, respectively) and hydraulic safety margin (*i.e.*, difference between typical minimum water potentials and P50 or P88) correlate with drought performance across global forests (Anderegg et al., 2016). However, these are time-consuming to measure and therefore infeasible for predicting or modeling drought responses in highly diverse forests (*e.g.*, in the tropics). More easily-measurable leaf drought tolerance traits that have direct linkage to plant hydraulic function can explain variation in plant distribution and function (Medeiros et al., 2019). These include leaf area shrinkage upon desiccation ( $PLA_{dry}$ ; Scoffoni et al., 2014) and the leaf water potential at turgor loss point ( $\pi_{tlp}$ ), *i.e.*, the water potential at which leaf wilting occurs (Bartlett et al., 2016a; Zhu et al., 2018). Both traits correlate with hydraulic vulnerability and drought tolerance as part of unified plant hydraulic systems (Scoffoni et al., 2014; Bartlett et al., 2016a; Zhu et al., 2018; Farrell et al., 2017). The abilities of both  $PLA_{dry}$  and  $\pi_{tlp}$  to explain tree drought resistance remains untested.

Here, we examine how tree height, microenvironment characteristics, and species’ traits collectively shape drought resistance, defined as the ratio of annual growth in a drought year to that which would be expected in the absence of drought based on previous years’ growth. We test a series of hypotheses and associated specific predictions (Table 1) based on the combination of tree-ring records from the three strongest droughts over a 60-year period (1950 - 2009), species trait measurements, and census and microenvironmental data from a large forest dynamics plot in Virginia, USA. First, we focus on how tree size, alone and in its interaction with microenvironmental gradients, influences drought resistance. We examine the contemporary relationship between tree height and microenvironment, including growing season meteorological conditions and crown exposure. We then test whether, consistent with most forests

globally, larger-diameter, taller trees tend to have lower drought resistance in this forest, which is in a region (eastern North America) represented by only two studies in the global review of Bennett et al. (2015). We also test for an influence of potential access to available soil water, which should be greater for larger trees in dry but not in perpetually wet microsites. Finally, we focus on the role of species' traits, testing the hypothesis that species' traits—particularly leaf drought tolerance traits—predict drought resistance. We test predictions that drought resistance is higher in ring-porous than semi-ring and diffuse-porous species and that it is correlated with wood density—either positively (Greenwood et al., 2017) or negatively (Hoffmann et al., 2011) and positively correlated with *LMA*. We further test predictions that species with low *PLA<sub>dry</sub>* have higher drought resistance, and that species whose leaves lose turgor lower water potentials (more negative  $\pi_{tlp}$ ) have higher resistance.

## Materials and Methods

### *Study site and microclimate*

Research was conducted at the 25.6-ha ForestGEO (Forest Global Earth Observatory) study plot at the Smithsonian Conservation Biology Institute (SCBI) in Virginia, USA (38°53'36.6"N, 78°08'43.4"W; Fig. S1) (Bourg et al., 2013; Anderson-Teixeira et al., 2015a). SCBI is located in the central Appalachian Mountains near the northern boundary of Shenandoah National Park. Elevations range from 273 to 338 m above sea level with a topographic relief of 65m (Bourg et al., 2013). Climate is humid temperate, with mean annual temperature of 12.7°C and precipitation of 1005 mm yr<sup>-1</sup> during our study period (1960-2009; source: CRU TS v.4.01; Harris et al., 2014). Dominant tree taxa within this secondary forest include *Liriodendron tulipifera*, oaks (*Quercus* spp.), and hickories (*Carya* spp.; Table 2).

### *Identifying drought years*

We identified the three largest droughts within the time period 1950-2009, defining drought (Slette et al., 2019) as events with anomalously dry peak growing season climatic conditions. Specifically, we used the metric of Palmer Drought Severity Index (PDSI) during May-August (MJJA; Table S1), which were identified by Helcoski et al. (2019) as the months of the current year to which annual tree growth was most sensitive at this site. PDSI divisional data for Northern Virginia were obtained from NOAA (<https://www7.ncdc.noaa.gov/CDO/CDODivisionalSelect.jsp>) in December 2017. Based on this, we identified the three strongest droughts during the study period (Figs. 1, S1; Table S1).

The droughts differed in intensity and antecedent moisture conditions (Fig. S1, Table S1). The 1966 drought was preceded by two years of moderate drought during the growing season and severe to extreme drought starting the previous fall. In August 1966, *PDSI* reached its lowest monthly value (-4.82) of the three droughts. The 1977 drought was the least intense throughout the growing season, and it was preceded by 2.5 years of near-normal conditions, making it the mildest of the three droughts. The 1999 drought was preceded by wetter than average conditions until the previous June, but *PDSI* plummeted below -3.0 in October 1998 and remained below this threshold through August 1999.

### *Data collection and preparation*

Within or just outside the ForestGEO plot, we collected data on a suite of variables including tree heights, microenvironment characteristics, and species traits (Table 3). The SCBI ForestGEO plot was censused in 2008, 2013, and 2018 following standard ForestGEO protocols, whereby all free-standing woody stems  $\geq$

1cm diameter at breast height (DBH) were mapped, tagged, measured at DBH, and identified to species (Condit, 1998). From these census data, we used measurements of DBH from 2008 to calculate historical DBH and data for all stems  $\geq 10$ cm to analyze functional trait composition relative to tree height (all analyses described below). Census data are available through the ForestGEO data portal ([www.forestgeo.si.edu](http://www.forestgeo.si.edu)).

We analyzed tree-ring data (xylem growth increment) from 571 trees representing the twelve dominant species (Table 2; Fig. **S2**). Selected species were those with the greatest contributions to woody aboveground net primary productivity ( $ANPP_{stem}$ ) and together comprised 97% of study plot  $ANPP_{stem}$  between 2008 and 2013 (Helcoski et al., 2019). Cores (one per tree) were collected within the ForestGEO plot at breast height (1.3m) in 2010-2011 or 2016-2017. In 2010-2011, cores were collected from randomly selected live trees of each species that had at least 30 individuals  $\geq 10$  cm DBH (Bourg et al., 2013). Annual tree mortality censuses were initiated in 2014 (Gonzalez-Akre et al., 2016), and in 2016-2017, cores were collected from all trees found to have died since the previous year’s census. We note that drought was probably not a cause of mortality for these trees, as monthly May-Aug  $PDSI$  did not drop below -1.75 in these years or the three years prior (2013-2017), and that trees cored dead displayed similar climate sensitivity to trees cored live (Helcoski et al., 2019). Cores were sanded, measured, and crossdated using standard procedures, as detailed in (Helcoski et al., 2019). The resulting chronologies (Fig. **1a**) were published in Zenodo (DOI: 10.5281/zenodo.2649302) in association with Helcoski et al. (2019).

For each cored tree, we combined tree-ring records and allometric equations of bark thickness to reconstruct DBH for the years 1950-2009. Prior  $DBH$  was estimated using the following equation:

$$DBH_Y = DBH_{2008} - 2 * \left[ r_{bark,2008} - r_{bark,Y} + \sum_{year=Y}^{2008} r_{ring,Y} \right]$$

Here,  $Y$  denotes the year of interest,  $r_{ring}$  denotes ring width derived from cores, and  $r_{bark}$  denotes bark thickness. Bark thickness was estimated from species-specific allometries based on the bark thickness data from the site (Anderson-Teixeira et al., 2015b). Specifically, we used linear regression on log-transformed data to relate  $r_{bark}$  to diameter inside bark from 2008 data (Table S2), which were then used to determine  $r_{bark}$  in the  $DBH$  reconstruction.

Tree heights ( $H$ ) were measured by several researchers for a variety of purposes between 2012 and 2019 (n=1,518 trees). Methods included direct measurements using a collapsible measurement rod on small trees (NEON, 2018) or a tape measure on recently fallen trees (this study); geometric calculations using clinometer and tape measure (Stovall et al., 2018a) or digital rangefinders (Anderson-Teixeira et al., 2015b; NEON, 2018); and ground-based LiDAR (Stovall et al., 2018b). Rangefinders used either the tangent method (Impulse 200LR, TruPulse 360R) or the sine method (Nikon ForestryPro) for calculating heights. Both methods are associated with some error (Larjavaara and Muller-Landau, 2013), but in this instance there was no clear advantage of one or the other. Measurements from the National Ecological Observatory Network (NEON) were collected near the ForestGEO plot following standard NEON protocol, whereby vegetation of short stature was measured with a collapsible measurement rod, and taller trees with a rangefinder (NEON, 2018). Species-specific height allometries were developed (Table S3) using log-log regression ( $\ln[H] \sim \ln[DBH]$ ). For species with insufficient height data to create reliable species-specific allometries (n=2, JUNI and FRAM), heights were calculated from an equation developed by combining the

height measurements across all species. We then used these allometries to estimate  $H$  for each drought year,  $Y$ , based on reconstructed  $DBH_Y$ . The distribution of  $H$  across drought years is shown in Fig. **S3**.

To characterize how environmental conditions vary with height, data were obtained from the NEON tower located <1km from the study area via the neonUtilities package (Lunch et al., 2020). We used wind speed, relative humidity, and air temperature data, all measured over a vertical profile spanning heights from 7.2 m to above the top of the tree canopy (31.0 or 51.8m, depending on censor), for the years 2016-2018 (NEON, 2018). After filtering for missing and outlier values, we determined the daily minima and maxima, which we then aggregated at the monthly scale.

Crown position—a categorical variable classifying trees based on exposure to sunlight—was recorded for all cored trees that remained standing during the growing season of 2018 following the protocol of Jennings et al. (1999). Trees were classified as follows: *dominant* trees were defined as those with crowns above the general level of the canopy, *co-dominant* trees as those with crowns within the the canopy; *intermediate* trees as those with crowns below the canopy level, but illuminated from above; and *suppressed* as those below the canopy and receiving minimal direct illumination from above.

Topographic wetness index (TWI), used here as a metric of long-term mean moisture availability, was calculated using the dynatopmodel package in R (Fig. **S2**) (Metcalf et al., 2018). Originally developed by Beven and Kirkby (1979), TWI was part of a hydrological run-off model and has since been used for a number of purposes in hydrology and ecology (Sørensen et al., 2006). TWI calculation depends on an input of a digital elevation model (DEM; ~3.7 m resolution from the elevatr package (Hollister, 2018)), and from this yields a quantitative assessment defined by how “wet” an area is, based on areas where run-off is more likely. From our observations in the plot, TWI performed better at categorizing wet areas than the Euclidean distance from the stream.

Species’ trait data were collected in August 2018 (Tables 2-3; Fig. **S4**). We sampled small, sun-exposed branches up to eight meters above the ground from three individuals of each species in and around the ForestGEO plot. Sampled branches were re-cut under water at least two nodes above the original cut and re-hydrated overnight in covered buckets under opaque plastic bags before measurements were taken. Rehydrated leaves taken towards the apical end of the branch (n=3 per individual: small, medium, and large) were scanned, weighed, dried at 60° C for  $\geq 48$  hours, and then re-scanned and weighed. Leaf area was calculated from scanned images using the LeafArea R package (Katabuchi, 2019).  $LMA$  was calculated as the ratio of leaf dry mass to fresh area.  $PLA_{dry}$  was calculated as the percent loss of area between fresh and dry leaves. Wood density was calculated for ~1cm diameter stem samples (bark and pith removed) as the ratio of dry weight to fresh volume, which was estimated using Archimedes’ displacement. We used the rapid determination method of Bartlett et al. (2012) to estimate osmotic potential at turgor loss point ( $\pi_{tlp}$ ). Briefly, two 4 mm diameter leaf discs were cut from each leaf, tightly wrapped in foil, submerged in liquid nitrogen, perforated 10-15 times with a dissection needle, and then measured using a vapour pressure osmometer (VAPRO 5520, Wescor, Logan, UT, USA). Osmotic potential ( $\pi_{osm}$ ) given by the osmometer was used to estimate ( $\pi_{tlp}$ ) using the equation  $\pi_{tlp} = 0.832\pi_{osm}^{-0.631}$  (Bartlett et al., 2012).

### Statistical Analysis

For each drought year, we calculated a metric drought resistance ( $Rt$ ) as the ratio of basal area increment ( $BAI$ ; *i.e.*, change in cross-sectional area) during the drought year to the mean  $BAI$  over the five years preceding the drought (Lloret et al., 2011). Thus,  $Rt$  values <1 and >1 indicate growth reductions and

increases, respectively. Because the  $Rt$  metric could be biased by directional pre-drought growth trends, we also tried an intervention time series analysis (ARIMA, (Hyndman et al., 2020)) that predicted mean drought-year growth based on trends over the preceding ten years and used this value in place of the five-year mean in calculations of resistance ( $Rt_{ARIMA} = \text{observed } BAI / \text{predicted } BAI$ ). The two metrics were strongly correlated (Fig. S5). Visual review of the individual tree-ring sequences with the largest discrepancies between these metrics revealed that  $Rt$  was less prone to unreasonable estimates than  $Rt_{ARIMA}$ , so we selected  $Rt$  as our focal metric, presenting parallel results for  $Rt_{ARIMA}$  in the Supplementary Info. In this study we focus exclusively on drought resistance metrics ( $Rt$  or  $Rt_{ARIMA}$ ), and not on the resilience metrics described in Lloret et al. (2011), because (1) we would expect resilience to be controlled by a different set of mechanisms, and (2) the findings of (DeSoto et al., 2020) suggest that  $Rt$  is a more important drought response metric for angiosperms in that low resistance to moderate droughts was a better predictor of mortality during subsequent severe droughts than the resilience metrics.

Analyses focused on testing the predictions presented in Table 1 with  $Rt$  as the response variable, and then repeated using  $Rt_{ARIMA}$  as the response variable. Models were run for all drought years combined and for each drought year individually. The general statistical model for hypothesis testing was a mixed effects model, implemented in the lme4 package in R (Bates et al., 2019). In the multi-year model, we included a random effect of tree nested within species and a fixed effect of drought year to represent the combined effects of differences in drought characteristics. Individual year models included a random effect of species. All models included fixed effects of independent variables of interest (Tables 1,3) as specified below. All variables across all best models had variance inflation factors  $< 1.2$  ( $1 \pm 0.019$ ). We used AICc to assess model selection, and conditional/marginal R-squared to assess model fit as implemented in the AICcmodavg package in R (Mazerolle and portions of code contributed by Dan Linden., 2019). AICc refers to a corrected version of AICc, and is best suited for small data sizes (see Brewer et al., 2016).

To avoid over-fitting models with five species traits (Table 3) across only 12 species, we did not include all traits as fixed effects in a single linear mixed model, but rather conducted individual tests of each species trait to determine the relative importance and appropriateness for inclusion in the main model. These tests followed the model structure specified above, then added  $\ln[H]$  and  $\ln[TWI]$  to create a base model against which we tested traits. Trait variables were considered appropriate for inclusion in the main model if they had a consistent direction of response across all droughts and if their addition to the base model improved fit (at  $\Delta AICc \geq 1.0$ ) in at least one drought year (Table S4). We note that we did not use the  $\Delta AICc \geq 1.0$  criterion as a test of significance, but rather of whether the variable had enough influence to be considered as a *candidate* variable in full models.

We then determined the top full models for predicting  $Rt$  (or  $Rt_{ARIMA}$ ). To do so, we compared models with all possible combinations of candidate variables, including  $\ln[H] * \ln[TWI]$  and species traits as specified above. We identified the full set of models within  $\Delta AICc = 2$  of the best model (that with lowest AICc). When a variable appeared in all of these models and the sign of the coefficient was consistent across models, we viewed this as support for the acceptance/rejection of the associated prediction (Table 1). If the variable appeared in some but not all of these models, and its sign was consistent across models, we considered this partial support/rejection. In presentation of the results below, we note instances where the  $Rt_{ARIMA}$  model disagreed with the  $Rt$  model, but otherwise do not discuss the  $Rt_{ARIMA}$  model.

All analysis beyond basic data collection was performed using R version 3.6.2 (R Core Team, 2019). Other R-packages used in analyses are listed in the Supplementary Information (*Appendix S1*). All data, code,



and results are available through the SCBI-ForestGEO organization on GitHub  
(<https://github.com/SCBI-ForestGEO/SCBI-ForestGEO-Data> and  
McGregor\_climate-sensitivity-variation repositories), with static versions corresponding to data and  
analyses presented here archived in Zenodo (DOIs: 10.5281/zenodo.3604993 and [TBD], respectively).

## Results

### *Tree height and microenvironment*

In the years for which we have vertical profiles in climate data (2016-2018), taller trees—or those in  
dominant crown positions—were generally exposed to higher evaporative demand during the peak growing  
season months (May-August; Fig. 2). Specifically, maximum daily wind speeds were significantly higher  
above the top of the canopy (40-50m) than within and below (10-30m) (Fig. 2a). Relative humidity was  
also somewhat lower during June-August, ranging from ~50-80% above the canopy and ~60-90% in the  
understory (Fig. 2b). Air temperature did not vary consistently across the vertical profile (Fig. 2c).

Crown position varied as expected with height (dominant > co-dominant > intermediate > suppressed),  
but with substantial variation (Fig. 2d). There were significant differences in height across all crown  
position classes (Fig. 2d). A comparison test between height and crown position data from the most recent  
ForestGEO census (2018) revealed a correlation of 0.73.

### *Community-level drought responses*

At the community level, cored trees showed substantial growth reductions in all three droughts, with a  
mean  $Rt$  of 0.86 in 1966 and 1999, and 0.84 in 1977 (Fig. 2b). Across the entire study period (1950-2009),  
the focal drought years were the three years with the largest fraction of trees exhibiting  $Rt \leq 0.7$ .  
Specifically, in each drought, roughly 30% of the cored trees had growth reductions of  $\geq 30\%$  ( $Rt \leq 0.7$ ):  
29% in 1966, 32% in 1977, and 27% in 1999. However, some individuals exhibited increased growth, *i.e.*,  
 $Rt > 1.0$ : 26% of trees in 1966, 22% in 1977, and 26% in 1999.

In the context of the multivariate model,  $Rt$  did not vary across drought years. That is, drought year as a  
variable did not appear in any of the top models – *i.e.*, models that were statistically indistinguishable  
( $\Delta AIC_c < 2$ ) from the best model.

### *Tree height, microenvironment, and drought resistance*

Taller trees (based on  $H$  in the drought year) showed stronger growth reductions during drought (Table 1;  
Figs. 4, S6). Specifically,  $\ln[H]$  appeared, with a negative coefficient, in the best model ( $\Delta AIC_c = 0$ ) and  
all top models when evaluating the three drought years together (Tables S6-S7). The same held true for  
1966 individually. For the 1977 drought,  $\ln[H]$  did not appear in the best model, but was included, with a  
negative coefficient, among the top models – *i.e.*, models that were statistically indistinguishable  
( $\Delta AIC_c < 2$ ) from the best model (Tables 1, S6-S7). For the 1999 drought,  $\ln[H]$  had no significant effect.

$Rt$  had a significantly negative response to  $\ln[TWI]$  across all drought years combined (Figs. 4, S6, Table  
S6-S7). The effect was also significant for 1977 and 1999 individually (Fig. 4, Table S6). When  $Rt_{ARIMA}$   
was used as the response variable, the effect was significant in 1977, and included in some of the top  
models in 1966 and 1999 (Table S7). This negates the idea that trees in moist microsites would be less  
affected by drought. Nevertheless, we tested for a  $\ln[H] * \ln[TWI]$  interaction, a negative sign of which

could indicate that smaller trees (presumably with smaller rooting volume) are more susceptible to drought in drier microenvironments with a deeper water table. This hypothesis was rejected, as the  $\ln[H] * \ln[TWI]$  interaction was never significant, and had a positive sign in any top models in which it appeared (Tables 1, S6-S7). This term did appear with a positive coefficient in the best  $Rt_{ARIMA}$  model for all years combined (Table S7), indicating that the negative effect of height on  $Rt$  was significantly stronger in wetter microhabitats.

#### *Species' traits and drought resistance*

Species, as a factor in ANOVA, had significant influence ( $p < 0.05$ ) on all traits (wood density,  $LMA$ ,  $PLA_{dry}$ , and  $\pi_{tlp}$ ), with more significant pairwise differences for wood density and  $PLA_{dry}$  than for  $LMA$  and  $\pi_{tlp}$  (Table 2, Fig. S4). Drought resistance also varied across species, overall and in each drought year (Fig. 3). Significant differences in  $Rt$  across species were most pronounced in 1966 with a total of seven distinct groupings, while 1977 had four and 1999 had two. Averaged across all droughts,  $Rt$  was lowest in *Liriodendron tulipifera* (mean  $Rt = 0.66$ ) and highest in *Fagus grandifolia* (mean  $Rt = 0.99$ ).

Wood density,  $LMA$ , and xylem porosity were all poor predictors of  $Rt$  (Tables 1, S4-S5). Wood density and  $LMA$  were never significantly associated with  $Rt$  in the single-variable tests and were therefore excluded from the full models. Xylem porosity was also excluded from the full models, as it had no significant influence for all droughts combined and had contrasting effects in the individual droughts: whereas ring-porous species had higher  $Rt$  than diffuse- and semi-ring-porous species in the 1966 and 1999 droughts, they had lower  $Rt$  in 1977 (Table S4). It is noteworthy that the two diffuse-porous species in our study, *Liriodendron tulipifera* and *Fagus grandifolia*, were at opposite ends of the  $Rt$  spectrum (Fig. 3), further refuting the idea that xylem porosity is a useful predictor of  $Rt$  in the context of this study.

In contrast,  $PLA_{dry}$ , and  $\pi_{tlp}$  were both negatively correlated to drought resistance (Figs. 4, S6; Tables 1, S4-S7). Both had consistent signs across all droughts, and their inclusion at least marginally improved the model ( $\Delta AICc > 1.0$ ) for at least one of the three droughts (Table S4), qualifying them as candidate variables for the full model.  $PLA_{dry}$  had a significant influence, with negative coefficient, in full models for the three droughts combined and for the 1966 drought individually (Fig. 4; Tables S6-S7). For 1977 and 1999, it was included with a negative coefficient in some of the top models (Tables S6-S7).  $\pi_{tlp}$  was included with a negative coefficient in the best model for both all droughts combined and for the 1977 drought individually (Fig. 4; Table S6). It was also included in some of the top models for 1999 (Tables S6-S7).

## **Discussion**

Tree height, microenvironment, and leaf drought tolerance traits shaped tree growth responses across three droughts at our study site (Table 1, Fig. 4). The greater susceptibility of larger trees to drought, similar to forests worldwide (Bennett et al., 2015), was driven primarily by their height (Stovall et al., 2019). Taller height was likely a liability in itself, and was also associated with greater exposure to conditions that would promote water loss and heat damage during drought (Fig. 2). There was no evidence that greater availability of, or access to, soil water availability increased drought resistance; in contrast, trees in wetter topographic positions had lower  $Rt$  (Zuleta et al., 2017; Stovall et al., 2019), and the larger potential rooting volume of large trees provided no advantage in the drier microenvironments. The negative effect of height on  $Rt$  held after accounting for species' traits, which is consistent with recent work finding height had a stronger influence on mortality risk than forest type during drought (Stovall et al., 2020). Drought

resistance was not consistently linked to species'  $LMA$ , wood density, or xylem type (ring- vs. diffuse porous), but was negatively correlated with leaf drought tolerance traits ( $PLA_{dry}$ ,  $\pi_{tlp}$ ). This is the first study to our knowledge linking  $PLA_{dry}$  and  $\pi_{tlp}$  to growth reduction during drought. The directions of these responses were consistent across droughts (Table S6), supporting the premise that they were driven by fundamental physiological mechanisms. However, the strengths of each predictor varied across droughts (Fig. 4; Tables S6-S7), indicating that drought characteristics interact with tree size, microenvironment, and traits to shape which individuals are most affected. These findings advance our knowledge of the factors that make trees vulnerable to growth declines during drought and, by extension, likely make them more vulnerable to mortality (Sapes et al., 2019).

The droughts considered here were of a magnitude that has occurred with an average frequency of approximately once every 10-15 years (Fig. 1a, Helcoski et al., 2019) and had substantial but not devastating impacts on tree growth (Figs. 1b). These droughts were classified as severe ( $PDSI < -3.0$ ; 1977) or extreme ( $PDSI < -4.0$ ; 1966, 1999) at our site and have been linked to tree mortality in the eastern United States (Druckenbrod et al., 2019). However, extreme, multiannual droughts such as the so-called “megadroughts” of this type that have triggered massive tree die-off in other regions (e.g., Allen et al., 2010; Stovall et al., 2019) have not occurred in the Eastern United States within the past several decades (Clark et al., 2016). Of the droughts considered here, the 1966 drought, which was preceded by two years of dry conditions (Fig. S1), severely stressed a larger portion of trees (Fig. 1b). The tendency for large trees to have lowest resistance was most pronounced in this drought, consistent with other findings that this physiological response increases with drought severity (Bennett et al., 2015; Stovall et al., 2019). Across all three droughts, the majority of trees experienced reduced growth, but a substantial portion had increased growth (Fig. 1b), consistent with prior observations that smaller trees can exhibit increased growth rates during drought (Bennett et al., 2015). It is likely because of the moderate impact of these droughts, along with other factors influencing tree growth (e.g., stand dynamics), that our best models characterize only a modest amount of variation in  $Rt$ : 11-12% for all droughts combined, and 18-25% for each individual drought (Fig. S6; Table S6).

Consistent with studies in other forests worldwide (Bennett et al., 2015), taller trees in this forest exhibited lower drought resistance. Mechanistically, this is consistent with, and reinforces, previous findings that biophysical constraints make it impossible for trees to efficiently transport water to great heights and simultaneously maintain strong resistance and resilience to drought-induced embolism (Olson et al., 2018; Couvreur et al., 2018; Roskilly et al., 2019). Taller trees also face dramatically different microenvironments (Fig. 2). They are exposed to higher wind speeds and lower humidity (Fig. 2a-b), resulting in higher evaporative demand. Unlike other temperate forests where modestly cooler understory conditions have been documented (Zellweger et al., 2019), particularly under drier conditions (Davis et al., 2019), we observed no significant variation in air temperatures across the vertical profile (Fig. 2c). More critically for tree physiology, leaf temperatures can become significantly elevated over air temperature under conditions of high solar radiation and low stomatal conductance (Campbell and Norman, 1998; Rey-Sánchez et al., 2016). Under drought, when air temperatures tend to be warmer, direct solar radiation tends to be higher (because of less cloud cover), and less water is available for evaporative cooling of the leaves, trees with sun-exposed crowns may not be able to simultaneously maintain leaf temperatures below damaging extremes and avoid drought-induced embolism. Indeed, previous studies have shown lower drought resistance in more exposed trees (Liu and Muller, 1993; Suarez et al., 2004; Scharnweber et al., 2019). Unfortunately, collinearity between height and crown exposure in this study (Fig. 2d) makes it impossible

to confidently partition causality. Additional research comparing drought responses of early successional and mature forest stands, along with short and tall isolated trees, would be valuable for more clearly disentangling the roles of tree height and crown exposure.

Belowground, taller trees would tend to have larger root systems (Enquist and Niklas, 2002), but this does not necessarily imply that they have greater access to or reliance on deep soil-water resources that may be critical during drought. While tree size can correlate with the depth of water extraction (Brum et al., 2019), the linkage is not consistent. Shorter trees can vary broadly in the depth of water uptake (Stahl et al., 2013), and larger trees may allocate more to abundant shallow roots that are beneficial for taking up water from rainstorms (Meinzer et al., 1999). Moreover, reliance on deep soil-water resources can actually prove a liability during severe and prolonged drought, as these can experience more intense water scarcity relative to non-drought conditions (Chitra-Tarak et al., 2018). In any case, the potentially greater access to water did not override the disadvantage conferred by height—and, in fact, greater moisture access in non-drought years (here, higher TWI) appears to make trees more sensitive to drought (Zuleta et al., 2017; Stovall et al., 2019). This may be because moister habitats would tend to support species and individuals with more mesophytic traits (Bartlett et al., 2016b; Mencuccini, 2003; Medeiros et al., 2019), potentially growing to greater heights (e.g., Detto et al. (2013)), and these are then more vulnerable when drought hits. The observed height-sensitivity of  $Rt$ , together with the lack of conferred advantage to large stature in drier topographic positions, agrees with the concept that physiological limitations to transpiration under drought shift from soil water availability to the plant-atmosphere interface as forests age (Bretfeld et al., 2018), such that tall, dominant trees are the most sensitive in mature forests. Again, additional research comparing drought responses across forests with different tree heights and water availability would be valuable for disentangling the relative importance of above- and belowground mechanisms across trees of different size.

The development of tree-ring chronologies for the twelve most dominant tree species at our site (Helcoski et al., 2019; Bourg et al., 2013) gave us the sample size to compare historical drought responses across species (Fig. 3) and associated traits at a single site (see also Elliott et al., 2015). Our study reinforced current understanding (see Introduction) that wood density and  $LMA$  are not reliably linked to drought resistance (Table 1). Contrary to previous studies in temperate deciduous forests, we did not find an association between xylem porosity and drought resistance, as the two diffuse-porous species, *Liriodendron tulipifera* and *Fagus grandifolia*, were at opposite ends of the  $Rt$  spectrum (Fig. 3). While the low  $Rt$  of *L. tulipifera* is consistent with other studies (Elliott et al., 2015), the high  $Rt$  of *F. grandifolia* contrasts with studies identifying diffuse porous species in general (Elliott et al., 2015; Kannenberg et al., 2019), and the genus *Fagus* in particular (Friedrichs et al., 2009), as drought sensitive. There are two potential explanations for this discrepancy. First, other traits can and do override the influence of xylem porosity on drought resistance. Ring-porous species are restricted mainly to temperate deciduous forests, while highly drought-tolerant diffuse-porous species exist in other biomes (Wheeler et al., 2007). *Fagus grandifolia* had intermediate  $\pi_{tlp}$  and low  $PLA_{dry}$  (Fig. S4), which would have contributed to its drought resistance (Fig. 4; see discussion below). A second explanation of why *F. grandifolia* trees at this particular site had higher  $Rt$  is that the sampled individuals, reflective of the population within the plot, are generally shorter and in less-dominant canopy positions compared to most other species (Fig. S4). The species, which is highly shade-tolerant, also has deep crowns (Anderson-Teixeira et al., 2015b), implying that a lower proportion of leaves would be affected by harsher microclimatic conditions at the top of the canopy under drought (Fig. 2). Thus, the high  $Rt$  of the sampled *F. grandifolia* population can be explained by a combination of fairly

drought-resistant leaf traits, shorter stature, and a buffered microenvironment.

Concerted measurement of tree-rings and leaf drought tolerance traits of emerging importance (Scoffoni et al., 2014; Bartlett et al., 2016a; Medeiros et al., 2019) allowed novel insights into the role of drought tolerance traits in shaping drought response. The finding that  $PLA_{dry}$  and  $\pi_{tlp}$  can be useful for predicting drought responses of tree growth (Fig. 4; Table 1) is both novel and consistent with previous studies linking these traits to habitat and drought tolerance. Previous studies have demonstrated that  $\pi_{tlp}$  and  $PLA_{dry}$  are physiologically meaningful traits linked to species distribution along moisture gradients (Maréchaux et al., 2015; Fletcher et al., 2018; Medeiros et al., 2019; Simeone et al., 2019; Rosas et al., 2019; Zhu et al., 2018), and our findings indicate that these traits also influence drought responses. Furthermore, the observed linkage of  $\pi_{tlp}$  to  $Rt$  in this forest aligns with observations in the Amazon that  $\pi_{tlp}$  is higher in drought-intolerant than drought-tolerant plant functional type. Further, it adds support to the idea that this trait is useful for categorizing and representing species' drought responses in models (Powell et al., 2017). Because both  $PLA_{dry}$  and  $\pi_{tlp}$  can be measured relatively easily (Bartlett et al., 2012; Scoffoni et al., 2014), they hold promise for predicting drought growth responses across diverse forests. The importance of predicting drought responses from species traits increases with tree species diversity; whereas it is feasible to study drought responses for all dominant species in most boreal and temperate forests (e.g., this study), this becomes difficult to impossible for species that do not form annual rings, and for diverse tropical forests. Although progress is being made for the tropics (Schöngart et al., 2017), a full linkage of drought tolerance traits to drought responses would be invaluable for forecasting how little-known species and whole forests will respond to future droughts (Christoffersen et al., 2016; Powell et al., 2017).

As climate change drives increasing drought in many of the world's forests (Trenberth et al., 2014; Intergovernmental Panel on Climate Change, 2015), the fate of forests and their climate feedbacks will be shaped by the biophysical and physiological drivers observed here. Our results show that taller, more exposed trees and species with less drought-tolerant leaf traits will be most affected, at least in terms of growth during the drought year. Resilience and survival are both linked to resistance (**REFS**), implying that the same factors may influence these. Indeed, while the influence of  $PLA_{dry}$  and  $\pi_{tlp}$  on drought resilience and survival remains to be tested, taller trees have lower resilience (Trugman et al., 2018; Gillerot et al., 2020) and survival (Bennett et al., 2015; Stovall et al., 2019). As climate change-driven droughts affect forests worldwide, there is likely to be a shift from mature forests with tall, buffering trees to forests with a shorter overall stature (McDowell et al., 2020). At this point, species whose drought resistance relies in part on existence within a buffered microenvironment (e.g., *F. grandifolia*) could in turn become more susceptible. Here, the relative importance of tree height *per se* versus crown exposure becomes crucial, shaping whether the dominant trees of shorter canopies are significantly more drought resistant because of their shorter stature, or whether high exposure makes them as vulnerable as the taller trees of the former canopy. Studies disentangling the influence of height and exposure on drought tolerance will be critical to answering this question. Ultimately, distributions of tree heights and drought tolerance traits across broad moisture gradients suggest that forests exposed to more drought will ultimately shift towards shorter stature (**REFS**) and be dominated by species with more drought-tolerant traits (**REFS**). Our study helps to elucidate the mechanisms behind these patterns, opening the door for more accurate forecasting of forest responses to future drought.

## Acknowledgements

We especially thank the numerous researchers who helped to collect the data used here, in particular Jennifer C. McGarvey, Jonathan R. Thompson, and Victoria Meakem for original collection and processing of cores. Thanks also to Camila D. Medeiros for guidance on leaf drought tolerance and functional trait measurements, Edward Brzostek’s lab for collaboration on leaf sampling, and Maya Prestipino for data collection. This manuscript was improved based on helpful reviews by Mark Olson and three anonymous reviewers. Funding for the establishment of the SCBI ForestGEO Large Forest Dynamics Plot was provided by the Smithsonian-led Forest Global Earth Observatory (ForestGEO), the Smithsonian Institution, and the HSBC Climate Partnership. This study was funded by ForestGEO, a Virginia Native Plant Society grant to KAT and AJT, and support from the Harvard Forest and National Science Foundation which supports the PaleON project (NSF EF-1241930) for NP.

## Author Contribution

KAT, IM, and AJT designed the research. Tree-ring chronologies were developed by RH under guidance of AJT and NP. Trait data were collected by IM, JZ under guidance of NK and LS. Other plot data were collected by IM, AS, EGA, and NB under guidance of EGA and WM. Data analyses were performed by IM under guidance of KAT and VH. KAT and IM interpreted the results. IM and KAT wrote the first draft of manuscript, and all authors contributed to revisions.

## Supplementary Information

Table S1: Monthly Palmer Drought Severity Index (PDSI), and its rank among all years between 1950 and 2009 (driest=1), for focal droughts.

Table S2: Species-specific bark thickness regression equations.

Table S3: Species-specific height regression equations.

Table S4. Individual tests of species traits as drivers of drought resistance, where  $Rt$  is used as the response variable.

Table S5. Individual tests of species traits as drivers of drought resistance, where  $Rt_{ARIMA}$  is used as the response variable.

Table S6. Summary of top full models for each drought instance, where  $Rt$  is used as the response variable.

Table S7. Summary of top models for each drought instance, where  $Rt_{ARIMA}$  is used as the response variable.

Figure S1. Time series of Palmer Drought Severity Index (PDSI) for the 2.5 years prior to each focal drought

Figure S2: Map of ForestGEO plot showing topographic wetness index and location of cored trees

Figure S3: Distribution of reconstructed tree heights across drought years.

Figure S4. Distribution of independent variables by species.

Figure S5. Comparison of  $R_t$  and  $R_{tARIMA}$  results, with residuals, for each drought scenario

Figure S6. Visualization of best model, with data, for all droughts combined.

## References

- Abrams, M. D. (1990). Adaptations and responses to drought in *Quercus* species of North America. *Tree Physiology*, 7(1-2-3-4):227–238.
- Allen, C. D., Breshears, D. D., and McDowell, N. G. (2015). On underestimation of global vulnerability to tree mortality and forest die-off from hotter drought in the Anthropocene. *Ecosphere*, 6(8):art129.
- Allen, C. D., Macalady, A. K., Chenchouni, H., Bachelet, D., McDowell, N., Vennetier, M., Kitzberger, T., Rigling, A., Breshears, D. D., Hogg, E. H. T., Gonzalez, P., Fensham, R., Zhang, Z., Castro, J., Demidova, N., Lim, J.-H., Allard, G., Running, S. W., Semerci, A., and Cobb, N. (2010). A global overview of drought and heat-induced tree mortality reveals emerging climate change risks for forests. *Forest Ecology and Management*, 259(4):660–684.
- Anderegg, W. R. L., Klein, T., Bartlett, M., Sack, L., Pellegrini, A. F. A., Choat, B., and Jansen, S. (2016). Meta-analysis reveals that hydraulic traits explain cross-species patterns of drought-induced tree mortality across the globe. *Proceedings of the National Academy of Sciences*, 113(18):5024–5029.
- Anderson-Teixeira, K. J., Davies, S. J., Bennett, A. C., Gonzalez-Akre, E. B., Muller-Landau, H. C., Wright, S. J., Salim, K. A., Zambrano, A. M. A., Alonso, A., Baltzer, J. L., Basset, Y., Bourg, N. A., Broadbent, E. N., Brockelman, W. Y., Bunyavejchewin, S., Burslem, D. F. R. P., Butt, N., Cao, M., Cardenas, D., Chuyong, G. B., Clay, K., Cordell, S., Dattaraja, H. S., Deng, X., Detto, M., Du, X., Duque, A., Erikson, D. L., Ewango, C. E. N., Fischer, G. A., Fletcher, C., Foster, R. B., Giardina, C. P., Gilbert, G. S., Gunatilleke, N., Gunatilleke, S., Hao, Z., Hargrove, W. W., Hart, T. B., Hau, B. C. H., He, F., Hoffman, F. M., Howe, R. W., Hubbell, S. P., Inman-Narahari, F. M., Jansen, P. A., Jiang, M., Johnson, D. J., Kanzaki, M., Kassim, A. R., Kenfack, D., Kibet, S., Kinnaird, M. F., Korte, L., Kral, K., Kumar, J., Larson, A. J., Li, Y., Li, X., Liu, S., Lum, S. K. Y., Lutz, J. A., Ma, K., Maddalena, D. M., Makana, J.-R., Malhi, Y., Marthens, T., Serudin, R. M., McMahon, S. M., McShea, W. J., Memiaghe, H. R., Mi, X., Mizuno, T., Morecroft, M., Myers, J. A., Novotny, V., Oliveira, A. A. d., Ong, P. S., Orwig, D. A., Ostertag, R., Ouden, J. d., Parker, G. G., Phillips, R. P., Sack, L., Sainge, M. N., Sang, W., Sri-ngernyung, K., Sukumar, R., Sun, I.-F., Sungpalee, W., Suresh, H. S., Tan, S., Thomas, S. C., Thomas, D. W., Thompson, J., Turner, B. L., Uriarte, M., Valencia, R., Vallejo, M. I., Vicentini, A., Vrška, T., Wang, X., Wang, X., Weiblen, G., Wolf, A., Xu, H., Yap, S., and Zimmerman, J. (2015a). CTFS-ForestGEO: a worldwide network monitoring forests in an era of global change. *Global Change Biology*, 21(2):528–549.
- Anderson-Teixeira, K. J., McGarvey, J. C., Muller-Landau, H. C., Park, J. Y., Gonzalez-Akre, E. B., Herrmann, V., Bennett, A. C., So, C. V., Bourg, N. A., Thompson, J. R., McMahon, S. M., and McShea, W. J. (2015b). Size-related scaling of tree form and function in a mixed-age forest. *Functional Ecology*, 29(12):1587–1602.

- Bartlett, M. K., Klein, T., Jansen, S., Choat, B., and Sack, L. (2016a). The correlations and sequence of plant stomatal, hydraulic, and wilting responses to drought. *Proceedings of the National Academy of Sciences*, 113(46):13098–13103.
- Bartlett, M. K., Scoffoni, C., Ardy, R., Zhang, Y., Sun, S., Cao, K., and Sack, L. (2012). Rapid determination of comparative drought tolerance traits: using an osmometer to predict turgor loss point. *Methods in Ecology and Evolution*, 3(5):880–888.
- Bartlett, M. K., Zhang, Y., Yang, J., Kreidler, N., Sun, S.-W., Lin, L., Hu, Y.-H., Cao, K.-F., and Sack, L. (2016b). Drought tolerance as a driver of tropical forest assembly: resolving spatial signatures for multiple processes. *Ecology*, 97(2):503–514.
- Bates, D., Maechler, M., Bolker, B., and Walker, S. (2019). *lme4: Linear Mixed-Effects Models using Eigen and S4*. R package version 1.1-21.
- Bennett, A. C., McDowell, N. G., Allen, C. D., and Anderson-Teixeira, K. J. (2015). Larger trees suffer most during drought in forests worldwide. *Nature Plants*, 1(10):15139.
- Beven, K. J. and Kirkby, M. J. (1979). A physically based, variable contributing area model of basin hydrology / Un modèle à base physique de zone d’appel variable de l’hydrologie du bassin versant. *Hydrological Sciences Bulletin*, 24(1):43–69.
- Bonan, G. B. (2008). Forests and Climate Change: Forcings, Feedbacks, and the Climate Benefits of Forests. *Science*, 320(5882):1444–1449.
- Bourg, N. A., McShea, W. J., Thompson, J. R., McGarvey, J. C., and Shen, X. (2013). Initial census, woody seedling, seed rain, and stand structure data for the SCBI SIGEO Large Forest Dynamics Plot. *Ecology*, 94(9):2111–2112.
- Bretfeld, M., Ewers, B. E., and Hall, J. S. (2018). Plant water use responses along secondary forest succession during the 2015–2016 El Niño drought in Panama. *New Phytologist*, 219(3):885–899.
- Brewer, M. J., Butler, A., and Cooksley, S. L. (2016). The relative performance of AIC, AICC and BIC in the presence of unobserved heterogeneity. *Methods in Ecology and Evolution*, 7(6):679–692.
- Brum, M., Vadeboncoeur, M. A., Ivanov, V., Asbjornsen, H., Saleska, S., Alves, L. F., Penha, D., Dias, J. D., Aragão, L. E. O. C., Barros, F., Bittencourt, P., Pereira, L., and Oliveira, R. S. (2019). Hydrological niche segregation defines forest structure and drought tolerance strategies in a seasonal Amazon forest. *Journal of Ecology*, 107(1):318–333.
- Campbell, G. S. and Norman, J. M. (1998). *An Introduction to Environmental Biophysics*, volume 2nd. Springer, New York.
- Chitra-Tarak, R., Ruiz, L., Dattaraja, H. S., Kumar, M. S. M., Riotte, J., Suresh, H. S., McMahon, S. M., and Sukumar, R. (2018). The roots of the drought: Hydrology and water uptake strategies mediate forest-wide demographic response to precipitation. *Journal of Ecology*, 106(4):1495–1507.
- Christoffersen, B. O., Gloor, M., Fauset, S., Fyllas, N. M., Galbraith, D. R., Baker, T. R., Kruijt, B., Rowland, L., Fisher, R. A., Binks, O. J., Sevanto, S., Xu, C., Jansen, S., Choat, B., Mencuccini, M., McDowell, N. G., and Meir, P. (2016). Linking hydraulic traits to tropical forest function in a size-structured and trait-driven model (TFS v.1-Hydro).



- Clark, J. S., Iverson, L., Woodall, C. W., Allen, C. D., Bell, D. M., Bragg, D. C., D'Amato, A. W., Davis, F. W., Hersh, M. H., Ibanez, I., Jackson, S. T., Matthews, S., Pederson, N., Peters, M., Schwartz, M. W., Waring, K. M., and Zimmermann, N. E. (2016). The impacts of increasing drought on forest dynamics, structure, and biodiversity in the United States. *Global Change Biology*, 22(7):2329–2352.
- Condit, R. (1998). *Tropical Forest Census Plots: Methods and Results from Barro Colorado Island, Panama and a Comparison with Other Plots*. Springer Berlin Heidelberg, Berlin, Heidelberg.
- Cook, B. I., Ault, T. R., and Smerdon, J. E. (2015). Unprecedented 21st century drought risk in the American Southwest and Central Plains. *Science Advances*, 1(1):e1400082.
- Couvreur, V., Ledder, G., Manzoni, S., Way, D. A., Muller, E. B., and Russo, S. E. (2018). Water transport through tall trees: A vertically explicit, analytical model of xylem hydraulic conductance in stems. *Plant, Cell & Environment*, 41(8):1821–1839.
- Dai, A., Zhao, T., and Chen, J. (2018). Climate Change and Drought: a Precipitation and Evaporation Perspective. *Current Climate Change Reports*, 4(3):301–312.
- Davis, K. T., Dobrowski, S. Z., Holden, Z. A., Higuera, P. E., and Abatzoglou, J. T. (2019). Microclimatic buffering in forests of the future: the role of local water balance. *Ecography*, 42(1):1–11.
- DeSoto, L., Cailleret, M., Sterck, F., Jansen, S., Kramer, K., Robert, E. M. R., Aakala, T., Amoroso, M. M., Bigler, C., Camarero, J. J., Čufar, K., Gea-Izquierdo, G., Gillner, S., Haavik, L. J., Hereş, A.-M., Kane, J. M., Kharuk, V. I., Kitzberger, T., Klein, T., Levanič, T., Linares, J. C., Mäkinen, H., Oberhuber, W., Papadopoulos, A., Rohner, B., Sangüesa-Barreda, G., Stojanovic, D. B., Suárez, M. L., Villalba, R., and Martínez-Vilalta, J. (2020). Low growth resilience to drought is related to future mortality risk in trees. *Nature Communications*, 11(1):545.
- Detto, M., Muller-Landau, H. C., Mascaro, J., and Asner, G. P. (2013). Hydrological Networks and Associated Topographic Variation as Templates for the Spatial Organization of Tropical Forest Vegetation. *PLOS ONE*, 8(10):e76296.
- Druckenbrod, D. L., Martin-Benito, D., Orwig, D. A., Pederson, N., Poulter, B., Renwick, K. M., and Shugart, H. H. (2019). Redefining temperate forest responses to climate and disturbance in the eastern United States: New insights at the mesoscale. *Global Ecology and Biogeography*, 28(5):557–575.
- Elliott, K. J., Miniati, C. F., Pederson, N., and Laseter, S. H. (2015). Forest tree growth response to hydroclimate variability in the southern Appalachians. *Global Change Biology*, 21(12):4627–4641.
- Enquist, B. J. and Niklas, K. J. (2002). Global Allocation Rules for Patterns of Biomass Partitioning in Seed Plants. *Science*, 295(5559):1517–1520.
- Farrell, C., Szota, C., and Arndt, S. K. (2017). Does the turgor loss point characterize drought response in dryland plants? *Plant, Cell & Environment*, 40(8):1500–1511.
- Fletcher, L. R., Cui, H., Callahan, H., Scoffoni, C., John, G. P., Bartlett, M. K., Burge, D. O., and Sack, L. (2018). Evolution of leaf structure and drought tolerance in species of Californian Ceanothus. *American Journal of Botany*, 105(10):1672–1687.

- Friedlingstein, P., Cox, P., Betts, R., Bopp, L., von Bloh, W., Brovkin, V., Cadule, P., Doney, S., Eby, M.,  
 Fung, I., Bala, G., John, J., Jones, C., Joos, F., Kato, T., Kawamiya, M., Knorr, W., Lindsay, K.,  
 Matthews, H. D., Raddatz, T., Rayner, P., Reick, C., Roeckner, E., Schnitzler, K.-G., Schnur, R.,  
 Strassmann, K., Weaver, A. J., Yoshikawa, C., and Zeng, N. (2006). Climate–Carbon Cycle Feedback  
 Analysis: Results from the C4MIP Model Intercomparison. *Journal of Climate*, 19(14):3337–3353.
- Friedrichs, D. A., Trouet, V., Büntgen, U., Frank, D. C., Esper, J., Neuwirth, B., and Löffler, J. (2009).  
 Species-specific climate sensitivity of tree growth in Central-West Germany. *Trees*, 23(4):729.
- Gillerot, L., Forrester, D. I., Bottero, A., Rigling, A., and Lévesque, M. (2020). Tree Neighbourhood  
 Diversity Has Negligible Effects on Drought Resilience of European Beech, Silver Fir and Norway  
 Spruce. *Ecosystems*.
- Gonzalez-Akre, E., Meakem, V., Eng, C.-Y., Tepley, A. J., Bourg, N. A., McShea, W., Davies, S. J., and  
 Anderson-Teixeira, K. (2016). Patterns of tree mortality in a temperate deciduous forest derived from a  
 large forest dynamics plot. *Ecosphere*, 7(12):e01595.
- Greenwood, S., Ruiz-Benito, P., Martínez-Vilalta, J., Lloret, F., Kitzberger, T., Allen, C. D., Fensham, R.,  
 Laughlin, D. C., Kattge, J., Bönisch, G., Kraft, N. J. B., and Jump, A. S. (2017). Tree mortality across  
 biomes is promoted by drought intensity, lower wood density and higher specific leaf area. *Ecology*  
*Letters*, 20(4):539–553.
- Guerfel, M., Baccouri, O., Boujnah, D., Chaïbi, W., and Zarrouk, M. (2009). Impacts of water stress on  
 gas exchange, water relations, chlorophyll content and leaf structure in the two main Tunisian olive  
 (*Olea europaea* L.) cultivars. *Scientia Horticulturae*, 119(3):257–263.
- Hacket-Pain, A. J., Cavin, L., Friend, A. D., and Jump, A. S. (2016). Consistent limitation of growth by  
 high temperature and low precipitation from range core to southern edge of European beech indicates  
 widespread vulnerability to changing climate. *European Journal of Forest Research*, 135(5):897–909.
- Harris, I., Jones, P. D., Osborn, T. J., and Lister, D. H. (2014). Updated high-resolution grids of monthly  
 climatic observations – the CRU TS3.10 Dataset. *International Journal of Climatology*, 34(3):623–642.
- Helcoski, R., Tepley, A. J., Pederson, N., McGarvey, J. C., Meakem, V., Herrmann, V., Thompson, J. R.,  
 and Anderson-Teixeira, K. J. (2019). Growing season moisture drives interannual variation in woody  
 productivity of a temperate deciduous forest. *New Phytologist*, 0(0).
- Hoffmann, W. A., Marchin, R. M., Abit, P., and Lau, O. L. (2011). Hydraulic failure and tree dieback are  
 associated with high wood density in a temperate forest under extreme drought. *Global Change Biology*,  
 17(8):2731–2742.
- Hollister, J. (2018). *elevatr: Access Elevation Data from Various APIs*. R package version 0.2.0.
- Hyndman, R., Athanasopoulos, G., Bergmeir, C., Caceres, G., Chhay, L., O’Hara-Wild, M., Petropoulos,  
 F., Razbash, S., Wang, E., and Yasmeen, F. (2020). *forecast: Forecasting Functions for Time Series and*  
*Linear Models*. R package version 8.12.
- Intergovernmental Panel on Climate Change (2015). *Climate Change 2014: Impacts, Adaptation and*  
*Vulnerability: Working Group II Contribution to the IPCC Fifth Assessment Report. Volume 2 Volume*  
*2*. OCLC: 900892773.

- Jennings, S. B., Brown, N. D., and Sheil, D. (1999). Assessing forest canopies and understorey illumination: canopy closure, canopy cover and other measures. *Forestry: An International Journal of Forest Research*, 72(1):59–74.
- Kannenbergh, S. A., Novick, K. A., Alexander, M. R., Maxwell, J. T., Moore, D. J. P., Phillips, R. P., and Anderegg, W. R. L. (2019). Linking drought legacy effects across scales: From leaves to tree rings to ecosystems. *Global Change Biology*, 0(ja).
- Katabuchi, M. (2019). *LeafArea: Rapid Digital Image Analysis of Leaf Area*. R package version 0.1.8.
- Kennedy, D., Swenson, S., Oleson, K. W., Lawrence, D. M., Fisher, R., Costa, A. C. L. d., and Gentine, P. (2019). Implementing Plant Hydraulics in the Community Land Model, Version 5. *Journal of Advances in Modeling Earth Systems*, 11(2):485–513.
- Koike, T., Kitao, M., Maruyama, Y., Mori, S., and Lei, T. T. (2001). Leaf morphology and photosynthetic adjustments among deciduous broad-leaved trees within the vertical canopy profile. *Tree Physiology*, 21(12-13):951–958.
- Kunert, N., Aparecido, L. M. T., Wolff, S., Higuchi, N., Santos, J. d., Araujo, A. C. d., and Trumbore, S. (2017). A revised hydrological model for the Central Amazon: The importance of emergent canopy trees in the forest water budget. *Agricultural and Forest Meteorology*, 239:47–57.
- Larjavaara, M. and Muller-Landau, H. C. (2013). Measuring tree height: a quantitative comparison of two common field methods in a moist tropical forest. *Methods in Ecology and Evolution*, 4(9):793–801.
- Liu, H., Gleason, S. M., Hao, G., Hua, L., He, P., Goldstein, G., and Ye, Q. (2019). Hydraulic traits are coordinated with maximum plant height at the global scale. *Science Advances*, 5(2):eaav1332.
- Liu, Y. and Muller, R. N. (1993). Effect of Drought and Frost on Radial Growth of Overstory and Understory Stems in a Deciduous Forest. *The American Midland Naturalist*, 129(1):19–25.
- Lloret, F., Keeling, E. G., and Sala, A. (2011). Components of tree resilience: effects of successive low-growth episodes in old ponderosa pine forests. *Oikos*, 120(12):1909–1920.
- Lunch, C., Laney, C., Mietkiewicz, N., Sokol, E., Cawley, K., and NEON (National Ecological Observatory Network) (2020). *neonUtilities: Utilities for Working with NEON Data*. R package version 1.3.5.
- Martin-Benito, D. and Pederson, N. (2015). Convergence in drought stress, but a divergence of climatic drivers across a latitudinal gradient in a temperate broadleaf forest. *Journal of Biogeography*, 42(5):925–937.
- Maréchaux, I., Bartlett, M. K., Sack, L., Baraloto, C., Engel, J., Joetzjer, E., and Chave, J. (2015). Drought tolerance as predicted by leaf water potential at turgor loss point varies strongly across species within an Amazonian forest. *Functional Ecology*, 29(10):1268–1277.
- Maréchaux, I., Saint-André, L., Bartlett, M. K., Sack, L., and Chave, J. (2019). Leaf drought tolerance cannot be inferred from classic leaf traits in a tropical rainforest. *Journal of Ecology*.
- Mazerolle, M. J. and portions of code contributed by Dan Linden. (2019). *AICcmodavg: Model Selection and Multimodel Inference Based on (Q)AIC(c)*. R package version 2.2-2.

- McDowell, N. G. and Allen, C. D. (2015). Darcy’s law predicts widespread forest mortality under climate warming. *Nature Climate Change*, 5(7):669–672.
- McDowell, N. G., Allen, C. D., Anderson-Teixeira, K., Aukema, B. H., Bond-Lamberty, B., Chini, L., Clark, J. S., Dietze, M., Grossiord, C., Hanbury-Brown, A., Hurtt, G. C., Jackson, R. B., Johnson, D. J., Kueppers, L., Lichstein, J. W., Ogle, K., Poulter, B., Pugh, T. A. M., Seidl, R., Turner, M. G., Uriarte, M., Walker, A. P., and Xu, C. (2020). Pervasive shifts in forest dynamics in a changing world. *Science*, 368(6494).
- McDowell, N. G., Bond, B. J., Dickman, L. T., Ryan, M. G., and Whitehead, D. (2011). Relationships Between Tree Height and Carbon Isotope Discrimination. In Meinzer, F. C., Lachenbruch, B., and Dawson, T. E., editors, *Size- and Age-Related Changes in Tree Structure and Function*, Tree Physiology, pages 255–286. Springer Netherlands, Dordrecht.
- Meakem, V., Tepley, A. J., Gonzalez-Akre, E. B., Herrmann, V., Muller-Landau, H. C., Wright, S. J., Hubbell, S. P., Condit, R., and Anderson-Teixeira, K. J. (2018). Role of tree size in moist tropical forest carbon cycling and water deficit responses. *New Phytologist*, 219(3):947–958.
- Medeiros, C. D., Scoffoni, C., John, G. P., Bartlett, M. K., Inman-Narahari, F., Ostertag, R., Cordell, S., Giardina, C., and Sack, L. (2019). An extensive suite of functional traits distinguishes Hawaiian wet and dry forests and enables prediction of species vital rates. *Functional Ecology*, 33(4):712–734.
- Meinzer, F. C., Andrade, J. L., Goldstein, G., Holbrook, N. M., Cavelier, J., and Wright, S. J. (1999). Partitioning of soil water among canopy trees in a seasonally dry tropical forest. *Oecologia*, 121(3):293–301.
- Mencuccini, M. (2003). The ecological significance of long-distance water transport: short-term regulation, long-term acclimation and the hydraulic costs of stature across plant life forms. *Plant, Cell & Environment*, 26(1):163–182.
- Metcalfe, P., Beven, K., and Freer, J. (2018). *dynatopmodel: Implementation of the Dynamic TOPMODEL Hydrological Model*. R package version 1.2.1.
- NEON (2018). National Ecological Observatory Network. 2016, 2017, 2018. Data Products: DP1.00001.001, DP1.00098.001, DP1.00002.001. Provisional data downloaded from <http://data.neonscience.org/> in May 2019. Battelle, Boulder, CO, USA.
- Olson, M., Rosell, J. A., Martínez-Pérez, C., León-Gómez, C., Fajardo, A., Isnard, S., Cervantes-Alcayde, M. A., Echeverría, A., Figueroa-Abundiz, V. A., Segovia-Rivas, A., Trueba, S., and Vázquez-Segovia, K. (2020). Xylem vessel-diameter–shoot-length scaling: ecological significance of porosity types and other traits. *Ecological Monographs*, n/a(n/a).
- Olson, M. E., Anfodillo, T., Rosell, J. A., Petit, G., Crivellaro, A., Isnard, S., León-Gómez, C., Alvarado-Cárdenas, L. O., and Castorena, M. (2014). Universal hydraulics of the flowering plants: vessel diameter scales with stem length across angiosperm lineages, habits and climates. *Ecology Letters*, 17(8):988–997.
- Olson, M. E., Soriano, D., Rosell, J. A., Anfodillo, T., Donoghue, M. J., Edwards, E. J., León-Gómez, C., Dawson, T., Martínez, J. J. C., Castorena, M., Echeverría, A., Espinosa, C. I., Fajardo, A., Gazol, A.,

- Isnard, S., Lima, R. S., Marcati, C. R., and Méndez-Alonzo, R. (2018). Plant height and hydraulic vulnerability to drought and cold. *Proceedings of the National Academy of Sciences*, 115(29):7551–7556.
- Phillips, N. G., Ryan, M. G., Bond, B. J., McDowell, N. G., Hinckley, T. M., and Čermák, J. (2003). Reliance on stored water increases with tree size in three species in the Pacific Northwest. *Tree Physiology*, 23(4):237–245.
- Powell, T. L., Wheeler, J. K., Oliveira, A. A. R. d., Costa, A. C. L. d., Saleska, S. R., Meir, P., and Moorcroft, P. R. (2017). Differences in xylem and leaf hydraulic traits explain differences in drought tolerance among mature Amazon rainforest trees. *Global Change Biology*, 23(10):4280–4293.
- Pretzsch, H., Schütze, G., and Biber, P. (2018). Drought can favour the growth of small in relation to tall trees in mature stands of Norway spruce and European beech. *Forest Ecosystems*, 5(1):20.
- R Core Team (2019). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria.
- Rey-Sánchez, A. C., Slot, M., Posada, J. M., and Kitajima, K. (2016). Spatial and seasonal variation in leaf temperature within the canopy of a tropical forest. *Climate Research*, 71(1):75–89.
- Rosas, T., Mencuccini, M., Barba, J., Cochard, H., Saura-Mas, S., and Martínez-Vilalta, J. (2019). Adjustments and coordination of hydraulic, leaf and stem traits along a water availability gradient. *New Phytologist*, 223(2):632–646.
- Roskilly, B., Keeling, E., Hood, S., Giuggiola, A., and Sala, A. (2019). Conflicting functional effects of xylem pit structure relate to the growth-longevity trade-off in a conifer species. *PNAS*. doi: /10.1073/pnas.1900734116.
- Ryan, M. G., Phillips, N., and Bond, B. J. (2006). The hydraulic limitation hypothesis revisited. *Plant, Cell & Environment*, 29(3):367–381.
- Sapes, G., Roskilly, B., Dobrowski, S., Maneta, M., Anderegg, W. R. L., Martinez-Vilalta, J., and Sala, A. (2019). Plant water content integrates hydraulics and carbon depletion to predict drought-induced seedling mortality. *Tree Physiology*, 39(8):1300–1312.
- Scharnweber, T., Heinze, L., Cruz-García, R., van der Maaten-Theunissen, M., and Wilmking, M. (2019). Confessions of solitary oaks: We grow fast but we fear the drought. *Dendrochronologia*, 55:43–49.
- Scholz, F. G., Phillips, N. G., Bucci, S. J., Meinzer, F. C., and Goldstein, G. (2011). Hydraulic Capacitance: Biophysics and Functional Significance of Internal Water Sources in Relation to Tree Size. In Meinzer, F. C., Lachenbruch, B., and Dawson, T. E., editors, *Size- and Age-Related Changes in Tree Structure and Function*, Tree Physiology, pages 341–361. Springer Netherlands, Dordrecht.
- Schöngart, J., Bräuning, A., Barbosa, A. C. M. C., Lisi, C. S., and de Oliveira, J. M. (2017). Dendroecological Studies in the Neotropics: History, Status and Future Challenges. In Amoroso, M. M., Daniels, L. D., Baker, P. J., and Camarero, J. J., editors, *Dendroecology: Tree-Ring Analyses Applied to Ecological Studies*, Ecological Studies, pages 35–73. Springer International Publishing, Cham.
- Scoffoni, C., Vuong, C., Diep, S., Cochard, H., and Sack, L. (2014). Leaf Shrinkage with Dehydration: Coordination with Hydraulic Vulnerability and Drought Tolerance. *Plant Physiology*, 164(4):1772–1788.

- Simeone, C., Maneta, M. P., Holden, Z. A., Sapes, G., Sala, A., and Dobrowski, S. Z. (2019). Coupled ecohydrology and plant hydraulics modeling predicts ponderosa pine seedling mortality and lower treeline in the US Northern Rocky Mountains. *New Phytologist*, 221(4):1814–1830.
- Slette, I. J., Post, A. K., Awad, M., Even, T., Punzalan, A., Williams, S., Smith, M. D., and Knapp, A. K. (2019). How ecologists define drought, and why we should do better. *Global Change Biology*, 0(0):1–8.
- Stahl, C., Hérault, B., Rossi, V., Burban, B., Bréchet, C., and Bonal, D. (2013). Depth of soil water uptake by tropical rainforest trees during dry periods: does tree dimension matter? *Oecologia*, 173(4):1191–1201.
- Stovall, A. E. L., Anderson-Teixeira, K. J., and Shugart, H. H. (2018a). Assessing terrestrial laser scanning for developing non-destructive biomass allometry. *Forest Ecology and Management*, 427:217–229.
- Stovall, A. E. L., Anderson-Teixeira, K. J., and Shugart, H. H. (2018b). Terrestrial LiDAR-derived non-destructive woody biomass estimates for 10 hardwood species in Virginia. *Data in Brief*, 19:1560–1569.
- Stovall, A. E. L., Shugart, H., and Yang, X. (2019). Tree height explains mortality risk during an intense drought. *Nature Communications*, 10(1):1–6.
- Stovall, A. E. L., Shugart, H. H., and Yang, X. (2020). Reply to “Height-related changes in forest composition explain increasing tree mortality with height during an extreme drought”. *Nature Communications*, 11(1):3401.
- Suarez, M. L., Ghermandi, L., and Kitzberger, T. (2004). Factors predisposing episodic drought-induced tree mortality in *Nothofagus*—site, climatic sensitivity and growth trends. *Journal of Ecology*, 92(6):954–966.
- Sørensen, R., Zinko, U., and Seibert, J. (2006). On the calculation of the topographic wetness index: evaluation of different methods based on field observations. *Hydrology and Earth System Sciences*, 10(1):101–112.
- Trenberth, K. E., Dai, A., van der Schrier, G., Jones, P. D., Barichivich, J., Briffa, K. R., and Sheffield, J. (2014). Global warming and changes in drought. *Nature Climate Change*, 4(1):17–22.
- Trugman, A. T., Detto, M., Bartlett, M. K., Medvigy, D., Anderegg, W. R. L., Schwalm, C., Schaffer, B., and Pacala, S. W. (2018). Tree carbon allocation explains forest drought-kill and recovery patterns. *Ecology Letters*, 21(10):1552–1560.
- Wheeler, E. A., Baas, P., and Rodgers, S. (2007). Variations In Dicot Wood Anatomy: A Global Analysis Based on the Insidewood Database. *IAWA Journal*, 28(3):229–258.
- Zach, A., Schuldt, B., Brix, S., Horna, V., Culmsee, H., and Leuschner, C. (2010). Vessel diameter and xylem hydraulic conductivity increase with tree height in tropical rainforest trees in Sulawesi, Indonesia. *Flora - Morphology, Distribution, Functional Ecology of Plants*, 205(8):506–512.
- Zellweger, F., Coomes, D., Lenoir, J., Depauw, L., Maes, S. L., Wulf, M., Kirby, K. J., Brunet, J., Kopecký, M., Máliš, F., Schmidt, W., Heinrichs, S., Ouden, J. d., Jaroszewicz, B., Buyse, G., Spicher, F., Verheyen, K., and Frenne, P. D. (2019). Seasonal drivers of understorey temperature buffering in temperate deciduous forests across Europe. *Global Ecology and Biogeography*, 28(12):1774–1786.

- 819 Zhu, S.-D., Chen, Y.-J., Ye, Q., He, P.-C., Liu, H., Li, R.-H., Fu, P.-L., Jiang, G.-F., and Cao, K.-F.  
820 (2018). Leaf turgor loss point is correlated with drought tolerance and leaf carbon economics traits. *Tree*  
821 *Physiology*, 38(5):658–663.
- 822 Zuleta, D., Duque, A., Cardenas, D., Muller-Landau, H. C., and Davies, S. J. (2017). Drought-induced  
823 mortality patterns and rapid biomass recovery in a terra firme forest in the Colombian Amazon. *Ecology*,  
824 98(10):2538–2546.