

Freshwater Science

Prioritizing management goals for stream biological integrity within the developed landscape context --Manuscript Draft--

Manuscript Number:	2018118
Corresponding Author:	Marcus Beck Southern California Coastal Water Research Project Costa Mesa, CA UNITED STATES
Other Authors:	Raphael D Mazor Scott Johnson Karin Wisenbaker Joshua Westfall Peter R Ode Ryan Hill Chad Loflen Martha Sutula Eric D Stein
Full Title:	Prioritizing management goals for stream biological integrity within the developed landscape context
Article Type:	Regular
Manuscript Region of Origin:	UNITED STATES
Abstract:	<p>Stream management goals for biological integrity may be difficult to achieve in developed landscapes where channel modification and other factors impose constraints on in-stream conditions. To evaluate potential constraints on biological integrity, we developed a statewide landscape model for California that estimates ranges of likely scores for a macroinvertebrate-based index that are typical at a site for the observed level of landscape alteration. This context can support prioritization decisions for stream management, like identifying reaches for restoration or enhanced protection based on how observed scores relate to the model expectations. Median scores were accurately predicted by the model for all sites in California with bioassessment data (Pearson correlation $r = 0.75$ between observed and predicted for calibration data, $r = 0.72$ for validation). The model also predicted that 15% of streams statewide are unlikely to achieve biological integrity within their present developed landscape, particularly for urban and agricultural areas in the South Coast, Central Valley, and Bay Area regions. We worked with a local stakeholder group from the San Gabriel River watershed (Los Angeles County, California) to evaluate how the statewide model could support local management decisions. To achieve this purpose, we created an interactive application, the Stream Classification and Priority Explorer (SCAPE), that compares observed scores with expectations from the landscape model to assign priorities. We observed model predictions that were consistent with the clear land use gradient from the upper to lower watershed, where potential limits to achieving biological integrity were more common in the heavily urbanized lower watershed. However, most of the sites in the lower watershed scored within their expected ranges, and were therefore given a low priority for restoration. In contrast, two low-scoring sites in the undeveloped upper watershed were prioritized for causal assessment and possible future restoration, whereas three high-scoring sites were prioritized for protection. The availability of geospatial and bioassessment data at the national level suggests that these tools can easily be applied to inform management decisions at other locations where altered landscapes may limit biological integrity.</p>



SOUTHERN CALIFORNIA COASTAL WATER RESEARCH PROJECT
A Public Agency for Environmental Research

September 10th, 2018

Dr. Charles Hawkins
Chief Editor
Freshwater Science

I am pleased to submit our manuscript, "Prioritizing management goals for stream biological integrity within the developed landscape context," to be considered as an original research article in Freshwater Science.

Many streams in urban and agricultural areas have degraded biological integrity and managing for reference conditions in developed landscapes may be a costly goal. This research addresses a critical need within the management community by providing a bioassessment tool that establishes a context of expectation for biological integrity in developed landscapes. Our model can be used to predict a range of expected scores for a biological index that can be compared to observed scores. Sites can then be ranked and prioritized relative to the expectation. We developed the landscape model for all stream reaches in California and worked with a regional monitoring program from a highly urbanized watershed to develop management priorities using results from the model. This model is an effective prioritization tool that can help managers identify stream sites for restoration, protection, or additional monitoring in the context of the developed landscape.

The data, text, and illustrations in this submission have not been used in existing or forthcoming papers or books. Our organization also agrees to submit payment for page charges if the paper is published. We are confident that readers of FWS will find this information useful and appreciate the opportunity to publish our work in this venue.

Marcus W. Beck (marcusb@sccwrp.org), 714-755-3217
S. California Coastal Water Research Project
3535 Harbor Blvd. Suite 110
Costa Mesa, CA 92626

On behalf of my co-authors:

Raphael D. Mazor (raphaelm@sccwrp.org), Scott Johnson (scott@aquaticbioassay.com), Karin Wisenbaker (karin@aquaticbioassay.com), Joshua Westfall (jwestfall@lacsdc.org), Peter R. Ode (peter.ode@wildlife.ca.gov), Ryan Hill (hill.ryan@epa.gov), Chad Loflen (Chad.Loflen@waterboards.ca.gov), Martha Sutula (marthas@sccwrp.org), Eric D. Stein (erics@sccwrp.org)

Running head: Stream priorities in developed landscapes

Prioritizing management goals for stream biological integrity within the developed landscape context

Marcus W. Beck (marcusb@sccwrp.org)^{1*}

Raphael D. Mazor (raphaelm@sccwrp.org)¹

Scott Johnson (scott@aquaticbioassay.com)²

Karin Wisenbaker (karin@aquaticbioassay.com)²

Joshua Westfall (jwestfall@lacsdsd.org)³

Peter R. Ode (peter.ode@wildlife.ca.gov)⁴

Ryan Hill (hill.ryan@epa.gov)⁵

Chad Loflen (Chad.Loflen@waterboards.ca.gov)⁶

Martha Sutula (marthas@sccwrp.org)¹

Eric D. Stein (erics@sccwrp.org)¹

* Corresponding author

¹ Southern California Coastal Water Research Project, 3535 Harbor Blvd., Costa Mesa, CA 92626

² Aquatic Bioassay & Consulting Laboratories, Inc. 29 North Olive, Ventura, CA 93001

³ Sanitation Districts of Los Angeles County, 1955 Workman Mill Road, Whittier 90601

⁴ California Department of Fish and Wildlife, Office of Spill Prevention and Response, 2005 Nimbus Road, Rancho Cordova, CA 95670

⁵ US Environmental Protection Agency, Office of Research and Development, National Health and Effects Research Laboratory, Western Ecology Division, 200 SW 35th Street, Corvallis, OR 97333

⁶ San Diego Regional Water Quality Control Board, Healthy Waters Branch, Monitoring and Assessment Research Unit, 2375 Northside Drive, San Diego, CA 92108

Abstract

Stream management goals for biological integrity may be difficult to achieve in developed landscapes where channel modification and other factors impose constraints on in-stream conditions. To evaluate potential constraints on biological integrity, we developed a statewide landscape model for California that estimates ranges of likely scores for a macroinvertebrate-based index that are typical at a site for the observed level of landscape alteration. This context can support prioritization decisions for stream management, like identifying reaches for restoration or enhanced protection based on how observed scores relate to the model expectations. Median scores were accurately predicted by the model for all sites in California with bioassessment data (Pearson correlation $r = 0.75$ between observed and predicted for calibration data, $r = 0.72$ for validation). The model also predicted that 15% of streams statewide are unlikely to achieve biological integrity within their present developed landscape, particularly for urban and agricultural areas in the South Coast, Central Valley, and Bay Area regions. We worked with a local stakeholder group from the San Gabriel River watershed (Los Angeles County, California) to evaluate how the statewide model could support local management decisions. To achieve this purpose, we created an interactive application, the Stream Classification and Priority Explorer (SCAPE), that compares observed scores with expectations from the landscape model to assign priorities. We observed model predictions that were consistent with the clear land use gradient from the upper to lower watershed, where potential limits to achieving biological integrity were more common in the heavily urbanized lower watershed. However, most of the sites in the lower watershed scored within their expected ranges, and were therefore given a low priority for restoration. In contrast, two low-scoring sites in the undeveloped upper watershed were prioritized for causal assessment and possible future

restoration, whereas three high-scoring sites were prioritized for protection. The availability of geospatial and bioassessment data at the national level suggests that these tools can easily be applied to inform management decisions at other locations where altered landscapes may limit biological integrity.

Key words: Bioassessment, biotic integrity, streams, urbanization, modified channels, landscape stressors, random forests, prioritization, data visualization, stakeholder group

Introduction

The widespread use of bioassessment data to assess ecological condition of aquatic environments is a significant advance over chemical or physical methods of assessment, yet managers and stakeholders require contextual information for synthesizing and interpreting biological information. The reference condition concept that is built into many biological indices provides a broad context for observed condition relative to unaltered habitats for a particular region (Reynoldson et al. 1997, Stoddard et al. 2006). However, achieving a reference condition of biological integrity (i.e., having structure and function comparable to natural habitat for the same region, Karr et al. 1986) may be challenging if site-specific conditions place limits on spatial and temporal scales that can be effectively managed (Chessman and Royal 2004, Chessman 2014). Use of bioassessment information to guide decisions that affect aquatic resources may also be challenging if the data are not accessible relative to the needs of local stakeholder groups. Accessibility can be limited from a contextual perspective of how likely a site is to achieve biological integrity, but also how bioassessment data collected over multiple locations and times can be used to support decisions or identify priorities. Explicit information is required to not only


synthesize site-level bioassessment data at the watershed scale, but also provide an assessment context that is sufficiently interpretable for prioritization.


In developed urban and agricultural landscapes, the majority of stream miles are in poor biotic condition and in need of some level of management (USGS 1999, Finkenbine, Atwater, and Mavinic 2000, Morgan and Cushman 2005). Conventional approaches to protect and restore biological integrity have commonly focused on direct improvements at the site level to mitigate instream stressors (Carline and Walsh 2007, Lester and Boulton 2008, Roni and Beechi 2012, Loflen et al. 2016), where stream preventive measures may be incentivized or enforced through regulation. Although these approaches can lead to improvements in ecological condition, there is no universal remedy for achieving biological integrity in streams. Restoring streams in urban or agricultural settings can be costly and it may be difficult to achieve regional reference-like conditions (Kenney et al. 2012, Shoredits and Clayton 2013). A confounding factor for managing streams in developed landscapes is the extensive modification to streams for flood control or water conveyance. In some cases, channel modification has been proposed as a basis for redefining water quality criteria or for re-evaluating use attainability goals (CRWQB 2014). For biological integrity, several states have implemented a tiered aquatic life use or alternative use designations to account for baseline shifts in ecosystem condition from channel modification (e.g., FDEP 2011, USEPA 2013, MBI 2016). Prioritizing among sites that are affected by landscape alteration is a critical challenge for managers in urban and agricultural settings (Walsh et al. 2005, Beechie et al. 2007, Paul et al. 2008).

The application of bioassessment data to inform management requires understanding the effects of multiple stressors acting at local, catchment, or watershed scales (Novotny et al. 2005, Townsend, Uhlmann, and Matthaei 2008, Leps et al. 2015). Nearly half of all stream-miles in the

93 USA are estimated to be in poor biotic condition based on macroinvertebrate bioassessment
94 index scores and ~~has been associated with~~ in-stream stressors, such as excess phosphorus,
95 nitrogen, or altered physical habitat (SEPA 2016). These ~~immediate~~ causes of poor biological
96 condition are often linked to landscape-level alterations that occur in the watershed. Consistent
97 and empirical links between land use thresholds and poor biotic integrity have been identified in
98 many cases (Allan, Erickson, and Fay 1997, Wang et al. 1997, Clapcott et al. 2011). Although
99 causal pathways linking land use and degraded biological condition have been described (e.g.,
100 Allan 2004, Riseng et al. 2011), not all pathways of stressors originating from the landscape are
101 clear (e.g., Cormier et al. 2013). Regardless, land use has long been used as a proxy for
102 environmental condition, and an associative link can be sufficient to predict condition as a
103 function of watershed activities.

104 Estimating the likely range of biological conditions as a function of historic alteration of the
105 landscape could help prioritize where management actions are most likely to achieve intended
106 outcomes, or conversely, where landscape alteration could limit management success in
107 achieving biological integrity. Here, we define constrained streams as those where reference
108 conditions for the biological community may be difficult to achieve with limited resources
109 because of large-scale, historical impacts from landscape alteration. Anthropogenic stressors that
110 constrain biology may originate from spatial or temporal scales that are difficult to address with
111 most management applications. Understanding limits to biological potential is one approach to
112 identify constraints, and is an important concept in bioassessment that has received some
113 attention. Analysis methods have been explored in a bioassessment context to characterize
114 environmental factors that limit assemblage composition (Chessman, Muschal, and Royal 2008,
115 Chessman 2014). This approach is based on the limiting factor theory that proposes the most

116 limiting biotic or abiotic factor as the primary regulator of species abundance and distribution.
117 Similar concepts have been applied in a landscape context to understand both variation in
118 bioassessment data at different spatial scales and **limits**  bioassessment tools with land use
119 gradients (Waite 2013, Waite et al. 2014). Applying these concepts in a predictive framework
120 could facilitate an expectation of bioassessment and management potential relative to a site-
121 specific context.

122 The development of modelling tools for understanding biological condition across landscape
123 gradients could provide a powerful approach to informing the use of limited resources to manage
124 stream integrity. Previous modelling efforts for bioassessment have successfully used geospatial
125 data to predict biological condition at regional or national scales (Vølstad et al. 2004, Carlisle,
126 Falcone, and Meador 2009, Brown et al. 2012, Hill et al. 2017), with the general purpose of
127 characterizing condition at unsampled locations. Macroinvertebrate communities can respond
128 predictably to landscape alteration (Sponseller, Benfield, and Valett 2001, Waite 2013) and
129 association of biological condition with landscape metrics that describe these changes could be
130 used to predict a range of expectations for biotic integrity as related to observed watershed
131 development. This approach differs fundamentally from previous efforts of estimating average
132 condition by providing an estimate of the minimum and maximum scores that are likely for the
133 landscape context nce the responses of macroinvertebrate communities to landscape changes
134 at large spatial scales are understood, expectations can be compared to field samples and sites
135 can be prioritized by local managers based on deviation from the expectation.

136 The goal of this study is to present the development and application of a landscape model to
137 classify and prioritize stream monitoring sites based on probable ranges of bioassessment scores
138 relative to landscape alteration. This model is presented as a screening tool for exploring

different priorities and is not intended for developing regulatory designations nor determining if a site can attain designated uses. The specific objectives were to 1) demonstrate development of a landscape model to predict expected ranges of biotic condition, 2) classify stream segments into biological constraint categories using modelling expectations, 3) assess the extent of stream classes and explore the sensitivity of the classifications to decision points in the model output, and 4) prioritize potential management decisions by comparing expectations to observed bioassessment scores. The model was developed and applied to all streams and rivers in California, specifically focusing on the potential of urban and agricultural land use to impact biological condition. We include a case study that demonstrates how the statewide model can be used to classify and prioritize in a regional context using guidance from a local stakeholder group from a heavily urbanized watershed where obstacles for achieving biological integrity have been encountered. In interactive software application, the Stream Classification and Priority Explorer (SCAPE), is also described that was developed to help choose management priorities using the landscape model.

Methods

Study area and data sources

The landscape model was developed for California using land use data, stream hydrography, and biological assessments. California covers 424,000 km² of land with extreme diversity in several environmental gradients, such as elevation, geology, and climate (Figure 1a, Ode et al. 2016). Temperate rainforests occur in the north (North Coast region), deserts and plateaus in the northeast and southeast (Deserts and Modoc Plateau region), and Mediterranean climates in

160 coastal regions (Chaparral and South Coast regions). The Central Valley region is largely
161 agricultural and drains a large mountainous area in the east-central region of the state (Sierra
162 Nevada region). Urban development is concentrated in coastal areas in the central (San Francisco
163 Bay Area, Chaparral region) and southern (Los Angeles, San Diego metropolitan area, South
164 Coast) regions of the state. California's stream network is approximately 280,000 km in length
165 and covers all of the major climate zones in the state. A high degree of endemism and
166 biodiversity occurs in these streams including nearly 4000 species of vascular plants,
167 macroinvertebrates, and vertebrates that depend on fresh water during their life history (Howard
168 and Revenga 2009, Howard et al. 2015). Approximately 30% of streams in California are
169 perennial with the remaining as intermittent or ephemeral.

170 Landscape alteration has been relatively recent, with one estimate showing that developed lands
171 have increased in California by 38% from 1973 to 2000 (Sleeter et al. 2011). Development prior
172 to 2001 was generally not required to incorporate stormwater structural mitigation measures,
173 such as site design and treatment controls, which are now required statewide to match hydrologic
174 flows and to treat and prevent pollutants from leaving developed areas (SDRWQB 2001). For
175 analysis, the state was evaluated as a whole and by major regions defined by hydrological and
176 geopolitical boundaries (Figure 1a): Central Valley (CV), Chaparral (CH), Deserts and Modoc
177 Plateau (DM), North Coast (NC), Sierra Nevada (SN), and South Coast (SC). Some of these
178 regions have large urban areas (SC, CH) or agriculture (CV), whereas others are largely forested,
179 but may be impacted by silviculture or logging (NC, SN).

180 Stream data from the National Hydrography Dataset Plus (NHD-plus) (McKay et al. 2012) were
181 used to identify stream segments in California for modelling biological integrity. The NHD-plus
182 is a surface water framework that maps drainage networks and associated features (e.g., streams,

lakes, canals, etc.) in the United States. Stream segments designated in the NHD-plus were used as the discrete spatial unit for modelling biological integrity. Here and throughout, “segment” is defined in the context of NHD-Plus flowlines. Hydrography data were combined with landscape metrics available from the StreamCat Dataset (Hill et al. 2016) to estimate land use at the riparian zone (i.e., a 100-m buffer on each side of the stream segment), the catchment (i.e., nearby landscape flowing directly into the immediate stream segment, excluding upstream segments), and the entire upstream watershed for each segment. Many of the metrics in StreamCat were derived from the 2006 National Land Cover Database (Fry et al. 2011).

The California Stream Condition Index (CSCI) (Mazor et al. 2016) was used as a measure of biological condition in California streams. The CSCI is a predictive index that compares the observed taxa and metrics at a site to those expected under reference conditions. Expected values at a site are based on models that estimate the likely macroinvertebrate community in relation to factors that naturally influence biology, e.g., watershed size, elevation, climate, etc. (Moss et al. 1987, Cao et al. 2007). The index score at a site can vary from 0 to ~ 1.4, with higher values indicating less deviation from reference state. Because the index was developed to minimize the influence of natural gradients, the index scores have consistent meaning across the state (Mazor et al. 2016). A CSCI threshold of 0.79, based on the tenth percentile of scores at all reference calibration sites, has been used to identify stream degradation by state regulatory agencies (SDRWQB 2016) and was used herein to represent a potential management target.

Benthic macroinvertebrate data were used to calculate 6270 individual CSCI scores at nearly 3400 unique sites between 2000 and 2016 (Figure 1b). Samples were collected during base flow conditions typically between May and July following methods in Ode et al. (2016).

Bioassessment sites were snapped to the closest NHD-plus stream segment in ArcGIS (ESRI

206 2016). In cases where multiple sites were located on the same segment, the most downstream site
207 was selected for further analysis under the assumption that the landscape data in StreamCat was
208 most relevant to this site. This created a final dataset of 2620 unique field observations used to
209 calibrate and validate the landscape model.

210 **Building and validating the landscape model**

211 A quantile random forest model was developed to estimate ranges of CSCI scores associated
212 with land use gradients, such as road density or urban and agricultural land use. Measures of land
213 use development were quantified for riparian, catchment, and watershed areas (as defined above)
214 of each stream segment in California using the StreamCat dataset (Hill et al. 2016). Expected
215 CSCI scores were modelled using estimates of canal/ditch density, imperviousness, road
216 density/crossings, and urban and agricultural land use for each stream segment (Table 1). These
217 variables were chosen specifically to model scores only in relation to potential impacts on
218 biological condition that are typically beyond the scope of management intervention or where
219 costs to mitigate are likely prohibitive. Potential effects on biological condition that may vary
220 through time or from stressors not associated with urban or agricultural land use were not
221 captured by the model (e.g., timber harvesting). Similarly, potential differences in the magnitude
222 of effects on stream condition for the chosen variables were also not explicitly evaluated, such
223 that all variables were given equal weighting in the models. Within these limits, we considered
224 deviation of observed scores from model predictions to be diagnostic of human activity not
225 related to anthropogenic stressors that can be measured on the landscape in addition to potential
226 model error. Methods for evaluating predictive performance of the model are described below.

The model was developed using quantile regression forests to estimate ranges of likely CSCI scores in different landscapes (Meinshausen 2006, 2017). Random forests are an ensemble learning approach to predictive modelling that aggregates information from a large number of regression trees and have been used extensively in bioassessment applications (Carlisle, Falcone, and Meador 2009, Chen et al. 2014, Mazor et al. 2016, Fox et al. 2017). Random forest models provide robust predictions by evaluating complex, non-linear relationships and interactions between variables relative to more commonly-used modelling approaches, such as multiple regression (Breiman 2001, Hastie, Tibshirani, and Friedman 2009). Quantile models, such as quantile regression forests, evaluate the conditional response across the range of values that are expected, in contrast to conventional models that provide only an estimate of the mean response (Cade and Noon 2003). This modelling approach allows use of prediction intervals to describe the range of likely scores, which can be used to identify sites where that range includes management targets. Quantile regression forests were used to predict CSCI scores in each stream segment at five percent increments (i.e., 5th, 10th, etc.) from the 5th to 95th percentile of expectations. The quantregForest package for the R Statistical Programming Language was used to develop the landscape model using the default settings, with the exception that out of bag estimates were used for model predictions (Meinshausen 2017, ~~RDCT 2018~~).

We stratified sample data to ensure sufficient representation of landscape gradients major region in the state and across percentiles of catchment imperviousness (Figure 1). Calibration data for the landscape model were obtained from a random selection of 75% of segments with observed CSCI scores across this stratification and where sufficient data were available in StreamCat (n = 1965 segments). The remaining sites were used for model validation (n = 655). Where multiple samples were available at a single site, one sample was selected at random for both calibration



and validation purposes. Model performance was assessed for the statewide dataset and within each major region by comparing differences between observed CSCI scores and median predictions at the same locations. Differences were evaluated using Pearson correlations and root mean squared errors (RMSE); high correlation coefficients and low RMSE values indicated good performance. Regression analysis between predicted and observed scores was used to assess potential bias based on intercept and slope values differing from 0 and 1, respectively. Collectively, the performance metrics were chosen to evaluate both predictive ability of the landscape model and potential for bias which may vary depending on different land use gradients across the state.

Statewide application of the landscape model

We applied the landscape model to 138,716 stream segments statewide to estimate the extent of streams in one of four different constraint classes: likely unconstrained, possibly unconstrained, possibly constrained, and likely constrained (Table 2). Here and throughout, constrained is defined as a biological community that is impacted by large-scale, historic alteration of the landscape. Consequently, achieving biological integrity in constrained communities may present management challenges given that many stressors in altered landscapes originate at spatial or temporal scales that are typically beyond the scope of most management applications or where resources for mitigation may be prohibitive.

The classification process is described in Figure 2a through c. Classifications were based on the comparison of a CSCI threshold representing a management goal and the predicted range or predicted median score at a segment. These two decision points (i.e., the threshold and the size of the predicted range) were critical in defining segment classifications. For most analyses, we used

a CSCI threshold of 0.79 (i.e., the 10th percentile of reference calibration sites) following previous examples (Mazor et al. 2016, SDRWQB 2016) and a prediction interval ranging from the 10th to the 90th percentiles. Stream segments with the range of CSCI score expectations entirely below the threshold were considered likely constrained whereas those with expectations entirely above were considered likely unconstrained (Figure 2c). The remaining sites were classified as possibly unconstrained or possibly constrained, based on whether the median expectation was above or below the threshold (respectively) (Table 2).

A sensitivity analysis was conducted to evaluate the influence of these key decision points on the extent of segment classifications created by the landscape model. Stream segment classifications depend on the chosen range of score expectations (or certainty) from the landscape model (Figure 2b) and the CSCI threshold for evaluating the overlap extent (Figure 2c). Eight different ranges of values for the score expectations from wide to narrow were evaluated at five percent intervals, i.e., 5th-95th, 10th-90th, ..., 45th-55th. Different CSCI thresholds were also evaluated using values of 0.63, 0.79, and 0.92, corresponding to the 1st, 10th, and 30th percentile of scores at reference calibration sites used to develop the CSCI (Figure 1b) (Mazor et al. 2016). The percentage of stream segments in each class statewide and by major regions were estimated for each of the twenty-four scenarios (width by threshold combinations) to evaluate sensitivity to changes in the decision points.

Sites were further classified by comparing observed CSCI scores from biomonitoring data to the range of expected scores (Figure 2d). Relative site scores were determined based on location of the observed score to the range of expected CSCI scores. Sites with observed scores above the upper limit of the segment expectation (e.g., above the 90th percentile of expected scores) were considered “over-scoring” and sites below the lower limit (e.g., 10th percentile) were considered

“under-scoring”. If neither “over-scoring” nor “under-scoring”, the relative site score was considered as “expected” within the context of the landscape model.

Defining management priorities in the San Gabriel River watershed

Site and stream classifications from the landscape model allowed a local stakeholder group to develop a framework for evaluating data from a watershed monitoring program to prioritize management actions. The San Gabriel River (SGR) Regional Monitoring Program (Los Angeles County, California) includes stakeholders from water quality regulatory agencies, municipalities, and non-governmental organizations that cooperatively work to manage aquatic resources in the watershed and improve coordination of compliance and ambient monitoring efforts. The workgroup met monthly over a six-month period to discuss model application and to refine the interpretation of results. The model was applied to 751 stream segments in the watershed, of which 147 samples at 75 segments were collected for bioassessment (Figure 3a). CSCI scores ranged from 0.2 to 1.23 and were averaged for repeat visits, of which sixty segments had only one visit. Fifty-six samples from the SGR watershed were used in the statewide dataset to develop the landscape model.

A strong land-use gradient occurs in the SGR watershed that creates challenges for managing stream condition (Figure 3b). The upper watershed in the San Gabriel mountains is largely undeveloped or protected for recreational use, whereas the lower watershed is in a heavily urbanized region of Los Angeles County. The SGR is dammed at four locations in the upper watershed for flood control. Spreading grounds in the middle of the watershed are used to recharge groundwater during high flow. As a result, the upper and lower watersheds are hydrologically disconnected when annual rainfall is normal. Nearly all of the stream segments in

the lower half of the watershed are channelized with concrete or other reinforcements. The majority of flow in the lower watershed is provided to the mainstem and major tributaries of the SGR by wastewater treatment plants releasing tertiary treated effluent. Approximately half of the monitored sites in the watershed are in poor biological condition, nearly all of which are in the lower watershed.

Stakeholders identified their relevant priorities by evaluating the different site types that were possible from the landscape model relative to the stream classes. The priorities defined by the group were generalized into three categories:

- Investigate: Conduct additional monitoring or review of supplementary data (e.g., field visits, review aerial imagery);
- Protect: Recommend additional scrutiny of any proposed development and/or projects;
- Restore: Pursue targeted action for causal assessment and/or restoration activity.

A template that showed the possible site scores relative to the segment classifications was given to the stakeholders (Figure S1, left side). The three priorities were then assigned a low, medium, or high importance for the scoring possibilities that could occur from the landscape model (Figure S1, right side). The assignments were made with the explicit recognition that any priority recommendations were in addition to baseline monitoring and maintenance that is currently provided by existing management programs. The final assignments were then mapped to each monitoring site in the watershed.

The outcomes of these assignments were visualized in an interactive and online application, the Stream Classification and Priority Explorer (SCAPE, Figure S2, <http://shiny.sccwrp.org/scape/>, Beck 2018). The application allowed stakeholders to provide input on the two key decision

points for classifying stream segments (i.e., choice of a threshold and a prediction interval), as well as to assign priorities to each management action described above. The application then allowed stakeholders to see the outcomes of these decisions. Specifically, SCAPE created maps showing the classifications for segments in the watershed, deviation of observed CSCI scores from the expectation, and maps of recommended priority actions that were assigned to each of the scoring possibilities. In addition, the application tabulated the extent of streams in each class, as well as the number of sites prioritized for each management action. Crucially, SCAPE allowed the stakeholders to modify key decisions points in the model and rapidly evaluate how these changes propagated to changes in recommended priorities for each site.

Results

Model performance

Model performance statewide indicated generally good agreement between observed CSCI scores and the median prediction for the associated stream segment (Table 3). Agreement between observed and predicted values for the entire calibration dataset was $r = 0.75$ (Pearson) and $RMSE = 0.17$. The intercept and slope for a regression between observed and predicted values were 0.34 and 0.60, suggesting a slight negative bias of predictions at lower scores and slight positive bias at higher scores. The statewide validation data showed similar results, with slightly smaller correlation ($r = 0.72$) and larger $RMSE$ (0.18) estimates.

Overall, the model performed well in regions with a mix of urban, agricultural, and open land (e.g., South Coast and Chaparral regions), whereas performance was weakest in regions without strong development gradients (e.g., Sierra Nevada and North Coast regions) (Table 3, Figure S3).

Performance for the Chaparral and South Coast regions were comparable or slightly improved compared to the statewide dataset for both the calibration ($r = 0.71, 0.75$, respectively) and validation ($r = 0.74, 0.72$) datasets. Model predictions for the Central Valley, ~~Desert/Modoc, and North Coast~~ regions had slightly lower performance compared to the statewide results, with correlations of approximately 0.57 with observed values in the calibration dataset and 0.53 in the validation dataset. Model performance was weakest for the Sierra Nevada and North Coast regions, where timber harvesting, rather than urban or agricultural development, is the most widespread stressor.

Statewide patterns in stream constraints

Statewide patterns in stream constraints were apparent from the results of the landscape model that were consistent with land use (Figure 4). A majority of stream segments statewide were classified as possibly constrained (11% of all stream length) or possibly unconstrained (46%), whereas a minority were likely constrained (4%) or likely unconstrained (39%) (Table 4). Large rivers across the state were more commonly classified as possibly constrained (e.g., Klamath, Owens, and Russian rivers). Overall, stream segments were more often constrained for biotic integrity in regions with more development, either as urban or agricultural land. For example, likely unconstrained streams were common in the Sierra Nevada (50%), North Coast (46%), and Desert/Modoc (46%) regions, whereas likely constrained were relatively abundant in the Central Valley (22%) and South Coast (15%) regions. However, constrained and unconstrained streams were both found in every region (Figure 4)

Observed CSCI scores were within the predicted range as often as expected (i.e., 80% statewide, based on the 10th and 90th prediction interval), and over-scoring sites were roughly as common

(9%) as under-scoring sites (10%) (Table 5). Similar patterns were observed within regions, although a slightly larger percentage of sites in the Central Valley were under-scoring compared to the other regions. Over-scoring sites were slightly more common in certain regions (i.e., the South Coast and Sierra Nevada regions) than others (i.e., the Chaparral, Central Valley, and Desert/Modoc regions).

Sensitivity analyses underscored the influence of key decision points of the landscape model on estimates of the extent of streams in each class (Figure 5). Unsurprisingly, decreasing the certainty of predictions from the landscape model by narrowing the prediction interval (5th-95th to 45th-55th) shifted a number of streams from the possible to likely category in both constrained and unconstrained segments. Similarly, changing the CSCI threshold from relaxed to more conservative (0.63 to 0.92) increased the number of streams classified as possibly or likely constrained and decreased the number of streams as possibly or likely unconstrained. However, the sensitivity to these decision points varied greatly by region. For example, over 80% of segments in the Central Valley were classified as likely constrained using a high CSCI threshold with the narrowest range of predictions, whereas less than 1% of segments were in this category using a low CSCI threshold with the widest range of predictions. Opposite trends were observed in regions with reduced land use pressures. For example, almost all stream segments in the North Coast and Sierra Nevada regions were classified as likely unconstrained using a low CSCI threshold and narrow range of predictions.

San Gabriel River Case study

Application of the landscape model results to the CSCI scores provided a context of expectations consistent with the strong land use gradient in the watershed (Figure 6). Stream segments in the

upper watershed were a mix of likely and possibly unconstrained (40% and 28%), whereas stream segments in the lower watershed were classified as likely and possibly constrained (25% and 7%). Several segments in the lower watershed had median CSCI scores that were very close to the 10th percentile (i.e., right-skewed) consistent with extreme landscape pressures (bottom left, Figure 6b).

Using the same classification decision points described above for the statewide model, only six sites were under-scoring (two likely unconstrained and four likely constrained) and eight sites were over-scoring (five likely constrained, one possibly unconstrained, and two likely unconstrained) (Figure 7, top). One of the under-scoring sites in the likely unconstrained class was below the CSCI threshold (Figure 6). One site scoring as expected in the possibly unconstrained class was below the chosen CSCI threshold, whereas none of the constrained (possibly or likely) sites were above the threshold.

The SCAPE application was effectively used to select management priorities for all monitoring sites in the SGR watershed. In general, the stakeholder group assigned high priority recommendations to over- and under-scoring sites in likely unconstrained segments or those below the biological threshold with possibly unconstrained classification (Figure S1). Continuing current practices (e.g., routine monitoring) were generally recommended at constrained sites or restoration actions were recommended as a lower priority despite low CSCI scores.

Recommended actions to investigate were more common for both over-scoring and under-scoring sites, protect was given a high priority exclusively at over-scoring sites, and restore was more common at under-scoring sites.

The SCAPE application also allowed the stakeholders to identify spatial patterns among the watershed priorities. For example, a clear distinction between low and high priority actions was

observed on the watershed map (Figure 7, bottom). Sites in the lower watershed were lower priority if an action was recommended, whereas the five high priority sites were in the upper watershed (multiple recommendations were assigned to the sites). The distinction between lower and higher priorities between the lower and upper watershed was driven exclusively by the segment classifications, where constrained segments were in the lower watershed and unconstrained segments were in the upper watershed. Several sites that were scoring as expected for likely and possibly unconstrained segments in the upper watershed were recommended as medium priority for protection.

Discussion

The prevalence of degraded streams in California requires the use of 1) assessment tools that can accurately evaluate condition, and 2) tools that can provide a context for evaluating the range of likely scores associated with different settings. The landscape model was developed with these needs in mind to better inform application of the CSCI for decision-making in the context of landscape constraints on biological condition. Statewide application of the model demonstrated where streams are likely constrained on a regional basis, whereas application to the SGR watershed demonstrated how the model can be used by local stakeholders to prioritize management actions that are informed by landscape context. Most importantly, the analyses underlying the model do not diagnose causes of impairment, nor do they justify by themselves an exemption from management intervention where constraints are high. The landscape model can inform the interpretation of biotic condition and is an exploratory tool that can help identify where management goals are more likely to be achieved.

Results from our analysis could be used for managing the biological integrity of streams under state or federal water quality mandates (e.g. “biological criteria” under the Clean Water Act). Regulatory management for biological integrity involves the protection of sites meeting biological objectives and the restoration of sites that do not meet biological objectives. The selection of appropriate regulatory management actions for streams requires the consideration of the physical and chemical condition of streams concurrent with biological monitoring results. The landscape model can evaluate sites that are or are not meeting biological objectives relative to their modeled condition. This information could provide flexibility in the selection of regulatory or management actions at specific sites or watershed scales (e.g., hydrologic subareas), and to further prioritize where and when actions should take place based on the temporal and spatial scale needed for protection or restoration actions. For example, for sites that meet biological objectives but where the models predict some degree of constraint (e.g., Figure S1, site types 5, 9, 10, or 13), regulatory actions may be associated with protecting that condition and could be implemented in the short-term to prevent degradation. This flexibility is not intended to exclude sites from consideration that are less likely to achieve biological objectives, but rather to facilitate the decision-making process through a more transparent application of the model in a regulatory context.

Non-regulatory applications of the landscape model are also possible by identifying where additional restoration, monitoring, or protection may have the most benefit. For example, landscape models could be used to support conservation planning, particularly at the watershed scale where land use practices can be a critical factor for decision-making. Ongoing work in California has focused on setting priorities for managing biodiversity that focus on watersheds within a conservation network (Howard et al. 2018). Results from the landscape model could be

used to enhance this network by providing supporting information on constraints in an assessment framework. More generally, these applications could represent a novel use of bioassessment data beyond the pass/fail paradigm of the regulatory context, for example, as tools for land use planning (Bailey et al. 2007). In many cases, including California, bioassessment indices have been sufficiently developed to allow large-scale condition assessment across regions, yet they are rarely used as planning tools to guide decisions on where resources should be focused (Nel et al. 2009). Our landscape model makes bioassessment data in California more accessible and identifies an appropriate context for the information, enabling the potential for both regulatory and non-regulatory applications.

The landscape model is a tool for exploring options



The primary objective of developing the landscape model was to provide a screening tool for exploring biological constraints to facilitate a discussion of management options relative to site contexts. This model by itself is not intended for direct application of regulatory designations at individual sites, nor is it fully adequate to assess whether a site can attain a particular use. Instead, the model can help identify patterns among monitoring sites where more intensive analyses may be appropriate or assist with decisions of where a use attainability assessment may be warranted. This application was effectively demonstrated through engagement of our local stakeholder group. Rather than identifying individual sites in need of specific management actions, the group used the landscape model to characterize patterns on the landscape that were consistent with the recommended management priorities. In doing so, the group was able to explore and discuss potential management actions relative to the landscape context of the watershed. The final decision by the group to prioritize management actions for the different

sites in broad categories of protect, restore, and investigate was based on an iterative process where ideas were discussed and shared freely among stakeholders. This approach ensured that stakeholders were generally in agreement with the final product and, therefore, potentially more likely to adopt the recommendations provided by these tools in formal decision-making (Stein et al. 2017). The recommended actions have relevance only in the context of interests of the SGR Regional Monitoring Program. Localized applications of the statewide model must engage stakeholders in a similar process to develop recommendations that are specific to regional needs at the watershed scale (Brody 2003, Reed 2008).

The development of the SCAPE application was also critical for applying the landscape model by synthesizing a large volume of bioassessment data. The application provided a means of demonstrating core concepts of the model and allowed stakeholders to explore the key decision points that affect the model output, specifically related to changing certainties in the CSCI score predictions and the ability to explore alternative thresholds for biological objectives. This functionality allowed the stakeholders to develop recommendations that were completely independent of the model, i.e., decisions were not hard-wired into the model nor SCAPE.

Because of this application, this stakeholder group has a better understanding of the potential impacts of biointegrity policies currently under review in California. Additionally, the SCAPE application provided assurance to the prioritization process by correctly identifying sites where discrepancies between CSCI scores and other measures of stream condition had been observed. Without this context (i.e., Figure 6a), stakeholders struggled to prioritize among sites, particularly for restoration activities. For example, some advocated that the lowest scoring sites should be prioritized, whereas others prioritized sites that scored just below the CSCI threshold.

515 Conflicting priorities were common in the absence of information about the range of scores
516 typical for these urban settings.

517 Several states have implemented alternative use designations for applying bioassessment criteria
518 in modified channels (FDEP 2011, USEPA 2013, MBI 2016). Although our results generally
519 support the link between impacted biology and channel modification, a regulatory framework
520 based on direct channel modification or other measures of channel morphology may be
521 insufficient by failing to recognize constraints on urban streams with natural morphology.  In the
522 context of the model, a constrained channel may or may not be engineered, but an engineered
523 channel will typically be constrained given the surrounding land use. For example, Tecolote
524 Creek (San Diego County, USA) was identified by our model as a constrained channel in an
525 urban landscape (Figure 8). The CSCI score is 0.61 indicating degraded biological integrity,
526 whereas the in-stream physical habitat is unaltered (Rehn, Mazor, and Ode 2018). Other stressors
527 originating at the landscape scale (e.g., water or sediment chemistry) have likely constrained the
528 biological community at this site independent of the physical habitat quality. Furthermore,
529 channel modification does not always result in biological degradation, particularly if the
530 contributing watershed is largely undeveloped. For example, Stein et al. (2013) observed
531 reference-like bioassessment index scores in armored reaches within national forest lands in
532 southern California. A classification framework for biological constraints using only channel
533 modification would provide incomplete and potentially misleading information on streams with
534 limited biological potential. Ideally, context from a landscape model, in conjunction with reach-
535 specific data on channel modification, should be used to determine where aquatic life uses may
536 be limited. 



Our approach to assessing constrained streams is readily transferable outside of California. The landscape model could be applied to other bioassessment methods, such as a multi-metric index (the most common bioassessment approach within the US, Buss et al. 2014), O/E assessments (Moss et al. 1987), biological condition gradients (Davies and Jackson 2006), or with other biological endpoints (e.g., fish or diatoms). More importantly, our use of national geospatial datasets (i.e., NHDPlus, McKay et al. 2012; StreamCat, Hill et al. 2016) means that these methods could be applied across the United States. National bioassessment indices have been developed and the landscape model could be developed as a national-scale product of constraints on biological condition to complement recent work that predicted probable biological conditions with the National Rivers and Streams Assessment (Hill et al. 2017). Global geospatial datasets of freshwater-specific environmental variables are also available and could be used to develop similar models outside of the United States (Domisch, Amatulli, and Jetz 2015).

Extension of the landscape models beyond California should also consider landscape stressors that are predictive of biotic condition in other regions. For example, urban and agricultural gradients were sufficient to characterize constraints in many regions of California, whereas Hill et al. (2017) found that the volume of water stored by dams was an important predictor of biological condition in the Northern Appalachian and Northern Plains regions of the US. In their paper, Hill et al. (2017) provided an example of how predictive models could be used to identify potential sites for restoration or conservation, however, their illustration did not explicitly identify sites that were over- or under-scoring relative to a biological endpoint. Doing so in California provided stakeholders with important context that helped establish management priorities, demonstrating the potential utility of this approach in other states.

Model assumptions and limitations

There are several characteristics of the landscape model that could affect its performance when applied outside of urban and agricultural settings. First, the model was developed with a focus on the needs of managers that apply bioassessment tools in developed landscapes where conditions are presumably constrained. As such, landscape variables were chosen to capture the effects of development on CSCI scores in these areas (Table 1). Application of the model in regions where different stressors have strong impacts on stream condition should consider the relevance of urban and agricultural stressors and if an alternative model that better captures other stressor gradients is needed. For example, our results suggest that streams in the North Coast and Sierra Nevada regions are largely unconstrained, but the landscape model was a poor predictor of CSCI scores in these areas. The dominant stressors likely to affect stream condition in these regions originate from sources that are less common in developed landscapes, such as silviculture and cannabis cultivation. The current landscape model does not adequately capture these impacts outside of urban and agricultural environments. Moreover, poor model predictions are compounded by low sensitivity of the CSCI to relevant stressor gradients in these regions (Mazor et al. 2016). Accurate data for quantifying these potential stressors are not available in StreamCat, but this is an area where investments in improving spatial data could yield significant improvements in further development of bioassessment indices and tools for their interpretation.

An additional assumption is that the landscape model can adequately discriminate between intractable constraints on biology that are spatially and temporally pervasive relative to more manageable constraints. That is, we assumed that the impacts of stressors included in the model, such as urbanization, are not manageable in the short term, whereas stressors associated with

581 deviations from model predictions can be mitigated.  These assumptions are not unique to our
582 model and have been used in other applications that have evaluated biological potential (Paul et
583 al. 2008, Chessman 2014, Waite et al. 2014  However, many stressors excluded from the model
584 can have long-lasting impacts, leading to potentially irreversible degradation or management
585 scenarios where long-term recovery may only be possible with sustained and costly application
586 of resources. For example, logging activities can impact benthic macroinvertebrate communities
587 for a decade or more after harvesting activities have stopped (Stone and Wallace 1998, Quinn
588 and Wright-Stow 2008). In urban areas, pervasive and profound alteration to groundwater and
589 hydrology is common and stream communities in groundwater fed systems may require
590 substantial time and resources for restoration. The potential legacy impacts of large-scale
591 alterations of the natural environment are not well-captured by the current model, neither from a
592 spatial nor temporal perspective. A more refined application of the landscape model would be
593 necessary to evaluate different scales of impact, which could include developing separate models
594 for each region, as well as more careful selection of model inputs to capture scales of interest for
595 potential impacts on stream condition.

596 The landscape model is associative by design and does not identify mechanistic links between
597 biological constraints and proximal causes. The model describes constraints at scales larger than
598 instream characteristics as a necessary approach to accurately predict bioassessment scores.
599 More comprehensive assessments at individual sites are needed to diagnose the immediate
600 causes of degraded condition. Further, a distinction between constraints on biological condition
601 and channel modification is implicit such that indication of the former by the model does not
602 explicitly indicate presence of the latter. As noted above, our results consistently indicated that
603 engineered channels are biologically constrained, but the model is based on an a priori selection

of land use variables to predict biotic integrity. A correspondence between habitat limitations and channel modification is likely in many cases but data are insufficient to evaluate biological effects statewide relative to land use constraints. Moreover, bioassessment scores can be similar in modified channels compared to natural streams independent of watershed land use, i.e., concordance between degraded stream condition and channel modification may not always be observed (Stein et al. 2013).

An additional consideration in using the landscape model is the meaning of biologically constrained in the context of whole stream communities. Biologically constrained sites were considered those where present landscapes were likely to limit CSCI scores that describe macroinvertebrate condition.

In many cases, poor biotic condition of the macroinvertebrate community translates to poor stream condition. However, a constrained macroinvertebrate community does not always mean other biological attributes of stream condition (e.g., fish assemblages) are also constrained. Urban streams sometimes support diverse algal assemblages such that algal-based measures of biotic condition may alternatively suggest good biotic condition relative to macroinvertebrate-based indices (Brown et al. 2009, Mazor, Beck, and Brown 2018). Broadening the landscape model to include multiple taxonomic assemblages or endpoints would allow a more complete assessment of how condition relates to landscape alteration.

Summary

The landscape model can be used to characterize the extent of biologically constrained channels in urban and agricultural landscapes. Our application to the SGR watershed demonstrated how the results of the model can be used at a spatial scale where many management decisions are

implemented through close interaction with a regional stakeholder group with direct interests in the local resources. Overall, the model provides a tool to determine how managers can best prioritize limited resources for stream management by focusing on segments where recommended actions are most likely to have the intended outcome of improving or protecting biological condition. The approach also leverages information from multiple sources to develop a context for biological assessment that provides an expectation of what is likely to be achieved based on current land use development. This can facilitate more targeted management actions that vary depending on the identified context and can also inform decisions on extent and effort for future monitoring locations.

Supplement

The SCAPE model application website: <http://shiny.sccwrp.org/scape/>, full source code accessible at Beck 2018. Additional figures and tables are available in the supplement.

Author contributions

MB, RM, SJ, KW, JW, PO, RH, CL, MS, and ES performed the research and analyzed the data. MB, RM, SJ, JW, PO, RH, and CL wrote the paper. RM, SJ, KW, and PO provided data. All authors discussed the methods and results and contributed to the development of the manuscript.

Acknowledgments

The authors acknowledge support from the San Gabriel River Regional Monitoring Program and the California State Water Resources Control Board. We thank Phil Markle and Lester Yuan for reviewing an earlier draft of the manuscript.

References

- Allan, D., D. Erickson, and J. Fay. 1997. The Influence of Catchment Land Use on Stream Integrity Across Multiple Spatial Scales. *Freshwater Biology* 37 (1):149–61. <https://doi.org/10.1046/j.1365-2427.1997.d01-546.x>.
- Allan, J. D. 2004. Landscapes and Riverscapes: The Influence of Land Use on Stream Ecosystems. *Annual Review of Ecology, Evolution, and Systematics* 35:257–64. <https://doi.org/10.1146/annurev.ecolsys.35.120202.110122>.
- Bailey, R. C., T. B. Reynoldson, A. G. Yates, J. Bailey, and S. Linke. 2007. Integrating Stream Bioassessment and Landscape Ecology as a Tool for Land Use Planning. *Freshwater Biology* 52 (5):908–17. <https://doi.org/10.1111/j.1365-2427.2006.01685.x>.
- Beck, M. W. 2018. SCCWRP/SCAPE: v1.0 (Version 1.0). Zenodo, <http://doi.org/10.5281/zenodo.1218121>.
- Beechie, T., G. Pess, P. Roni, and G. Giannico. 2007. Setting River Restoration Priorities: A Review of Approaches and General Protocol for Identifying and Prioritizing Actions. *North American Journal of Fisheries Management* 28 (3):891–905. <https://doi.org/10.1577/M06-174.1>.
- Breiman, L. 2001. Random Forests. *Machine Learning* 45:5–32.

662 Brody, S. D. 2003. Measuring the Effects of Stakeholder Participation on the Quality of Local
 663 Plans Based on the Principles of Collaborative Ecosystem Management. *Journal of Planning*
 664 *Education and Research* 22 (4):407–19. <https://doi.org/10.1177/0739456X03022004007>.

665 Brown, L. R., T. F. Cuffney, J. F. Coles, F. Fitzpatrick, G. McMahon, J. Steuer, A. H. Bell, and
 666 J. T. May. 2009. Urban Streams Across the USA: Lessons Learned from Studies in Nine
 667 Metropolitan Areas. *Journal of the North American Benthological Society* 28 (4):1051–69.
 668 <https://doi.org/10.1899/08-153.1>.

669 Brown, L. R., J. T. May, A. C. Rehn, P. R. Ode, I. R. Waite, and J. G. Kennen. 2012. Predicting
 670 Biological Condition in Southern California Streams. *Landscape and Urban Planning* 108
 671 (1):17–27. <https://doi.org/10.1016/j.landurbplan.2012.07.009>.

672 Buss, D. F., D. M. Carlisle, T. -S. Chon, J. Culp, J. s. Harding, H. E. Keizer-Vlek, W. A.
 673 Robinson, S. Strachan, C. Thirion, and R. M. Hughes. 2014. Stream Biomonitoring Using
 674 Macroinvertebrates Around the Globe: A Comparison of Large-Scale Programs. *Environmental*
 675 *Monitoring and Assessment* 187:4132. <https://doi.org/10.1007/s10661-014-4132-8>.

676 Cade, B. S., and B. R. Noon. 2003. A Gentle Introduction to Quantile Regression for Ecologists.
 677 *Frontiers in Ecology and the Environment* 1 (8):412–20.

678 Cao, Y., C. P. Hawkins, J. Olson, and M. A. Kosterman. 2007. Modeling Natural Environmental
 679 Gradients Improves the Accuracy and Precision of Diatom-Based Indicators. *Journal of the*
 680 *North American Benthological Society* 26 (3):566–85. <https://doi.org/10.1899/06-078.1>.

681 Carline, R. F., and M. C. Walsh. 2007. Responses to Riparian Restoration in the Spring Creek
 682 Watershed, Central Pennsylvania. *Restoration Ecology* 15 (4):731–42.
 683 <https://doi.org/10.1111/j.1526-100X.2007.00285.x>.

684 Carlisle, D. M., J. Falcone, and M. R. Meador. 2009. Predicting the Biological Condition of
 685 Streams: Use of Geospatial Indicators of Natural and Anthropogenic Characteristics of
 686 Watersheds. *Environmental Monitoring and Assessment* 151 (1-4):143–60.
 687 <https://doi.org/10.1007/s10661-008-0256-z>.

688 Chen, K., R. M. Hughes, S. Xu, J. Zhang, D. Cai, and B. Wang. 2014. Evaluating Performance
 689 of Macroinvertebrate-Based Adjusted and Unadjusted Multi-Metric Indices (MMI) Using Multi-
 690 Season and Multi-Year Samples. *Ecological Indicators* 36:142–51.
 691 <https://doi.org/10.1016/j.ecolind.2013.07.006>.

692 Chessman, B. C. 2014. Predicting Reference Assemblages for Freshwater Bioassessment with
 693 Limiting Environmental Difference Analysis. *Freshwater Science* 33 (4):1261–71.
 694 <https://doi.org/10.1086/678701>.

695 Chessman, B. C., M. Muschal, and M. J. Royal. 2008. Comparing Apples with Apples: Use of
 696 Limiting Environmental Differences to Match Reference and Stressor-Exposure Sites for
 697 Bioassessment of Streams. *River Research and Applications* 24 (1):103–17.
 698 <https://doi.org/10.1002/rra.1053>.

699 Chessman, B. C., and M. J. Royal. 2004. Bioassessment Without Reference Sites: Use of
 700 Environmental Filters to Predict Natural Assemblages of River Macroinvertebrates. *Journal of*
 701 *the North American Benthological Society* 23 (3):599–615. [https://doi.org/10.1899/0887-](https://doi.org/10.1899/0887-3593(2004)023%3C0599:BWRSUO%3E2.0.CO;2)
 702 [3593\(2004\)023%3C0599:BWRSUO%3E2.0.CO;2](https://doi.org/10.1899/0887-3593(2004)023%3C0599:BWRSUO%3E2.0.CO;2).

703 Clapcott, J. E., K. J. Collier, R. G. Death, E. O. Goodwin, J. S. Harding, D. Kelly, J. R.
 704 Leathwick, and R. G. Young. 2011. Quantifying Relationships Between Land-Use Gradients and
 705 Structural and Functional Indicators of Stream Ecological Integrity. *Freshwater Biology* 57
 706 (1):74–90. <https://doi.org/10.1111/j.1365-2427.2011.02696.x>.

707 Cormier, S. M., G. W. Suter II, L. Zhang, and G. J. Pond. 2013. Assessing Causation of the
 708 Extirpation of Stream Macroinvertebrates by a Mixture of Ions. *Environmental Toxicology and*
 709 *Chemistry* 32 (2):277–87. <https://doi.org/10.1002/etc.2059>.

710 CRWQB (California Regional Water Quality Control Board, Los Angeles Region). 2014. Basin
 711 Plan for the Coastal Watersheds of Los Angeles and Ventura Counties. California Regional
 712 Water Quality Control Board, Los Angeles, California.

713 Davies, S. P., and S. K. Jackson. 2006. The Biological Condition Gradient: A Descriptive Model
 714 for Interpreting Change in Aquatic Ecosystems. *Ecological Applications* 16 (4):1251–66.

715 Domisch, S., G. Amatulli, and W. Jetz. 2015. Near-Global Freshwater-Specific Environmental
 716 Variables for Biodiversity Analyses in 1 Km Resolution. *Scientific Data* 2:150073.
 717 <https://doi.org/10.1038/sdata.2015.73>.

718 ESRI (Environmental Systems Research Institute). 2016. ArcGIS v10.5. Redlands, California.

719 FDEP (Florida Department of Environmental Protection). 2011. Development of aquatic life use
 720 support attainment thresholds for Florida’s Stream Condition Index and Lake Vegetation Index.
 721 DEP-SAS-003/11. Tallahassee, Florida: FDEP Standards; Assessment Section, Bureau of
 722 Assessment; Restoration Support.

723 Finkenbine, J. K., J. W. Atwater, and D. S. Mavinic. 2000. Stream Health After Urbanization.
 724 *Journal of the American Water Resources Association* 36 (5):1149–60.
 725 <https://doi.org/10.1111/j.1752-1688.2000.tb05717.x>.

726 Fox, E. W., R. A. Hill, S. G. Leibowitz, A. R. Olsen, D. J. Thornbrugh, and M. H. Weber. 2017.
 727 Assessing the Accuracy and Stability of Variable Selection Methods for Random Forest
 728 Modeling in Ecology. *Environmental Monitoring and Assessment* 189:316.
 729 <https://doi.org/10.1007/s10661-017-6025-0>.

730 Fry, J., G. Xian, S. Jin, J. Dewitz, C. Homer, L. Yang, C. Barnes, N. Herold, and J. Wickham.
 731 2011. Completion of the 2006 National Land Cover Database for the Conterminous United
 732 States. *Photogrammetric Engineering and Remote Sensing* 77 (9):858–64.

733 Hastie, T., R. Tibshirani, and J. Friedman. 2009. *The Elements of Statistical Learning: Data*
 734 *Mining, Inference, and Prediction*. 2nd ed. New York: Springer.

735 Hill, R. A., E. W. Fox, S. G. Leibowitz, A. R. Olsen, D. J. Thornbrugh, and M. H. Weber. 2017.
 736 Predictive Mapping of the Biotic Condition of Conterminous U.S. Rivers and Streams.
 737 *Ecological Applications* 27 (8):2397–2415. <https://doi.org/10.1002/eap.1617>.

738 Hill, R. A., M. H. Weber, S. G. Leibowitz, A. R. Olsen, and D. J. Thornbrugh. 2016. The
 739 Stream-Catchment (StreamCat) Dataset: A Database of Watershed Metrics for the Conterminous
 740 United States. *Journal of the American Water Resources Association* 52:120–28.
 741 <https://doi.org/10.1111/1752-1688.12372>.

742 Howard, J. K., K. A. Fesenmyer, T. E. Grantham, J. H. Viers, P. R. Ode, P. B. Moyle, S. J.
 743 Kupferburg, et al. 2018. A Freshwater Conservation Blueprint for California: Prioritizing

744 Watersheds for Freshwater Biodiversity. *Freshwater Science* 37 (2):417–31.
745 <https://doi.org/10.1086/697996>.

746 Howard, J. K., K. R. Klausmeyer, K. A. Fesenmyer, J. Furnish, T. Gardali, T. Grantham, J. V. E.
747 Katz, et al. 2015. Patterns of Freshwater Species Richness, Endemism, and Vulnerability in
748 California. *PLOS ONE* 10 (7):e0130710. <https://doi.org/10.1371/journal.pone.0130710>.

749 Howard, J., and C. Revenga. 2009. California’s Freshwater Biodiversity in a Continental
750 Context. Science for Conservation Technical Brief Series. San Francisco, CA: The Nature
751 Conservancy of California.

752 Karr, J. R., K. D. Fausch, P. L. Angermeier, P. R. Yant, and I. J. Schlosser. 1986. Assessing
753 Biological Integrity in Running Waters: A Method and Its Rationale. Special Publication 5.
754 Champaign, Illinois: Illinois Natural History Survey.

755 Kenney, M. A., P. R. Wilcock, B. F. Hobbs, N. E. Flores, and D. C. Martínez. 2012. Is Urban
756 Stream Restoration Worth It? *Journal of the American Water Resources Association* 48 (3):603–
757 15. <https://doi.org/10.1111/j.1752-1688.2011.00635.x>.

758 Leps, M., J. D. Tonkin, V. Dahm, P. Haase, and A. Sundermann. 2015. Disentangling
759 Environmental Drivers of Benthic Invertebrate Assemblages: The Role of Spatial Scale and
760 Riverscape Heterogeneity in a Multiple Stressor Environment. *Science of the Total Environment*
761 536:546–56. <https://doi.org/10.1016/j.scitotenv.2015.07.083>.

762 Lester, R. E., and A. J. Boulton. 2008. Rehabilitating Agricultural Streams in Australia with
763 Wood: A Review. *Environmental Management* 42 (2):310–26.
764 <https://doi.org/10.1007%2Fs00267-008-9151-1>.

765 Loflen, C., H. Hettesheimer, L. B. Busse, K. Watanabe, R. M. Gersberg, and V. Lüderitz. 2016.
 766 Inadequate Monitoring and Inappropriate Project Goals: A Case Study on the Determination of
 767 Success for the Forester Creek Improvement Project. *Ecological Restoration* 34 (2):124–34.
 768 <https://doi.org/10.3368/er.34.2.124>.

769 Mazor, R. D., M. W. Beck, and J. Brown. 2018. 2017 Report on the SMC Regional Stream
 770 Survey. 1029. Costa Mesa, California: Southern California Coastal Water Research Project.

771 Mazor, R. D., A. C. Rehn, P. R. Ode, M. Engeln, K. C. Schiff, E. D. Stein, D. J. Gillett, D. B.
 772 Herbst, and C. P. Hawkins. 2016. Bioassessment in Complex Environments: Designing an Index
 773 for Consistent Meaning in Different Settings. *Freshwater Science* 35 (1):249–71.

774 MBI (Midwest Biodiversity Institute). 2016. Identification of predictive habitat attributes for
 775 Minnesota streams to support tiered aquatic life uses. MBI Technical Report
 776 MBI/OHPAN1518840. Columbus, Ohio: Midwest Biodiversity Institute, prepared on behalf of
 777 the Minnesota Pollution Control Agency.

778 McKay, L., T. Bondelid, T. Dewald, J. Johnston, R. Moore, and A. Reah. 2012. NHDPlus
 779 Version 2: User Guide.

780 Meinshausen, N. 2006. Quantile Regression Forests. *Journal of Machine Learning Research*
 781 7:983–99.

782 Meinshausen, Nicolai. 2017. *QuantregForest: Quantile Regression Forests*. [https://CRAN.R-](https://CRAN.R-project.org/package=quantregForest)
 783 [project.org/package=quantregForest](https://CRAN.R-project.org/package=quantregForest).

784 Morgan, R. P., and S. E. Cushman. 2005. Urbanization Effects on Stream Fish Assemblages in
 785 Maryland, USA. *Journal of the North American Benthological Society* 24 (3):643–55.

786 Moss, D., M. T. Furse, J. F. Wright, and P. D. Armitage. 1987. The Prediction of the Macro-
 787 Invertebrate Fauna of Unpolluted Running-Water Sites in Great Britain Using Environmental
 788 Data. *Freshwater Biology* 17 (1):41–52. <https://doi.org/10.1111/j.1365-2427.1987.tb01027.x>.

789 Nel, J. L., D. J. Roux, R. Abell, P. J. Ashton, R. M. Cowling, J. V. Higgins, M. Thieme, and J. H.
 790 Viers. 2009. Progress and Challenges in Freshwater Conversation Planning. *Aquatic*
 791 *Conservation: Marine and Freshwater Ecosystems* 19 (4):474–85.
 792 <https://doi.org/10.1002/aqc.1010>.

793 Novotny, V., A. Bartosová, N. O'Reilly, and T. Ehlinger. 2005. Unlocking the Relationship of
 794 Biotic Integrity of Impaired Waters to Anthropogenic Stresses. *Water Research* 39 (1):184–98.
 795 <https://doi.org/10.1016/j.watres.2004.09.002>.

796 Ode, P. R., A. C. Rehn, R. D. Mazor, K. C. Schiff, E. D. Stein, J. T. May, L. R. Brown, et al.
 797 2016. Evaluating the Adequacy of a Reference-Site Pool for Ecological Assessments in
 798 Environmentally Complex Regions. *Freshwater Science* 35 (1):237–48.

799 Paul, M. J., D. W. Bressler, A. H. Purcell, M. T. Barbour, E. T. Rankin, and V. H. Resh. 2008.
 800 Assessment Tools for Urban Catchments: Defining Observable Biological Potential. *Journal of*
 801 *the American Water Resources Association* 45 (2):320–30. [https://doi.org/10.1111/j.1752-](https://doi.org/10.1111/j.1752-1688.2008.00280.x)
 802 [1688.2008.00280.x](https://doi.org/10.1111/j.1752-1688.2008.00280.x).

803 Quinn, J. M., and A. E. Wright-Stow. 2008. Stream Size Influences Stream Temperature Impacts
 804 and Recovery Rates After Clearfell Logging. *Forest Ecology and Management* 256 (12):2101–9.
 805 <https://doi.org/10.1016/j.foreco.2008.07.041>.

806 RDCT (R Development Core Team). 2018. R: A language and environment for statistical
 807 computing, v3.4.4. R Foundation for Statistical Computing, Vienna, Austria.

808 Reed, M. S. 2008. Stakeholder Participation for Environmental Management: A Literature
 809 Review. *Biological Conservation* 141 (10):2417–31.
 810 <https://doi.org/10.1016/j.biocon.2008.07.014>.

811 Rehn, A. C., R. D. Mazor, and P. R. Ode. 2018. An Index to Measure the Quality of Physical
 812 Habitat in California Wadeable Streams. SWAMP Technical Memorandum, SWAMP-TM-2018-
 813 0005. Sacramento, California: California Water Boards, Surface Water Ambient Monitoring
 814 Program, California Department of Fish; Wildlife, Southern California Coastal Water Research
 815 Project.
 816 [https://www.waterboards.ca.gov/water_issues/programs/swamp/bioassessment/docs/physical_ha](https://www.waterboards.ca.gov/water_issues/programs/swamp/bioassessment/docs/physical_habitat_index_technical_memo.pdf)
 817 [bitat_index_technical_memo.pdf](https://www.waterboards.ca.gov/water_issues/programs/swamp/bioassessment/docs/physical_habitat_index_technical_memo.pdf).

818 Reynoldson, T. B., R. H. Norris, V. H. Resh, K. E. Day, and D. M. Rosenberg. 1997. The
 819 Reference Condition: A Comparison of Multimetric and Multivariate Approaches to Assess
 820 Water-Quality Impairment Using Benthic Macroinvertebrates. *Journal of the North American*
 821 *Benthological Society* 16 (4):833–52. <https://doi.org/10.2307/1468175>.

822 Riseng, C. M., M. J. Wiley, R. W. Black, and M. D. Munn. 2011. Impacts of Agricultural Land
 823 Use on Biological Integrity: A Causal Analysis. *Ecological Applications* 21 (8):3128–46.
 824 <https://doi.org/10.1890/11-0077.1>.

825 Roni, P., and T. Beechi. 2012. *Stream and Watershed Restoration Guide: A Guide to Restoring*
 826 *Riverine Processes and Habitats*. First. Hoboken, New Jersey: John Wiley & Sons.

827 SDRWQB (San Diego Regional Water Quality Control Board). 2001. Waste discharge
828 requirements for discharges of urban runoff from the municipal separate storm sewer systems
829 (MS4s) draining the watersheds of the County of San Diego, the incorporated cities of San Diego
830 County, and the San Diego unified port district. Sacramento, California: California
831 Environmental Protection Agency.

832 SDRWQB (San Diego Regional Water Quality Control Board). 2016. Clean Water Act Sections
833 305(B) and 303(D) Integrated Report for the San Diego Region. Sacramento, California:
834 California Environmental Protection Agency.
835 https://www.waterboards.ca.gov/sandiego/water_issues/programs/303d_list/docs/Staff_Report_1
836 [01216.pdf](https://www.waterboards.ca.gov/sandiego/water_issues/programs/303d_list/docs/Staff_Report_101216.pdf).

837 Shoredits, A. S., and J. A. Clayton. 2013. Assessing the Practice and Challenges of Stream
838 Restoration in Urbanized Environments of the USA. *Geography Compass* 7 (5):358–72.
839 <https://doi.org/10.1111/gec3.12039>.

840 Sleeter, B. M., T. S. Wilson, C. E. Soulard, and J. Liu. 2011. Estimation of the Late Twentieth
841 Century Land-Cover Change in California. *Environmental Monitoring and Assessment* 173 (1-
842 4):251–66. <https://doi.org/10.1007/s10661-010-1385-8>.

843 Sponseller, R. A., E. F. Benfield, and H. M. Valett. 2001. Relationships Between Land Use,
844 Spatial Scale and Stream Macroinvertebrate Communities. *Freshwater Biology* 46 (10):1409–24.
845 <https://doi.org/10.1046/j.1365-2427.2001.00758.x>.

846 Stein, E. D., M. R. Cover, A. E. Fetscher, C. O'Reilly, R. Guardado, and C. W. Solek. 2013.
847 Reach-Scale Geomorphic and Biological Effects of Localized Streambank Armoring. *Journal of*
848 *the American Water Resources Association* 49 (4):780–92. <https://doi.org/10.1111/jawr.12035>.

849 Stein, E. D., A. Sengupta, R. D. Mazon, K. McCune, B. P. Bledsoe, and S. Adams. 2017.
 850 Application of Regional Flow-Ecology Relationships to Inform Watershed Management
 851 Decisions: Applications of the ELOHA Framework in the San Diego River Watershed,
 852 California, USA. *Ecohydrology* 10 (7):e1869. <https://doi.org/10.1002/eco.1869>.

853 Stoddard, J. L., D. P. Larsen, C. P. Hawkins, R. K. Johnson, and R. H. Norris. 2006. Setting
 854 Expectations for the Ecological Condition of Streams: The Concept of Reference Condition.
 855 *Ecological Applications* 16 (4):1267–76. [https://doi.org/10.1890/1051-](https://doi.org/10.1890/1051-0761(2006)016[1267:SEFTEC]2.0.CO;2)
 856 [0761\(2006\)016\[1267:SEFTEC\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2006)016[1267:SEFTEC]2.0.CO;2).

857 Stone, M. K., and J. B. Wallace. 1998. Long-Term Recovery of a Mountain Stream from Clear-
 858 Cut Logging: The Effects of Forest Succession on Benthic Invertebrate Community Structure.
 859 *Freshwater Biology* 39 (1):151–69. <https://doi.org/10.1046/j.1365-2427.1998.00272.x>.

860 Townsend, C. R., S. S. Uhlmann, and C. D. Matthaei. 2008. Individual and Combined Responses
 861 of Stream Ecosystems to Multiple Stressors. *Journal of Applied Ecology* 45 (6):1810–9.
 862 <https://doi.org/10.1111/j.1365-2664.2008.01548.x>.

863 USEPA (US Environmental Protection Agency, Region 10). 2013. Technical Support Document
 864 for EPA’s Action on the State of Oregon’s Revised Water Quality Standards for the West
 865 Division Main Canal. USEPA Region 10, Seattle, Washington.

866 USEPA (US Environmental Protection Agency). 2016. National Rivers and Streams Assessment
 867 2008-2009: A Collaborative Survey. EPA-841-R-16-007. Washington, DC.

868 USGS (US Geological Survey). 1999. The quality of our nation’s waters: nutrients and
 869 pesticides. Reston, Virginia.

870 Vølstad, J. H., N. E. Roth, G. Mercurio, M. T. Southerland, and D. E. Strebel. 2004. Using
871 Environmental Stressor Information to Predict the Ecological Status of Maryland Non-Tidal
872 Streams as Measured by Biological Indicators. *Environmental Monitoring and Assessment* 84
873 (3):219–42. <https://doi.org/10.1023/A:1023374524254>.

874 Waite, I. R. 2013. Development and Application of an Agricultural Intensity Index to
875 Invertebrate and Algal Metrics from Streams at Two Scales. *Journal of the American Water*
876 *Resources Association* 49 (2):431–48. <https://doi.org/10.1111/jawr.12032>.

877 Waite, I. R., J. G. Kennen, J. T. May, L. R. Brown, T. F. Cuffney, K. A. Jones, and J. L.
878 Orlando. 2014. Stream Macroinvertebrate Response Models for Bioassessment Metrics:
879 Addressing the Issue of Spatial Scale. *PLOS ONE* 9 (3):e90944.
880 <https://doi.org/10.1371/journal.pone.0090944>.

881 Walsh, C. J., A. H. Roy, J. w. Feminella, P. D. Cottingham, P. M. Groffman, and R. P. Morgan.
882 2005. The Urban Stream Syndrome: Current Knowledge and the Search for a Cure. *Journal of*
883 *the North American Benthological Society* 24 (3):706–23. <https://doi.org/10.1899/04-028.1>.

884 Wang, L., J. Lyons, P. Kanehl, and R. Gatti. 1997. Influences of Watershed Land Use on Habitat
885 Quality and Biotic Integrity in Wisconsin Streams. *Fisheries* 22 (6):6–12.
886 [https://doi.org/10.1577/1548-8446\(1997\)022%3C0006:IOWLUO%3E2.0.CO;2](https://doi.org/10.1577/1548-8446(1997)022%3C0006:IOWLUO%3E2.0.CO;2).

887

888

Figure captions

Figure 1 Urban and agricultural land use (a) and distribution of observed stream CSCI scores (b) in California. Cover of urban and agricultural land use in stream watersheds was used to develop a landscape model for stream segment expectations of bioassessment scores. Breakpoints for CSCI scores are the 1st, 10th, and 30th percentile of scores at least-disturbed, reference sites throughout the state. Altered and intact refers to biological condition (Mazor et al. 2016). Grey lines are major environmental regions in California defined by ecoregional and watershed boundaries, CV: Central Valley, CH: Chaparral, DM: Deserts and Modoc Plateau, NC: North Coast, SN: Sierra Nevada, SC: South Coast.

Figure 2 Application of the landscape model to identify site expectations and bioassessment performance for sixteen example stream segments. A range of CSCI scores is predicted from the model (a) and the lower and upper limits of the expectations are cut to define a certainty range for the predictions (b). Overlap of the certainty range at each segment with a chosen CSCI threshold (c) defines the stream segment classification as likely unconstrained, possibly unconstrained, possibly constrained, and likely constrained. The observed bioassessment scores are described relative to the classification as over scoring (above the certainty threshold), expected (within), and under scoring (below) for each of four stream classes (d).

Figure 3 San Gabriel River watershed in southern California. Land cover is shown in plot (a) and the predicted median CSCI scores at each stream segment and observed CSCI scores are shown in (b).

Figure 4 Statewide application of the landscape model showing the stream segment classifications. Major regional boundaries are also shown (see Figure 1).

Figure 5 Changes in stream segment classes by region and statewide for different scenarios used to define biological constraints. Twenty-seven scenarios were tested that evaluated different combinations of certainty in the CSCI predictions (nine scenarios from wide to narrow prediction intervals as identified by the tail cutoff for the expected range) and potential CSCI thresholds (three scenarios from low to high). The percentage of total stream length for likely unconstrained and likely constrained is shown for each scenario. Stream classifications as possibly unconstrained or possibly constrained are not shown but can be inferred from the area of white space above or below each bar. The solid black line indicates the percentage division between unconstrained and constrained classifications. CV: Central Valley, CH: Chaparral, DM: Deserts and Modoc Plateau, NC: North Coast, SN: Sierra Nevada, SC: South Coast.

Figure 6 Application of the landscape model to stream segments in the San Gabriel River watershed, Los Angeles County, California. CSCI scores with (a) no context from the model are on the left and (b) scores with context from the model are on the right. Relative site scores as under-scoring, expected, or over-scoring are based on observed scores given the segment class as likely constrained, possibly constrained, possibly unconstrained, and likely unconstrained. Segment classes are based on overlap of the expectations with a biological threshold for the CSCI (0.79, dashed lined) and location of the median expectation (white ticks).

Figure 7 Relative site scores and recommended management actions for locations with CSCI scores in the San Gabriel River watershed. Relative site scores as under scoring, expected, or over scoring are based on observed scores given the segment class as likely constrained, possibly constrained, possibly unconstrained, and likely unconstrained. Recommended management actions were defined by a local stakeholder group (see Figure S1) and are ranked

933 *by priority for actions to investigate, protect, and restore a site. No recommended actions*
934 *assume baseline maintenance and monitoring is sufficient.*

935 *Figure 8 Tecolote Creek (San Diego County, USA) is a constrained channel in an urban*
936 *landscape (a, Source: 32.81736, -117.19986. Google Earth. November 8, 2016. Accessed July*
937 *20, 2018.). Physical habitat (b, Source: R. Mazor) at the sample site suggests no channel*
938 *alteration. The CSCI was scored at 0.61 indicating degraded biological integrity.*

Tables

Table 1 Land use variables used to develop the landscape model of stream bioassessment scores. All variables were obtained from StreamCat (Hill et al. 2016) and applied to stream segments in the National Hydrography Dataset Plus (NHD-plus) (McKay et al. 2012). The measurement scales for each variable are at the riparian (100 m buffer), catchment, and/or watershed, scale relative to a stream segment. Combined scales for riparian measurements (e.g., riparian + catchment, riparian + watershed) are riparian estimates for the entire catchment or watershed area upstream, as compared to only the individual segment. Total urban and agriculture land use variables were based on sums of individual variables in StreamCat as noted in the description. Rp100: riparian, Cat: catchment, Ws: watershed

Name	Scale	Description	Unit
CanalDens	Cat, Ws	Density of NHDPlus line features classified as canal, ditch, or pipeline	km/sq km
PctImp2006	Cat, Ws, Cat + Rp100, Ws + Rp100	Mean imperviousness of anthropogenic surfaces (NLCD 2006)	%
TotUrb2011	Cat, Ws, Cat + Rp100, Ws + Rp100	Total urban land use as sum of developed open, low, medium, and high intensity (NLCD 2011)	%
TotAg2011	Cat, Ws, Cat + Rp100, Ws +	Total agricultural land use as sum of hay and crops (NLCD 2011)	%

	Rp100		
RdDens	Cat, Ws, Cat + Rp100, Ws + Rp100	Density of roads (2010 Census Tiger Lines)	km/sq km
RdCrs	Cat, Ws	Density of roads-stream intersections (2010 Census Tiger Lines-NHD stream lines)	crossings/sq km

950

951

952 *Table 2 Stream class definitions describing potential biological constraints. Classes are based*
 953 *on the overlap of the range of likely bioassessment scores with a potential threshold for a*
 954 *biological objective. Identifying stream classes requires selecting the cutoff range of likely*
 955 *scores from the landscape model and a chosen threshold for the objective.*

Class	Definition	Example
Likely unconstrained	Lower bound of prediction interval is above threshold	10 th percentile > 0.79
Possibly unconstrained	Lower bound of prediction interval is below threshold, but median prediction is above	50 th percentile > 0.79
Possibly constrained	Upper bound of prediction interval is above threshold, but median prediction is below	50 th percentile < 0.79
Likely constrained	Upper bound of prediction interval is below threshold	90 th percentile < 0.79

956

957

958 *Table 3 Performance of the landscape model by calibration (Cal) and validation (Val) datasets*
959 *in predicting CSCI scores. The statewide dataset (Figure 4) and individual regions of California*
960 *(Figure 1) are evaluated. Averages and standard deviations (in parentheses) for observed and*
961 *predicted CSCI values of each dataset are shown. Pearson correlations (r), root mean squared*
962 *errors (RMSE), intercept, and slopes are for comparisons of predicted and observed values to*
963 *evaluate model performance. All correlations, intercepts, and slopes are significant at alpha =*
964 *0.05. CV: Central Valley, CH: Chaparral, DM: Deserts and Modoc Plateau, NC: North Coast,*
965 *SN: Sierra Nevada, SC: South Coast.*

Dataset	Location	n	Observed	Predicted	r	RMSE	Intercept	Slope
Cal	Statewide	1965	0.82 (0.26)	0.83 (0.20)	0.75	0.17	0.34	0.60
	CH	512	0.76 (0.27)	0.79 (0.21)	0.71	0.19	0.38	0.54
	CV	116	0.51 (0.18)	0.57 (0.15)	0.66	0.15	0.29	0.54
	DM	86	0.87 (0.22)	0.91 (0.14)	0.50	0.20	0.63	0.31
	NC	208	0.92 (0.20)	0.94 (0.13)	0.55	0.17	0.61	0.36
	SC	631	0.79 (0.24)	0.78 (0.21)	0.75	0.16	0.27	0.65
	SN	412	0.98 (0.18)	0.98 (0.09)	0.45	0.16	0.75	0.23
Val	Statewide	655	0.82 (0.25)	0.84 (0.20)	0.72	0.18	0.36	0.58
	CH	172	0.76 (0.27)	0.81 (0.21)	0.74	0.19	0.39	0.56
	CV	40	0.52 (0.19)	0.59 (0.16)	0.49	0.19	0.38	0.40
	DM	28	0.84 (0.17)	0.93 (0.11)	0.55	0.17	0.63	0.36
	NC	71	0.94 (0.19)	0.96 (0.11)	0.55	0.16	0.67	0.31

SC	208	0.80 (0.24)	0.78 (0.21)	0.72	0.17	0.27	0.63
SN	136	0.97 (0.17)	0.98 (0.09)	0.21	0.17	0.88	0.11

966

967

968 *Table 4 Summary of stream length for each stream class statewide and major regions of*
969 *California (Figures 1, 4). Lengths are in kilometers with the percentage of the total length in a*
970 *region in parentheses. All lengths are based on a CSCI threshold of 0.79 and the 10th to 90th*
971 *percentile of expected scores from the landscape model. CV: Central Valley, CH: Chaparral,*
972 *DM: Deserts and Modoc Plateau, NC: North Coast, SN: Sierra Nevada, SC: South Coast.*

	constrained		unconstrained	
Region	likely	possibly	possibly	likely
Statewide	8150 (4)	24735 (11)	101591 (46)	85317 (39)
CV	3356 (22)	8010 (52)	3202 (21)	951 (6)
CH	1642 (3)	7840 (13)	30693 (50)	21206 (35)
DM	255 (0)	3395 (6)	27194 (47)	26479 (46)
NC	108 (0)	1442 (5)	14152 (49)	13286 (46)
SN	20 (0)	1067 (3)	18228 (48)	19032 (50)
SC	2770 (15)	2981 (16)	8122 (45)	4363 (24)

973



974

975 *Table 5 Summary of CSCI scores by relative expectations for each stream class statewide and in*
976 *each major region of California (Figures 1, 4). Average CSCI scores (standard deviation) and*
977 *counts (percent) of the number of monitoring stations in each relative score category and region*
978 *are shown. Sites are over-scoring if the observed scores are above the range of expectations at a*
979 *segment, expected if within the range, or under-scoring if below the range. CV: Central Valley,*
980 *CH: Chaparral, DM: Deserts and Modoc Plateau, NC: North Coast, SN: Sierra Nevada, SC:*
981 *South Coast.*



	under-scoring		expected		over-scoring	
Region	CSCI	n (%)	CSCI	n (%)	CSCI	n (%)
Statewide	0.54 (0.21)	267 (10)	0.83 (0.23)	2041 (80)	1.08 (0.17)	242 (9)
CH	0.47 (0.18)	89 (13)	0.79 (0.24)	535 (80)	1.08 (0.17)	45 (7)
CV	0.34 (0.12)	25 (17)	0.54 (0.17)	118 (81)	0.63 (0.25)	2 (1)
DM	0.6 (0.17)	15 (14)	0.9 (0.17)	89 (80)	1.15 (0.08)	7 (6)
NC	0.66 (0.17)	28 (10)	0.93 (0.16)	228 (82)	1.15 (0.08)	22 (8)
SC	0.54 (0.22)	56 (7)	0.78 (0.22)	656 (81)	1.02 (0.2)	97 (12)
SN	0.67 (0.16)	54 (10)	0.99 (0.11)	415 (77)	1.16 (0.06)	69 (13)

982


(a) Land use and regions




(b) CSCI scores




Figure



(a) Segment median predictions and observed scores for the CSCI




(b) Land cover in the San Gabriel watershed




Figure

Segment classification


- | | |
|--------------------------|------------------------|
| — likely unconstrained | — possibly constrained |
| — possibly unconstrained | — likely constrained |




Figure



Figure



Figure



(a) Satellite view of Tecolote Creek and surrounding land use, San Diego County, USA



(b) Physical habitat conditions at Tecolote Creek



